Math. Struct. in Comp. Science (2018), *vol.* 28, *pp.* 1639–1694. © Cambridge University Press 2017
doi:10.1017/S0960129517000196 First published online 28 September 2017

On geometry of interaction for polarized linear logic

MASAHIRO HAMANO[†] and PHILIP SCOTT[‡]

[†]*PRESTO, Japan Science and Technology Agency (JST),*
4-1-8 Honcho Kawaguchi, Saitama 332-0012, Japan
Email: hamano@jaist.ac.jp
[‡]*Department of Mathematics & Statistics, University of Ottawa,*
585 King Edward, Ottawa, Ontario, K1N 6N5, Canada
Email: phil@site.uottawa.ca

Received 25 October 2013; revised 4 August 2016

We present Geometry of Interaction (GoI) models for Multiplicative Polarized Linear Logic, MLLP, which is the multiplicative fragment of Olivier Laurent's Polarized Linear Logic. This is done by uniformly adding *multi-points* to various categorical models of GoI. Multi-points are shown to play an essential role in semantically characterizing the dynamics of proof networks in polarized proof theory. For example, they permit us to characterize the key feature of polarization, *focusing*, as well as being fundamental to our construction of concrete polarized GoI models.
Our approach to polarized GoI involves following two independent studies, based on different categorical perspectives of GoI:

(i) Inspired by the work of Abramsky, Haghverdi and Scott, a *polarized GoI situation* is defined in which multi-points are added to a traced monoidal category equipped with a reflexive object U. Using this framework, categorical versions of Girard's execution formula are defined, as well as the GoI interpretation of MLLP proofs. Running the execution formula is shown to characterize the focusing property (and thus polarities) as well as the dynamics of cut elimination.
(ii) The Int construction of Joyal–Street–Verity is another fundamental categorical structure for modelling GoI. Here, we investigate it in a multi-pointed setting. Our presentation yields a compact version of Hamano–Scott's polarized categories, and thus denotational models of MLLP. These arise from a contravariant duality between monoidal categories of *positive* and *negative* objects, along with an appropriate bimodule structure (representing 'non-focused proofs') between them.

Finally, as a special case of (ii) above, a compact model of MLLP is also presented based on Rel (the category of sets and relations) equipped with multi-points.

1. Introduction

Linear Logic, introduced by Girard (1987), originated from a profound analysis of the proof theory of traditional logic. In particular, linear logic involves a fine-grained study of how rules and connectives manipulate resources. This important development is by now quite familiar to researchers in many areas of logic and computer science. What is somewhat less familiar is that shortly after the introduction of linear logic, Andreoli (1992, 2001) pointed out a different approach to the fundamental connectives of linear

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

logic; namely, to classify the connectives according to whether their introduction rules are reversible (*negative*) or irreversible (*positive*). Positive connectives are the foundation of Andreoli's influential notion of *focusing* in proof search. The fundamental role of focusing in logic programming has been actively explored in numerous recent works (for example, in papers of Chaudhuri 2008, Chaudhuri *et al.* 2013, Liang and Miller 2009, Miller 2004, 2011).

The Andreoli view also led to intrinsic studies of polarity and polarized logics, first taken up by Girard (1991, 1999), and systematically studied by Laurent (1999, 2002). Such logics are also related to Girard's Ludics games Girard (2001) and other games-semantics models (cf. the dialogue games and recent dialogue categories with chiralities of Melliès 2006–2012 and 2013). In related categorical proof theory, we should mention the polarized categories and proof theory of Cockett and Seely (2007), which influenced our own (Hamano and Scott 2007), as well as proof-theoretical papers of one of us (Hamano and Takemura 2008, 2010).

In this paper, we begin a study of the dynamics of cut elimination for the multiplicative fragment MLLP of O. Laurent's polarized linear logic, using categorical versions of Girard's Geometry of Interaction (GoI) programme (Girard 1989, 1995); here, we follow the categorical GoI literature (Abramsky *et al.* 2002; Haghverdi and Scott 2006). A common theme to the different parts of this paper is a fundamental new semantical idea first discussed in Hamano and Takemura (2008): the addition of *multi-points*. These multi-points have no syntactic counterpart but nevertheless provide a new understanding of the dynamics of cut elimination in the presence of polarities.

— In Section 3, following the methods of Abramsky *et al.* (2002), we introduce *polarized GoI situations with multi-points* as an appropriate but simple categorical setting for studying Girard's execution formula in the polarized setting. The version of the execution formula we use is inspired by the general categorical execution formula in Haghverdi and Scott (2006, 2011) for linear logic, but now extended to the multi-pointed setting. For the polarized multiplicative system MLLP, the execution formula becomes a *two-layered* pair of execution formulas, one at the usual reflexive object level (as in Haghverdi and Scott 2006) and a similar one at the multi-point level. The usual GoI properties for the dynamics of cut elimination (Girard 1989; Haghverdi and Scott 2006, 2011) turn out to be much stronger for MLLP. The execution formula(s) form *full invariants* for normalization[†], which is well known to fail in full linear logic (see Abramsky *et al.* 2002 and Section 3.4 below). In fact, the polarized execution formula(s) satisfy a fundamental additional property. Namely, in Proposition 3.28 below, we characterize focusing, which is intrinsic to MLLP, as preservation of multi-points under the (polarized) execution formulas. Thus, in a precise sense, the execution formulas give rise to the polarities.

— In the next Sections 4 and 5, which are independent of Section 3, we consider general multi-pointed traced monoidal categories (TMCs) and study the Int construction of Joyal *et al.* (1996) in this setting. As is well known (Abramsky *et al.* 2002;

[†] In the sense that if $\pi \to_* \pi'$ by MLLP cut elimination, then $\mathsf{Ex}(\llbracket\pi\rrbracket, \sigma) = \mathsf{Ex}(\llbracket\pi'\rrbracket, \sigma')$.

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

Haghverdi and Scott 2006), the Int construction is an essential feature of all the different categorical approaches to GoI. It yields a kind of 'compact closure' of a TMC; moreover, composition in the Int category leads to categorical versions of Girard's execution formula. As in Section 3 above, we investigate a two-layered Int construction, the upper layer for general objects and the lower one restricted to multi-points.

In Section 4, we study a general categorical semantics for certain polarized linear logics, a simplified version of the bi-module duality framework in our paper (Hamano and Scott 2007). This is related to more general polarized categories introduced for modelling polarities (see Cockett and Seely 2007, Hamano and Scott 2007, and Melliès 2006–2012; Melliès 2013). Our goal is to use GoI and the Int construction to build compact polarized categorical models of MLLP.

In these sections, multi-points give a semantical framework for explaining the idea of bidirectional dataflow implicit in the Int construction (see the discussion preceding Proposition 4.8 below). In this setting, multi-points satisfy a certain commutativity condition – corresponding to the focusing condition discussed in Section 3 – which we show is compatible with this construction. As one expects, this yields an appropriate compact closed version of a denotational model for MLLP.

Finally, Section 5 constructs a concrete instance of the Int construction of Section 4, when specialized to a multi-pointed version of Rel. It may be read independently of the previous section and uses the relational calculus of Joyal *et al.* (1996).

— In order to make the paper self-contained, the Appendices include some supplemental material. In Appendix A, we briefly recall the original *GoI situations* in the sense of Abramsky *et al.* (2002), as well as the categorical approach to GoI (for a survey, see Haghverdi and Scott 2011). In Appendix B, we recall Haghverdi's Unique Decomposition Categories (UDCs) which provide a general framework (and matrix calculus) for 'particle-style' GoI, familiar from Girard's GoI 1 (Girard 1989; Haghverdi and Scott 2006). The other Appendices include some detailed but routine proofs.

2. Polarized multiplicative linear logic MLLP

Following the work of Andreoli, polarities naturally arise within the proof theory of linear logic, LL. We can further divide the connectives according to whether their introduction rules are reversible or not (Girard 1999; Laurent 2002). Those connectives which are reversible are called *negative*; those which are not are called *positive*. Positive connectives are the foundation of Andreoli's influential *focusing property* in proof search for linear logic (Andreoli 1992, 2001; Miller 2004). Focusing is a property dual to reversibility.

We recall Olivier Laurent's theory of *polarized multiplicative linear logic*, MLLP. The theory MLLP is a fragment (without structural rules) of Laurent's full polarized linear logic LLP Girard (2001).

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

Definition 2.1 (Polarized MLL). The theory MLLP is defined as follows.

Syntax: *Positive* and *Negative* formulas are given by the following BNF notation:

$$\begin{array}{rcl} P & ::= & X \mid P \otimes P \mid \mathbf{1} \mid \downarrow N \\ N & ::= & X^{\perp} \mid N \parr N \mid \perp \mid \uparrow P \end{array}$$

Here, X is an atom, $\uparrow$ and $\downarrow$ are called *polarity shifting* operations. Note that $\mathbf{1}$ is the unit of $\otimes$ (and dually for $\perp$ with respect to $\parr$). In our categorical models introduced later, $\uparrow$ and $\downarrow$ will be functorial operators weaker than the traditional exponentials of linear logic.

Syntactic negation: Following O. Laurent, we adjoin to MLLP a syntactic strictly involutive negation $(-)^{\perp}$, defined by general de Morgan duality. Thus, we extend the negation on positive atoms to all formulas, as follows: $X^{\perp\perp} = X$ for atoms X, and we assume $\{\otimes, \parr\}$ are de Morgan duals, as in linear logic. Finally, de Morgan duals for polarity changing operators are $(\downarrow A)^{\perp} = \uparrow A^{\perp}$ and $(\uparrow A)^{\perp} = \downarrow A^{\perp}$ for any formula A. Positivity and negativity of formulas may be defined as before, after cancelling any occurrences of double negations.

Rules of MLLP

In the following rules, M and N (resp. $\mathcal{M}$ and $\mathcal{N}$) range over negative formulas (resp. sequences of negative formulas) and P and Q over positive formulas. Γ contains *at most one* positive formula. We assume the rule of exchange, so sequents are closed under permutation.

$$\frac{}{\vdash N, N^{\perp}}\ \text{Axiom} \qquad \frac{\vdash \mathcal{M}, P \quad \vdash \mathcal{N}, Q}{\vdash \mathcal{M}, \mathcal{N}, P \otimes Q}\ \otimes \qquad \frac{\vdash \Gamma, N, M}{\vdash \Gamma, N \parr M}\ \parr$$

$$\frac{\vdash N, \mathcal{N}}{\vdash \downarrow N, \mathcal{N}}\ \downarrow \qquad \frac{\vdash P, \mathcal{N}}{\vdash \uparrow P, \mathcal{N}}\ \uparrow \qquad \frac{\vdash \Gamma, N \quad \vdash \mathcal{M}, N^{\perp}}{\vdash \Gamma, \mathcal{M}}\ \text{cut}$$

The following theorem is an important proof-theoretical property of MLLP, proved in Laurent (1999, 2002):

Proposition 2.2 (Focalization property). *If* $\vdash \Gamma$ *is provable in* MLLP*, then the sequence* Γ *contains at most one positive formula.*


A *focused sequent* is one of the form $\vdash P, \mathcal{N}$, while a *non-focused sequent* has the form $\vdash \mathcal{N}$, where $\mathcal{N}$ is a finite sequence of negative formulas and P is positive. Proposition 2.2 says that every provable sequent of MLLP is either focused or non-focused. We say a proof is focused if the sequent it proves is focused. We say a proof *has the focusing property* if it is focused.

Notation: For the discussion of GoI and cut elimination, we use the Girard notation (Girard 1989) for sequent calculus proofs: a proof of the sequent $\vdash [\Delta], \Gamma$ denotes a proof of the sequent $\vdash \Gamma$ appended with the list Δ of all pairs of cut formulas $A, A^{\perp}$ used in the proof. Here, $|\Gamma| = n$ and $|\Delta| = 2m$, for some m, n. This is further described in the categorical GoI papers of Haghverdi–Scott (see Appendix A and Figure 6 there.)

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

Cut elimination for MLLP**:** MLLP is the subsystem of polarized linear logic LLP (Laurent 2002) without additive connectives and structural rules (where ↓ and ↑ replace, respectively, ! and ?). The cut-elimination theorem for MLLP is obtained by restricting the one for LLP (cf. Definition 5.27 of Laurent 2002). For this, the interpretation using polarized proof-nets (e.g. Laurent 1999) is essential. The crucial ingredient for our subsystem MLLP is the use of *boxes* to interpret the ↓ rules. Each box has a principal door and an auxiliary door on which occur, respectively, the principal formula $\downarrow N$ and the side formulas $\mathcal{M}$.

The crucial step for cut elimination is the following (cf. the !/! case of Definition 5.27 of Laurent 2002): reduction of a cut against a side formula of the ↓ rule. Here, a proof ending with a cut against a formula at an auxiliary door of a box is reduced to that ending with the ↓ rule, whose box is enlarged to contain all rules including the original cut. This is illustrated as follows:




Example 2.3 (Extrusion of ↓ boxes through cut elimination). Consider the following three proofs, as instances of normalization: $\pi_1 \succ \pi_2 \succ \pi_3$.

$$\pi_1 = \dfrac{\dfrac{\dfrac{\vdash X_1^\perp, X_1}{\vdash X_1^\perp, \uparrow X_1}}{\vdash \downarrow X_1^\perp, \uparrow X_1}\downarrow \qquad \vdots \; \vdash \downarrow X_2^\perp, \uparrow X_2}{\vdash [\uparrow X_1, \downarrow X_2^\perp], \downarrow X_1^\perp, \uparrow X_2}\,\text{cut} \qquad \vdots \; \vdash \downarrow X_3^\perp, \uparrow X_3}{\vdash [\uparrow X_1, \downarrow X_2^\perp, \uparrow X_2, \downarrow X_3^\perp], \downarrow X_1^\perp, \uparrow X_3}\,\text{cut}$$

$$\pi_2 = \dfrac{\dfrac{\dfrac{\vdash X_1^\perp, X_1}{\vdash X_1^\perp, \uparrow X_1} \qquad \vdots \; \vdash \downarrow X_2^\perp, \uparrow X_2}{\vdash [\uparrow X_1, \downarrow X_2^\perp], X_1^\perp, \uparrow X_2}\,\text{cut}}{\vdash [\uparrow X_1, \downarrow X_2^\perp], \downarrow X_1^\perp, \uparrow X_2}\downarrow \qquad \vdots \; \vdash \downarrow X_3^\perp, \uparrow X_3}{\vdash [\uparrow X_1, \downarrow X_2^\perp, \uparrow X_2, \downarrow X_3^\perp], \downarrow X_1^\perp, \uparrow X_3}\,\text{cut}$$

$$\pi_3 = \dfrac{\dfrac{\dfrac{\vdash X_1^\perp, X_1}{\vdash X_1^\perp, \uparrow X_1} \qquad \vdots \; \vdash \downarrow X_2^\perp, \uparrow X_2}{\vdash [\uparrow X_1, \downarrow X_2^\perp], X_1^\perp, \uparrow X_2}\,\text{cut} \qquad \vdots \; \vdash \downarrow X_3^\perp, \uparrow X_3}{\vdash [\uparrow X_1, \downarrow X_2^\perp, \uparrow X_2, \downarrow X_3^\perp], X_1^\perp, \uparrow X_3}\,\text{cut}}{\vdash [\uparrow X_1, \downarrow X_2^\perp, \uparrow X_2, \downarrow X_3^\perp], \downarrow X_1^\perp, \uparrow X_3}\downarrow$$



https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

During normalization, boxes enter outer boxes and the scope of the outer box is extruded. That is, for $i = 1, 2, 3$, we represent proof π_i by the proof-net obtained by choosing box_i for the left-most $\downarrow$. The box_i corresponds to the indicated $\downarrow$ rule in π_i. Normalization gives rise to extrusions of box_1 into box_2 to include the middle box of π_2 and then into box_3 to include both the middle and right boxes of π_3.

3. Polarized GoI and GoI situations

GoI situations were first introduced in Abramsky *et al.* (2002) as an algebraic framework for Girard's GoI for full linear logic. Later, Haghverdi and Scott (2006, 2011) gave a detailed categorical analysis of Girard's original GoI for MELL in GoI situations associated to UDCs, augmented with abstract (Hyland–Schalk) orthogonality relations. For a quick summary of this categorical GoI for the original (non-polarized) setting, see Appendix A (and its references) for GoI and Appendix B for UDCs. In particular, we will make use of the categorical version of the execution formula.

In this section, we develop an appropriate notion of *polarized GoI situation with multi-points* for MLLP. Multi-points will enable us to give a detailed analysis of the dynamics and information flow in cut elimination, via polarized (two-layered) execution formulas. It will also enable us to give a characterization of focusing. Our main result in this Section 3 is to semantically characterize the proof-theoretical notion of focusing in terms of the polarized execution formulas. This goes as follows. First, in Theorem 3.26, the polarized execution formulas for focused sequents are shown to satisfy an invariance property preserving the multi-points. Conversely in Proposition 3.28, this invariance property is shown to be sufficient to distinguish focused sequents. Thus, in a precise sense, polarized GoI execution formulas give rise to the polarities.

Notation: In a category $\mathcal{C}$, we denote identity arrows $X \to X$ either as Id_X or just X, depending on context. Let $Y \underset{f}{\overset{g}{\rightleftarrows}} X$ be a pair of morphisms in a category. We write $g : Y \rhd X : f$ to mean $g \circ f = \mathrm{Id}_X$ and we say X *is a retract of* Y *with respect to* (g, f). This is often abbreviated by $Y \rhd_{(g,f)} X$. We say X *is a retract of* Y if there is a pair (g, f) as above such that $Y \rhd_{(g,f)} X$.

For simplicity, we assume all categories $\mathcal{C}$ below are locally small, so we can speak of hom-sets (rather than hom-classes).

3.1. *Polarized GoI situations*

We introduce a polarized analogue of GoI situations from Abramsky *et al.* (2002) (see Appendix A for a summary), which is suitable for polarized multiplicative linear logic. The full exponential operators T of GoI situations for (non-polarized) linear logic LL are here replaced by much weaker, functorial *polarity shifters:* $\uparrow$ and $\downarrow$, as well as by introducing the critical notion of multi-points.

Definition 3.1 (Polarized GoI situation). A *polarized GoI situation* is a tuple

$$(\mathcal{C}, \otimes, I, s, U, 1, (\mathfrak{e}, \mathfrak{m}), 1 \xrightarrow{\alpha} U, (\mathfrak{e}_\alpha, \mathfrak{m}_\alpha), 0),$$

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

where

1. $(\mathcal{C}, \otimes, I, s)$ is a traced symmetric monoidal category (with symmetry maps $s = \{s_{A,B} : A \otimes B \to B \otimes A\}$), and with unit object I, satisfying the usual identities (Mac Lane 1971). (A *trace* is given by a family of functions $\mathsf{Tr}^Z_{X,Y} : \mathcal{C}(X \otimes Z, Y \otimes Z) \longrightarrow \mathcal{C}(X, Y)$ subject to three naturalities: (Natural in X), (Natural in Y), (Dinatural in Z) and three axioms (Vanishing) (Superposing) and (Yanking). We refer to Joyal *et al.* (1996), Abramsky *et al.* (2002) and Haghverdi and Scott (2006) for detailed treatment.) In the monoidal structure, X^m and f^m denote, respectively, m-ary tensors (also called *tensor foldings*) $\overbrace{X \otimes \cdots \otimes X}^{m}$ and $\overbrace{f \otimes \cdots \otimes f}^{m}$.
2. U is an object of $\mathcal{C}$ with a retraction $U \rhd_{(k,j)} U \otimes U$, i.e.

$$U \underset{j}{\overset{k}{\rightleftarrows}} U \otimes U \quad \text{such that } k \circ j = \mathrm{Id}_{U \otimes U}.$$

3. (*The Object* 1) 1 is an object of $\mathcal{C}$ with a retraction $1 \otimes 1 \rhd_{(\mathfrak{e},\mathfrak{m})} 1$, i.e.

$$1 \otimes 1 \underset{\mathfrak{m}}{\overset{\mathfrak{e}}{\rightleftarrows}} 1 \quad \text{such that } \mathfrak{e} \circ \mathfrak{m} = \mathrm{Id}_1.$$


   Note: In general, $1 \neq I$.
4. (*Distinguished points*) A morphism $1 \longrightarrow U$ is called a *point* of U. A polarized GoI situation always has a distinguished point $1 \xrightarrow{\alpha} U$ among the points of U. Later (in Definition 3.5), we introduce inductively a subclass $\mathbb{P}$ of distinguished points generated from α. From now on, the word 'point' refers to any distinguished point.
5. (*Uniformity of trace*)
   The trace of $\mathcal{C}$ is *uniform* (cf. Hasegawa 2004 and Simpson and Plotkin 2000) over points; i.e. every distinguished point $p : 1 \longrightarrow U \in \mathbb{P}$ satisfies the following condition, which says points are trace invariant.
   For any morphisms f and g,

$$\begin{array}{ccc} X \otimes U & \xrightarrow{\quad f \quad} & Y \otimes U \\ {\scriptstyle X \otimes p}\uparrow & & \uparrow{\scriptstyle Y \otimes p} \\ X \otimes 1 & \xrightarrow[\quad g \quad]{} & Y \otimes 1 \end{array} \qquad \text{implies} \qquad \mathsf{Tr}^U_{X,Y}(f) = \mathsf{Tr}^1_{X,Y}(g).$$


   The final axioms 6, 7 and 9 in the definition concern properties satisfied by the distinguished point $1 \xrightarrow{\alpha} U$.
6. (*Lifting Property* $U \otimes 1 \rhd_{(\mathfrak{e}_\alpha,\mathfrak{m}_\alpha)} U$ *along* α)
   For the distinguished point $1 \xrightarrow{\alpha} U$, there exists a pair $(\mathfrak{e}_\alpha, \mathfrak{m}_\alpha)$ giving a retraction $U \otimes 1 \rhd_{(\mathfrak{e}_\alpha,\mathfrak{m}_\alpha)} U$ which lifts the retraction $1 \otimes 1 \rhd_{(\mathfrak{e},\mathfrak{m})} 1$ along the point α. This means the

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

following diagram commutes (in all possible ways) with $\mathfrak{e}_\alpha \circ \mathfrak{m}_\alpha = \mathrm{Id}_U$ and $\mathfrak{e} \circ \mathfrak{m} = \mathrm{Id}_1$:




7. (*Semi-invertibility of* α)
   The point $\alpha : 1 \longrightarrow U$ is *semi-invertible*. That is, there exists $\alpha^* : U \longrightarrow 1$ such that $\alpha^* \circ \alpha = \mathrm{Id}_1$.
8. (0 *morphisms*)
   The category $\mathcal{C}$ has *zero morphisms*. This means for every pair of objects $X, Y \in \mathcal{C}$, there is an assigned map $0_{XY} : X \to Y$ such that the family $\{0_{XY} \mid X, Y \in \mathcal{C}\}$ satisfies: for every $f : W \to X, g : Y \to Z$,




   Note: If f or g equals the identity, this amounts to the fact that any composition with one factor zero is itself zero (see also Manes and Arbib 1986). For simplicity, 0 denotes the zero morphism $0_{11} : 1 \to 1$. Note that $f \otimes 0_{Y,Z}$ is not in general $0_{W \otimes Y, X \otimes Z}$ for any objects X, Y, W, Z[‡].
9. ($(\mathfrak{e}_\alpha, \mathfrak{m}_\alpha)$ *and* 0)
   For every morphism $f : V \otimes X \longrightarrow W \otimes X$ with $X \in \{U, 1\}$ and $0 : 1 \longrightarrow 1$,

$$(\mathrm{Id}_W \otimes \mathfrak{e}_\alpha) \circ (f \otimes 0) \circ (\mathrm{Id}_V \otimes \mathfrak{m}_\alpha) = f \quad \text{and} \quad (\mathrm{Id}_W \otimes \mathfrak{e}) \circ (f \otimes 0) \circ (\mathrm{Id}_V \otimes \mathfrak{m}) = f$$

   Diagrammatically, the following are the respective equations when $X = U$ and $X = 1$:



$= f$ and



$= f$

This ends the definition of a polarized GoI situation.

The following Examples 3.2 and 3.3 of polarized GoI situations are built from the category Rel of sets and relations. Rel has two standard traced-monoidal structures (see Abramsky

‡ Hence, zero morphisms are absorbing with respect to composition, but not with respect to tensor.

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

et al. 2002 and Joyal *et al.* 1996 for details), one with $\otimes = \times$ (cartesian product), the other $\otimes = +$ (disjoint union). These two categories are denoted by $\mathsf{Rel}_\times$ and Rel_+, respectively. We do not discuss $\mathsf{Rel}_\times$ in this paper, so in what follows we often abbreviate Rel_+ to Rel. Below, we emphasize the additional polarized structure.

Example 3.2 (Polarized GoI situation Rel_+).

$$(\mathsf{Rel}, \otimes, I, s, \mathbb{N}, 1, (\mathfrak{e}, \mathfrak{m}), 1 \xrightarrow{\alpha} U, (\mathfrak{e}_\alpha, \mathfrak{m}_\alpha), 0)$$

is a polarized GoI situation, denoted Rel_+ (or just Rel), where we define

- the objects of Rel_+ are sets and morphisms are (binary) relations between them;
- $\otimes$ is the disjoint union $+$, where $A + B := (\{1\} \times A) \cup (\{2\} \times B)$;
- $I := \varnothing$, $U := \mathbb{N}$, $1 := \{*\}$;
- the retraction $k : \mathbb{N} \triangleright \mathbb{N} + \mathbb{N} : j$ is a standard one often used in GoI (Abramsky *et al.* 2002; Girard 1989; Haghverdi 2000):
$$j(1,n) = 2n\ ,\ j(2,n) = 2n+1, \text{ and } k(n) = \begin{cases} (1, \frac{n}{2}) & n \text{ even} \\ (2, \frac{n-1}{2}) & n \text{ odd} \end{cases};$$
- $\mathfrak{m} : 1 \to 1 + 1$ is the maximal relation, and $\mathfrak{e} : 1 + 1 \to 1$ is its converse;
- $\alpha : \{*\} \longrightarrow \mathbb{N}$ is a *non-empty relation* that determines a distinguished *singleton* subset $\{n_\alpha\}$ of $\mathbb{N}$;
- $\mathfrak{e}_\alpha : U + 1 \longrightarrow U$ is the relation whose restriction on U (resp. on 1) is Id_U (resp. α) and $\mathfrak{m}_\alpha$ is the converse relation of $\mathfrak{e}_\alpha$;
- the zero morphism 0_{XY} is the empty relation, for all X, Y.

Example 3.3 (Pfn and PInj as degenerate polarized GoI situations). Starting with the category Rel above, we consider the two major subcategories Pfn (partial functions) and PInj (partial injective functions) from Abramsky *et al.* (2002), with the following choices of $\mathfrak{e}, \mathfrak{m}, \mathfrak{e}_\alpha, \mathfrak{m}_\alpha$: for both Pfn and PInj , $\mathfrak{m} : 1 \to 1 + 1$ is the left (or right) inclusion and $\mathfrak{e}$ is its inverse. $\mathfrak{m}_\alpha : U \to U + 1$ is the left embedding.

- Pfn: $\mathfrak{e}_\alpha : U + 1 \longrightarrow U$ is the total function whose restriction on U (resp. on 1) is Id_U (resp. α).
- PInj: $\mathfrak{e}_\alpha : U + 1 \longrightarrow U$ is the partial injection determined by the identity on U.

We say these models are *degenerate* since in Rel, $\mathfrak{m}[1] := \{y \mid (*, y) \in \mathfrak{m}\} = 1 + 1$, whereas in the other models Pfn and PInj, $\mathfrak{m}[1] = 1$. This latter property of 1 causes the interpretation of polarized proofs in MLLP to become degenerate (see Remark 3.20 below).

We now make some remarks and observations about the axioms for polarized GoI situations in Definition 3.1 above. We generalize Axioms **2** and **3** to m-ary tensors, and denote them by Axioms **2′** and **3′**, respectively. We will apply these remarks in Proposition 3.8 below, where Axioms **6** and **9** are generalized, resulting in Axioms **6′** and **9′**, respectively.

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

• *Strengthening of Axiom 5 (Uniformity of Trace)*:

Proposition 3.4 (Strong uniformity of trace:). *For any morphisms*

$$X_2 \otimes U \xrightarrow{f} Y_2 \otimes U\,,\ X_1 \otimes 1 \xrightarrow{g} Y_1 \otimes 1\,,\ X_1 \xrightarrow{a} X_2\,,\ Y_1 \xrightarrow{b} Y_2,$$

and point $p : 1 \to U$, *we have*

$$\begin{array}{ccc} X_2 \otimes U & \xrightarrow{f} & Y_2 \otimes U \\ {\scriptstyle a\otimes p}\uparrow & & \uparrow{\scriptstyle b\otimes p} \\ X_1 \otimes 1 & \xrightarrow[g]{} & Y_1 \otimes 1 \end{array} \quad \textit{implies} \quad \begin{array}{ccc} X_2 & \xrightarrow{\mathsf{Tr}^U_{X_2,Y_2}(f)} & Y_2 \\ {\scriptstyle a}\uparrow & & \uparrow{\scriptstyle b} \\ X_1 & \xrightarrow[\mathsf{Tr}^1_{X_1,Y_1}(g)]{} & Y_1 \end{array}$$

Proof. Define

$$f' = f\circ(a \otimes U) : X_1 \otimes U \longrightarrow Y_2 \otimes U \ \text{ and } \ g' = (b \otimes 1)\circ g : X_1 \otimes 1 \longrightarrow Y_2 \otimes 1.$$

Apply Axiom 5 to f' and g' with $X = X_1$ and $Y = Y_2$, then

$$\mathsf{Tr}^U_{X_1,Y_2}(f \circ (a \otimes U)) \ = \ \mathsf{Tr}^1_{X_1,Y_2}((b \otimes 1) \circ g).$$

By naturality, the L.H.S (resp. R.H.S) is equal to $\mathsf{Tr}^U_{X_2,Y_2}(f) \circ a$ (resp. $b \circ \mathsf{Tr}^1_{X_1,Y_1}(g)$), which proves the assertion. □

Observe, strong uniformity of trace implies the original version (Axiom 5) by setting a and b to be the appropriate identity arrows.

• *Generalizing Axiom 2 to the case of m-ary tensors.*

Axiom **2′.** *For any natural numbers* $m \geqslant 2$, *there is a retraction* $U \rhd_{(k_m,j_m)} U^m$

$$U \underset{j_m}{\overset{k_m}{\rightleftarrows}} U^m \quad \text{such that } k_m \circ j_m = \mathrm{Id}_{U^m}.$$

We define j_m and k_m as follows. This will be our fixed choice for the retraction structure for the rest of the paper.

$$j_m = j\circ(j \otimes U)\circ\cdots\circ(j \otimes U^{m-1}), \quad k_m = (k \otimes U^{m-1})\circ\cdots\circ(k \otimes U)\circ k. \tag{1}$$

When $m = 1$, under the convention that $j_m = U = k_m$, the retraction of 2′ becomes the trivial identity. Hence in what follows, (k_m, j_m) is used for any non-zero natural number m. When $m = 2$, we get the original Axiom 2.

• *An m-fold tensor version of Axiom 3 :* $1^m \otimes 1^m \rhd_{(\mathfrak{e}^m,\mathfrak{m}^m)} 1^m$.

It is straightforward to show the following:

Axiom **3′.** $\mathfrak{e}^m$ *has right inverse* $\mathfrak{m}^m$, i.e.

$$1^m \otimes 1^m \underset{\mathfrak{m}^m}{\overset{\mathfrak{e}^m}{\rightleftarrows}} 1^m \ \text{ satisfying } \mathfrak{e}^m \circ \mathfrak{m}^m = \mathrm{Id}_{1^m},$$

where $\mathfrak{e}^m$ and $\mathfrak{m}^m$ are m-ary tensors of $\mathfrak{e}$ and r, respectively.

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

Finally, the reader may wonder why, in our definition of polarized GoI situation (Definition 3.1), we introduced opposite directions for the retractions on U and on 1. This will be explained in Appendix E below.

3.2. *Multi-points in polarized GoI situations*

In this section, we introduce the key notion of *multi-points* for interpreting the weak exponentials $\uparrow$ and $\downarrow$ (polarity shifting) of MLLP in polarized GoI situations. We then generalize Axioms 6 and 9 of a polarized GoI situation to the level of multi-points. In the following Section 3.3, we shall make use of multi-points to translate polarized formulas, and then extend this to a polarized GoI interpretation of MLLP proofs. This will be used later (in Sections 3.4 and 3.5 below) to find new invariants for cut elimination, and to characterize focussing, hence positivity and negativity, in polarized logics.

Definition 3.5 (Distinguished points and multi-points).
Multi-points in a polarized GoI situation form a certain class (denoted $\mathbb{MP}$) of morphisms $1^m \longrightarrow U$ for natural numbers m defined below. The *distinguished points* are the subclass (denoted $\mathbb{P}$) of multi-points in which $m = 1$. We define both classes inductively as follows.

First, $\mathbb{P}$ is constructed by the following BNF construction:

$$\mathbb{P} := \quad \alpha \quad | \quad j \circ (\mathbb{P} \otimes 0_{I,U}) \quad | \quad j \circ (0_{I,U} \otimes \mathbb{P})$$

That is, the following takes place:

1. The distinguished point $\alpha : 1 \longrightarrow U$ is a point.
2. If $\beta : 1 \longrightarrow U$ is a point, so are
$$1 \cong 1 \otimes I \xrightarrow{\beta \otimes 0_{I,U}} U \otimes U \xrightarrow{j} U \quad \text{and} \quad 1 \cong I \otimes 1 \xrightarrow{0_{I,U} \otimes \beta} U \otimes U \xrightarrow{j} U\,.$$

Second, $\mathbb{MP}$ is constructed from $\mathbb{P}$ and j_m ranging over natural numbers $m \geqslant 2$, as follows:

$$\mathbb{MP} := \quad \mathbb{P} \quad | \quad j_m \circ \tau \circ (\overbrace{\mathbb{P} \otimes \cdots \otimes \mathbb{P}}^{m}), \qquad \text{where } \tau \text{ ranges over the permutations of } U^m.$$

That is, while all (distinguished) points are multi-points, the second construction stipulates in addition of the following:

3. If $p_i : 1 \longrightarrow U$ are points for $i = 1, \ldots, m$, and τ is a permutation of U^m, then the following is a multi-point:
$$j \circ \tau \circ \bigotimes_{i=1}^{m} p_i : \quad 1^m \xrightarrow{\bigotimes_{i=1}^{m} p_i} U^m \xrightarrow{\tau} U^m \xrightarrow{j_m} U\,. \tag{2}$$

Remark 3.6 (Various contractions $j_m \circ \tau$ arising from permutations τ.). We make no assumptions on commutativity nor on associativity axioms for the monoidal j and the comonoidal k. Instead, we adopt a minimal categorical setting for a GoI situation, as suggested by the referee. There are various ways of contracting U^m to U, depending on the choice of U (to apply j to) in each of the $m - 1$ steps of contraction. Our specific choice of j_m in Equations (1) determines *one* such choice. Precomposing each permutation

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press


τ of U^m with j_m determines a different choice of contraction. Correspondingly, the left inverse of $j_m \circ \tau$ is given by the comonoidal $\tau^{-1} \circ k_m$, and the pair gives a different retraction $U \rhd_{(j_m \circ \tau,\, \tau^{-1} \circ k_m)} U^m$. Later in Section 3.3, when we interpret every polarized formula as a multi-point, the permutation τ will be explicitly specified by the syntactic tree of the formula.

Example 3.7. In the Rel model of Example 3.2 where the distinguished α is taken to be a singleton subset $\{n_\alpha\}$ of $U = \mathbb{N}$, a multi-point $1^m \longrightarrow U$ (generated by this α) determines a finite indexed family of cardinality m of U.

Later we shall see how polarized formulas can be interpreted in a multi-pointed setting (Definition 3.13 in Section 3.3 below). Together with a two-layered GoI interpretation of MLLP proofs (Definition 3.18), this will turn out to be essential in our characterization of focusing (see Theorem 3.26).

We now prove generalizations of Axioms **6** and **9** of a polarized GoI situation to the level of multi-points, using the m-fold setting of **3**′. We call these **Axioms 6**′ and **9**′, respectively.

Proposition 3.8 (Axiom 6′: Lifting $U \otimes 1^m \rhd_{(\mathfrak{e}_{\boldsymbol{p}}, \mathfrak{m}_{\boldsymbol{p}})} U$ along a multi-point $\boldsymbol{p}$).
For any multi-point $\boldsymbol{p} : 1^m \longrightarrow U$, there exists a pair $(\mathfrak{e}_{\boldsymbol{p}}, \mathfrak{m}_{\boldsymbol{p}})$ giving a retraction which lifts the retraction $1^m \otimes 1^m \rhd_{(\mathfrak{e}^m, \mathfrak{m}^m)} 1^m$ along $\boldsymbol{p}$. This means the following diagram commutes (in all possible ways), with $\mathfrak{e}_{\boldsymbol{p}} \circ \mathfrak{m}_{\boldsymbol{p}} = \mathrm{Id}_U$:



Proof. See Appendix D. □

The next Proposition discusses the lifted retraction pair $(\mathfrak{e}_{\boldsymbol{p}}, \mathfrak{m}_{\boldsymbol{p}})$ of the above Axiom **6**′.

Proposition 3.9 (Axiom 9′: On the lifted retraction pair $(\mathfrak{e}_{\boldsymbol{p}}, \mathfrak{m}_{\boldsymbol{p}})$).
For any multi-point $\boldsymbol{p} : 1^m \longrightarrow U$, any morphism $f : V \otimes X \longrightarrow W \otimes X$ with $X \in \{U, 1^m\}$ and $0 : 1^m \longrightarrow 1^m$,

$$(\mathrm{Id} \otimes \mathfrak{e}_{\boldsymbol{p}}) \circ (f \otimes 0) \circ (\mathrm{Id} \otimes \mathfrak{m}_{\boldsymbol{p}}) = f \quad \textit{and} \quad (\mathrm{Id} \otimes \mathfrak{e}^m) \circ (f \otimes 0) \circ (\mathrm{Id} \otimes \mathfrak{m}^m) = f$$

This is illustrated in Figure 1 for the respective equations when $X = U$ and $X = 1^m$.

Proof. Straightforward induction on the given construction of $\boldsymbol{p}$. □

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press




Fig. 1. Axiom **9′**.

The above definition of multi-points is compatible with Axiom **5** (uniformity of trace) in Definition 3.1, in the sense that uniformity of trace generalizes to multi-points (see Axiom **5′** in Proposition 3.11 below).

Lemma 3.10 (Invariance of traces under conjugate actions of the retractions). *For any non-zero natural number m and a multi-point $\boldsymbol{p} : 1^m \longrightarrow U$, the retractions (k_m, j_m) and $(\mathfrak{e}_{\boldsymbol{p}}, \mathfrak{m}_{\boldsymbol{p}})$, respectively act by conjugation on morphisms $f : X \otimes U^m \longrightarrow Y \otimes U^m$ and $g : X \otimes U \longrightarrow Y \otimes U$ as follows:*

$$\begin{aligned} f^{(k_m, j_m)} &:= (X \otimes j_m) \circ f \circ (X \otimes k_m) \quad : X \otimes U \longrightarrow Y \otimes U \\ g^{(\mathfrak{e}_{\boldsymbol{p}}, \mathfrak{m}_{\boldsymbol{p}})} &:= (X \otimes \mathfrak{m}_{\boldsymbol{p}}) \circ g \circ (X \otimes \mathfrak{e}_{\boldsymbol{p}}) \quad : X \otimes U \otimes 1^m \longrightarrow Y \otimes U \otimes 1^m \end{aligned}$$

Then the following invariant equations hold:

$$\mathsf{Tr}^{U^m}_{X,Y}(f) = \mathsf{Tr}^{U}_{X,Y}\left(f^{(k_m, j_m)}\right), \tag{3}$$


$$\mathsf{Tr}^{U}_{X,Y}(g) = \mathsf{Tr}^{U \otimes 1^m}_{X,Y}\left(g^{(\mathfrak{e}_{\boldsymbol{p}}, \mathfrak{m}_{\boldsymbol{p}})}\right). \tag{4}$$

The two equations guarantee the invariance of taking traces along U (instead of U^m) and along $U \otimes 1^m$ (instead of $U \otimes 1$).

Proof. See Appendix D. □

The following Axiom **5′** generalizes Axiom **5**:

Proposition 3.11 (Axiom 5′ (uniformity of trace on multi-points)). *Every multi-point $\boldsymbol{p} : 1^m \longrightarrow U$ satisfies the following condition: for any morphisms f and g,*



commuting implies

$$\mathsf{Tr}^{1^m}_{X,Y}(g) = \mathsf{Tr}^{U}_{X,Y}\left(f^{(k_m, j_m)}\right) = \mathsf{Tr}^{U^m}_{X,Y}(f).$$

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

Note:


i. The composition of the top horizontal arrows above is $f^{(k_m, j_m)}$ of Lemma 3.10.
ii. Axiom **5** is a special case, by setting $m = 1$ and using our convention that $j_1 = U = k_1$.

Proof. We first note that the second equation of the assertion is by Equation (3) of Lemma 3.10.

We prove the assertion by induction on $m \geqslant 1$; i.e. by induction on the construction of a multi-point $\boldsymbol{p} : 1^m \longrightarrow U$.

(Base Case): $m = 1$. The assertion is the original Axiom 5 of Definition 3.1.


(Induction Case for $m + 1$)
The given commutative diagram with $m + 1$ factors as follows from the construction of $\boldsymbol{p}$ by Definition 3.5 for some permutation τ on U^m. Typographically, we write τ^- for τ^{-1}.



where $\boldsymbol{p} = j_{m+1} \circ \tau \circ \otimes_{i=1}^{m+1} p_i$ and $\boldsymbol{p}' = j_m \circ \tau \circ \otimes_{i=1}^{m} p_i$.

Suppose the outer trapezium commutes. We note that the inner trapezium commutes by both precomposing $X \otimes U \overset{X\otimes j}{\longleftarrow} X \otimes U \otimes U$ and postcomposing $Y \otimes U \otimes U \overset{Y\otimes k}{\longleftarrow} Y \otimes U$, respectively on the top left-most and right-most horizontal arrows, because $k \circ j = \mathrm{Id}_{U\otimes U}$ and since the left and right triangles commute.

Then applying the I.H. to this inner trapezium, we have



The upper horizontal arrow is equal to

$$\mathsf{Tr}^U_{X\otimes U, Y\otimes U}\left(f^{(j_m \circ \tau,\ \tau^- \circ k_m)}\right) = \mathsf{Tr}^U_{X\otimes U, Y\otimes U}\left(f^{(j_m, k_m)}\right) = \mathsf{Tr}^{U^m}_{X\otimes U, Y\otimes U}(f)$$

The first equation is by dinaturality on τ, cancelling with its inverse τ^-, and the second equation is by Equation (3).

Then applying the original Axiom 5 (uniformity of trace) to the above square, we obtain


$$\mathsf{Tr}^U_{X,Y}\left(\mathsf{Tr}^{U^m}_{X\otimes U, Y\otimes U}(f)\right) = \mathsf{Tr}^1_{X,Y}\left(\mathsf{Tr}^m_{X\otimes 1, Y\otimes 1}(g)\right)$$

By Vanishing applied to both sides of the equation, we have $\mathsf{Tr}^{U^{m+1}}_{X,Y}(f) = \mathsf{Tr}^{1^{m+1}}_{X,Y}(g)$. □

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

As in the case for points in Proposition 3.4, a stronger version of **5′** can be derived for multi-points.

Corollary 3.12 (Strong uniformity of trace on multi-points).
Every multi-point $\boldsymbol{p} : 1^m \longrightarrow U$ *satisfies the following condition: for any morphisms* f, g, a, b,



Proof. Same as Proposition 3.11, using naturality. □

3.3. *The GoI interpretation of* MLLP *proofs*

We now define one of the central notions of this paper: the GoI interpretation of MLLP proofs in polarized GoI situations. We shall begin with a detailed discussion of how to interpret multi-points of polarized formulas. We then present a categorical approach to GoI in the polarized case, influenced by the categorical approach to ordinary GoI of Haghverdi and Scott (2006, 2011) in (ordinary) GoI situations, as summarized in Appendix A below.

Definition 3.13 (Multi-points associated with formulas). Given a polarized GoI situation and a polarized MLLP formula A, we will inductively construct below a morphism $\mathsf{mp}(A)$ together with its domain $\mathbb{1}_A$ and codomain U_A, where $U_A := U$.

$$\mathsf{mp}(A) : \mathbb{1}_A \longrightarrow U_A.$$

More generally, for a sequence $\mathcal{M} = A_1, \ldots, A_n$ of polarized formulas, we will construct a morphism

$$\mathsf{mp}(\mathcal{M}) : \mathbb{1}_{\mathcal{M}} \longrightarrow U_{\mathcal{M}} := U_{A_1} \otimes \cdots \otimes U_{A_n}$$

from the constructed domain $\mathbb{1}_{\mathcal{M}}$ to the codomain $U_{\mathcal{M}}$. The arrows $\mathsf{mp}(A)$ (resp. $\mathsf{mp}(\mathcal{M})$) defined below are called *the multi-point associated with formula A (resp. with sequence* $\mathcal{M}$*)*. All these multi-points are elements of the distinguished class $\mathbb{MP}$ of Definition 3.5.

Construction of multi-points.

First, to each positive (resp. negative) formula P (resp. N) of MLLP, we associate an object $\mathbb{1}_P$ (resp. $\mathbb{1}_N$), which is a tensor product of 1s, defined inductively as follows:

$$\begin{array}{llll} \mathbb{1}_X := 1 & \mathbb{1}_{P \otimes Q} := \mathbb{1}_P \otimes \mathbb{1}_Q & \mathbb{1}_{\downarrow N} := 1 \\ \mathbb{1}_{X^\perp} := 1 & \mathbb{1}_{N \parr M} := \mathbb{1}_N \otimes \mathbb{1}_M & \mathbb{1}_{\uparrow P} := 1 \end{array}$$

For a sequence $\mathcal{M} = A_1, \ldots, A_n$ of polarized formulas, $\mathbb{1}_{\mathcal{M}} := \mathbb{1}_{A_1} \otimes \cdots \otimes \mathbb{1}_{A_n}$. Then $\mathbb{1}_{\mathcal{M}}$ is 1^m for a certain natural number m, which we denote $\kappa(\mathcal{M})$.

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

Second, with each positive (resp. negative) formula P (resp. N), we associate three objects $\boldsymbol{U}_P$, $P^\flat$ and $P^\sharp$ (resp. $\boldsymbol{U}_N$, $N^\flat$ and $N^\sharp$) so that

$$\boldsymbol{U}_A \cong A^\sharp \otimes A^\flat \tag{5}$$

inductively as follows:

$$\begin{array}{lll} \boldsymbol{U}_X := U & \boldsymbol{U}_{P\otimes Q} := \boldsymbol{U}_P \otimes \boldsymbol{U}_Q & \boldsymbol{U}_{\downarrow N} := U \otimes \boldsymbol{U}_N \\ X^\flat := I & (P \otimes Q)^\flat := P^\flat \otimes Q^\flat & (\downarrow N)^\flat := \boldsymbol{U}_N \\ X^\sharp := U & (P \otimes Q)^\sharp := P^\sharp \otimes Q^\sharp & (\downarrow N)^\sharp := U \end{array}$$

$$\begin{array}{lll} \boldsymbol{U}_{X^\perp} := U & \boldsymbol{U}_{N \parr M} := \boldsymbol{U}_N \otimes \boldsymbol{U}_M & \boldsymbol{U}_{\uparrow P} := \boldsymbol{U}_P \otimes U \\ (X^\perp)^\flat := I & (N \parr M)^\flat := N^\flat \otimes M^\flat & (\uparrow P)^\flat := \boldsymbol{U}_P \\ (X^\perp)^\sharp := U & (N \parr M)^\sharp := N^\sharp \otimes M^\sharp & (\uparrow P)^\sharp := U \end{array}$$

All these objects are isomorphic to tensor products of Us. For a sequence $\mathcal{M} = A_1, \ldots, A_n$ of polarized formulas, we define the object $\boldsymbol{U}_\mathcal{M} := \boldsymbol{U}_{A_1} \otimes \cdots \otimes \boldsymbol{U}_{A_n}$ and for $\star \in \{\flat, \sharp\}$, we define the object $\mathcal{M}^\star := A_1^\star \otimes \cdots \otimes A_n^\star$.

Note that the following hold for any polarized formula A:

– If A is a literal, $U_A = \boldsymbol{U}_A$. Otherwise,

$$U_A \quad \rhd_{(\tau \circ k_{n-1},\ j_{n-1} \circ \tau)} \quad \boldsymbol{U}_A, \tag{6}$$

where n is the number logical connectives in A so that $\boldsymbol{U}_A = U^n$, as in Axiom **2′**, and τ is a permutation of U^n.

– There exists a natural number m such that

$$A^\sharp \cong U^m \text{ and } \mathbb{1}_A \cong 1^m. \tag{7}$$

Finally, using Equations (5), (6) and (7), we define $\mathsf{mp}(A)$ by the following composition (we give a simpler, equivalent definition in Remark 3.14 below):

$$\mathsf{mp}(A) :\ \mathbb{1}_A \cong \mathbb{1}_A \otimes I \xrightarrow{\alpha^m \otimes 0_{I,A^\flat}} A^\sharp \otimes A^\flat \xrightarrow{\tau} U^n \xrightarrow{j_{n-1}} U_A, \tag{8}$$

where the permutation τ on the tensor folding $\boldsymbol{U}_A$ is determined by the syntax tree of the polarized formula A. Finally, in the case of a sequence $\mathcal{M} = A_1, \ldots, A_n$,

$$\mathsf{mp}(\mathcal{M}) := \mathsf{mp}(A_1) \otimes \cdots \otimes \mathsf{mp}(A_n) : \mathbb{1}_\mathcal{M} \longrightarrow U_\mathcal{M}. \tag{9}$$

Remark 3.14. Multi-points $\mathsf{mp}(A)$ can be *inductively* defined as follows, where the first, second and third constructions correspond, respectively, to 1, 2 and 3 of Definition 3.5:

– $\mathsf{mp}(X) := \alpha$ and $\mathsf{mp}(X^\perp) := \alpha$.
– $\mathsf{mp}(\downarrow N) := j \circ (\alpha \otimes 0_{I,U_N})$ and $\mathsf{mp}(\uparrow P) := j \circ (0_{I,U_P} \otimes \alpha)$.
– $\mathsf{mp}(P \otimes Q) := j \circ (\mathsf{mp}(P) \otimes \mathsf{mp}(Q))$ and $\mathsf{mp}(N \parr M) := j \circ (\mathsf{mp}(N) \otimes \mathsf{mp}(M))$.

We observe that $\mathsf{mp}(A)$ so inductively defined uniquely factors as Equation (8), in which the permutation τ is determined by the syntax tree of the MLLP formula A.

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

In what follows, we make the convention that $U_{\downarrow}$ (resp. $U_{\uparrow}$) denotes the codomain U of α in the above construction of $\mathsf{mp}(\downarrow N)$ (resp. of $\mathsf{mp}(\uparrow P)$). That is, $U_{\downarrow}$ (resp. $U_{\uparrow}$) denotes U of $(\downarrow N)^{\sharp}$ (resp. of $(\uparrow P)^{\sharp}$).

Example 3.15. In the Rel polarized GoI situation,

- $\mathbb{1}_{\uparrow X} = 1 = \{n_\alpha\}$ and $\mathsf{mp}(\uparrow X) : 1 \cong I + 1 \xrightarrow{0 \otimes \alpha} U_X + U_{\uparrow} \xrightarrow{j} U_{\uparrow X}$ is a singleton subset consisting of the element $j((2, n_\alpha)) = 2n_\alpha + 1 \in \mathbb{N}$.
- $\mathbb{1}_{Y^{\perp} \parr X^{\perp}} = 1^2 = \{n_\alpha\} + \{n_\alpha\}$ and $\mathsf{mp}(Y^{\perp} \parr X^{\perp}) : 1^2 \xrightarrow{\alpha^2} U_{Y^{\perp}} + U_{X^{\perp}} \xrightarrow{j} U_{Y^{\perp} \parr X^{\perp}}$ is a subset of cardinality 2 consisting of the elements $j((1, n_\alpha)) = 2n_\alpha$ and $j((2, n_\alpha)) = 2n_\alpha + 1$ in $\mathbb{N}$.

Definition 3.16 (retractions $(\mathfrak{e}_A, \mathfrak{m}_A)$). For a polarized formula A, we define two morphisms $\mathfrak{m}_A$ and $\mathfrak{e}_A$ to give a retraction

$$A^{\sharp} \otimes \mathbb{1}_A \rhd_{(\mathfrak{e}_A, \mathfrak{m}_A)} A^{\sharp} \text{ so that } \mathfrak{e}_A \circ \mathfrak{m}_A = A^{\sharp}.$$

These are defined inductively on A as follows:


$\mathfrak{m}_X$ and $\mathfrak{m}_{X^{\perp}}$ are $\mathfrak{m}_\alpha$, $\mathfrak{m}_{P \otimes Q} := \mathfrak{m}_P \otimes \mathfrak{m}_Q$, $\mathfrak{m}_{N \parr M} := \mathfrak{m}_N \otimes \mathfrak{m}_M$, $\mathfrak{m}_{\downarrow N}$ and $\mathfrak{m}_{\uparrow P}$ are $\mathfrak{m}_\alpha$,

$\mathfrak{e}_X$ and $\mathfrak{e}_{X^{\perp}}$ are $\mathfrak{e}_\alpha$, $\mathfrak{e}_{P \otimes Q} := \mathfrak{e}_P \otimes \mathfrak{e}_Q$, $\mathfrak{e}_{N \parr M} := \mathfrak{e}_N \otimes \mathfrak{e}_M$, $\mathfrak{e}_{\downarrow N}$ and $\mathfrak{m}_{\uparrow N}$ are $\mathfrak{e}_\alpha$.

For a sequence $\mathcal{M} = A_1, \ldots, A_n$ of polarized formulas, define $\mathfrak{m}_{\mathcal{M}}$ (resp. $\mathfrak{e}_{\mathcal{M}}$) to be $\mathfrak{m}_{A_1} \otimes \cdots \otimes \mathfrak{m}_{A_n}$ (resp. $\mathfrak{e}_{A_1} \otimes \cdots \otimes \mathfrak{e}_{A_n}$).

Then the lifting property (Axiom 6′ of Proposition 3.8) has the following variant:

Proposition 3.17 (Lifting property $A^{\sharp} \otimes \mathbb{1}_A \rhd_{(\mathfrak{e}_A, \mathfrak{m}_A)} A^{\sharp}$ along α^m).
Suppose m is a natural number satisfying Equation (7). Then, $(\mathfrak{e}_A, \mathfrak{m}_A)$ gives a retraction which lifts the retraction $\mathbb{1} \otimes \mathbb{1} \rhd_{(\mathfrak{e}^m, \mathfrak{m}^m)} \mathbb{1}$ along α^m. This means the following diagram commutes with $\mathfrak{e}_A \circ \mathfrak{m}_A = \mathrm{Id}_{A^{\sharp}}$:



Proof. Straightforward by noting that composing the right (resp. left) vertical arrow with the second and third maps of Equation (8) (resp. $\otimes 1^m$ of these maps) gives rise to Axiom 6′. □

In what follows, we introduce the GoI interpretation for MLLP. The original GoI situations in Abramsky *et al.* (2002) (summarized in Appendix A below) form a very basic framework for interpreting GoI. For example, their exponential structure, which

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

is sufficient for defining linear combinatory algebras on $\mathrm{End}_{\mathcal{C}}(U)$, for a reflexive object U, does not include the more elaborate categorical structure of the exponentials (e.g. cocommutative coalgebras, comonoids, etc.) in genuine models of linear logic (Melliès 2009).

Recall that the GoI interpretation of an MLL proof π of the sequent $\vdash [\Delta], \Gamma$ in an (ordinary) GoI situation $\mathcal{C}$ yields an endomorphism $[\![\pi]\!] \in \mathrm{End}_{\mathcal{C}}(U^{2m+n})$, for the reflexive object U. We now introduce the analogue for the polarized case of MLLP.

Let us sketch the framework, before going into details. Consider a *polarized* GoI situation $\mathcal{C}$, with a reflexive object U and an object 1. The *polarized GoI Interpretation* in Definition 3.18 below will yield *a pair* of endomorphisms, $([\![\pi]\!], f_\pi) \in \mathrm{End}_{\mathcal{C}}(U^{2m+n}) \times \mathrm{End}_{\mathcal{C}}(1^{2m'+n'})$.

We think of the two endomorphisms as layers: an 'upper' and a 'lower' GoI interpretation.

- The *upper* interpretation, $[\![\pi]\!] \in \mathrm{End}_{\mathcal{C}}(U^{2m+n})$, is on the level of the reflexive object U. It is analogous to the non-polarized GoI interpretation, using the polarized retraction structure $U \triangleright U \otimes U$ for coding 'untyped' GoI, by folding tensors of U's into a single U. Here, U^{2m+n} comes from $U_{\Delta,\Gamma}$ of Definition 3.13; n (resp. $2m$) is the number of formulas in Γ (resp. Δ). At this level, the polarity will be handled by the retraction $U \otimes 1 \triangleright U$. Hence, both the retraction structures 3 and 6 of Definition 3.1 are used.
- The *lower* interpretation, $f_\pi \in \mathrm{End}_{\mathcal{C}}(1^{2m'+n'})$, is a similar GoI formula to $[\![\pi]\!]$, but without assuming any reflexivity on 1. It is defined on the level of multi-points. Here, $1^{2m'+n'}$ comes from $\mathbb{1}_{\Delta,\Gamma}$ of Definition 3.13; n' (resp. $2m'$) is the sum, over formulas in Γ (resp. Δ), of the number of MLLP logical connectives (including literals) not bounded by polarity shifting operations. At this level, the polarity will be handled by the retraction $1 \otimes 1 \triangleright 1$. Hence, only retraction 3 of Definition 3.1 is used.

Definition 3.18 (The two-layered polarized GoI interpretation of MLLP proofs).
An MLLP proof π of $\vdash [\Delta], \Gamma$ is interpreted by two endomorphisms:

$$[\![\pi]\!] \in \mathcal{C}(U_{\Delta,\Gamma}, U_{\Delta,\Gamma}) \text{ and } f_\pi \in \mathcal{C}(\mathbb{1}_{\Delta,\Gamma}, \mathbb{1}_{\Delta,\Gamma}).$$

We see the polarized view as a *two-layered interpretation:* an *upper* layer $[\![\pi]\!]$ at the level of reflexive objects U, a *lower* layer f_π at the level of multi-points 1.

We define simultaneously[§] $[\![\pi]\!]$ and f_π by induction on π.



1. (*Axiom*): π is $\vdash P^\perp, P$.
Remember that $U_P = U_{P^\perp} = U$ and $\mathbb{1}_P \cong \mathbb{1}_{P^\perp} :\cong 1^n$ for a certain natural number n.

$$[\![\pi]\!] := s_{U,U} : U_P \otimes U_{P^\perp} \longrightarrow U_P \otimes U_{P^\perp}$$

[§] **Notation:** In Appendix A for traditional GoI (e.g. see Figure 6), we illustrate proofs of sequents $\vdash [\Delta], \Gamma$ as I/O boxes with labelled wires as interface. In what follows below, for typographical reasons, we often omit the wires and just write I/O labels for the interface (e.g. see the cut rule below).
On the I/O box, interface formulas are ordered (from top to bottom) as follows: first, the unique focused (positive) formula (if it exists), then the negative formulas and finally the sequence Δ of cut formulas.

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

$$f_\pi := s_{1^n,1^n} : \ \mathbb{1}_P \otimes \mathbb{1}_{P^\perp} \longrightarrow \mathbb{1}_P \otimes \mathbb{1}_{P^\perp}$$

where $s_{U,U} =$  and $s_{1^n,1^n} =$ 

Each arrow consisting of the crossing $s_{1^n,1^n}$ denotes the permutation (between factors of the tensor foldings $\mathbb{1}_P$ and $\mathbb{1}_{P^\perp}$) which is induced by De Morgan duality between polarized formulas P and $P^\perp$.


2. (*Cut*):


$$\pi \text{ is } \frac{\begin{array}{cc}\vdots\ \pi' & \vdots\ \pi'' \\ \vdash [\Delta'], P, \mathcal{N} & \vdash [\Delta''], P^\perp, \Xi\end{array}}{\vdash [\Delta', \Delta'', P, P^\perp], \mathcal{N}, \Xi}\ \text{cut}$$

We define

$$[\![\pi]\!] := \tau^-([\![\pi']\!] \otimes [\![\pi'']\!])\ \tau \qquad\qquad f_\pi := \tau^-(f_{\pi'} \otimes f_{\pi''})\ \tau$$

where τ denotes the indicated exchange for the conclusions and the cut formulas. Here, τ and its inverse τ^- are simply permutations of the interface (denoted by $U_{(\ )}$ and $\mathbb{1}_{(\ )}$).

$[\![\pi]\!] :=$ 

$f_\pi :=$ 

In the following cases, we define $[\![\pi]\!]^\dagger \in \mathcal{C}(\boldsymbol{U}_{\Delta,\Gamma}, \boldsymbol{U}_{\Delta,\Gamma})$ so that $[\![\pi]\!] = j_\ell \circ [\![\pi]\!]^\dagger \circ k_\ell$, where j_ℓ and k_ℓ are the retractions for $U_{\Delta,\Gamma} \rhd_{(k_\ell, j_\ell)} \boldsymbol{U}_{\Delta,\Gamma}$ in equation (6) of the construction of multi-points after Definition 3.13. Note that ℓ is a number of logical connectives of formulas contained in Γ, Δ.

3. (*Linear connectives*): For a ⅋ rule, the interpretation remains that of the premise proof. For a ⊗-rule, the interpretations are the same as those of the cut rule defined above. In the above diagrams, τ is a permutation of the interface. To be precise

(⅋ rule) we define

$$[\![\pi]\!]^\dagger = [\![\pi']\!]^\dagger \qquad \text{and} \qquad f_\pi = f_{\pi'},$$

(⊗ rule) we define

$$[\![\pi]\!]^\dagger = \tau^- [\![\pi']\!]^\dagger \tau \qquad \text{and} \qquad f_\pi = \tau^- f_{\pi'} \tau,$$

where τ for $[\![\pi]\!]^\dagger$ (resp. for f_π) is $\boldsymbol{U}_P \otimes \boldsymbol{U}_Q \otimes \boldsymbol{U}_{\Gamma'} \otimes \boldsymbol{U}_{\Gamma''} \otimes \boldsymbol{U}_{\Delta'} \otimes \boldsymbol{U}_{\Delta''} \longrightarrow \boldsymbol{U}_P \otimes \boldsymbol{U}_{\Gamma'} \otimes \boldsymbol{U}_{\Delta'} \otimes \boldsymbol{U}_Q \otimes \boldsymbol{U}_{\Gamma''} \otimes \boldsymbol{U}_{\Delta''}$ (resp. $\mathbb{1}_P \otimes \mathbb{1}_Q \otimes \mathbb{1}_{\Gamma'} \otimes \mathbb{1}_{\Gamma''} \otimes \mathbb{1}_{\Delta'} \otimes \mathbb{1}_{\Delta''} \longrightarrow \mathbb{1}_P \otimes \mathbb{1}_{\Gamma'} \otimes \mathbb{1}_{\Delta'} \otimes \mathbb{1}_Q \otimes \mathbb{1}_{\Gamma''} \otimes \mathbb{1}_{\Delta''}$).

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

Note that in the above Cases 1–3, $[\![\pi]\!]^\dagger$ and f_π are defined in the same way: replace $\boldsymbol{U}$'s by $\mathbb{1}$'s. However, this is no longer true in the following Cases 4 and 5 (cf. the paragraph above Definition 3.18).

4. (*Polarity Changing* $\uparrow$):

$$\pi \quad \text{is} \quad \dfrac{\begin{array}{c}\vdots\ \pi' \\ \vdash [\Delta], \mathcal{M}, P\end{array}}{\vdash [\Delta], \mathcal{M}, \uparrow P}\ \uparrow$$

We define

$$[\![\pi]\!]^\dagger = [\![\pi']\!]^\dagger \otimes 0_{U,U} \quad \text{with } 0_{U,U} : U_\uparrow \longrightarrow U_\uparrow.$$
$$f_\pi = ((0_{\mathbb{1}_P, I} \otimes \mathbb{1}_\mathcal{M} \otimes \mathbb{1}_\Delta) \circ f_{\pi'} \circ (0_{I, \mathbb{1}_P,} \otimes \mathbb{1}_\mathcal{M} \otimes \mathbb{1}_\Delta)) \otimes 0_{1,1} \quad \text{with } 0_{1,1} : 1 \longrightarrow 1.$$

Diagrammatically,

$$[\![\pi]\!]^\dagger := \begin{array}{ccc} \boldsymbol{U}_\Delta & & \boldsymbol{U}_\Delta \\ \boldsymbol{U}_\mathcal{M} & [\![\pi']\!]^\dagger & \boldsymbol{U}_\mathcal{M} \\ \boldsymbol{U}_P & & \boldsymbol{U}_P \\ \otimes & & \otimes \\ U_\uparrow & \xrightarrow{\quad 0_{U,U} \quad} & U_\uparrow \end{array} \qquad \boldsymbol{U}_{\uparrow P} \cong$$

$$f_\pi := \begin{array}{ccc} \mathbb{1}_\Delta & & \mathbb{1}_\Delta \\ \mathbb{1}_\mathcal{M} & f_{\pi'} & \mathbb{1}_\mathcal{M} \\ \mathbb{1}_P & & \mathbb{1}_P \\ I \xrightarrow{0_{I,\mathbb{1}_P}} & & \mathbb{1}_P \xrightarrow{0_{\mathbb{1}_P, I}} I \\ \otimes & & \otimes \\ 1 & \xrightarrow{\quad 0_{1,1} \quad} & 1 \end{array} \qquad \mathbb{1}_{\uparrow P} \cong$$

5. (*Polarity changing* $\downarrow$):

$$\pi \quad \text{is} \quad \dfrac{\begin{array}{c}\vdots\ \pi' \\ \vdash [\Delta], \mathcal{M}, N\end{array}}{\vdash [\Delta], \mathcal{M}, \downarrow N}\ \downarrow$$

For the interpretation of this case, $\mathfrak{m}_\mathcal{M}$ and $\mathfrak{e}_\mathcal{M}$ of Definition 3.16 are used. Let m be $\kappa(\mathcal{M})$ of Definition 3.13 so that $\mathbb{1}_\mathcal{M} = 1^m$. Depending on the value of m, two morphisms g_m and h_m are defined as follows, modulo associativity of the monoidal (resp. comonoidal) structure of j (resp. k) (cf. Axiom 2'): Let

$$g_m := \mathfrak{e} \circ (\mathfrak{e} \otimes 1) \circ \cdots \circ (\mathfrak{e} \otimes 1^{m-2}) \quad \text{and} \quad h_m := (\mathfrak{m} \otimes 1^{m-2}) \circ \cdots \circ (\mathfrak{m} \otimes 1) \circ \mathfrak{m} \tag{10}$$

to yield a retraction $1^m \rhd_{(g_m, h_m)} 1$. When m is 1, both g_m and h_m are defined to be Id_1.

For an exchange τ of the interface, the conjugate actions $(\ \ )^\tau$ and ${}^\tau(\ \ )$ are defined as follows: $x^\tau = (\tau^- \otimes 1^m) \circ x \circ \tau$ and ${}^\tau x = \tau^- \circ x \circ (\tau \otimes 1^m)$. In what follows in the definition, the compositions $\circ$'s are modulo permutation of $\otimes$[¶].


We define

$$[\![\pi]\!]^\dagger := \theta^- \circ (h_m \otimes [\![\pi']\!]^\dagger \otimes g_m) \circ \theta,$$

[¶] That is, in composing two maps between tensors of objects, the matching of the codomain of one with the domain of the other is only up to permutation of the tensor factors.

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press


where $\theta^- = \alpha \otimes {}^{\mathfrak{r}}(\boldsymbol{U}_N \otimes \boldsymbol{U}_\Delta \otimes \mathcal{M}^\flat \otimes \mathfrak{e}_\mathcal{M})$ and $\theta = \alpha^* \otimes (\boldsymbol{U}_N \otimes \boldsymbol{U}_\Delta \otimes \mathcal{M}^\flat \otimes \mathfrak{m}_\mathcal{M})^{\mathfrak{r}}$. See the following:



We define

$$f_\pi := \eta^- \circ (h_m \otimes f_{\pi'} \otimes g_m) \circ \eta$$

where $\eta^- = 1 \otimes 0_{\mathbb{1}_N, I} \otimes {}^{\rho}(\mathbb{1}_\Delta \otimes \mathfrak{e}^m)$ and $\eta = 1 \otimes 0_{I,\mathbb{1}_N} \otimes (\mathbb{1}_\Delta \otimes \mathfrak{m}^m)^\rho$ for an exchange ρ of the interface. See the following:



Example 3.19 ($[\![\pi]\!]$ and f_π of Definition 3.18). The following two examples are interpretations of proofs in the GoI situation Rel of Examples 3.2 and 3.15, in which α is identified with the singleton subset $\{n_\alpha\}$ of $\mathbb{N} = U$. In the following, matrices are UDC representations of Rel morphisms (see Appendix B), the blank elements denote 0 morphisms (i.e. empty relations) of appropriate types and g_m and h_m are from Equation (10).

(i) Let π be the unique cut-free proof for $\vdash \downarrow X^\perp, \uparrow X$ (the η-expansion of $\vdash X^\perp, X$). The side formula of the $\downarrow$-rule of π is $\uparrow X$ so that $\mathbb{1}_{\uparrow X} = 1$. It also holds that $\mathbb{1}_{\downarrow X^\perp} = 1$.

Then,

$$[\![\pi]\!]^\dagger = \begin{array}{c} \\ U_\downarrow \\ U_{X^\perp} \\ U_X \\ U_\uparrow \end{array} \begin{array}{c} \begin{array}{cccc} U_\downarrow & U_{X^\perp} & U_X & U_\uparrow \end{array} \\ \begin{pmatrix} & & & \{(n_\alpha, n_\alpha)\} \\ & & \mathrm{Id}_U & \\ & \mathrm{Id}_U & & \\ \{(n_\alpha, n_\alpha)\} & & & \end{pmatrix} \end{array} \quad \text{and} \quad f_\pi = \begin{array}{c} \\ \mathbb{1}_{\downarrow X^\perp} \\ \mathbb{1}_{\uparrow X} \end{array} \begin{array}{c} \begin{array}{cc} \mathbb{1}_{\downarrow X^\perp} & \mathbb{1}_{\uparrow X} \end{array} \\ \begin{pmatrix} & \mathrm{Id}_1 \\ \mathrm{Id}_1 & \end{pmatrix} \end{array}.$$

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

They are obtained, respectively, from

$$\begin{array}{c} \\ U_{\downarrow} \\ U_{X^\perp} \\ U_X \\ U_{\uparrow} \\ 1 \end{array}\!\!\begin{array}{c} \begin{array}{ccccc} U_{\downarrow} & U_{X^\perp} & U_X & U_{\uparrow} & 1 \end{array} \\ \begin{pmatrix} & & & & \alpha\circ g_1 \\ & & \mathrm{Id}_U & & \\ & \mathrm{Id}_U & & & \\ & & & & \\ h_1\circ\alpha^* & & & & \end{pmatrix} \end{array} \quad \text{and} \quad \begin{array}{c} \\ \mathbb{1}_{\downarrow X^\perp} \\ \mathbb{1}_{\uparrow X} \\ 1 \end{array}\!\!\begin{array}{c} \begin{array}{ccc} \mathbb{1}_{\downarrow X^\perp} & \mathbb{1}_{\uparrow X} & 1 \end{array} \\ \begin{pmatrix} & & g_1 \\ & & \\ & & \\ h_1 & & \end{pmatrix} \end{array}$$

when contracting the last two columns (resp. rows), respectively, by $\mathfrak{m}_{\uparrow X}$ (resp. $\mathfrak{e}_{\uparrow X}$) and by r (resp. $\mathfrak{e}$).

(ii) Let π be the unique cut-free proof of $\vdash \downarrow\uparrow(X \otimes Y), Y^\perp \parr X^\perp$.
The side formula of the $\downarrow$ rule of π is $Y^\perp \parr X^\perp$ so that $\mathbb{1}_{Y^\perp \parr X^\perp} = 1^2$. Both $\mathbb{1}_{\downarrow\uparrow(X\otimes Y)}$ and $\mathbb{1}_{\uparrow(X\otimes Y)}$ are 1. Note that $\boldsymbol{U}_{Y^\perp \parr X^\perp} = (Y^\perp \parr X^\perp)^\sharp = U + U$.

Then,

$$[\![\pi]\!]^\dagger = \begin{array}{c} \\ U_{\uparrow} \\ \\ \boldsymbol{U}_{Y^\perp \parr X^\perp} \end{array}\!\!\begin{array}{c} \begin{array}{cc} U_{\downarrow} & \boldsymbol{U}_{X\otimes Y} \end{array} \\ \begin{pmatrix} & \\ & \\ \{(n_\alpha, n_\alpha)\} & \\ +\{(n_\alpha, n_\alpha)\} & \mathrm{Id}_{U+U} \end{pmatrix} \end{array} \otimes \begin{array}{c} \\ U_{\downarrow} \\ \\ \boldsymbol{U}_{X\otimes Y} \end{array}\!\!\begin{array}{c} \begin{array}{cc} U_{\uparrow} & \boldsymbol{U}_{Y^\perp \parr X^\perp} \end{array} \\ \begin{pmatrix} & \{(n_\alpha, n_\alpha)\} \\ & +\{(n_\alpha, n_\alpha)\} \\ & \\ & \mathrm{Id}_{U+U} \end{pmatrix} \end{array}$$

and

$$f_\pi = \begin{array}{c} \\ \mathbb{1}_{\downarrow\uparrow(X\otimes Y)} \\ \mathbb{1}_{Y^\perp \parr X^\perp} \end{array}\!\!\begin{array}{c} \begin{array}{cc} \mathbb{1}_{\downarrow\uparrow(X\otimes Y)} & \mathbb{1}_{Y^\perp \parr X^\perp} \end{array} \\ \begin{pmatrix} & \mathrm{Id}_1 + \mathrm{Id}_1 \\ \mathrm{Id}_1 + \mathrm{Id}_1 & \end{pmatrix} \end{array}.$$

They are obtained, respectively, from

$$\begin{array}{c} \\ U_{\downarrow} \\ \boldsymbol{U}_{X\otimes Y} \\ U_{\uparrow} \\ \boldsymbol{U}_{Y^\perp \parr X^\perp} \\ 1^2 \end{array}\!\!\begin{array}{c} \begin{array}{ccccc} U_{\downarrow} & \boldsymbol{U}_{X\otimes Y} & U_{\uparrow} & \boldsymbol{U}_{Y^\perp \parr X^\perp} & 1^2 \end{array} \\ \begin{pmatrix} & & & & \alpha\circ g_2 \\ & & & \mathrm{Id}_{U+U} & \\ & & & & \\ & \mathrm{Id}_{U+U} & & & \\ h_2\circ\alpha^* & & & & \end{pmatrix} \end{array} \quad \text{and} \quad \begin{array}{c} \\ \mathbb{1}_{\downarrow\uparrow(X\otimes Y)} \\ \mathbb{1}_{Y^\perp \parr X^\perp} \\ 1^2 \end{array}\!\!\begin{array}{c} \begin{array}{ccc} \mathbb{1}_{\downarrow\uparrow(X\otimes Y)} & \mathbb{1}_{Y^\perp \parr X^\perp} & 1^2 \end{array} \\ \begin{pmatrix} & & g_2 \\ & & \\ h_2 & & \end{pmatrix} \end{array}$$

when contracting the last two columns (resp. rows), respectively, by $\mathfrak{m}_{Y^\perp \parr X^\perp}$ (resp. $\mathfrak{e}_{Y^\perp \parr X^\perp}$) and by $\mathfrak{m}^2$ (resp. $\mathfrak{e}^2$).

Remark 3.20 (On degenerate GoI situations). In Example 3.3, we noted that the two polarized GoI situations Pfn and Plnj are degenerate. We can now say why we chose this terminology. The reader can check that in both Pfn and Plnj , the interpretation above of the polarity-changing rule $\downarrow$ has no effect; that is, $[\![\pi]\!] = [\![\pi']\!]$ for the conclusion of this rule. This is definitely *not* the case in the Rel model.

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

Definition 3.21 (Polarized execution formulas). Let $\sigma = \otimes^m s$, the m-fold tensor product of the symmetry $s = s_{U,U}$, and $\sigma_* = \otimes_{i=1}^m s_i$ where each s_i is the symmetry $s_{1^{\ell_i},1^{\ell_i}}$ for certain ℓ_i. The s corresponds to the permutation between dual cut formulas so that $U = U_P = U_{P^\perp}$ and the s_i corresponds to the permutation induced by De Morgan duality for dual cut formulas so that $1^{\ell_i} \cong \mathbb{1}_P \cong \mathbb{1}_{P^\perp}$. Then *polarized execution formulas* are defined both on $[\![\pi]\!]$ and on f_π as follows:

$$\mathsf{Ex}([\![\pi]\!],\sigma) := \mathsf{Tr}^{U_\Delta}_{U_\Gamma,U_\Gamma}\left((\mathrm{Id}_{U_\Gamma}\otimes\sigma)\circ[\![\pi]\!]\right) = \mathsf{Tr}^{U^{2m}}_{U^n,U^n}\left((\mathrm{Id}_{U^n}\otimes\sigma)\circ[\![\pi]\!]\right)$$

$$\mathsf{Ex}(f_\pi,\sigma_*) := \mathsf{Tr}^{\mathbb{1}_\Delta}_{\mathbb{1}_\Gamma,\mathbb{1}_\Gamma}\left((\mathrm{Id}_{\mathbb{1}_\Gamma}\otimes\sigma_*)\circ[\![\pi]\!]\right) = \mathsf{Tr}^{1^{2n'}}_{1^{n'},1^{n'}}\left((\mathrm{Id}_{1^{n'}}\otimes\sigma_*)\circ f_\pi\right)$$

Note: $\mathsf{Ex}([\![\pi]\!],\sigma) \in \mathrm{End}_{\mathcal{C}}(U_\Gamma) = \mathrm{End}_{\mathcal{C}}(U^n)$ and $\mathsf{Ex}(f_\pi,\sigma_*) \in \mathrm{End}_{\mathcal{C}}(\mathbb{1}_\Gamma) = \mathrm{End}_{\mathcal{C}}(1^{n'})$, as pictured below in Figure 2.




Fig. 2. Execution formulas on $[\![\pi]\!]$ and on f_π of polarized proofs of $\vdash [\Delta],\Gamma$.

3.4. *Polarized execution formulas are an invariant for cut elimination*

In MELL, the execution formula is only an invariant of normalization for sequents not containing '?' (see Haghverdi and Scott 2006, 2011). But in MLLP, $\uparrow$ and $\downarrow$ are logically weak (although still functorial) operators. In the case of the exponentials in Abramsky *et al.* (2002), semantical axioms inspired directly from the syntactical rules of ! and ? in linear logic are imposed on top of the GoI situation. Here, for the weaker polarized case, we instead use the multi-point semantical structure to simulate the logically weak $\uparrow$ and $\downarrow$. That is the purpose of the rather subtle retraction structure of multi-points. The simulation is indirect in the sense that the notion of multi-point does not live in the syntax of MLLP. But semantically the notion is fine grained enough to sufficiently simulate $\uparrow$ and $\downarrow$. This allows us to prove the execution formula is a full invariant for cut elimination, in the sense of the following Proposition.

Proposition 3.22 (Ex is an invariant). If $\pi \to_* \pi'$ by MLLP cut elimination, then $\mathsf{Ex}([\![\pi]\!],\sigma) = \mathsf{Ex}([\![\pi']\!],\sigma')$ and $\mathsf{Ex}(f_\pi,\sigma_*) = \mathsf{Ex}(f_{\pi'},(\sigma')_*)$. In particular, if π' is a (cut free) normal form of π, then (since σ' and $(\sigma')_*$ are Id_I), we have $\mathsf{Ex}([\![\pi]\!],\sigma) = [\![\pi']\!]$ and $\mathsf{Ex}(f_\pi,\sigma_*) = f_{\pi'}$.

Proof. The crucial polarized case is where π contains cut formulas $\downarrow P^\perp$ and $\uparrow P$, which are transformed into the cut formulas $P^\perp$ and P of π' by cut elimination. The other cases

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

are similar to the non-polarized MLL case. So the crucial case is that of a proof π, ending with a cut between $\uparrow P$ and $\downarrow P^{\perp}$, which has the following form. A one-step reduction then gives rise to the proof ending with a cut between P in π_1' and $P^{\perp}$ in π_2':

$$\dfrac{\dfrac{\begin{array}{c}\vdots\ \pi_1'\\ \vdash [\cdots], P, \cdots\end{array}}{\vdash [\cdots], \uparrow P, \cdots}\uparrow \quad \dfrac{\begin{array}{c}\vdots\ \pi_2'\\ \vdash [\cdots], P^{\perp}, \mathcal{M}\end{array}}{\vdash [\cdots], \downarrow P^{\perp}, \mathcal{M}}\downarrow}{\vdash [\cdots, \uparrow P, \downarrow P^{\perp}], \cdots, \mathcal{M}}\ \text{cut}$$

Recall the definition of the interpretations of polarity changing $\uparrow$ (see 4 of Definition 3.18)), where $U_{\uparrow} \to U_{\uparrow}$ occurring in the construction of $[\![\pi]\!]$ (for the principal $\uparrow$) is the zero morphism. This directly means that in $(\mathrm{Id} \otimes s^m) \circ [\![\pi]\!]$, there arise no new loops via $U_{\uparrow}$ and $U_{\downarrow}$ for the $\uparrow P$ and $\downarrow P$ in the last cut. Hence, the trace of $(\mathrm{Id} \otimes s^m) \circ [\![\pi']\!] =$ the trace of $(\mathrm{Id} \otimes s^m) \circ [\![\pi]\!]$. Let us calculate this precisely using the trace axioms. Let π_1 (resp. π_2) denotes π_1' (resp. π_2') followed by a $\uparrow$ (resp. by $\downarrow$) rule applied to P (resp. to $P^{\perp}$). Let Γ_i (resp. Γ) be conclusions with the list of cut formulas of π_i' (resp. π).

$$\begin{aligned}
[\![\pi_1]\!]^{\dagger} &= [\![\pi_1']\!]^{\dagger} \otimes (c : U_{\uparrow} \to U_{\uparrow}) \ \text{ with } c = 0\\
[\![\pi_2]\!]^{\dagger} &= (\mathfrak{e}_{\mathcal{M}} \otimes \boldsymbol{U}_{\Gamma_2' \setminus \mathcal{M}}) \circ \{[\![\pi_2']\!] \otimes (a : U_{\downarrow} \to 1^m) \otimes (b : 1^m \to U_{\downarrow})\} \circ (\mathfrak{m}_{\mathcal{M}} \otimes \boldsymbol{U}_{\Gamma_2' \setminus \mathcal{M}})\\
&\quad \text{where } \mathbb{1}_{\mathcal{M}} = 1^m,\ a = h_m \circ \alpha^* \text{ and } b = \alpha \circ g_m.\\
[\![\pi]\!]^{\dagger} &= [\![\pi_1]\!]^{\dagger} \otimes [\![\pi_2]\!]^{\dagger}\\
&= (\hat{\mathfrak{e}}_{\mathcal{M}} \otimes \boldsymbol{U}_{\Gamma \setminus \mathcal{M}}) \circ \{([\![\pi_1']\!] \otimes c) \otimes ([\![\pi_2']\!] \otimes a \otimes b)\} \circ (\hat{r}_{\mathcal{M}} \otimes \boldsymbol{U}_{\Gamma \setminus \mathcal{M}})
\end{aligned}$$

Note that the compositions and precompositions of the middle arrows occurring in the above (R.H.S) are modulo permutations of their domains and codomains.

In the definition of $[\![\pi]\!]^{\dagger}$, we use the following retraction $(\hat{\mathfrak{e}}_A, \hat{\mathfrak{m}}_A)$ derivable from Definition 3.16 using the canonical isomorphism (5). We define $\hat{\mathfrak{e}}_A := \mathfrak{e}_A \otimes A^{\flat}$ and $\hat{\mathfrak{m}}_A := \mathfrak{m}_A \otimes A^{\flat}$:

$$\boldsymbol{U}_A \otimes \mathbb{1}_A \rhd_{(\hat{\mathfrak{e}}_A, \hat{\mathfrak{m}}_A)} \boldsymbol{U}_A \ \text{ satisfying } \hat{\mathfrak{e}}_A \circ \hat{\mathfrak{m}}_A = \boldsymbol{U}_A$$

The same definitions as above also apply for $\hat{\mathfrak{e}}_{\mathcal{M}}$ and $\hat{\mathfrak{m}}_{\mathcal{M}}$ for a sequence $\mathcal{M}$.

In the following proof, $\boldsymbol{U}_x$ is simply abbreviated by x: it is either a formula A or a sequence $\mathcal{M}$. We abbreviate $U_{\uparrow}$ and $U_{\downarrow}$ by $\uparrow$ and $\downarrow$. For the proof, it suffices to show

$$\mathsf{Tr}^{\uparrow \otimes \downarrow}_{\Gamma \setminus \{\downarrow, \uparrow\}, \Gamma \setminus \{\downarrow, \uparrow\}}((\mathrm{Id} \otimes s_{\uparrow,\downarrow} \otimes s_{P,P^{\perp}}) \circ [\![\pi]\!]^{\dagger}) = (\mathrm{Id} \otimes s_{P,P^{\perp}}) \circ ([\![\pi_1']\!]^{\dagger} \otimes [\![\pi_2']\!]^{\dagger})$$
$$\text{where } \mathrm{Id} = \mathrm{Id}_{\Gamma \setminus \{P, P^{\perp}\}} \text{ and } [\![\pi']\!]^{\dagger} = [\![\pi_1']\!]^{\dagger} \otimes [\![\pi_2']\!]^{\dagger}. \quad (11)$$

For proving Equation (11), we start with observing the following identity derivable by generalized yanking (Appendix C), vanishing and dinaturality in a TMC.


(Iterated generalized yanking)

For any morphisms $f : X \longrightarrow U$, $g : U \longrightarrow V$ and $h : V \longrightarrow Y$ in a TMC,

$$\mathsf{Tr}^{V \otimes U}_{X,Y}((s_{Y,U} \otimes V) \circ (Y \otimes s_{U,V}) \circ (h \otimes f \otimes g)) = h \circ g \circ f \quad (12)$$

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

Pictorially, this says



The L.H.S of Equation (11), by naturality w.r.t $\hat{\mathfrak{m}}_{\mathcal{M}}$ and $\hat{\mathfrak{e}}_{\mathcal{M}}$, is equal to

$$(\hat{\mathfrak{e}}_{\mathcal{M}} \otimes \mathrm{Id}_{\Gamma\setminus\{\mathcal{M},\downarrow,\uparrow\}})\circ \mathsf{Tr}^{\downarrow}_{1^m\otimes\Gamma\setminus\{\downarrow,\uparrow\},\,1^m\otimes\Gamma\setminus\{\downarrow,\uparrow\}} \{(\mathrm{Id}\otimes\sigma)\circ([\![\pi_1']\!]^\dagger \otimes c \otimes [\![\pi_2']\!]^\dagger \otimes a \otimes b)\}\circ(\hat{\mathfrak{m}}_{\mathcal{M}} \otimes \mathrm{Id}_{\Gamma\setminus\{\mathcal{M},\downarrow,\uparrow\}}) \quad (13)$$

The middle arrow of Equation (13) is, by superposing, equal to

$$(\mathrm{Id}\otimes s_{\uparrow,\downarrow}\otimes s_{P,P^\perp})\circ([\![\pi_1']\!]^\dagger \otimes [\![\pi_2']\!]^\dagger) \otimes \mathsf{Tr}^{\uparrow\otimes\downarrow}_{1^m,1^m}(a\otimes b\otimes c)$$

which by Equation (12), is equal to $(\mathrm{Id}\otimes s_{\uparrow,\downarrow}\otimes s_{P,P^\perp})\circ([\![\pi_1']\!]^\dagger \otimes [\![\pi_2']\!]^\dagger) \otimes a\circ c\circ b$.
This amounts to showing that Equation (13) is equal to the R.H.S of Equation (11) by Axiom 9′ since $a\circ c\circ b = 0$ when $c = 0$. See Appendix F for a pictorial representation of the above calculations.
The analogous assertion for f_{π_1}, f_{π_2} and f_π is easily checked in the bottom layer. □

Example 3.23 (Extrusion of ↓ boxes by expanding Ex). In the dynamics of normalization, *extrusion* of ↓ boxes depicted in Example 2.3 is captured by running the execution formula. Let us illustrate this phenomenon in Rel, considered as a polarized GoI situation arising from Haghverdi's notion of a UDC (see Appendix B). In this case, the execution formulas (defined via traces) can be written in Girard's original form of a power series, as in Equation (24) of Definition A.1 in Appendix A, and have a matrix representation.

This phenomenon is sufficiently explained by examining the lower layer (= multipoint) interpretation f_{π_i} of the three proofs ($i = 1,2,3$) together with permutation $\sigma_* = s_{\mathbb{1}_{\uparrow_1},\mathbb{1}_{\downarrow_2}} \otimes s_{\mathbb{1}_{\downarrow_3},\mathbb{1}_{\uparrow_2}}$ of cuts, which are common to the three proofs:

$$f_{\pi_i} = \begin{array}{c|cc|ccc} & \mathbb{1}_{\downarrow_1} & \mathbb{1}_{\uparrow_3} & \mathbb{1}_{\uparrow_1} & \mathbb{1}_{\downarrow_2} & \mathbb{1}_{\downarrow_3} \quad \mathbb{1}_{\uparrow_2} \\ \mathbb{1}_{\downarrow_1} & & \delta_{i3} & \delta_{i1} & & \delta_{i2} \\ \mathbb{1}_{\uparrow_3} & \delta_{i3} & & & 1 & \\ \hline \mathbb{1}_{\uparrow_1} & \delta_{i1} & & & & \\ \mathbb{1}_{\downarrow_2} & & & & & 1 \\ \mathbb{1}_{\downarrow_3} & & & 1 & & \\ \mathbb{1}_{\uparrow_2} & \delta_{i2} & & & 1 & \end{array}$$

$$\sigma_* = \begin{array}{c|cc} & \mathbb{1}_{\uparrow_1} & \mathbb{1}_{\downarrow_2} \\ \mathbb{1}_{\uparrow_1} & & 1 \\ \mathbb{1}_{\downarrow_2} & 1 & \end{array} \otimes \begin{array}{c|cc} & \mathbb{1}_{\downarrow_3} & \mathbb{1}_{\uparrow_2} \\ \mathbb{1}_{\downarrow_3} & & 1 \\ \mathbb{1}_{\uparrow_2} & 1 & \end{array}$$

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

In the above, 1 abbreviates Id_1 and δ_{ij} is the Kronecker delta. Note that the normal form π of π_1, π_2 and π_3 is the η-expansion of the axiom link, hence is interpreted by the 2×2 anti-diagonal identity matrix f_π indexed with $\mathbb{1}_{\downarrow_1}$ and $\mathbb{1}_{\uparrow_3}$.

The scope extrusion of Example 2.3 is represented by the expansion of the execution formula $\mathrm{Ex}(f_{\pi_1}, \sigma_*)$ (i.e. taking the trace of $(\mathrm{Id} \otimes \sigma_*) \circ f_{\pi_i}$). Note that the box_j with $j \in \{1,2,3\}$ is represented by the two paired elements δ_{ij} in the symmetric matrix f_{π_i}, while the symmetric pair of 1s at $(\mathbb{1}_{\downarrow_i}, \mathbb{1}_{\uparrow_i})$ and $(\mathbb{1}_{\uparrow_i}, \mathbb{1}_{\downarrow_i})$ with $i \in \{2,3\}$ represents the box introduced by $\downarrow_i$. Then, the calculation passes through $\mathrm{Ex}(f_{\pi_2}, \sigma_*)$, then $\mathrm{Ex}(f_{\pi_3}, \sigma_*)$ and finally will terminate in f_π. Here, the lower left δ_{i1} moves to the lower left δ_{i2} by the action of $(f_{\pi_1})_{22}\ \sigma_*(f_{\pi_1})_{21}$. Then it moves finally to the lower left δ_{i3} by the action of $(f_{\pi_1})_{12}\ \sigma_*(f_{\pi_1})_{22}\ \sigma_*(f_{\pi_1})_{21}$. This says that as j decreases, the sum for $\mathrm{Ex}(f_{\pi_j}, \sigma_*)$ becomes 'finer grained'; that is, the information flow realized in the sum of $\mathrm{Ex}(f_{\pi_j}, \sigma_*)$ can be retrieved from that in $\mathrm{Ex}(f_{\pi_i}, \sigma_*)$ with $i > j$. For the information flow arising from the execution formula in a UDC, the reader is referred to Appendices A.1, B and Haghverdi's thesis (Haghverdi 2000).

3.5. *Polarized execution formulas characterize focusing*

As we saw above in MLLP, the execution formula yields invariants of the dynamical process of cut elimination. In this section, we give a second property peculiar to the polarized execution formula. As far as we know it has no analogue in traditional linear logic.

Our main result is that the execution formula in polarized GoI situations is able to characterize the focusing property, which is the fundamental characteristic of polarized logics. Observe that in a polarized GoI situation, a proof π of an MLLP sequent is interpreted as a pair $([\![\pi]\!], f_\pi)$. This interpretation does not capture the focusing property. Instead, the GoI situation only provides an interpretation of polarities in terms of multi-points arising from the retractions $U \otimes 1^m \rhd U$ for $[\![\pi]\!]$ and $1^m \otimes 1^m \rhd 1^m$ for f_π. We may now ask how do we semantically obtain the stronger notion of positivity/negativity? We show that this stronger property can be characterized in terms of *preservation of multi-points* through running the execution formulas. These results are described in Theorem 3.26 below. For this proposition, we introduce the following definition:

Definition 3.24 (Restriction and range of morphisms).
Let $\mathcal{C}$ be a polarized GoI situation.

– In the presence of 0 morphisms of $\mathcal{C}$, the two morphisms ι_j and ρ_j are derivable, as follows, where $\check{K} := K \setminus \{j\}$:

$$\iota_j : \quad X_j \simeq X_j \otimes (\bigotimes_{i \in \check{K}} I) \xrightarrow{X_j \otimes (\bigotimes 0_{I, X_i})} X_j \otimes (\bigotimes_{i \in \check{K}} X_i) \simeq \bigotimes_{i \in K} X_i$$

$$\rho_j : \quad \bigotimes_{i \in K} X_i \simeq X_j \otimes (\bigotimes_{i \in \check{K}} X_i) \xrightarrow{X_j \otimes (\bigotimes 0_{X_i, I})} X_j \otimes (\bigotimes_{i \in \check{K}} I) \simeq X_j$$

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

so that $\rho_k \iota_j = X_i$ if $j = k$ and $0_{X_j,X_k}$ otherwise (These are called quasi-injections and quasi-projections in Haghverdi 2000 and Haghverdi and Scott 2011).
- For a morphism f with domain $\bigotimes_{i\in K} X_i$, its *restriction* to X_j is the morphism $f\circ\iota_j$.
- For a morphism f with codomain $\bigotimes_{i\in K} X_i$, f *ranges over* X_j if $(\iota_j\circ\rho_j)\circ f = f$.

The following lemma will be used for Theorem 3.26 (Case 4).


Lemma 3.25 (Trace on zero morphisms).
Zero morphisms satisfy the following property on tracing. For any morphism $f : X \otimes U \longrightarrow Y \otimes U$*:*

$$\mathsf{Tr}^U_{X,Y}\left(f\circ(X\otimes 0_{U,U})\right) \quad = \quad (X \otimes 0_{U,I})\circ f\circ(X \otimes 0_{I,U}) \quad = \quad \mathsf{Tr}^U_{X,Y}\left((X\otimes 0_{U,U})\circ f\right)$$

Proof. We prove the first equation (a similar calculation goes for the second).
First, decompose $0_{U,U} = 0_{I,U}\circ 0_{U,I}$.
$\mathsf{Tr}^U_{X,Y}\left(f\circ(X\otimes 0_{I,U}\circ 0_{U,I})\right) = \mathsf{Tr}^I_{X,Y}\left((X\otimes 0_{U,I})\circ f\circ(X\otimes 0_{I,U})\right) = (X\otimes 0_{U,I})\circ f\circ(X\otimes 0_{I,U})$
The first equation is by dinaturailty and the second one is by vanishing. □

Notation: for a sequence Γ containing a formula A, consider the composition $\mathbb{1}_A \xrightarrow{\mathsf{mp}(A)} U_A \xrightarrow{\iota} U_\Gamma$, where ι is the quasi-injection induced by the embedding of A into Γ. This is denoted (by abuse of notation) by $\mathsf{mp}(A) : \mathbb{1}_A \longrightarrow U_\Gamma$, leaving the ι implicit.

Theorem 3.26 (focusing = invariance of mp**).** *Let* π *be an* MLLP *proof of a sequent* $\vdash [\Delta], \mathcal{N}, P$*, where* P *is a positive formula. Then the execution formulas over* $[\![\pi]\!]$ *and over* f_π *give rise to the following commutative diagram:*




The bottom arrow of this diagram indicates that $\mathsf{Ex}(f_\pi,\sigma_*)$*, when its domain is restricted to* $\mathbb{1}_P$*, ranges over* $\mathbb{1}_{\mathcal{N}}$.

Proof. We prove the proposition directly for Case 1 (axiom) and Case 4 ($\downarrow$ rule) and by induction on the size of π for the other cases. *Note:* Symbols used in the proof are the same as those in Definition 3.18.

1. *Axiom*:
The following diagram commutes because multi-points associated with De Morgan dual formulas are equal by the duality-induced permutation on their domains (which are

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

foldings of tensors of 1's).



2. *Linear connectives:*
We prove it for a proof π ending with the $\otimes$ rule (the result is obvious for the $\parr$ rule). Suppose π is


$$\dfrac{\begin{array}{cc}\vdots\ \pi_1 & \vdots\ \pi_2\\ \vdash [\Delta_1], \mathcal{M}_1, P_1 & \vdash [\Delta_2], \mathcal{M}_2, P_2\end{array}}{\vdash [\Delta_1, \Delta_2], \mathcal{M}_1, \mathcal{M}_2, P_1 \otimes P_2}\ \otimes$$

First, note that

$$\begin{aligned}\mathsf{Ex}([\![\pi_1]\!] \otimes [\![\pi_2]\!], \sigma_1 \otimes \sigma_2) &= \mathsf{Ex}([\![\pi_1]\!], \sigma_1) \otimes \mathsf{Ex}([\![\pi_2]\!], \sigma_2)\\ \mathsf{Ex}(f_{\pi_1} \otimes f_{\pi_2}, (\sigma_1)_* \otimes (\sigma_2)_*) &= \mathsf{Ex}(f_{\pi_1}, (\sigma_1)_*) \otimes \mathsf{Ex}(f_{\pi_2}, (\sigma_2)_*)\end{aligned}$$

where the σ_i are iterated tensors of symmetries s, representing the cut formulas in π_i, for $i = 1, 2$.

Then commutativity of the first diagram by I.H. yields directly that of the second where $n = n_1 + n_2$:




3. *Polarity Changing* $\uparrow$*:*
This case never happens since a conclusion of any such proof does not contain a positive formula.

4. *Polarity Changing* $\downarrow$*:* This case does not use the I.H. Instead it uses Lemma 3.25 directly. This lemma connects traces with compositions of zero morphisms (see the discussion after Equation (16) below). We begin with an equation which is entailed from the commutativity of the following diagram (to be discussed below)[‖]:

$$\alpha^m \circ f_\pi = [\![\pi]\!]^\dagger \circ \alpha, \qquad \text{where } f_\pi \text{ is restricted to } \mathbb{1}_{\downarrow N} = 1. \tag{14}$$

[‖] For typographical simplicity, we rotated the diagram 90° clockwise.

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press



In the diagram, we let $\mathbb{1}_\mathcal{M} := 1^m$ and Ξ denotes Δ, N. Why does this diagram commutes? The diagram consists of four regions from left to right: modulo symmetry, (i) a leftmost pentagon bordered by f_π, (ii) a middle upper hexagon bordered by $1^m \otimes \alpha^m \otimes V$, (iii) a middle lower square bordered by $1^m \otimes \alpha^m \otimes V$ and (iv) a rightmost square bordered by $[\![\pi]\!]^\dagger$. The left pentagon commutes from the definition of f_π. The upper middle hexagon commutes because $\alpha^* \circ \alpha = \mathrm{Id}_1$. The lower middle square is $V\otimes$ lifting along α^m of Proposition 3.17, hence commutes. The rightmost square commutes by definition of $[\![\pi]\!]^\dagger$. These commutativities imply the outermost hexagon commutes.

Then the following is a commutative square for Equation (14).

$$\begin{array}{ccc} \boldsymbol{U}_{\Delta,\mathcal{M},\downarrow N} & \xrightarrow{[\![\pi]\!]^\dagger} & \boldsymbol{U}_{\Delta,\mathcal{M},\downarrow N} \\ {\scriptstyle \alpha = \mathsf{mp}(\downarrow N)} \uparrow & & \uparrow {\scriptstyle \alpha^m = \mathsf{mp}(\mathcal{M})} \\ \mathbb{1}_{\downarrow N} & \xrightarrow[f_\pi]{} & 1^m \end{array} \tag{15}$$

Both $\sigma \otimes \boldsymbol{U}_{\mathcal{M},\downarrow N}$ and $0_{\boldsymbol{U}_\Delta} \otimes \boldsymbol{U}_{\mathcal{M},\downarrow N}$ (resp. both $\sigma_* \otimes \mathbb{1}_{\mathcal{M},\downarrow N}$ and $0_{\mathbb{1}_\Delta} \otimes \mathbb{1}_{\mathcal{M},\downarrow N}$) act identically on the component $\boldsymbol{U}_\mathcal{M}$ of the codomain (resp. on the codomain 1^m) of α^m. Thus, we have

$$\begin{aligned} (\sigma \otimes \boldsymbol{U}_{\mathcal{M},\downarrow N}) \circ [\![\pi]\!]^\dagger \circ \alpha &= (0_{\boldsymbol{U}_\Delta} \otimes \boldsymbol{U}_{\mathcal{M},\downarrow N}) \circ [\![\pi]\!]^\dagger \circ \alpha, \qquad \text{and} \\ \alpha^m \circ (\sigma_* \otimes \mathbb{1}_{\mathcal{M},\downarrow N}) \circ f_\pi &= \alpha^m \circ (0_{\mathbb{1}_\Delta} \otimes \mathbb{1}_{\mathcal{M},\downarrow N}) \circ f_\pi. \end{aligned} \tag{16}$$

By composing and precomposing the upper (resp. the lower) horizontal arrow of Equation (15) with $0_{\boldsymbol{U}_\Delta} \otimes \boldsymbol{U}_{\mathcal{M},\downarrow N}$ (resp. with $0_{\mathbb{1}_\Delta} \otimes \mathbb{1}_{\mathcal{M},\downarrow N}$), we have, by Lemma 3.25,

$$\mathsf{Tr}^{\boldsymbol{U}_\Delta}_{\boldsymbol{U}_{\mathcal{M},\downarrow N},\boldsymbol{U}_{\mathcal{M},\downarrow N}} \left((\sigma \otimes \boldsymbol{U}_{\mathcal{M},\downarrow N}) \circ [\![\pi]\!]^\dagger \right) \circ \alpha = \alpha^m \circ \mathsf{Tr}^{\mathbb{1}_\Delta}_{\mathbb{1}_{\mathcal{M},\downarrow N},\mathbb{1}_{\mathcal{M},\downarrow N}} \left((\sigma_* \otimes \mathbb{1}_{\mathcal{M},\downarrow N}) \right) \circ f_\pi.$$

This is the assertion $\mathsf{Ex}([\![\pi]\!]^\dagger, \sigma) \circ \alpha = \alpha^m \circ \mathsf{Ex}(f_\pi, \sigma_*)$.

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

5. *Cut*[††]:
This case uses the following property of the associativity of cut (cf. Haghverdi and Scott 2011):

$$\mathsf{Ex}(\llbracket\pi\rrbracket, \sigma \otimes s) = \mathsf{Ex}(\mathsf{Ex}(\llbracket\pi\rrbracket, \sigma), s) \quad \text{and} \quad \mathsf{Ex}(f_\pi, \sigma_* \otimes s_*) = \mathsf{Ex}(\mathsf{Ex}(f_\pi, \sigma_*), s_*)$$

where s corresponds to the last cut of π.

The sequence Ξ of Case 2 of Definition 3.18 must be of the form $Q, \mathcal{M}$ with a positive Q: i.e. π is

$$\dfrac{\begin{array}{cc}\vdots\ \pi' & \vdots\ \pi'' \\ \vdash [\Delta'], P, \mathcal{N} & \vdash [\Delta''], P^\perp, Q, \mathcal{M}\end{array}}{\vdash [\Delta', \Delta'', P, P^\perp], \mathcal{N}, Q, \mathcal{M}}\ \text{cut}$$

Note first that $\sigma = \sigma' \otimes \sigma''$ with σ' (resp. σ'') representing all the cuts of π' (resp. π'').

The I.H. implies the following two diagrams commute:

$$\begin{array}{ccc} U_{P,\mathcal{N}} & \xrightarrow{\mathsf{Ex}(\llbracket\pi'\rrbracket,\sigma')} & U_{P,\mathcal{N}} \\ {\scriptstyle \mathsf{mp}(P)}\uparrow & & \uparrow{\scriptstyle \mathsf{mp}(\mathcal{N})} \\ \mathbb{1}_P & \xrightarrow[\mathsf{Ex}(f_{\pi'},\sigma'_*)]{} & \mathbb{1}_{\mathcal{N}} \end{array} \qquad \begin{array}{ccc} U_{P^\perp,Q,\mathcal{M}} & \xrightarrow{\mathsf{Ex}(\llbracket\pi''\rrbracket,\sigma'')} & U_{P^\perp,Q,\mathcal{M}} \\ {\scriptstyle \mathsf{mp}(Q)}\uparrow & & \uparrow{\scriptstyle \mathsf{mp}(P^\perp,\mathcal{M})} \\ \mathbb{1}_Q & \xrightarrow[\mathsf{Ex}(f_{\pi''},\sigma''_*)]{} & \mathbb{1}_{P^\perp} \otimes \mathbb{1}_{\mathcal{M}} \end{array}$$

Since $\llbracket\pi\rrbracket \cong \llbracket\pi'\rrbracket \otimes \llbracket\pi''\rrbracket$ modulo permutation of interface,

$$\mathsf{Ex}(\llbracket\pi\rrbracket, \sigma) \cong \mathsf{Ex}(\llbracket\pi'\rrbracket, \sigma') \otimes \mathsf{Ex}(\llbracket\pi''\rrbracket, \sigma'')$$

Then,

$$(U_{\mathcal{N},Q,\mathcal{M}} \otimes s_{U_P,U_{P^\perp}}) \circ \mathsf{Ex}(\llbracket\pi\rrbracket, \sigma) \cong (U_{\mathcal{N},Q,\mathcal{M}} \otimes s_{U_P,U_{P^\perp}}) \circ (\mathsf{Ex}(\llbracket\pi'\rrbracket, \sigma') \otimes \mathsf{Ex}(\llbracket\pi''\rrbracket, \sigma''))$$


This composing morphisms is realized by the upper horizontal morphisms of the following composition of the two commutative squares whose left one is obtained by tensoring the above two squares:

$$\begin{array}{ccccc} U_{P,\mathcal{N}} \otimes U_{P^\perp,Q,\mathcal{M}} & \xrightarrow{\mathsf{Ex}(\llbracket\pi'\rrbracket,\sigma')\otimes\mathsf{Ex}(\llbracket\pi''\rrbracket,\sigma'')} & U_{P,\mathcal{N}} \otimes U_{P^\perp,Q,\mathcal{M}} & \xrightarrow{U_{\mathcal{N},Q,\mathcal{M}}\otimes s_{U_P,U_{P^\perp}}} & U_{\mathcal{N},Q,\mathcal{M}} \otimes U_{P,P^\perp} \\ \uparrow{\scriptstyle \mathsf{mp}(P)\otimes\mathsf{mp}(Q)} & & \uparrow{\scriptstyle \mathsf{mp}(P^\perp)\otimes\mathsf{mp}(\mathcal{N},\mathcal{M})} & & \uparrow{\scriptstyle \mathsf{mp}(P)\otimes\mathsf{mp}(\mathcal{N},\mathcal{M})} \\ \mathbb{1}_P \otimes \mathbb{1}_Q & \xrightarrow[\mathsf{Ex}(f(\pi'),\sigma'_*)\otimes\mathsf{Ex}(f(\pi''),\sigma''_*)]{} & \mathbb{1}_{\mathcal{N}} \otimes \mathbb{1}_{P^\perp} \otimes \mathbb{1}_{\mathcal{M}} & \xrightarrow[\mathbb{1}_{\mathcal{N},Q,\mathcal{M}}\otimes s_{\mathbb{1}_P,\mathbb{1}_{P^\perp}}]{} & \mathbb{1}_{\mathcal{N}} \otimes \mathbb{1}_{\mathcal{M}} \otimes \mathbb{1}_P \end{array}$$

In the outermost square, taking trace along $U_P \otimes U_{P^\perp}$ (resp. along $\mathbb{1}_P \otimes \mathbb{1}_{P^\perp}$) of the upper (resp. the lower) horizontal morphism, by Corollary 3.12 (strong uniformity), we have the following diagram, which is the assertion by virtue of the associativity mentioned at the beginning of this case.

[††] This is the case where Corollary 3.12 (strong uniformity) is used.

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press



□

Example 3.27 (Invariance of Theorem 3.26). We give two examples of Theorem 3.26 for the interpretations shown in Example 3.19. In these examples, $\mathsf{Ex}([\![\pi]\!],\sigma)$ and $\mathsf{Ex}(f_\pi,\sigma_*)$ are $[\![\pi]\!]$ and f_π, respectively, since π is cut free.

i. In this case, $P = \downarrow X^\perp$ and $\mathcal{M} = \uparrow X$.
Since f_π restricted to $\mathbb{1}_{\downarrow X^\perp} := 1$ ranges over $\mathbb{1}_{\uparrow X}$, the Proposition follows from

$$[\![\pi]\!]^\dagger \circ (\alpha : \mathbb{1}_{\downarrow X^\perp} \longrightarrow U_\downarrow) = (\alpha : \mathbb{1}_{\uparrow X} \longrightarrow U_\uparrow) \circ f_\pi \quad \text{under this restriction.}$$


Recall that in the Rel examples, the equated morphisms are identified with $\{n_\alpha\}$.
ii. In this case, $P = \downarrow\uparrow(X \otimes Y)$ and $\mathcal{M} = Y^\perp \parr X^\perp$.
Since f_π restricted to $\mathbb{1}_{\downarrow\uparrow(X\otimes Y)}$ ranges over $\mathbb{1}_{Y^\perp} \otimes \mathbb{1}_{X^\perp}$, the commutativity follows from

$$[\![\pi]\!]^\dagger \circ (\alpha : \mathbb{1}_{\downarrow\uparrow(X\otimes Y)} \longrightarrow U_\downarrow) = (\alpha^2 : \mathbb{1}_{Y^\perp \parr X^\perp} \longrightarrow \boldsymbol{U}_{Y^\perp \parr X^\perp}) \circ f_\pi \quad \text{under the restriction.}$$

The equated morphisms are identified with $\{n_\alpha\} + \{n_\alpha\}$.

We now show there is a kind of 'converse': any MLLP-provable sequent $\vdash [\Delta], \mathcal{N}, A$ with A invariant in the sense of the diagram in Theorem 3.26 must have A positive. More precisely:

Proposition 3.28 (Converse of focusing). *Let $\vdash [\Delta], \mathcal{M}, A$ be a sequent provable in MLLP such that A contains a polarity changing connective and $\mathcal{M}$ is a sequence of negative formulas. If the following diagram non-trivially commutes in all models*[‡‡]



then A is a positive formula.

Proof. It suffices to prove the assertion for cut-free proofs of $\vdash \mathcal{M}, A$ so that $\mathsf{Ex}([\![\pi]\!],\sigma)$ and $\mathsf{Ex}(f_\pi,\sigma_*)$ become, respectively, $[\![\pi]\!]$ and f_π because of the invariance of cut elimination

‡‡ We say a commutative diagram *trivially commutes* if the unique arrow in it is the zero morphism, and *non-trivially commutes* if it is not the zero morphism.

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

of Proposition 3.22. For the proof, we make the following observation, which is proved for f_π by induction on the construction of proofs π:

For a positive formula Q and any negative formula M in a conclusion of a cut-free focused proof π, the following inequality holds in any Rel *polarized GoI situation:*

$$[\mathbb{1}_M] f_\pi \cap \mathbb{1}_Q \neq \varnothing \tag{17}$$

(cf. Notation 5.1 for Rel in Section 5 below).

Proof of (17): Note first that the last rule of π must not be an $\uparrow$ rule. Since the induction is straightforward for the axiom and linear connectives $\otimes$ and $\parr$, we prove the case when the last rule is $\downarrow$, so that $Q = \downarrow N$. The corresponding case of Definition 3.18 says that $[\mathbb{1}_{\mathcal{M}}] f_\pi \supseteq \mathbb{1}_{\downarrow N}$, where $\mathbb{1}_{\mathcal{M}} = 1^m$ and $\mathbb{1}_{\downarrow N} = 1$, which directly implies Equation (17). (See the last I/O diagram of Definition 3.18 for the inclusion, where $\mathbb{1}_{\mathcal{M}}$ of the input on the left first goes via ρ to 1^m, which splits via $\mathfrak{m}^m$, and the bottom output goes to the 1 of the (right-hand) side, by g_m.) *End of Proof of (17)*

We prove Proposition 3.28 by contradiction; suppose that A is negative and consider the polarized GoI situation Rel. From the assumption, the given cut-free proof π has a bottom most $\uparrow$ rule, say I, whose principal formula is denoted by $\uparrow Q$. So any inference (if it exists) below I is either the $\parr$ rule or exchange. Let π' be the subproof of π ending at the premise of I. See the proof figure below for π', where $\mathcal{N}$ is a certain sequence (including the empty one) of negative formulas. π' is focused with the formula Q positive.

$$\pi = \dfrac{\dfrac{\dfrac{\vdots\ \pi'}{\vdash \mathcal{N}, Q}}{\vdash \mathcal{N}, \uparrow Q}\, I}{\dfrac{\parr\text{'s and exchanges}}{\vdash \mathcal{M}, A}}$$

We have the following equation:

$$f_\pi \cong f_{\pi'} \otimes 0_{\mathbb{1}_{\uparrow Q}, \mathbb{1}_{\uparrow Q}} \qquad \text{modulo exchange rules below } I. \tag{18}$$

First, we observe the following claim used in (Case 1) and (Case 2.2) below:

(**Claim**) *The inequality (17) for $M = A$ and the condition $\mathbb{1}_Q \cap \mathbb{1}_{\mathcal{M}} = \varnothing$ yield a contradiction.*

The contradiction is that f_π restricted to $\mathbb{1}_A$ ranges over $\mathbb{1}_{\mathcal{M}}$ (the commutative diagram in the assertion).

(Case 1) The case where the distinguished occurrence $\uparrow Q$ occurs in $\mathcal{M}$. In this case, $\mathcal{M}$ is constructed from negative formulas and the $\uparrow Q$ via $\parr$ connectives and commas. The conclusion of π' has occurrences Q and a factorization A' of A by means of $\parr$ rules below I: That is, A is either A' or $A' \parr \check{A}$ for some subformula $\check{A}$. Note that $\mathbb{1}_{A'} \subseteq \mathbb{1}_A$ because of the factorization. Because π' is focused, we have $[\mathbb{1}_{A'}] f_{\pi'} \cap \mathbb{1}_Q \neq \varnothing$ by Equation (17). This directly implies $[\mathbb{1}_A] f_\pi \cap \mathbb{1}_Q \neq \varnothing$ by Equation (18). On the other hand, since $\uparrow Q$

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

occurs in $\mathcal{M}$ but $\mathbb{1}_{\uparrow Q} \cap \mathbb{1}_Q = \varnothing$, we have $\mathbb{1}_Q \cap \mathbb{1}_{\mathcal{M}} = \varnothing$. Thus, we have a contradiction by the above claim.

(Case 2) The case where the distinguished occurrence $\uparrow Q$ occurs in A. In this case, A is constructed from negative formulas and the $\uparrow Q$ via $⅋$ connectives.

(Case 2.1) The case where $A = \uparrow Q$. From the definition of the interpretation of the $\uparrow$-rule, f_π restricted to $\mathbb{1}_A = 1$ is 0, hence the diagram trivially commutes in the sense of footnote 7.

(Case 2.2) Otherwise. In this case, Q and another factor A' both occur in the conclusion of π', where A' as well as $\uparrow Q$ is a factorization of A by means of $⅋$ rules below I. That is, A is (modulo commutativity of $⅋$) either $\uparrow Q ⅋ A'$ or $\uparrow Q ⅋ A' ⅋ \check{A}$ for some subformula $\check{A}$. By the same argument for Q and A' as Case 1, we have $[\mathbb{1}_{A'}]f_{\pi'} \cap \mathbb{1}_Q \neq \varnothing$ by Equation (17). This directly implies $[\mathbb{1}_A]f_\pi \cap \mathbb{1}_Q \neq \varnothing$ by Equation (18). On the other hand, since this Q does not occur in $\mathcal{M}$, $\mathbb{1}_Q \cap \mathbb{1}_{\mathcal{M}} = \varnothing$. Thus, we have a contradiction by the above claim. □

4. Constructing a compact polarized category via GoI

In this section, we describe certain kinds of polarized categories arising from a different view of GoI; namely, from Joyal–Street–Verity's Int-construction in the multi-pointed setting. Although the material of this section is formally independent of the Section 3, it highlights different aspects of GoI arising from Int-constructions, in particular the construction of compact polarized models in the sense of our previous paper (Hamano and Scott 2007).

4.1. *Polarized logic and polarized categories*

In Section 2, we described O. Laurent's polarized multiplicative linear logic MLLP. For references to polarized categories, the reader is referred to our own paper (Hamano and Scott 2007), as well as the original sources referred to there. We begin with a very general definition of a polarized categorical model for MLLP. Our models are included among Cockett–Seely polarized categories (Cockett and Seely 2007), although theirs are considerably more general. The definition below is simpler than assumed in our previous paper (Hamano and Scott 2007), which emphasized full completeness theorems. As also mentioned there, our previous categorical models (as well as the categorical models below) are related to the dialogue categories and dialogue chiralities of Melliès (2006–2012); Melliès (2013), although our motivation arises from the proof theory of MLLP.

Definition 4.1 (Polarized categories). A *polarized category* $(\langle \mathcal{C}_+, \mathcal{C}_- \rangle, \widehat{\mathcal{C}})$ consists of the following data:

— A pair of monoidal categories $(\mathcal{C}_+, \otimes)$ and $(\mathcal{C}_-, ⅋)$ called *positive* (resp. *negative*).
— A contravariant monoidal equivalence $(\ )^\perp$ of the two categories:

$$(-)^\perp : (\mathcal{C}_+)^{op} \xrightarrow{\cong} \mathcal{C}_-$$

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

— (*Polarity changing functors*) A pair of adjoint functors $\uparrow \dashv \downarrow$, where $\uparrow : \mathcal{C}_+ \longrightarrow \mathcal{C}_-$ and $\downarrow : \mathcal{C}_- \longrightarrow \mathcal{C}_+$. Diagrammatically,

$$\mathcal{C}_+ \underset{\downarrow}{\overset{\uparrow}{\rightleftarrows}} \mathcal{C}_- \quad (\perp)$$

— De Morgan duality for $\downarrow$ and $\uparrow$ (w.r.t. the monoidal equivalence):

$$(\downarrow X)^\perp \cong \uparrow X^\perp \text{ and } (\uparrow X)^\perp \cong \downarrow X^\perp$$

— A bimodule[§§] $\widehat{\mathcal{C}} : \mathcal{C}_+^{op} \times \mathcal{C}_- \to \mathbf{Set}$ satisfying that there is a natural equivalence:

$$\widehat{\mathcal{C}}(P, N) \cong \mathcal{C}_+(P, \downarrow N) \quad \text{for all } P \in \mathcal{C}_+, N \in \mathcal{C}_-.$$

This may be written as a reversible 'rule':

$$\frac{\widehat{\mathcal{C}}(P, N)}{\mathcal{C}_+(P, \downarrow N)} \downarrow$$

Remark 4.2. By duality, there is a dual rule:


$$\frac{\mathcal{C}_-(\uparrow P, N)}{\widehat{\mathcal{C}}(P, N)} \uparrow$$

In the language of distributors (profunctors), we are demanding that $\widehat{\mathcal{C}}$ be left and right representable (Joyal 2011) and compatible with $(\ )^\perp$.

At the *-autonomous category level $(\otimes, \parr, (\ )^\perp)$, Blute and Scott (2004) the various coherence theorems in Cockett *et al.* (2006) guarantee negation can be taken to be strictly involutive (along with strict monoidal structure for each monoidal product). We have not, however, examined the more general question of strictness for polarized categories, Mac Lane (1971).

Remark 4.3 (The case where the profunctor $\widehat{\mathcal{C}}$ is a hom functor $\mathcal{C}(-,-)$).

A polarized category arises from the following framework of adjoint functors $L \dashv \downarrow$ and $\uparrow \dashv R$ with contravariant equivalence $(\ )^\perp$, as shown in the following diagram:



[§§] Intuitively, families of *non-focused proofs*

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

These functors are subject to a natural isomorphism $(\uparrow L(\ ))^{\perp} \simeq \downarrow R(\ )^{\perp}$ between two endofunctors on $\mathcal{C}_+$. In this framework, the polarity changing functors for Definition 4.1 are defined by $\uparrow \circ L$ and $\downarrow \circ R$.

The following definition of compactness is sufficient for the purposes of this paper, although more general frameworks are possible.

Definition 4.4 (Compact polarized category). We say a polarized category is *compact* if

— $\mathcal{C}_+^{op} = \mathcal{C}_-$.

— $(A \otimes B)^{\perp} = B^{\perp} \otimes A^{\perp}$ for all objects A, B.

A compact polarized category is *degenerate* if $\downarrow$ (equivalently $\uparrow$) is the identity functor.

Examples of compact polarized categories include various categories of multi-pointed relations (arising in work of Hamano and Takemura 2008), as well as various polarized Int-categories arising from GoI, to be discussed below. The reader is referred to Section 5 for the above-mentioned category of multi-pointed relations. An analogous approach to pointed relations is seen in Ehrhard's category PpL of preorders with projections (Ehrhard 2012).

Remark 4.5. Although the notion of compact polarized category may appear to be 'degenerate' in some informal sense, nevertheless the notion is sufficiently robust to distinguish the two key proofs in our paper (Hamano and Scott 2007) (Example 2.2). In other words, compact polarized categories are adequate to account for $\downarrow$ boxes in MLLP. These boxes are intrinsic for MLLP, but not for the weaker logic $\mathsf{MLL}^{\downarrow\uparrow}$ of Laurent (2002).

4.2. *The* Int *construction*

The original connection of GoI to categories was realized by several researchers (e.g. M. Hyland and S. Abramsky) as being related to the so-called Int-construction in the original papers of Joyal *et al.* (1996). Further history and related notions are discussed in the paper of Abramsky *et al.* (2002), Haghverdi and Scott (2010), Blute and Scott (2004).

Given a TMC $\mathcal{C}$, we can define a compact closed category $\mathsf{Int}(\mathcal{C})$ as follows: an object is a pair (A^+, A^-) of $\mathcal{C}$-objects and a morphism $f : (A^+, A^-) \longrightarrow (B^+, B^-)$ is a $\mathcal{C}$-morphism $f : A^+ \otimes B^- \longrightarrow B^+ \otimes A^-$. The composition of $\mathsf{Int}(\mathcal{C})$ is defined by $g \circ f :=$

$$\mathsf{Tr}^{B^-}_{A^+ \otimes C^-,\, C^+ \otimes A^-}((C^+ \otimes s_{B^-,A^-}) \circ (g \otimes A^-) \circ (B^+ \otimes s_{A^-,C^-}) \circ (f \otimes C^-) \circ (A^+ \otimes s_{C^-,B^-})). \tag{19}$$

This composition is shown in Figure 3 below, after Proposition 4.8. An arrow $(A^+, A^-) \xrightarrow{f} (B^+, B^-) \in \mathsf{Int}(\mathcal{C})$ is really an arrow $A^+ \otimes B^- \xrightarrow{f} B^+ \otimes A^- \in \mathcal{C}$. We picture the arrow pointing upwards and denote it by a box, with the four objects at the corners. The domain $A^+ \otimes B^-$ is denoted by the lower edge, the codomain $B^+ \otimes A^-$ denoted by the upper edge.

We would like to think of an $\mathsf{Int}(\mathcal{C})$ morphism $f : (A^+, A^-) \longrightarrow (B^+, B^-)$ intuitively as a bidirectional data flow: a *pair* of arrows, one from A^+ to B^+ and the other 'backwards' from B^- to A^-. Unfortunately, this view of f is only heuristic; officially, f is not a

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

tensor of two maps going in opposite directions, i.e. $f \neq g \otimes h$, where $g : A^+ \to B^+$ and $h : B^- \to A^-$. However, in the following subsections, we shall explicitly model this bidirectional dataflow by using multi-points.


4.3. *A polarized* Int *construction*

In what follows in this subsection, let $\mathcal{C}$ be a TMC

$$(\mathcal{C}, \otimes, I, s, 1, (\mathfrak{e}, \mathfrak{m}), 0)$$

with 0 morphisms (Axiom 8) and a distinguished object 1 satisfying the retraction $1 \otimes 1 \rhd_{(\mathfrak{e},r)} 1$ (Axiom 3) of Definition 3.1 of the previous Section. Moreover, $\mathcal{C}$ is supposed to satisfy the following variations of the lifting properties (Axiom 6′) and (Axiom 9) in which U can be taken to be *any* object A of $\mathcal{C}$ and β can be *any* morphism $1^m \longrightarrow A$[¶¶]:


Axiom 6‴ :(Lifting Property along β)
For any object A of $\mathcal{C}$, any $m \in \mathbb{Z}^+$ and any morphism $\beta : 1^m \longrightarrow A$, there exists a retraction pair $A \otimes 1^m \rhd_{(\mathfrak{e}_\beta, \mathfrak{m}_\beta)} A$ lifting the retraction $1^m \otimes 1^m \rhd_{(\mathfrak{e}^m, \mathfrak{m}^m)} 1^m$ along β:



Axiom 9″
For any morphism $f : V \otimes X_1 \longrightarrow W \otimes X_2$ with $X_i = A_i$ or $X_i = 1^{m_i}$ $(i = 1, 2)$ for any non-zero natural number m_i, and any morphism $\beta_i : 1^{m_i} \longrightarrow A_i$,

$$(W \otimes \mathfrak{e}_{\beta_2}) \circ (f \otimes 0_{1^{m_1}, 1^{m_2}}) \circ (V \otimes \mathfrak{m}_{\beta_1}) = f \quad \text{and} \quad (W \otimes \mathfrak{e}) \circ (f \otimes 0_{1^{m_1}, 1^{m_2}}) \circ (V \otimes \mathfrak{m}) = f$$

See the following figure for the respective equations when $X = A$ and $X = 1$:



Notation 4.6 (Morphism f_{X_i, Y_j}). For a morphism $f : X_0 \otimes X_1 \longrightarrow Y_0 \otimes Y_1$, we denote by $f_{X_i, Y_j} : X_i \longrightarrow Y_j$ the following composition:

[¶¶] Note: Unlike in the previous Section, here A is not assumed to have any retraction structure and we also assume $A \otimes 1^m \rhd_{(\mathfrak{e}_\beta, \mathfrak{m}_\beta)} A$ lifts the m-fold retraction structure of $1 \otimes 1 \rhd_{(\mathfrak{e},r)} 1$ on 1.

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

$$X_i \simeq X_i \otimes I \xrightarrow{X_i \otimes 0_{I,X_{1-i}}} X_i \otimes X_{1-i} \simeq X_0 \otimes X_1 \xrightarrow{f} Y_0 \otimes Y_1 \simeq Y_j \otimes Y_{1-j} \xrightarrow{Y_j \otimes 0_{Y_{1-j},I}} Y_j \otimes I \simeq Y_j$$

Definition 4.7 (Positive category $\mathsf{Int}_P(\mathcal{C})$). The positive category $\mathsf{Int}_P(\mathcal{C})$ consists of the following data:

— Objects are multi-pointed objects of $\mathsf{Int}_P(\mathcal{C})$:

$$(A^+_{\alpha^+}, A^-_{\alpha^-})$$

where (A^+, A^-) is an object of $\mathsf{Int}(\mathcal{C})$ and for $\star \in \{+,-\}$, $\alpha^\star$ is a morphism $1^{m^\star} \xrightarrow{\alpha^\star} A^\star$, whose domain is the $m^\star$-ary tensor folding of 1. Here, $m^\star$ is a natural number associated to $\alpha^\star$ and the * refers to the sign of the codomain of α. We call $\alpha^\star$ *a multi-point of* $A^\star$.

— Morphisms are three-tuples:

$$(A^+_{\alpha^+}, A^-_{\alpha^-}) \xrightarrow{(f,f_+,f_-)} (B^+_{\beta^+}, B^-_{\beta^-})$$

where

– $f : (A^+, A^-) \longrightarrow (B^+, B^-)$ is a morphism in $\mathsf{Int}(\mathcal{C})$.


– f_+ and f_- are morphisms in $\mathcal{C}$ making the following respective diagrams commute (see Notation 4.6):

$$\begin{array}{ccc} A^+ & \xrightarrow{f_{A^+,B^+}} & B^+ \\ {\scriptstyle\alpha^+}\uparrow & & \uparrow{\scriptstyle\beta^+} \\ 1^{m^+} & \xrightarrow[f_+]{} & 1^{n^+} \end{array} \qquad \begin{array}{ccc} B^- & \xrightarrow{f_{B^-,A^-}} & A^- \\ {\scriptstyle\beta^-}\uparrow & & \uparrow{\scriptstyle\alpha^-} \\ 1^{n^-} & \xrightarrow[f_-]{} & 1^{m^-} \end{array}$$

– f makes the following diagram commute:

$$\begin{array}{ccc} B^- & \xrightarrow{f_{B^-,B^+}} & B^+ \\ {\scriptstyle\beta^-}\uparrow & & \uparrow{\scriptstyle\beta^+} \\ 1^{n^-} & \xrightarrow[0]{} & 1^{n^+} \end{array}$$

Diagrammatically, a morphism (f, f_+, f_-) of $\mathsf{Int}_P(\mathcal{C})$ is described as follows:



In the above diagram, we say f_+ and f_- *represent* the bidirectional dataflow implicit in the upper arrow f.

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press




Fig. 3. Composition in $\mathsf{Int}(\mathcal{C})$.




Fig. 4. Composition in $\mathsf{Int}_P(\mathcal{C})$ for multi-points.

Proposition 4.8. $\mathsf{Int}_P(\mathcal{C})$ *forms a monoidal category.*

Proof. $\mathrm{Id}_{(A^+_{\alpha^+}, A^-_{\alpha^-})}$ is defined by $(\mathrm{Id}_{(A^+,A^-)}, \mathrm{Id}_+, \mathrm{Id}_-)$, where $\mathrm{Id}_\star$ is Id on the domain of $\alpha^\star$ with $\star \in \{+,-\}$. Since the first element of the tuple is $\mathrm{Id}_{A^+} \otimes \mathrm{Id}_{A^-}$ in $\mathcal{C}$, $\mathrm{Id}_{(A^+_{\alpha^+}, A^-_{\alpha^-})}$ belongs to $\mathsf{Int}_P(\mathcal{C})$.

It is immediate that the tensor product of $\mathsf{Int}(\mathcal{C})$ restricts to a tensor product in $\mathsf{Int}_P(\mathcal{C})$, forming a monoidal subcategory. We show that $\mathsf{Int}(\mathcal{C})$ composition preserves the positivity of morphisms. Recall the composition of two $\mathsf{Int}(\mathcal{C})$ morphisms $(A^+, A^-) \xrightarrow{f} (B^+, B^-)$ and $(B^+, B^-) \xrightarrow{g} (C^+, C^-)$ given in Equation (19) and Figure 3 above.

The general $\mathsf{Int}_P(\mathcal{C})$ composition with multi-points is shown in Figure 5 below. It represents the morphism $(g, g_+, g_-) \circ (f, f_+, f_-)$. Here, the top plane corresponds to the ordinary $\mathsf{Int}(\mathcal{C})$ composition in Figure 3 below. The bottom plane represents the analogous composition at the level of multi-points, where the composition coincides more specifically with $(g_+ \circ f_+, f_- \circ g_-)$ by generalized yanking, and is illustrated separately in Figure 4 above.

Figures 3 and 4 are the upper and lower surfaces of a three-dimensional diagram pictured in Figure 5 below. Consider the two central cubes in Figure 5. The top and bottom squares of these cubes compose because of Figures 3 and 4. The main question is the composition of the two left and, respectively, the two right vertical faces of the two central cubes. The left vertical two squares obviously compose to form a commutative square. The right (rearmost) vertical two squares in the cubes compose to be a commutative square because of generalized yanking in Appendix C. □

Obviously, there is a forgetful functor $|\ \ | : \mathsf{Int}_P(\mathcal{C}) \longrightarrow \mathsf{Int}(\mathcal{C})$.

Definition 4.9 (functor $\downarrow$). The functor $\downarrow : \mathsf{Int}(\mathcal{C}) \longrightarrow \mathsf{Int}_P(\mathcal{C})$ is defined as follows:

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press





Fig. 5. Composition $(g, g_+, g_-) \circ (f, f_+, f_-)$ in $\mathsf{Int}_P(\mathcal{C})$ (the top plane is from Figure 3 and the bottom plane is from Figure 4. The rightmost feedback arrow on the lower level maps 1^{n^-} (below the upper B^-) to 1^{n^-}.

- On objects: $\downarrow(A^+, A^-) := ((A^+ \otimes 1)_1, (A^- \otimes 1)_1)$, where $(A^\star \otimes 1)_1$ denotes *adjoining the point* $1 \cong I \otimes 1 \overset{0_{I,A^\star} \otimes 1}{\longrightarrow} A^\star \otimes 1$, $\star \in \{+, -\}$.
- On morphisms: for $f : (A^+, A^-) \longrightarrow (B^+, B^-)$, define

$$\downarrow f : ((A^+ \otimes 1)_1, (A^- \otimes 1)_1) \longrightarrow ((B^+ \otimes 1)_1, (B^- \otimes 1)_1)$$

as $\downarrow f := (s_{1,B^+} \otimes A^- \otimes 1) \circ (1 \otimes f \otimes 1) \circ (s_{A^+,1} \otimes B^- \otimes 1)$ and f_+ and f_- are Id's on 1. Diagrammatically,



Proposition 4.10 (Adjunction).
$\downarrow$ *is right adjoint to the forgetful functor* $|\ \ |$, *i.e.*

$$\mathsf{Int}(\mathcal{C})((A^+, A^-), (B^+, B^-)) \cong \mathsf{Int}_P(\mathcal{C})((A^+_{\alpha^+}, A^-_{\alpha^-}), ((B^+ \otimes 1)_1, (B^- \otimes 1)_1)). \tag{20}$$

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

Proof. Note first Equation (20) gives the required adjunction, because $(A^+, A^-) = |(A^+_{\alpha^+}, A^-_{\alpha^-})|$ and $\downarrow(B^+, B^-) = ((B^+ \otimes 1)_1, (B^- \otimes 1)_1)$. The natural bijection in Equation (20) is given by the following:


— **(From right to left)**:
Define $\epsilon_{(B^+,B^-)} : |\downarrow(B^+, B^-)| = (B^+ \otimes 1, B^- \otimes 1) \longrightarrow (B^+, B^-)$ by

$$\epsilon_{(B^+,B^-)} : B^+ \otimes 1 \otimes B^- \xrightarrow{B^+ \otimes 0_1 \otimes B^-} B^+ \otimes 1 \otimes B^- \xrightarrow{B^+ \otimes s_{1,B^-}} B^+ \otimes B^- \otimes 1$$


modulo canonical associativity isomorphisms of $\otimes$. Then, ϵ is the counit of the adjunction. Given a morphism g of the R.H.S, we obtain a morphism of the L.H.S. by composing with $\epsilon_{(B^+,B^-)}$. That is, by the composition of $\mathsf{Int}(\mathcal{C})$,

$$\begin{aligned}
&\epsilon_{(B^+,B^-)} \circ |g| := \\
&\mathsf{Tr}^{B^- \otimes 1}_{A^+ \otimes B^-, B^+ \otimes A^-} \left(\begin{array}{l} (B^+ \otimes s_{B^- \otimes 1, A^-}) \circ (\epsilon_{(B^+,B^-)} \otimes A^-) \circ \\ \qquad (B^+ \otimes 1 \otimes s_{A^-,B^-}) \circ (|g| \otimes B^-) \circ (A^+ \otimes s_{B^-, B^- \otimes 1}) \end{array} \right) \\
&= \mathsf{Tr}^{B^- \otimes 1}_{A^+ \otimes B^-, B^+ \otimes A^-} \left(((B^+ \otimes s_{1,A^-}) \circ (B^+ \otimes A^- \otimes 0_1) \circ |g|) \otimes B^- \right) \\
&\overset{\text{vanishing}}{=} \mathsf{Tr}^{1}_{A^+ \otimes B^-, B^+ \otimes A^-} \left((B^+ \otimes s_{1,A^-}) \circ (B^+ \otimes 0_1 \otimes A^-) \circ |g| \right). \qquad (21)
\end{aligned}$$

— **(From left to right):** This is the part where certain commutativity conditions will be used (to compare the two layers).
Given a morphism $f : (A^+, A^-) \longrightarrow (B^+, B^-)$ in $\mathsf{Int}(\mathcal{C})((A^+, A^-), (B^+, B^-))$, i.e.



$f' : (A^+_{\alpha^+}, A^-_{\alpha^-}) \longrightarrow \downarrow(B^+, B^-) := ((B^+ \otimes 1)_1, (B^- \otimes 1)_1)$ is constructed by the following diagram:



Note that in the diagram the domain 1 of α^- is hidden, situated behind A^+.
In the diagram,enlargethispage2pt the upper (outer) square denotes the morphism f' being constructed and the parallelogram (inside the square) with the vertices $A^+ \otimes 1^{m^+}, B^- \otimes 1, A^- \otimes 1^{m^-}, B^+ \otimes 1$ denote the following morphism:
with $g_{m^+} : 1^{m^+} \longrightarrow 1$ and $h_{m^-} : 1 \longrightarrow 1^{m^-}$ of Equation (10):

$$\begin{aligned}
&(s_{1,B^+} \otimes A^- \otimes 1^{m^-}) \circ (g_{m^+} \otimes f \otimes h_{m^-}) \circ (s_{A^+,1^{m^+}} \otimes B^- \otimes 1) : \\
&\qquad (A^+ \otimes 1^{m^+}) \otimes (B^- \otimes 1) \longrightarrow (B^+ \otimes 1) \otimes (A^- \otimes 1^{m^-}). \qquad (22)
\end{aligned}$$

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

The vertical square on the right front face and the left rear face are lifting properties (Axiom 6‴) over α^+ and α^-, respectively. Hence, as the upper surface of the diagram depicts,

$$\begin{aligned} f' :=\ & (B^+ \otimes 1 \otimes \mathfrak{e}_{\alpha^-}) \circ (22) \circ (\mathfrak{m}_{\alpha^+} \otimes B^- \otimes 1), \\ =\ & (s_{1,B^+} \otimes \mathfrak{e}_{\alpha^-}) \circ (g_{m^+} \otimes f \otimes h_{m^-}) \circ (\mathfrak{m}_{\alpha^+} \otimes B^- \otimes 1). \end{aligned} \tag{23}$$

In the bottom surface, f'_+ and f'_- are constructed by composing the bottom arrows in the diagram. The right and left faces are shown to be commutative by virtue of the fact that the two morphisms $g_{2m} \circ \mathfrak{m}^m$ and $\mathfrak{e}^m \circ h_{2m}$ give the retraction structure $1^m \rhd 1$.

Finally, we show that when the f' of Equation (23) is applied to the above 'from right to left' construction, then the original f is recovered.

$$\begin{aligned} & \epsilon_{(B^+,B^-)} \circ f' \overset{(21)}{=} \mathsf{Tr}^1_{A^+ \otimes B^-, B^+ \otimes A^-} \left((B^+ \otimes s_{1,A^-}) \circ (B^+ \otimes 0_1 \otimes A^-) \circ f' \right) \\ =\ & \text{modulo symmetry} \\ & \mathsf{Tr}^1_{A^+ \otimes B^-, B^+ \otimes A^-} \left(\begin{array}{r} (B^+ \otimes \mathfrak{e}_{\alpha^-} \otimes 1) \circ (B^+ \otimes A^- \otimes s_{1,1^{m^-}}) \circ (s_{1,B^+ \otimes A^-} \otimes 1^{m^-}) \circ \\ (0_1 \circ g_{m^+} \otimes f \otimes h_{m^-}) \circ (\mathfrak{m}_{\alpha^+} \otimes B^- \otimes 1) \end{array} \right) \\ =\ & \text{naturalities} \\ & (B^+ \otimes \mathfrak{e}_{\alpha^-}) \circ \\ & \mathsf{Tr}^1_{1^{m^+} \otimes A^+ \otimes B^-, B^+ \otimes A^- \otimes 1^{m^-}} \left(\begin{array}{r} (B^+ \otimes A^- \otimes s_{1,1^{m^-}}) \circ (s_{1,B^+ \otimes A^-} \otimes 1^{m^-}) \circ \\ (0_{1^{m^+},1} \otimes f \otimes h_{m^-}) \end{array} \right) \\ & \qquad \circ (\mathfrak{m}_{\alpha^+} \otimes B^-), \end{aligned}$$

whose Tr part is

$$\begin{aligned} & \mathsf{Tr}^1_{1^{m^+} \otimes A^+ \otimes B^-, B^+ \otimes A^- \otimes 1^{m^-}} \left(\begin{array}{r} (B^+ \otimes A^- \otimes s_{1,1^{m^-}}) \circ (f \otimes 0_{1^{m^+},1} \otimes h_{m^-}) \circ \\ (s_{1^{m^+},A^+ \otimes B^-} \otimes 1) \end{array} \right) \\ =\ & \text{naturality} \\ & \mathsf{Tr}^1_{A^+ \otimes B^- \otimes 1^{m^+}, B^+ \otimes A^- \otimes 1^{m^-}} \left((B^+ \otimes A^- \otimes s_{1,1^{m^-}}) \circ (f \otimes 0_{1^{m^+},1} \otimes h_{m^-}) \right) \circ s_{1^{m^+},A^+ \otimes B^-} \\ =\ & \text{superposing} \\ & f \otimes \mathsf{Tr}^1_{1^{m^+},1^{m^-}} \left(s_{1,1^{m^-}} \circ (0_{1^{m^+},1} \otimes h_{m^-}) \right) \circ s_{1^{m^+},A^+ \otimes B^-} \\ =\ & \text{generalized yanking} \\ & f \otimes ((h_{m^-} \circ 0_{1^{m^+},1}) \circ s_{1^{m^+},A^+ \otimes B^-}) \quad = \quad f \otimes (0_{1^{m^+},1^{m^-}} \circ s_{1^{m^+},A^+ \otimes B^-}) \text{zero absorbing} \end{aligned}$$

We conclude $\epsilon_{(B^+,B^-)} \circ f' = (B^+ \otimes \mathfrak{e}_{\alpha^-}) \circ (f \otimes 0_{1^{m^+},1^{m^-}}) \circ s_{1^{m^+},A^+ \otimes B^-} \circ (\mathfrak{m}_{\alpha^+} \otimes B^-) = f$, by Axiom 9″. □


Recall that the duality $(\ )^\perp$ of $\mathsf{Int}(\mathcal{C})$ is a contravariant endofunctor such that $(A^+, A^-)^\perp := (A^-, A^+)$ and $f^\perp := s_{B^+,A^-} \circ f \circ s_{B^-,A^+}$ for $f : (A^+, A^-) \longrightarrow (B^+, B^-)$. This duality $(\ )^\perp$ acts on $\mathsf{Int}_P(\mathcal{C})$ to yield the following dual category $\mathsf{Int}_N(\mathcal{C})$.

Definition 4.11 (Negative category $\mathsf{Int}_N(\mathcal{C})$). The negative category $\mathsf{Int}_N(\mathcal{C})$ consists of the following data:


— Objects: those of $\mathsf{Int}_P(\mathcal{C})$.

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

— Morphisms: those of $\mathsf{Int}_P(\mathcal{C})$ but the last condition on f is replaced by



Diagrammatically, a morphism (f, f_+, f_-) of $\mathsf{Int}_N(\mathcal{C})$ is described as follows:




Note that the 0 morphism occurring in the bottom level is antidiagonal to that of the 0 morphism of $\mathsf{Int}_P(\mathcal{C})$.

Hence, the positive and the negative categories are contravariantly equivalent. The functor $\uparrow : \mathsf{Int}(\mathcal{C}) \longrightarrow \mathsf{Int}_P(\mathcal{C})$ is defined by de Morgan duality $\uparrow(\ ) := (\downarrow(\ \ )^{\perp})^{\perp}$.

Thus, we obtain a compact polarized category (Definition 4.4), in the style of Remark 4.3:

Theorem 4.12 (A compact polarized category).
$(\langle \mathsf{Int}_P(\mathcal{C}), \mathsf{Int}_N(\mathcal{C})\rangle, \widehat{\mathsf{Int}(\mathcal{C})})$ *is a polarized category such that* $\downarrow$ *(resp.* $\uparrow$*) is right (resp. left) adjoint to the forgetful functor* $|\ \ |$. *The polarized category is compact so that* $(\mathsf{Int}_P(\mathcal{C}))^{op} = \mathsf{Int}_N(\mathcal{C})$. *In diagrammatic form,*



5. A polarized Int construction using Rel with multi-points

In this section, we show how to build a concrete instance of the previous polarized Int-construction using Rel with multi-points to construct the associated commutativity

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

conditions compatible with these multi-points. Thus, we obtain a concrete compact polarized model of MLLP.

We make the following observations. First, this section is a relational instance of Section 4 and can be read independently of it. Second, in Sections 2 and 3 of this paper, we often use the matrix formalism of Haghverdi's UDC's (see Appendix B). This agrees with the usual matrix calculus in linear algebra. In what follows, we adopt instead the matrix notation of Joyal *et al.* (1996) for Int(Rel), since these authors do similar calculations to those below. We introduce the following standard notions, *cf.* Abramsky *et al.* (2002) and Hamano and Scott (2007).

Notation 5.1.

— For a relation $R : A \to B$, and subsets $X \subseteq A$ and $Y \subseteq B$,

$$[Y]R := \{x \mid \exists b \in Y.(x,b) \in R\} \subseteq A \qquad R[X] := \{y \mid \exists a \in X.(a,y) \in R\} \subseteq B$$

We write R^* for the smallest reflexive and transitive relation containing R.

— (The category Rel)
Rel denotes the category of sets and relations. Relational composition of $R : A \longrightarrow B$ and $S : B \longrightarrow C$ is written from right to left. We write $SR : A \longrightarrow C$ where $SR = \{(x,z) \in A \times C \mid \exists y \in B.\ (x,y) \in R \text{ and } (y,z) \in S\}$ and omit the $\circ$ symbol. Rel becomes monoidal with disjoint union $A + B$ of sets as the tensor product. The empty set $\varnothing$ is the tensor unit.

— A morphism $R : (A^+, A^-) \to (B^+, B^-)$ of Int(Rel) is represented by the following matrix (where the border objects represent appropriate domains and codomains of the entries, as shown below):

$$\begin{array}{c} \\ B^+ \\ A^- \end{array} \begin{array}{c} \begin{array}{cc} A^+ & B^- \end{array} \\ \begin{pmatrix} R_{12} & R_{22} \\ R_{11} & R_{21} \end{pmatrix} \end{array}$$

The entries are relations $R_{11} : A^+ \to A^-$, $R_{12} : A^+ \to B^+$, $R_{21} : B^- \to A^-$, $R_{22} : B^- \to B^+$.

E.g. $\mathrm{Id}_{(A^+,A^-)}$ is represented by $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$, where 1 denotes the singleton $\{*\}$.

— The dual morphism $R^\perp : (B^-, B^+) \to (A^-, A^+)$ in Int(Rel) is represented by


$$R^\perp := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} R \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} R_{21} & R_{11} \\ R_{22} & R_{12} \end{pmatrix}$$

Fact 5.2 (Composition in Int(Rel)). Given morphisms $R : (A^+, A^-) \to (B^+, B^-)$ and $S : (B^+, B^-) \to (C^+, C^-)$ of Int(Rel) represented by $\begin{array}{c} \\ B^+ \\ A^- \end{array} \begin{array}{c} \begin{array}{cc} A^+ & B^- \end{array} \\ \begin{pmatrix} R_{12} & R_{22} \\ R_{11} & R_{21} \end{pmatrix} \end{array}$ and $\begin{array}{c} \\ C^+ \\ B^- \end{array} \begin{array}{c} \begin{array}{cc} B^+ & C^- \end{array} \\ \begin{pmatrix} S_{12} & S_{22} \\ S_{11} & S_{21} \end{pmatrix} \end{array}$, the composition $S \circ R$ in Int(Rel) is given by the following relation:

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

$$\begin{array}{c} \\ C^+ \\ A^- \end{array}\!\!\begin{array}{c} \begin{array}{cc} A^+ & C^- \end{array} \\ \begin{pmatrix} \varnothing & S_{22} \\ R_{11} & \varnothing \end{pmatrix} \end{array} \cup \begin{array}{c} \\ C^+ \\ A^- \end{array}\!\!\begin{array}{c} \begin{array}{cc} A^+ & C^- \end{array} \\ \begin{pmatrix} S_{12}(R_{22}S_{11})^*R_{12} & S_{12}R_{22}(S_{11}R_{22})^*S_{21} \\ R_{21}S_{11}(R_{22}S_{11})^*R_{12} & R_{21}(S_{11}R_{22})^*S_{21} \end{pmatrix} \end{array}$$

which we can write as



.

In general in Rel, a multi-point $1^m \longrightarrow A$ is an m-indexed family of subsets of A (cf. Example 3.7). In what follows, we denote a multi-point of a set A by $\mathsf{mp}(A)$ and think of it as a subset $\mathsf{mp}(A) \subseteq A$.


Definition 5.3 (Positive category Pos).

— Objects are $(A^+_{\mathsf{mp}(A^+)}, A^-_{\mathsf{mp}(A^-)})$ where (A^+, A^-) is an object of Int(Rel) and $\mathsf{mp}(A^+) \subseteq A^+$ and $\mathsf{mp}(A^-) \subseteq A^-$ are multi-points, respectively, of A^+ and A^-.

— Arrows (also called *positive maps*) are morphisms $R : (A^+, A^-) \to (B^+, B^-)$ in Int(Rel) satisfying the following three conditions:

1. $[\mathsf{mp}(B^+)]R_{12} = \mathsf{mp}(A^+)$.
2. $R_{21}[\mathsf{mp}(B^-)] = \mathsf{mp}(A^-)$.
3. $[\mathsf{mp}(B^+)]R_{22} = \varnothing = R_{22}[\mathsf{mp}(B^-)]$.

The above three conditions are instances of the three conditions on morphisms of Definition 4.7. The following figure explains how the conditions (1) and (2) correspond to the left and right vertical squares, respectively, of that definition, and that the condition (3) corresponds to the diagonal 0 morphism.



Proposition 5.4. Pos *forms a category.*

Proof. Id's are Pos maps since Id on (A^+, A^-) is given by the matrix s.t. $\mathrm{Id}_{12} = \mathrm{Id}_{A^+}$, $\mathrm{id}_{21} = \mathrm{Id}_{A^-}$ and $\mathrm{Id}_{11} = \mathrm{Id}_{22} = \varnothing$. We check that the composition of Int(Rel) preserves Pos maps. The computation below is essentially a concrete instance of Figure 5.

Given two morphisms $R : (A^+, A^-) \to (B^+, B^-)$ and $S : (B^+, B^-) \to (C^+, C^-)$ of Int(Rel).

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

$$
\begin{aligned}
1.\ [\mathsf{mp}(C^+)](SR)_{12} &= [\mathsf{mp}(C^+)](S_{12}(R_{22}S_{11})^*R_{12}) \\
&= [\mathsf{mp}(B^+)]((R_{22}S_{11})^*R_{12}) \\
&= [\mathsf{mp}(B^+)]R_{12} \cup [\mathsf{mp}(B^+)]((R_{22}S_{11})(R_{22}S_{11})^*R_{12}) \\
&= \mathsf{mp}(A^+) \cup \varnothing \\
2.\ (SR)_{21}[\mathsf{mp}(C^-)] &= (R_{21}(S_{11}R_{22})^*S_{21})[\mathsf{mp}(C^-)] \\
&= (R_{21}(S_{11}R_{22})^*)[\mathsf{mp}(B^-)] \\
&= R_{21}[\mathsf{mp}(B^-)] \cup (R_{21}(S_{11}R_{22})^*S_{11})(R_{22}[\mathsf{mp}(B^-)]) \\
&= \mathsf{mp}(A^-) \cup \varnothing \\
3.\ [\mathsf{mp}(C^+)](SR)_{22} &= [\mathsf{mp}(C^+)]S_{22} \cup [\mathsf{mp}(C^+)](S_{12}R_{22}(S_{11}R_{22})^*S_{21}) \\
&= \varnothing \cup [\mathsf{mp}(B^+)](R_{22}(S_{11}R_{22})^*S_{21}) \\
&= \varnothing \cup \varnothing \\
(SR)_{22}[\mathsf{mp}(C^-)] &= S_{22}[\mathsf{mp}(C^-)] \cup (S_{12}R_{22}(S_{11}R_{22})^*S_{21})[\mathsf{mp}(C^-)] \\
&= \varnothing \cup (S_{12}R_{22}(S_{11}R_{22})^*)[\mathsf{mp}(B^-)] \\
&= \varnothing \cup (S_{12}R_{22})[\mathsf{mp}(B^-)] \cup (S_{12}R_{22}(S_{11}R_{22})^*S_{11}R_{22})[\mathsf{mp}(B^-)] \\
&= \varnothing \cup \varnothing \cup \varnothing
\end{aligned}
$$

□

Proposition 5.5. *The category* Pos *is monoidal with respect to the tensor product of* Int(Rel).

Proof. Given Pos maps $R : (A^+_{\mathsf{mp}(A^+)}, A^-_{\mathsf{mp}(A^-)}) \to (B^+_{\mathsf{mp}(B^+)}, B^-_{\mathsf{mp}(B^-)})$ and $S : (C^+_{\mathsf{mp}(C^+)}, C^-_{\mathsf{mp}(C^-)}) \to (D^+_{\mathsf{mp}(D^+)}, D^-_{\mathsf{mp}(D^-)})$, the tensor product

$$
\begin{aligned}
R \otimes S : ((A^+ + C^+)_{\mathsf{mp}(A^+)+\mathsf{mp}(C^+)}, (A^- + C^-)_{\mathsf{mp}(A^-)+\mathsf{mp}(C^-)}) \longrightarrow \\
((B^+ + D^+)_{\mathsf{mp}(B^+)+\mathsf{mp}(D^+)}, (B^- + D^-)_{\mathsf{mp}(B^-)+\mathsf{mp}(D^-)})
\end{aligned}
$$

is given by

$$
\begin{array}{c} \\ D^+ \\ B^+ \\ C^- \\ A^+ \end{array}
\begin{array}{c}
\begin{array}{cccc} A^+ & C^+ & B^- & D^- \end{array} \\
\begin{pmatrix} \varnothing & S_{12} & \varnothing & S_{22} \\ R_{12} & \varnothing & R_{22} & \varnothing \\ \varnothing & S_{11} & \varnothing & S_{21} \\ R_{11} & \varnothing & R_{21} & \varnothing \end{pmatrix}
\end{array}
$$


It is straightforward that $\otimes$ preserves positivity. □

Dually in Int(Rel), negative maps are defined so that they form a monoidal category Neg.

Definition 5.6 (Negative category Neg). The objects of Neg are the same as those of Pos and the morphisms (*negative maps*) are dual; that is, they satisfy the following three conditions:

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

1. $R_{12}[\mathsf{mp}(A^+)] = \mathsf{mp}(B^+)$
2. $[\mathsf{mp}(A^-)]R_{21} = \mathsf{mp}(B^-)$
3. $[\mathsf{mp}(A^-)]R_{11} = \varnothing = R_{11}[\mathsf{mp}(A^+)]$



Proposition 5.7 (Negative category). Neg *forms a monoidal category.*

Remark 5.8 (Positive ≠ negative). Pos maps and Neg maps are different, so the categories are different and the model is in this sense non-degenerate.

The following functor ↓ is a special instance of the previously defined functor in Definition 4.9 and dually for ↑.


Definition 5.9 (Functors ↓ and ↑).
Functors $\downarrow\colon \mathsf{Int}(\mathsf{Rel}) \longrightarrow \mathsf{Pos}$, $\uparrow\colon \mathsf{Int}(\mathsf{Rel}) \longrightarrow \mathsf{Neg}$ are defined as follows:

— On objects: $\downarrow(A^+, A^-) := \uparrow(A^+, A^-) := ((A^+ + 1)_1, (A^- + 1)_1)$.
— On morphisms: For $R : (A^+, A^-) \to (B^+, B^-)$, $\downarrow R$ and $\uparrow R$ are defined by

$$\downarrow R := \uparrow R := \begin{array}{c} \\ 1 \\ B^+ \\ A^- \\ 1 \end{array} \begin{array}{c} \begin{array}{cccc} 1 & A^+ & B^- & 1 \end{array} \\ \begin{pmatrix} (\star,\star) & \varnothing & \varnothing & \varnothing \\ \varnothing & R_{12} & R_{22} & \varnothing \\ \varnothing & R_{11} & R_{21} & \varnothing \\ \varnothing & \varnothing & \varnothing & (\star,\star) \end{pmatrix} \end{array}.$$

Note that the functors ↑ and ↓ are not full since Pos ≠ Neg.

Proposition 5.10 (Adjunctions).
↓ *(resp.* ↑*) is right (resp. left) adjoint to the forgetful functor* | |*:*



Proof. We show the following:

$$\begin{aligned} \mathsf{Int}(\mathsf{Rel})((A^+, A^-), (B^+, B^-)) &\cong \mathsf{Pos}((A^+_{\mathsf{mp}(A^+)}, A^-_{\mathsf{mp}(A^-)}), ((B^+ + 1)_1, (B^- + 1)_1)) \\ &\cong \mathsf{Neg}(((A^+ + 1)_1, (A^- + 1)_1), (B^+_{\mathsf{mp}(B^+)}, B^-_{\mathsf{mp}(B^-)})), \end{aligned}$$

where $((B^+ + 1)_1, (B^- + 1)_1) = \downarrow(B^+, B^-)$ and $((A^+ + 1)_1, (A^- + 1)_1) = \uparrow(A^+, A^-)$.

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

Every positive map R' from the R.H.S is of the form

$$\begin{array}{c} \\ 1 \\ B^+ \\ A^- \end{array} \begin{array}{c} \begin{array}{ccc} A^+ & B^- & 1 \end{array} \\ \begin{pmatrix} \mathsf{mp}(A^+) \times 1 & \varnothing & \varnothing \\ R_{12} & R_{22} & \varnothing \\ R_{11} & R_{21} & 1 \times \mathsf{mp}(A^-) \end{pmatrix} \end{array}$$

so that $R' = R \cup \mathsf{mp}(A^+) \times 1 \cup 1 \times \mathsf{mp}(A^-)$ with R from the L.H.S. This gives a natural bijective correspondence. □

6. Conclusion and future work

In this paper, we presented two independent studies of GoI for multiplicative polarized linear logic (MLLP): one based on the notion of GoI situations (Abramsky *et al.* 2002) and the other based on a direct application of Joyal–Street–Verity's Int construction (Joyal *et al.* 1996). Both modellings use the idea of adjoining multi-points to account for polarities, hence focusing. In polarized GoI situations, preservation of multi-points via the execution formula allows us to characterize focusing semantically. In the case of the Int construction, the goal was instead to construct compact polarized denotational models. This involved adding multi-points to the Int construction so as to be compatible with those commutativity conditions previously discussed.

Finally, in the last section, we construct a concrete example of such a polarized category, based on the Int construction applied to the category of multi-pointed relations. For future studies, we leave open the following questions:

1. What is the logical status of multi-points? For example, multi-points have no counterpart in the syntax: they are an additional structure added to a non-polarized (although somewhat 'degenerate') compact closed model. Yet, multi-points allow us to characterize syntactic questions of polarized logic, for example, characterizing focusing.

2. One question of interest is how Sections 3 and 4 of this paper are related. We note that our main theorem characterizing focusing (Theorem 3.26) involves commutative squares which can also be shown to be weak pullbacks. Thus, weak pullbacks arise from the termination of the execution formula given by traces. In Section 4, we start from weak pullbacks in the definition of morphisms in $\mathsf{Int}_P(C)$ (below Definition 4.7), where we see the property that the two-side vertical faces are actually weak pullbacks. The main results of Section 4 show these weak pullbacks Lambek and Scott (1986); Mac Lane (1971) are preserved not only under composition but more strongly under tracing. We have used the fact that the squares that arise in both sections are analogous. We hope to give a categorical characterization of such analogies.

3. This paper is restricted to the multiplicative fragment. It would be interesting to extend this to the full MALLP level, which is the language of Girard's Ludics, Girard (2001) as studied in our paper (Hamano and Scott 2007). This seems more promising compared to non-polarized additive models because the additive connectives are less complicated in the polarized setting, as mentioned in the work of O. Laurent (for example, Laurent 2002). This future work may relate our work to Laurent's GoI model for additives (Laurent 2001).

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

Acknowledgements

We wish to thank the referee for detailed and very helpful comments that have greatly improved the presentation. Masahiro Hamano's Research was supported by a PRESTO grant from JST. Philip Scott's Research was supported in part by an NSERC Discovery Grant.

Appendix A. GoI Situations and Execution Formulas for MELL

In Abramsky *et al.* (2002), the authors introduced a general algebraic framework for studying Girard's GoI (Girard 1989, 1995) for multiplicative-exponential linear logic, MELL. This framework, called a *GoI Situation*, contains an underlying TMC $\mathcal{C}$, along with a reflexive object U and an endofunctor T used to represent the exponentials of linear logic. It is shown in that paper how to interpret GoI as yielding linear combinatory algebras on the endomorphism monoids $\mathrm{End}_{\mathcal{C}}(U)$.

In later work (Haghverdi and Scott 2006, 2011), the underlying traced category $\mathcal{C}$ of a GoI situation was specialized to a traced UDC (e.g. Rel, Pfn , PInj) (see Appendix B below) which is equipped with a standard *particle-style* trace, as developed in Haghverdi's thesis (Haghverdi 2000), together with an abstract categorical execution formula given in terms of that trace (see below). The categorical GoI interpretation of Haghverdi–Scott (which captures Girard's GoI 1 (Girard 1989)) contains three components: (i) an interpretation of proofs in $\mathrm{End}_{\mathcal{C}}(U)$, (ii) an interpretation of formulas as types (= bi-orthogonally closed subsets of $\mathrm{End}_{\mathcal{C}}(U)$, with respect to an appropriate Girard–Hyland–Schalk orthogonality) Hyland and Schalk (2003), and (iii) an analysis of the dynamics of cut elimination via the execution formula. For the polarized system MLLP studied in this paper, we only discuss (i) and (iii).

Definition A.1. *A GoI Situation is a triple* $(\mathcal{C}, T, U)$*, where*

(1) $\mathcal{C}$ *is a traced symmetric monoidal category and* $T : \mathcal{C} \longrightarrow \mathcal{C}$ *is a traced symmetric monoidal functor with the following monoidal retractions (i.e. the retraction pairs are monoidal natural transformations):*

(i) $e' : T \rhd TT : e$ *(comultiplication)*

(ii) $d' : T \rhd \mathrm{Id} : d$ *(dereliction)*


(iii) $c' : T \rhd T \otimes T : c$ *(contraction)*

(iv) $w' : T \rhd \mathcal{K}_I : w$ *(weakening). Here,* $\mathcal{K}_I$ *is the constant* I *functor.*

(2) U *is an object of* $\mathcal{C}$*, called a reflexive object, with retractions:*
(a) $k : U \rhd U \otimes U : j$ *(b)* $U \rhd I$ *, and (c)* $v : U \rhd TU : u$.

Here, $e' : T \rhd TT : e$ means that $e_X : TTX \longrightarrow TX$ and $e'_X : TX \longrightarrow TTX$ are monoidal natural transformations such that $e'e = \mathrm{Id}_{TT}$. We say that TT *is a (monoidal) retract of* T. Similarly for the other items. Following the presentation in Haghverdi and Scott (2006, 2011), given a GoI situation $(\mathcal{C}, T, U)$, the GoI interpretation of a proof π of an MELL sequent (with explicit cuts) $\vdash [\Delta], \Gamma$ (where Δ denotes the set of all pairs of cut formulas $A, A^{\perp}$ used in π) is determined by a pair of morphisms $(\llbracket \pi \rrbracket, \sigma)$ as in Figure 6,

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press



$(\llbracket \pi \rrbracket, \sigma)$ $\qquad$ $Ex(\llbracket \pi \rrbracket, \sigma) = \mathbf{Tr}^{\otimes\Delta}_{\otimes\Gamma,\otimes\Gamma}((1_{\otimes\Gamma} \otimes \sigma)\llbracket \pi \rrbracket)$

Fig. 6. Proofs of $\vdash [\Delta], \Gamma$ as I/O boxes and the execution formula.

where σ represents the cuts Δ. If $|\Delta| = 2m$ and $|\Gamma| = n$, these data are given by $\mathcal{C}$-arrows, $\sigma : U^{2m} \to U^{2m}$, $\llbracket \pi \rrbracket : U^{n+2m} \to U^{n+2m}$. Finally, Girard's *execution formula* determines an arrow $\mathsf{Ex}(\llbracket \pi \rrbracket, \sigma) : U^n \to U^n$, where $U^k = U \otimes \cdots \otimes U$ (k times). If $\mathcal{C}$ is a Haghverdi traced UDC with a standard (particle-style) trace (as in the Rel-based models in this paper: see Appendix B below), we can write the execution Formula in the more familiar form:

$$\mathsf{Ex}(\llbracket \pi \rrbracket, \sigma) \;=\; \pi_{11} + \sum_{n \geqslant 0} \pi_{12}(\sigma\pi_{22})^n(\sigma\pi_{21}), \tag{24}$$

where $[\pi_{ij}]$ is the matrix representation of $\llbracket \pi \rrbracket$; this was shown in Haghverdi and Scott (2006) to agree with Girard's original execution formula (Girard 1989) in his model Hilb_2 ($= \ell_2[\mathsf{PInj}]$). Such UDC models also support a robust matrix calculus to represent morphisms, which agrees with the usual matrix representation of relations used in this paper (see Proposition B.1 below).

Appendix B. Unique Decomposition Categories (UDCs)

E. Haghverdi, in his thesis (Haghverdi 2000), introduced UDCs. These were specifically developed for modelling 'particle-style' GoI as in GoI 1 (Girard 1989; Haghverdi and Scott 2006).

Briefly, UDCs are symmetric monoidal categories with the following additional structure:

— The homsets are enriched with a Σ-monoid additive structure, such that composition distributes over addition, both from the left and the right. For the precise Σ-monoid axioms, we refer to Haghverdi's thesis, Chapter 4. In particular, there are zero morphisms $0_{XY} : X \to Y$ between any two objects X, Y.

— For a finite set I and for each $j \in I$, there are *quasi injections* $\iota_j : X_j \to \otimes_I X_i$, and *quasi projections* $\rho_j : \otimes_I X_i \to X_j$ such that

(i) $\rho_k \iota_j = \mathrm{Id}_{X_j}$ if $j = k$ and $0_{X_j X_k}$ otherwise;
(ii) $\sum_{i \in I} \iota_i \rho_i = \mathrm{Id}_{\otimes_I X_i}$.

Examples of UDC's (for GoI) include variations of Rel_+, for example, the categories Pfn and PInj of *partial functions* (resp. *partial injective functions*).

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

The main theorem on UDC's, which is used in various places in this paper, is the representation of morphisms as matrices, with an associated full matrix calculus for computations. This can be summarized as follows (see Haghverdi 2000, Proposition 4.0.6):


Proposition B.1 (Matricial representation). *Given a morphism* $f : \otimes_J X_j \to \otimes_I Y_i$ *in a UDC, with* $|I| = m, |J| = n$, *there exists a unique family* $\{f_{ij}\}_{i\in I, j\in J} : X_j \to Y_i$ *with* $f = \sum_{i\in I, j\in J} \iota_i f_{ij} \rho_j$, *where* $f_{ij} = \rho_i f \iota_j$. *Moreover, composition of morphisms in a UDC corresponds to matrix multiplication of their associated matrices.*

Appendix C. Generalized Yanking for a Traced Monoidal Category

The following identity is frequently used in calculating traces (see Proposition 2.4, in Abramsky *et al.* 2002),

$$\mathsf{Tr}^U_{X,Y}(s\circ(f \otimes g)) = g\circ f$$

Pictorially, this says




Appendix D. Omitted Proofs

Proof of Proposition 3.8. By induction on the construction of a multi-point $\boldsymbol{p}$.
(Base Case): This is when $\boldsymbol{p}$ is the distinguished point α, hence the property is the original lifting property 6′.
(Induction Case): $\boldsymbol{p}$ is constructed either by 2 or 3 of Definition 3.5 (Note in 2, the $\boldsymbol{p}$ is a point.): We define

$\boldsymbol{p}$ constructed by 3	$\boldsymbol{p}$ constructed by 2
$\mathfrak{m}_{\boldsymbol{p}} := ((j_m \circ \tau) \otimes 1^m) \circ \bigotimes_i \mathfrak{m}_{\boldsymbol{p}_i} \circ \tau^- \circ k_m$	$\mathfrak{m}_{\boldsymbol{p}} := (j \otimes 1) \circ (U \otimes \mathfrak{m}_{\boldsymbol{p}'}) \circ k$
$\mathfrak{e}_{\boldsymbol{p}} := j_m \circ \tau \circ \bigotimes_i \mathfrak{e}_{\boldsymbol{p}_i} \circ (k_m \otimes 1^m)$	$\mathfrak{e}_{\boldsymbol{p}} := j \circ (U \otimes \mathfrak{e}_{\boldsymbol{p}'}) \circ (k \otimes 1)$

Then the commutativity property follows from I.H.'s (the lower square of the following diagram) and the retractions $U \triangleright U^m$ and $U \otimes 1 \triangleright U^m \otimes 1^m$ with k either $m_1 + m_2$ or 1 by Axiom 2 (the upper right and left vertical arrows, respectively). Note especially that the lower square of the left case is the m-fold tensoring of Axiom 6:

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press



We check one commutativity for each of the constructions 2 and 3 of $\boldsymbol{p}$.

$\boldsymbol{p}$ constructed by 3	$\boldsymbol{p}$ constructed by 2	
$\mathfrak{m}_{\boldsymbol{p}} \circ j_m \circ \tau \circ \otimes \boldsymbol{p}_i \circ \mathfrak{e}^m$	$\mathfrak{m}_{\boldsymbol{p}} \circ j \circ (0 \otimes \boldsymbol{p}') \circ (I \otimes \mathfrak{e})$	
$= (j_m \otimes 1^m) \circ \otimes \mathfrak{m}_{\boldsymbol{p}_i} \circ \tau^- \circ k_m \circ j_m \circ \tau \circ \otimes \boldsymbol{p}_i \circ \mathfrak{e}^m$	$= (j \otimes 1) \circ (U \otimes \mathfrak{m}_{\boldsymbol{p}'}) \circ k \circ j \circ (0 \otimes \boldsymbol{p}') \circ (I \otimes \mathfrak{e})$	by retractions:
$= (j_m \otimes 1^m) \circ \otimes \mathfrak{m}_{\boldsymbol{p}_i} \circ \otimes \boldsymbol{p}_i \circ \mathfrak{e}^m$	$= (j \otimes 1) \circ (U \otimes \mathfrak{m}_{\boldsymbol{p}'}) \circ (0 \otimes \boldsymbol{p}') \circ (I \otimes \mathfrak{e})$	In 3: $U \rhd_{(k_m, j_m)} U^m$ In 2: $U \rhd U^2$
$= (j_m \otimes 1^m) \circ (\otimes \boldsymbol{p}_i \otimes 1^m)$	$= (j \otimes 1) \circ (0 \otimes \boldsymbol{p}' \otimes 1)$	by commutativity of lower square
$= \boldsymbol{p} \otimes 1^m$	$= \boldsymbol{p} \otimes 1$	by definition of $\boldsymbol{p}$

□

Proof of Lemma 3.10. These retractions are compatible with TMCs (Abramsky *et al.* 2002) by virtue of dinaturality and the directions of the retractions (k_m, j_m) and $(\mathfrak{e}_{\boldsymbol{p}}, \mathfrak{m}_{\boldsymbol{p}})$, respectively, as follows:

For (3);		For (4);	
$\mathsf{Tr}^U_{X,Y}((j \otimes Y) \circ f \circ (k \otimes X))$		$\mathsf{Tr}^{U\otimes 1^m}_{X,Y}((\mathfrak{m}_{\boldsymbol{p}} \otimes Y) \circ g \circ (\mathfrak{e}_{\boldsymbol{p}} \otimes X))$	
$=$	dinaturality	$=$	dinaturality
$\mathsf{Tr}^{U^m}_{X,Y}(f \circ (k \otimes X)) \circ (j \otimes X))$		$\mathsf{Tr}^U_{X,Y}(g \circ (\mathfrak{e}_{\boldsymbol{p}} \otimes X) \circ (\mathfrak{m}_{\boldsymbol{p}} \otimes X))$	
$=$		$=$	
$\mathsf{Tr}^{U^m}_{X,Y}(f \circ (k \circ j \otimes X))$		$\mathsf{Tr}^U_{X,Y}(g \circ (\mathfrak{e}_{\boldsymbol{p}} \circ \mathfrak{m}_{\boldsymbol{p}} \otimes X))$	
$=$	$k \circ j = \mathrm{Id}_{U^m}$	$=$	$\mathfrak{e}_{\boldsymbol{p}} \circ \mathfrak{m}_{\boldsymbol{p}} = \mathrm{Id}_U$
$\mathsf{Tr}^{U^m}_{X,Y}(f)$		$\mathsf{Tr}^U_{X,Y}(g)$	

See the following pictures in Figure 7, in which Tr^{U^m} is described via Vanishing II.

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press




Fig. 7. Invariance of traces under conjugate actions.

□

Appendix E. Remarks on Retractions $U \rhd_{(k,j)} U \otimes U$ and $U \otimes 1 \rhd_{(\mathfrak{e}_z, \mathfrak{m}_z)} U$

The reader may wonder about the opposite directions of the retractions $U \rhd_{(k,j)} U \otimes U$ and $U \otimes 1 \rhd_{(\mathfrak{e}_z, \mathfrak{m}_z)} U$ in the two-layered GoI interpretation $[\![\pi]\!]$ and f_π.

i. (On the retraction $U \rhd_{(k,j)} U \otimes U$)
The retract $U \otimes U$ (of U) follows the form of the logical constructions. In (untyped) GoI, where there is a reflexive object U, the interpretations of $\otimes$ and $\parr$ are indistinguishable. In general, for any formula A, U_A is identified with U. So to make sense of the logical connectives, via reflexivity of U, we use the retraction. Thus, $U_A \otimes U_B$, which is $U \otimes U$ is faithfully projected to U which is defined to be both $U_{A \otimes B}$ as well as $U_{A \parr B}$. Similarly, letting $U_\downarrow = U_\uparrow = U$, we have $U_\downarrow \otimes U_A$ (resp. $U_A \otimes U_\uparrow$) is faithfully projected to U, which itself is defined to be $U_{\downarrow A}$ (resp. $U_{\uparrow A}$). The faithfulness is guaranteed by $k \circ j = \mathrm{Id}_{U \otimes U}$. Note that at the level of the U's, the dual logical connectives $\downarrow$ and $\uparrow$ are not distinguishable, just as $\otimes$ and $\parr$ are not distinguishable. However, to account for the asymmetry in logical rules for the polarities, we will need to employ a new ingredient, the object 1.

ii. On the retraction $U \otimes 1 \rhd_{(\mathfrak{e}_z, \mathfrak{m}_z)} U$
The retract 1 (of $1 \otimes 1$) is for the sake of realizing the retract U (of $U \otimes 1$) of

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

the lifting property (along α) of Definition 3.1. Then what is the meaning of the retraction $U \otimes 1 \rhd_{(\mathfrak{e}_\alpha, \mathfrak{m}_\alpha)} U$? Tensoring 1 with U in the construction of $\mathfrak{m}_\alpha : U \longrightarrow U \otimes 1$ corresponds to making the point $\alpha : 1 \longrightarrow U$ explicitly appear. Conversely, $\mathfrak{e}_\alpha : U \otimes 1 \longrightarrow U$ hides the point α (i.e. makes it implicit). The faithfulness of making the point α explicit is guaranteed by $\mathfrak{e}_\alpha \circ \mathfrak{m}_\alpha = \mathrm{Id}_U$, i.e. intuitively, making α explicit, then hiding it gives the identity.


For a multi-point $\mathsf{mp}(A) : 1^m \xrightarrow{p} U$ for any given polarized formula A so that $\mathbb{1}_A \cong 1^m$, the retraction $1 \otimes 1 \rhd_{(\mathfrak{e}, \mathfrak{m})} 1$ and its lifting $U \otimes 1 \rhd_{(\mathfrak{e}_\alpha, \mathfrak{m}_\alpha)} U$ are correspondingly generalized into the retraction $1^m \otimes 1^m \rhd_{(\mathfrak{e}^m, \mathfrak{m}^m)} 1^m$ and the lifting $U \otimes 1^m \rhd_{(\mathfrak{e}_p, \mathfrak{m}_p)} U$ of Axiom 6′ of Proposition 3.8. These retractions are in order to accommodate polarities in U and in 1^m.

iii. Note also the opposite directions of retractions between (i) and (ii) above are compatible with the conjugate actions of Lemma 3.10.

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press


Appendix F. Pictorial Proof for Proposition 3.22 (Ex is an Invariant)

L.H.S. of (11)




naturality

=



superposing


=



⋆ is L.H.S of Fig for (12) when $f = b, g = c$ and $h = a$.


(12) and $c = 0$

=



Ax 9'

=



R.H.S. of (11)

References

Abramsky, S., Haghverdi, E. and Scott, P.J. (2002). Geometry of interaction and linear combinatory algebras. *Mathematical Structures in Computer Science* **12** (5) 1–40.

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

Andreoli, J.-M. (1992). Logic programming with focusing proofs in linear logic. *Journal of Logic and Computation* **2** (3) 297–347.

Andreoli, J.-M. (2001). Focussing and proof construction. *Annals of Pure and Applied Logic* **107** (1) 131–163.

Blute, R.F. and Scott, P.J. (2004). Category theory for linear logicians. In: Ehrhard, T., Girard, J.-Y., Ruet, P., Scott, P. (eds.) *Linear Logic and Computer Science*, London Mathematical Society Lecture Note Series, vol. 316, C.U.P.

Chaudhuri, K. (2008). Focusing strategies in the sequent calculus of synthetic connectives. In: *Logic for Programming, Artificial Intelligence and Reasoning (LPAR-15)*, Doha, Qatar, Lecture Notes in Computer Science, vol. 5330, Springer, 467–481.

Chaudhuri, K., Hetzl, S. and Miller, D. A (2013). Multi-focused proof system isomorphic to expansion proofs. *Journal of Logic and Computation* **26** (2) 577–603.

Cockett, J.R.B., Hasegawa, M. and Seely, R.A.G. (2006). Coherence of the double involution on *-autonomous categories. *TAC* **17** 17–29.

Cockett, J.R.B. and Seely, R.A.G. (2007). Polarized category theory, modules and game semantics. *TAC* **18** 4–101.

Ehrhard, T. (2012). The Scott model of linear logic is the extensional collapse of its relational model. *Theoretical Computer Science* **424** (23) 20–45.

Girard, J.-Y. (1987). Linear logic. *Theoretical Computer Science* **50** 1–102.

Girard, J.-Y. (1989). Geometry of interaction I: Interpretation of system F. In: Ferro, R. *et al.* (eds.) *Logic Colloquium '88*, North-Holland, 221–260.

Girard, J.-Y. (1991). A new constructive logic: Classical logic. *Mathematical Structures in Computer* **1** (3) 255–296.

Girard, J.-Y. (1995). Geometry of interaction III: Accommodating the additives. In: *Advances in Linear Logic,* Lecture Notes in Computer Science, vol. 222, CUP, 329–389.

Girard, J.-Y. (1999). On the meaning of logical rules I: Syntax vs. semantics. In Berger, U. and Schwichtenberg, H. (eds.) *Computational Logic*, NATO ASI Series, vol. 165, Springer, 215–272.

Girard, J.-Y. (2001). Locus solum: From the rules of logic to the logic of rules. *Mathematical Structures in Computer Science* **11** 301–506.

Girard, J.-Y. (2011). *The Blind Spot: Lectures in Logic*, European Mathematical Society, 550.

Haghverdi, E. (2000). *A Categorical Approach to Linear Logic, Geometry of Proofs and Full Completeness.* PhD Thesis, University of Ottawa, Canada.

Haghverdi, E. and Scott, P. (2006). A categorical model for the geometry of interaction. *Theoretical Computer Science* **350** (2–3) 252–274.

Haghverdi, E. and Scott, P.J. (2010). Towards a typed geometry of interaction. *Mathematical Structures in Computer Science*, **20** 473–521.

Haghverdi, E. and Scott, P.J. (2011). Geometry of interaction and the dynamics of proof reduction: A tutorial, new structures in physics. In: Coecke, R. (ed.) *Springer Lectures Notes in Physics*, vol. 813, Oxford.

Hamano, M. and Scott, P. (2007). A categorical semantics for polarized MALL. *Annals of Pure & Applied Logic* **145** 276–313.

Hamano, M. and Takemura, R. (2008) An indexed system for multiplicative additive polarized linear logic. In: *Proceedings of 17th Annual Conference on Computer Science Logic (CSL'08)*, Lecture Notes in Computer Science, vol. 5213, 262–277.

Hamano, M. and Takemura, R. (2010). A phase semantics for polarized linear logic and second order conservativity. *The Journal of Symbolic Logic* **75** (1) 77–102.

Hasegawa, M. (2004). The uniformity principle on traced monoidal categories. *Publications of the Research Institute for Mathematical Sciences* **40** (3) 991–1014.

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press

Hyland, J.M.E. and Schalk, A. (2003). Glueing and orthogonality for models of linear logic. *Theoretical Computer Science* **294** 183–231.

Joyal, A., Street, R. and Verity, D. (1996). Traced monoidal categories. *Mathematical Proceedings of the Cambridge Philosophical Society* **119** 447–468.

Joyal, A. (Nov. 2011) Joyal's CatLab, Distributors and Barrels. Available at: `http://ncatlab.org/joyalscatlab/show/Distributors+and+barrels#distributors_6`

Lambek, J. and Scott, P.J. (1986). *Introduction to Higher Order Categorical Logic*, Cambridge Studies in Advanced Mathematics, vol. 7, Cambridge University Press.

Laurent, O. (1999). Polarized proof-nets: Proof-nets for LC (Extended Abstract). In: *LNCS 1581 (TLCA '99)*, 213–217.

Laurent, O. (2001). A token machine for full geometry of interaction (Extended Abstract). In: *LNCS 2044 (TLCA '01)*, 283–297.

Laurent, O. (March 2002). Étude de la polarisation en logique. (A study of polarization in logic.) Thèse de Doctorat. Institut de Mathématiques de Luminy – Université Aix-Marseille II.

Liang, C. and Miller, D. (2009). Focusing and polarization in linear, intuitionistic, and classical logic. *Theoretical Computer Science* **410** (46) 4747–4768.

Mac Lane, S. (1971). *Categories for the Working Mathematician*, Springer.

Manes, E.G. and Arbib, M.A. (1986). *Algebraic Approaches to Program Semantics*, Springer-Velag.

Melliès, P.-A. (2009). Categorical semantics of linear logic. Published in: 'Interactive models of computation and program behaviour'. Pierre-Louis Curien, Hugo Herbelin, Jean-Louis Krivine, Paul-André Melliès. *Panoramas et Synthèses* 27, Société Mathématique de France.

Melliès, P.-A. (2013). Dialogue categories and frobenius monoids. In: Coecke, B., Ong, L., and Panangaden, P. (eds.) *Computation, Logic, Games, and Quantum Foundations – The Many Facets of Samson Abramsky*, Lecture Notes in Computer Science, vol. 7860, Springer, 197–224.

Melliès, P.-A. Dialogue categories and chiralities, manuscript. 2006–2012. Available at: http://www.pps.univ-paris-diderot.fr/∼mellies/tensorial-logic.html

Miller, D. (2004). An overview of linear logic programming. In: Ehrhard, T., Girard, J.-Y., Ruet, P. and Scott, P. (eds.) *Linear Logic and Computer Science*, LMS Lecture Note Series, vol. 316, C.U.P., 119–150.

Miller, D. Tutorial: Sequent calculus: Overview and recent developments. In: *8th Panhellenic Logic Symposium Ioannina*, Greece, July 4–8, 2011 (slides available: http://www.lix.polytechnique.fr/Labo/Dale.Miller/papers/pls8.html)

Simpson, A. and Plotkin, G. (2000) Complete axioms for categorical fixed-point operators. In: *Proceedings of the 15th Annual IEEE Symposium on Logic in Computer Science (LICS)*, IEEE Computer Society, 30–41.

https://doi.org/10.1017/S0960129517000196 Published online by Cambridge University Press