Proceedings of the Edinburgh Mathematical Society (2025) **68**, 705–711
doi:10.1017/S0013091525000045

A NOTE ON SEPARABILITY IN OUTER AUTOMORPHISM GROUPS

FRANCESCO FOURNIER-FACIO

Department of Pure Mathematics and Mathematical Statistics, University of Cambridge, UK
Email: ff373@cam.ac.uk

(Received 4 November 2024)

Abstract We give a criterion for separability of subgroups of certain outer automorphism groups. This answers questions of Hagen and Sisto, by strengthening and generalizing a result of theirs on mapping class groups.

Keywords: separable subgroup; outer automorphism; acylindrically hyperbolic; mapping class group

1. Introduction

A subgroup $H \leqslant G$ is called *separable* if for all $g \in G \setminus H$ there exists a finite quotient $q\colon G \to Q$ such that $q(g) \notin q(H)$. A question of Reid [26, Question 3.5] asks whether all convex-cocompact subgroups of mapping class groups (as defined in [9], see also [19] for a characterization) are separable. This was first verified for virtually cyclic subgroups [21], and the full conjecture is known conditionally on the residual finiteness of hyperbolic groups [3]. Hagen and Sisto show in [17] that certain examples of free convex-cocompact subgroups, constructed by Mj [22], are separable. In order to do so, they prove the following criterion.

Theorem ([17, Theorem 1.1]). *Let $g \geqslant 2$, and let $H \leqslant \mathrm{MCG}(\Sigma_g)$. Suppose that H is torsion-free, malnormal, and convex-cocompact. If the preimage of H under the natural quotient map $\mathrm{MCG}(\Sigma_g \setminus \{p\}) \to \mathrm{MCG}(\Sigma_g)$, for some point $p \in \Sigma_g$, is conjugacy separable, then H is separable.*

Remark. The theorem is stated for $g \geqslant 1$; however, in that case, $\mathrm{MCG}(\Sigma_g)$ is virtually free so every finitely generated subgroup is separable. We focus on the case $g \geqslant 2$ for coherence with the results of this note.

© The Author(s), 2025. Published by Cambridge University Press on Behalf of The Edinburgh Mathematical Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (https://creativecommons.org/licenses/by/4.0/), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

https://doi.org/10.1017/S0013091525000045 Published online by Cambridge University Press

A group G is said to be *conjugacy separable* if for all non-conjugate elements $g, h \in G$ there exists a finite quotient $q\colon G \to Q$ such that $q(g)$ and $q(h)$ are non-conjugate. Our first result removes the other hypotheses on H, answering [17, Question 1.4].

Theorem A. *Let $g \geqslant 2$, and let $H \leqslant \mathrm{MCG}(\Sigma_g)$. If the preimage of H under the natural quotient map $\mathrm{MCG}(\Sigma_g \setminus \{p\}) \to \mathrm{MCG}(\Sigma_g)$, for some point $p \in \Sigma_g$, is conjugacy separable, then H is separable.*

We can replace the hypothesis of conjugacy separability by a more geometric one, which goes in the direction of Reid's question [26, Question 3.5]. This is the same way that Hagen and Sisto verify that their criterion holds for the groups constructed by Mj.

Corollary B. *Let $g \geqslant 2$, and let $H \leqslant \mathrm{MCG}(\Sigma_g)$ be a convex-cocompact subgroup. If the preimage of H under the natural quotient map $\mathrm{MCG}(\Sigma_g \setminus \{p\}) \to \mathrm{MCG}(\Sigma_g)$, for some point $p \in \Sigma_g$, acts properly and cocompactly on a $\mathrm{CAT}(0)$ cube complex, then H is separable.*

Notice that Corollary B applies to *all* groups constructed in [22], without needing to modify the construction to ensure malnormality, as in [17, Section 5].

Theorem A will follow from a more general result.

Theorem C. *Let G be a finitely generated group with trivial centre, and let $H \leqslant \mathrm{Out}(G)$. Suppose that $\mathrm{Aut}(G)$ is acylindrically hyperbolic and has no non-trivial finite normal subgroups. If the preimage of H under the natural quotient map $\mathrm{Aut}(G) \to \mathrm{Out}(G)$ is conjugacy separable, then H is separable.*

Thanks to recent works proving acylindrical hyperbolicity of automorphism groups [8, 10–13], this can be applied in several contexts. We isolate two instances.

Corollary D. *Let G be a torsion-free hyperbolic group, and let $H \leqslant \mathrm{Out}(G)$. If the preimage of H under the natural quotient map $\mathrm{Aut}(G) \to \mathrm{Out}(G)$ is conjugacy separable, then H is separable.*

Corollary E. *Let Γ be a finite simplicial graph that does not decompose as a join of two non-empty subgraphs. Let A_Γ be the corresponding right-angled Artin group, and let $H \leqslant \mathrm{Out}(A_\Gamma)$. If the preimage of H under the natural quotient map $\mathrm{Aut}(A_\Gamma) \to \mathrm{Out}(A_\Gamma)$ is conjugacy separable, then H is separable.*

Both of these corollaries apply to $\mathrm{Out}(F_n)$. In this case, there are notions of convex-cocompact subgroups [6, 7, 16]; the corresponding preimage in $\mathrm{Aut}(F_n)$ is hyperbolic [7], but we do not know of instances in which conjugacy separability of such groups is known (besides the case of virtually cyclic subgroups).

Question F. Are there examples of subgroups $H \leqslant \mathrm{Out}(F_n)$ that are separable, not virtually cyclic, and satisfy some version of convex-cocompactness?

We remark that free-by-cyclic groups $F_n \rtimes \mathbb{Z}$ often have non-separable subgroups [20].

The proof of Theorem C involves an element of G that recognises non-inner automorphisms. The existence of such an element relies on the acylindrical hyperbolicity of

https://doi.org/10.1017/S0013091525000045 Published online by Cambridge University Press

$\mathrm{Aut}(G)$, and is what allows to strengthen and generalise the criterion of Hagen and Sisto, with a shorter proof. It partially answers [17, Questions 1.5] (Proposition 2.6) and [17, Question 1.6] (Proposition 2.8).

2. Proofs

We will use π to denote the quotient map $\pi\colon \mathrm{Aut}(G) \to \mathrm{Out}(G)$, and for a subgroup $H \leqslant \mathrm{Out}(G)$ we denote $\widetilde{H} := \pi^{-1}H \leqslant \mathrm{Aut}(G)$. The starting point is the following criterion for separability, which in turn is based on Grossman's criterion for residual finiteness [14].

Proposition 2.1. ([17, Proposition 2.5]). *Let G be a finitely generated group with trivial centre. Let $H \leqslant \mathrm{Out}(G)$ and $\alpha \in \mathrm{Aut}(G)$. Suppose that:*

1. *There exists $x \in G$ such that $\alpha(x) \neq h(x)$ for all $h \in \widetilde{H}$;*
2. *$\widetilde{H}$ is conjugacy separable.*

Then there exists a finite quotient $q\colon \mathrm{Out}(G) \to Q$ such that $q(\pi(\alpha)) \notin q(H)$.

Let us formulate the special case that we will use:

Corollary 2.2. *Let G be a finitely generated group with trivial centre and let $H \leqslant \mathrm{Out}(G)$. Suppose that:*

1. *There exists $\gamma \in \mathrm{Inn}(G)$ such that the centraliser of γ in $\mathrm{Aut}(G)$ is $\langle\gamma\rangle$;*
2. *$\widetilde{H}$ is conjugacy separable.*

Then H is separable.

Proof. We show that Proposition 2.1 holds for H and an arbitrary $\alpha \in \mathrm{Aut}(G) \setminus \widetilde{H}$. Choose $x \in G$ such that the corresponding inner automorphism γ_x is as in the first assumption of the corollary. Suppose that $\alpha(x) = h(x)$ for some $h \in \widetilde{H}$. Then $h^{-1}\alpha(x) = x$, and so $h^{-1}\alpha$ belongs to the centraliser of γ_x in $\mathrm{Aut}(G)$. By the choice of x, we have $h^{-1}\alpha \in \langle\gamma_x\rangle \leqslant \mathrm{Inn}(G) \leqslant \widetilde{H}$, which contradicts $\alpha \notin \widetilde{H}$. □

Acylindrical hyperbolicity of $\mathrm{Aut}(G)$ ensures that the first item holds.

Lemma 2.3. *Let G be a group such that $\mathrm{Inn}(G)$ is infinite and $\mathrm{Aut}(G)$ is acylindrically hyperbolic and has no non-trivial finite normal subgroups. Then there exists $\gamma \in \mathrm{Inn}(G)$ such that the centraliser of γ in $\mathrm{Aut}(G)$ is $\langle\gamma\rangle$.*

In fact, such a γ can be found by performing a simple random walk on G: see Proposition 2.8 and its proof.

Proof. Recall that a subgroup of an acylindrically hyperbolic group is called *suitable* if it is non-elementary and does not normalise any non-trivial finite normal subgroup. In an acylindrically hyperbolic group with no non-trivial finite normal subgroups, every infinite normal subgroup is suitable [25, Lemma 2.4]. In particular, $\mathrm{Inn}(G) \leqslant \mathrm{Aut}(G)$

https://doi.org/10.1017/S0013091525000045 Published online by Cambridge University Press

is suitable. Therefore, there exists an inner automorphism $\gamma \in \mathrm{Inn}(G)$ such that the elementary closure of γ in $\mathrm{Aut}(G)$ is reduced to $\langle\gamma\rangle$ [18, Lemma 5.6]. In particular, the centraliser of γ in $\mathrm{Aut}(G)$ is reduced to $\langle\gamma\rangle$ [4, Corollary 6.6]. □

Proof of Theorem C. Combine Lemma 2.3 and Corollary 2.2. □

For the next applications, we will use the following criterion to check that an automorphism group has no non-trivial finite normal subgroups. We say that G has the *unique root property* if $x^n = y^n$ for some $x, y \in G, n \geqslant 1$ implies $x = y$.

Lemma 2.4. *Let G be a group with trivial centre and with the unique root property. Then $\mathrm{Aut}(G)$ has no non-trivial finite normal subgroups.*

Proof. Suppose that $N \leqslant \mathrm{Aut}(G)$ is a finite normal subgroup. The action of $G \cong \mathrm{Inn}(G)$ on N by conjugacy has a finite index kernel K. Then every element of K commutes with every element of N, in other words, automorphisms in N fix K pointwise. Now let $\alpha \in N$ and $x \in G$. Let $n \geqslant 1$ be such that $x^n \in K$, so x^n is fixed by α. Then $x^n = \alpha(x^n) = \alpha(x)^n$, so by the unique root property $\alpha(x) = x$. This shows that α fixes every element of G and we conclude. □

Proof of Corollary D. If G is torsion-free elementary hyperbolic, then G is either trivial or isomorphic to $\mathbb{Z}$, and in both cases $\mathrm{Out}(G)$ is finite, so all subgroups are separable.

If G is torsion-free non-elementary hyperbolic, then $\mathrm{Aut}(G)$ is acylindrically hyperbolic [12, Theorem 1.3]. Moreover, G has trivial centre and the unique root property [2, Lemma 2.2] and so $\mathrm{Aut}(G)$ has no non-trivial finite normal subgroups by Lemma 2.4. Therefore, Theorem C applies. □

Proof of Corollary E. Let $G = A_\Gamma$ be as in the statement. If Γ has at most one vertex, then G is either trivial or isomorphic to $\mathbb{Z}$, and in both cases $\mathrm{Out}(G)$ is finite, so all subgroups are separable.

If Γ has at least two vertices, then $\mathrm{Aut}(G)$ is acylindrically hyperbolic [11, Theorem 1.5]. Moreover, G has trivial centre and the unique root property [5, 3-2) and 3-3)] and so $\mathrm{Aut}(G)$ has no non-trivial finite normal subgroups by Lemma 2.4. Therefore, Theorem C applies. □

For the results on mapping class groups, we apply Corollary D to the special case of surface groups.

Proof of Theorem A. Recall the Dehn–Nielsen–Baer Theorem: $\mathrm{MCG}(\Sigma_g)$ can be identified with an index-2 subgroup of $\mathrm{Out}(\pi_1(\Sigma_g))$, and $\mathrm{MCG}(\Sigma_g \setminus \{p\})$ is the corresponding index-2 subgroup of $\mathrm{Aut}(\pi_1(\Sigma_g))$. Under these identifications, given a subgroup $H \leqslant \mathrm{MCG}(\Sigma_g)$, its preimage in $\mathrm{MCG}(\Sigma_g \setminus \{p\})$ is the same as its preimage in $\mathrm{Aut}(\pi_1(\Sigma_g))$. Since $\pi_1(\Sigma_g)$ is torsion-free non-elementary hyperbolic, we can apply Corollary D to get separability of H in $\mathrm{Out}(\pi_1(\Sigma_g))$, which then implies separability in $\mathrm{MCG}(\Sigma_g)$. □

Proof of Corollary B. By Theorem A, it suffices to show that, under the hypotheses, the preimage $\widetilde{H}$ is conjugacy separable. By convex-cocompactness of H, $\widetilde{H}$ is hyperbolic

https://doi.org/10.1017/S0013091525000045 Published online by Cambridge University Press

[9, 15]. By assumption, $\widetilde{H}$ acts properly and cocompactly on a CAT(0) cube complex, and so it is virtually compact special [1, 27]. It follows that $\widetilde{H}$ is conjugacy separable [24]. □

Let us end by addressing two questions from [17], which ask for elements that recognise non-inner automorphisms. The observation behind Lemma 2.3 allows to answer both, under the assumptions of Theorem C. The two questions are asked for torsion-free acylindrically hyperbolic groups, with the case of hyperbolic groups being singled out. It is an open question whether the automorphism group of a finitely generated acylindrically hyperbolic group is always acylindrically hyperbolic [10, Question 1.1].

Question 2.5. ([17, Question 1.5]). Let G be a torsion-free acylindrically hyperbolic group, and let $\phi_1, \dots, \phi_n$ be non-inner automorphisms of G. Does there exist $x \in G$ with x and $\phi_i(x)$ non-conjugate for all i?

Proposition 2.6. *Let G be a group such that $\mathrm{Inn}(G)$ is infinite and $\mathrm{Aut}(G)$ is acylindrically hyperbolic and has no non-trivial finite normal subgroups: for instance, a torsion-free non-elementary hyperbolic group. Then there exists $x \in G$ with the following property: for every non-inner automorphism ϕ, x and $\phi(x)$ are non-conjugate.*

Recall that torsion-free non-elementary hyperbolic groups do indeed satisfy the hypotheses, as we saw in the proof of Corollary D.

Proof. Let x be such that γ_x satisfies the statement of Lemma 2.3. Suppose that $\phi(x) = hxh^{-1}$. Then $\gamma_h^{-1}\phi$ fixes x, so it centralises γ_x. By the choice of x, we have $\gamma_h^{-1}\phi \in \langle \gamma_x \rangle \leqslant \mathrm{Inn}(G)$ and so $\phi \in \mathrm{Inn}(G)$. □

Question 2.7. ([17, Question 1.6]). Let G be a torsion-free acylindrically hyperbolic group, let ϕ be a non-inner automorphism of G, and let (w_n) be a simple random walk on G. Is it true that, with probability going to 1 as n goes to infinity, w_n is not conjugate to $\phi(w_n)$?

Note that finite generation is implicit in this question, as *simple* random walks are not defined over infinitely generated groups.

Proposition 2.8. *Let G be a finitely generated group with trivial centre such that $\mathrm{Aut}(G)$ is acylindrically hyperbolic and has no non-trivial finite normal subgroups: for instance, a torsion-free non-elementary hyperbolic group. Let (w_n) be a simple random walk on G. Then, with probability going to 1 as n goes to infinity, w_n has the following property: for every non-inner automorphism ϕ, w_n and $\phi(w_n)$ are non-conjugate.*

Proof. We will use a result from [23], from which we recall some terminology, in the special case we are interested in. We call an element $\alpha \in \mathrm{Aut}(G)$ *asymmetric* if its elementary closure is $\langle \alpha \rangle$. Let μ be a probability distribution on $\mathrm{Aut}(G)$. We say that μ is *admissible* (with respect to a fixed acylindrical action) if the support of μ is bounded and generates a non-elementary subgroup containing an asymmetric element. Let ν be the uniform measure on a finite generating set of G, and let μ be the pushforward of ν under the map $G \to \mathrm{Aut}(G)$. The simple random walk (w_n) is generated by ν, and it induces

https://doi.org/10.1017/S0013091525000045 Published online by Cambridge University Press

a random walk (γ_{w_n}) generated by μ. Since the support of μ is finite, and it generates $\mathrm{Inn}(G)$ which is non-elementary and contains an asymmetric element (Lemma 2.3), μ is admissible. The cyclic subgroups $\langle\gamma_{w_n}\rangle$ are called *random subgroups* of $\mathrm{Aut}(G)$ (for $k=1$) in the language of [23].

Now we can apply [23, Theorem 2.5], which states that with probability going to 1 as n goes to infinity, γ_{w_n} is an asymmetric element of $\mathrm{Aut}(G)$. This implies that the centraliser of γ_{w_n} is $\langle\gamma_{w_n}\rangle$ [4, Corollary 6.6], and we conclude as in Proposition 2.6. □

Acknowledgements. The author thanks Mark Hagen, Jonathan Fruchter, Monika Kudlinska, Alessandro Sisto, Ric Wade and Henry Wilton for useful comments on a first version; and the anonymous referee for more useful comments, especially for suggesting Proposition 2.8. The author is supported by the Herchel Smith Postdoctoral Fellowship Fund.

References

(1) I. Agol, The virtual Haken conjecture, *Doc. Math.* **18** (2013), 1045–1087, With an appendix by Agol, D. Groves, and J. Manning.
(2) L. Bartholdi and O. Bogopolski, On abstract commensurators of groups, *J. Group Theory* **13** (2010), 6, 903–922.
(3) J. Behrstock, M. Hagen, A. Martin and A. Sisto, A combinatorial take on hierarchical hyperbolicity and applications to quotients of mapping class groups, *J. Topol.* **17** (2024), 3, 94.
(4) F. Dahmani, V. Guirardel and D. Osin, Hyperbolically embedded subgroups and rotating families in groups acting on hyperbolic spaces, *Mem. Amer. Math. Soc.* **245** (2017), 1156, v+152.
(5) G. Duchamp and D. Krob, Partially commutative Magnus transformations, *Internat. J. Algebra Comput.* **3** (1993), 1, 15–41.
(6) S. Dowdall and S. J. Taylor, The co-surface graph and the geometry of hyperbolic free group extensions, *J. Topol.* **10** (2017), 2, 447–482.
(7) S. Dowdall and S. J. Taylor, Hyperbolic extensions of free groups, *Geom. Topol.* **22** (2018), 1, 517–570.
(8) A. Escalier and C. Horbez, Graph products and measure equivalence: classification, rigidity, and quantitative aspects, arXiv preprint arXiv:2401.04635.
(9) B. Farb and L. Mosher, Convex cocompact subgroups of mapping class groups, *Geom. Topol.* **6** (2002), 91–152.
(10) A. Genevois, Negative curvature in automorphism groups of one-ended hyperbolic groups, *J. Comb. Algebra* **3** (2019), 3, 305–329.
(11) A. Genevois, Automorphisms of graph products of groups and acylindrical hyperbolicity, *Mem. Amer. Math. Soc.* **301** (2024), 1509.
(12) A. Genevois and C. Horbez, Acylindrical hyperbolicity of automorphism groups of infinitely ended groups, *J. Topol.* **14** (2021), 3, 963–991.
(13) A. Genevois and A. Martin, Automorphisms of graph products of groups from a geometric perspective, *Proc. Lond. Math. Soc. (3)* **119** (2019), 6, 1745–1779.
(14) E. K. Grossman, On the residual finiteness of certain mapping class groups, *J. London Math. Soc. (2)* **9** (1974/75), 160–164.
(15) U. Hamenstädt, Word hyperbolic extensions of surface groups, arXiv preprint math/0505244.

https://doi.org/10.1017/S0013091525000045 Published online by Cambridge University Press

(16) U. Hamenstädt and S. Hensel, Stability in outer space, *Groups Geom. Dyn.* **12** (2018), 1, 359–398.

(17) M. Hagen and A. Sisto, Some examples of separable convex-cocompact subgroups, *Bull. Lond. Math. Soc.* **55** (2023), 5, 2242–2257.

(18) M. Hull, Small cancellation in acylindrically hyperbolic groups, *Groups Geom. Dyn.* **10** (2016), 4, 1077–1119.

(19) R. P. Kent IV and C. J. Leininger, Shadows of mapping class groups: capturing convex cocompactness, *Geom. Funct. Anal.* **18** (2008), 4, 1270–1325.

(20) M. Kudlinska, On subgroup separability of free-by-cyclic and deficiency 1 groups, *Bull. Lond. Math. Soc.* **56** (2024), 1, 338–351.

(21) C. J. Leininger and D. B. McReynolds, Separable subgroups of mapping class groups, *Topology Appl.* **154** (2007), 1, 1–10.

(22) M. Mj, Cubulating surface-by-free groups, With an appendix by J. F. Manning, Mj and M. Sageev. arXiv:1908.03545, to appear in J. Topol.

(23) J. Maher and A. Sisto, Random subgroups of acylindrically hyperbolic groups and hyperbolic embeddings, *Int. Math. Res. Not. IMRN* (2019), 13, 3941–3980.

(24) A. Minasyan and P. Zalesskii, Virtually compact special hyperbolic groups are conjugacy separable, *Comment. Math. Helv.* **91** (2016), 4, 609–627.

(25) D. Osin, $Out(F_n)$-invariant probability measures on the space of n-generated marked groups, arXiv preprint arXiv:2207.03659.

(26) A. W. Reid, *Surface subgroups of mapping class groups*, Problems on mapping class groups and related topics, Proc. Sympos. Pure Math., 74, Amer. Math. Soc., Providence, RI, 2006, 257–268.

(27) D. T. Wise, *The structure of groups with a quasiconvex hierarchy*, Annals of Mathematics Studies, 209, Princeton University Press, Princeton, NJ, (2021) © 2021.

https://doi.org/10.1017/S0013091525000045 Published online by Cambridge University Press