Math. Struct. in Comp. Science (2018), *vol.* 28, *pp.* 45–76. © Cambridge University Press 2016
doi:10.1017/S0960129516000050 First published online 3 May 2016

Realizability algebras III: some examples

JEAN-LOUIS KRIVINE

University Paris-Diderot - C.N.R.S., France
Email: jean-louis.krivine@pps.univ-paris-diderot.fr

We use the technique of "classical realizability" to build new models of ZF + DC in which R is not well ordered. This gives new relative consistency results, which are not obtainable by forcing. This gives also a new method to get programs from proofs of arithmetical formulas with dependent choice.

1. Introduction

The notion of *realizability algebra*, which was introduced in Krivine (2011; 2012), is a tool to study the proof-program correspondence and to build new models of set theory, which we call *realizability models of ZF*.

It is a variant of the well-known notion of *combinatory algebra*, with a new instruction cc, and a new type for the *environments*.

The *sets of forcing conditions*, in common use in set theory, are (very) particular cases of realizability algebras; and the forcing models of ZF are very particular cases of realizability models.

We show here how to extend an arbitrary realizability algebra, by means of a certain set of conditions, so that the axiom DC of dependent choice is realized.

In order to avoid introducing new instructions, we use an idea of Miquel (2011).

This technique has applications of two kinds:

1. Construction of models of ZF + DC.
   When the initial realizability algebra is not trivial (that is, if we are not in the case of forcing or equivalently, if the associated Boolean algebra $\mathbb{2}$ is $\neq \{0,1\}$), then we always obtain in this way a model of ZF *which satisfies DC + there is no well ordering of* $\mathbb{R}$.
   By suitably choosing the realizability algebras, we can get, for instance, the relative consistency over ZF of the following two theories:
   i. ZF + DC + there exists an increasing function $i \mapsto X_i$, from the countable atomless Boolean algebra $\mathcal{B}$ into $\mathcal{P}(\mathbb{R})$ such that:
   $X_0 = \{0\}$; $i \neq 0 \Rightarrow X_i$ is uncountable;
   $X_i \cap X_j = X_{i \wedge j}$;
   if $i \wedge j = 0$ then $X_{i \vee j}$ is equipotent with $X_i \times X_j$;
   $X_i \times X_i$ is equipotent with X_i;
   there exists a surjection from X_1 onto $\mathbb{R}$;
   if there exists a surjection from X_j onto X_i, then $i \leqslant j$;
   if $i, j \neq 0, i \wedge j = 0$, there is no surjection from $X_i \uplus X_j$ (*disjoint union*) onto

https://doi.org/10.1017/S0960129516000050 Published online by Cambridge University Press

$X_i \times X_j$;
more generally, if $J \subset \mathcal{B}$ and if there exists a surjection from $\bigcup_{j\in J} X_j$ onto X_i, then $i \leqslant j$ for some $j \in J$.

In particular, there exists a sequence of subsets of $\mathbb{R}$, the cardinals of which are not comparable, and also a decreasing sequence of subsets of $\mathbb{R}$, the cardinals of which are strictly decreasing.

ii. ZF + DC + there exists $X \subset \mathbb{R}$ such that:
X is uncountable and there is no surjection from X onto $\aleph_1$
(and therefore, every well-orderable subset of X is countable);
$X \times X$ is equipotent with X;
there exists a total order on X, every proper initial segment of which is countable;
there exists a surjection from $X \times \aleph_1$ onto $\mathbb{R}$;
there exists an injection from $\aleph_1$ (thus also from $X \times \aleph_1$) into $\mathbb{R}$.

2. Curry–Howard correspondence.
With this technique of extension of realizability algebras, we can obtain a program from any proof, in ZF + DC, of an arithmetical formula F, which is a λ_c-term, that is, a λ-term containing cc, *but no other new instruction.*
This is a notable difference with the method given in Krivine (2003; 2009), where we use the instruction `quote` and which is, on the other hand, simpler and not limited to arithmetical formulas.

It is important to observe that the program we get in this way does not really depend on the given proof of DC $\to F$ in ZF, but only on the *program P extracted from this proof*, which is a closed λ_c-term. Indeed, we obtain this program by means of an operation of *compilation* applied to P (look at the remark at the end of the introduction of Krivine (2011)).

Finally, apart from applications 1 and 2, we may notice Theorem 27, which gives an interesting property of *every* realizability model: as soon as the Boolean algebra $\beth 2$ is not trivial (i.e. if the model is not a forcing model), there exists a non-well-orderable individual.

2. Generalities

2.1. *Realizability algebras*

It is a first-order structure, which is defined in Krivine (2011). We recall here briefly the definition (with a little change) and some essential properties:

A *realizability algebra* $\mathcal{A}$ is made up of three sets: Λ (the set of *terms*), Π (the set of *stacks*), $\Lambda \star \Pi$ (the set of *processes*) with the following operations:

$(\xi, \eta) \mapsto (\xi)\eta$ from Λ^2 into Λ (*application*);
$(\xi, \pi) \mapsto \xi \cdot \pi$ from $\Lambda \times \Pi$ into Π (*push*);
$(\xi, \pi) \mapsto \xi \star \pi$ from $\Lambda \times \Pi$ into $\Lambda \star \Pi$ (*process*);
$\pi \mapsto \mathsf{k}_\pi$ from Π into Λ (*continuation*).

https://doi.org/10.1017/S0960129516000050 Published online by Cambridge University Press

There are, in Λ, distinguished elements $\mathsf{B}, \mathsf{C}, \mathsf{I}, \mathsf{K}, \mathsf{W}, \mathsf{cc}$, called *elementary combinators* or *instructions.*

Notation.
The term $(\ldots(((\xi)\eta_1)\eta_2)\ldots)\eta_n$ will be also written as $(\xi)\eta_1\eta_2\ldots\eta_n$ or $\xi\eta_1\eta_2\ldots\eta_n$.
For instance: $\xi\eta\zeta = (\xi)\eta\zeta = (\xi\eta)\zeta = ((\xi)\eta)\zeta$.

We define a pre-order on $\Lambda \star \Pi$, denoted by $\succ$, which is called *execution*;
$\xi \star \pi \succ \xi' \star \pi'$ is read as: *the process* $\xi \star \pi$ *reduces to* $\xi' \star \pi'$.
It is the smallest reflexive and transitive binary relation, such that, for any $\xi, \eta, \zeta \in \Lambda$ and $\pi, \varpi \in \Pi$, we have:

$(\xi)\eta \star \pi \succ \xi \star \eta \cdot \pi$.
$\mathsf{I} \star \xi \cdot \pi \succ \xi \star \pi$.
$\mathsf{K} \star \xi \cdot \eta \cdot \pi \succ \xi \star \pi$.
$\mathsf{W} \star \xi \cdot \eta \cdot \pi \succ \xi \star \eta \cdot \eta \cdot \pi$.
$\mathsf{C} \star \xi \cdot \eta \cdot \zeta \cdot \pi \succ \xi \star \zeta \cdot \eta \cdot \pi$.
$\mathsf{B} \star \xi \cdot \eta \cdot \zeta \cdot \pi \succ \xi \star (\eta)\zeta \cdot \pi$.
$\mathsf{cc} \star \xi \cdot \pi \succ \xi \star \mathsf{k}_\pi \cdot \pi$.
$\mathsf{k}_\pi \star \xi \cdot \varpi \succ \xi \star \pi$.

We are also given a subset $\perp\!\!\!\perp$ of $\Lambda \star \Pi$ such that:

$$\xi \star \pi \succ \xi' \star \pi',\ \xi' \star \pi' \in \perp\!\!\!\perp \ \Rightarrow\ \xi \star \pi \in \perp\!\!\!\perp.$$

Given two processes $\xi \star \pi, \xi' \star \pi'$, the notation $\xi \star \pi \gg \xi' \star \pi'$ means:

$$\xi \star \pi \notin \perp\!\!\!\perp \Rightarrow \xi' \star \pi' \notin \perp\!\!\!\perp.$$

Therefore, obviously, $\xi \star \pi \succ \xi' \star \pi' \ \Rightarrow\ \xi \star \pi \gg \xi' \star \pi'$.

Finally, we choose a set of terms $\mathsf{PL}_{\mathcal{A}} \subset \Lambda$, containing the elementary combinators: $\mathsf{B}, \mathsf{C}, \mathsf{I}, \mathsf{K}, \mathsf{W}, \mathsf{cc}$ and closed by application. Its elements are called the *proof-like terms of the algebra* $\mathcal{A}$. We write also PL instead of $\mathsf{PL}_{\mathcal{A}}$ if there is no ambiguity about $\mathcal{A}$.

The algebra $\mathcal{A}$ is called *coherent* if, for every proof-like term $\theta \in \mathsf{PL}_{\mathcal{A}}$, there exists a stack π such that $\theta \star \pi \notin \perp\!\!\!\perp$.

Remark. The *sets of forcing conditions* can be considered as degenerate cases of realizability algebras, if we present them in the following way: an inf-semi-lattice P, with a greatest element $\mathbf{1}$ and an initial segment $\perp\!\!\!\perp$ of P (the set of *false* conditions). Two conditions $p, q \in P$ are called *compatible* if their g.l.b. $p \wedge q$ is not in $\perp\!\!\!\perp$.
We get a realizability algebra if we set $\Lambda = \Pi = \Lambda \star \Pi = P$; $\mathsf{B} = \mathsf{C} = \mathsf{I} = \mathsf{K} = \mathsf{W} = \mathsf{cc} = \mathbf{1}$ and $\mathsf{PL} = \{\mathbf{1}\}$; $(p)q = p \cdot q = p \star q = p \wedge q$ and $\mathsf{k}_p = p$. The pre-order $p \succ q$ is defined as $p \leqslant q$, i.e. $p \wedge q = p$. The condition of coherence is $\mathbf{1} \notin \perp\!\!\!\perp$.

2.2. c-*terms and* λ-*terms*

The terms of the language of combinatory algebra, which are built with variables, the constant symbols $\mathsf{B}, \mathsf{C}, \mathsf{I}, \mathsf{K}, \mathsf{W}, \mathsf{cc}$ and the application (binary operation), will be called *combinatory terms* or c-*terms*, in order to distinguish them from the terms of the algebra $\mathcal{A}$, which are elements of Λ.

https://doi.org/10.1017/S0960129516000050 Published online by Cambridge University Press

Each closed c-term (i.e. without variable) takes a value in the algebra $\mathcal{A}$, which is a proof-like term of $\mathcal{A}$.

Let us call *atom* a c-term of length 1, i.e. a constant symbol $\mathsf{B}, \mathsf{C}, \mathsf{I}, \mathsf{K}, \mathsf{W}, \mathsf{cc}$ or a variable.

Lemma 1. Every c-term t can be written, in a unique way, in the form $t = (a)t_1 \dots t_k$ where a is an atom and $t_1, \dots, t_k$ are c-terms.

Proof. Immediate, by recurrence on the length of t. □

The result of the *substitution of* $\xi_1, \dots, \xi_n \in \Lambda$ *to the variables* $x_1, \dots, x_n$ in a c-term t, is a *term* (i.e. an element of Λ) denoted by $t[\xi_1/x_1, \dots, \xi_n/x_n]$ or, more briefly, $t[\vec{\xi}/\vec{x}]$.
The inductive definition is as follows:

$a[\vec{\xi}/\vec{x}] = \xi_i$ if $a = x_i (1 \leqslant i \leqslant n)$;
$a[\vec{\xi}/\vec{x}] = a$ if a is an atom $\neq x_1, \dots, x_n$;
$(tu)[\vec{\xi}/\vec{x}] = (t[\vec{\xi}/\vec{x}])u[\vec{\xi}/\vec{x}]$.

Given a c-term t and a variable x, we define inductively on t, a new c-term denoted by $\lambda x\, t$, which does not contain x. To this aim, we apply the first possible case in the following list:

1. $\lambda x\, t = (\mathsf{K})t$ if t does not contain x.
2. $\lambda x\, x = \mathsf{I}$.
3. $\lambda x(t)u = (\mathsf{C}\lambda x\, t)u$ if u does not contain x.
4. $\lambda x(t)x = t$ if t does not contain x.
5. $\lambda x(t)x = (\mathsf{W})\lambda x\, t$ (if t contains x).
6. $\lambda x(t)(u)v = \lambda x(((\mathsf{B})t)u)v$ (if uv contains x).

Lemma 2. This re-writing is finite for every c-term.

Proof. For each c-term t, define inductively as follows the integers $l[t]$, $v_0[t], v_1[t], v_2[t]$:
$l[a] = 1$ for every atom a; $l[(t)u] = l[t] + l[u] + 2$;
$v_0[a] = 0$ for every atom a; $v_0[(t)u] = l[u]$;
$v_1[x] = 1$; $v_1[a] = 0$ for every atom $a \neq x$; $v_1[(t)u] = v_1[t] + v_1[u]$;
$v_2[t] = 0$ if x is not in t; $v_2[x] = 1$; $v_2[(t)u] = v_2[u]$ if x appears in u;
$v_2[(t)u] = v_2[t] + l[u] + 1$ if x is not in u (and then x appears in t).

$l[t]$ is the length of t;
$v_0[t]$ is the length of the argument, when t is an application;
$v_1[t]$ is the number of occurrences of x in t;
$v_2[t]$ is the distance to the end of the term t, of the last occurrence of x in t.

Rule 5 strictly decreases $v_1(t)$; rule 3 does not change $v_1[t]$ and strictly decreases $v_2[t]$; rule 6 does not change $v_1[t]$ and $v_2[t]$ and strictly decreases $v_0[t]$.
Therefore, after a finite number of applications of rules 3, 5 and 6, we must apply one of the rules 1, 2 or 4, and the re-writing stops. □

Given a c-term t and a variable x, we now define the c-term $\lambda x\, t$ by setting:
$\lambda x\, t = \lambda x\, (\mathsf{I})t$.

https://doi.org/10.1017/S0960129516000050 Published online by Cambridge University Press

This enables us to translate every λ-term into a c-term. In the sequel, almost all c-terms will be written as λ-terms.

The fundamental property of this translation is given by Theorem 3:

Theorem 3. Let t be a c-term with the only variables $x_1,\dots,x_n$; let $\xi_1,\dots,\xi_n \in \Lambda$ and $\pi \in \Pi$. Then $\lambda x_1 \dots \lambda x_n\, t \star \xi_1 \cdot \dots \cdot \xi_n \cdot \pi \succ t[\xi_1/x_1,\dots,\xi_n/x_n] \star \pi$.

Lemma 4. Let a be an atom, $t = (a)t_1 \dots t_k$ a c-term with the only variables $x, y_1,\dots,y_n$ and $\xi, \eta_1,\dots,\eta_n \in \mathbf{\Lambda}$; then

$$(\mathcal{K}x\, t)[\vec{\eta}/\vec{y}] \star \xi \cdot \pi \succ a[\xi/x,\vec{\eta}/\vec{y}] \star t_1[\xi/x,\vec{\eta}/\vec{y}] \cdot \dots \cdot t_k[\xi/x,\vec{\eta}/\vec{y}] \cdot \pi.$$

Proof. By induction on the number of rules 1 to 6 used to translate the term $\mathcal{K}x\, t$. Consider the rule used first.

- Rule 1: we have $(\mathcal{K}x\, t)[\vec{\eta}/\vec{y}] \star \xi \cdot \pi \equiv (\mathsf{K})t[\vec{\eta}/\vec{y}] \star \xi \cdot \pi \succ \mathsf{K} \star t[\vec{\eta}/\vec{y}] \cdot \xi \cdot \pi \succ t[\vec{\eta}/\vec{y}] \star \pi \equiv t[\xi/x,\vec{\eta}/\vec{y}] \star \pi$ because x is not in t. The result follows immediately.
- Rule 2: we have $t = x, \mathcal{K}x\, t = \mathsf{I}$ and the result is trivial.

In rules 3, 4, 5 or 6, we have $t = ut_k$ with $u = at_1 \dots t_{k-1}$, by Lemma 1.

- Rule 3: $(\mathcal{K}x\, t)[\vec{\eta}/\vec{y}] \star \xi \cdot \pi \equiv ((\mathsf{C}\mathcal{K}x\, u)t_k)[\vec{\eta}/\vec{y}] \star \xi \cdot \pi \succ \mathsf{C} \star (\mathcal{K}x\, u)[\vec{\eta}/\vec{y}] \cdot t_k[\vec{\eta}/\vec{y}] \cdot \xi \cdot \pi$
  $\succ (\mathcal{K}x\, u)[\vec{\eta}/\vec{y}] \star \xi \cdot t_k[\vec{\eta}/\vec{y}] \cdot \pi$
  $\succ a[\xi/x,\vec{\eta}/\vec{y}] \star t_1[\xi/x,\vec{\eta}/\vec{y}] \cdot \dots \cdot t_{k-1}[\xi/x,\vec{\eta}/\vec{y}] \cdot t_k[\vec{\eta}/\vec{y}]\pi$ by the induction hypothesis
  $\equiv a[\xi/x,\vec{\eta}/\vec{y}] \star t_1[\xi/x,\vec{\eta}/\vec{y}] \cdot \dots \cdot t_{k-1}[\xi/x,\vec{\eta}/\vec{y}] \cdot t_k[\xi/x,\vec{\eta}/\vec{y}] \cdot \pi$ since x is not in t_k.

In rules 4 and 5, we have $t_k = x$, i.e. $t = (u)x$.

- Rule 4: we have $(\mathcal{K}x\, t)[\vec{\eta}/\vec{y}] \star \xi \cdot \pi \equiv u[\vec{\eta}/\vec{y}] \star \xi \cdot \pi \equiv u[\xi/x,\vec{\eta}/\vec{y}] \star \xi \cdot \pi$ because x is not in u. Since $u = at_1 \dots t_{k-1}$ and $t_k = x$, the result follows immediately.
- Rule 5: we have $t_k = x$ and $(\mathcal{K}x\, t)[\vec{\eta}/\vec{y}] \star \xi \cdot \pi \equiv (\mathsf{W}\mathcal{K}x\, u)[\vec{\eta}/\vec{y}] \star \xi \cdot \pi$
  $\succ \mathsf{W} \star (\mathcal{K}x\, u)[\vec{\eta}/\vec{y}] \cdot \xi \cdot \pi \succ (\mathcal{K}x\, u)[\vec{\eta}/\vec{y}] \star \xi \cdot \xi \cdot \pi$
  $\succ a[\xi/x,\vec{\eta}/\vec{y}] \star t_1[\xi/x,\vec{\eta}/\vec{y}] \cdot \dots \cdot t_{k-1}[\xi/x,\vec{\eta}/\vec{y}] \cdot \xi \cdot \pi$ by the induction hypothesis
  $\equiv a[\xi/x,\vec{\eta}/\vec{y}] \star t_1[\xi/x,\vec{\eta}/\vec{y}] \cdot \dots \cdot t_k[\xi/x,\vec{\eta}/\vec{y}] \cdot \pi$.
- Rule 6: we have $t_k = (v)w$ and $(\mathcal{K}x\, t)[\vec{\eta}/\vec{y}] \star \xi \cdot \pi \equiv (\mathcal{K}x(\mathsf{B})uvw)[\vec{\eta}/\vec{y}] \star \xi \cdot \pi$
  $\succ \mathsf{B} \star u[\xi/x,\vec{\eta}/\vec{y}] \cdot v[\xi/x,\vec{\eta}/\vec{y}] \cdot w[\xi/x,\vec{\eta}/\vec{y}] \cdot \pi$ (by the induction hypothesis)
  $\succ u[\xi/x,\vec{\eta}/\vec{y}] \star t_k[\xi/x,\vec{\eta}/\vec{y}] \cdot \pi$
  $\succ a[\xi/x,\vec{\eta}/\vec{y}] \star t_1[\xi/x,\vec{\eta}/\vec{y}] \cdot \dots \cdot t_{k-1}[\xi/x,\vec{\eta}/\vec{y}] \cdot t_k[\xi/x,\vec{\eta}/\vec{y}] \cdot \pi$.

□

Lemma 5. $(\lambda x\, t)[\vec{\eta}/\vec{y}] \star \xi \cdot \pi \succ t[\xi/x,\vec{\eta}/\vec{y}] \star \pi$.

Proof. Immediate by Lemma 4 and the definition of $\lambda x\, t$ which is $\mathcal{K}x(\mathsf{I})t$. □

Proof of Theorem 3. By induction on n; the case $n = 0$ is trivial. Now, we have:
$\lambda x_1 \dots \lambda x_{n-1}\lambda x_n\, t \star \xi_1 \cdot \dots \cdot \xi_{n-1} \cdot \xi_n \cdot \pi \succ (\lambda x_n t)[\xi_1/x_1,\dots,\xi_{n-1}/x_{n-1}] \star \xi_n \cdot \pi$
(by induction hypothesis) $\succ t[\xi_1/x_1,\dots,\xi_{n-1}/x_{n-1},\xi_n/x_n] \star \pi$ by Lemma 5. □

2.3. *The formal system*

We write formulas and proofs in the language of first-order logic. This formal language consists of:

https://doi.org/10.1017/S0960129516000050 Published online by Cambridge University Press

- *individual variables* $x, y, \ldots$;
- *function symbols* $f, g, \ldots$ of various arities; function symbols of arity 0 are called *constant symbols.*
- *relation symbols*; there are three binary relation symbols: $\not\in, \notin, \subset$.

The terms of this first-order language will be called *ℓ-terms*; they are built in the usual way with individual variables and function symbols.

Remark. Thus, we use four expressions with the word *term*: term, c-term, λ-term and ℓ-term.

The *atomic formulas* are the expressions $\top, \bot, t \not\in u, t \notin u, t \subset u$, where t, u are ℓ-terms.

Formulas are built as usual, from atomic formulas, *with the only logical symbols* $\to, \forall$:

- each atomic formula is a formula;
- if A, B are formulas, then $A \to B$ is a formula;
- if A is a formula and x an individual variable, then $\forall x\, A$ is a formula.

Notations. Let $A_1, \ldots, A_n, A, B$ be formulas. Then:
$A \to \bot$ is written $\neg A$;
$A_1 \to (A_2 \to \cdots \to (A_n \to B)\cdots)$ is written $A_1, A_2, \ldots, A_n \to B$;
$\neg A_1, \ldots, \neg A_n \to \bot$ is written $A_1 \vee \ldots \vee A_n$;
$(A_1, \ldots, A_n \to \bot) \to \bot$ is written $A_1 \wedge \ldots \wedge A_n$;
$\neg\forall x(A_1, \ldots, A_n \to \bot)$ is written $\exists x\{A_1, \ldots, A_n\}$.

The *rules of natural deduction* are the following (the A_i's are formulas, the x_i's are variables of c-term, t, u are c-terms, written as λ-terms):

1. $x_1 : A_1, \ldots, x_n : A_n \vdash x_i : A_i$.
2. $x_1 : A_1, \ldots, x_n : A_n \vdash t : A \to B, \quad x_1 : A_1, \ldots, x_n : A_n \vdash u : A \quad \Rightarrow$
   $x_1 : A_1, \ldots, x_n : A_n \vdash tu : B$.
3. $x_1 : A_1, \ldots, x_n : A_n, x : A \vdash t : B \quad \Rightarrow \quad x_1 : A_1, \ldots, x_n : A_n \vdash \lambda x\, t : A \to B$.
4. $x_1 : A_1, \ldots, x_n : A_n \vdash t : A \quad \Rightarrow \quad x_1 : A_1, \ldots, x_n : A_n \vdash t : \forall x\, A$ where x is an individual variable which does not appear in $A_1, \ldots, A_n$.
5. $x_1 : A_1, \ldots, x_n : A_n \vdash t : \forall x\, A \quad \Rightarrow \quad x_1 : A_1, \ldots, x_n : A_n \vdash t : A[\tau/x]$
   where x is an individual variable and τ is a ℓ-term.
6. $x_1 : A_1, \ldots, x_n : A_n \vdash \mathsf{cc} : ((A \to B) \to A) \to A$ (law of Peirce).
7. $x_1 : A_1, \ldots, x_n : A_n \vdash t : \bot \quad \Rightarrow \quad x_1 : A_1, \ldots, x_n : A_n \vdash t : A$ for every formula A.

2.4. *Realizability models*

We formalize set theory with the first-order language described above. We write, in this language, the axioms of a theory named ZF_ε, which are given in Krivine (2012).
The usual set theory ZF is supposed written with the only relation symbols $\notin, \subset$.
Then, ZF_ε is a *conservative extension* of ZF, which is proved in Krivine (2012).

Remark. In ZF_ε, we have two membership relations: $\in$ which the *weak* or *extensional* one, and ε which is the *strong* one and does not satisfy extensionality.

https://doi.org/10.1017/S0960129516000050 Published online by Cambridge University Press

Therefore, we have also two relations of inclusion: $x \subset y \equiv \forall z(z \notin y \to z \notin x)$ ($\in$-inclusion or ZF-inclusion) and $x \subseteq y \equiv \forall z(z \not\in y \to z \not\in x)$ (ε-inclusion). Finally, we have three notions of equality:
the weakest one is $(x \sim y) \equiv x \subset y \wedge y \subset x$ ($\in$-equality);
the strongest one is $(x = y) \equiv \forall z(x \not\in z \to y \not\in z)$ (Leibniz equality);
the middle one is $(x \simeq y) \equiv x \subseteq y \wedge y \subseteq x$ (ε-equality).

Let us consider a *coherent* realizability algebra $\mathcal{A}$, defined in a model $\mathcal{M}$ of ZF + V = L, which is called the *ground model*. The elements of $\mathcal{M}$ will be called *individuals* (in order to avoid the word *set*, as far as possible).

We defined, in Krivine (2012), a *realizability model*, denoted by $\mathcal{N}_{\mathcal{A}}$ (or even $\mathcal{N}$, if there is no ambiguity about the algebra $\mathcal{A}$).
It has the same domain (the same individuals) as $\mathcal{M}$ and the interpretation of the function symbols is the same as in $\mathcal{M}$.

Each closed formula F of $\mathrm{ZF}_{\varepsilon}$ with parameters in $\mathcal{M}$, has *two truth values* in $\mathcal{N}$, which are denoted by $\|F\|$ (which is a subset of Π) and $|F|$ (which is a subset of Λ).
Here are their definitions:

$|F|$ is defined immediately from $\|F\|$ as follows:

$$\xi \in |F| \;\Leftrightarrow\; (\forall \pi \in \|F\|)\, \xi \star \pi \in \perp\!\!\!\perp.$$

We shall write $\xi \Vdash F$ (read 'ξ *realizes* F') for $\xi \in |F|$.

$\|F\|$ is now defined by recurrence on the length of F:

- F is atomic;
  then F has one of the forms $\top$, $\bot$, $a \not\in b$, $a \subset b$, $a \notin b$ where a, b are parameters in $\mathcal{M}$. We set:
  $\|\top\| = \varnothing$; $\|\bot\| = \Pi$; $\|a \not\in b\| = \{\pi \in \Pi;\ (a, \pi) \in b\}$.
  $\|a \subset b\|, \|a \notin b\|$ are defined simultaneously by induction on $(\mathrm{rk}(a) \cup \mathrm{rk}(b), \mathrm{rk}(a) \cap \mathrm{rk}(b))$ ($\mathrm{rk}(a)$ being the ordinal rank of a in $\mathcal{M}$).
  $$\|a \subset b\| = \bigcup_{c} \{\xi \cdot \pi;\ \xi \in \Lambda,\ \pi \in \Pi,\ (c, \pi) \in a,\ \xi \Vdash c \notin b\};$$
  $$\|a \notin b\| = \bigcup_{c} \{\xi \cdot \xi' \cdot \pi;\ \xi, \xi' \in \Lambda,\ \pi \in \Pi,\ (c, \pi) \in b,\ \xi \Vdash a \subset c,\ \xi' \Vdash c \subset a\}.$$
- $F \equiv A \to B$; then $\|F\| = \{\xi \cdot \pi;\ \xi \Vdash A,\ \pi \in \|B\|\}$.
- $F \equiv \forall x\, A$: then $\|F\| = \bigcup_{a} \|A[a/x]\|$.

The following theorem, proved in Krivine (2012), is an essential tool:

Theorem 6 (Adequacy lemma).
Let $A_1, \ldots, A_n, A$ be closed formulas of $\mathrm{ZF}_{\varepsilon}$; suppose that $x_1 : A_1, \ldots, x_n : A_n \vdash t : A$.
If $\xi_1 \Vdash A_1, \ldots, \xi_n \Vdash A_n$ then $t[\xi_1/x_1, \ldots, \xi_n/x_n] \Vdash A$.
In particular, if $\vdash t : A$, then $t \Vdash A$.

Let F be a closed formula of $\mathrm{ZF}_{\varepsilon}$, with parameters in $\mathcal{M}$. We say that $\mathcal{N}_{\mathcal{A}}$ *realizes* F or that F *is realized in* $\mathcal{N}_{\mathcal{A}}$ (which is written $\mathcal{N}_{\mathcal{A}} \Vdash F$ or even $\Vdash F$), if there exists a proof-like term θ such that $\theta \Vdash F$.

https://doi.org/10.1017/S0960129516000050 Published online by Cambridge University Press

It is shown in Krivine (2012) that *all the axioms of ZF_ε are realized in $\mathcal{N}_\mathcal{A}$,* and thus also all the axioms of ZF.

Remarks. $\mathcal{N}_\mathcal{A}$ is not a Tarski model of ZF_ε, because the truth values of formulas are not 0 or 1 but subsets of Π. We can obtain Tarski models by using the completeness theorem, or else an ultrafilter on the set of truth values equipped with a suitable structure of Boolean algebra (see Krivine 2012). Note that, doing this, we introduce new individuals in the model, i.e. individuals which are not in $\mathcal{M}$.
An individual a of $\mathcal{M}$ is, in general, not related with its interpretation in $\mathcal{N}_\mathcal{A}$. For instance, if no element of a is an ordered pair (b,π) with $\pi \in \Pi$, then a is interpreted as $\varnothing$ in $\mathcal{N}_\mathcal{A}$. Indeed, we have $\|\forall x(x \notin a)\| = \varnothing = \|\top\|$.

Definitions. Given a set of terms $X \subset \Lambda$ and a formula F, we shall use the notation $X \to F$ as an *extended formula*; its truth value is $\|X \to F\| = \{\xi \cdot \pi;\ \xi \in X,\ \pi \in \|F\|\}$.
Two formulas $F[x_1,\dots,x_n]$ and $G[x_1,\dots,x_n]$ of ZF_ε will be called *interchangeable* if the formula $\forall x_1 \dots \forall x_n(F[x_1,\dots,x_n] \leftrightarrow G[x_1,\dots,x_n])$ is realized.
That is, for instance, the case if $\|F[a_1,\dots,a_n]\| = \|G[a_1,\dots,a_n]\|$
or also if $\|F[a_1,\dots,a_n]\| = \|\neg\neg G[a_1,\dots,a_n]\|$
for every $a_1,\dots,a_n \in \mathcal{M}$.

The following lemma gives a useful example:

Lemma 7. For every formula A, define $\overline{\neg}A \subset \Lambda$ by $\overline{\neg}A = \{\mathsf{k}_\pi\ ;\ \pi \in \|A\|\}$.
Then $\neg A \to B$ and $\overline{\neg}A \to B$ are interchangeable, for every formula B.

Proof. We have immediately $\mathsf{k}_\pi \Vdash \neg A$ for every $\pi \in \|A\|$.
Therefore, $\|\overline{\neg}A \to B\| \subset \|\neg A \to B\|$ and it follows that $\mathsf{I} \Vdash (\neg A \to B) \to (\overline{\neg}A \to B)$.
Conversely, let $\xi,\eta \in \Lambda$, $\xi \Vdash \overline{\neg}A \to B$, $\eta \Vdash \neg B$ and let $\pi \in \|A\|$.
We have $\xi\mathsf{k}_\pi \Vdash B$, thus $(\eta)(\xi)\mathsf{k}_\pi \Vdash \bot$ and therefore $(\eta)(\xi)\mathsf{k}_\pi \star \pi \in \perp\!\!\!\perp$.
It follows that $\theta \star \xi \cdot \eta \cdot \pi \in \perp\!\!\!\perp$ with $\theta = \lambda x\lambda y(\mathsf{cc})\lambda k(y)(x)k$. Finally, we have shown that $\theta \Vdash (\overline{\neg}A \to B) \to (\neg B \to A)$, from which the result follows. □

2.5. *Equality and type-like sets*

The formula $x = y$ is, by definition, $\forall z(x \notin z \to y \notin z)$ *(Leibniz equality).*
If t,u are ℓ-terms and F is a formula of ZF_ε, with parameters in $\mathcal{M}$, we define the formula $t = u \hookrightarrow F$. When it is closed, its truth value is:
$\|t = u \hookrightarrow F\| = \|\top\| = \varnothing$ if $\mathcal{M} \models t \neq u$; $\|t = u \hookrightarrow F\| = \|F\|$ if $\mathcal{M} \models t = u$.
The formula $t = u \hookrightarrow \bot$ is written $t \neq u$.
The formula $t_1 = u_1 \hookrightarrow (t_2 = u_2 \hookrightarrow \cdots \hookrightarrow (t_n = u_n \hookrightarrow F)\cdots)$ is written:
$t_1 = u_1, t_2 = u_2, \dots, t_n = u_n \hookrightarrow F$.
The formulas $t = u \to F$ and $t = u \hookrightarrow F$ are interchangeable, as is shown in the:

Lemma 8.

i. $\mathsf{CII} \Vdash \forall x\forall y\,((x = y \to F) \to (x = y \hookrightarrow F))$;
ii. $\mathsf{CI} \Vdash \forall x\forall y\,((x = y \hookrightarrow F) \to (x = y \to F))$.

https://doi.org/10.1017/S0960129516000050 Published online by Cambridge University Press

Proof.

i. Trivial.

ii. Let a, b be individuals; let $\xi \Vdash a = b \hookrightarrow F$, $\eta \Vdash a = b$ and $\pi \in \|F\|$.
We show that $\eta \star \xi \cdot \pi \in \bot\!\!\!\bot$.
Let $c = \{(b, \pi)\}$; by hypothesis on η, we have $\eta \Vdash a \notin c \to b \notin c$. Since $\pi \in \|b \notin c\|$, it suffices to show that $\xi \Vdash a \notin c$. This is clear if $a \neq b$, since $\|a \notin c\| = \varnothing$ in this case.
If $a = b$, then $\xi \Vdash F$, by hypothesis on ξ, thus $\xi \star \pi \in \bot\!\!\!\bot$; but $\|a \notin c\| = \{\pi\}$ in this case, and therefore $\xi \Vdash a \notin c$.

□

We set $\daleth X = X \times \Pi$ for every individual X of $\mathcal{M}$; we define the quantifier $\forall x^{\daleth X}$ as follows:

$$\|\forall x^{\daleth X} F[x]\| = \bigcup_{a \in X} \|F[a]\|.$$

Of course, we set $\exists x^{\daleth X} F[x] \equiv \neg \forall x^{\daleth X} \neg F[x]$.

The quantifier $\forall x^{\daleth X}$ has the intended meaning, which is that the formulas $\forall x^{\daleth X} F[x]$ and $\forall x (x \, \varepsilon \daleth X \to F[x])$ are interchangeable. This is shown by the following:

Lemma 9. $\mathsf{CI} \Vdash \forall x^{\daleth X} F[x] \to \forall x^{\daleth X} \neg\neg F[x]$;
$\mathsf{cc} \Vdash \forall x^{\daleth X} \neg\neg F[x] \to \forall x^{\daleth X} F[x]$;
$\|\forall x^{\daleth X} \neg\neg F[x]\| = \|\forall x (\neg F[x] \to x \notin \daleth X)\|$.

Proof. Immediate. □

Each *functional* $f : \mathcal{M}^n \to \mathcal{M}$, defined in $\mathcal{M}$ by a formula of ZF with parameters, gives a function symbol, that we denote also by f, and which has the same interpretation in the realizability model $\mathcal{N}_\mathcal{A}$.

Proposition 10.
Let $t, t_1, \ldots, t_n, u, u_1, \ldots, u_n$ be ℓ-terms, built with variables $x_1, \ldots, x_k$ and function symbols of $\mathcal{M}$.

i. If $\mathcal{M} \models \forall x_1 \ldots \forall x_k (t_1 = u_1, \ldots, t_k = u_k \to t = u)$, then
$\mathsf{I} \Vdash \forall x_1 \ldots \forall x_k (t_1 = u_1, \ldots, t_k = u_k \hookrightarrow t = u)$.

ii. If $\mathcal{M} \models (\forall x_1 \in X_1) \ldots (\forall x_k \in X_k)(t_1 = u_1, \ldots, t_k = u_k \to t = u)$, then
$\mathsf{I} \Vdash \forall x_1^{\daleth X_1} \ldots \forall x_k^{\daleth X_k} (t_1 = u_1, \ldots, t_k = u_k \hookrightarrow t = u)$.

Proof. Trivial. □

Proposition 11. If $f : X_1 \times \cdots \times X_n \to Y$ is a function in $\mathcal{M}$, its interpretation in $\mathcal{N}_\mathcal{A}$ is a function $f : \daleth X_1 \times \cdots \times \daleth X_n \to \daleth Y$.

Proof. Indeed, let $f', f'' : \mathcal{M}^n \to \mathcal{M}$ be any two functionals which are extensions of the function f to the whole of $\mathcal{M}^n$. By Proposition 10(ii), we have:
$\mathsf{I} \Vdash \forall x_1^{\daleth X_1} \ldots \forall x_k^{\daleth X_k} (f'(x_1, \ldots, x_k) = f''(x_1, \ldots, x_k))$. □

An important example is the set $2 = \{0, 1\}$ equipped with the trivial boolean functions, written $\wedge, \vee, \neg$. The extension to $\mathcal{N}_\mathcal{A}$ of these operations gives a structure of Boolean algebra on $\daleth 2$. It is called the *characteristic Boolean algebra* of the model $\mathcal{N}_\mathcal{A}$.

https://doi.org/10.1017/S0960129516000050 Published online by Cambridge University Press

Conservation of well-foundedness

Theorem 12 below says that every well-founded relation in the ground model $\mathcal{M}$, gives a well-founded relation in the realizability model $\mathcal{N}$.

Theorem 12.

Let $f : \mathcal{M}^2 \to 2$ be a functional defined in the ground model $\mathcal{M}$ such that $f(x,y) = 1$ is a well-founded relation on $\mathcal{M}$.

Then, for every formula $F[x]$ of ZF_ε with parameters in $\mathcal{M}$:

$\mathsf{Y} \Vdash \forall y\,(\forall x(f(x,y) = 1 \hookrightarrow F[x]) \to F[y]) \to \forall y\, F[y]$

with $\mathsf{Y} = AA$ and $A = \lambda a \lambda f(f)(a)af$ (or $A = (\mathsf{W})(\mathsf{B})(\mathsf{BW})(\mathsf{C})\mathsf{B}$).

Proof. Let us fix $b \in X$ and let $\xi \Vdash \forall y\,(\forall x(f(x,y) = 1 \hookrightarrow F[x]) \to F[y])$. We show, by induction on b, following the well-founded relation $f(x,y) = 1$, that $\mathsf{Y} \star \xi \cdot \pi \in \bot\!\!\!\bot$ for every $\pi \in \|F[b]\|$.

Thus, suppose that $\pi \in \|F[b]\|$; since $\mathsf{Y} \star \xi \cdot \pi \succ \xi \star \mathsf{Y}\xi \cdot \pi$, we need to show that $\xi \star \mathsf{Y}\xi \cdot \pi \in \bot\!\!\!\bot$. By hypothesis, we have $\xi \Vdash \forall x(f(x,b) = 1 \hookrightarrow F[x]) \to F[b]$; thus, it suffices to show that $\mathsf{Y}\xi \Vdash f(a,b) = 1 \hookrightarrow F[a]$ for every $a \in X$.

This is clear if $f(a,b) \neq 1$, by definition of $\hookrightarrow$.

If $f(a,b) = 1$, we must show $\mathsf{Y}\xi \Vdash F[a]$, i.e. $\mathsf{Y} \star \xi \cdot \varpi \in \bot\!\!\!\bot$ for every $\varpi \in \|F[a]\|$. But this follows from the induction hypothesis. □

Remarks.

i. If the function f is only defined on a set X in the ground model $\mathcal{M}$, we can apply Theorem 12 to the extension f' of f defined by $f'(x,y) = 0$ if $(x,y) \notin X^2$. This shows that, in the realizability model $\mathcal{N}$, the binary relation $f(x,y) = 1$ is well founded on $\mathbb{1}X$.

ii. We can use Theorem 12 to show that the axiom of foundation of ZF_ε is realized in $\mathcal{N}_\mathcal{A}$.

Indeed, let us define $f : \mathcal{M}^2 \to 2$ by setting $f(x,y) = 1 \Leftrightarrow \exists z((x,z) \in y)$. The binary relation $f(x,y) = 1$ is obviously well founded in $\mathcal{M}$. Now, we have $\mathsf{I} \Vdash \forall x \forall y(f(x,y) \neq 1 \to x \notin y)$ because $\pi \in \|x \notin y\| \Rightarrow f(x,y) = 1$. Thus, the relation $x \varepsilon y$ is stronger than the relation $f(x,y) = 1$, which is well founded in $\mathcal{N}_\mathcal{A}$ by Theorem 12.

2.6. *Integers*

Let $\phi, \alpha \in \Lambda$ and $n \in \mathbb{N}$; we define $(\phi)^n\alpha \in \Lambda$ by setting $(\phi)^0\alpha = \alpha$; $(\phi)^{n+1}\alpha = (\phi)(\phi)^n\alpha$.

For $n \in \mathbb{N}$, we define $\underline{n} = (\sigma)^n\underline{0}$ with $\underline{0} = \mathsf{KI}$ and $\sigma = (\mathsf{BW})(\mathsf{B})\mathsf{B}$;

$\underline{n}$ is 'the integer n' and σ the 'successor' in combinatory logic.

The essential property of $\underline{0}$ and σ is: $\underline{0} \star \phi \cdot \alpha \cdot \pi \succ \alpha \star \pi$; $\sigma \star v \cdot \phi \cdot \alpha \cdot \pi \succ v \star \phi \cdot \phi\alpha \cdot \pi$.

We set $\mathbb{N}_\mathcal{A} = \{(n, \underline{n} \cdot \pi)\ ;\ n \in \mathbb{N}, \pi \in \Pi\}$; it is shown below that $\mathbb{N}_\mathcal{A}$ is the set of integers of the realizability model $\mathcal{N}_\mathcal{A}$.

We define the quantifier $\forall x^{\mathrm{int}}$ as follows:

$$\|\forall x^{\mathrm{int}} F[x]\| = \{\underline{n} \cdot \pi\ ;\ n \in \mathbb{N},\ \pi \in \|F[n]\|\}.$$

which means intuitively

$$\|\forall x^{\mathrm{int}} F[x]\| = \|\forall n^{\mathbb{1}\mathbb{N}}(\{\underline{n}\} \to F[n])\|.$$

https://doi.org/10.1017/S0960129516000050 Published online by Cambridge University Press

The formulas $\forall x^{\mathrm{int}} F[x]$ and $\forall x(x\,\varepsilon\,\mathbb{N}_{\mathcal{A}} \to F[x])$ are interchangeable, as is shown in the:

Lemma 13.
$\lambda x\lambda n\lambda y(y)(x)n \Vdash \forall x^{\mathrm{int}} F[x] \to \forall x^{\mathrm{int}} \neg\neg F[x]$;
$\lambda x\lambda n(\mathsf{cc})(x)n \Vdash \forall x^{\mathrm{int}} \neg\neg F[x] \to \forall x^{\mathrm{int}} F[x]$;
$\|\forall x^{\mathrm{int}} \neg\neg F[x]\| = \|\forall x(\neg F[x] \to x \notin \mathbb{N}_{\mathcal{A}})\|$.

Proof. Immediate. □

Lemma 14.

i. $\mathsf{K} \Vdash \forall x(x \notin \daleth\mathbb{N} \to x \notin \mathbb{N}_{\mathcal{A}})$.
ii. $\lambda x(x)\underline{0} \Vdash 0 \notin \mathbb{N}_{\mathcal{A}} \to \bot$; $\lambda f\lambda x(f)(\sigma)x \Vdash \forall y^{\daleth\mathbb{N}}((y+1) \notin \mathbb{N}_{\mathcal{A}} \to y \notin \mathbb{N}_{\mathcal{A}})$.
iii. $\mathsf{I} \Vdash \forall x^{\mathrm{int}} \left(\forall y^{\daleth\mathbb{N}}(F[y] \to F[y+1]), F[0] \to F[x]\right)$ for every formula $F[x]$ of $\mathrm{ZF}_{\varepsilon}$.

Proof. (i) and (ii) Immediate.
(iii) Let $n \in \mathbb{N}$, $\phi \Vdash \forall y^{\daleth\mathbb{N}}(F[y] \to F[y+1]), \alpha \Vdash F[0]$ et $\pi \in \|F[n]\|$. We must show: $\underline{n} \star \phi \cdot \alpha \cdot \pi \in \perp\!\!\!\perp$ i.e., by Lemma 15, $(\phi)^n\alpha \star \pi \in \perp\!\!\!\perp$.
But it is clear, by recurrence on n, that $(\phi)^n\alpha \Vdash F[n]$ for every $n \in \mathbb{N}$. □

Lemma 14(i) shows that $\mathbb{N}_{\mathcal{A}}$ is a subset of $\daleth\mathbb{N}$.
But it is clear that $\daleth\mathbb{N}$ contains 0 and is closed by the function $n \mapsto n+1$.
Now, by Lemma 14(ii) and (iii), $\mathbb{N}_{\mathcal{A}}$ is the smallest subset of $\daleth\mathbb{N}$ which contains 0 and is closed by the function $n \mapsto n+1$. Therefore:

$$\mathbb{N}_{\mathcal{A}} \textit{ is the set of integers of the model } \mathcal{N}_{\mathcal{A}}.$$

The following Lemmas 15 and 16 will be used in Section 4 (proof of Lemma 39).

Lemma 15.
Let $O, \varsigma \in \Lambda$ be such that: $O \star \phi \cdot \alpha \cdot \pi \succgtr \alpha \star \pi$ and $\varsigma \star v \cdot \phi \cdot \alpha \cdot \pi \succgtr v \star \phi \cdot \phi\alpha \cdot \pi$
for every $\alpha, v, \phi \in \Lambda$ and $\pi \in \Pi$.
Then, for every $n \in \mathbb{N}, \alpha, \zeta, \phi \in \Lambda$ and $\pi \in \Pi$:

i. $(\varsigma)^n O \star \phi \cdot \alpha \cdot \pi \succgtr (\phi)^n\alpha \star \pi$; in particular, $\underline{n} \star \phi \cdot \alpha \cdot \pi \succgtr (\phi)^n\alpha \star \pi$
ii. $(\varsigma)^n O \star \mathsf{CB}\phi \cdot \zeta \cdot \alpha \cdot \pi \succgtr \zeta \star (\phi)^n\alpha \cdot \pi$.

Proof.

i. Proof by recurrence on n; this is clear if $n = 0$; if $n = m+1$, we have:
$\varsigma \star (\varsigma)^m O \cdot \phi \cdot \alpha \cdot \pi \succgtr (\varsigma)^m O \star \phi \cdot \phi\alpha \cdot \pi \succgtr (\phi)^m(\phi)\alpha \star \pi$ by the recurrence hypothesis.
The particular case is $O = \underline{0}, \varsigma = \sigma$.
ii. By (i), we have $(\varsigma)^n O \star \mathsf{CB}\phi \cdot \zeta \cdot \alpha \cdot \pi \succgtr (\mathsf{CB}\phi)^n\zeta \star \alpha \cdot \pi$.
We now show, by recurrence on n, that $(\mathsf{CB}\phi)^n\zeta \star \alpha \cdot \pi \succgtr \zeta \star (\phi)^n\alpha \cdot \pi$.
This is clear if $n = 0$; if $n = m+1$, we have:
$(\mathsf{CB}\phi)^n\zeta \star \alpha \cdot \pi \succgtr \mathsf{CB}\phi \star (\mathsf{CB}\phi)^m\zeta \cdot \alpha \cdot \pi \succgtr \mathsf{C} \star \mathsf{B} \cdot \phi \cdot (\mathsf{CB}\phi)^m\zeta \cdot \alpha \cdot \pi$
$\succgtr \mathsf{B} \star (\mathsf{CB}\phi)^m\zeta \cdot \phi \cdot \alpha \cdot \pi \succgtr (\mathsf{CB}\phi)^m\zeta \star \phi\alpha \cdot \pi \succgtr \zeta \star (\phi)^m(\phi)\alpha \cdot \pi$ (by the recurrence hypothesis).

□

https://doi.org/10.1017/S0960129516000050 Published online by Cambridge University Press

Lemma 16.
Let $\Omega, \Sigma \in \Lambda$ be such that:
$\Omega \star \delta \cdot \phi \cdot \alpha \cdot \pi \succgtr \alpha \star \pi$ and $\Sigma \star \nu \cdot \delta \cdot \phi \cdot \alpha \cdot \pi \succgtr \nu \star \delta \cdot \phi \cdot \phi\alpha \cdot \pi$
for every $\alpha, \delta, \nu, \phi \in \Lambda$ and $\pi \in \Pi$. For instance: $\Omega = (\mathsf{K})(\mathsf{K})\mathsf{I}$; $\Sigma = (\mathsf{B})(\mathsf{BW})(\mathsf{B})\mathsf{B}$.
Then, for every $n \in \mathbb{N}, \alpha, \delta, \zeta, \phi \in \Lambda$ and $\pi \in \Pi$:

i. $(\Sigma)^n\Omega \star \delta \cdot \phi \cdot \alpha \cdot \pi \succ (\phi)^n\alpha \star \pi$.
ii. $(\Sigma)^n\Omega \star \delta \cdot \mathsf{CB}\phi \cdot \zeta \cdot \alpha \cdot \pi \succ \zeta \star (\phi)^n\alpha \cdot \pi$.

Proof. Same as Lemma 15. □

3. The characteristic Boolean algebra $\beth 2$

3.1. *Function symbols*

Let us now define the principal function symbols commonly used in the sequel. As explained before, they are defined in the ground model $\mathcal{M}$ and are immediately extended to the realizability model $\mathcal{N}_{\mathcal{A}}$.

- The projections $pr_0 : X \times Y \to X$ and $pr_1 : X \times Y \to Y$ defined by:
  $pr_0(x, y) = x$, $pr_1(x, y) = y$
  give, in $\mathcal{N}_{\mathcal{A}}$, a *bijection* from $\beth(X \times Y)$ onto $\beth X \times \beth Y$.
- We define, in $\mathcal{M}$, the function app $: Y^X \times X \to Y$ (read *application*) by setting:
  $\mathrm{app}(f, x) = f(x)$ for $f \in Y^X$ and $x \in X$.
  This gives, in $\mathcal{N}_{\mathcal{A}}$, an application app $: \beth(Y^X) \times \beth X \to \beth Y$.
  We shall write $f(x)$ for $\mathrm{app}(f, x)$.

Theorem 17.
If $X \neq \varnothing$, then $\mathsf{I} \Vdash \forall f^{\beth(Y^X)} \forall g^{\beth(Y^X)} \left(\forall x^{\beth X} (\mathrm{app}(f, x) = \mathrm{app}(g, x)) \to f = g \right)$.
In other words: $\mathsf{I} \Vdash$ the function *app* gives an injection from $\beth(Y^X)$ into $(\beth Y)^{\beth X}$.

Proof. Let $f, g \in Y^X$, $\xi \Vdash \forall x^{\beth X}(\mathrm{app}(f, x) \neq \mathrm{app}(g, x) \to \bot)$ and $\pi \in \|f \neq g \to \bot\|$.
We must show $\xi \star \pi \in \perp\!\!\!\perp$. We choose $a \in X$; then $\xi \Vdash (f(a) \neq g(a) \to \bot)$.
If $f = g$, we have $\|f(a) \neq g(a) \to \bot\| = \|f \neq g \to \bot\| = \|\bot \to \bot\|$. Hence the result.
If $f \neq g$, we could choose a such that $f(a) \neq g(a)$.
Then, $\|f(a) \neq g(a) \to \bot\| = \|f \neq g \to \bot\| = \|\top \to \bot\|$. Hence the result. □

- Let sp $: \mathcal{M} \to \{0, 1\}$ (read *support*) the unary function symbol defined by:
  $\mathrm{sp}(\varnothing) = 0$; $\mathrm{sp}(x) = 1$ if $x \neq \varnothing$.
  In the realizability model $\mathcal{N}_{\mathcal{A}}$, we have sp $: \mathcal{N} \to \beth 2$.
- Let $\mathsf{P} : \{0, 1\} \times \mathcal{M} \to \mathcal{M}$ (read *projection*) the binary function symbol defined by:
  $\mathsf{P}(0, x) = \varnothing$; $\mathsf{P}(1, x) = x$.
  In the realizability model $\mathcal{N}_{\mathcal{A}}$, we have $\mathsf{P} : \beth 2 \times \mathcal{N} \to \mathcal{N}$.
  In the following, we shall write ix instead of $\mathsf{P}(i, x)$.
  When t, u are ℓ-terms with values in $\beth 2$, we write $t \leqslant u$ for $t \wedge u = t$.

Proposition 18.

i. $\mathsf{I} \Vdash \forall i^{\beth 2} \forall x (i(jx) = (i \wedge j)x)$.

https://doi.org/10.1017/S0960129516000050 Published online by Cambridge University Press

ii. $\mathsf{I} \Vdash \forall i^{\beth 2} \forall x (ix = x \rightleftharpoons \mathrm{sp}(x) \leqslant i)$.
iii. If $\varnothing \in E$, then $\mathsf{I} \Vdash \forall i^{\beth 2} \forall x^{\beth E} (ix \,\varepsilon\, \beth E)$.
iv. If $f : \mathcal{M}^n \to \mathcal{M}$ is a function symbol such that $f(\varnothing, \ldots, \varnothing) = \varnothing$, then:
$\mathsf{I} \Vdash \forall j^{\beth 2} \forall x_1 \ldots \forall x_n (jf(x_1, \ldots, x_n) = f(jx_1, \ldots, jx_n))$.
v. $\mathsf{I} \Vdash \forall i^{\beth 2} \forall x \left(i \neq 1 \to \forall y (y \not\in ix)\right)$ and therefore $\mathsf{K}^2 \mathsf{I} \Vdash \forall i^{\beth 2} \forall x \left(i \neq 1 \to \forall y (y \not\in ix)\right)$.

Proof. Trivial. □

Remark. Proposition 18(v) shows that, in the realizability model $\mathcal{N}$, every non-empty individual has support 1.

Because of property (iv), we shall define, as far as possible, each function symbol f in $\mathcal{M}$, so that to have $f(\varnothing, \ldots, \varnothing) = \varnothing$.

- Thus, let us change the ordered pair (x, y) by setting $(\varnothing, \varnothing) = \varnothing$. Then, we have: $\mathsf{I} \Vdash \forall i^{\beth 2} \forall x \forall y \, (i(x, y) = (ix, iy))$.
- We define the binary function symbol $\sqcup : \mathcal{M}^2 \to \mathcal{M}$ by setting: $a \sqcup b = a \cup b$.

Remark. The extension to $\mathcal{N}$ of this operation is neither the union for the $\in$-membership, nor the union for the ε-membership.

The operation $\beth_i$
Let $E \in \mathcal{M}$ be such that $\varnothing \in E$. In $\mathcal{M}$, we define $\beth_i E$ for $i \in 2$ by setting:
$\beth_0 E = \beth\{\varnothing\} = \{\varnothing\} \times \Pi$; $\beth_1 E = \beth E = E \times \Pi$.
In this way, we have now defined $\beth_i E$ in $\mathcal{N}$, for every $i \,\varepsilon\, \beth 2$.

Proposition 19.

i. $\mathsf{I} \Vdash \forall i^{\beth 2} \forall x \forall y (i(x \sqcup y) = ix \sqcup iy)$.
ii. $\mathsf{I} \Vdash \forall i^{\beth 2} \forall j^{\beth 2} \forall x ((i \vee j)x = ix \sqcup jx)$.
iii. $\mathsf{I} \Vdash \forall i^{\beth 2} \forall j^{\beth 2} \forall x \forall y \forall z (i \wedge j = 0, z = ix \sqcup jy \hookrightarrow iz = ix)$.
$\mathsf{I} \Vdash \forall i^{\beth 2} \forall j^{\beth 2} \forall x \forall y \forall z (i \wedge j = 0, z = ix \sqcup jy \hookrightarrow jz = jy)$.
iv. $\mathsf{I} \Vdash \forall i^{\beth 2} \forall j^{\beth 2} \forall x^{\beth E} \forall y^{\beth E} \forall z \left(i \wedge j = 0, z = ix \sqcup jy \hookrightarrow z \,\varepsilon\, \beth_{i \vee j} E\right)$.

Proof. Trivial. □

Proposition 20.
If $\varnothing \in E, E'$, the following formulas are realized:

i. $\beth_i E$ increases with i. In particular, $\beth_i E \subseteq \beth E$.
ii. The ε-elements of $\beth_i E$ are the ix for $x \,\varepsilon\, \beth E$.
iii. The ε-elements of $\beth_i E$ are those of $\beth E$ such that $\mathrm{sp}(x) \leqslant i$.
iv. The only ε-element common to $\beth_i E$ and $\beth_{1-i} E$ is $\varnothing$.
v. If $i \wedge j = 0$, then the application $x \mapsto (ix, jx)$ is a bijection from $\beth_{i \vee j} E$ onto $\beth_i E \times \beth_j E$. The inverse function is $(x, y) \mapsto x \sqcup y$.
vi. $\beth_i (E \times E') = \beth_i E \times \beth_i E'$.

Proof. We check immediately (i), (ii), (iii), (iv) below:

i. $\mathsf{I} \Vdash \forall i^{\beth 2} \forall j^{\beth 2} \forall x (i \wedge j = i \hookrightarrow (x \not\in \beth_j E \to x \not\in \beth_i E))$.

https://doi.org/10.1017/S0960129516000050 Published online by Cambridge University Press

ii. $\mathsf{I} \Vdash \forall i^{\rightarrow 2}\forall x^{\rightarrow E}(ix\,\varepsilon\,\rightarrow_i E)$; $\mathsf{I} \Vdash \forall i^{\rightarrow 2}\forall x^{\rightarrow E}(ix \neq x \rightarrow x \notin \rightarrow_i E)$.

iii. $\mathsf{I} \Vdash \forall i^{\rightarrow 2}\forall x^{\rightarrow E}(x \notin \rightarrow_i E \rightarrow i\wedge \mathrm{sp}(x) \neq \mathrm{sp}(x))$; $\mathsf{I} \Vdash \forall i^{\rightarrow 2}\forall x^{\rightarrow E}(i\wedge \mathrm{sp}(x) \neq \mathrm{sp}(x) \rightarrow x \notin \rightarrow_i E)$.

iv. $\mathsf{I} \Vdash \forall i^{\rightarrow 2}\forall x^{\rightarrow E}\forall y^{\rightarrow E}(ix = (1-i)y \hookrightarrow ix = \varnothing)$.

v. By Proposition 19(ii), we have $ix \sqcup jx = (i\vee j)x = x$ if $x\,\varepsilon\,\rightarrow_{i\vee j} E$.
By Proposition 19(iii,iv), if $x, y\,\varepsilon\,\rightarrow E$, there exists $z\,\varepsilon\,\rightarrow_{i\vee j} E$ such that $iz = ix, jz = jy$, namely $z = ix \sqcup jy$.

vi. By Proposition 18(iv), we have $\mathsf{I} \Vdash \forall i^{\rightarrow 2}\forall x\forall y(i(x,y) = (ix, iy))$.

□

Proposition 21. Let $E, E' \in \mathcal{M}$ be such that $\varnothing \in E, E'$ and E is equipotent with E'. Then $\Vdash \forall i^{\rightarrow 2}(\rightarrow_i E$ is equipotent with $\rightarrow_i E')$.

Proof. Let ϕ be, in $\mathcal{M}$, a bijection from E onto E', such that $\phi(\varnothing) = \varnothing$. Then ϕ is, in $\mathcal{N}$, a bijection from $\rightarrow E$ onto $\rightarrow E'$. But we have immediately: $\mathsf{I} \Vdash \forall i^{\rightarrow 2}\forall x^{\rightarrow E}(\phi(ix) = i\phi(x))$. This shows that ϕ is a bijection from $\rightarrow_i E$ onto $\rightarrow_i E'$. □

3.2. *Some general theorems*

Theorems 22 to 30, which are shown in this section, are valid *in every realizability model.*

In the ground model $\mathcal{M}$, which satisfies ZF + V = L, we denote by κ the cardinal of $\Lambda \cup \Pi \cup \mathbb{N}$ (which we shall also call the *cardinal of the algebra* $\mathcal{A}$) and by $\kappa_+ = \mathcal{P}(\kappa)$ the power set of κ.

Theorem 22.
Let $\forall \vec{x}\forall y F[\vec{x}, y]$ be a closed formula of ZF_ε with parameters in $\mathcal{M}$ (with $\vec{x} = (x_1, \ldots, x_n)$). Then, there exists in $\mathcal{M}$, a functional $f_F : \kappa \times \mathcal{M}^n \to \mathcal{M}$ such that:

i. If $\vec{a}, b \in \mathcal{M}$ and $\xi \Vdash F[\vec{a}, b]$, then there exists $\alpha \in \kappa$ such that $\xi \Vdash F[\vec{a}, f_F(\alpha, \vec{a})]$.

ii. $\mathsf{C}\,\mathsf{I} \Vdash \forall \vec{x}\forall y \left(F[\vec{x}, y] \rightarrow \exists v^{\rightarrow \kappa} F[\vec{x}, f_F(v, \vec{x})]\right)$.

Proof.

i. Let $\xi \mapsto \alpha_\xi$ be an injection from Λ into κ. Using the principle of choice in $\mathcal{M}$ (which satisfies V = L), we can define a functional $f_F : \kappa \times \mathcal{M}^n \to \mathcal{M}$ such that, in $\mathcal{M}$, we have $\forall \vec{x}\forall y(\forall \xi \in \Lambda)\left(\xi \Vdash F[\vec{x}, y] \Rightarrow \xi \Vdash F[\vec{x}, f_F(\alpha_\xi, \vec{x})]\right)$.

ii. Let $\xi \Vdash F[\vec{a}, b]$, $\eta \Vdash \forall v^{\rightarrow \kappa} \neg F[\vec{a}, f_F(v, \vec{a})]$ and $\pi \in \Pi$. Therefore, we have: $\eta \Vdash \neg F[\vec{a}, f_F(\alpha_\xi, \vec{a})]$; now, by definition of f_F, we have $\xi \Vdash F[\vec{a}, f_F(\alpha_\xi, \vec{a})]$. Therefore $\eta \star \xi \cdot \pi \in \perp\!\!\!\perp$ and $\mathsf{C}\,\mathsf{I} \star \xi \cdot \eta \cdot \pi \in \perp\!\!\!\perp$.

□

Remark. The function f_F is a kind of *weak choice function* for the relation $F[\vec{x}, y]$; given $\vec{x}$, it does not give exactly a witness y such that $F[\vec{x}, y]$, but a family indexed by $\rightarrow \kappa$ in which there is such a witness.

https://doi.org/10.1017/S0960129516000050 Published online by Cambridge University Press

Subsets of $\beth\kappa_+$

Theorem 23. Let $\forall x\forall y\forall z\, F[x,y,z]$ be a closed formula of ZF_ε, with parameters in $\mathcal{M}$. Then, there exists, in $\mathcal{M}$, a functional $\beta_F : \mathcal{M} \to \kappa_+$ such that:
$\mathsf{W} \Vdash \forall z(\forall x\forall y\forall y'(F[x,y,z], F[x,y',z] \to y = y')$
$\to \forall i^{\beth 2}\forall x(F[x, i\beta_F(z), z] \to \mathrm{sp}(x) \geqslant i))$.

Proof. By Theorem 22(i), there exists, in $\mathcal{M}$, a functional $g : \kappa \times \mathcal{M}^2 \to \mathcal{M}$ such that:
(*) If $a,b,c \in \mathcal{M}$ and $\xi \Vdash F[a,b,c]$, there exists $\alpha \in \kappa$ such that $\xi \Vdash F[a, g(\alpha,a,c), c]$.

Using the principle of choice in $\mathcal{M}$, we define a functional $\beta_F : \mathcal{M} \to \kappa_+$ such that:
for every $\alpha \in \kappa$ and $c \in \mathcal{M}$, we have $\beta_F(c) \neq g(\alpha, \varnothing, c)$.
This is possible since κ_+ is of cardinal $> \kappa$.

Now let: $a,c \in \mathcal{M}$, $i \in \{0,1\}$, $\phi \Vdash \forall x\forall y\forall y'(F[x,y,c], F[x,y',c], y \neq y' \to \bot)$,
$\xi \Vdash F[a, i\beta_F(c), c]$, $\eta \Vdash \mathrm{sp}(a)i \neq i$ and $\pi \in \Pi$.

We must show that $\mathsf{W} \star \phi \cdot \xi \cdot \eta \cdot \pi \in \perp\!\!\!\perp$, that is $\phi \star \xi \cdot \xi \cdot \eta \cdot \pi \in \perp\!\!\!\perp$.
We set $b = i\beta_F(c)$ and therefore, we have $\xi \Vdash F[a,b,c]$.
Thus, by (*), we have $\xi \Vdash F[a, g(\alpha,a,c), c]$ for some $\alpha \in \kappa$.

Let us show that $\|b \neq g(\alpha,a,c)\| \subset \|\mathrm{sp}(a)i \neq i\|$; there are three possible cases:
If $i = 0$, then $\|\mathrm{sp}(a)i \neq i\| = \|0 \neq 0\| = \Pi$, hence the result.
If $i = 1$ and $a \neq \varnothing$, then $\|\mathrm{sp}(a)i \neq i\| = \|1 \neq 1\| = \Pi$, hence the result.
If $i = 1$ and $a = \varnothing$, then:
$\|b \neq g(\alpha,a,c)\| = \|i\beta_F(c) \neq g(\alpha,a,c)\| = \|\beta_F(c) \neq g(\alpha,\varnothing,c)\| = \|\top\| = \varnothing$, by definition of β_F, hence the result.

It follows that $\eta \Vdash b \neq g(\alpha,a,c)$. Now, we have seen that:
$\xi \Vdash F[a,b,c]$ and $\xi \Vdash F[a, g(\alpha,a,c), c]$.
Therefore, by hypothesis on ϕ, we have $\phi \star \xi \cdot \xi \cdot \eta \cdot \pi \in \perp\!\!\!\perp$. □

Remark. Theorem 23 is crucial for the applications in the sequel. A perhaps more intuitive reformulation is given in Corollary 24(i).

Corollary 24. If $\varnothing \in E$, then the following formulas are realized:
i. $\forall i^{\beth 2}$(there is no surjection from $\bigcup\{\beth_j E \;;\; j \varepsilon \beth 2, j \not\geqslant i\}$ onto $\beth_i\kappa_+$).
ii. $\forall i^{\beth 2}\forall j^{\beth 2}$(if there exists a surjection from $\beth_j E$ onto $\beth_i\kappa_+$ then $j \geqslant i$).
iii. $\forall i^{\beth 2}\forall j^{\beth 2}(i, j \neq 0, i \wedge j = 0$
$\to$ (there is no surjection from $\beth_i\kappa_+ \uplus \beth_j\kappa_+$ onto $\beth_i\kappa_+ \times \beth_j\kappa_+$)).

Remarks. $\uplus$ is the symbol for *disjoint union.*
The notation $\bigcup\{\beth_j E \;;\; j \varepsilon \beth 2, j \not\geqslant i\}$ denotes any individual X of $\mathcal{N}$ such that:
$\mathcal{N} \models \forall x(x \varepsilon X \leftrightarrow \exists j^{\beth 2}(j \not\geqslant i \wedge x \varepsilon \beth_j E))$.

Proof.

i. We apply Theorem 23, with the formula $F[x,y,z] \equiv (x,y) \varepsilon z$.
In the realizability model $\mathcal{N}$, we have $\beta_F : \mathcal{N} \to \beth\kappa_+$.
Let z_0 be, in $\mathcal{N}$, a surjective function onto $\beth_i\kappa_+$.
We have $\beta_F(z_0) \varepsilon \beth\kappa_+$, and therefore $i\beta_F(z_0) \varepsilon \beth_i\kappa_+$.

https://doi.org/10.1017/S0960129516000050 Published online by Cambridge University Press

If x_0 is such that $(x_0, i\beta_F(z_0))\,\varepsilon\, z_0$, then $\mathrm{sp}(x_0) \geqslant i$ by Theorem 23.
Therefore, we have $x_0\,\varepsilon\, \rrbracket_j E \Rightarrow j \geqslant i$, by Proposition 20(iii).

ii. It is a trivial consequence of (i).

iii. We take $E = \kappa_+$; since $i, j \neq 0, i \wedge j = 0$, we have $i, j \not\geqslant i \vee j$; by (i), there is no surjection from $\rrbracket_i \kappa_+ \cup \rrbracket_j \kappa_+$ onto $\rrbracket_{i\vee j}\kappa_+$.
Now, since $i \wedge j = 0$, $\rrbracket_{i\vee j}\kappa_+$ is equipotent with $\rrbracket_i\kappa_+ \times \rrbracket_j\kappa_+$ by Proposition 20(v).
Moreover, $\varnothing$ is the only ε-element common to $\rrbracket_i\kappa_+$ and $\rrbracket_j\kappa_+$ by proposition20(iv).
But these sets contain a countable subset by Theorem 26. It follows that $\rrbracket_i\kappa_+ \cup \rrbracket_j\kappa_+$ is equipotent with $\rrbracket_i\kappa_+ \uplus \rrbracket_j\kappa_+$.

□

Theorem 25. There is a function symbol f such that $\Vdash (\forall y \subseteq \rrbracket\kappa)(\exists x\,\varepsilon\, \rrbracket\kappa_+)(y \simeq f(x))$.
Roughly: $\Vdash (f$ is a surjection from $\rrbracket\kappa_+$ onto $2^{\rrbracket\kappa})$.

Reminder. $x \subseteq y$ is $\forall z(z \notin y \to z \notin x)$; $x \simeq y$ is $x \subseteq y \wedge y \subseteq x$.

Proof. In the ground model $\mathcal{M}$, we have $\kappa \geqslant \mathrm{card}(\Pi)$; thus, there exists a bijection f from $\kappa_+ = 2^\kappa$ onto $\mathcal{P}(\kappa \times \Pi)$. Therefore, f is a function symbol which is a bijection from $\rrbracket\kappa_+$ onto $\rrbracket\mathcal{P}(\kappa\times\Pi)$.
In the ground model $\mathcal{M}$, we define $\phi : \mathcal{M} \to \mathcal{P}(\kappa\times\Pi)$ by $\phi(x) = x \cap (\kappa \times \Pi)$.
In $\mathcal{N}$, we have $\phi : \mathcal{N} \to \rrbracket\mathcal{P}(\kappa\times\Pi)$; $\Vdash \forall x(\phi(\phi(x)) = \phi(x))$.

Now, we check immediately that:

i. $\Vdash \forall v \forall x^{\rrbracket\mathcal{P}(\kappa\times\Pi)}(v \notin \rrbracket\kappa \to v \notin x)$ because $\|v \notin a\| \subset \|v \notin \rrbracket\kappa\|$ for all $a \in \mathcal{P}(\kappa\times\Pi)$.

ii. $\Vdash \forall x \forall v^{\rrbracket\kappa}(v \notin x \rightleftharpoons v \notin \phi(x))$ because $\|v \notin a\| = \|v \notin \phi(a)\|$ for all $v \in \kappa$.

From (i), it follows that $\rrbracket\mathcal{P}(\kappa\times\Pi)$ is, in $\mathcal{N}$, a set of ε-subsets of $\rrbracket\kappa$;
from (ii), it follows that it contains at least one representative for each equivalence class for $\simeq$.
It follows that f has the desired properties. □

Theorem 26. Let $E \in \mathcal{M}$ be infinite and such that $\varnothing \in E$. Then we have:
$\Vdash \forall i^{\rrbracket 2}(i \neq 0 \to$ there exists an injection from $\mathbb{N}$ into $\rrbracket_i E)$.

Proof. In $\mathcal{M}$, let $\phi : \mathbb{N} \to (E \setminus \{\varnothing\})$ be injective. In $\mathcal{N}$, we have $\phi : \rrbracket\mathbb{N} \to \rrbracket E$.
The desired function is $n \mapsto i\phi(n)$. Indeed, we have:
$\Vdash \forall i^{\rrbracket 2} \forall m^{\rrbracket\mathbb{N}} \forall n^{\rrbracket\mathbb{N}}(i \neq 0 \to i\phi(m+n+1) \neq i\phi(m))$.
This shows that the restriction of this function to $\mathbb{N}_{\mathcal{A}}$ (the set of integers of $\mathcal{N}_{\mathcal{A}}$) is injective. □

Theorem 27. $\Vdash \forall i^{\rrbracket 2}\,(i \neq 0, i \neq 1 \to (\rrbracket\kappa_+$ cannot be well ordered$))$.

Proof. Let $i \in \rrbracket 2, i \neq 0, 1$; then, $\rrbracket_i\kappa_+$ and $\rrbracket_{1-i}\kappa_+$ are infinite (Theorem 26) and $\subseteq \rrbracket\kappa_+$ by Proposition 20(i). But there exists no surjection from $\rrbracket_i\kappa_+$ onto $\rrbracket_{1-i}\kappa_+$, neither from $\rrbracket_{1-i}\kappa_+$ onto $\rrbracket_i\kappa_+$, by Corollary 24. □

Remark. It is probably useful to make clear what a well ordering is in this non-extensional context. The formula (R is a strict well ordering on X) is the conjunction of:

https://doi.org/10.1017/S0960129516000050 Published online by Cambridge University Press

$(\forall x, y, z \,\varepsilon\, X)((x, y) \,\varepsilon\, R, (y, z) \,\varepsilon\, R \to (x, z) \,\varepsilon\, R)$, $(\forall x, y \,\varepsilon\, X)(x \neq y \leftrightarrow (x, y) \,\varepsilon\, R \vee (y, x) \,\varepsilon\, R)$
(i.e. R is a strict total order); $(\forall Y \subseteq X)(\forall x \,\varepsilon\, Y)(\exists y \,\varepsilon\, Y)(\forall z \,\varepsilon\, Y)((z, y) \notin R)$ (i.e. R is well founded).
Theorem 27 says that, if the Boolean algebra $\beth 2$ is not trivial, then $\beth \kappa_+$ is not well orderable. On the other hand, it can be shown that, if this Boolean algebra is trivial, then the realizability model $\mathcal{N}$ is an extension by forcing of the ground model $\mathcal{M}$. In this case, it is well known that $\mathcal{N}$ itself can be well ordered (we suppose that the ground model $\mathcal{M}$ satisfies ZF + V = L).
Note also that, even if there is a non-well-orderable set in $\mathcal{N}$, we do not know if the underlying model of ZF satisfies AC or not.

A strict order on $\beth \kappa_+$ A binary relation $<$ on X is a *strict order* if it is transitive $(x < y, y < z \Rightarrow x < z)$ and antireflexive $(x \not< x)$. This strict order is called *total* if we have:

$$x < y \text{ or } y < x \text{ or } x = y.$$

If $(X_0, <_0), (X_1, <_1)$ are two strictly ordered sets, then the *strict order product* $<$ on $X_0 \times X_1$ is defined by: $(x_0, x_1) < (y_0, y_1) \Leftrightarrow x_0 < y_0$ and $x_1 < y_1$.

Lemma 28. The strict order product of $<_0, <_1$ is well founded if and only if one of the strict orders $<_0, <_1$ is well founded.

Proof. Proof of $\Rightarrow$: by contradiction; if $<_0$ and $<_1$ are not well founded, we have:
$\forall y_0\,(\forall x_0(x_0 <_0 y_0 \to F_0[x_0]) \to F_0[y_0])$; $\neg F_0[b_0]$;
$\forall y_1\,(\forall x_1(x_1 <_1 y_1 \to F_1[x_1]) \to F_1[y_1])$; $\neg F_1[b_1]$;
for some formulas F_0, F_1 and some individuals b_0, b_1. It follows that:
$\forall y_0 \forall y_1\,(\forall x_0 \forall x_1(x_0 <_0 y_0, x_1 <_1 y_1 \to G[x_0, x_1]) \to G[y_0, y_1])$; $\neg G[b_0, b_1]$
where $G[x_0, x_1] \equiv F_0[x_0] \vee F_1[x_1]$.

Proof of $\Leftarrow$: suppose that $<_0$ is well founded and let $G[x_0, x_1]$ be any formula.
Let $F[x_0] \equiv \forall x_1 G[x_0, x_1]$. We have to prove $\forall y_0 \forall y_1 G[y_0, y_1]$, i.e. $\forall y_0 F[y_0]$ with the hypothesis $\forall y_0 \forall y_1\,(\forall x_0 \forall x_1(x_0 <_0 y_0, x_1 <_1 y_1 \to G[x_0, x_1]) \to G[y_0, y_1])$. But this implies:
$\forall y_0\,(\forall x_0(x_0 <_0 y_0 \to F[x_0]) \to F[y_0])$ and the result follows, because $<_0$ is well founded. □

We denote by $\lhd$ a strict well ordering on κ_+, in $\mathcal{M}$; we suppose that its least element is $\varnothing$ and that the cardinal of each proper initial segment is $\leqslant \kappa$.

This gives a binary function from κ_+^2 into $\{0, 1\}$, denoted by $(x \lhd y)$, which is defined as follows: $(x \lhd y) = 1 \Leftrightarrow x \lhd y$.
We can extend it to the realizability model $\mathcal{N}_{\mathcal{A}}$, which gives a function from $(\beth \kappa_+)^2$ into $\beth 2$.

Lemma 29. The following propositions are realized:
If $i \,\varepsilon\, \beth 2, i \neq 0$, then $(x \lhd y) = i$ is a strict ordering of $\beth_i \kappa_+$, which we denote by $\lhd_i$.
If i is an atom of the Boolean algebra $\beth 2$, then this ordering is total.

Proof. We have immediately:
i. $\mathbb{1} \Vdash \forall x^{\beth \kappa_+} \forall y^{\beth \kappa_+} \forall z^{\beth \kappa_+}((x \lhd y) \wedge (y \lhd z) \leqslant (x \lhd z))$; $\mathbb{1} \Vdash \forall x^{\beth \kappa_+}((x \lhd x) = 0)$.

https://doi.org/10.1017/S0960129516000050 Published online by Cambridge University Press

ii. $\mathsf{I} \Vdash \forall i^{\beth 2} \forall x^{\beth \kappa_+} \forall y^{\beth \kappa_+} ((ix \lhd iy) \leqslant i)$.

iii. $\mathsf{I} \Vdash \forall x^{\beth \kappa_+} \forall y^{\beth \kappa_+} ((x \lhd y) = 0, (y \lhd x) = 0 \hookrightarrow x = y)$.

It follows from (i) that, if $i \neq 0$, then $(x \lhd y) \geqslant i$ is a strict ordering relation on $\beth \kappa_+$. It follows from (ii), that this relation, restricted to $\beth_i \kappa_+$, is equivalent to $(x \lhd y) = i$. Finally, it follows from (iii), that the relation $(x \lhd y) = i$, restricted to $\beth_i \kappa_+$, is total when i is an atom of $\beth 2$. □

Lemma 30. The following propositions are realized:

i. $\forall i^{\beth 2}$(the application $x \mapsto (ix, (1-i)x)$ is an isomorphism of strictly ordered sets from $(\beth \kappa_+, \lhd)$ onto $(\beth_i \kappa_+, \lhd_i) \times (\beth_{1-i} \kappa_+, \lhd_{1-i})$).

ii. $\forall i^{\beth 2}$(either $\beth_i \kappa_+$ or $\beth_{1-i} \kappa_+$ is a well-founded ordered set).

Proof.

i. It follows from Proposition 20(v), that the application $x \mapsto (ix, (1-i)x)$ is a bijection from $\beth \kappa_+$ onto $\beth_i \kappa_+ \times \beth_{1-i} \kappa_+$. In fact, it is an *isomorphism* of ordered sets, since we have: $\mathsf{I} \Vdash \forall i^{\beth 2} \forall x^{\beth \kappa_+} \forall y^{\beth \kappa_+} ((x \lhd y) = (ix \lhd iy) \vee ((1-i)x \lhd (1-i)y))$ and therefore:
$\Vdash \forall i^{\beth 2} \forall x^{\beth \kappa_+} \forall y^{\beth \kappa_+} ((x \lhd y) = 1 \leftrightarrow (ix \lhd iy) = i \wedge ((1-i)x \lhd (1-i)y) = 1 - i)$.

ii. By Theorem 12, the relation $(x \lhd y) = 1$ is well founded on $\beth \kappa_+$. Thus, the result follows immediately from (i) and Lemma 28.

□

3.3. $\beth \kappa$ *countable*

In this section, we consider some consequences of the hypothesis: ($\beth \kappa$ is countable).

Non-extensional choice (NEAC) and dependent choice (DC)
The formula $\forall x \forall y \forall y' ((x, y) \varepsilon f, (x, y') \varepsilon f \rightarrow y = y')$ will be written Func(f)
(read: f *is a function*).
We recall that $x \subseteq y$ is the formula $\forall z (z \not\varepsilon y \rightarrow z \not\varepsilon x)$.
The formula $\forall z \exists f \left(f \subseteq z \wedge \text{Func}(f) \wedge \forall x \forall y \exists y' ((x, y) \varepsilon z \rightarrow (x, y') \varepsilon f) \right)$
is called the *non-extensional axiom of choice* and denoted by NEAC.

It is easily shown (Krivine 2012) that $\text{ZF}_\varepsilon + \text{NEAC} \vdash \text{DC}$ (axiom of dependent choice). On the other hand, we have built, in Krivine (2012), a model of $\text{ZF}_\varepsilon + \text{NEAC} + \neg\text{AC}$; and other such models will be given in the present paper. In all these models, $\mathbb{R}$ is not well orderable.

Theorem 31.
There exists a closed c-term H such that $\mathsf{H} \Vdash (\beth \kappa \text{ is countable} \rightarrow \text{NEAC})$.

Proof. We apply Theorem 22(ii) to the formula $(x, y) \varepsilon z$. We get a function symbol g such that $\mathsf{C}\mathsf{I} \Vdash \forall x \forall y \forall z ((x, y) \varepsilon z \rightarrow \exists v^{\beth \kappa} (x, g(v, x, z)) \varepsilon z)$.
Therefore, it suffices to prove NEAC in ZF_ε, by means of this formula and the additional hypothesis: ($\beth \kappa$ is countable). Now, from this hypothesis, it follows that there exists a strict well ordering $<$ on $\beth \kappa$. Then, we can define the desired function f by means of the

https://doi.org/10.1017/S0960129516000050 Published online by Cambridge University Press

comprehension scheme:
$(x,y)\,\varepsilon\, f \leftrightarrow (x,y)\,\varepsilon\, z \wedge \exists v^{\beth\kappa}(y = g(v,x,z) \wedge \forall\alpha^{\beth\kappa}(\alpha < v \rightarrow (x, g(\alpha,x,z))\notin z)$.
Intuitively, $f(x) = g(v,x,z)$ for the least $v\,\varepsilon\,\beth\kappa$ such that $(x,g(v,x,z))\,\varepsilon\, z$. □

Subsets of $\mathbb{R}$

Theorem 32. $\Vdash$ ($\beth\kappa$ is countable) $\rightarrow$
every bounded above subset of the ordered set $(\beth\kappa_+, \lhd)$ is countable.

Proof. Every proper initial segment of the well ordering $\lhd$ on κ_+ is of cardinal κ. Thus, there exists a function $\phi : \kappa\times\kappa_+ \rightarrow \kappa_+$ such that, for each $x \in \kappa_+, x \neq \varnothing$, the function $\alpha \mapsto \phi(\alpha,x)$ is a surjection from κ onto $\{y \in \kappa_+ \,;\, y \lhd x\}$. Then, we have immediately:
$\mathbf{I} \Vdash \forall x^{\beth\kappa_+}\forall y^{\beth\kappa_+}\Big((y \lhd x) = 1 \hookrightarrow (\forall\alpha^{\beth\kappa}(y \neq \phi(\alpha,x)) \rightarrow \bot)\Big)$.
This shows that, in $\mathcal{N}$, there exists a surjection from $\beth\kappa$, onto every subset of $\beth\kappa_+$ which is bounded from above for the strict ordering $\lhd$.
Thus, all these subsets of $\beth\kappa_+$ are countable, since $\beth\kappa$ is. □

Theorem 33. $\Vdash$ ($\beth\kappa$ is countable) $\rightarrow$ there exists an injection from $\beth\kappa_+$ into $\mathbb{R}$.

Proof. We have obviously $\Vdash$ ($\beth\kappa$ is countable $\rightarrow \beth 2$ is countable), and therefore:
$\Vdash$ ($\beth\kappa$ is countable $\rightarrow (\beth 2)^{\beth\kappa}$ is equipotent to $\mathbb{R}$).
Now, by Theorem 17, we have: $\Vdash$ (there is an injection from $\beth\kappa_+ = \beth(2^\kappa)$ into $(\beth 2)^{\beth\kappa}$). □

Theorem 34. The following formula is realized:
($\beth\kappa$ is countable) $\rightarrow$ there exists an application $i \mapsto X_i$ from the countable Boolean algebra $\beth 2$ into $\mathcal{P}(\mathbb{R})$ such that:

i. $X_0 = \{\varnothing\}$; $i \neq 0 \rightarrow X_i$ is uncountably infinite;
ii. $X_i \times X_i$ is equipotent with X_i;
iii. $X_i \cap X_j = X_{i\wedge j}$ and therefore $i \leqslant j \rightarrow X_i \subset X_j$;
iv. $i\wedge j = 0 \rightarrow X_{i\vee j}$ is equipotent with $X_i \times X_j$;
v. there exists a surjection from X_1 onto $\mathbb{R}$.
vi. if J is a subset of $\beth 2$ and if there is a surjection from $\bigcup_{j\varepsilon J} X_j$ onto X_i, then $i \leqslant j$ for some $j\,\varepsilon\, J$.
vii. if there is a surjection from X_j onto X_i, then $i \leqslant j$;
viii. if $i, j \neq 0, i\wedge j = 0$, then there is no surjection from $X_i \uplus X_j$ onto $X_i \times X_j$.

Proof. For each $i\,\varepsilon\,\beth 2$, let us denote by X_i the image of $\beth_i\kappa_+$ by the injection from $\beth\kappa_+$ into $\mathbb{R}$, given by Theorem 33.

i. The fact that X_i is infinite for $i \neq 0$ is a consequence of Theorem 26.
If $i = 1$, X_i is uncountable by (v). If $i \neq 0, 1$ and X_i is countable, then X_{1-i} is infinite and thus, there exists a surjection from X_{1-i} onto X_i. This contradicts Corollary 24.
ii. By Proposition 20(vi), $\beth_i\kappa_+ \times \beth_i\kappa_+$ is equipotent with $\beth_i(\kappa_+^2)$, thus also with $\beth_i\kappa_+$ by Proposition 21.
iii. If $a\,\varepsilon\,\beth_i\kappa_+$ and $a\,\varepsilon\,\beth_j\kappa_+$, then $ia = a$, and therefore $(i\wedge j)a = ja = a$.
iv. This is Proposition 20(v).

https://doi.org/10.1017/S0960129516000050 Published online by Cambridge University Press

v. Application of Theorem 25.
vi., vii), viii) Applications of Corollary 24.

□

Theorem 34 is interesting only if the countable Boolean algebra $\daleth 2$ is not trivial.
In this case, by Theorems 27 and 33, $\mathbb{R}$ *cannot be well ordered*; the underlying model of ZF does not satisfy AC.

4. Collapsing $\daleth\kappa$

In this section, we start with an *arbitrary* realizability algebra $\mathcal{A}$ and we build a new algebra $\mathcal{B}$ such that:

- $\mathcal{N}_{\mathcal{B}}$ realizes the formula: ($\daleth\kappa$ is countable).
- The (countable) Boolean algebra $\daleth 2$ of the model $\mathcal{N}_{\mathcal{B}}$ is elementarily equivalent (for the language of rings) to the algebra $\daleth 2$ of $\mathcal{N}_{\mathcal{A}}$.

In the sequel, we shall consider two interesting cases: $\daleth 2$ is atomless; $\daleth 2$ has four ε-elements.

4.1. *Extending a realizability algebra*

In the ground model $\mathcal{M}$, we consider a realizability algebra $\mathcal{A} = (\Lambda, \Pi, \Lambda \star \Pi, \perp\!\!\!\perp)$, the elementary combinators of which are denoted by $\mathsf{B}, \mathsf{C}, \mathsf{I}, \mathsf{K}, \mathsf{W}, \mathsf{cc}$ and the continuations k_π for $\pi \in \Pi$.

Let κ be an infinite cardinal of $\mathcal{M}$, $\kappa \geqslant \mathrm{card}(\Lambda \cup \Pi)$; we consider the tree (usually denoted as $\kappa^{<\omega}$) of functions, the domain of which is an integer, with values in κ.
Let P be the ordered set obtained by adding a least element $\mathbb{O}$ to this tree.
P is an inf-semi-lattice, the greatest element $\mathbf{1}$ of which is the void function $\varnothing$.
The greatest lower bound of $p, q \in P$, denoted by pq, is p (resp. q) if $p, q \neq \mathbb{O}$ and $q \subset p$ (resp. $p \subset q$). It is $\mathbb{O}$ in every other case.

Remark. $P \setminus \{\mathbb{O}\} = \kappa^{<\omega}$ is the ordered set used, in the method of forcing, to *collapse* (i.e. make countable) the cardinal κ.

We define a new realizability algebra $\mathcal{B}$ by setting:
$\mathbf{\Lambda} = \Lambda \times P$; $\mathbf{\Pi} = \Pi \times P$; $\mathbf{\Lambda} \star \mathbf{\Pi} = (\Lambda \star \Pi) \times P$;
$(\xi, p) \cdot (\pi, q) = (\xi \cdot \pi, pq)$; $(\xi, p) \star (\pi, q) = (\xi \star \pi, pq)$; $(\xi, p)(\eta, q) = (\mathsf{C}\xi\eta, pq)$.
$\mathbf{B} = (\mathsf{B}^*, \mathbf{1})$; $\mathbf{C} = (\mathsf{C}^*, \mathbf{1})$; $\mathbf{I} = (\mathsf{I}^*, \mathbf{1})$; $\mathbf{K} = (\mathsf{K}^*, \mathbf{1})$; $\mathbf{W} = (\mathsf{W}^*, \mathbf{1})$; $\mathbf{cc} = (\mathsf{cc}^*, \mathbf{1})$; $\mathbf{k}_{(\pi,p)} = (\mathsf{k}^*_\pi, p)$.

The combinators $\mathsf{B}^*, \mathsf{C}^*, \mathsf{I}^*, \mathsf{K}^*, \mathsf{W}^*, \mathsf{cc}^*$, and the continuations k^*_π are defined in such a way that we have, for every $\nu, \xi, \eta, \zeta \in \Lambda$ and $\pi \in \Pi$:

$\mathsf{B}^* \star \nu \cdot \xi \cdot \eta \cdot \zeta \cdot \pi \succgtr \xi \star \nu \cdot \mathsf{C}\eta\zeta \cdot \pi$;
$\mathsf{C}^* \star \nu \cdot \xi \cdot \eta \cdot \zeta \cdot \pi \succgtr \xi \star \nu \cdot \zeta \cdot \eta \cdot \pi$;
$\mathsf{I}^* \star \nu \cdot \xi \cdot \pi \succgtr \xi \star \nu \cdot \pi$;
$\mathsf{K}^* \star \nu \cdot \xi \cdot \eta \cdot \pi \succgtr \xi \star \nu \cdot \pi$;
$\mathsf{W}^* \star \nu \cdot \xi \cdot \eta \cdot \pi \succgtr \xi \star \nu \cdot \eta \cdot \eta \cdot \pi$;

https://doi.org/10.1017/S0960129516000050 Published online by Cambridge University Press

$\mathsf{k}_\pi^* \star v \cdot \xi \cdot \varpi \gg \xi \star v \cdot \pi$;
$\mathsf{cc}^* \star v \cdot \xi \cdot \pi \gg \xi \star v \cdot \mathsf{k}_\pi^* \cdot \pi$.
(reminder: the notation $\xi \star \pi \gg \xi' \star \pi'$ means $\xi \star \pi \notin \bot\!\!\!\bot \Rightarrow \xi' \star \pi' \notin \bot\!\!\!\bot$).

Therefore, we set:

$\mathsf{B}^* = \lambda n \lambda x \lambda y \lambda z (xn)(\mathsf{C})yz = \big((\mathsf{C})(\mathsf{BC})(\mathsf{C})(\mathsf{B})(\mathsf{BB})\mathsf{B}\big)\mathsf{C}$;
$\mathsf{C}^* = \lambda n \lambda x \lambda y \lambda z (x)nzy = (\mathsf{C})(\mathsf{B})\mathsf{C}$;
$\mathsf{I}^* = \lambda n \lambda x (x)n = \mathsf{C}\,\mathsf{I}$;
$\mathsf{K}^* = \lambda n \lambda x \lambda y (x)n$ or $(\mathsf{C})(\mathsf{B})\mathsf{K}$;
$\mathsf{W}^* = \lambda n \lambda x \lambda y (x)nyy = (\mathsf{C})(\mathsf{B})\mathsf{W}$;
$\mathsf{k}_\pi^* = \lambda n \lambda x (\mathsf{k}_\pi)(x)n = (\mathsf{C})(\mathsf{B})\mathsf{k}_\pi$;
$\mathsf{cc}^* = \lambda n \lambda x (\mathsf{cc}) \lambda k (xn) \lambda n \lambda x (k)(x)n$
$= \big((\mathsf{C})\big((\mathsf{C})\big((\mathsf{B})\big((\mathsf{B})(\mathsf{B})\mathsf{C}\big)\mathsf{C}\big)(\mathsf{C})(\mathsf{B})\big((\mathsf{B})(\mathsf{B})\big((\mathsf{B})(\mathsf{B})\mathsf{cc}\big)\mathsf{B}\big)\mathsf{B}\big)\mathsf{C}\big)\mathsf{B}$.

We define, in $\mathcal{M}$, a function symbol from $P \times \mathbb{N}$ into $\{0,1\}$, denoted by $(p \ll n)$, by setting: $(p \ll n) = 1 \Leftrightarrow p \neq \mathbb{O}$ and the domain of p is an integer $\leqslant n$.

We define $\bot\!\!\!\bot_{\mathcal{B}}$, that we shall denote also by $\bot\!\!\!\bot$, as follows:

$(\xi \star \pi, p) \in \bot\!\!\!\bot \Leftrightarrow (\forall n \in \mathbb{N})((p \ll n) = 1 \Rightarrow \xi \star \underline{n} \cdot \pi \in \bot\!\!\!\bot)$ for $p \in P$, $\xi \in \Lambda$ and $\pi \in \Pi$.
In particular, we have $(\xi \star \pi, \mathbb{O}) \in \bot\!\!\!\bot$ for any $\xi \in \Lambda, \pi \in \Pi$.

We check now that $\mathcal{B}$ is a realizability algebra.

- $(\xi, p)(\eta, q) \star (\pi, r) \notin \bot\!\!\!\bot \Rightarrow (\xi, p) \star (\eta, q) \cdot (\pi, r) \notin \bot\!\!\!\bot$:
  Suppose that $(\xi \star \eta \cdot \pi, pqr) \in \bot\!\!\!\bot$; we must show $(\mathsf{C}\xi\eta \star \pi, pqr) \in \bot\!\!\!\bot$ i.e. $\mathsf{C}\xi\eta \star \underline{n} \cdot \pi \in \bot\!\!\!\bot$ for $(pqr \ll n) = 1$. Now, we have $\mathsf{C}\xi\eta \star \underline{n} \cdot \pi \gg \xi \star \underline{n} \cdot \eta \cdot \pi$ which is in $\bot\!\!\!\bot$ by hypothesis.
- $(\mathsf{B}^*, \mathbf{1}) \star (\xi, p) \cdot (\eta, q) \cdot (\zeta, r) \cdot (\pi, s) \notin \bot\!\!\!\bot \Rightarrow (\xi, p) \star (\eta, q)(\zeta, r) \cdot (\pi, s) \notin \bot\!\!\!\bot$:
  Suppose that $(\xi, p) \star (\eta, q)(\zeta, r) \cdot (\pi, s) \in \bot\!\!\!\bot$ i.e. $(\xi \star \mathsf{C}\eta\zeta \cdot \pi, pqrs) \in \bot\!\!\!\bot$; we must show: $(\mathsf{B}^* \star \xi \cdot \eta \cdot \zeta \cdot \pi, pqrs) \in \bot\!\!\!\bot$ i.e. $\mathsf{B}^* \star \underline{n} \cdot \xi \cdot \eta \cdot \zeta \cdot \pi \in \bot\!\!\!\bot$ for $(pqrs \ll n) = 1$.
  Now, we have: $\mathsf{B}^* \star \underline{n} \cdot \xi \cdot \eta \cdot \zeta \cdot \pi \gg \xi \star \underline{n} \cdot \mathsf{C}\eta\zeta \cdot \pi$ which is in $\bot\!\!\!\bot$ by hypothesis.
- $(\mathsf{C}^*, \mathbf{1}) \star (\xi, p) \cdot (\eta, q) \cdot (\zeta, r) \cdot (\pi, s) \notin \bot\!\!\!\bot \Rightarrow (\xi, p) \star (\zeta, r) \cdot (\eta, q) \cdot (\pi, s) \notin \bot\!\!\!\bot$:
  Suppose that $(\xi \star \zeta \cdot \eta \cdot \pi, pqrs) \in \bot\!\!\!\bot$; we must show: $(\mathsf{C}^* \star \xi \cdot \eta \cdot \zeta \cdot \pi, pqrs) \in \bot\!\!\!\bot$ i.e. $\mathsf{C}^* \star \underline{n} \cdot \xi \cdot \eta \cdot \zeta \cdot \pi \in \bot\!\!\!\bot$ for $(pqrs \ll n) = 1$.
  Now, we have: $\mathsf{C}^* \star \underline{n} \cdot \xi \cdot \eta \cdot \zeta \cdot \pi \gg \xi \star \underline{n} \cdot \zeta \cdot \eta \cdot \pi$ which is in $\bot\!\!\!\bot$ by hypothesis.
- $(\mathsf{I}^*, \mathbf{1}) \star (\xi, p) \cdot (\pi, q) \notin \bot\!\!\!\bot \Rightarrow (\xi, p) \star (\pi, q) \notin \bot\!\!\!\bot$:
  Suppose that $(\xi \star \pi, pq) \in \bot\!\!\!\bot$; we must show: $(\mathsf{I}^* \star \xi \cdot \pi, pq) \in \bot\!\!\!\bot$ i.e. $\mathsf{I}^* \star \underline{n} \cdot \xi \cdot \pi \in \bot\!\!\!\bot$ for $(pq \ll n) = 1$.
  Now, we have: $\mathsf{I}^* \star \underline{n} \cdot \xi \cdot \pi \gg \xi \star \underline{n} \cdot \pi$ which is in $\bot\!\!\!\bot$ by hypothesis.
- $(\mathsf{K}^*, \mathbf{1}) \star (\xi, p) \cdot (\eta, q) \cdot (\pi, r) \notin \bot\!\!\!\bot \Rightarrow (\xi, p) \star (\pi, r) \notin \bot\!\!\!\bot$:
  Suppose that $(\xi \star \pi, pr) \in \bot\!\!\!\bot$; we must show: $(\mathsf{K}^* \star \xi \cdot \eta \cdot \pi, pqr) \in \bot\!\!\!\bot$ i.e. $\mathsf{K}^* \star \underline{n} \cdot \xi \cdot \eta \cdot \pi \in \bot\!\!\!\bot$ for $(pqr \ll n) = 1$. Now, we have: $\mathsf{K}^* \star \underline{n} \cdot \xi \cdot \eta \cdot \pi \gg \xi \star \underline{n} \cdot \pi$ which is in $\bot\!\!\!\bot$ by hypothesis.
- $(\mathsf{W}^*, \mathbf{1}) \star (\xi, p) \cdot (\eta, q) \cdot (\pi, r) \notin \bot\!\!\!\bot \Rightarrow (\xi, p) \star (\eta, q) \cdot (\eta, q) \cdot (\pi, r) \notin \bot\!\!\!\bot$:
  Suppose that $(\xi \star \eta \cdot \eta \cdot \pi, pqr) \in \bot\!\!\!\bot$; we must show: $(\mathsf{W}^* \star \xi \cdot \eta \cdot \eta \cdot \pi, pqr) \in \bot\!\!\!\bot$ i.e. $\mathsf{W}^* \star \underline{n} \cdot \xi \cdot \eta \cdot \eta \cdot \pi \in \bot\!\!\!\bot$ for $(pqr \ll n) = 1$.
  Now, we have: $\mathsf{W}^* \star \underline{n} \cdot \xi \cdot \eta \cdot \eta \cdot \pi \gg \xi \star \underline{n} \cdot \eta \cdot \eta \cdot \eta \cdot \pi$ which is in $\bot\!\!\!\bot$ by hypothesis.
- $(\mathsf{cc}^*, \mathbf{1}) \star (\xi, p) \cdot (\pi, q) \notin \bot\!\!\!\bot \Rightarrow (\xi, p) \star (\mathsf{k}_\pi^*, q) \cdot (\pi, q) \notin \bot\!\!\!\bot$:
  Suppose that $(\xi \star \mathsf{k}_\pi^* \cdot \pi, pq) \in \bot\!\!\!\bot$; we must show:

https://doi.org/10.1017/S0960129516000050 Published online by Cambridge University Press

$(\mathsf{cc}^* \star \xi \cdot \pi, pq) \in \Vdash$ i.e. $\mathsf{cc}^* \star \underline{n} \cdot \xi \cdot \pi \in \bot\!\!\!\bot$ for $(pq \ll n) = 1$.
Now, we have: $\mathsf{cc}^* \star \underline{n} \cdot \xi \cdot \pi \succgtr \xi \star \underline{n} \cdot \mathsf{k}_\pi^* \cdot \pi$ which is in $\bot\!\!\!\bot$ by hypothesis.

- $(\mathsf{k}_\pi^*, p) \star (\xi, q) \cdot (\varpi, r) \notin \Vdash \Rightarrow (\xi, q) \star (\pi, p) \notin \Vdash$:
Suppose that $(\xi \star \pi, pq) \in \Vdash$; we must show:
$(\mathsf{k}_\pi^* \star \xi \cdot \varpi, pqr) \in \Vdash$ i.e. $\mathsf{k}_\pi^* \star \underline{n} \cdot \xi \cdot \varpi \in \bot\!\!\!\bot$ for $(pqr \ll n) = 1$.
Now, we have: $\mathsf{k}_\pi^* \star \underline{n} \cdot \xi \cdot \varpi \succgtr \xi \star \underline{n} \cdot \pi$ which is in $\bot\!\!\!\bot$ by hypothesis.

For each closed c-term τ (built with the elementary combinators and the application), we define τ^* by recurrence, as follows:
if τ is an elementary combinator, τ^* is already defined;
we set $(tu)^* = \mathsf{C}t^*u^*$.

In the algebra $\mathcal{B}$, the value of the combinator τ is $\tau_{\mathcal{B}} = (\tau_{\mathcal{A}}^*, \mathbf{1})$.
In particular, the integer n of the algebra $\mathcal{B}$ is $\underline{n}_{\mathcal{B}} = (\underline{n}^*, \mathbf{1})$.
We have $\underline{0}_{\mathcal{B}} = (\underline{0}^*, \mathbf{1}) = (\mathsf{K}^*, \mathbf{1})(\mathsf{I}^*, \mathbf{1})$; therefore: $\underline{0}^* = \mathsf{CK}^*\mathsf{I}^*$.
We have $(\underline{n+1})_{\mathcal{B}} = ((\underline{n+1})^*, \mathbf{1}) = (\sigma^*, \mathbf{1})(\underline{n}^*, \mathbf{1})$; therefore: $(\underline{n+1})^* = \mathsf{C}\sigma^*\underline{n}^*$.

Thus, we have, for every $n \in \mathbb{N}$: $\underline{n}^* = (\mathsf{C}\sigma^*)^n\underline{0}^*$.

We define the proof-like terms of the algebra $\mathcal{B}$ as the terms of the form $(\theta, \mathbf{1})$ where θ is a proof-like term of the algebra $\mathcal{A}$. The condition of coherence for $\mathcal{B}$ is therefore:
If θ is a proof-like term of $\mathcal{A}$, there exist $n \in \mathbb{N}$ and $\pi \in \Pi$ such that $\theta \star \underline{n} \cdot \pi \notin \bot\!\!\!\bot$.
If $\mathcal{A}$ is coherent, then so is $\mathcal{B}$: indeed, if θ is a proof-like term of $\mathcal{A}$, then so is $\theta\underline{0}$; this gives a stack π such that $\theta\underline{0} \star \pi \notin \bot\!\!\!\bot$.

Notations.
The realizability models associated with the algebras $\mathcal{A}$ and $\mathcal{B}$ are denoted, respectively, by $\mathcal{N}_{\mathcal{A}}$ and $\mathcal{N}_{\mathcal{B}}$.
The truth value of a formula F in the realizability model $\mathcal{N}_{\mathcal{B}}$ will be denoted by $\|F\|_{\mathcal{B}}$ or also $|||F|||$.
We write $(\xi, p) \Vdash_{\mathcal{B}} F$ or $(\xi, p) \Vdash\!\!\!| F$ to say that (ξ, p) realizes the formula F in the realizability model $\mathcal{N}_{\mathcal{B}}$.

4.2. *The collapsing function*

We now define $\mathcal{G} \in \mathcal{M}$ in the following way:
$\mathcal{G} = \{((m, \alpha), (\pi, p)) \ ; \ m \in \mathbb{N}, \alpha \in \kappa, \pi \in \Pi, p \in P \setminus \{\mathbb{O}\}, p(m) \text{ is defined and } p(m) = \alpha\}$.

Theorem 35.
The formula ($\mathcal{G}$ is a surjection from $\mathbb{N}$ onto $\beth\kappa$) is realized in the model $\mathcal{N}_{\mathcal{B}}$.
More precisely, we have:

i. $(\theta_0, \mathbf{1}) \Vdash\!\!\!| \forall x \forall y \forall y' \left((x, y) \varepsilon \mathcal{G}, y \neq y' \rightarrow (x, y') \notin \mathcal{G}\right)$ with $\theta_0 = \lambda n \lambda k \lambda x (x) n$;

ii. $(\theta_1, \mathbf{1}) \Vdash\!\!\!| \forall y^{\beth\kappa} [\forall x^{\mathrm{int}} ((x, y) \notin \mathcal{G}) \rightarrow \bot]$ with $\theta_1 = \lambda n \lambda x ((((n)(\mathsf{CB})(\mathsf{C})\sigma^*)(\mathsf{C})x)\underline{0}^*)(\sigma)n$, and $\sigma = (\mathsf{BW})(\mathsf{B})\mathsf{B}$ (successor).

https://doi.org/10.1017/S0960129516000050 Published online by Cambridge University Press

Proof.

i. Let $m \in \mathbb{N}, \alpha, \alpha' \in \kappa, (\pi, p) \in |||(m,\alpha)\not\in\mathcal{G}|||$, $(\pi', p') \in |||(m,\alpha')\not\in\mathcal{G}|||$ and $(\xi, q) \Vdash \alpha \neq \alpha'$.
Thus, we have $m \in \mathrm{dom}(p), m \in \mathrm{dom}(p'), p(m) = \alpha$ and $p'(m) = \alpha'$.
By Lemma 7, we can replace the formula $(m,\alpha)\,\varepsilon\,\mathcal{G}$, which is $\neg((m,\alpha)\not\in\mathcal{G})$, with the set of terms $\ulcorner((m,\alpha)\not\in\mathcal{G})$, which is $\{\mathsf{k}_{(\pi,p)}\ ;\ (\pi,p) \in |||(m,\alpha)\not\in\mathcal{G}|||\}$.
Therefore, we have to show that:
$(\theta_0, \mathbf{1}) \star \mathsf{k}_{(\pi,p)} \cdot (\xi, q) \cdot (\pi', p') \in \perp\!\!\!\perp$ that is $(\theta_0 \star \mathsf{k}_\pi^* \cdot \xi \cdot \pi', pp'q) \in \perp\!\!\!\perp$.
This is obvious if $pp'q = \mathbb{O}$. Otherwise, p and p' are compatible, thus $\alpha = \alpha'$.
Let n be such that $(pp'q \ll n) = 1$; we must show that $\theta_0 \star \underline{n} \cdot \mathsf{k}_\pi^* \cdot \xi \cdot \pi' \in \perp\!\!\!\perp$ i.e. $\xi \star \underline{n} \cdot \pi' \in \perp\!\!\!\perp$.
Now, we have $(\xi, q) \Vdash \bot$ by hypothesis on (ξ, q), thus $(\xi, q) \star (\pi', \mathbf{1}) \in \perp\!\!\!\perp$.
Since $(q \ll n) = 1$, it follows that $\xi \star \underline{n} \cdot \pi' \in \perp\!\!\!\perp$.

ii. Let us first show that $\theta_1 \star \underline{n} \cdot \eta \cdot \varpi \gg \eta \star \underline{n+1} \cdot \underline{n}^* \cdot \varpi$ for each $n \in \mathbb{N}, \eta \in \Lambda$ and $\varpi \in \Pi$.
We have $\theta_1 \star \underline{n} \cdot \eta \cdot \varpi \gg \underline{n} \star (\mathsf{CB})(\mathsf{C})\sigma^* \cdot \mathsf{C}\eta \cdot \underline{0}^* \cdot \underline{n+1} \cdot \varpi$. By Lemma 15(ii), in which we set $\zeta = \mathsf{C}\eta, \phi = \mathsf{C}\sigma^*, \alpha = \underline{0}^*, \varsigma = \sigma, O = \underline{0}$ and $\pi = \underline{n+1} \cdot \varpi$, we obtain:
$\theta_1 \star \underline{n} \cdot \eta \cdot \varpi \gg \mathsf{C}\eta \star \underline{n}^* \cdot \underline{n+1} \cdot \varpi$ (since $\underline{n}^* = (\mathsf{C}\sigma^*)^n \underline{0}^*$) $\gg \eta \star \underline{n+1} \cdot \underline{n}^* \cdot \varpi$.

We prove now that $(\theta_1, \mathbf{1}) \Vdash \forall y^{\beth_\kappa}[\forall x^{\mathrm{int}}((x,y)\not\in\mathcal{G}) \to \bot]$.
Let $\alpha \in \kappa$, $(\eta, p_0) \Vdash \forall x^{\mathrm{int}}((x,\alpha)\not\in\mathcal{G})$ and $(\varpi, q_0) \in \Pi \times P$;
we show that $(\theta_1, \mathbf{1}) \star (\eta, p_0) \cdot (\varpi, q_0) \in \perp\!\!\!\perp$.
This is trivial if $p_0 q_0 = \mathbb{O}$; otherwise, let $n \in \mathbb{N}$ be such that $(p_0 q_0 \ll n) = 1$.
We must show that $\theta_1 \star \underline{n} \cdot \eta \cdot \varpi \in \perp\!\!\!\perp$, that is $\eta \star \underline{n+1} \cdot \underline{n}^* \cdot \varpi \in \perp\!\!\!\perp$.
But we have $(\eta, p_0) \Vdash \{(\underline{n}^*, \mathbf{1})\} \to (n,\alpha)\not\in\mathcal{G}$ by hypothesis on η.
Since $(p_0 q_0 \ll n) = 1$, we can define $q \in P$ with domain $n+1$ such that $q \supset p_0 q_0$ and $q(n) = \alpha$. Then, we have $(\varpi, q) \in |||(n,\alpha)\not\in\mathcal{G}|||$ by definition of $\mathcal{G}$.
We have thus $(\eta, p_0) \star (\underline{n}^*, \mathbf{1}) \cdot (\varpi, q) \in \perp\!\!\!\perp$ that is $(\eta \star \underline{n}^* \cdot \varpi, p_0 q) \in \perp\!\!\!\perp$.
But we have $p_0 q = q$, and therefore $(\eta \star \underline{n}^* \cdot \varpi, q) \in \perp\!\!\!\perp$.
Since $(q \ll n+1) = 1$, it follows that $\eta \star \underline{n+1} \cdot \underline{n}^* \cdot \varpi \in \perp\!\!\!\perp$. □

Corollary 36. $\mathcal{N}_\mathcal{B}$ realizes the non-extensional axiom of choice and thus also DC.

Proof. Indeed, by Theorem 35, the model $\mathcal{N}_\mathcal{B}$ realizes the formula: ($\beth\kappa$ is countable).
But we have $\kappa = \mathrm{card}(\mathbf{\Lambda} \cup \mathbf{\Pi} \cup \mathbb{N})$, since $\kappa \geqslant \mathrm{card}(\Lambda \cup \Pi \cup \mathbb{N})$ and $\kappa = \mathrm{card}(P)$.
Therefore $\mathcal{N}_\mathcal{B}$ realizes NEAC, by Theorem 31. □

Remark. Intuitively, the model $\mathcal{N}_\mathcal{B}$ is an extension of the model $\mathcal{N}_\mathcal{A}$ obtained by forcing, by collapsing $\beth\kappa$. We cannot apply directly the usual theory of forcing, because $\beth\kappa$ is not defined in ZF.

4.3. *Elementary formulas*

Elementary formulas are defined as follows, where t, u are ℓ-terms, i.e. terms built with variables, individuals, and function symbols defined in $\mathcal{M}$:

- $\top, \bot$ are elementary formulas;

https://doi.org/10.1017/S0960129516000050 Published online by Cambridge University Press

- if U is an elementary formula, then $t = u \hookrightarrow U$ and $\forall x\, U$ are too;
- if U, V are elementary formulas, then $U \to V$ too;
- if U is an elementary formula, then $\forall n^{\mathrm{int}} U$ too.

Remark. $t \neq u$ is an elementary formula, and also $t \not\in u$. Indeed, this formula can be written $f(t,u) \neq 1$ where f is the function symbol defined in $\mathcal{M}$ by: $f(a,b) = 1$ iff $a \in b$. If U is an elementary formula, then $\forall x^{\in t} U$ is too: indeed, it is written $\forall x(f(x,t) = 1 \hookrightarrow U)$.

For each elementary formula U, we define two formulas U_p and U^p, with one additional free variable p, by the conditions below.
Condition 1 defines U^p by means of U_p; conditions 2 to 5 define U_p by recurrence:

1. $U^p \equiv \forall q^{\in P} \forall n^{\mathrm{int}}((pq \ll n) = 1 \hookrightarrow U_q)$;
2. $\bot_p \equiv \bot$ and $\top_p \equiv \top$;
3. $(t = u \hookrightarrow U)_p \equiv (t = u \hookrightarrow U_p)$; $(\forall x\, U[x])_p \equiv \forall x\, U_p[x]$;
4. $(U \to V)_p \equiv \forall q^{\in P} \forall r^{\in P}\, (p = qr \hookrightarrow (U^q \to V_r))$;
5. $(\forall n^{\mathrm{int}} U[n])_p \equiv \forall n^{\in \mathbb{N}}(\{\underline{n}^*\} \to U_p[n])$, in other words:
   $\|(\forall n^{\mathrm{int}} U[n])_p\| = \{\underline{n}^* \cdot \pi \ ;\ n \in \mathbb{N}, \pi \in \|U_p[n]\|\}$.

With the help of these formulas U_p, U^p, we can interpret realizability in the algebra $\mathcal{B}$ in terms of realizability in the algebra $\mathcal{A}$, at least for elementary formulas, as shown in the following Lemma 37.

Lemma 37. For each closed elementary formula U, we have:
$(\pi, p) \in |||U||| \Leftrightarrow \pi \in \|U_p\|$; $(\xi, p) \Vdash U \Leftrightarrow \xi \Vdash U^p$.

Proof. By recurrence on the length of the formula U.

1. We have $(\xi, p) \Vdash U \Leftrightarrow (\xi, p) \star (\pi, q) \in \perp\!\!\!\perp$ for $(\pi, q) \in |||U|||$, that is:
$(\xi \star \pi, pq) \in \perp\!\!\!\perp$ for every $\pi \in \|U_q\|$, by the recurrence hypothesis, or also:
$(\forall q \in P)(\forall \pi \in \|U_q\|)(\forall n \in \mathbb{N})((pq \ll n) = 1 \Rightarrow \xi \star \underline{n} \cdot \pi \in \bot\!\!\!\bot)$ which is equivalent to:
$\xi \Vdash \forall q^{\in P} \forall n^{\mathrm{int}}((pq \ll n) = 1 \hookrightarrow U_q)$ that is $\xi \Vdash U^p$.

2 and 3. Obvious.

4. Any element of $|||U \to V|||$ has the form $(\xi, q) \cdot (\pi, r)$, i.e. $(\xi \cdot \pi, p)$, with $p = qr$, $(\xi, q) \Vdash U$ and $(\pi, r) \in |||V|||$; by the recurrence hypothesis, this is equivalent to:
$\xi \cdot \pi \in \|U^q \to V_r\|$.

5. We have $|||\forall n^{\mathrm{int}} U[n]||| = |||\forall n^{\in \mathbb{N}}(\{(\underline{n}^*, \mathbf{1})\} \to U[n])|||$
$= \{(\underline{n}^*, \mathbf{1}) \cdot (\pi, p)\ ;\ n \in \mathbb{N}, (\pi, p) \in |||U[n]|||\} = \{(\underline{n}^* \cdot \pi, p)\ ;\ n \in \mathbb{N}, (\pi, p) \in |||U[n]|||\}$.
Thus, by the recurrence hypothesis, it is $\{(\underline{n}^* \cdot \pi, p)\ ;\ n \in \mathbb{N}, \pi \in \|U_p[n]\|\}$. □

Lemma 38.
For each elementary formula U, there exist two proof-like terms θ_U^0, θ_U^1, such that:

i. $\theta_U^0 \Vdash \forall p^{\in P} \forall n^{\mathrm{int}}((p \ll n) = 1 \hookrightarrow (U \to U_p))$;

ii. $\theta_U^1 \Vdash \forall p^{\in P} \forall n^{\mathrm{int}}((p \ll n) = 1 \hookrightarrow (U_p \to U))$;

https://doi.org/10.1017/S0960129516000050 Published online by Cambridge University Press

iii. $\tau_U^0 \Vdash \forall p^{\neg P} \forall n^{\mathrm{int}}((p \ll n) = 1 \hookrightarrow (U \to U^p))$;

iv. $\tau_U^1 \Vdash \forall p^{\neg P} \forall n^{\mathrm{int}}((p \ll n) = 1 \hookrightarrow (U^p \to U))$;

with $\tau_U^0 = \lambda n \lambda x \lambda m (\theta_U^0) m x$ and $\tau_U^1 = \lambda n \lambda x (\theta_U^1 n)(x) n$.

Proof. We first show (iii) and (iv) from (i) and (ii).

(i)⇒(iii)
Let $p \in P$ and $n \in \mathbb{N}$ be such that $(p \ll n) = 1$; let $\xi \Vdash U$ and $\pi \in \|U^p\|$. We have to show: $\lambda n \lambda x \lambda m (\theta_U^0) m x \star \underline{n} \cdot \xi \cdot \pi \in \perp\!\!\!\perp$.
Now, by the definition (1) of U^p, there exist $q \in P$, $m \in \mathbb{N}$ and $\varpi \in \|U_q\|$ such that $(pq \ll m) = 1$ and $\pi = \underline{m} \cdot \varpi$. Therefore, we have $(q \ll m) = 1$ and, by (i):
$\theta_U^0 \star \underline{m} \cdot \xi \cdot \varpi \in \perp\!\!\!\perp$, hence $\lambda n \lambda x \lambda m (\theta_U^0) m x \star \underline{n} \cdot \xi \cdot \underline{m} \cdot \varpi \in \perp\!\!\!\perp$.

(ii)⇒(iv)
Let $p \in P$, $n \in \mathbb{N}, \xi \in \Lambda$ and $\pi \in \|U\|$ such that $(p \ll n) = 1$ and $\xi \Vdash U^p$.
We have to show: $\lambda n \lambda x (\theta_U^1 n)(x) n \star \underline{n} \cdot \xi \cdot \pi \in \perp\!\!\!\perp$ i.e. $\theta_U^1 \star \underline{n} \cdot \xi \underline{n} \cdot \pi \in \perp\!\!\!\perp$.
But, by the definition (1) of U^p, in which we set $q = p$, we have $\xi \underline{n} \Vdash U_p$; therefore, the desired result follows from (ii).

We now show (i) and (ii) by recurrence on the length of U.

- If U is $\perp$ or $\top$, we take $\theta_U^0 = \theta_U^1 = \lambda n \lambda x\, x$.
- If $U \equiv (t = u \hookrightarrow V)$ or $U \equiv \forall x\, V$, then $\theta_U^0 = \theta_V^0$ and $\theta_U^1 = \theta_V^1$ by (3).
- If $U \equiv V \to W$, let $q, r \in \mathbb{N}$ and $p = qr$; let $n \in \mathbb{N}$ such that $(p \ll n) = 1$. We have:
  $\tau_V^0 \underline{n} \Vdash V \to V^q$; $\tau_V^1 \underline{n} \Vdash V^q \to V$; $\theta_W^0 \underline{n} \Vdash W \to W_r$; $\theta_W^1 \underline{n} \Vdash W_r \to W$.
  Let $\xi \Vdash V \to W$; then, by the recurrence hypothesis, we have:
  $\lambda y (\theta_W^0 \underline{n})(\xi) y \Vdash V \to W_r$ and $\lambda z (\theta_W^0 \underline{n})(\xi)(\tau_V^1 \underline{n}) z \Vdash V^q \to W_r$.
  Thus, by (4), we obtain $\theta_U^0 = \lambda n \lambda x \lambda z (\theta_W^0 n)(x)(\tau_V^1 n) z$.
  Now, let $\xi \Vdash V^q \to W_r$; then, by the recurrence hypothesis, we have:
  $\lambda y (\theta_W^1 \underline{n})(\xi) y \Vdash V^q \to W$ and $\lambda z (\theta_W^1 \underline{n})(\xi)(\tau_V^0 \underline{n}) z \Vdash V \to W$.
  Thus, by (4), we obtain $\theta_U^1 = \lambda n \lambda x \lambda z (\theta_W^1 n)(x)(\tau_V^0 n) z$.
- If $U \equiv \forall m^{\mathrm{int}} V[m]$, we first prove:

Lemma 39.
There exist two proof-like terms T_0, T_1 such that, for every closed formula F of ZF_ε:

i. $T_0 \Vdash \forall n^{\neg \mathbb{N}}((\{\underline{n}^*\} \to F) \to (\{\underline{n}\} \to F))$.

ii. $T_1 \Vdash \forall n^{\neg \mathbb{N}}((\{\underline{n}\} \to F) \to (\{\underline{n}^*\} \to F))$.

iii. For every elementary formula $V[n]$, we have:
$T_0 \Vdash (\forall n^{\mathrm{int}} V[n])_p \to \forall n^{\mathrm{int}} V_p[n]$ and $T_1 \Vdash \forall n^{\mathrm{int}} V_p[n] \to (\forall n^{\mathrm{int}} V[n])_p$.

Proof.
i. We apply Lemma 15(ii) to the realizability algebra $\mathcal{A}$, with:
$\varsigma = \sigma$, $O = \underline{0}$, $\phi = \mathsf{C}\sigma^*$ and $\alpha = \underline{0}^*$. For every $n \in \mathbb{N}, \zeta \in \Lambda$ and $\pi \in \Pi$, we obtain:
$\underline{n} \star (\mathsf{CB})(\mathsf{C})\sigma^* \cdot \zeta \cdot \underline{0}^* \cdot \pi \gg \zeta \star \underline{n}^* \cdot \pi$, since $\underline{n}^* = (\mathsf{C}\sigma^*)^n \underline{0}^*$.

https://doi.org/10.1017/S0960129516000050 Published online by Cambridge University Press

Therefore, if we set $T_0 = \lambda f \lambda n((n)(\mathsf{CB})(\mathsf{C})\sigma^*)f\underline{0}^*$, we have $T_0 \star \zeta \cdot \underline{n} \cdot \pi \gg \zeta \star \underline{n}^* \cdot \pi$.
Thus, we have $T_0 \Vdash \forall n^{\exists\mathbb{N}}((\{\underline{n}^*\} \to F) \to (\{\underline{n}\} \to F))$.

ii. We apply now Lemma 15(i) to the realizability algebra $\mathcal{B}$, with:
$\varsigma = \sigma_{\mathcal{B}}$, $O = \underline{0}_{\mathcal{B}}$, $\phi = (\mathsf{C}\Sigma, \mathbf{1})\,\alpha = (\Omega, \mathbf{1})$ and $\Omega = \lambda d \lambda f \lambda a\, a$; $\Sigma = \lambda n \lambda d \lambda f \lambda a (ndf)(f)a$.
Since $\underline{n}_{\mathcal{B}} = (\sigma_{\mathcal{B}})^n \underline{0}_{\mathcal{B}} = (\underline{n}^*, \mathbf{1})$, we get, by setting $\Sigma_2 = (\mathsf{C})^2\Sigma$:
$(\underline{n}^*, \mathbf{1}) \star (\mathsf{C}\Sigma, \mathbf{1}) \cdot (\Omega, \mathbf{1}) \cdot (\varpi, \mathbf{1}) \gg ((\Sigma_2)^n\Omega, \mathbf{1}) \star (\varpi, \mathbf{1})$
because $((\mathsf{C}\Sigma, \mathbf{1}))^n(\Omega, \mathbf{1}) = ((\Sigma_2)^n\Omega, \mathbf{1})$. We write this as:
$(\underline{n}^* \star \mathsf{C}\Sigma \cdot \Omega \cdot \varpi, \mathbf{1}) \gg ((\Sigma_2)^n\Omega \star \varpi, \mathbf{1})$.
It follows that $\underline{n}^* \star \underline{0} \cdot \mathsf{C}\Sigma \cdot \Omega \cdot \varpi \gg (\Sigma_2)^n\Omega \star \underline{d} \cdot \varpi$ for some $d \in \mathbb{N}$.
Let us take $\varpi = \mathsf{CB}\sigma \cdot \zeta \cdot \underline{0} \cdot \pi$. We obtain:
$\underline{n}^* \star \underline{0} \cdot \mathsf{C}\Sigma \cdot \Omega \cdot \mathsf{CB}\sigma \cdot \zeta \cdot \underline{0} \cdot \pi \gg (\Sigma_2)^n\Omega \star \underline{d} \cdot \mathsf{CB}\sigma \cdot \zeta \cdot \underline{0} \cdot \pi$.
Now, we apply Lemma 16(ii), with $\phi = \sigma$ and $\alpha = \underline{0}$ (note that $\Sigma_2 = (\mathsf{C})^2\Sigma$ satisfies the hypothesis of Lemma 16). We obtain $(\Sigma_2)^n\Omega \star \underline{d} \cdot \mathsf{CB}\sigma \cdot \zeta \cdot \underline{0} \cdot \pi \succ \zeta \star (\sigma)^n\underline{0} \cdot \pi$ and therefore: $\underline{n}^* \star \underline{0} \cdot \mathsf{C}\Sigma \cdot \Omega \cdot \mathsf{CB}\sigma \cdot \zeta \cdot \underline{0} \cdot \pi \gg \zeta \star \underline{n} \cdot \pi$.
Finally, if we set $T_1 = \lambda f \lambda n((((n\underline{0})(\mathsf{C})\Sigma)\Omega)(\mathsf{C})\mathsf{B}\sigma)f\underline{0}$, we have:
$T_1 \star \zeta \cdot \underline{n}^* \cdot \pi \gg \zeta \star \underline{n} \cdot \pi$ and therefore $T_1 \Vdash \forall n^{\exists\mathbb{N}}((\{\underline{n}\} \to F) \to (\{\underline{n}^*\} \to F))$.

iii. This follows immediately from (i) and (ii), by definition of $(\forall n^{\mathrm{int}} V[n])_p$. □

We can now finish the proof of Lemma 38, considering the last case which is:

- $U \equiv \forall m^{\mathrm{int}} V[m]$.

We show that $\theta^0_U = \lambda n \lambda x (T_1) \lambda m (\theta^0_V n)(x)m$.
By the recurrence hypothesis, we have $\theta^0_V \Vdash \forall p^{\exists P} \forall n^{\mathrm{int}}((p \ll n) = 1 \hookrightarrow (V[m] \to V_p[m]))$.
Let $p \in P$, $n \in \mathbb{N}$, $\xi \in \Lambda$ be such that $(p \ll n) = 1$ and $\xi \Vdash \forall m^{\mathrm{int}} V[m]$.
Then, for every $m \in \mathbb{N}$, we have $\xi\underline{m} \Vdash V[m]$; thus $(\theta^0_V \underline{n})(\xi)m \Vdash V_p[m]$ and therefore:
$\lambda m(\theta^0_V \underline{n})(\xi)m \Vdash \forall m^{\mathrm{int}} V_p[m]$. By Lemma 39(iii), we get:
$(T_1)\lambda m(\theta^0_V \underline{n})(\xi)m \Vdash (\forall m^{\mathrm{int}} V[m])_p$ and therefore:
$\lambda x (T_1) \lambda m (\theta^0_V \underline{n})(x)m \Vdash \forall m^{\mathrm{int}} V[m] \to (\forall m^{\mathrm{int}} V[m])_p$. Finally:
$\lambda n \lambda x (T_1) \lambda m (\theta^0_V n)(x)m \Vdash \forall p^{\exists P} \forall n^{\mathrm{int}}((p \ll n) = 1 \hookrightarrow (\forall m^{\mathrm{int}} V[m] \to (\forall m^{\mathrm{int}} V[m])_p))$.

We show now that $\theta^1_U = \lambda n \lambda x \lambda m (\theta^1_V n)(T_0)xm$.
By the recurrence hypothesis, we have $\theta^1_V \Vdash \forall p^{\exists P} \forall n^{\mathrm{int}}((p \ll n) = 1 \hookrightarrow (V_p[m] \to V[m]))$;
Let $p \in P$, $n \in \mathbb{N}$, $\xi \in \Lambda$ be such that $(p \ll n) = 1$ and $\xi \Vdash (\forall m^{\mathrm{int}} V[m])_p$.
By Lemma 39(iii), we have $T_0\xi \Vdash \forall m^{\mathrm{int}} V_p[m]$, thus $T_0\xi\underline{m} \Vdash V_p[m]$.
Therefore $(\theta^1_V \underline{n})(T_0)\xi\underline{m} \Vdash V[m]$, and $\lambda m(\theta^1_V \underline{n})(T_0)\xi m \Vdash \forall m^{\mathrm{int}} V[m]$, hence the result. □

Theorem 40.
The same closed elementary formulas, with parameters in $\mathcal{M}$, are realized in the models $\mathcal{N}_{\mathcal{A}}$ and $\mathcal{N}_{\mathcal{B}}$.

Proof. Let U be a closed elementary formula, which is realized in $\mathcal{N}_{\mathcal{A}}$ and let θ be a proof-like term such that $\theta \Vdash U$. Then, we have $(\tau^0_U)\underline{n}\theta \Vdash U^p$ for $(p \ll n) = 1$, by Lemma 38(iii); therefore, setting $p = \varnothing = \mathbf{1}$, we have $((\tau^0_U)\underline{0}\theta, \mathbf{1}) \Vdash U$ by Lemma 37.
Therefore, the formula U is also realized in the model $\mathcal{N}_{\mathcal{B}}$.
Conversely, if $(\theta, \mathbf{1}) \Vdash U$ with $\theta \in \mathsf{PL}$, we have $\theta \Vdash U^{\mathbf{1}}$, by Lemma 37. Thus $\tau^1_U \underline{0}\,\theta \Vdash U$ by Lemma 38(iv). □

https://doi.org/10.1017/S0960129516000050 Published online by Cambridge University Press

Remark. For instance:

- If the Boolean algebra $\beth 2$ has four ε-elements or if it is atomless, in the model $\mathcal{N}_{\mathcal{A}}$, the same goes for the model $\mathcal{N}_{\mathcal{B}}$.
- Arithmetical formulas are elementary. Therefore, by Theorem 40, the models $\mathcal{N}_{\mathcal{A}}$ and $\mathcal{N}_{\mathcal{B}}$ realize the same arithmetical formulas. In fact, this was already known, because they are the same as the arithmetical formulas which are true in $\mathcal{M}$ (Krivine 2009; 2007).

4.4. *Arithmetical formulas and dependent choice*

In this section, we obtain, by means of the previous results, a technique to transform into a program, a given proof, in ZF + DC, of an arithmetical formula F.
We notice that this program is a closed c-term, written with the elementary combinators $\mathsf{B}, \mathsf{C}, \mathsf{I}, \mathsf{K}, \mathsf{W}, \mathsf{cc}$ *without any other instruction.*

Thus, let us consider a proof of $\mathrm{ZF}_{\varepsilon} \vdash \mathrm{NEAC} \to F$ (NEAC is the *non-extensional axiom of choice*, see Section 3.3).
It gives us a closed c-term Φ_0 such that $\Phi_0 \Vdash \mathrm{NEAC} \to F$, in every realizability algebra.
We now describe a re-writing on closed c-terms, which will transform Φ_0 into a closed c-term Φ such that $\Phi \Vdash F$ in every realizability algebra $\mathcal{A}$.

By Theorem 31, we have $\Phi_1 \Vdash (\beth\kappa \text{ is countable}) \to F$ with $\Phi_1 = \lambda x(\Phi_0)(\mathsf{H})x$.
We apply this result in the algebra $\mathcal{B}$, which gives:
$(\Phi_1^*, \mathbf{1}) \Vdash (\beth\kappa \text{ is countable} \to F)$.
Now, Theorem 35 gives a closed c-term Δ such that $(\Delta, \mathbf{1}) \Vdash (\beth\kappa \text{ is countable})$.
It follows that $(\Phi_1^*, \mathbf{1})(\Delta, \mathbf{1}) \Vdash F$, i.e. $(\Psi, \mathbf{1}) \Vdash F$, with $\Psi = \mathsf{C}\Phi_1^*\Delta$.
Since F is an arithmetical formula, it is an elementary formula.
Therefore, by Lemma 37, we have $\Psi \Vdash F^{\mathbf{1}}$. Now, by Lemma 38(iv), we have:
$\tau_F^1 \Vdash \forall p^{\beth P} \forall n^{\mathrm{int}}((p \ll n) = 1 \hookrightarrow (F^p \to F))$.
We set $p = \mathbf{1}$ and $n = 0$, and we obtain $\tau_F^1 \underline{0} \Vdash F^{\mathbf{1}} \to F$.

Finally, by setting $\Phi = (\tau_F^1)\underline{0}\Psi$, we have $\Phi \Vdash F$.

4.5. *A relative consistency result*

In Krivine (2012), we have defined a countable realizability algebra $\mathcal{A}$ such that the characteristic Boolean algebra $\beth 2$ of the model $\mathcal{N}_{\mathcal{A}}$ is atomless (in this example, we have $\kappa = \mathbb{N}$).
If we apply the technique of Section 4, in order to collapse $\beth\kappa$, we obtain a realizability algebra $\mathcal{B}$ and a model $\mathcal{N}_{\mathcal{B}}$, the characteristic Boolean algebra of which is also atomless.
Indeed, the property: ($\beth 2$ is atomless) is expressed by an elementary formula.
But now $\beth 2$ is *the countable atomless Boolean algebra* (they are all isomorphic). Therefore, by applying Theorems 31 and 34, we obtain the relative consistency result (i) announced in the introduction.

https://doi.org/10.1017/S0960129516000050 Published online by Cambridge University Press

Remark. We note that this method applies to every realizability algebra such that we have:
$\Vdash$ ($\daleth 2$ is an atomless Boolean algebra).

5. A two-threads model ($\daleth 2$ with four elements)

In this section, we suppose that $\mathcal{A}$ is a *standard realisability algebra* (Krivine 2012). This means, by definition, that the terms and the stacks are finite sequences, built with:

the alphabet B, C, I, K, W, cc, k, $\cdot$, (,), [,]
a countable set of *term constants* (also called *instructions*),
a countable set of *stack constants*

and that they are defined by the following rules:

B, C, I, K, W, cc and all the term constants are terms;
if t, u are terms, the sequence $(t)u$ is a term;
if π is a stack, the sequence $\mathsf{k}[\pi]$ is a term (denoted by k_π);
each stack constant is a stack;
if t is a term and π is a stack, then $t \cdot \pi$ is a stack.

If t is a term and π is a stack, then the ordered pair (t, π) is a *process*, denoted by $t \star \pi$.

A proof-like term of $\mathcal{A}$ is a term which does not contain the symbol k; or, which is the same, a term which does not contain any stack constant.

We now build a realizability model in which $\daleth 2$ has exactly 4 elements.

We suppose that there are exactly two stack constants π^0, π^1 and one term constant d.
For $i \in \{0,1\}$, let Λ^i (resp. Π^i) be the set of terms (resp. stacks) which contain the only stack constant π^i.
For $i, j \in \{0,1\}$, define $\perp\!\!\!\perp^i_j$ as the least set $P \subset \Lambda^i \star \Pi^i$ of processes such that:

1. $\mathsf{d} \star \underline{j} \cdot \pi \in P$ for every $\pi \in \Pi^i$.
2. $\xi \star \pi \in \Lambda^i \star \Pi^i$, $\xi' \star \pi' \in P$, $\xi \star \pi \succ \xi' \star \pi' \Rightarrow \xi \star \pi \in P$
3. If at least two out of three processes $\xi \star \pi$, $\eta \star \pi$, $\zeta \star \pi$ are in P, then $\mathsf{d} \star \underline{2} \cdot \xi \cdot \eta \cdot \zeta \cdot \pi \in P$.

Remarks. The pre-order $\succ$ on $\Lambda \star \Pi$ was defined at the beginning of Section 2.
We express condition 2 by saying that P *is saturated in* $\Lambda^i \star \Pi^i$.
Following this definition of $\succ$, the constant d is a *halting instruction*. Indeed, we have:

$$\mathsf{d} \star \pi \succ \xi \star \varpi \;\Leftrightarrow\; \xi \star \varpi = \mathsf{d} \star \pi.$$

We define $\perp\!\!\!\perp$ by: $\Lambda \star \Pi \setminus \perp\!\!\!\perp = (\Lambda^0 \star \Pi^0 \setminus \perp\!\!\!\perp^0_0) \cup (\Lambda^1 \star \Pi^1 \setminus \perp\!\!\!\perp^1_1)$
In other words, a process is in $\perp\!\!\!\perp$ if and only if
either it is in $\perp\!\!\!\perp^0_0 \cup \perp\!\!\!\perp^1_1$ or it contains both stack constants π^0, π^1.

Lemma 41. If $\xi \star \pi \in \perp\!\!\!\perp^i_j$ and $\xi \star \pi \succ \xi' \star \pi'$ then $\xi' \star \pi' \in \perp\!\!\!\perp^i_j$ (closure by reduction).

Proof. Suppose that $\xi_0 \star \pi_0 \succ \xi'_0 \star \pi'_0$, $\xi_0 \star \pi_0 \in \perp\!\!\!\perp^i_j$ and $\xi'_0 \star \pi'_0 \notin \perp\!\!\!\perp^i_j$. We may suppose that:

(*) $\xi_0 \star \pi_0 \succ \xi'_0 \star \pi'_0$ in exactly one step of reduction.

https://doi.org/10.1017/S0960129516000050 Published online by Cambridge University Press

Let us show that $\perp\!\!\!\perp^i_j \setminus \{\xi_0 \star \pi_0\}$ has properties 1,2,3 defining $\perp\!\!\!\perp^i_j$, which will contradict the definition of $\perp\!\!\!\perp^i_j$:

1. If $\xi_0 \star \pi_0 = \mathsf{d} \star \underline{j} \cdot \pi$, with $\pi \in \Pi^i$, then $\mathsf{d} \star \underline{j} \cdot \pi \succ \xi'_0 \star \pi'_0$, thus $\xi'_0 \star \pi'_0 = \mathsf{d} \star \underline{j} \cdot \pi$. Therefore $\xi'_0 \star \pi'_0 \in \perp\!\!\!\perp^i_j$, which is false.
2. Suppose $\xi \star \pi \in \Lambda^i \star \Pi^i$, $\xi \star \pi \succ \xi' \star \pi' \in \perp\!\!\!\perp^i_j$, $\xi' \star \pi' \neq \xi_0 \star \pi_0$. Then $\xi \star \pi \in \perp\!\!\!\perp^i_j$, by (2).
   If $\xi \star \pi = \xi_0 \star \pi_0$, then $\xi_0 \star \pi_0 \succ \xi' \star \pi'$; since $\xi' \star \pi' \neq \xi_0 \star \pi_0$, it follows from (*) that $\xi'_0 \star \pi'_0 \succ \xi' \star \pi'$ and therefore $\xi'_0 \star \pi'_0 \in \perp\!\!\!\perp^i_j$, which is false.
3. Suppose that two out of the processes $\xi \star \pi$, $\eta \star \pi$, $\zeta \star \pi$ are in $\perp\!\!\!\perp^i_j \setminus \{\xi_0 \star \pi_0\}$, but $\mathsf{d} \star \underline{2} \cdot \xi \cdot \eta \cdot \zeta \cdot \pi$ is not. From (3), it follows that $\mathsf{d} \star \underline{2} \cdot \xi \cdot \eta \cdot \zeta \cdot \pi = \xi_0 \star \pi_0$.
   Thus, $\mathsf{d} \star \underline{2} \cdot \xi \cdot \eta \cdot \zeta \cdot \pi \succ \xi'_0 \star \pi'_0$, and therefore $\xi'_0 \star \pi'_0 = \mathsf{d} \star \underline{2} \cdot \xi \cdot \eta \cdot \zeta \cdot \pi$.

Therefore $\xi'_0 \star \pi'_0 \in \perp\!\!\!\perp^i_j$, which is false. □

Lemma 42. $\perp\!\!\!\perp^i_0 \cap \perp\!\!\!\perp^i_1 = \varnothing$.

Proof. We prove that $(\Lambda^i \star \Pi^i \setminus \perp\!\!\!\perp^i_1) \supset \perp\!\!\!\perp^i_0$ by showing that $\Lambda^i \star \Pi^i \setminus \perp\!\!\!\perp^i_1$ has properties 1, 2, 3 which define $\perp\!\!\!\perp^i_0$.

1. $\mathsf{d} \star \underline{0} \cdot \pi \notin \perp\!\!\!\perp^i_1$ because $\perp\!\!\!\perp^i_1 \setminus \{\mathsf{d} \star \underline{0} \cdot \pi\}$ has properties 1, 2, 3 defining $\perp\!\!\!\perp^i_1$.
2. Follows from Lemma 41.
3. Suppose $\xi_0 \star \pi_0$, $\eta_0 \star \pi_0 \notin \perp\!\!\!\perp^i_1$; we show that $\mathsf{d} \star \underline{2} \cdot \xi_0 \cdot \eta_0 \cdot \zeta_0 \cdot \pi_0 \notin \perp\!\!\!\perp^i_1$ by showing that $\perp\!\!\!\perp^i_1 \setminus \{\mathsf{d} \star \underline{2} \cdot \xi_0 \cdot \eta_0 \cdot \zeta_0 \cdot \pi_0\}$ has properties 1, 2, 3 defining $\perp\!\!\!\perp^i_1$.
   1. Clearly, $\mathsf{d} \star \underline{1} \cdot \pi' \in (\perp\!\!\!\perp^i_1 \setminus \{\mathsf{d} \star \underline{2} \cdot \xi_0 \cdot \eta_0 \cdot \zeta_0 \cdot \pi_0\})$ for every $\pi' \in \Pi^i$.
   2. Suppose that $\xi \star \pi \in \Lambda^i \star \Pi^i$, $\xi \star \pi \succ \xi' \star \pi' \in \perp\!\!\!\perp^i_1$, $\xi' \star \pi' \neq \mathsf{d} \star \underline{2} \cdot \xi_0 \cdot \eta_0 \cdot \zeta_0 \cdot \pi_0$ and that $\xi \star \pi \notin (\perp\!\!\!\perp^i_1 \setminus \{\mathsf{d} \star \underline{2} \cdot \xi_0 \cdot \eta_0 \cdot \zeta_0 \cdot \pi_0\})$.
      From (2), it follows that $\xi \star \pi = \mathsf{d} \star \underline{2} \cdot \xi_0 \cdot \eta_0 \cdot \zeta_0 \cdot \pi_0$ which contradicts $\xi \star \pi \succ \xi' \star \pi'$.
   3. Suppose that two out of the processes $\xi \star \pi, \eta \star \pi, \zeta \star \pi$ are in $\perp\!\!\!\perp^i_1 \setminus \{\mathsf{d} \star \underline{2} \cdot \xi_0 \cdot \eta_0 \cdot \zeta_0 \cdot \pi_0\}$ but that $\mathsf{d} \star \underline{2} \cdot \xi \cdot \eta \cdot \zeta \cdot \pi$ is not. It follows from (3) that:
      $\mathsf{d} \star \underline{2} \cdot \xi \cdot \eta \cdot \zeta \cdot \pi = \mathsf{d} \star \underline{2} \cdot \xi_0 \cdot \eta_0 \cdot \zeta_0 \cdot \pi_0$, i.e. $\xi = \xi_0, \eta = \eta_0, \zeta = \zeta_0$ and $\pi = \pi_0$.
      But this contradicts the hypothesis $\xi_0 \star \pi_0, \eta_0 \star \pi_0 \notin \perp\!\!\!\perp^i_1$. □

Theorem 43. This realizability algebra is coherent.

Proof. Let $\theta \in \mathsf{PL}$ be such that $\theta \star \pi^0 \in \perp\!\!\!\perp^0_0$ and $\theta \star \pi^1 \in \perp\!\!\!\perp^1_1$. Then $\theta \star \pi^0 \in \perp\!\!\!\perp^0_0 \cap \perp\!\!\!\perp^0_1$ which contradicts Lemma 42. □

Lemma 44. $\mathsf{d}\underline{2} \Vdash$ (the boolean algebra $\mathbb{B}2$ has at most four ε-elements).

Proof. We show that $\mathsf{d}\,\underline{2} \Vdash \forall x^{\mathbb{B}2} \forall y^{\mathbb{B}2} (x \neq 0, y \neq 1, x \neq y \to x \wedge y \neq x)$.
Let $i, j \in \{0,1\}$, $\xi \Vdash i \neq 0, \eta \Vdash j \neq 1, \zeta \Vdash i \neq j$ and $\pi \in \|i \wedge j \neq i\|$.
Since $\|i \wedge j \neq i\| \neq \varnothing$, we have $i \leqslant j$. Thus, there are three possibilities for (i, j):
$i = j = 0$; $i = j = 1$; $i = 0, j = 1$.
In each case, two out of the terms ξ, η, ζ realize $\perp$. Thus, we have $\mathsf{d} \star \underline{2} \cdot \xi \cdot \eta \cdot \zeta \cdot \pi \in \perp\!\!\!\perp$. □

Remark. If $\pi \in \Pi \setminus (\Pi^0 \cup \Pi^1)$, then $\xi \star \pi \in \perp\!\!\!\perp$ for every term ξ. Thus, we can remove these stacks and consider only $\Pi^0 \cup \Pi^1$.

https://doi.org/10.1017/S0960129516000050 Published online by Cambridge University Press

We define two individuals in this realizability model:
$\gamma_0 = (\{0\} \times \Pi^0) \cup (\{1\} \times \Pi^1)$; $\gamma_1 = (\{1\} \times \Pi^0) \cup (\{0\} \times \Pi^1)$.
Obviously, $\gamma_0, \gamma_1 \subset \mathbb{B}2 = \{0,1\} \times \Pi$. Now we have:
$\|\forall x^{\mathbb{B}2}(x \not\varepsilon \gamma_0)\| = \Pi^0 \cup \Pi^1 = \|\bot\|$ and therefore $\mathsf{l} \Vdash \neg\forall x^{\mathbb{B}2}(x \not\varepsilon \gamma_0)$.
$\mathsf{d}\underline{0} \Vdash 0 \not\varepsilon \gamma_0$ and $\mathsf{d}\underline{1} \Vdash 1 \not\varepsilon \gamma_0$.
It follows that γ_0, γ_1 are not ε-empty and that every ε-element of γ_0, γ_1 is $\neq 0, 1$.
Therefore:

The Boolean algebra $\mathbb{B}2$ *has exactly four* ε*-elements.*

We have $\xi \Vdash \forall x^{\mathbb{B}2}(x \varepsilon \gamma_0, x \varepsilon \gamma_1 \to \bot)$ for *every* term ξ:
Indeed, let $i, j \in \{0,1\}$; using Lemma 7, we replace the formula $i \varepsilon \gamma_j$, i.e. $\neg(i \not\varepsilon \gamma_j)$, with $\neg(i \not\varepsilon \gamma_j)$ which is $\{\mathsf{k}_\pi \ ; \ \pi \in \Pi^{i+j}\}$. Therefore, we have to check:
$\rho_0 \in \Pi^0, \rho_1 \in \Pi^1 \Rightarrow \xi \star \mathsf{k}_{\rho_0} \cdot \mathsf{k}_{\rho_1} \cdot \pi \in \perp\!\!\!\perp$ which is clear.

In the same way, we get:
$\lambda x \lambda y \lambda z \, z \Vdash \forall x \forall y (x \varepsilon \gamma_i, y \varepsilon \gamma_i, x \neq y \to \bot)$.
It follows that γ_0, γ_1 are singletons and that their ε-elements are the two atoms of $\mathbb{B}2$.

5.1. $\mathbb{B}2$ *has four* ε*-elements and* $\mathbb{B}\kappa$ *is countable*

We now apply to the algebra $\mathcal{A}$ the technique expounded in Section 4, in order to make $\mathbb{B}\kappa$ countable; this gives a realizability algebra $\mathcal{B}$.
In this case, we have $\kappa = \mathbb{N}$, and therefore $\kappa_+ = \mathcal{P}(\kappa) = \mathbb{R}$.

Now, there is an elementary formula which express that the Boolean algebra $\mathbb{B}2$ has four ε-elements, for instance: $\exists x^{\mathbb{B}2}\{x \neq 0, x \neq 1\} \wedge \forall x^{\mathbb{B}2} \forall y^{\mathbb{B}2}(x \neq 1, y \neq 1, x \neq y \to xy = 0)$.
Therefore, the realizability model $\mathcal{N}_\mathcal{B}$ realizes the following two formulas:

($\mathbb{B}2$ has four ε-elements); ($\mathbb{B}\kappa$ is countable);
and therefore also NEAC by Theorem 31.

Let us denote by i_0, i_1 the two atoms of $\mathbb{B}2$; thus, we have $i_1 = 1 - i_0$.

We suppose that $\mathcal{M} \models \mathrm{V} = \mathrm{L}$; thus, there exists on $\kappa_+ = \mathcal{P}(\mathbb{N}) = \mathbb{R}$ a strict well ordering $\lhd$ of type $\aleph_1$. This gives a function from $\mathbb{R}^2$ into $\{0,1\}$, denoted by $(x \lhd y)$, which is defined as follows: $(x \lhd y) = 1 \Leftrightarrow x \lhd y$.
We can extend it to $\mathcal{N}_\mathcal{A}$ and $\mathcal{N}_\mathcal{B}$, which gives a function from $(\mathbb{B}\mathbb{R})^2$ into $\mathbb{B}2$.
From Lemmas 29 and 30, we get:

For $i = i_0$ or i_1, the relation $(x \lhd y) = i$ is a strict total ordering on $\mathbb{B}_i\mathbb{R}$ and one of these two relations is a well ordering;
in order to fix the ideas, we shall suppose that it is for $i = i_0$.
The relation $(x \lhd y) = 1$ is a strict order relation on $\mathbb{B}\mathbb{R}$, which is well founded.
The application $x \mapsto (i_0 x, i_1 x)$ from $\mathbb{B}\mathbb{R}$ onto $\mathbb{B}_{i_0}\mathbb{R} \times \mathbb{B}_{i_1}\mathbb{R}$ is an isomorphism of strictly ordered sets.

It follows from Theorem 26, that each of the sets $\mathbb{B}_{i_0}\mathbb{R}$, $\mathbb{B}_{i_1}\mathbb{R}$ contains a countable subset.
By Corollary 24, there is no surjection from each one of the sets $\mathbb{B}_{i_0}\mathbb{R}, \mathbb{B}_{i_1}\mathbb{R}$ onto the other.
Thus, there is no surjection from $\mathbb{N}$ onto $\mathbb{B}_{i_0}\mathbb{R}$ or onto $\mathbb{B}_{i_1}\mathbb{R}$.
Therefore, the well ordering on $\mathbb{B}_{i_0}\mathbb{R}$ has, at least, the order type $\aleph_1$ in $\mathcal{N}_\mathcal{B}$.

https://doi.org/10.1017/S0960129516000050 Published online by Cambridge University Press

Now, by Theorem 32, every subset of $\beth\mathbb{R}$, which is bounded from above for the ordering $\lhd$, is countable; thus, the same goes for the proper initial segments of $\beth_{i_0}\mathbb{R}$ and $\beth_{i_1}\mathbb{R}$, since these sets are totally ordered and $\beth\mathbb{R}$ is isomorphic to $\beth_{i_0}\mathbb{R}\times\beth_{i_1}\mathbb{R}$.

It follows that the well ordering on $\beth_{i_0}\mathbb{R}$ is at most $\aleph_1$, and therefore exactly $\aleph_1$.
Moreover, *there exists, on* $\beth_{i_1}\mathbb{R}$*, a total ordering, every proper initial segment of which is countable*.

Then, we can apply Theorem 34, to the sets X_{i_0}, X_{i_1} which are the images of $\beth_{i_0}\mathbb{R}, \beth_{i_1}\mathbb{R}$ by the injection from $\beth\kappa_+$ into $\mathbb{R}$, which is given by Theorem 33. By setting $X = X_{i_1}$, we obtain exactly the result (ii) of relative consistency announced in the introduction.

I want to thank the referees, especially the second one for numerous pertinent and helpful remarks and suggestions.

References

Berardi, S., Bezem, M. and Coquand, T. (1998). On the computational content of the axiom of choice. *Journal of Symbolic Logic* **63** (2) 600–622.

Curry, H.B. and Feys, R. (1958). *Combinatory Logic,* North-Holland.

Friedman, H. (1973). The consistency of classical set theory relative to a set theory with intuitionistic logic. *Journal of Symbolic Logic* **38** (2) 315–319.

Friedman, H. (1977). Classically and intuitionistically provably recursive functions. In: Muller, G.H. and Scott, D.S. (eds.) *Higher Set Theory*. Springer Lect. Notes in Math., volume 669 21–27.

Girard, J.-Y. (1971). *Une extension de l'interprétation fonctionnelle de Gödel à l'analyse*. Proc. 2nd Scand. Log. Symp., North-Holland 63–92.

Griffin, T. (1990). A formulæ-as-type notion of control. In: *Conf. record 17th A.C.M. Symp. on Principles of Progr. Languages*, San Francisco, A.C.M. Press 47–58.

Howard, W. (1980). The formulas–as–types notion of construction. In: Seldin, J.P. and Hindley, J.R. (eds.) *Essays on Combinatory Logic, λ-Calculus, and Formalism*, Acad. Press 479–490.

Hyland, J.M.E. (1981). The effective topos. *The L.E.J. Brouwer Centenary Symposium*, Noordwijkerhout 165–216, Stud. Logic Foundations Math., 110, North-Holland, Amsterdam-New York, 1982.

Kreisel, G. (1951). On the interpretation of non-finitist proofs I. *Journal of Symbolic Logic* **16** 248–26.

Kreisel, G. (1952). On the interpretation of non-finitist proofs II. *Journal of Symbolic Logic* **17** 43–58.

Krivine, J.-L. (2001). Typed lambda-calculus in classical Zermelo-Fraenkel set theory. *Archive for Mathematical Logic* **40** (3) 189–205. http://www.pps.univ-paris-diderot.fr/~krivine/articles/zf_epsi.pdf

Krivine, J.-L. (2003). Dependent choice, 'quote' and the clock. *Theoretical Computer Science* **308** 259–276. http://www.pps.univ-paris-diderot.fr/~krivine/articles/quote.pdf

Krivine, J.-L. (2007). Realizability: A machine for Analysis and set theory. Geocal'06 (febr. 2006 - Marseille); Mathlogaps'07 (June 2007 - Aussois). http://www.pps.univ-paris-diderot.fr/~krivine/articles/Mathlog07.pdf

Krivine, J.-L. (2009). Realizability in classical logic. In: *Interactive Models of Computation and Program Behaviour*. Panoramas et synthèses, volume 27, Société Mathématique de France 197–229. http://www.pps.univ-paris-diderot.fr/~krivine/articles/Luminy04.pdf

https://doi.org/10.1017/S0960129516000050 Published online by Cambridge University Press

Krivine, J.-L. (2011). Realizability algebras: A program to well order $\mathbb{R}$. *Logical Methods in Computer Science* **7** (3:02) 1–47.

Krivine, J.-L. (2012). Realizability algebras II: New models of ZF + DC. *Logical Methods in Computer Science* **8** (1:10) 1–28.

Miquel, A. (2011). Forcing as a program transformation. In: *Logic in Computer Science (LICS'11)*, Toronto, Canada 197–206.

https://doi.org/10.1017/S0960129516000050 Published online by Cambridge University Press