Bull. Aust. Math. Soc. **112** (2025), 387–388
doi:10.1017/S0004972725100117

CHARACTER VARIETIES, EARTHQUAKES AND ESSENTIAL SURFACES

GRACE SHANTI GARDEN

(Received 24 April 2025; first published online 27 June 2025)

2020 *Mathematics subject classification*: primary 20H10; secondary 30F60, 57K31, 57K32, 57M05.

Keywords and phrases: character varieties, Teichmüller theory, earthquakes, essential surfaces.

We study low-dimensional manifolds through their character variety from two perspectives. Character varieties of low-dimensional manifolds are a rich area of study that reflect geometric and topological information of the underlying manifold. Given a manifold M and an algebraic group G, the G-character variety is loosely the space of conjugacy classes of representations from the fundamental group of M to G, with some conjugacy classes identified so that the resulting space is a variety. We are particularly interested in the situation when M is a hyperbolic surface or 3-manifold and G is SL_2. This is because of the relationship between elements of the character variety and the orientation preserving isometry groups of the hyperbolic plane ($\mathrm{PSL}_2(\mathbb{R})$) and hyperbolic space ($\mathrm{PSL}_2(\mathbb{C})$).

The first perspective uses Teichmüller space and earthquakes. Teichmüller space of a surface is the space of all finite-area complete hyperbolic structures of the surface and can be naturally identified with a component of the associated $\mathrm{SL}_2(\mathbb{R})$-character variety [5]. There is a natural action on Teichmüller space by earthquakes, which have proved to be powerful in the exploration of these hyperbolic structures [6]. We derive explicit forms of the earthquake deformations on Teichmüller space associated with simple closed curves of the once-punctured torus with both algebraic and geometric interpretations [2]. Examining the limiting behaviour gives insight into earthquakes about measured geodesic laminations, of which simple closed curves are a special case [2].

The second perspective uses essential surfaces and the theory of Culler and Shalen. Essential surfaces are orientable, incompressible and boundary-incompressible surfaces in a 3-manifold, and tell us about the inherent geometry and topology of the manifold. Thurston's hyperbolisation theorem implies these essential surfaces

Thesis submitted to the University of Sydney in July 2024; degree approved on 6 November 2024; supervisor Stephan Tillman, auxiliary supervisor Anne Thomas.

© The Author(s), 2025. Published by Cambridge University Press on behalf of Australian Mathematical Publishing Association Inc.

https://doi.org/10.1017/S0004972725100117 Published online by Cambridge University Press

are intimately connected with hyperbolic structures. In the seminal work of Culler and Shalen, essential surfaces in 3-manifolds are associated to ideal points of their $SL_2(\mathbb{C})$-character varieties [1]. We lay a general foundation for this theory in arbitrary characteristic by using the same approach instead over the $SL_2(\mathbb{F})$-variety of characters for $\mathbb{F}$ an arbitrary algebraically closed field [4]. We provide several applications of the extended theory, including bounds on dimension of the character variety, the A-polynomial, prime decomposition and JSJ decomposition [3, 4]. We give examples where new essential surfaces are detected in positive characteristic and where an essential surface is never detected in any characteristic.

References

[1] M. Culler and P. B. Shalen, 'Varieties of group representations and splittings of 3-manifolds', *Ann. of Math. (2)* **117**(1) (1983), 109–146.
[2] G. S. Garden, 'Earthquakes on the once-punctured torus', *Exp. Math.* **33**(4) (2024), 781–907.
[3] G. S. Garden, B. Martin and S. Tillmann, 'Essential tori in 3-manifolds not detected in any characteristic', *Geom. Dedicata* **219** (2025), Article no. 64.
[4] G. S. Garden and S. Tillmann, 'An invitation to Culler–Shalen theory in arbitrary characteristic', Preprint, 2024, arXiv:2411.06859.
[5] W. Goldman, 'Topological components of spaces of representations', *Invent. Math.* **93** (1988), 557–607.
[6] W. P. Thurston, 'Earthquakes in two-dimensional hyperbolic geometry', in: *Low-Dimensional Topology and Kleinian Groups (Coventry/Durham, 1984)*, London Mathematical Society Lecture Note Series, 112 (ed. D. B. A. Epstein) (Cambridge University Press, Cambridge, 1986), 91–112.

GRACE SHANTI GARDEN,
Institut de Mathématiques de Jussieu – Paris Rive Gauche,
Sorbonne Université, Paris, France
e-mail: grace.garden@imj-prg.fr

https://doi.org/10.1017/S0004972725100117 Published online by Cambridge University Press