Zygote

cambridge.org/zyg

Inseminating dose for the artificial fertilization of *Brycon amazonicus* (Teleostei: Characidae)

Rosilane Gomes de Souza de Oliveira and Marle Angélica Villacorta-Correa

Federal University of Amazonas (UFAM), Faculty of Agricultural Sciences, Department of Fisheries Sciences, Manaus, Amazonas, Brazil

Summary

Knowledge of the sperm–oocyte ratio in fish fertilization serves as the basis for studies on artificial reproduction and gamete manipulation. The aim of this study was to determine the minimum insemination dose for *Brycon amazonicus* oocyte fertilization. Female and male gametes were used and tested with the following doses of spermatozoa oocyte^{-1} ml^{-1}: 10,000, 20,000, 40,000, 60,000 and 80,000 (in triplicate). Fertilization rates were calculated and estimated from the regression equation by applying the segmented regression model 'Linear Response Plateau' to determine the appropriate proportion of gametes. Based on the equation $\hat{Y} = 14.3415 + 0.0007836X$, the fertilization rate increased up to 63.34% as it reached a plateau with a proportion of 62,524 spermatozoa oocyte^{-1} ml^{-1}, which is the minimum insemination dose recommended for artificial insemination of the species.

Short Communication

Cite this article: Oliveira RGS and Villacorta-Correa MA (2021) Inseminating dose for the artificial fertilization of *Brycon amazonicus* (Teleostei: Characidae). *Zygote*. **29**: 252–255. doi: 10.1017/S096719942000074X

Received: 17 April 2020
Revised: 29 September 2020
Accepted: 21 November 2020
First published online: 15 January 2021

Keywords:
Artificial reproduction; Fertilization rate; Matrinxã; Oocytes; Spermatozoa

Author for correspondence:
Rosilane Gomes de Souza de Oliveira. Federal University of Amazonas (UFAM), Faculty of Agricultural Sciences, Department of Fisheries Sciences, Av. Gen. Rodrigo Octávio Jordão Ramos, 3000, CEP: 69077-000, Manaus, Amazonas, Brazil. Tel: +55 923305 1797. Fax: +55 923305 1797.
E-mail: rosilanegso@gmail.com

Introduction

The purpose of artificial insemination in fish is to successfully fertilize the largest number of fertile oocytes with the minimum number of males, bearing in mind the need to maintain a satisfactory gender balance (Billard *et al.*, 1995). Therefore, there is a need to establish an inseminating dose (or sperm ratio per oocyte) for each species that can be defined as the amount of sperm needed to fertilize an oocyte (Chereguini *et al.*, 1999). This dose can improve the use of breeders by increasing their reproductive efficiency through artificial reproduction, as it allows rational management of semen, reducing the number of animals and their maintenance cost.

Use of small or excessive sperm concentrations results in lower fertilization rates (Levanduski and Cloud, 1988; Ninhaus-Silveira *et al.*, 2006; Shimoda *et al.*, 2007). The insemination dose for fish, unlike terrestrial animals, is relatively higher and varies between species (Beirão *et al.*, 2019). In addition, variations in sperm to oocyte ratio may occur based on oocyte characteristics or semen quality (Suquet *et al.*, 1995), which directly interfere with fertilization potential (Rurangwa *et al.*, 2004).

Brycon amazonicus, know as matrinxã, is a neotropical fish whose reproductive biology is described, taking into account that many aspects are species specific. Reproduction in *B. amazonicus* does not always follow a standardized hormone induction protocol, with frequent failures in spawning, fertilization and hatching rates within and between reproductive periods (Romagosa *et al.*, 2001; Pardo-Carrasco *et al.*, 2002; Arias, 2006; Hainfellner *et al.*, 2012). Females have high relative fecundity (Arias, 2006) and the same female can spawn at least twice in the same reproductive cycle when given hormonal therapies (Honczaryk and Inoue, 2009). However, their large-scale production is limited by their intense cannibalistic behaviour in the early stages of development, with losses of up to 99% (Bernardino *et al.*, 1993) and a reproductive seasonality that restricts the availability of juveniles to a certain period of the year (Zaniboni-Filho *et al.*, 1988). Therefore, it is important to maximize the use of available gametes to produce a greater number of viable embryos, especially considering that *B. amazonicus* is an aggressive fish and highly sensitive to the entire process of artificial reproduction, and which justifies avoidance of gamete wastage and unnecessary use of breeders.

There have been studies for this species on its reproductive biology (Zaniboni-Filho *et al.*, 1988; Romagosa *et al.*, 1999), covering artificial reproduction (Bernardino *et al.*, 1993; Romagosa *et al.*, 2001), cryoprotectants, cryopreservation and conservation of gametes (Cruz-Casallas *et al.*, 2006; Ninhaus-Silveira *et al.*, 2006; Velasco-Santamaría *et al.*, 2006; Bashiyo-Silva *et al.*, 2015). However, the minimum insemination dose for the species has not yet been reported and unspecified volumes are being used for artificial fertilization, without any concern for better use of gametes. We highlight the importance of this work not only for artificial reproduction of the species, but also for tests in applied biotechnology, as manipulation of gametes, control of fertilization, polyploidy and androgenesis. The objective of this work was to determine the insemination dose for artificial fertilization of *B. amazonicus* oocytes, to attain a higher reproductive efficiency through the rational use of breeders.

© The Author(s), 2021. Published by Cambridge University Press.

CAMBRIDGE UNIVERSITY PRESS

Materials and methods

Broodstock maintenance and hormonal induction

We used 4-year-old wild specimens that had been maintained since 6 months of age at the Experimental Farm Aquaculture Station of the Federal University of Amazonas (UFAM) Manaus, Amazonas, Brazil (2°38′56.1′′S, 60°03′14.7′′W). They were placed in earthen ponds (550 m^2, ca. 0.1 fish m^2) and fed twice daily to their apparent satiation with a commercial fish feed that contained 32% crude protein.

Four breeders were selected in December 2014. Two males that emitted semen under mild abdominal pressure and two females with hyperemic genital papilla, soft, bulging abdomens (Woynarovich and Horváth, 1983). They had >50% oocytes that presented eccentric germinal vesicles (Arias, 2006), which were observed after ovarian biopsy by insertion of a flexible catheter into the genital pore, followed by gentle aspiration and immersion of oocytes in Serra's fixative solution for a few minutes (Stoeckel, 2000). The animals were separated by sex in two circular fibreglass tanks with a capacity of 3000 L each. Water was constantly renewed and aerated, and an average temperature of 27.7 ± 1.0°C and pH 7.9 ± 0.2 was maintained.

Females (1.66 ± 0.16 kg) received 5.5 mg kg^{-1} carp pituitary extract divided into three doses and males (1.31 ± 0.69 kg) received a single dose of 2.0 mg kg^{-1} carp pituitary extract (Zaniboni-Filho and Barbosa, 1996), as intramuscular injections in the base of the dorsal fin. The animals had never been previously subjected to hormone therapy.

Gamete collection and evaluation

Each specimen was removed from the tank, and placed in a container containing 50 mg L^{-1} eugenol solution (Vidal *et al.*, 2007) to reduce the stress associated with this procedure and to collect the gametes.

Semen were collected 12 h after hormonal injection in ml-graduated tubes (16.5 ± 7.0 ml). Sperm motility was evaluated subjectively (94.16 ± 2.5% of mobile sperm and 33.66 ± 2.9 s of post-activation motility) for each sample in two steps as described by Billard and Cosson (1992). The first step consisted of diluting the semen in a 1% NaCl solution (1:40, semen:diluter) and, in the second step, the previously diluted sample was activated with distilled water (1:20, diluted sample:water), on a microscope slide coated with 0.05% bovine serum albumin to prevent sperm from adhering to the slide, under an optical microscope, ×40 magnification objective. Subsequently, a semen pool was constituted. A Neubauer haematimetric chamber was used to determine the pool sperm concentration (1.63×10^{10} sperm ml^{-1}), as recommended by the Brazilian College of Animal Reproduction (Colégio Brasileiro de Reprodução Animal, 2013).

Oocytes were released 5 h after the last hormone dose (ca. 140 accumulated thermal units) into plastic containers.

Insemination doses, artificial fertilization and fertilization rate

The insemination doses were prepared according to the methodology used for *Misgurnus anguillicaudatus* (Yasui *et al.*, 2009), with some dosage modifications for *B. amazonicus*, based on two previous experiments. Sperm were diluted in 1% NaCl solution to attain concentrations of 0.75×10^7, 1.51×10^7, 3.02×10^7, 4.53×10^7 and 6.04×10^7 spermatozoa ml^{-1}. This solution keeps the sperm still and viable until insemination. We pipetted 150 µl of the oocytes (755.2 ± 19.6 oocytes) into 18 glass beakers (50 ml) using a clipped pipette tip. Each oocyte mass was immediately inseminated with 50 µl of the previously diluted sperm solution. After insemination, gametes were activated with 800 µl distilled water, homogenized for 1 min and then hydrated with 20 ml of incubator water (28.9°C). These dilutions resulted in insemination doses of 10,000, 20,000, 40,000, 60,000 and 80,000 spermatozoa $oocyte^{-1}$ ml^{-1} (three replicates for each dose were made). A mass of oocytes mixed with only 50 µl of 1% NaCl solution was maintained as a control group.

The fertilized oocytes were transferred to 18 cylindrical incubators (10 × 12 cm), with screened bottoms, properly identified and packed in a 3000 L fibreglass water tank with constant water and aeration renewal, at a temperature of 28.9 ± 1.0°C and a pH of 7.8 ± 0.1. Fertilization rates were calculated relative to the total amount of eggs ca. 6 h after artificial insemination, under a stereomicroscope, when the eggs were in the blastopore closure phase (Nakaghi *et al.*, 2014). In this phase of development, eggs that were whitish or broken were considered unfertilized or dead, and eggs that were translucent with perfect embryos inside the chorion were considered fertilized.

Statistical analysis

The regression equation that fitted the data for the variation in fertilization rates was estimated as a function of spermatozoa count per oocyte by applying the segmented regression model 'Linear Response Plateau' in the R statistical program. The appropriate ratio of spermatozoa to oocyte was calculated using this equation. Data were first tested for normality (Shapiro–Wilk test). Data are presented as the mean ± standard error. A P-value < 0.05 was considered to be significant.

Results

The estimated regression equation was $\hat{Y} = 14.3415 + 0.0007836X$ ($P < 0.05$), where $\hat{Y}$ = fertilization rate and X = spermatozoa per oocyte ($P < 0.05$; $R^2 = 0.827$). The fertilization rate increased linearly until it reached the maximum rate of 63.34%, remaining constant and starting to plateau at a ratio of 62,524 spermatozoa $oocyte^{-1}$ ml^{-1} (Fig. 1). This starting point of the plateau represents the minimum ratio required to achieve maximum fertility. Although the insemination dose increased after this point, the fertilization rate remained constant on a plateau of 63.34%.

Discussion

Knowledge of the appropriate inseminating dose is important for the development of semen and/or oocyte cryopreservation programmes, intended either for the conservation of genetic biodiversity of the species or even for breeding programmes in farms, in addition to the possibility of gamete optimization in aquaculture stations (Denniston *et al.*, 2000).

The quality of gametes observed in this study corroborates the results of other authors (Cruz-Casallas *et al.*, 2006, Ninhaus-Silveira *et al.*, 2006) and fertilization rates were similar to those reported by other authors (Romagosa *et al.*, 2001; Arias 2006). Therefore, the animals used were fit and the gamete quality parameters can be considered satisfactory for the artificial reproduction process.

According to Rurangwa *et al.* (2004), fertilization rates are related to sperm motility and may be influenced by the inseminating dose used, but it is poorly reported for fish species and presents

https://doi.org/10.1017/S096719942000074X Published online by Cambridge University Press




Figure 1. Values for oocyte fertilization rates of *Brycon amazonicus* as a function of spermatozoa oocyte^{-1} ml^{-1}.

great variation between species, which can probably be related to the characteristics of gametes, such as oocyte diameter, speed of spermatozoa, uptime and the distance travelled by the spermatozoa to reach the micropyle of the oocyte (Suquet *et al.*, 1995; Lahnsteiner 2000; Gage *et al.*, 2004; Bombardelli *et al.*, 2006).

Suquet *et al.* (1995) found that the insemination dose for *Scophthalmus maximus* was 6000 spermatozoa oocyte^{-1} with 87.3% fertilization. These authors concluded that this species needed a small amount of semen because of the size of its oocytes (0.91–1.2 mm) and also the semen had a long activation time. *B. amazonicus* oocytes are small (1.11–1.30 mm in diameter; Nakaghi *et al.*, 2014), as reported for another species of the same genus, *B. insignis* (1.3 mm; Shimoda *et al.*, 2007). However, the insemination dose for fertilization of *B. insignis* oocytes (314,481 spermatozoa oocyte^{-1}) was much higher than that found in our study, which suggested that other factors may influence interspecifically, such as contact time between gametes and the fertilization protocol used (Rurangwa *et al.*, 2004).

According to Cruz-Casallas and Velasco-Santamaría (2004) a proportion of 50,000 fresh sperm or 75,000 thawed sperm per oocyte is sufficient to obtain maximum fertilization percentages in artificial insemination of *Brycon siebenthalae*. In this study, the fertilization results showed effect pertinent to the treatments, presenting positive linear behaviour up to the proportion of 62,524 spermatozoa oocyte^{-1}. The fertilization rates reached a plateau at this ratio, and had no effect at the higher doses, remaining constant thereafter.

For *Brycon insignis* (Shimoda *et al.*, 2007), the fertilization rate increased to 87.8% and remained on this plateau at a ratio of 314,481 spermatozoa oocyte^{-1}. Other authors have also found this plateau from a given ratio of spermatozoa to oocyte on the fertilization rate of some species, such as *Astyanax altiparanae* (75.4%, 2390 spermatozoa oocyte^{-1}, Pereira-Santos *et al.*, 2017), *Colossoma macropomum* (84%, 102,486 spermatozoa oocyte^{-1}; Leite *et al.*, 2013), *Misgurnus anguillicaudatus* (62.84%, 684.42 spermatozoa oocyte^{-1}; Yasui *et al.*, 2009) *Salminus brasiliensis* (57.1%, 30,722 spermatozoa oocyte^{-1}; Sanches *et al.*, 2009) *Rhamdia quelen* (86.68%, 89,472 spermatozoa oocyte^{-1}; Bombardelli *et al.*, 2006).

Determination of insemination dose is important for evaluation of semen cryopreservation procedures and better economical use of large-scale artificial reproduction procedures, as lower insemination doses may trigger lower fertilization rates, and excess sperm reflects the inefficient use breeders. Therefore, fertilization tests, if performed with excessive amounts of semen, may not provide suitable conditions to identify the best treatment to be used, as eventual differences between methodologies may be attenuated by increase in the ratio of sperm to oocyte (Shimoda *et al.*, 2007). In addition, sperm economy will multiply the reproductive potential of higher performing or genetic breeding stock and, as a result, will benefit semen banks for species conservation and will also enable higher profitability for breeding companies in aquaculture.

Considering that the sperm concentration of *B. amazonicus* is ca. 1.4×10^9 ml^{-1} (Cruz-Casallas *et al.*, 2006, Ninhaus-Silveira *et al.*, 2006) and the number of oocytes per gram is estimated at 1500 cells and, based on the results found, it can be recommended that 7 ml of semen per 100 g of oocytes would be sufficient for good fertilization rates in this species. However, the use of a higher proportion of sperm per oocyte should be considered, if the semen has been cryopreserved, and to evaluate sperm concentration and quality under favourable incubation conditions. The sperm to oocyte ratio should only be considered as a reference value and not as an absolute or unalterable value for fertilization trials (Beirão *et al.*, 2019).

The insemination dose for fertilization of *B. amazonicus* oocytes will be useful for optimizing the use of breeding stock in the artificial propagation process, even more considering that this species is an extremely aggressive fish during reproductive management. This study will be the basis for conducting future studies of induced reproduction and gamete manipulation of fish.

In conclusion, the highest oocyte fertilization rates for *B. amazonicus* may be obtained with the use of a proportion of 62,524 spermatozoa oocyte^{-1} ml^{-1}.

Acknowledgements. The authors thank George Shigueki Yasui (University of São Paulo – USP) for his scientific and methodological contributions. We are also grateful to Eduardo Antônio Sanches and Rafael Vilhena Reis Neto (São Paulo State University – UNESP) for contribution in data analysis.

Data availability statement. Research data are not shared.

Financial support. This study was funded by Fundação de Amparo à Pesquisa no Amazonas – FAPEAM (project grant no. 1271/2012) and Financiadora de Estudos e Projetos – FINEP/Projeto Desenvolvimento da Aquicultura e Recursos Pesqueiros da Amazônia – DARPA (protocol no. 01.09.0472.00).

Conflict of interest. The authors report no conflict of interest

Ethical approval. All procedures performed in studies involving animals were in accordance with the ethical standards of the institution or practice at which the studies were conducted (Ethics Committee on Animal Use 'CEUA' of Federal University of Amazonas, protocol no. 019/2014). The Brazilian College of Animal Experimentation (COBEA) guidelines for the care and use of laboratory animals were followed.

References

Arias JA (2006). Estado actual del conocimiento sobre el yamú, *Brycon amazonicus*. *Rev Colomb Cienc Pec* **19**, 125–33.

Bashiyo-Silva C, da Silva Costa R, de Castro Ribeiro D, Senhorini JA, Veríssimo-Silveira R and Ninhaus-Silveira A (2015). Hormonal induction of *Brycon cephalus* (Characiformes, Characidae) to spermiation using D-ala6, pro9net-mGnRH + metoclopramide. *Zygote* **24**, 319–25.

Beirão J, Boulais M, Gallego V, O'Brien JK, Peixoto S, Robeck TR and Cabrita E (2019). Sperm handling in aquatic animals for artificial reproduction. *Theriogenology* **133**, 161–78.

Bernardino G, Senhorini JA, Fontes NA, Bock CL and Mendonça JOJ (1993). Propagação artificial do matrinchã *Brycon cephalus* (Günther, 1869) (Teleostei, Characidae). *Boletim Técnico CEPTA* **6**, 1–9.

Billard R and Cosson MP (1992). Some problems related to the assessment of sperm motility in freshwater fish. *J Exp Zool* **261**, 122–31.

https://doi.org/10.1017/S096719942000074X Published online by Cambridge University Press

Billard R, Cosson J, Perchec G and Linhart O (1995). Biology of sperm and artificial reproduction in carp. *Aquaculture* **129**, 95–112.

Bombardelli RA, Mörschbächer EF, Campagnolo R, Sanches EA and Syperreck MA (2006). Dose inseminante para fertilização artificial de ovócitos de jundiá cinza, *Rhamdia quelen* (Quoy & Gaimardm, 1824). *Rev Bras Zootecn* **35**, 1251–7.

Chereguini O, De La Banda IG, Rasines I and Fernandez A (1999). Artificial fertilization in turbot, *Scophthalmus maximus* (L.): different methods and determination of the optimal sperm–egg ratio. *Aquac Res* **30**, 319–24.

Colégio Brasileiro de Reprodução Animal (2013). *Manual para exames andrológicos e avaliação de sêmen animal*. 3rd edn. Belo Horizonte: CBRA. 104 pp.

Cruz-Casallas PE, Medina-Robles VM and Velasco-Santamaría YM (2006). Evaluación de diferentes crioprotectores para la crioconservación de espermatozoides de yamú (*Brycon amazonicus*). *Rev Colomb Cienc Pec* **19**, 152–9.

Cruz-Casallas PE and Velasco-Santamaría YM (2004). Crioconservación de semen de yamú (*Brycon siebenthalae*): calidad seminal, dosis inseminante y sistemas de empaque. In Memorias II Congreso Colombiano de Acuicultura. Universidad de los Llanos (IALL). Villavicencio, Colombia. 27–34 pp.

Denniston RS, Michelet S and Godke RA (2000). Principles of cryopreservation. In: Tiersch TR and Mazik PM (eds). *Cryopreservation in aquatic species*. World Aquaculture Society, Baton Rouge, pp 59–74.

Gage MJ, Macfarlane CP, Yeates S, Ward RG, Searle JB and Parker GA (2004). Spermatozoal traits and sperm competition in Atlantic salmon: relative sperm velocity is the primary determinant of fertilization success. *Curr Biol* **14**, 44–7.

Hainfellner P, De Souza TG, Muñoz ME, Freitas GA and Batlouni SR (2012). Spawning failure in *Brycon amazonicus* may be associated with ovulation and not with final oocyte maturation. *Arq Bras Med Vet Zoo* **64**, 515–7.

Honczaryk A and Inoue LAKA (2009). Produção comercial de alevinos de matrinxã na Amazônia Ocidental. *Embrapa Amazônia Ocidental-Circular Técnica* (INFOTECA-E).

Lahnsteiner F (2000). Semen cryopreservation in the Salmonidae and in the Northern pike. *Aquac Res* **31**, 245–58.

Leite LV, Melo MA, Oliveira FC, Pinheiro JP, Campello CC, Nunes JF and Salmito-Vanderley CS (2013). Determination of insemination dose and embryonic development in the artificial fertilization of tambaqui (*Colossoma macropomum*). *Arq Bras Med Vet Zoo* **65**, 421–9.

Levanduski MJ and Cloud JG (1988). Rainbow trout (*Salmo gairdneri*) semen: effect of non-motile sperm on fertility. *Aquaculture* **75**, 171–9.

Nakaghi LSO, Neumann E, Faustino F, Mendes JMR and de Braga FM (2014). Moments of induced spawning and embryonic development of *Brycon amazonicus* (Teleostei, Characidae). *Zygote* **22**, 549–57.

Ninhaus-Silveira A, Foresti F, Veríssimo-Silveira R and Senhorini JA (2006). Seminal analysis, cryogenic preservation, and fertility in matrinxã fish, *Brycon cephalus* (Günther, 1869). *Braz Arch Biol Techn* **49**, 651–9.

Pardo-Carrasco S, Suarez-Mahecha H, Muñoz-Lara D, Arias-Castellanos J and Hernando GIL (2002). Inducción de la ovulación y del desove del yamú, *Brycon siebenthale*, con implantes de mGnRH-a. *Bol Inst Pesca* **28**, 19–24.

Pereira-Santos M, Shimoda E, de Andrade AFC, Silva LA, Fujimoto T, Senhorini JA, Yasui GS and Nakaghi LSO (2017). Grooves surrounding the micropyle decrease the inseminating dose in fish. *Zygote* **25**, 731–9.

Romagosa E, Narahara MY, Borella MI, Parreira SF and Fenerich-Verani N (1999). Ultrastructure of the germ cells in the testis of matrinxã, *Brycon cephalus* (Teleostei, Characidae). *Tissue Cell* **31**, 540–4.

Romagosa E, Narahara M, Borella M and Fenerich-Verani N (2001). Seleção e caracterização de fêmeas de matrinxã, *Brycon cephalus*, induzidas a reprodução. *Bol Inst Pesca* **27**, 139–47.

Rurangwa E, Kime DE, Ollevier F and Nash JP (2004). The measurement of sperm motility and factors affecting sperm quality in cultured fish. *Aquaculture* **234**, 1–28.

Sanches EA, Bombardelli RA, Baggio DM and Souza BED (2009). Dose inseminante para fertilização artificial de ovócitos de dourado. *Rev Bras Zootecn* **38**, 2091–8.

Shimoda E, Andrade DR, Vidal Júnior MV, Godinho HP and Yasui GS (2007). Determination of the optimum ratio of spermatozoa per oocyte of the piabanha *Brycon insignis* (Pisces-Characidae). *Arq Bras Med Vet Zoo* **59**, 877–82.

Stoeckel JN (2000). A method for viewing the germinal vesicle in oocytes of commercial catfishes. *N Am J Aquacult* **62**, 240–7.

Suquet M, Billard R, Cosson J, Normant Y and Fauvel C (1995). Artificial insemination in turbot (*Scophthalmus maximus*): determination of the optimal sperm to egg ratio and time of gamete contact. *Aquaculture* **133**, 83–90.

Velasco-Santamaría YM, Medina-Robles VM and Cruz-Casallas PE (2006). Cryopreservation of yamú (*Brycon amazonicus*) sperm for large scale fertilization. *Aquaculture* **256**, 264–71.

Vidal LVO, Furuya WMM, Graciano TS, Schamber CR, da Silva LCR, dos Santos LD and de Souza SR (2007). Eugenol como anestésico para juvenis de matrinxã ('*Brycon cephalus*'). *Rev Bras Saúd Prod Anim* **8**, 335–42.

Woynarovich E and Horváth L (1983). *A propagação artificial de peixes de águas tropicais; manual de extensão*. Brasília: FAO/CODEVASF/CNPq.

Yasui GS, Arias-Rodriguez L, Fujimoto T and Arai K (2009). A sperm cryopreservation protocol for the loach *Misgurnus anguillicaudatus* and its applicability for other related species. *Anim Reprod Sci* **116**, 335–45.

Zaniboni-Filho E and Barbosa NDC (1996). Priming hormone administration to induce spawning of some Brazilian migratory fish. *Rev Bras Biol* **56**, 655–9.

Zaniboni-Filho E, Carvalho JL, Villacorta-Correa MA and Rezende EK (1988). Caracterização morfológica da matrinxã, *Brycon cephalus* (Günther, 1869) (Teleostei: Characidae). *Revista Brasileira Biologia* **48**, 41–50.

https://doi.org/10.1017/S096719942000074X Published online by Cambridge University Press