The Journal of Agricultural Science

www.cambridge.org/ags

Crops and Soils Review

Cite this article: Ablormeti FK, Honger JO, Lutuf H, Obeng J, Muntala A, Osabutey AF, Ochar K, Otoo BA, Sossah FL, Ndede Yankey E, Dofuor AK, and Aidoo OF (2025). Threats to global food security from emerging phytoplasma crop diseases. *The Journal of Agricultural Science* **163**, 365–380. https://doi.org/10.1017/S0021859625100178

Received: 2 December 2024
Revised: 9 June 2025
Accepted: 25 June 2025

Keywords:
detection methods; food security; Phytoplasmas; plant diseases; surveillance

Corresponding author: Aboagye Kwarteng Dofuor; Email: akdofuor@uesd.edu.gh

© The Author(s), 2025. Published by Cambridge University Press.

Threats to global food security from emerging phytoplasma crop diseases

Fred Kormla Ablormeti[1], Joseph Okani Honger[2], Hanif Lutuf[3], Joshua Obeng[3,4], Abdulai Muntala[5], Angelina Fathia Osabutey[6], Kingsley Ochar[7], Bernice Araba Otoo[8], Frederick Leo Sossah[1], Egya Ndede Yankey[1], Aboagye Kwarteng Dofuor[9] and Owusu Fordjour Aidoo[9,10]

[1]Council for Scientific and Industrial Research, Oil Palm Research Institute, Coconut Research Programme, P. O. BOX, 245, Sekondi, Ghana; [2]Soil and Irrigation Research Centre, College of Basic and Applied Sciences, School of Agriculture, University of Ghana, Accra, Ghana; [3]Crop Protection Division, Oil Palm Research Institute, Council for Scientific and Industrial Research, Kade, Ghana; [4]College of Agriculture, Tennessee State University, Nashville, TN, USA; [5]Department of Horticultural and Crop Production, University of Energy and Natural Resources, Sunyani, Ghana; [6]Department of Entomology, Agricultural Research Organization, The Volcani Institute, Israel; [7]Council for Scientific and Industrial Research-Plant Genetic Resources Research Institute, Bunso, Eastern Region, Ghana; [8]Department of Physical and Mathematical Sciences, University of Environment and Sustainable Development, Somanya, Ghana; [9]Department of Biological Sciences, University of Environment and Sustainable Development, Somanya, Ghana and [10]Department of Entomology, Washington State University, Pullman, WA, 99164, USA

Abstract

Extensive damage to over 1000 plant species, including food crops, oil and industrial crops, vegetables, fruit trees, ornamentals, fodder species and weeds, has been caused by emerging phytoplasma-mediated diseases, thereby posing significant threat to global food security. Multiple factors, including environmental changes, invasion pathways, vector transmission and the emergence of new pathogen lineages, contribute to the spread of these diseases. Effective management requires stable, long-term strategies to safeguard plant health. Key approaches include comprehensive loss assessments, integration of climate change impacts, predictive modelling, enhanced disease surveillance, and improved detection techniques targeting phytoplasmas. This review highlights phytoplasma-associated plant diseases, emerging pathogen threats, and the factors facilitating their spread, alongside methods for surveillance and detection. In addition, case studies and global collaborative efforts are discussed. Finally, we outline future research priorities aimed at improving the management of phytoplasma-induced plant diseases.

Introduction

The United Nations Department of Economic and Social Affairs, Population Division, projected in 2017 that the world's population, currently 7.6 billion, will rise to 8.6 billion in 2030, 9.8 billion in 2050 and 11.2 billion in 2100 (UN Department of Economic and Social Affairs, 2021). A meta-analysis revealed that global food demand will increase by 35–56% from 2010 to 2050, while the number of people at risk of hunger could fluctuate between −91% and +8% (van Dijk *et al.*, 2021). Substantial yield reductions due to crop pests and diseases are evident, with average losses of approximately 40% in wheat, rice, maize, potato, and soybean, significantly contributing to food insecurity (Savary *et al.*, 2019).Yield losses of up to 100% are possible without effective control measures, leading to widespread destruction. Globally, plant diseases caused by viruses, nematodes, bacteria, and fungi result in annual losses of approximately $220 billion (Savary *et al.*, 2019). In addition, Germany lost €25 million, and Italy lost approximately €100 million due to a phytoplasma epidemic in apple trees in 2001 (Strauss, 2009).

Phytoplasmas are wall-less bacteria belonging to the class Mollicutes that reside in the phloem sieve elements of diseased plants (Lee *et al.*, 1998). Taxonomy and identification of phytoplasmas rely on molecular methods and gene sequences, as these pathogens cannot be cultured in cell-free conditions. Currently, 48 '*Candidatus* Phytoplasma' species have been named based on several criteria, including <98.65% sequence identity in the 16S rRNA gene, <95–96% whole genome similarity, or ecological separation (Bertaccini *et al.*, 2022; Wei and Zhao, 2022). Wei and Zhao (2022) classified genetically diverse phytoplasmas into 37 groups and over 150 subgroups using restriction fragment length polymorphism (RFLP) profiles of the F2nR2 region of the 16S rRNA gene. Many phytoplasmas have been linked to newly emerging diseases worldwide in recent years (Huang *et al.*, 2023).

Phytoplasmas-associated diseases affect over 1000 plant species worldwide (Hiruki and Wang, 2004; Maejima *et al.*, 2014; Wang *et al.*, 2024). As a threat to food security, their infestation may cause substantial yield losses of up to 100% in cucumber, tomato, pepper, potato and Navratil crops (Bogoutdinov *et al.*, 2008; Rao and Kumar, 2017). Over the last four decades, millions of coconut palm trees in the Caribbean have been wiped out by a deadly lethal yellowing disease (LYD) associated with phytoplasmas (Brown *et al.*, 2006). In Jamaica alone, more than seven million palm trees died by 1980 due to LYD (Roca de Doyle, 2001). In Africa, similar diseases were observed and collectively referred to as lethal yellowing-like disease (Eden-Green, 1997). Eight million coconut palms, or 38 % of Tanzania's total hectarage, have been destroyed due to phytoplasma-associated-lethal disease since the 1960s (Mugini, 2002). Furthermore, an outbreak of LYD in Côte d'Ivoire damaged 350 hectares of coconut and destroyed 12,000 metric tonnes of copra annually, with an additional 7,000 hectares at risk (Arocha-Rosete *et al.*, 2014).

The incidence of phytoplasmas-associated plant diseases has been on the rise (Kumari *et al.*, 2019). Experts predict that this trend will continue in the future due to climate change in the geographic distribution of phytoplasmas and the increased international trade of host plants for planting (Al Ruheili *et al.*, 2021; Aidoo *et al.*, 2021; EFSA Panel on Plant Health (PLH), 2020). Moreover, efficient management of phytoplasmas can improve and sustainably increase agricultural yields (Bertaccini, 2021), thereby contributing to global food security. This review sheds light on phytoplasmas as plant pathogens, the threats they pose, factors contributing to their spread, global collaborative research efforts, management strategies and surveillance and detection. We present case studies that highlight management practices, lessons learned, and future research directions for phytoplasma-mediated diseases.

Understanding phytoplasma plant pathogens

Phytoplasmas, previously identified as mycoplasma-like organisms (MLOs) are one of the smallest known pathogens infecting several plant species worldwide. They are prokaryotic plant pathogenic bacteria, with size ranging from 200 to 1000 nm (McCoy *et al.*, 1989). The phytoplasma's cell lacks a wall and their outer covering is made up of a triple layered single unit membrane (Lee and Davis, 1992). They have genomic sizes ranging from 0.53 to 1.2 kb (Bai *et al.*, 2006; Oshima *et al.*, 2004), with a low G+C content (Kollar and Seemuller, 1989), being descended from an ancestral Gram-positive bacteria in the Bacillus – Clostridium group (Zhao *et al.*, 2009). They live and multiply in the functional phloem sieve tube elements of their hosts, producing disease symptoms such as virescence, phyllody and witches' broom (Bertaccini *et al.*, 2022; Lee *et al.*, 2000). Until their first isolation in pure culture, they were previously thought to be obligate parasites. Moreover, there is a gradual improvement in the methods for applicability to a wider range of '*Ca.* Phytoplasma' species (Contaldo *et al.*, 2012; Contaldo *et al.*, 2016).

The inability to obtain axenic culture of phytoplasmas in the past made their identification using cultural, morphological and biochemical methods challenging (Makarova *et al.*, 2012). Consequently, their identification has relied primarily on molecular methods (Bertaccini and Duduk, 2009) particularly using the 16S ribosomal RNA (16S rRNA) gene. This approach has been used to create two parallel classification systems. In one system all strains of the pathogen are placed in the provisional genus '*Ca.* Phytoplasma' and are separated into species based on variations in the nucleotide sequences of the 16S rRNA gene (IRPCM, 2004; Harrison *et al.*, 2011). In the second system, phytoplasmas are classified into groups and subgroups based on RFLP patterns obtained from PCR-amplified products of unknown isolates (Zhao *et al.*, 2009; Lee and Davis, 1992). Other investigators prefer to first sequence the PCR-amplified 16S rRNA-encoding gene products of the phytoplasma, after which the assembled nucleotides are used in a BLAST search to identify the species and phylogenetic analysis used to assign them to groups and sub-groups. In some instances, the gene sequences can be digested *in silico* and computer programs such as the *i*Phyclassifier can be used to delineate the 16Sr group and subgroups of the unknown isolates (Zhao *et al.*, 2009; Wei *et al.*, 2008). Multilocus analysis involving the 16S ribosomal gene, the 16S–23S intergenic spacer region and *secA* and *groEL* genes have been used to identify the species status of some phytoplasma isolated from palms in Florida (Soto *et al.*, 2021).

Phytoplasmas can infect a wide range of plant species (Seemüller *et al.*, 2002; Lee *et al.*, 2000; Olivier *et al.*, 2009). According to Hemmati *et al.* (2021a), more than 164 plant species made up of fruit crops, vegetables, cereal and oilseed crops, trees, ornamental plants and weeds in the Middle East region have been associated with fourteen 16Sr phytoplasma strains. In Mexico, phytoplasmas have been associated with diseases of different species of palm causing wilting of fronds and eventual death of plants (Hernandez *et al.*, 2020).

Transmission of phytoplasmas occurs via vegetative propagation, dodder (*Cuscuta* spp.), and insect vectors (Aryan *et al.*, 2016; Kaminska and Korbin, 1999). There have been studies on possible transmission of phytoplasma through alfafa seeds and the embryo of coconut fruits obtained from infected palms (Kahn *et al.*, 2002; Cordova *et al.*, 2003). Though some studies could not confirm seed transmission of phytoplasmas (Nipah *et al.*, 2007; Cordova *et al.*, 2003), recent studies have confirmed seed transmission of phytoplasmas in coconut (Narvaez *et al.*, 2022). Worldwide, leafhoppers, planthoppers and psyllids are known to transmit phytoplasmas (Weintraub and Beanland, 2006). These insects feed on plant phloem, ingest the pathogen, which then colonizes their guts and salivary glands, multiplies, and is subsequently released into new hosts during feeding (Ammar and Hogenhout, 2006). Demonstrating insect transmission of phytoplasmas has been challenging due to the difficulty of producing pure cultures. However, recent methodological advances have enabled transmission studies linking specific insect species to pathogen spread. For example, *Austroagallia sinuata* collected from infected fields transmitted '*Ca.* Phytoplasma aurantifolia' to *Aerva javanica* in periwinkle cages (Hemmati *et al.*, 2019). Similarly, the pathogen responsible for lethal bronzing disease (LB) of palm was successfully transmitted from infected spear leaves to a sucrose medium by *Haplaxius crudus* (Mou *et al.*, 2022), and *H. crudus* has recently been experimentally confirmed to transmit LYD (Narvaez *et al.*, 2022). As axenic culture methods for phytoplasmas continue to improve, more studies on insect transmissibility are expected in the future.

Threats posed by emerging phytoplasma pathogens

Evidence first appeared in 1967 linking prokaryotes that morphologically resembled mycoplasmas colonising phloem tissue (then termed mycoplasma-like organisms, MLOs) to yellowing-type plant diseases previously assumed to be caused by viruses (Doi

https://doi.org/10.1017/S0021859625100178 Published online by Cambridge University Press

et al., 1967). Today, phytoplasmas are recognised as major plant pathogens associated with diseases that have serious environmental and economic consequences. A wide range of plant species, including ornamentals, timber trees, shade trees, and economically important food, vegetable, and fruit crops, are affected by phytoplasma-associated diseases (Bertaccini *et al.* 2014; Gasparich, 2010). Varying degrees of phytoplasma incidence on vegetables have been documented in 47 nations, spanning five continents (Kumari *et al.*, 2019). Phytoplasmas have been linked to 164 plant hosts, including feed crops, cereals, fruit crops, medicinal plants and shade trees. Fourteen of the 34 identified phytoplasma ribosomal groups have been reported across the Middle East and other regions worldwide (Hemmati *et al.*, 2021b). While specific phytoplasmas may have limited host ranges, phytoplasma-related disorders impact a diverse range of crops worldwide. Examples include pigeon pea witches' broom (16SrIX) in Brazil (Chen *et al.*, 2008) and the citrus huanglongbing disease in China, which is linked to the aster yellows phytoplasmas (16SrI) (Teixeira *et al.*, 2009). Over 300 different plant diseases that have been connected to phytoplasmas have impacted hundreds of plant taxa (Bertaccini and Duduk, 2009). Woody plant diseases such as coconut lethal yellowing, peach X-disease, grapevine yellows (GY) and apple proliferation, are particularly important due to their commercial significance. Notable phytoplasma diseases that have been recorded in Southeast Asia include rice yellow dwarf, peanut witches' broom, Bermuda grass white leaf and sugarcane white leaf and grassy shoot (Win and Jung, 2012). The phytoplasma group 16SrIX, which affects almond crops, is particularly problematic in the Middle East. Since the 1990s, almond production in Lebanon and Iran has been severely impacted by a fatal disease associated with this group. Thousands of almond trees have been lost in Lebanon since the initial outbreak in the south of the country 15 years ago (Abou-Jawdah *et al.*, 2003). Diseases linked to phytoplasma have long been known to cause significant financial harm to a range of domestic and wild plants. The threat posed by phytoplasma diseases is growing on a global scale due to two main causes: severe epidemics in the rest of the world that affect grapevines, citrus, forest trees, oil-seed crops, alfalfa, stone and pome fruits; and emerging diseases in Latin America, Asia, Africa and the Caribbean that primarily affect sugarcane, corn, cassava, coconuts, papaya and vegetables. In both scenarios, these diseases have the potential to expand to new crop species and significantly affect international trade. Although phytoplasmas are highly metabolically dependent on their host plant, they generally do not cause rapid death. However, in exceptionally cold climates, infected plants die, while in tropical climates, asymptomatic plant presence is common and can have serious epidemiological repercussions (Bertaccini, 2008).

Similar to the identification of phytoplasma strains on the American continent, which comprised strains from 12 subgroups within 10 ribosomal groups, strains from ten (10) ribosomal groups and 16 subgroups on the Asian continent have been documented. The phytoplasma group 16SrIII affects 10 types of vegetable crops, whereas the group 16SrI affects 14. Apart from the few incidences of the pathogen on tomatoes in Mexico, the 16SrIII phytoplasma group seems to be limited to countries such as Bolivia, Brazil, Argentina, Chile and Costa Rica. The 16SrI phytoplasmas group of the aster yellows category is the most common in various genera, followed by '*Ca.* Phytoplasma solani' (Stolbur phytoplasma) (16SrXIIA), clover proliferation (16SrVI) and the 16SrII group of peanut witches' broom (Kumari *et al.*, 2019). Analyses of NCBI database records show that phytoplasma groups 16SrI, 16SrII, 16SrIV, 16SrV, 16SrVI, 16SrIX, 16SrXI, 16SrXII, and 16SrXIV are prevalent in India (Ayman *et al.*, 2010; Priya *et al.*, 2010). In both Iran and India, 16SrI and 16SrII phytoplasmas significantly reduce squash yields (Salehi *et al.*, 2015; Rao *et al.*, 2017b).

Phytoplasmas also represent a major limiting factor for several economically important crops in Europe and North America. For instance, in North America and Europe, the aster yellow phytoplasma significantly reduces the value of ornamental plants such as gladiolus, hydrangea, purple coneflower and China aster, as well as vegetable crops such as lettuce, carrots and celery (Bertaccini and Duduk, 2009). Apple proliferation, European stone fruit yellows and pear decline are fruit tree phytoplasma diseases that are economically significant in Europe (Marcone *et al.*, 2023). Interestingly, in some cases, phytoplasmas infection in ornamental plants may provide desirable and valuable traits, such as the free-branching phenotypes in most commercial poinsettia varieties resulting from infection by phytoplasmas (Lee *et al.*, 2021).

Apple proliferation is widespread across Europe, where affected *Malus domestica* Borkh. trees produce undersized, unmarketable apples. Fruit quality is diminished, size is reduced by approximately 50%, and weight losses range from 63% to 74%. Additionally, reduced tree vigour increases susceptibility to powdery mildew. Three subgroups (16SrII-A, -C and -D) are widely distributed over the African continent and infect faba beans, squash, tomatoes, brinjal and chiles (Omar and Foissac, 2012; Alfaro-Fernández *et al.*, 2011, 2012). In Australia, only three (16SrII, 16SrV and 16SrXII) out of the seven phytoplasma groups (16SrI, 16SrII, 16SrIII, 16SrV, 16SrX, 16SrXI and 16SrXII) have been recorded, and these are known to infect vegetable crops. Nevertheless, there are few reports of phytoplasma disease affecting vegetable crops in Australia. The global analysis of phytoplasma and their threat to food security around the globe is presented in Table 1.

Factors contributing to spread of phytoplasma

Biotic and abiotic factors contribute immensely to phytoplasma transmission. The biotic factors transmit phytoplasma in persistent-propagative mechanism through multiplication within the vector after acquisition (Christensen *et al.*, 2005; Weintraub and Beanland, 2006; Jarausch and Weintraub, 2013; Mou *et al.*, 2022). The transmission of phytoplasmas by insect vectors begins with the acquisition of the pathogen from an infected plant via feeding. The pathogen spreads from the vector's gut to its salivary glands and reproduces within the vector. After reproduction, the vector transfers the pathogen to another plant via feeding thereby infecting the plant. Insects, specifically the phloem-feeding insects such as leafhoppers, planthoppers and psyllids are major vectors of phytoplasma spread (Weintraub and Beanland, 2006). For example, the planthopper *Haplaxius crudus* is a confirmed vector of '*Ca.* Phytoplasma aculeata', a phytoplasma responsible for lethal bronzing of palms in Mexico and Florida (Halbert *et al.*, 2014; Narváez *et al.*, 2018; Mou *et al.*, 2020a; Dzido *et al.*, 2020). Recent studies have also demonstrated transovarial transmission, where offspring of infected vectors carry the pathogen. For instance, the progeny of *Matsumuratettix hiroglyphicus* were found to harbour the sugarcane white leaf phytoplasma and could transmit it to healthy plants (Hanboonsong *et al.*, 2002; Weintraub and

https://doi.org/10.1017/S0021859625100178 Published online by Cambridge University Press

Table 1. The global analysis of phytoplasma and their threat to food security

Strains of phytoplasma	Host range	Distribution of phytoplasma (Country)	References
16SrII,16SrVI	Cabbage	China Iran	Salehi *et al.*, 2007; Cai *et al.*, 2016
16SrIV-D	Canary Island Date palms	USA	Harrision *et al.*, 2008
16SrIV-D	Edible date palms	USA	Harrision *et al.*, 2008
16SrIV-D	Wild date palms	USA	Harrision *et al.*, 2008
16SrIV-D	Pygmy date palms	USA	Jeyaprakash *et al.*, 2011
16SrII, 16SrVI, 16SrI, 16SrIX, and 16SrXII	Tomato	Iran	Zibadoost *et al.*, 2016; Salehi *et al.*, 2016
16SrVI, 16SrII	Cucumber	Iran	Zibadoost *et al.*, 2016
16SrII, 16SrVI	Potato	Iran	Zibadoost *et al.*, 2016
16SrII, 16SrVI	Chili	Iran	Zibadoost *et al.* 2016
16SrII, 16SrVI	Lettuce	Iran	Zibadoost *et al.*, 2016
16SrII, 16SrVI	Spinach	Iran	Zibadoost *et al.* 2016
16SrII, 16SrVI	Squash	Iran	Zibadoost *et al.*, 2016
16SrXIV	Bermuda grass	Iran, Turkey, Iraq, Saudi Arabia	Salehi *et al.*, 2009; Çağlar *et al.*, 2013; Omar, 2016; Alkuwaiti *et al.*, 2017
16SrII-D	Squash	Egypt	El-Sisi *et al.*, 2017

Beanland, 2006). In addition, phytoplasma can spread through agricultural and horticultural planting materials such as rootstocks, cuttings and grafting materials, especially in woody plants. Plant shoots and roots, such as basal shoots, stems, rhizomes, tubers, stolons, corms, buds and bulbs, can all vegetatively spread phytoplasma (Caglayan *et al.*, 2019; 2023). Seed transmission has been reported in both herbaceous and woody plant species (Satta *et al.*, 2019). The seeds from phytoplasma-infected alfalfa (*Medicago sativa*), lime (*Citrus aurantiaca*) and tomato (*Lycopersicum esculentum*) from Oman and Italy were found to contain phytoplasmas belonging to ribosomal groups 16SrI, 16SrXII and 16SrII (Khan *et al.*, 2002; Botti and Bertaccini, 2006).

Abiotic factors also facilitate the spread and prevalence of phytoplasmas. The weather influences the life cycle, behaviour and abundance of insect vectors responsible for phytoplasma transmission. Galetto *et al.* (2011) found that two phytoplasma – chrysanthemum yellows (vector: Euscelidius variegatus, host: daisy) and 'Flavescence dorée' (vector: *Scaphoideus titanus*, host: broad bean) multiplied faster in insects when it was cooler and in plants when it was warmer. Similarly, Maggi *et al.* (2014) discovered that the epidemics of chrysanthemum yellows phytoplasma in *Chrysanthemum carinatum* plants were faster at higher temperatures, with a linear increase in spreading rate from 0.2 plants infected per day at 15°C to about 0.7 plants per day at 30° *Ca.* Phytoplasma infections are also influenced by prevailing winds and geographical factors, such as mountain ranges, which determine their spread and direction of lethal yellowing of coconut palm (Mpunami *et al.*, 2000; Mora-Aguillera, 2002).

Surveillance and detection of phytoplasmas

The surveillance of phytoplasmas encompasses various techniques aimed at monitoring their presence and distribution in plant populations. Historically, methods like symptom profiling, microscopy, serology and dodder transmission studies have been employed for disease detection (Nair and Manimekalai, 2021; Gupta *et al.*, 2023). However, recent advances in molecular techniques, such as PCR and loop-mediated isothermal amplification (LAMP), have revolutionized early detection capabilities (Jawhari *et al.*, 2015; Parnell *et al.*, 2017). Additionally, biosensing techniques and innovative surveillance methods like remote sensing and citizen science initiatives have enhanced the ability to monitor phytoplasma populations (Dyussembayev *et al.*, 2021; Parnell *et al.*, 2017). These developments have significantly improved early detection and monitoring strategies. Various surveillance strategies, such as airborne surveillance and habitat monitoring, offer valuable insights into vulnerable plant hosts and habitats, aiding in resource allocation and decision-making for habitat restoration and biosecurity (Mitchell, 2024). Moreover, effective monitoring and surveillance methods, informed by statistical analyses such as geographic information systems (GIS), are critical for timely detection and management of phytoplasma diseases (Mitchell, 2024; Parnell *et al.*, 2017).

The implementation of PCR and nested-PCR assays enables the broad detection of phytoplasma presence, including instances of mixed infection, in field-collected samples (Bertaccini *et al.*, 2014). Utilizing conserved gene sequences has represented a significant breakthrough in detecting, identifying and categorizing phytoplasmas. A barcode system was previously utilized for the detection and identification of phytoplasmas (Bertaccini *et al.*, 2014). Additionally, the introduction of diagnostic tests based on quantitative PCR assays (qPCR) has proven highly sensitive, reducing the risk of amplicon contamination and eliminating the need for gel-based post-PCR product analysis, thus establishing qPCR as a reliable alternative method to nested-PCR assays in routine testing (Pérez-López *et al.*, 2017). PCR assays employing primers sourced from phytoplasma-specific DNA probes or sequences from the 16S rRNA gene have demonstrated superior sensitivity in detecting phytoplasmas within infected plant or insect hosts. Identification and classification have typically involved the use of RFLP analysis of this genetic locus, leading to the delineation of over thirty 16Sr groups designated as 16SrI –

https://doi.org/10.1017/S0021859625100178 Published online by Cambridge University Press

16SrXXXIII (Pérez-López *et al.*, 2017). Using a primer pair (P1/Tint) identified in a portion of the $tRNA^{Ile}$ region within the spacer region resulted in a universal phytoplasma detection involving a secondary PCR product of approximately 200 bp (Smart *et al.*, 1996). Universal PCR primers targeting the 16S rRNA gene have also facilitated the detection of known phytoplasma strains (Gundersen and Lee, 1996; EPPO, 2018). However, alternative genes, such as *groEL* which is also known as *cpn60*, have been utilized as supplementary markers for phytoplasma identification and classification (Pérez-López *et al.*, 2017). The *cpn60* universal target (cpn60 UT), spanning approximately 550 bp, resides within the Cpn60-encoding gene, recognized as a molecular barcode for the Bacteria domain, and serving as a taxonomic marker for characterizing microbial communities (Pérez-López *et al.*, 2017).

Despite significant progress, persistent limitations primarily stem from the intricate nature of phytoplasma infections and their accelerated spread through global plant trade (Pierro *et al.*, 2019). Agricultural practices face substantial consequences due to challenges in early identification (Dyussembayev *et al.*, 2021). Statistical methodologies play a crucial role in guiding surveillance endeavours, aiding in resource allocation and decision-making for habitat restoration and establishment (Mitchell, 2024; Parnell *et al.*, 2017). Nonetheless, the challenge of detecting phytoplasmas in late spring may be attributed to their proliferation time on stems, branches and new foliage (Gupta *et al.*, 2023). Addressing these constraints necessitates ongoing research and innovative detection technologies to mitigate the impact of phytoplasma diseases on agriculture and ecosystems. Despite advancements in surveillance and detection techniques, several challenges persist, particularly in early identification due to inconspicuous symptoms and the erratic distribution of phytoplasmas within infected plants (Dyussembayev *et al.*, 2021; Gupta *et al.*, 2023). Moreover, predicting phytoplasma effectors remains challenging, hindering effectorome comparisons (Carreón-Anguiano *et al.*, 2023). Table 2 shows techniques and principles of phytoplasma detection. The difficulty in detecting phytoplasmas during late spring may be attributed to their proliferation on stems, branches and new leaves (Gupta *et al.*, 2023). When examining a pear tree, symptoms indicative of pear decline might easily be misinterpreted as signs of other problems like graft-incompatibility, chlorosis, virus-related diseases, or even drought (Errea *et al.*, 2002). However, enhancing early detection capabilities necessitates reliable detection methods and a deeper comprehension of phytoplasma biology. Table 3 depicts the challenges and solutions for early detection of phytoplasma.

Case studies on selected phytoplasma-related disease outbreaks

The global impact of phytoplasma diseases on crops and other environmentally significant plants is profoundly concerning (Bertaccini, 2021). These pathogens have caused substantial economic losses and continue to pose significant threats to agricultural sustainability (Pierro *et al.*, 2019). Multiple cases of devastating consequences of these diseases on various categories of crop species including tree crops (e.g. grapevines, apples, coconuts), vegetables (tomatoes, potatoes, carrots), cereals (maize) and legumes have been reported across different regions of the world (Pierro *et al.*, 2019; Siampour *et al.*, 2019). The outbreak of phytoplasma-related diseases highlights the high vulnerability of global cropping sysems and stresses the importance of continuous research in developing resistant varieties and innovative control methods (Hemmati *et al.*, 2021a). Currently, there is a continuous effort to control the impact of phytoplasma diseases in several crop cultivation areas across the globe (Wang *et al.*, 2024). Knowledge about previous outbreaks of the disease, management practices and current status of these diseases is essential in facilitating research and development efforts. A case study of selected phytoplasma diseases is briefly presented below:

Grapevine yellows diseases

The Grapevine yellow (GY) diseases are a group of phytoplasma-associated diseases affecting grapevines worldwide. Twelve '*Ca.* Phytoplasma' species, distributed across 17 16SrRNA subgroups, are reported to be associated with these diseases (Bertaccini *et al.*, 2022). In Europe, the most significant GY diseases are Flavescence dorée and Bois noir (BN), both of which have caused severe damage to viticulture over several decades (Jarausch *et al.*, 2021; Belli *et al.*, 2010). Flavescence dorée is linked to phytoplasmas belonging to the 16SrV-C and 16SrV-D subgroups, while BN disease is associated with phytoplasmas of the 16SrXII-A subgroup (Bertaccini *et al.*, 2007; 2014). GY outbreaks, particularly in Europe, have led to significant economic losses, reduced grape, apple, and peach yields, and quality deterioration, adversely affecting the wine industry (Ember *et al.*, 2018). Severe outbreaks in Italy, France, and Spain have produced symptoms such as leaf rolling, vine decline, shrivel.

Management strategies

Effective management of GY disease involves a comprehensive approach, including cultural practices such as removing infected vines planting disease-resistant cultivars, and implementing rigorous monitoring using molecular diagnostic tools (Oliveira *et al.*, 2019). Chemical control is limited due to the feeding behaviour of insect vectors (primarily leafhoppers), but integrated pest management strategies, incorporating biological controls such as predatory insects, are being explored, though further research is needed for widespread application (Bianco *et al.*, 2019; Belli *et al.*, 2010; Boudon-Padieu, 2003). Early detection and removal of infected vines, combined with strict quarantine measures, remain crucial for preventing further spread (Krüger *et al.*, 2022). Current research focuses on pathogen diversity, vector ecology, and host-pathogen interactions, aiming to develop sustainable management strategies and resilient grape cultivars through coordinated global efforts (Krüger *et al.*, 2022; Zambon *et al.*, 2018; Abu Alloush *et al.*, 2023; Bertaccini, 2021).

Witches' broom disease of lime

Witches' broom disease of lime (WBDL), caused by '*Ca.* Phytoplasma aurantifolia' (16Sr II-B), has severly affected Mexican lime production. The disease originated in Oman and subsequently spread to Brazil, the Middle East, and India (Hemmati *et al.*, 2021a; Al-Yahyai *et al.*, 2015). The outbreak destroyed over 50% of lime cultivation in some areas, leading to significant economic losses. Early symptoms include proliferation of multiple small pale-green leaves, which later dry and drop, leaving dead twigs (Donkersley *et al.*, 2018a).

https://doi.org/10.1017/S0021859625100178 Published online by Cambridge University Press

Table 2. Techniques and principles of phytoplasma detection

Detection techniques	Advantages/Principles	Reference
Symptom profiling, electron microscopy, DAPI (4',6-diamidino-2-phenylindole) staining, serology (monoclonal and polyclonal antisera)	Earlier and simple methods for phytoplasma detection.	Gupta *et al.*, 2023
Loop-mediated isothermal amplification (LAMP)	Early detection, rapid and sensitive identification of phytoplasma DNA. Minimal equipment requirement and amenable to in-field use.	Jawhari *et al.*, 2015; Nair and Manimekalai, 2021; Mitchell, 2024
Quantitative real-time PCR (QPCR)	High-throughput, rapid and sensitive detection of phytoplasmas; low concentration of phytoplasma detection is possible.	Satta *et al.*, 2017
Electrochemical and optical techniques such as the use of pathogen biosensors, anti-body biosensors, and DNA probe.	High sensitivity	Khater *et al.*, 2017; Dyussembayev *et al.*, 2021
Remote sensing	Phytoplasma surveillance	Parnell *et al.*, 2017
Citizen science initiatives	Phytoplasma surveillance	Parnell *et al.*, 2017
Diagnostic networks	Phytoplasma surveillance	Parnell *et al.*, 2017

Management strategies

Control measures for WBDL involve a combination of cultural, biological, and chemical approaches (Donkersley *et al.*, 2018b). Cultural measures include pruning infected branches, removing severely affected trees, and controlling weed hosts. Biological control methods, including the use of natural enemies of the psyllid vector and entomopathogenic fungi, have shown promise (Siampour *et al.*, 2019). However, insecticide-based chemical control is challenging due to the rapid reproduction of psyllids. Complete removal of infected trees in endemic areas is recommended to reduce disease spread (Hemmati *et al.*, 2021a). WBDL management underscores the importance of early detection and rapid response. The persistence of '*Ca.* Phytoplasma aurantifolia' in alternative hosts and wild plant species complicates eradication efforts (Bertaccini, 2023). Despite extensive control strategies, challenges persist due to the pathogen's wide host range, difficulty in controlling psyllid populations, and its capacity to infect multiple citrus species (Golmohammadi *et al.*, 2023). Research on WBDL focuses on understanding pathogen epidemiology, transmission dynamics, and genetic diversity (Bertaccini, 2023). Studies of psyllid vector behaviour also support efforts to breed resistant cultivars and develop biocontrol agents (Mankin and Rohde, 2020). Advances in diagnostic tools and epidemiological modelling are contributing to improved management strategies (Santos *et al.*, 2023). Collaboration among researchers, growers, and regulatory agencies remains essential in combating WBDL.

Lethal yellowing diseases

Lethal yellowing disease (LYD) is a devastating phytoplasma-mediated condition affecting about 35 palm species and characterized by similar symptoms (Gurr *et al.*, 2016). LYD was first observed in the Caribbean Island at the end of the 19th century. In Jamaica and the Americas, the disease is simply referred to as LYD, and as lethal yellowing-like diseases in other parts of the globe. LYD poses major impacts on palm trees, especially coconut trees, thus affecting global coconut production (Oropeza-Salín *et al.*, 2020). LYD and related diseases lead to rapid palm death, causing premature fruit shedding, necrosis, and leaf yellowing, with no known cure (Gurr *et al.*, 2016; Dollet *et al.*, 2009). These outbreaks, spanning over five decades in various countries worldwide, have caused extensive losses, even affecting previously resistant coconut cultivars in some regions, threatening coconut cultivation globally (Gurr *et al.*, 2016; Eziashi and Omamor, 2010). The disease is associated with '*Ca.* Phytoplasma palmae' (16SrIV-A, -B, -D, E and -F) in the Caribbeans and Americas (EFSA PLH Panel, 2017). '*Ca.* Phytoplasma palmicola' (16SrXXII-A, -B) in West Africa and Mozambique (Harrison *et al.*, 2014) and '*Ca.* Phytoplasma cocostanzaniae' (16SrIV-C) in Kenya and Tanzania. In Papua New Guinea, '*Ca.* Phytoplasma noviguineense' (16SrIV) is associated with Bogia Cococnut Syndrome (Miyazaki *et al.*, 2018).

Management strategies

LYD outbreaks have had devastating impacts on coconut palms and other susceptible species (Gurr *et al.*, 2016). Management of LYD involves various approaches, including removal and destruction of infected palms to reduce spread of disease, controlling insect vectors and employing resistant palm species or varieties (Gurr *et al.*, 2016). The persistence of phytoplasmas in alternative hosts and weed reservoirs complicates disease management. Application of antibiotics such as oxytetracycline and cultural practices via trunk injections have been attempted, though with limited success (Soto *et al.*, 2021).

'Candidatus Phytoplasma solani'

The outbreak of '*Ca.* Phytoplasma solani' in Serbia and neighbouring regions posed a severe threat to crops, such as potatoes and tomatoes, causing stunting and leaf deformation (Mitrović *et al.*, 2022; Kosovac *et al.*, 2018). The pathogen's prevalence in Europe results in substantial economic losses, despite control efforts involving quarantine measures, crop rotation and insecticides application (Kosovac *et al.*, 2023; Jakovljević *et al.*, 2020; Quaglino *et al.*, 2013). Challenges persist due to its varied host range, making vector control complex and necessitating separate disease management strategies in affected regions (Kosovac *et al.*, 2023; Pierro *et al.*, 2020).

https://doi.org/10.1017/S0021859625100178 Published online by Cambridge University Press

Table 3. Challenges and solutions for early detection of phytoplasma

Challenges of early detection	Solutions	References
Lack of visible symptoms during initial stages	Molecular techniques such as PCR and LAMP for rapid and sensitive identification	Parnell *et al.*, 2017; Dyussembayev *et al.*, 2021
Low concentration in woody hosts	Proper sampling techniques and reliable nucleic acid extraction methods	Jawhari *et al.*, 2015; EPPO, 2018
Erratic distribution within infected plants	Use of biosensing techniques and advanced tools for plant pathogen detection	Nair and Manimekalai, 2021; Gupta *et al.*, 2023
Diverse range of plant hosts	Universal PCR primers designed for the amplification of the 16S rRNA gene of any known phytoplasma	EPPO, 2018
Phytoplasmas disguising in plant's vascular system	New forms of surveillance, including remote sensing, citizen science initiatives and diagnostic networks	Parnell *et al.*, 2017
Difficulty in predicting effectors	Efforts towards reliable comparison of effectoromes, overcoming prediction challenges	Carreón-Anguiano *et al.*, 2023
Late spring detection challenges	Understanding the duration required for phytoplasma growth on stems, branches and new leaves	Gupta *et al.*, 2023

Management strategies

Managing outbreaks of '*Ca.* Phytoplasma solani' involves integrated pest management practices, including the removal and destruction of infected plants, controlling insect vectors and implementing strict quarantine measures to limit disease spread (Carminati *et al.*, 2021). Cultural practices such as crop rotation, planting disease-resistant cultivars and the use of healthy planting material, contribute to disease reduction (Sémétey *et al.*, 2018). Chemical control targeting insect vectors may be employed, but effectiveness varies due to the diverse nature of the vectors and their habitats (Mitrović *et al.*, 2022). Lessons learned from '*Ca.* Phytoplasma solani' outbreaks stress early detection, rapid response and preventive measures (Nutricati *et al.*, 2023). Understanding vector behaviour aids in targeted control. Quarantine and biosecurity are crucial (Chalam *et al.*, 2023). The current situation regarding '*Ca.* Phytoplasma solani' outbreaks varies across affected regions, with sporadic occurrences reported in some areas while persistent outbreaks continue in others (Mitrović *et al.*, 2022). Despite control efforts, challenges remain due to the complexity of vector-pathogen interactions, multiple potential reservoir hosts and environmental factors influencing disease spread (Kosovac *et al.*, 2023). Research on '*Ca.* Phytoplasma solani' emphasizes pathogen diversity, genetic studies and rapid diagnostics (Nutricati *et al.*, 2023; Çağlar *et al.*, 2021).

Focus areas includes vector biology, biocontrol agents and breeding resistant plants. The outbreak highlights the critical importance of proactive disease management through surveillance, early detection, and eradication. Integrated methods and collaborations have proven crucial for successful control in vineyards (Kosovac *et al.*, 2023; Mitrović *et al.*, 2022). Figure 1 indicates representative phytoplasma diseases, countries of major reported occurrence, management strategies, lessons learnt, and current situations and research efforts. The integration of various control methods, including both cultural practices and chemical treatments, was vital for effective disease management. Furthermore, collaboration between researchers, vineyard owners and governmental agencies was essential to implement and sustain these control measures, successfully.

International cooperation for phytoplasma disease control

The International Plant Protection Convention (IPPC) has outlined measures to curb the global outbreak of plant pests, aiming to mitigate negative impacts on food security, biodiversity, and economic prosperity. Currently, there is no effective curative treatment for LYD; however, thanks to the efforts of Michael Black, outbreaks have been maintained at manageable levels. As a palm grower in Jamaica, Black has significantly reduced the incidence of the disease on his farm by employing an integrated pest management approach. This includes on-farm quarantine measures, rigorous weekly surveillance, the removal and burning of palms exhibiting LYD symptoms, and the replanting of disease-resistant varieties. Also, by managing weeds and applying fertilizers effectively, as shown in the studies by Myrie *et al.* (2011), Serju (2012), and CARDI (2013), farmers can improve the health and yield of palm crops.

A large group of plant pests threaten global food production, forest productivity, biodiversity, and natural flora. Therefore, preventing the spread and establishment of these harmful pests in new countries and regions is the main task of national plant protection organizations (NPPOs) and the IPPC. These NPPOs are reliable entities entrusted with responsibilities to provide and receive government-to-government phytosanitary assurances, and should be resourced to accomplish this core mandate successfully. Myrie *et al.* (2011) and Serju (2012) documented that an extensive comparison of seven selected farms over several years revealed a significant reduction in LYD incidence on four farms that implemented Black's management practices. In contrast, three farms that did not adopt any management strategies continued to suffer severe losses due to the disease. Globally, it is recommended that infected palms be promptly removed and destroyed. Eradication programs should be enhanced with natural barriers to prevent vector movement, along with the application of insecticides and antibiotic treatments via trunk injection using tetracycline products (McCoy *et al.*, 1976; Eziashi and Omamor, 2010; Wang *et al.*, 2024). Figure 2 shows a network diagram showing international research collaboration under EU-funded TROPICSAFE Project (Adapted from Final Handbook with Innovation Factsheets).

Research initiatives and funding

Phytoplasma research projects have received a substantial amount of financial support from funding agencies to study pathogen identification on different crops across the globe. The European Union's Horizon 2020 programme (2014–2020) allocated nearly €80 billion to research and innovation. The TROPICSAFE Project, involving 22 partners from 12 countries, received €4 million of this

https://doi.org/10.1017/S0021859625100178 Published online by Cambridge University Press




Figure 1. Representative phytoplasma diseases, countries of major reported occurrence, management strategies, lessons learnt and current situation and research effort.

funding. The project focuses on identifying insect-borne prokaryote-associated diseases in tropical and subtropical perennial crops (www.tropicsafe.eu). Major diseases under this funding include lethal yellowing in palms ('*Ca.* Phytoplasma' species), yellows in grapevines ('*Ca.* Phytoplasma' species), and Huanglongbing in citrus ('*Ca.* Liberibacter' species). The aims include: (1) generating data on epidemiological cycles, insect vectors, and alternative host plants; (2) developing rapid, reliable detection methods and holistic management approaches; and (3) scaling up demonstration activities and field trials to improve farmers' livelihoods. Researchers from Canada and Côte d'Ivoire are also exploring ways to reduce coconut crop losses from lethal yellowing, which devastates plantations in West Africa. Improved understanding of the disease, plant breeding, and replanting efforts will help preserve the livelihoods of Côte d'Ivoire's coconut farmers. Table 4 shows funding allocation for phytoplasma research. The Canadian International Food Security Research Fund (CIFSRF) has contributed CA$2.57 million to this work.

Management strategies of phytoplasma-mediated diseases

The lack of a cell wall by phytoplasmas makes them difficult to target with traditional bactericides (Hogenhout *et al.*, 2008). Consequently, managing phytoplasma-mediated diseases primarily involves targeting the insect vectors responsible for transmitting these pathogens, rather than directly treating infected plants (Bianco *et al.*, 2019). As there is currently no effective cure, mitigation strategies emphasize preventive cultural practices (Kumari *et al.*, 2019). Cultural practices play a pivotal role in managing phytoplasma diseases and include adopting resistant plant varieties, rogueing (removing and destroying infected plants), ensuring the use of clean propagating material, and controlling insect vectors known to transmit phytoplasmas (Bertaccini, 2021).

Antibiotics and other molecules

Even though antibiotics such as tetracyclines have been used in the field to control phytoplasma diseases in some high-value crops, their extensive use is limited by costs, emergence of resistant microbial strains, potential hazards to humans and concerns with environmental pollution (Tanno *et al.*, 2018). Other molecules such as ribosome-inactivating proteins, plant hormones and resistance inducers have been tested and shown to exhibit some level of efficacy (Bertaccini, 2021). Additionally, essential oils and ribosome-inactivating proteins have demonstrated promising results in eliminating phytoplasmas in micro-propagated infected plant shoots (Bertaccini, 2021). In vitro systems have also been utilized to produce phytoplasma-free germplasm, which can be further propagated in insect-proof conditions before deployment in open fields (Hogenhout, 2009). These diverse strategies offer a multifaceted approach to mitigate the impact of phytoplasma diseases on crops while minimizing reliance on traditional antibiotics.

Resistant plant varieties

Resistant plant varieties play a crucial role in reducing the susceptibility of crops to phytoplasma infections (Gorshkov and Tsers, 2022). Several species have been identified for their resistance

https://doi.org/10.1017/S0021859625100178 Published online by Cambridge University Press




Figure 2. A Network Map showing international research collaboration involved in the TROPICSAFE Project (Adapted from Final Handbook with Innovation Factsheets, www.tropicsafe.eu). A=The Americas, B=Africa, C=Europe.

to specific phytoplasma infections, contributing to crucial insights for disease management and breeding programs (Wei *et al.*, 2021). For example, *S. mulayanum* and *S. alatum* have been identified as being resistant to sesame phyllody, with *S. alatum* specifically noted for possessing a dominant gene linked to phyllody resistance (Yadav *et al.*, 2022). In temperate fruit trees, various *Prunus* species, including European plum, sour and sweet cherry, demonstrated reduced susceptibility to European Stone Fruit Yellows (ESFY) phytoplasma infections, with some exhibiting tolerance and resistance (Cieślińska, 2011). Additionally, several *Prunus* species, such as *P. betulifolia, P. calleryana, P. nivalis, P. elaeagrifolia, P. syriaca, P. pashia* and *P. dimorphophylla,* were found to be resistant to Pear Decline phytoplasma infections (Marcone *et al.*, 2023). These resistant cultivars are essential for long-term disease management and breeding efforts aimed at developing elite plant lines with enhanced resistance (Roy *et al.*, 2023). Natural resistance mechanisms, including non-host resistance, are also valuable in managing phytoplasma diseases (Hogenhout, 2009). While resistant plant varieties play a crucial role in reducing the susceptibility of crops to phytoplasma diseases, their effectiveness may diminish over time. This may can be due to the emergence of new phytoplasma strains or changes in environmental conditions that can overcome these resistance mechanisms.

Use of disease-free materials

The use of disease-free planting material is crucial to minimizing phytoplasma contamination (Kumari *et al.*, 2019). This approach aligns with the principles of integrated pest management, emphasizing sustainable and holistic practices in agriculture (Deguine *et al.*, 2021). Ongoing research and adaptive strategies remain crucial to staying ahead of evolving challenges posed by phytoplasma diseases and their vectors.

Rogueing

Rogueing, the removal and destruction of infected plants, helps prevent the spread of phytoplasmas to healthy plants (Jeger and Gilligan, 2007). The primary objective of rogueing is twofold. Firstly, it aims to suppress the disease by preventing the dissemination of phytoplasmas (Bertaccini, 2021). In doing so, it disrupts the transmission cycle, especially considering the common mode of transmission through insect vectors. Secondly, rogueing serves as a vigilance monitoring tool, enabling the early detection of infected plants (Sisterson and Stenger, 2013). Early identification is crucial for swift action, preventing the disease from establishing and spreading extensively within the crop. However, rogueing is labour-intensive and costly, especially for small-scale farmers (Welbaum, 2017). Training and awareness programmes are therefore essential to ensure accurate symptom recognition.

Control of insect vectors

Phytoplasma diseases are primarily transmitted by insect vectors, particularly leafhoppers, planthoppers and psyllids (Kumari *et al.*,

https://doi.org/10.1017/S0021859625100178 Published online by Cambridge University Press

Table 4. Funding allocation for Phytoplasma research

	Funding agency	Amount	Time span	Location of collaborators	Reference
1.	European Union (TROPICSAFE)	€4 million	2014–2020	Global	www.tropicsafe.eu
2.	CIFSRF-IDRC	CA$2.57 million	2014–2017	West Africa & Canada	idrc-crdi.ca
3.	Citrus Research and Development Foundation	US$124 million	2010–2017	Florida, USA	The National Academies Press. https://doi.org/10.17226/25026
4.	European Union	€52.3 million	2013–2015	Europe	https://www.hrpub.org/download/20160930/UJAR5-10407286.pdf
5.	Kaken	¥7.9 Million	1995–1996	Asia	https://kaken.nii.ac.jp/en/grant/KAKENHI-PROJECT-07456026/
6.	Washington Tree Fruit Research commission	$230,561	2024–2027	North America	https://treefruit.wsu.edu/article/2024-washington-tree-fruit-research-commission-grant-awards-for-cherry

2019). Identification of vector species transmitting specific phytoplasma strains is essential for targeted control strategies. For example, the cixiid planthopper *Hyalesthes obsoletus* is the main vector of the 16SrI-B phytoplasma strain causing sesame phyllody (Kosovac *et al.*, 2018). In the case of ESFY, the psyllid *Cacopsylla pruni* is the primary vector transmitting the 16SrX-B phytoplasma to temperate fruit trees (Riedle-Bauer *et al.*, 2019). The control of insect vectors in the context of phytoplasma diseases requires a multifaceted and integrated approach (Weintraub and Beanland, 2006). By combining insect vector surveillance, biological control, and when necessary, targeted chemical interventions, farmers can effectively manage vector populations and minimize the risk of phytoplasma transmission (Skendžić *et al.*, 2021). Sustainable and environmentally friendly strategies are essential to protect agricultural ecosystems. Chemical vector control should be used cautiously due to potential ecological and environmental impacts (Gurr *et al.*, 2015). Nevertheless, in certain situations, specific classes of pesticides may be considered. These include neonicotinoids, pyrethroids, organophosphates, insect growth regulators (IGRs) and botanical insecticides (Lalah *et al.*, 2022). Important considerations include managing resistance, minimizing harm to non-target organisms, environmental impact, adherence to legal regulations and precise application timing (Damalas and Eleftherohorinos, 2011).

Good Agricultural Practices (GAPs)

Good sanitation practices, such as proper pruning and propagation techniques, are important in preventing the spread of phytoplasma diseases (Bertaccini, 2021). Timely planting is also critical, as susceptibility can vary with planting dates (Marcone *et al.*, 2023). To effectively manage phytoplasma diseases, it is crucial to combine multiple cultural practices, as relying on a single method may prove insufficient. Figure 3 is an infographic showing phytoplasmas disease management approaches.

Future perspectives

Human activities, such as monoculture farming and global trade, facilitate the rapid evolution of phytoplasmas, posing significant threats to global food security. Monitoring genetic diversity, mutation rates and potential host shifts is crucial for understanding and tackling emerging challenges. Expected trends include the development of more virulent strains and changes in host ranges (Venbrux *et al.*, 2023). Research into vector ecology, particularly the role of insect species, is essential to understand the factors influencing pathogen evolution and gene flow.

Technological advancement is vital for enhancing detection and control of phytoplasma infections (Carvajal-Yepes *et al.*, 2022). The collective employment of rapid and reliable diagnostic tools, molecular techniques such as PCR-based assays, advanced imaging technologies and omics Sciences such as genomics and proteomics offer promise for improved and accurate detection (Carvajal-Yepes *et al.*, 2022). Control strategies should explore new antimicrobial agents, resistant crop varieties and environmentally sustainable practices. Collaboration among researchers, agriculturists and industry players is crucial for practical solutions.

Developing resilient agricultural systems is imperative to mitigate the impact of emerging phytoplasma on global food security (Altieri, 1999). Integrated pest management, crop diversification and rotation practices are essential for enhancing crop resilience. Fostering resilient farming communities through education and extension services, empowering farmers with knowledge on disease prevention and sustainable practices; these contribute to building a robust defence against phytoplasma threats.

Recent breakthroughs in machine learning, geospatial analytics and big data mining, present exciting possibilities in battling phytoplasma-mediated diseases (Buja *et al.*, 2021; Kasinathan *et al.*, 2021; Ristaino *et al.*, 2021; Silva *et al.*, 2021; Zhang *et al.*, 2019). AI-driven monitoring systems can provide real-time alerts, contributing to proactive responses. Precision agriculture, guided by AI, optimizes resource use and enhances farm efficiency, providing adaptive management strategies against phytoplasma threats. These technologies also have the capacity to establish and strengthen global surveillance networks for phytoplasma for early detection and timely response (Carvajal-Yepes *et al.*, 2019). International collaboration, standardized surveillance protocols and remote sensing technologies are essential to prevent transboundary spread and ensure coordinated control measures.

In the long term, climate change may influence the spread of plant diseases, including phytoplasma infections (Chaloner *et al.*, 2021; Delgado-Baquerizo, 2020; Dudney *et al.*, 2021). Understanding the potential impact of climate change on the distribution and prevalence of phytoplasmas and their vectors is

https://doi.org/10.1017/S0021859625100178 Published online by Cambridge University Press




Figure 3. An infographic showing Integrated Pest Management Approaches.

critical. Climate modelling combined with epidemiological studies can predict areas at heightened risk, enabling proactive mitigation strategies.

Public–private partnerships are effective for accomplishing extension and outreach objectives in plant pathology (Markell *et al.*, 2020). Collaboration between academia, government, NGOs and agribusinesses expedites the development and implementation of new technologies and control strategies. Public-private collaborations enhance the accessibility and affordability of diagnostic tools, treatments and preventive measures, benefiting farmers globally.

Investing in capacity building and education programs is fundamental for building resilience plant protection (Gervais, 2004). Empowering individuals with knowledge and skills to identify, manage and prevent phytoplasma infections ensures a robust and adaptive agricultural sector. Training initiatives that cover integrated pest management, sustainable agriculture, and responsible technology use ensure a robust and adaptive agricultural sector capable of responding effectively to phytoplasma threats.

Conclusion

This review highlights the rising incidence of phytoplasma infections across diverse crops, ranging from staple grains to high-value cash crops, and their potential impact on food production. The adaptability of phytoplasmas to new environments and hosts, coupled with climate change, underscores the urgency of addressing this emerging agricultural challenge. The immediate and long-term implications for global food security necessitate a collaborative, multi-faceted approach to mitigate phytoplasma threats and safeguard future food supplies.

Data availability statement. All the data used and generated in the study can be found in this manuscript.

Author contributions. OFA: Study-conceived and designed, Writing-original draft, Review & Editing; AKD, JOH: Writing-original draft, Review & Editing; FKA, JO, AM, AFO, KO, FLS, NEY, HL: Writing-original draft; BAO: Formatting, Review & Editing.

Funding statement. Not applicable.

Competing interests. The authors affirm there were no financial or commercial ties that might be viewed as having a potential conflict of interest.

Ethics standards. Not applicable.

References

Abou-Jawdah Y, Dakhil H, El-Mehtar S and Lee IM (2003) Almond witches'-broom phytoplasma: a potential threat to almond, peach, and nectarine. *Canadian Journal of Plant Pathology* **25**, 28–32.

Abu Alloush AH, Bianco PA, Busato E, Alkhawaldeh Y, Alma A, Tedeschi R and Quaglino F (2023) Grapevine yellows in Jordan: associated phytoplasmas, putative insect vectors and reservoir plants. *Plant Pathology* **72**, 1380–1392.

Aidoo OF, Cunze S, Guimapi RA, Arhin L, Ablormeti FK, Tettey E, ... and Yankey N (2021) Lethal yellowing disease: insights from predicting potential distribution under different climate change scenarios. *Journal of Plant Diseases and Protection* **128**, 1313–1325. https://doi.org/10.1007/s41348-021-00488-1

Al Ruheili AM, Boluwade A and Al Subhi AM (2021) Assessing the impact of climate change on the distribution of lime (16SRII-b) and alfalfa (16srii-d) phytoplasma disease using maxent. *Plants* **10**, 460.

Al-Yahyai RA, Al-Sadi AM, Al-Said FA, Al-Kalbani ZH, Carvalho CM, Elliot SL and Bertaccini A (2015) Development and morphological changes in leaves and branches of acid lime (Citrus aurantifolia) affected by witches' broom. *Phytopathologia Mediterranea* **54**, 133–139.

Alfaro-Fernández A, Ali MA, Abdelraheem FM, Saeed EAE and FontSan-Ambrosio MI (2012) Molecular identification of 16SrII-D subgroup phytoplasmas associated with chickpea and faba bean in Sudan. *European Journal of Plant Pathology* **133**, 791–795. https://doi.org/10.1007/s10658-012-9975-7

Alfaro-Fernández A, Cebrián MDC, Villaescusa FJ and Font-San–Ambrosio IM (2011) Detection and identification of aster yellows and stolbur phytoplasmas in various crops in Spain. *Bulletin of Insectology* **64**, S63–S64.

Alkuwaiti NAS, Kareem TA and Sabier LJ (2017) Molecular detection of 'Candidatus Phytoplasma australasia' and 'Ca. P. cynodontis' in Iraq. *Agriculture* **63**, 112–119.

Altieri MA (1999) The ecological role of biodiversity in agroecosystems. In *Invertebrate Biodiversity as Bioindicators of Sustainable Landscapes*. Amsterdam: Elsevier, pp. 19–31.

Ammar E-D and Hogenhout SA (2006) Mollicutes associated with arthropods and plants. In Kostas B. and Miller T. (eds.), *Insect Symbiosis*. Vol. 2. FL, USA: CRC Press, Taylor and Francis Group, Boca Raton, pp. 97–118.

https://doi.org/10.1017/S0021859625100178 Published online by Cambridge University Press

Arocha-Rosete Y, Konan Konan JL, Diallo AH, Allou K and Scott JA (2014) Identification and molecular characterization of the phytoplasma associated with a lethal yellowing-type disease of coconut in Côte d'Ivoire. *Canadian Journal of Plant Pathology* **36**, 141–150.

Aryan P, Musetti R, Riedle-Bauer M and Brader G (2016) Pytoplasma transmission by heterologous grafting influences viability of the scion and results in early symptoms development in periwinkle rootstocks. *Journal of Phytopathology* **164**, 631–640.

Ayman OF, Kumar Y, Hallan V and Zaidi AA (2010) Molecular characterization of the phytoplasmas associated with toon trees and periwinkle in India. *The Journal of General Plant Pathology* **76**, 351–354.

Bai X, Zhang J, Ewing A, Miller SA, Jancso Radek A, Shevchenko DV, Tsukerman K, Walunas T, Lapidus A, Campbell JW and Hogenhout SA (2006) Living with genome instability: the adaptation of phytoplasmas to diverse environments of their insect and plant hosts. *Journal of Bacteriology* **188**, 3682–3696.

Belli G, Bianco PA and Conti M (2010) Grapevine yellows in Italy: past, present and future. *Journal of Plant Pathology* **92**, 303–326.

Bertaccini A (2008) Phytoplasma diseases and climate change. II Taller Internacional de Fitoplasmas, 23 y 25 de Septiembre, Las Habana, Cuba, CD Rom.

Bertaccini A (2021) Containment of phytoplasma-associated plant diseases by antibiotics and other antimicrobial molecules. *Antibiotics* **10**, 1398. https://doi.org/10.3390/antibiotics10111398

Bertaccini A (2023) 'Candidatus Phytoplasma aurantifolia' and 'Candidatus Phytoplasma australasia': epidemiology meets quarantine. *Phytopathogenic Mollicutes* **13**, 77–78. https://doi.org/10.5958/2249-4677.2023.00039.7

Bertaccini A, Arocha-Rosete Y, Contaldo N, Duduk B, Fiore N, Montano HG ... and Zamorano A (2022) Revision of the 'Candidatus Phytoplasma'species description guidelines. *International Journal of Systematic and Evolutionary Microbiology* **72**, 005353. https://doi.org/10.1099/ijsem.0.005353

Bertaccini A, Calari A and Felker P (2007) Developing a method for phytoplasma identification in cactus pear samples from California. *Bulletin of Insectology* **60**, 257.

Bertaccini A and Duduk B (2009) Phytoplasma and phytoplasma diseases: a review of recent research. *Phytopathologia Mediterranea* **48**, 355–378.

Bertaccini A, Duduk B, Paltrinieri S and Contaldo N (2014) Phytoplasmas and phytoplasma diseases: a severe threat to agriculture. *American Journal of Plant Sciences* **5**, 1763–1788. https://doi.org/10.4236/ajps.2014.512191

Bianco PA, Romanazzi G, Mori N, Myrie W and Bertaccini A (2019) Integrated management of phytoplasma diseases. In Bertaccini A, Weintraub P, Rao G and Mori N (eds.), *Phytoplasmas: Plant Pathogenic Bacteria - II*. Singapore: Springer. pp. 193–210.

Bogoutdinov DZ, Valyunas D, Navalinskene M and Samuitene M (2008) About specific identification of phytoplasmas in Solanaceae crops. *Agricultural Biology* **1**, 77–80.

Botti S and Bertaccini A (2006) Phytoplasma infection trough seed transmission: further observations. In: *16th International Congress of the International Organization of Mycoplasmology*, Cambridge, UK, 9–14 July 2006, 113.

Boudon-Padieu E (2003, September) The situation of grapevine yellows and current research directions: distribution, diversity, vectors, diffusion and control. In *Proceedings of XV International Conference of Virus and Virus-like diseases of Grapevine* (pp. 47–53). Locorotondo (Italy).

Brown SE, Been BO and McLaughlin WA (2006) Detection and variability of the lethal yellowing group (16Sr IV) phytoplasmas in the *Cedusa* sp. (Hemiptera: Auchenorrhyncha: Derbidae) in Jamaica. *Annals of Applied Biology* **149**, 53–62.

Buja I, Sabella E, Monteduro AG, Chiriacò MS, De Bellis L, Luvisi A and Maruccio G (2021) Advances in plant disease detection and monitoring: from traditional assays to in-field fiagnostics. *Sensors* **21**, 2129. https://doi.org/10.3390/s21062129

Çağlar BK, Satar S and Elbeaino T (2013) Detection and molecular characterization of bermuda grass (Cynodon dactylon) white leaf phytoplasma from Turkey. *International Journal of Agriculture and Biology* **15**, 90–94.

Çağlar BK, Şimşek E, Dikilitas M and Bertaccini A (2021) Characterization of 'Candidatus Phytoplasma solani'associated with a maize leaf reddening disease in Turkey. *Journal of Phytopathology* **169**, 658–666.

Çağlayan K (2023) Diversity, distribution, and status of phytoplasmas diseases in Turkey. In *Diversity, Distribution, and Current Status*. Cambridge, MA: Academic Press, pp. 249–267.

Caglayan K, Gazel M and Škorić D (2019) Transmission of phytoplasmas by agronomic practices. In Bertaccini A, Weintraub P, Rao G and Mori N (eds.), *Phytoplasmas: Plant Pathogenic Bacteria - II*. Singapore: Springer, pp. 149–163.

Cai H, Wang L, Mu W, Wan Q, Wei W and Davis RE, *et al.* (2016) Multilocus genotyping of a 'Candidatus Phytoplasma aurantifolia'-related strain associated with cauliflower phyllody disease in China. *Annals of Applied Biology* **169**, 64–74.

CARDI (2013) *Coconut Industry Development for the Caribbean: Towards a Shared Vision and Road Map*. Georgetown Guyana: CARDI.

Carminati G, Brusa V, Loschi A, Ermacora P and Martini M (2021) Spatiotemporal and Quantitative Monitoring of the Fate of 'Candidatus Phytoplasma Solani' in Tomato Plants Infected by Grafting. *Pathogens* **10**, 811. https://doi.org/10.3390/pathogens10070811

Carreón-Anguiano KG, Vila-Luna SE, Sáenz-Carbonell L and Canto-Canche B (2023) PhyEffector, the first algorithm that identifies classical and non-classical effectors in phytoplasmas. *Biomimetics* **8**, 550. https://doi.org/10.3390/biomimetics8070550

Carvajal-Yepes M, Cardwell K, Nelson A, Garrett KA, Giovani B, Saunders DGO, Kamoun S, Legg JP, Verdier V, Lessel J, Neher RA, Day R, Pardey P, Gullino ML, Records AR, Bextine B, Leach JE, Staiger S and Tohme J (2019) A global surveillance system for crop diseases. *Science* **364**, 1237–1239. https://doi.org/10.1126/science.aaw1572

Carvajal-Yepes M, Cuervo M, Kreuze JF, Alakonya A, Kumar P, Onaga G ... and Bui T (2022) *Report of the Global Online Survey to Identify Key Knowledge and Capacity Gaps on Diagnostics and Surveillance of Pests & Diseases in Targeted Countries*. Rome, Italy: CGIAR Plant Health Initiative / International Potato Center (CIP).

Chalam VC, Kumari P, Deepika DD, Yadav P, Kalaiponmani K and Maurya AK (2023) Cross-boundary movement of phytoplasmas in Asia and status of plant quarantine. In *Characterization, Epidemiology, and Management*. Cambridge, MA: Academic Press, pp. 85–96.

Chaloner TM, Gurr SJ and Bebber DP (2021) Plant pathogen infection risk tracks global crop yields under climate change. *Nature Climate Change* **11**, 710–715.

Chen J, Pu X, Deng X, Liu S, Li H and Civerolo E (2008) A phytoplasma closely related to the pigeon pea witches'-broom phytoplasma (16SrIX) is associated with citrus huanglongbing symptoms in the state of São Paulo, Brazil. *Phytopathology* **98**, 977–984.

Christensen NM, Axelsen KB, Nicolaisen M and Schulz A (2005) Phytoplasmas and their interactions with hosts. *Trends in Plant Science* **10**, 526–535.

Cieślińska M (2011) European stone fruit yellows disease and its causal agent 'Candidatus Phytoplasma prunorum'. *Journal of Plant Protection Research* **51**, 441–447. https://doi.org/10.2478/v10045-011-0073-1

Constable FE (2009) Phytoplasma epidemiology: Grapevines as a model. In *Phytoplasmas: Genomes, Plant Hosts And Vectors*. Wallingford, UK: CAB International, pp. 188–212.

Contaldo N, Bertaccini A, Paltrinieri S, Windsor HM and Windsor DG (2012) Axenic culture of plant pathogenic phytoplasmas. *Phytopathologia Mediterranea* **51**, 607–617.

Contaldo N, Satta E, Zambon Y, Paltrinieri S and Bertaccini A (2016) Development and evaluation of different complex media for phytoplasma isolation and growth. *Journal of Microbiological Methods* **127**, 105–110. https://doi.org/10.1016/j.mimet.2016.05.031

Cordova I, Jones P, Harrison NA and Oropeza C (2003) In situ PCR detection of phytoplasma DNA in embryos from coconut palms with lethal yellowing disease. *Molecular Plant Pathology* **4**, 99–108.

Damalas CA and Eleftherohorinos IG (2011) Pesticide exposure, safety issues, and risk assessment indicators. *International Journal of Environmental Research and Public Health* **8**, 1402–1419. https://doi.org/10.3390/ijerph8051402

https://doi.org/10.1017/S0021859625100178 Published online by Cambridge University Press

Deguine J-P, Aubertot J-N, Flor RJ, Lescourret F, Wyckhuys KAG and Ratnadass A (2021). Integrated pest management: Good intentions, hard realities. *Agronomy for Sustainable Development* **41**, 66. https://doi.org/10.1007/s13593-021-00689-w

Delgado-Baquerizo M (2020) The proportion of soil-borne pathogens increases with warming at the global scale. *Nature Climate Change* **10**, 550–554.

Doi Y, Teranaka M, Yora K and Asuyama H (1967) Mycoplasma or PLT grouplike microrganisms found in the phloem elements of plants infected with mulberry dwarf, Potato Witches' Broom, Aster Yellows or Pawlownia Witches' Broom. *Japanese Journal of Phytopathology* **33**, 259–266.

Dollet M, Quaicoe R and Pilet F (2009). Review of coconut "Lethal Yellowing" type diseases diversity, variability and diagnosis. *Oléagineux, Corps Gras, Lipides* **16**, 97–101. https://doi.org/10.1051/ocl.2009.0246

Donkersley P, Blanford JM, Queiroz RB, Silva FWS, Carvalho CM, Al-Sadi AM and Elliot SL (2018b) Witches' Broom Disease of Lime (Candidatus Phytoplasma aurantifolia): Identifying High-Risk Areas by climatic mapping. *Journal of Economic Entomology* **111**, 2553–2561. https://doi.org/10.1093/jee/toy248

Donkersley P, Silva FW, Carvalho CM, Al-Sadi AM and Elliot SL (2018a) Biological, environmental and socioeconomic threats to citrus lime production. *Journal of Plant Diseases and Protection* **125**, 339–356.

Dudney J, Willing CE, Das AJ, Latimer AM, Nesmith JC and Battles JJ (2021) Nonlinear shifts in infectious rust disease due to climate change. *Nature Communications* **12**, 5102.

Dyussembayev K, Sambasivam P, Bar I, Brownlie JC, Shiddiky MJA and Ford R (2021) Biosensor technologies for early detection and quantification of plant pathogens. *Frontiers in Chemistry* **9**, 636245. https://doi.org/10.3389/fchem.2021.636245

Dzido J-L, Sánchez R, Dollet M, Julia J-F, Narvaez M, Fabre S and Oropeza C (2020) Haplaxius crudus (Hemiptera: Cixiidae) transmits the lethal yellowing phytoplasmas, 16SrIV, to Pritchardia pacifica Seem. & H. Wendl (Arecaceae) in Yucatan, Mexico. *Neotropical Entomology* **49**, 795–805. https://doi.org/10.1007/s13744-020-00799-2

Eden-Green S (1997) History, distribution and research on coconut lethal yellowing-like diseases of palms. In *International Workshop on Lethal Yellowing-like Diseases of Coconut. Chatham: Natural Resources Institute*. Chatham, UK: Natural Resources Institute, pp. 9–25.

EFSA Panel on Plant Health (PLH), Jeger M, Bragard C, Candresse T, Chatzivassiliou E,Dehnen-Schmutz K … and Caffier D (2017) Pest categorisation of Palm lethal yellowing phytoplasmas. *EFSA Journal* **15**, e05028. https://doi.org/10.2903/j.efsa.2017.5028

EFSA Panel on Plant Health (PLH), Bragard C, Dehnen-Schmutz K, Gonthier P, Jaques Miret JA, Justesen AF … and Jacques MA (2020) Pest categorisation of the non-EU phytoplasmas of tuber-forming Solanum spp. *EFSA Journal* **18**, e06356.

El-SisiOmar YAF; Sidaros SA; ElSharkawy MM (2017) Characterization of 16SrII-D subgroup associated phytoplasmas in new host plants in Egypt. *Archives of Phytopathology and Plant Protection* **50**, 504–513.

Ember I, Bodor P, Zsófi Z, Pálfi Z, Ladányi M, Pásti G … and Bisztray GD (2018) Bois noir affects the yield and wine quality of Vitis vinifera L. cv.'Chardonnay'. *European Journal of Plant Pathology* **152**, 185–197.

EPPO (2018) PM 7/133 (1) Generic detection of phytoplasmas. *European and Mediterranean Plant Protection Organization* **48**, 414–424. https://doi.org/10.1111/epp.12541

Errea P, Aguelo V and Hormaza JI (2002) Seasonal variations in detection and transmission of pear decline phytoplasma. *Journal of Phytopathology* **150**, 439–443.

Eziashi E and Omamor I (2010) Lethal yellowing disease of the coconut palms (Cocos nucifera L.): An overview of the crises. *African Journal of Biotechnology* **9**, 9122–9127.

Galetto L, Marzachì C, Marques R, Graziano C and Bosco D (2011) Effects of temperature and CO 2 on phytoplasma multiplication pattern in vector and plant. *Bulletin of Insectology* **64**, 201–202.

Gasparich GE (2010) Spiroplasmas and phytoplasmas: microbes associated with plant hosts. *Biologicals* **38**, 193–203.

Gervais S (2004) *Local Capacity Building in Title II Food Security Projects: A Framework and Best Practices*. Washington, DC: United States Agency for International Development (USAID).

Golmohammadi M, Khankahdani HH and Rastegar S (2023) Reaction of some Persian lime accessions on different rootstocks to witches' broom disease of lime. *Phytopathogenic Mollicutes* **13**, 65–66. https://doi.org/10.5958/2249-4677.2023.00033.6

Gorshkov V and Tsers I (2022) Plant susceptible responses: The underestimated side of plant–pathogen interactions. *Biological Reviews of the Cambridge Philosophical Society* **97**, 45–66. https://doi.org/10.1111/brv.12789

Gundersen DE and Lee I-M (1996) Ultrasensitive detection of phytoplasmas by nested-PCR assays using two universal primer pairs. *Phytopathologia Mediterranea* **35**, 144–151. http://www.jstor.org/stable/42685262

Gupta S, Handa A, Brakta A, Negi G, Tiwari RK, Lal MK and Kumar R (2023) First report of 'Candidatus Phytoplasma asteris' associated with yellowing, scorching and decline of almond trees in India. *Peer-jay*, **11**, e15926. https://doi.org/10.7717/peerj.15926

Gurr GM, Bertaccini A, Gopurenko D, Krueger RR, Alhudaib KA, Liu J and Fletcher MJ (2015) Phytoplasmas and their insect vectors: Implications for date palm. In Wakil W, *et al.* (eds.), *Sustainable Pest Management in Date Palm: Current Status and Emerging Challenges*. Cham, Switzerland: Springer International Publishing Switzerland, p. 10.

Gurr GM, Johnson AC, Ash GJ, Wilson BA, Ero MM, Pilotti CA … and You MS (2016) Coconut lethal yellowing diseases: a phytoplasma threat to palms of global economic and social significance. *Frontiers in Plant Science*, **7**, 1521.

Halbert SE, Wilson SW, Bextine B and Youngblood SB (2014) Potential planthopper vectors of palm phytoplasmas in Florida with a description of a new species of the genus Omolicna (Hemiptera: Fulgoroidea). *Florida Entomologist* **97**, 90–97. https://doi.org/10.1653/024.097.0112

Hanboonsong Y, Choosai C, Panyim S and Damak S (2002) Transovarial transmission of sugarcane white leaf phytoplasma in the insect vector Matsumuratettix hiroglyphicus (Matsumura). *Insect Molecular Biology* **11**, 97–103. https://doi.org/10.1046/j.0962-1075.2001.00314.x

Harrison NA, Davis RE, Oropeza C, Helmick EE, Narvaez M, Eden-Green S… and Dickinson M (2014) 'Candidatus Phytoplasma palmicola', associated with a lethal yellowing-type disease of coconut (Cocos nucifera L.) in Mozambique. *International Journal of Systematic and Evolutionary Microbiology* **64**, 1890–1899. https://doi.org/10.1099/ijs.0.060053-0

Harrison NA, Gundersen-Rindal D and Davis RE (2011) Genus I. "Candidatus Phytoplasma" gen.nov. IRPCM Phytoplasma/Spiroplasma working Team, 2004, 1244. In Krieg NR, Staley JT, Brown DR, Hedlund BP, Paster BJ, Ward NL, Ludwig W and Whitman WB (eds.), *Bergey's Manual of Systematic Bacteriology*. Second edition, Volume 4. New York, NY: Springer, pp. 696–719.

Harrison NA, Helmick EE and Elliott ML (2008) Lethal yellowing-type diseases of palms associated with phytoplasmas newly identified in Florida, USA. *Annals of Applied Biology* **153**, 85–94. https://doi.org/10.1111/j.1744-7348.2008.00240.x

Hemmati C, Nikooei M and Al-Sadi AM (2021a) "Candidatus Phytoplasma aurantifolia" increased the fitness of Hishimonus phycitis; the vector of lime witches' broom disease. *Crop Protection* **142**, 105532.

Hemmati C, Nikooei M, Al-Subhi AM and Al-Sadi AM (2021b) History and current status of phytoplasma diseases in the Middle East. *Biology* **10**, 226.

Hemmati C, Nikooei M and Bertaccini A (2019) Identification and transmission of phytoplasma and their impact on essential oil composition in Aerva javanica. *3biotech* **9**, 299.

Hernandez ER, Gordillo JML, Oropeza Saın C, Ortiz Garcıa CF, Magana Alejandro MA, Sanchez Soto S and Garcıa Estrada Y (2020) Detection and identification of phytoplasmas in the 16SrIV-A, -B, and -Dsubgroups in palms in Tabasco, Mexico. *Plant Disease* **104**, 2606–2612.

Hiruki C and Wang K (2004) Clover proliferation phytoplasma:'Candidatus Phytoplasma trifolii'. *International Journal of Systematic and Evolutionary Microbiology* **54**, 1349–1353. https://doi.org/10.1099/ijs.0.02842-0

Hogenhout S (2009) Plant pathogens, minor (phytoplasmas). In *Encyclopedia of Microbiology* (Third Edition). Oxford, UK: Academic Press, pp. 678–688.

Hogenhout SA, Oshima K, Ammar D, Kakizawa S, Kingdom HN and Namba S (2008). Phytoplasmas: Bacteria that manipulate plants and insects.

https://doi.org/10.1017/S0021859625100178 Published online by Cambridge University Press

Molecular Plant Pathology **9**, 403–423. https://doi.org/10.1111/J.1364-3703.2008.00472.X

Huang YK, Wang XY, Zhang RY, Li J, Li YH, Shan HL and Wang CM (2023) The diversity, distribution, and status of phytoplasma diseases in China. In *Diversity, Distribution, and Current Status*. Cambridge, MA: Academic Press, pp. 121–147.

IRPCM Phytoplasma/Spiroplasma Working Team—Phytoplasma taxonomy group (2004) 'Candidatus Phytoplasma', a taxon for the wall-less, non-helical prokaryotes that colonize plant phloem and insects. *International Journal of Systematic and Evolutionary Microbiology* **54**, 1243–1255. https://doi.org/10.1099/ijs.0.02854-0

Jakovljević M, Jović J, Krstić O, Mitrović M, Marinković S, Toševski I and Cvrković T (2020) Diversity of phytoplasmas identified in the polyphagous leafhopper Euscelis incisus (Cicadellidae, Deltocephalinae) in Serbia: Pathogen inventory, epidemiological significance and vectoring potential. *European Journal of Plant Pathology* **156**, 201–221. https://doi.org/10.1007/s10658-019-01878-w

Jarausch B, Biancu S, Kugler S, Wetzel T, Baumann M, Winterhagen P, … and Maixner M (2021) First report of Flavescence Dorée-related phytoplasma in a productive vineyard in Germany. *Plant Disease* **105**, 3285.

Jarausch B and Weintraub P (2013) Spread of phytoplasmas by insect vectors: an introduction. In Weintraub PG and Jones P (eds.), *New Perspectives in Phytoplasma Disease Management*. Bangkok, Thailand: Food and Agriculture Organization of the United Nations (FAO) and International Society for Phytoplasma Research, pp. 17–22.

Jawhari M, Abrahamian P, Sater AA, Sobh H, Tawidian P and Abou-Jawdah Y (2015) Specific PCR and real-time PCR assays for detection and quantitation of 'Candidatus Phytoplasma phoenicium'. *Molecular and Cellular Probes* **29**, 63–70. https://doi.org/10.1016/j.mcp.2014.12.003

Jeger MJ and Gilligan CA (2007) Disease control and its selection for damaging plant virus strains in vegetatively propagated staple food crops; a theoretical assessment. *Proceedings of the Royal Society B: Biological Sciences* **274**, 11–18. https://doi.org/10.1098/rspb.2006.3715

Jeyaprakash A, Sutton BD, Halbert SE and Schubert TS (2011). First report of a 16SrIV-D phytoplasma associated with Texas Phoenix palm decline on pigmy date palm (Phoenix roebelenii) in Florida. *Plant Disease* **95**, 1475–1475.

Kaminska M and Korbin M (1999) Graft and dodder transmission of phytoplasma affecting lily to experimental hosts. *Acta Physiologiae Plantarum* **21**, 21–26.

Kasinathan T, Singaraju D and Uyyala SR (2021) Insect classification and detection in field crops using modern machine learning techniques. *Information Processing in Agriculture* **8**, 446–457. https://doi.org/10.1016/j.inpa.2020.09.006

Khan AJ, Botti S, Paltrinieri S, Al-Subhi AM and Bertaccini AF (2002) Phytoplasmas in alfalfa seedlings: infected or contaminated seeds? In *Abstracts, 14th International Organization of Mycoplasmology Conference*, p. 148. Vienna, Austria.

Khater M, De La Escosura-Muñiz A and Merkoçi A (2017) Biosensors for plant pathogen detection. *Biosensors and Bioelectronics* **93**, 72–86. https://doi.org/10.1016/j.bios.2016.09.091

Kollar A and Seemuller E (1989) Base composition of the DNA mycoplasma-like organisms associated with various plant diseases. *Phytopathology* **127**, 177–186.

Kosovac A, Ćurčić Ž, Stepanović J, *et al.* (2023) Epidemiological role of novel and already known 'Ca. P. solani' cixiid vectors in rubbery taproot disease of sugar beet in Serbia. *Scientific Reports* **13**, 1433. https://doi.org/10.1038/s41598-023-28562-8

Kosovac A, Johannesen J, Krstić O, Mitrović M, Cvrković T, Toševski I and Jović J (2018) Widespread plant specialization in the polyphagous planthopper Hyalesthes obsoletus (Cixiidae), a major vector of stolbur phytoplasma: evidence of cryptic speciation. *PLOS ONE* **13**, e0196969. https://doi.org/10.1371/journal.pone.0196969

Krüger K, Pietersen G, Pietersen G, Stiller M, Engelbrecht G, Rensburg RJ and Bertaccini A (2022) Aster yellows phytoplasma in grapevine in South Africa: insect vector management and alternative host plants. *Phytopathogenic Mollicutes* **12**, 62–62. https://doi.org/10.5958/2249-4677.2022.00027.5

Kumari S, Nagendran K, Rai AB, Singh B, Rao GP and Bertaccini A (2019) Global status of phytoplasma diseases in vegetable crops. *Frontiers in Microbiology* **10**, 1349. https://doi.org/10.3389/fmicb.2019.01349

Lalah JO, Otieno PO, Odira Z and Ogunah JA (2022) *Pesticides: Chemistry, Manufacturing, Regulation, Usage and Impacts on Population in Kenya*. London, UK: IntechOpen.

Lee I, Davis RE and Gundersen-Rindal DE (2000) Phytopathogenic mollicutes. *Annual Review of Microbiology* **54**, 221–255.

Lee IM and Davis RE (1992) Mycoplasma which infect plant and insects. In Maniloff J, McElhaney RN, Finch LR and Baseman JB (eds.), *Mycoplasmas: Molecular Biology and Pathogenesis*. Washington, DC: American Society for Microbiology, pp. 379–390.

Lee IM, Gundersen-Rindal DE, Davis RE and Bartoszyk IM (1998) Revised classification scheme of phytoplasmas based on RFLP analyses of 16S rRNA and ribosomal protein gene sequences. *International Journal of Systematic and Evolutionary Microbiology* **48**, 1153–1169. https://doi.org/10.1099/00207713-48-4-1153

Lee S, Chu C-Y and Chu C-C (2021) Variability of phytoplasma infection density in poinsettia and evaluation of its association with the level of branching in host plants. *Plant Disease* **105**, 1539–1545. https://doi.org/10.1094/PDIS-10-20-2304-RE

Maejima K, Oshima K and Namba S (2014) Exploring the phytoplasmas, plant pathogenic bacteria. *Journal of General Plant Pathology* **80**, 210–221.

Maggi F, Galetto L, Marzachì C and Bosco D (2014) Temperature-dependent transmission of candidatus phytoplasma asteris by the vector leafhopper macrosteles quadripunctulatus kirschbaum. *Entomologia* **2**, 202. https://doi.org/10.4081/entomologia.2014.202

Makarova O, Contaldo N, Paltrinieri S, Kawube G, Bertaccini A and Nicolaisen M (2012) DNA barcoding for identification of 'Candidatus Phytoplasmas' using a fragment of the elongation factor Tu Gene. *PLoS ONE* **7**, e52092. https://doi.org/10.1371/journal.pone.0052092

Mankin RW and Rohde B (2020) Mating behavior of the Asian citrus psyllid. In *Asian Citrus Psyllid: Biology, Ecology and Management of the Huanglongbing Vector*. Wallingford, UK: CABI, pp. 30–42.

Marcone C, Valiunas D, Salehi M, Mondal S and Sundararaj R (2023) Phytoplasma diseases of trees. In *Forest Microbiology*. Oxford, UK: Elsevier, pp. 99–120.

Markell SG, Tylka GL, Anderson EJ and van Esse HP (2020) Developing public–private partnerships in Plant Pathology extension: case studies and opportunities in the United States. *Annual Review of Phytopathology* **58**, 161–180.

McCoy R, Carroll V, Poucher C and Gwin G (1976) Field control of lethal yellowing with oxytetracycline hydrochloride. *Phytopathology* **66**, 1148–1150. https://doi.org/10.1094/Phyto-66-1148

McCoy RE, Caudwell A, Chang CJ, Chen TA and Chiykowski LN, *et al.* (1989) Plant diseases associated with mycoplasma-like organisms. In Whitcomb RF and Tulley JG (eds.), *The Mycoplasmas. Volume v. Spiroplasmas, Acholeplasmas and Mycoplasmas of Plants and Arthropods*. New York: Academic Press, pp. 545–640.

Mitchell RJ (2024) A host-based approach for the prioritisation of surveillance of plant pests and pathogens in wild flora and natural habitats in the UK. *Biological Invasions* **26**, 151–168. https://doi.org/10.1007/s10530-023-03233-x

Mitrović M, Marinković S, Cvrković T, Jović J, Krstić O and Jakovljević M (2022) Framework for risk assessment of 'Candidatus Phytoplasma solani'associated diseases outbreaks in agroecosystems in Serbia. *Journal of Plant Pathology* **104**, 537–552.

Miyazaki A, Shigaki T, Koinuma H, Iwabuchi N, Rauka GB, Kembu A and Namba S (2018) 'Candidatus Phytoplasma noviguineense', a novel taxon associated with Bogia coconut syndrome and banana wilt disease on the island of New Guinea. *International Journal of Systematic and Evolutionary Microbiology* **68**, 170–175. https://doi.org/10.1099/ijsem.0.002480

Mora-Aguillera G (2002) "Dispersal potential of lethal yellowing of the coconut palm," in Proceedings of the Expert Consultation on Sustainable Coconut Production through Control of Lethal Yellowing Disease, CFC Technical Paper No. 18. (Kingston: Common Fund for Commodities), 128–130.

Mou D, Di-Lella B, Herbert SE, Bextine B, Helnick EE and Bahder BW (2022) Acquisition and transmission of the lethal bronzing phytoplasma by

https://doi.org/10.1017/S0021859625100178 Published online by Cambridge University Press

Haplaxius crudus using infected palm spear leaves and srtificial feeding media. *Phytopathology* **112**, 2407–2413.

Mou D-F, Humphries AR, Soto N, Helmick EE, Ascunce MS, Goss EM and Bahder BW (2020a) A survey of auchenorrhynchan insects for identification of potential vectors of the 16SrIV-D phytoplasma in Florida. *Florida Entomologist* **103**, 344–352. https://doi.org/10.1653/024.103.0306

Mpunami A, Tymon A, Jones P and Dickinson MJ (2000) Identification of potential vectors of the coconut lethal disease phytoplasma. *Plant Pathology* **49**, 355–361.

Mugini J (2002) Current status of coconut lethal disease research in Tanzania. In Eden-Green S and Ofori F (eds.), *Proceedings of the Expert Consultation on Sustainable Coconut Production through Control of Lethal Yellowing Disease*. Amsterdam, Netherlands: Common Fund for Commodities, pp. 134–142.

Myrie W, Oropeza C, Saenz L, Harrison N, Roca MM, Córdova I . . . and Douglas L (2011) Reliable improved molecular detection of coconut lethal yellowing phytoplasma and reduction of associated disease through field management strategies. *Bulletin of Insectology* **64**, S203–4.

Nair S and Manimekalai R (2021) Phytoplasma diseases of plants: Molecular diagnostics and way forward. *World Journal of Microbiology and Biotechnology* **37**, 102. https://doi.org/10.1007/s11274-021-03061-y

Narváez M, Nic-Matos G and Oropeza C (2022) In vitro transmission of 16SrIV phytoplasmas to coconut plants by Haplaxius crudus in Yucatan, Mexico. *3 Biotech* **12**, 5.

Narváez M, Vázquez-Euán R, Harrison NA, Nic-Matos G, Julia JF, Dzido JL, Fabre S, Dollet M and Oropeza C (2018) Presence of 16SrIV phytoplasmas of subgroups A, D and E in planthopper Haplaxius crudus Van Duzee insects in Yucatán, Mexico. *3 Biotech* **8**, 61. https://doi.org/10.1007/s13205-018-1094-5

Nipah JO, Jones P and Dickinson MJ (2007) Detection of lethal yellowing phytoplasma in embryos from coconut palms infected with Cape St Paul wilt disease in Ghana.. *Plant Pathology* **56**, 777–784. https://doi.org/10.1111/j.1365-3059.2007.01623.x

Nutricati E, De Pascali M, Negro C, Bianco PA, Quaglino F, Passera A . . . and Luvisi A (2023) Signaling cross-talk between salicylic and gentisic acid in the 'Candidatus Phytoplasma Solani'interaction with Sangiovese Vines. *Plants* **12**, 2695.

Oliveira MJRA, Roriz M, Vasconcelos MW, Bertaccini A and Carvalho SMP (2019) Conventional and novel approaches for managing "flavescence dorée" in grapevine: knowledge gaps and future prospects. *Plant Pathology* **68**, 3–17.

Olivier CY, Lowery DT and Stobbs LW (2009) Phytoplasma diseases and their relationships with insect and plant hosts in Canadian horticultural and field crops. *Canadian Entomologist* **141**, 425–462. https://doi.org/10.4039/n08-CPA02

Omar AF (2016) Association of 'Candidatus phytoplasma cynodontis' with Bermuda grass white leaf disease and its new hosts in Qassim province, Saudi Arabia. *Journal of Plant Interactions* **11**, 101–107.

Omar AF and Foissac X (2012) Occurrence and incidence of phytoplasmas of the 16SrII-D subgroup on solanaceous and cucurbit crops in Egypt. *European Journal of Plant Pathology* **133**, 353–360. https://doi.org/10.1007/s10658-011-9908-x

Oropeza-Salín C, Sáenz L, Narvaez M, Nic-Matos G, Córdova I, Myrie W, Ortíz CF and Ramos E (2020) Dealing with lethal yellowing and related diseases in coconut. In Oropeza C, Verdeil JL, Ashburner GR, Cardeña R and Santamaría JM (eds.), *Coconut Biotechnology: Towards the Sustainability of the 'Tree of Life*. Cham, Switzerland: Springer, pp. 169–197.

Oshima K, Kakizawa S, Nishigawa H Jung HY, Wei W, Suzuki S, Arashida R, Nakata D, Miyata S, Ugaki M and Namba S (2004) Reductive evolution suggested from the complete genome sequence of a plant-pathogenic phytoplasma. *Nature Genetics* **36**, 27–29. https://doi.org/10.1038/ng1277

Parnell S, van den Bosch F, Gottwald T and Gilligan CA (2017) Surveillance to inform control of emerging plant diseases: an epidemiological perspective. *Annual Review of Phytopathology* **55**, 591–610. https://doi.org/10.1146/annurev-phyto-080516-035334.

Pérez-López E, Rodríguez-Martínez D, Olivier CY, Luna-Rodríguez M and Dumonceaux TJ (2017) Molecular diagnostic assays based on cpn60 UT sequences reveal the geographic distribution of subgroup 16SrXIII-(A/I) I phytoplasma in Mexico. *Scientific Reports* **7**, 950. https://doi.org/10.1038/s41598-017-00895-1

Pierro R, Panattoni A, Passera A, Materazzi A, Luvisi A, Loni A, Ginanni M, Lucchi A, Bianco PA and Quaglino F (2020) Proposal of a new bois noir epidemiological pattern related to 'Candidatus Phytoplasma Solani' strains characterized by a possible moderate virulence in Tuscany. *Pathogens (Basel, Switzerland)* **9**, 268. https://doi.org/10.3390/pathogens9040268

Pierro R, Semeraro T, Luvisi A, Garg H, Vergine M, De Bellis L and Gill HK (2019) The distribution of phytoplasmas in south and east Asia: an emerging threat to grapevine cultivation. *Frontiers in Plant Science* **10**, 467526. https://doi.org/10.3389/fpls.2019.01108

Priya M, Chaturvedi Y, Rao GP and Raj SK (2010) First report of phytoplasma Candidatus Phytoplasma trifolii (16SrVI) group associated with leaf yellows of Calotropis gigantean in India. *New Disease Reports* **22**, 29. https://doi.org/10.5197/j.2044-0588.2010.022.029

Quaglino F, Zhao Y, Casati P, Bulgari D, Bianco PA, Wei W and Davis RE (2013) 'Candidatus Phytoplasma solani', a novel taxon associated with stolbur- and bois noir-related diseases of plants. *International Journal of Systematic and Evolutionary Microbiology* **63**, 2879–2894.

Rao GP, Gopala GS and Rao A (2017b) First report of a 'Candidatus Phytoplasma asteris'-related strain (16SrI-B subgroup) associated with witches'broom disease in Cucurbita pepo in India. *New Disease Reports* **35**, 33. https://doi.org/10.5197/j.2044-0588.2017.035.033

Rao GP and Kumar M (2017) World status of phytoplasma diseases associate with eggplant. *Crop Protection* **96**, 22–29. https://doi.org/10.1016/j.cropro.2017.01.005

Riedle-Bauer M, Paleskić C, Schwanzer J, Kölber M, Bachinger K, Schönhuber C, Elek R, Stradinger J, Emberger M, Engel C, Makay M, Zajcsek F and Brader G (2019) Epidemiological and molecular study on '*Candidatus* Phytoplasma prunorum' in Austria and Hungary. *Annals of Applied Biology* **175**, 400–414. https://doi.org/10.1111/aab.12541

Ristaino JB, Anderson PK, Bebber DP, Brauman KA, Cunniffe NJ, Fedoroff NV, Finegold C, Garrett KA, Gilligan CA, Jones CM, Martin MD, MacDonald GK, Neenan P, Records A, Schmale DG, Tateosian L and Wei Q (2021) The persistent threat of emerging plant disease pandemics to global food security. *Proceedings of the National Academy of Sciences* **118**, e2022239118. https://doi.org/10.1073/pnas.2022239118

Roca De Doyle MM (2001) Crisis research: managing lethal yellowing disease. *Biotechnology Development Monitoring* **44**, 12–16.

Roy A, Sahu PK, Das C, Bhattacharyya S, Raina A and Mondal S (2023) Conventional and new-breeding technologies for improving disease resistance in lentil (Lens culinaris Medik). *Frontiers in Plant Science* **13**, 1001682. https://doi.org/10.3389/fpls.2022.1001682

Salehi M, Esmailzadeh Hosseini SA and Salehi E (2016) First report of a 'Candidatus phytoplasma asteris' related phytoplasma associated with eucalyptus little leaf disease in Iran. *Journal of Plant Pathology* **98**, 175.

Salehi M, Izadpanah K, Nejat N and Siampour M (2007) Partial characterization of phytoplasmas associated with lettuce and wild lettuce phyllodies in Iran. *Plant Pathology* **56**, 669–676. https://doi.org/10.1111/j.1365-3059.2007.01616.x

Salehi M, Izadpanah K, Siampour M and Taghizadeh M (2009) Molecular characterization and transmission of bermuda grass white leaf phytoplasma in Iran. *Journal of Plant Pathology* **91**, 655–661.

https://doi.org/10.1017/S0021859625100178 Published online by Cambridge University Press

Salehi M, Siampour M, Alireza S, Hosseini E and Bertaccini A (2015) Characterization and vector identification of phytoplasmas associated with cucumber and squash phyllody in Iran. *Bulletin of Insectology* **68**, 311–319.

Santos AS, Mora-Ocampo IY, de Novais DPS, Aguiar ERGR and Pirovani CP (2023) State of the art of the molecular biology of the interaction between cocoa and witches' broom disease: a systematic review. *International Journal of Molecular Sciences* **24**, 5684. https://doi.org/10.3390/ijms24065684

Satta E, Nanni IM, Contaldo N, Collina M, Poveda JB, Ramírez AS and Bertaccini A (2017) General phytoplasma detection by a q-PCR method using mycoplasma primers. *Molecular and Cellular Probes* **35**, 1–7. https://doi.org/10.1016/j.mcp.2017.05.008

Satta E, Paltrinieri S and Bertaccini A (2019) Phytoplasma transmission by seed. In Bertaccini A, Weintraub P, Rao G and Mori N (eds.), *Phytoplasmas: Plant Pathogenic Bacteria - II*. Singapore: Springer, pp. 131–147.

Savary S, Willocquet L, Pethybridge SJ, Esker P, McRoberts N and Nelson A (2019) The global burden of pathogens and pests on major food crops. *Nature Ecology & Evolution* **3**, 430–439.

Seemüller E, Garnier M and Schneider B (2002) Mycoplasmas of plants and insects. In Razin S and Herrmann R (eds.), *Molecular Biology and Pathogenicity of Mycoplasmas*. Dordrecht, Netherlands: Kluwer Academic/Plenum Publishers, pp. 91–116.

Sémétey O, Gaudin J, Danet JL, Salar P, Theil S, Fontaine M … **and Foissac X** (2018) Lavender decline in France is associated with chronic infection by lavender-specific strains of "Candidatus Phytoplasma solani". *Applied and Environmental Microbiology* **84**, e01507–18.

Serju C (2012) 30 years dedicated to coconut cultivation. In *Jamaica Gleaner*. Kingston: The Gleaner Company.

Siampour M, Izadpanah K, Salehi M and Afsharifar A (2019) Occurrence and distribution of phytoplasma diseases in Iran. In Olivier C, Dumonceaux T and Pérez-López E (eds.), *Sustainable Management of Phytoplasma Diseases in Crops Grown in the Tropical Belt. Sustainability in Plant and Crop Protection*. vol 12. Cham: Springer.

Silva G, Tomlinson J, Onkokesung N, Sommer S, Mrisho L, Legg J, Adams IP, GutierrezVazquez Y, Howard TP, Laverick A, Hossain O, Wei Q, Gold KM and Boonham N (2021) Plant pest surveillance: From satellites to molecules. *Emerging Topics in Life Sciences* **5**, 275–287. https://doi.org/10.1042/etls20200300

Sisterson MS and Stenger DC (2013) Roguing with replacement in perennial crops: Conditions for successful disease management. *Phytopathology* **103**, 117–128. https://doi.org/10.1094/PHYTO-05-12-0101-R

Skendžić S, Zovko M, Živković IP, Lešić V and Lemić D (2021) The impact of climate change on agricultural insect pests. *Insects* **12**, 440. https://doi.org/10.3390/insects12050440

Smart CD, Schneider B, Blomquist CL, Guerra LJ, Harrison NA, Ahrens UKH Lorenz KH, Seemuller E and Kirkpatrick BC (1996) Phytoplasma-specific PCR primers based on sequences of the 16S-23S rRNA spacer region. *Applied and Environmental Microbiology* **62**, 2988–2993.

Soto N, Helmick EE, Harrison NA and Bahder BW (2021) Genetic variability of palm lethal decline phytoplasmas in the Caribbean basin and Florida, USa, based on a multilocus analysis. *Phytopathology* **111**, 2203–2212. https://doi.org/10.1094/PHYTO-04-21-0130-R

Strauss E (2009) Phytoplasma research begins to bloom. *Science* **325**, 388–390.

Tanno K, Maejima K, Miyazaki A, Koinuma H, Iwabuchi N, Kitazawa Y, Nijo T, Hashimoto M, Yamaji Y and Namba S (2018) Comprehensive screening of antimicrobials to control phytoplasma diseases using an in vitro plant-phytoplasma co-culture system. *Microbiology (Reading)* **164**, 1048–1058. https://doi.org/10.1099/mic.0.000681

Teixeira DC, Wulff NA, Martins EC, Kitajima EW, Bassanezi R, Ayres AJ, Eveillard S, Saillard C and Bové JM (2009) A phytoplasma related to 'Candidatus Phytoplasma asteri' detected in citrus showing huanglongbing (yellow shoot disease) symptoms in Guangdong, P. R. China. *Phytopathology* **99**, 236–242.

United Nations Department of Economic and Social Affairs (2021) *World Population Prospects 2017*. New York, USA: United Nations.

Van Dijk M, Morley T, Rau ML and Saghai Y (2021) A meta-analysis of projected global food demand and population at risk of hunger for the period 2010–2050. *Nature Food* **2**, 494–501.

Venbrux M, Crauwels S and Rediers H (2023) Current and emerging trends in techniques for plant pathogen detection. *Frontiers in Plant Science* **14**, 1120968.

Wang R, Bai B, Li D, Wang J, Huang W, Wu Y and Zhao L (2024) Phytoplasma: a plant pathogen that cannot be ignored in agricultural production—Research progress and outlook. *Molecular Plant Pathology* **25**, e13437. https://doi.org/10.1111/mpp.13437

Wei W, Davis RE, Jomantiene R and Zhao Y (2008) Ancient, recurrent phage attacks and recombination shaped dynamic sequence-variable mosaics at the root of phytoplasma genome evolution. *Proceedings of the National Academy of Sciences* **105**, 11827–11832. https://doi.org/10.1073/pnas.0805237105

Wei W, Trivellone V, Dietrich CH, Zhao YDK and Ivanauskas A (2021) Identification of phytoplasmas representing multiple new genetic lineages from phloem-feeding leafhoppers highlights the diversity of phytoplasmas and their potential vectors. *Pathogens* **10**, 352. https://doi.org/10.3390/pathogens10030352

Wei W and Zhao Y (2022) Phytoplasma taxonomy: nomenclature, classification, and identification. *Biology* **11**, 1119. https://doi.org/10.3390/biology11081119

Weintraub PG and Beanland L (2006) Insect vectors of phytoplasmas. *Annual Review of Entomology* **51**, 91–111. https://doi.org/10.1146/annurev.ento.51.110104.151039

Welbaum G (2017) Seed production. In Thomas B, Murray BG and Murphy DJ (eds.), *Encyclopedia of Applied Plant Sciences*. Oxford, UK: Academic Press (Elsevier), pp. 546–552.

Win NKK and Jung HY (2012) The distribution of phytoplasmas in Myanmar. *Journal of Phytopathology* **160**, 139–145.

Yadav R, Kalia S, Rangan P, Pradheep K, Rao GP, Kaur V, Pandey R, Rai V, Vasimalla CC, Langyan S, Sharma S, Thangavel B, Rana VS, Vishwakarma H, Shah A, Saxena A, Kumar A, Singh K and Siddique KH (2022) Current research trends and prospects for yield and quality improvement in sesame, an important oilseed crop. *Frontiers in Plant Science* **13**, 863521. https://doi.org/10.3389/fpls.2022.863521

Zambon Y, Canel A, Bertaccini A and Contaldo N (2018) Molecular diversity of phytoplasmas associated with grapevine yellows disease in North-Eastern Italy. *Phytopathology* **108**, 206–214. https://doi.org/10.1094/phyto-07-17-0253-r

Zhang J, Huang Y, Pu R, Gonzalez-Moreno P, Yuan L, Wu K and Huang W (2019) Monitoring plant diseases and pests through remote sensing technology: a review. *Computers and Electronics in Agriculture* **165**, 104943. https://doi.org/10.1016/j.compag.2019.104943

Zhao Y, Wei W, Lee IM, Shao J, Suo X and Davis RE (2009) Construction of an interactive online phytoplasma classification tool, iPhyClassifier, and its application in analysis of the peach X-disease phytoplasma group (16SrIII). *International Journal of Systematic and Evolutionary Microbiology* **59**, 2582–2593. https://doi.org/10.1099/ijs.0.010249-0

Zibadoost S, Rastgou M and Tazehkand SA (2016) Detection and molecular identification of 'Candidatus phytoplasma trifoli'infecting some cultivated crops and vegetables in West Azarbaijan province, Iran. *Australasian Plant Disease Notes* **11**, 3. https://doi.org/10.1007/s13314-015-0188-8

https://doi.org/10.1017/S0021859625100178 Published online by Cambridge University Press