Ergod. Th. & Dynam. Sys. (2011), **31**, 449–457 doi:10.1017/S0143385710000064
© *Cambridge University Press, 2010*

An example of a non-uniquely ergodic lamination

ÁLVARO LOZANO-ROJO

Universidad del País Vasco, Facultad de CC, Económicas y Empresariales, Dpto. de Economía Aplicada III, Av. Lehendakari Aguirre 83, 48015 Bilbao, Spain (e-mail: alvaro.lozano@ehu.es)

(*Received* 15 *May* 2009 *and accepted in revised form* 13 *January* 2010)

Abstract. This paper presents an example of Riemann surface lamination with at least two ergodic invariant measures. The generic leaves for those measures are of different growth and have different numbers of ends.

1. *Introduction*

A *foliation* is a decomposition of a compact manifold into submanifolds of a given dimension such that it looks locally like a product. The study of foliations leads naturally to the theory of minimal saturated sets which are examples of compact laminations. A *lamination* is a decomposition of a locally compact Hausdorff space into manifolds of a given dimension with the same triviality condition as foliations.

One way to study laminations is to pay attention to the behavior of 'most' of their leaves, that is, to properties shared by a *generic* set of leaves. There are two kinds of genericity in this context: (1) *topological*, that is, a subset of a compact set is generic if it is a countable intersection of open dense sets (a residual set); (2) *measurable*, that is, a subset of a probability space is generic if it is a Borel set of full measure.

Ghys [**8**] showed that generic leaves for ergodic harmonic measure have the same number of ends: 0, 1, 2 or a Cantor set of ends. Later [**4**], Cantwell and Conlon proved the analogous result in the topological setting.

Hence, a natural question arises: does there exist a minimal lamination which exhibits different generic properties depending on the invariant measure considered? In his PhD thesis [**3**], Blanc constructs such a *mixed* minimal lamination. It is a subspace of the Gromov–Hausdorff graph foliated space of the subtrees of the Cayley graph of the free group with three generators, a huge space with lots of leaves of exponential growth. In fact, Blanc's example contains a generic set of leaves with this growth.

Exponential growth is very permissive in terms of the kinds of phenomena it allows. So, is it possible to construct a similar example dropping exponential growth, allowing only polynomial growth? The aim of this paper is to construct such an example of mixed lamination in this more restrictive setting. In fact, the growth allowed will be sub-quadratic.

https://doi.org/10.1017/S0143385710000064 Published online by Cambridge University Press

THEOREM 1.1. *There exists a minimal Riemann surface lamination with two transverse invariant measures ν and ν' such that:*

(1) *the generic leaves for ν are the two-ended linear ones;*

(2) *the generic leaves for ν' are the one-ended ones with growth $f(x) = x^{\ln 5/\ln 3}$.*

Moreover, the set of one-ended leaves is residual.

Whether such spaces can be realized as a minimal set in a foliation is also discussed. This will lead directly to proof of the following result.

COROLLARY 1.1. *There is a foliation containing a minimal set like that of Theorem 1.1.*

The paper is organized as follows. Section 2 presents the construction of the *Gromov–Hausdorff foliated space*. Section 3 is devoted to the construction of two trees on the Gromov–Hausdorff foliated space. The minimal lamination generated by these trees proves Theorem 3.1. In §4 the transverse dynamics of the Gromov–Hausdorff foliated space is reduced to the one given by a group action on the Cantor set. This fact allows one to prove Theorem 4.1.

The results in this paper are part of the author's PhD thesis [**12**] defended at Universidad del País Vasco/Euskal Herriko Unibertsitatea on 6 June 2008.

2. *The Gromov–Hausdorff foliated space*

2.1. *The Gromov–Hausdorff foliated space.* Let $\mathcal{Z}^2$ be the Cayley graph of $\mathbb{Z}^2$ with respect to the usual generating system $S = \{(\pm 1, 0), (0, \pm 1)\}$. Denote by $\mathcal{T}$ the set of infinite subtrees of $\mathcal{Z}^2$ containing the origin as a vertex. Write $B_T(x, r)$ for the open ball in the graph T centered at x with radius r with respect to the usual distance on a graph: the distance between two vertices is the least number of edges one has to go through to get from one vertex to another, and the edges (without their extremes) are isometric to the open unit interval $(0, 1)$. The *Gromov–Hausdorff distance* between two trees T and $T' \in \mathcal{T}$ is given by $d(T, T') = e^{-R(T,T')}$, where $R(T, T') = \sup\{N \geq 1 \mid B_T(0, N) = B_{T'}(0, N)\}$ or $R(T, T') = 0$ if the supremum is not defined. With this distance, $\mathcal{T}$ becomes a Cantor set (see [**7, 12**] for details).

The action of $\mathbb{Z}^2$ over $\mathcal{Z}^2$ induces a pseudogroup of transformations over $\mathcal{T}$: each $v \in \mathbb{Z}^2$ defines a partial transformation $\tau_v : T \mapsto T - v$ defined on the open set of all trees having v as vertex. All these maps generate a pseudogroup of transformations Γ over $\mathcal{T}$. As $\mathbb{Z}^2$ is generated by S, each element $\gamma \in \Gamma$ can be locally written as $\gamma = \tau_{s_1} \circ \cdots \circ \tau_{s_k}$, with $s_1, \ldots, s_k \in S$. Suppose that τ_s is defined over the clopen set $D_s = \{T \in \mathcal{T} \mid \text{the edge } [0, s] \subset T\}$; then the local writing holds, so Γ is finitely generated by $\{\tau_s\}$. The choice of a generating set defines a graph structure on $\mathcal{T}$ in the following way: there is a edge (labeled by $s \in S$) from $T \in D_s$ to $\tau_s(T)$.

As the maps τ_s generate Γ, the connected components of the graph structure on $\mathcal{T}$ are the orbits $\Gamma[T] = \{T - v \mid v \in T\}$. Each $T' \in \Gamma[T]$ can be thought as 'T rooted at v instead of 0'. With this idea of *change of root* in mind, it is obvious that the geometric realization $\overline{\Gamma}[T]$ of $\Gamma[T]$ can be identified with the quotient graph $T/\mathrm{Iso}(T)$, where $\mathrm{Iso}(T) \subset \mathbb{Z}^2$ denotes the *isotropy group of T*.

https://doi.org/10.1017/S0143385710000064 Published online by Cambridge University Press

Recall that the *degree of a vertex* v in a graph is the number of edges incident with it. In this case, each tree T is a vertex in the graph $\Gamma[T]$, so it makes sense to consider its degree, for which one has $\deg(T) = \deg_{\Gamma[T]}(T) = \deg_T(0)$. Hence the degree is constant on the ball $B_{\mathcal{T}}(T, e^{-r}) = \{T' \in \mathcal{T} \mid B_T(0, r) = B_{T'}(0, r)\}$, so $\deg : \mathcal{T} \to \mathbb{N}$ is a continuous map. Ghys [**7**] implicitly used the continuity of this map to construct a Riemann surface lamination which has $\mathcal{T}$ as complete transversal and Γ as holonomy pseudogroup. This construction can be divided into two steps. First, construct the *geometric realization* of all the graphs $\Gamma[T]$ at once, obtaining a graph foliated space (Theorem 2.1). Then replace the local graph structure by a thickened version of the local graph structure (Theorem 2.2).

THEOREM 2.1. (Geometric realization theorem [**12, 13**]) *There exists a compact transversely zero-dimensional graph foliated space* $(\mathcal{G}_{\mathcal{T}}, \mathcal{F})$ *such that (a)* $\mathcal{T}$ *is the complete closed transversal of the vertices, (b) the holonomy pseudogroup of* $\mathcal{F}$ *on* $\mathcal{T}$ *is* Γ*, and (c) the leaf* L_T *through* T *is the graph* $\overline{\Gamma}[T]$*. The graph foliated space* $(\mathcal{G}_{\mathcal{T}}, \mathcal{F})$ *is called the* Gromov–Hausdorff *foliated space.*

THEOREM 2.2. (Thickening theorem [**7, 12, 13**]) *There exists a compact Riemann surface lamination* $(M_{\mathcal{T}}, \mathcal{L})$ *of bounded geometry (see* [**10**]*) such that* $\mathcal{T}$ *is a closed complete transversal of* $\mathcal{L}$ *and the holonomy pseudogroup restricted to* $\mathcal{T}$ *is* Γ*. Additionally, the leaves of* $\mathcal{L}$ *are quasi-isometric (in a uniform way) to the corresponding leaves of* $\mathcal{F}$*.*

See [**12, 13**] for a complete proof of these results in a more general setting.

2.2. *Minimal sets in the Gromov–Hausdorff foliated space.* Roughly speaking, a graph is *repetitive* when one cannot know what point one is at, taking only into account a finite neighborhood of that point. This property is based on the analogous properties of repetitiveness [**2**] or local isomorphism [**15**] for tilings. The notion for graphs appears implicitly in Blanc's PhD thesis [**3**].

Given two trees $T, T' \in \mathcal{T}$, the ball $B_T(x, r)$ is said to *embed faithfully* into T', written as $B_T(x, r) \hookrightarrow T'$, if $B_T(x, r) + v = B_{T'}(x + v, r)$ for some $v \in \mathbb{Z}^2$. If additionally $B_{T'}(x + v, r) \subset B_{T'}(x', r')$, $B_T(x, r)$ is said to *embed faithfully* into $B_{T'}(x', r')$, written as $B_T(x, r) \hookrightarrow B_{T'}(x', r')$.

Definition 2.1. A tree $T \in \mathcal{T}$ is *repetitive* if for all $r > 0$, there exists $R(r) > 0$ such that $B_T(0, r) \hookrightarrow B_T(x, R)$ for any $x \in T$.

Remark 2.1. There is a *uniform* notion of repetitiveness. In that case, one requires the existence of $R > 0$, for a given $r > 0$, such that any ball of radius r can be embedded faithfully into any ball of radius R. This notion is equivalent to repetitiveness (see [**1, 12**]).

The following theorem gives a very convenient criterion for minimality in the Gromov–Hausdorff space in terms of repetitive trees.

PROPOSITION 2.1. [**1, 3, 12**] *A subset of* $\mathcal{T}$ *is minimal if and only if it is the closure of the orbit of a repetitive tree.*

COROLLARY 2.1. *A subspace of* $(\mathcal{G}_{\mathcal{T}}, \mathcal{F})$ *or* $(M_{\mathcal{T}}, \mathcal{L})$ *is minimal if and only if it is the closure of the leaf through a repetitive tree.*

https://doi.org/10.1017/S0143385710000064 Published online by Cambridge University Press




FIGURE 1. The tree G_1.




FIGURE 2. Constructing G_2 with G_1 and P_1.

3. *A non-uniquely ergodic example*

This section is devoted to the construction of two aperiodic subtrees of $\mathcal{Z}^2$. These trees will be created using the same finite patterns, so they will be locally indistinguishable, but their global properties will be different.

As they share the same building blocks, they belong to the same minimal set. On the other hand, the different properties of the trees will translate into two distinct measures with different generic leaves. In fact, those measures will be constructed by means of the averaging sequences used on the construction.

3.1. *Construction of the trees.* The construction is done inductively. Let G_1 be the cross-shaped tree of width two contained in a square of side $l_1 = 3$, like the one in Figure 1. Let T_1 be the infinite horizontal straight line through the origin. Let P_1 be the ball in T_1 centered at the origin of radius $(l_1 - 1)/2 = 1$.

The tree G_2 is constructed mixing G_1 and P_2: put four copies of G_1 around G_1 itself and glue all their edges. In this way a bigger cross-shaped tree is obtained. On both sides of each vertical arm add a copy of P_1, joined with an edge. The resulting tree is G_2, which is contained in a square of side $l_2 = 3l_1 = 3^2$. The process is shown in Figure 2.

Let us define T_2. Choose an even integer r_1 such that

$$\frac{\#G_1}{r_1} \leq \frac{1}{l_1}.$$

Now insert copies of G_1 periodically on intervals of length r_1 along T_1, in the way shown in Figure 3. The resulting tree is T_2. As before, P_2 denotes the ball $\overline{B}_{T_2}(\mathbf{0}, (l_2 - 1)/2)$.

Continue in this way: construct G_3 from P_2 and G_2, putting five copies of G_2 to make a cross, and then gluing four copies of P_2 on the vertical arms, as in the case of G_2 (see Figure 4). Obviously G_3 is contained in a square of side $l_3 = 3l_2 = 3^3$. To construct T_3

https://doi.org/10.1017/S0143385710000064 Published online by Cambridge University Press




FIGURE 3. Constructing the infinite tree T_2, adding copies of G_1.




FIGURE 4. The tree G_3.




FIGURE 5. Detail in the construction of T_3. A copy of G_2 replaces one of G_1.

choose r_2, a multiple of r_1, such that

$$\frac{\#G_2}{r_2} \leq \frac{1}{l_2}.$$

Add copies of G_2 to T_2, but now with periodicity r_2. Some copies of G_2 will replace copies of G_1, as in Figure 5. Notice that in this step (and in the next steps) there is no need to add more edges to connect the copies of G_2 to T_2.

Recursively one gets two increasing sequences of finite trees $\{P_n\}$ and $\{G_n\}$, converging to $P_\infty = \bigcup_{n\geq 1} P_n = \bigcup_{n\geq 1} T_n$ and $G_\infty = \bigcup_{n\geq 1} G_n$ respectively. As these trees share all of their finite patches, it is straightforward to prove the following lemma.

https://doi.org/10.1017/S0143385710000064 Published online by Cambridge University Press

LEMMA 3.1. *Both P_∞ and G_∞ are repetitive and $\mathcal{X} = \overline{\Gamma[P_\infty]} = \overline{\Gamma[G_\infty]}$.*

3.2. *Inequalities derived from the construction.* The construction implies some inequalities for the cardinals of P_n and G_n. In a horizontal straight segment of length l_{n-1} there exist at most $(l_{n-1} - r_1)/r_1 \leq l_{n-1}/r_1$ copies of G_1 within P_∞. For G_2 there are at most $(l_{n-1} - r_2)/r_2 \leq l_{n-1}/r_2$ copies, and so on. So

$$\begin{aligned} l_{n-1} + 1 \leq \#P_n &\leq l_{n-1} + 1 + \sum_{i=1}^{\infty} \frac{l_{n-1}}{r_i} \#G_i \\ &\leq l_{n-1} + 1 + l_{n-1} \sum_{i=1}^{\infty} \frac{1}{3^i} \leq \frac{3}{2} l_{n-1} + 1. \end{aligned} \tag{1}$$

On the other hand, it is clear that

$$\#G_n = 5\#G_{n-1} + 4\#P_{n-1} = \cdots = 5^{n-1}\#G_1 + 4 \sum_{i=0}^{n-2} 5^i \#P_{n-i}. \tag{2}$$

Using (1) in (2), it follows that

$$5^n \leq \#G_n \leq 5^n \left(1 + 6 \sum_{i=0}^{n-2} \frac{3^{n-i-1} + 1}{5^{n-i}} \right) \leq 16 \cdot 5^{n-1}. \tag{3}$$

Inequality (1) shows that P_∞ has linear growth. From (3) it is also obvious that the growth of G_∞ is like that of $f(x) = x^{\ln 5/\ln 3}$.

3.3. *From the trees to a non-uniquely ergodic lamination.* As Lemma 3.1 says, the trees P_∞ and G_∞ are repetitive and belong to the same minimal set $\mathcal{X} \subset \mathcal{T}$. Obviously one gets the same result for the geometric realization and thickening of $\mathcal{X}$. Hence, a minimal graph foliated space $(\mathcal{G}_\mathcal{X}, \mathcal{F})$ and a minimal Riemann surface lamination $(M_\mathcal{X}, \mathcal{L})$ are obtained, which are the closure of the leaf through P_∞ or G_∞ in $(\mathcal{G}_\mathcal{T}, \mathcal{F})$ and $(M_\mathcal{T}, \mathcal{L})$ respectively. By definition, the spaces of transverse invariant measures of those spaces agree with the space $\mathfrak{M}(\Gamma)$ of Γ-invariant measures over X.

PROPOSITION 3.1. *The space $\mathfrak{M}(\Gamma)$ contains at least two invariant measures.*

Proof. It is obvious that $\mathfrak{G} = \{G_n\}_{n\geq 1}$ and $\mathfrak{P} = \{P_n\}_{n\geq 1}$ are two Følner sequences. Therefore, they define two Γ-invariant measures $\mu_\mathfrak{G}$ and $\mu_\mathfrak{P}$ (see [**9**]).

Let us show that there exists a Borel set where these two measures differ. The sets $\mathcal{X}_{G_n} = \{T \in \mathcal{X} \mid G_n - p \subset T \text{ for some } p \in G_n\}$ and $\mathcal{X}_{G_n,0} = \{T \in \mathcal{X} \mid G_n \subseteq T\} \subseteq \mathcal{X}_{G_n}$ are clopen sets. Define $A_n = \bigcap_{i\leq n} \mathcal{X}_{G_i}$. These sets form a decreasing family of clopen sets of non-empty intersection: $G_\infty \in \bigcap_n A_n$. As

$$A_n = \{T \in \mathcal{X} \mid G_n - p \subseteq T \text{ with } p \text{ in the copies of } G_1 \text{ within } G_n\},$$

it is clear that $\mu(A_n) = 5^n \mu(\mathcal{X}_{G_n,0})$ for any $\mu \in \mathfrak{M}(\Gamma)$.

https://doi.org/10.1017/S0143385710000064 Published online by Cambridge University Press

CLAIM. $\mu_{\mathfrak{P}}(\bigcap_n A_n) = 0$. Within P_p there exist at most l_{p-1}/r_n copies of G_n. There are also at most l_{p-1}/r_{n+1} copies of G_{n+1} which has five copies of G_n, and so on. Therefore

$$\mu_{\mathfrak{P}}(\mathcal{X}_{G_n,0}) = \lim_{p\to\infty} \frac{\#(\mathcal{X}_{G_n,0} \cap P_p)}{\#P_p} \leq \lim_{p\to\infty} \frac{l_{p-1}\sum_{i\geq n} 5^{i-n}/r_i}{l_{p-1}+1} \leq \sum_{i\geq n} \frac{5^{i-n}}{r_i}.$$

Using this, inequality (1), the property about the measure of A_n and the relations between r_i, $\#G_n$ and l_i, one has that $\mu_{\mathfrak{P}}(A_n) \leq \sum_{i\geq n} 1/l_i$. Hence $\mu_{\mathfrak{P}}(\bigcap_n A_n) = 0$.

CLAIM. $\mu_{\mathfrak{G}}(\bigcap_n A_n) > 0$. If $p \geq n$, the number of copies of G_n in G_p is at least 5^{p-n}. From (3),

$$\mu_{\mathfrak{G}}(\mathcal{X}_{G_n,0}) = \lim_{p\to\infty} \frac{\#(\mathcal{X}_{G_n,0} \cap G_p)}{\#G_p} \geq \lim_{p\to\infty} \frac{5^{p-n}}{16\cdot 5^{p-1}} = \frac{1}{16\cdot 5^{1-n}}.$$

Therefore $\mu_{\mathfrak{G}}(A_n) \geq 5/16$. So $\mu_{\mathfrak{G}}(\bigcap_n A_n) > 0$, thus $\mu_{\mathfrak{P}}$ and $\mu_{\mathfrak{G}}$ are different. □

3.4. *Generic leaves.* Consider the following tree of $\bigcap_n A_n$:

$$T = \bigcup_n \left(G_n - \sum_{i=1}^{n-1} (l_i, 0) \right).$$

This is a one-ended tree which covers the right half-plane and the horizontal right ray defines its end. The corresponding leaf has one end, therefore the set of all one-ended leaves is residual [**4**]. From the measurability point of view there are various sorts of generic leaves, depending on the measure considered.

THEOREM 3.1. *There exist two transverse invariant measures $\nu_{\mathfrak{G}}$ and $\nu_{\mathfrak{P}}$ on $(\mathcal{X}, \mathcal{L})$ such that:*

(1) *the generic leaves for $\nu_{\mathfrak{G}}$ are the one-ended ones with growth $f(x) = x^{\ln 5/\ln 3}$;*

(2) *the generic leaves for $\nu_{\mathfrak{P}}$ are the two-ended ones with linear growth.*

Proof. (1) Using the notation above, the saturation A of $\bigcap_n A_n$ has positive $\mu_{\mathfrak{G}}$-measure. Obviously, all the leaves $L \subseteq A$ have the same growth as G_∞, namely $f(x) = x^{\ln 5/\ln 3}$. So, there exits an invariant measure $\nu_{\mathfrak{G}}$ whose generic leaves are of growth $f(x) = x^{\ln 5/\ln 3}$. A result of Levitt says that the set of nonlinear two-ended leaves is of null measure [**11**]. Hence, the residual leaves of $\nu_{\mathfrak{G}}$ have one end.

(2) This is just an implication of a general result of Gaboriau [**6**]: *if there is a linear growth leaf, there exists an invariant measure with the two-ended leaves as generic*. In fact, the measure $\nu_{\mathfrak{P}}$ can be constructed from $\mu_{\mathfrak{P}}$ in a similar way to $\nu_{\mathfrak{G}}$ from $\mu_{\mathfrak{G}}$ [**3, 6**]. □

4. *Realization of the example as a minimal set in a foliation*

The aim of this section is to build a surface foliation of codimension two having a minimal set with the same transverse dynamics (i.e. with equivalent holonomy pseudogroup) and coarse geometry as $(\mathcal{G}_{\mathcal{T}}, \mathcal{F})$. Therefore any foliated subspace of $(\mathcal{G}_{\mathcal{T}}, \mathcal{F})$ can be *found* (in the previous sense) within a codimension-two foliation. In fact, the construction can be applied as is to any transversally Cantor compact graph foliated space, because in such a space the vertex transversal is always homeomorphic to the Cantor set.

https://doi.org/10.1017/S0143385710000064 Published online by Cambridge University Press




FIGURE 6. How the graph structure changes from $\mathcal{G}_\mathcal{T}$ to $\mathcal{G}_\mathcal{Y}$.

4.1. *The transverse dynamics of $\mathcal{G}_\mathcal{T}$ are given by a group action.* Consider the graph foliated space $(\mathcal{G}_\mathcal{T}, \mathcal{F})$. Define $\mathcal{Y} = \mathcal{T} \sqcup E$ where E is the set of edges of the graphs in $\mathcal{F}$. Identifying each edge in E with its midpoint, $\mathcal{Y}$ becomes a closed complete transversal of $(\mathcal{G}_\mathcal{T}, \mathcal{F})$.

PROPOSITION 4.1. *The holonomy pseudogroup of $\mathcal{F}$ over $\mathcal{Y}$ is the pseudogroup of a continuous group action of the free group with $2S$ generators $\mathbf{F}_{2S}$.*

Proof. The proof is similar to Feldman and Moore's construction for Borel equivalence relations (see [**5**]). Each translation $\tau_s : D_s \subset \mathcal{T} \to R_s \subset \mathcal{T}$ can be written as $\tau_s = \tau_s^2 \circ \tau_s^1$ where $\tau_s^1 : D_s \subset \mathcal{T} \to E_s \subset E$ and $\tau_s^2 : E_s \subset E \to R_s \subset \mathcal{T}$ are two maps constructed as follows: the image of $T \in D_s$ through τ_s^1 is the edge $(T, s, T - s) \in E$, and the image of the edge $(T, s, T - s) \in E$ is $T - s$, and E_s is $\{(T, s, T - s) \mid T \in D_s\}$. Obviously, τ_s^1 and τ_s^2 are homeomorphisms between clopen disjoint subset of $\mathcal{Y}$. So, for each $s \in S$ and $i = 1, 2$ one can define the homeomorphism $f_s^i : \mathcal{Y} \to \mathcal{Y}$ given by

$$f_s^i(y) = \begin{cases} \tau_s^i(y) & \text{if } y \in D_s, \\ (\tau_s^i)^{-1}(y) & \text{if } y \in E_s, \\ y & \text{in any other case.} \end{cases}$$

These homeomorphisms define an obvious action of the free group $\mathbf{F}_{2S}$ on Y which induces the pseudogroup of holonomy on Y. □

Remark 4.1. The free group $\mathbf{F}_{2S}$ can be replaced by $*_{i=1}^{2S}\mathbb{Z}/2\mathbb{Z}$, because $(f_s^i)^2 = 1_Y$.

The usual graph structure on $\mathbf{F}_{2S}$ induces another on the orbits in the obvious way. Applying the geometric realization theorem, one gets a graph foliated space $(\mathcal{G}_\mathcal{Y}, \mathcal{F})$ whose pseudogroup of holonomy restricted to the complete closed transversal $\mathcal{T}$ is Γ. This foliated space is obtained by substituting each edge of $(\mathcal{G}_\mathcal{T}, \mathcal{F})$ by a flower-shaped graph and adding some cycles corresponding to the action of elements fixing those points (see Figure 6). It is straightforward to prove the following lemma.

LEMMA 4.1. *The inclusion $\iota : \mathcal{T} \to \mathcal{Y}$ is a Lipschitz map, when $\mathcal{T}$ and $\mathcal{Y}$ have the metric induced by the leaves of $\mathcal{G}_\mathcal{T}$ and $\mathcal{G}_\mathcal{Y}$ respectively.*

Applying the thickening theorem to $(\mathcal{G}_\mathcal{Y}, \mathcal{F})$, one obtains a Riemann surface lamination $(M_\mathcal{Y}, \mathcal{L})$ with these properties: it has $\mathcal{T}$ and $\mathcal{Y}$ as a complete closed transversals; the holonomy pseudogroup reduced to $\mathcal{T}$ is Γ; the leaf through a tree $T \in \mathcal{T}$ is quasi-isometric to $\overline{\mathcal{R}}[T]$; and its dynamics reduced to $\mathcal{Y}$ are given by a group action.

https://doi.org/10.1017/S0143385710000064 Published online by Cambridge University Press

4.2. *Realization in a foliation.*

THEOREM 4.1. *There exists a Riemann surface foliation* $(M, \mathcal{F})$ *of class* $C^{\infty,0}$ *and codimension two containing* $(M_{\mathcal{Y}}, \mathcal{L})$ *as a closed saturated set. Therefore, there exists a foliation containing a minimal set with the transverse dynamic and coarse geometric properties of the example of §3.*

Proof. The set $\mathcal{Y}$ is a Cantor set, so the homeomorphisms $f_s^i : \mathcal{Y} \to \mathcal{Y}$ can be extended to homeomorphisms $\widetilde{f}_s^i : \mathbb{S}^2 \to \mathbb{S}^2$ (see [**14**, Ch. 13]). Therefore an action of the free group $\mathbf{F}_{2S}$ on the sphere $\mathbb{S}^2$ appears. Again, the graph structure of $\mathbf{F}_{2S}$ defines a graph structure of constant degree on the orbits given by a generating set. Applying the thickening theorem, a codimension-two Riemann surface foliation of class $C^{\infty,0}$ is obtained, which obviously has $(M_{\mathcal{Y}}, \mathcal{L})$ as a minimal saturated set. □

Acknowledgements. The author was partially supported by UPV 00127.310-E-15916, MEC MTM2004-08214, EHU 06/05, MTM2007-66262 and a grant of the UPV/EHU.

REFERENCES

[1] F. Alcalde Cuesta, Á. Lozano Rojo and M. Macho Stadler. Dynamique transverse de la lamination de Ghys–Kenyon. *Astérisque* **323** (2009), 1–16.

[2] J. Bellissard, R. Benedetti and J.-M. Gambaudo. Spaces of tilings, finite telescopic approximations and gap-labelling. *Comm. Math. Phys.* **261**(1) (2006), 1–41.

[3] E. Blanc. Propriétés génériques des laminations, *Thèse UCB-Lyon* 1, 2001. Available at http://www.umpa.ens-lyon.fr/∼eblanc/.

[4] J. Cantwell and L. Conlon. Generic leaves. *Comment. Math. Helv.* **73** (1998), 306–336.

[5] J. Feldman and C. Moore. Ergodic equivalence relations, cohomology and Von Neumann algebras I. *Trans. Amer. Math. Soc.* **234**(2) (1977), 289–324.

[6] D. Gaboriau. Dynamique des systèmes d'isométries: Sur les bouts des orbites. *Invent. Math.* **126** (1996), 297–318.

[7] É. Ghys. Laminations par surfaces de Riemman. *Panor. Synthèses* **8** (1999), 49–95.

[8] É. Ghys. Topologie des feuilles génériques. *Ann. of Math.* (2) **141** (1995), 387–422.

[9] S. E. Goodman and J. F. Plante. Holonomy and averaging in foliated sets. *J. Differential Geom.* **14** (1979), 401–407.

[10] M. Gromov. Asymptotic invariants for infinite groups. *Geometric Group Theory,* Vol. 2 *(Sussex, 1991) (London Mathematical Society Lecture Note Series, 182)*. Cambridge University Press, Cambridge, 1993.

[11] G. Levitt. On the cost of generating an equivalence relation. *Ergod. Th. & Dynam. Sys.* **15**(6) (1995), 1173–1181.

[12] Á. Lozano Rojo. Laminaciones definidas por grafos repetitivos. *PhD Thesis*, University of Basque Country, 2008.

[13] Á Lozano Rojo. The Cayley foliated space of a graphed pseudogroup. *Proceedings of the XIV Fall Workshop on Geometry and Physics (Publ. de la RSME, 10)*. RSME, Madrid, 2006, pp. 267–272.

[14] E. Moise. *Geometric Topology in Dimensions 2 and 3 (Graduate Texts in Mathematics, 47)*. Springer, Berlin, 1977.

[15] C. Radin and M. Wolff. Space tilings and local isomorphism. *Geom. Dedicata* **42** (1992), 355–360.

https://doi.org/10.1017/S0143385710000064 Published online by Cambridge University Press