Canad. Math. Bull. Vol. 67 (3), 2024, pp. 860–871
http://dx.doi.org/10.4153/S0008439524000250
© The Author(s), 2024. Published by Cambridge University Press on behalf of Canadian Mathematical Society

# Li coefficients and the quadrilateral zeta function

Kajtaz H. Bllaca, Kamel Mazhouda, and Takashi Nakamura

*Abstract.* In this note, we study the Li coefficients $\lambda_{n,a}$ for the quadrilateral zeta function. Furthermore, we give an arithmetic and asymptotic formula for these coefficients. Especially, we show that for any fixed $n \in \mathbb{N}$, there exists $a > 0$ such that $\lambda_{2n-1,a} > 0$ and $\lambda_{2n,a} < 0$.

## 1 Introduction and statement of main results

### 1.1 Li coefficients

The Riemann hypothesis (RH) is a critical question in analytic number theory. As such, it is interesting to examine different ways to frame it, which may shed more light on its resolution. In 1997, Xian-Jin Li has discovered a new positivity criterion for the RH. In [10], he defined the Li coefficients for the Riemann zeta function as

$$\lambda_n = \frac{1}{(n-1)!}\frac{d^n}{ds^n}\left[s^{n-1}\log \xi(s)\right]_{s=1},$$

where $\xi$ is the completed Riemann zeta function defined by

$$\xi(s) = s(s-1)\pi^{-s/2}\Gamma(s/2)\zeta(s),$$

which satisfies $\xi(s) = \xi(1-s)$ and gave a simple equivalence criterion for the RH: RH is true if and only if these coefficients are nonnegative for every positive integer $n$. The Li coefficients $\lambda_n$ can be written as follows:

$$\lambda_n = \sum_{\rho}^{*}\left[1-\left(1-\frac{1}{\rho}\right)^n\right] = \lim_{T\to\infty}\sum_{\rho;|\mathrm{Im}(\rho)|\le T}\left[1-\left(1-\frac{1}{\rho}\right)^n\right],$$

where the sum runs over the nontrivial zeros of the Riemann zeta function counted with multiplicity. This criterion is generalized by Bombieri and Lagarias [4] for any arbitrarily multiset of numbers assuming certain convergence conditions. Voros [19, Section 3.3] has proved that the RH true is equivalent to the growth of $\lambda_n$ as $\frac{1}{2}n\log n$ determined by its archimedean part, while the RH false is equivalent to the oscillations of $\lambda_n$ with exponentially growing amplitude, determined by its finite part. The Li

Received by the editors May 21, 2023; revised March 25, 2024; accepted March 26, 2024.
Published online on Cambridge Core April 8, 2024.
The third author was partially supported by JSPS grant 22K03276.
AMS subject classification: **11M26, 11M35**.
Keywords: Li's coefficients, the quadrilateral zeta function, Riemann Hypothesis.

https://doi.org/10.4153/S0008439524000250 Published online by Cambridge University Press

coefficients were generalized in two ways: by generalizing these coefficients to various sets of functions (the Selberg class, the class of automorphic $L$-functions, zeta function on function fields,...[8, 11, 17]) and by introducing new parameter in its definition (see [12]). The Li coefficients (and its generalizations) have generated a lot of research interest due to its applicability and simplicity.

## 1.2 Quadrilateral zeta function

Recall the definitions of Hurwitz and periodic zeta functions. The Hurwitz zeta function $\zeta(s,a)$ is defined by the series

$$\zeta(s,a) := \sum_{n=0}^{\infty} \frac{1}{(n+a)^s}, \qquad \sigma > 1, \quad 0 < a \le 1.$$

The function $\zeta(s,a)$ is a meromorphic function with a simple pole at $s=1$ whose residue is 1 (see, for example, [1, Section 12]). The periodic zeta function $\mathrm{Li}_s(e^{2\pi ia})$ is defined by

$$\mathrm{Li}_s(e^{2\pi ia}) := \sum_{n=1}^{\infty} \frac{e^{2\pi ina}}{n^s}, \qquad \sigma > 1, \quad 0 < a \le 1$$

(see, for instance, [1, Exercise 12.2]). Note that the function $\mathrm{Li}_s(e^{2\pi ia})$ with $0 < a < 1$ is analytically continuable to the whole complex plane since $\mathrm{Li}_s(e^{2\pi ia})$ does not have any pole, that is shown by the fact that the Dirichlet series of $\mathrm{Li}_s(e^{2\pi ia})$ converges uniformly in each compact subset of the half-plane $\sigma > 0$ when $0 < a < 1$ (see, for example, [9, p. 20]). For $0 < a \le 1/2$, we define zeta functions

$$Z(s,a) := \zeta(s,a) + \zeta(s,1-a), \qquad P(s,a) := \mathrm{Li}_s(e^{2\pi ia}) + \mathrm{Li}_s(e^{2\pi i(1-a)}),$$
$$2Q(s,a) := Z(s,a) + P(s,a), \qquad \xi_Q(s,a) := s(s-1)\pi^{-s/2}\Gamma(s/2)Q(s,a).$$

We can see that $Q(s,a)$ is meromorphic functions with a simple pole at $s=1$. In addition, we have $Q(0,a) = -1/2 = \zeta(0)$ and $\xi_Q(s,a) = \xi_Q(1-s,a)$, which is proved by

$$Q(1-s,a) = \Gamma_{\cos}(s)Q(s,a), \qquad \Gamma_{\cos}(s) := \frac{2\Gamma(s)}{(2\pi)^s}\cos\Big(\frac{\pi s}{2}\Big) \tag{1.1}$$

(see [13, Theorem 1.1]). Moreover, the function $Q(s,a)$ has the following properties. When $a = 1/6, 1/4, 1/3$, and $1/2$, the RH holds true if and only if all nonreal zeros of $Q(s,a)$ are on the line $\mathrm{Re}(s) = 1/2$ (see [14, Proposition 1.3]). Let $N_{\mathrm{Q}}^{\mathrm{CL}}(T)$ be the number of the zeros of $Q(s,a)$ on the line segment from $1/2$ to $1/2 + iT$. In [13, Theorem 1.2], the third author proved that for any $0 < a \le 1/2$, there exist positive constants $A(a)$ and $T_0(a)$ such that

$$N_{\mathrm{Q}}^{\mathrm{CL}}(T) \ge A(a)T \quad \text{whenever} \quad T \ge T_0(a).$$

Next, let $N_F(T)$ count the number of nonreal zeros of a function $F(s)$ having $|\mathrm{Im}(s)| < T$. Then, for any $0 < a \le 1/2$,

$$N_\zeta(T) - N_Q(T) = O_a(T),$$

https://doi.org/10.4153/S0008439524000250 Published online by Cambridge University Press

and the third author [14, Proposition 1.8] proved that

$$N_Q(T) = \frac{T}{\pi}\log T - \frac{T}{\pi}\log(2e\pi a^2) + O_a(\log T).$$

Furthermore, he [14, Theorem 1.1] proved that there is a unique absolute $a_0 \in (0,1/2)$ such that

$$Q(1/2,a) > 0 \iff 0 < a < a_0.$$

In addition, it is proved in [14, Corollary 1.2] that all real zeros of $Q(s,a)$ are simple and are located only at the negative even integers just like $\zeta(s)$ if and only if $a_0 < a \le 1/2$. Let us note by $Z_Q$ the set of all nontrivial zeros $\rho_a$ of $\xi_Q(s,a)$. Since it is an entire function of order 1, one has

$$\xi_Q(s,a) = e^{A+Bs}\prod_{\rho_a\in Z_Q}\left(1-\frac{s}{\rho_a}\right)e^{\frac{s}{\rho_a}} = \xi_Q(0,a)\prod_{\rho_a\in Z_Q}\left(1-\frac{s}{\rho_a}\right), \tag{1.2}$$

where $e^A = 1/2$, $B = \frac{Q'}{Q}(0,a) - 1 - \frac{\gamma+\log\pi}{2}$, and $\gamma$ denotes the Euler constant. Note that $Q'(0,a)$ is given explicitly in [14, Theorem 1.5].

## 1.3 Main results

Recall that $\zeta(1-s) = \Gamma_{\cos}(s)\zeta(s)$ and $Q(1-s,a) = \Gamma_{\cos}(s)Q(s,a)$ by (1.1). However, the function $Q(s,a)$ does not have an Euler product except for $a = 1/6, 1/4, 1/3$, and $1/2$. Hence, the function $Q(s,a)$ is a suitable object to consider the influence of not Riemann's functional equation but an Euler product to zeros of zeta functions. We show a criterion for nonvanishing of $Q(s,a)$ in terms of the positivity of the Li coefficients, an arithmetic and asymptotic formula for these coefficients in Theorems 1.1, 1.2, and 1.4, respectively. It should be emphasized that $\lambda_{n,a}$ defined in (1.3) are the first Li coefficients that we can explicitly give $n \in \mathbb{N}$ such that $\lambda_{n,a} < 0$. There is a possibility that this fact would give an idea to find negative Li coefficients for $\zeta(s)$ if they would exist.

For $n \neq 0$, the Li coefficients attached to $Q(s,a)$ nonvanishing at zero are defined by the sum

$$\lambda_{n,a} := \sum_{\rho_a\in Z_Q}^{*}\left(1-\left(1-\frac{1}{\rho_a}\right)^n\right) = \lim_{T\longrightarrow\infty}\sum_{|\mathrm{Im}(\rho_a)|\le T}^{*}\left(1-\left(1-\frac{1}{\rho_a}\right)^n\right).$$

The symmetry $\rho_a \longmapsto 1-\rho_a$ in the set $Z_Q$ of nontrivial zeros of $Q(s,a)$ implies that $\lambda_{-n,a} = \overline{\lambda_{n,a}} = \lambda_{n,a}$ for all $n \in \mathbb{N}$. So, $\lambda_{n,a}$ are real. We have also

$$\lambda_{n,a} := \frac{1}{(n-1)!}\frac{d^n}{ds^n}\left[s^{n-1}\log\xi_Q(s,a)\right]_{s=1}. \tag{1.3}$$

Moreover, from (1.2), we have (see [4, Equations (2.3) and (2.4)] or [17, Appendix A])

$$\sum_{n=0}^{\infty}\lambda_{n+1,a}s^n = \frac{d}{ds}\log\left[\xi_Q\left(\frac{1}{1-s},a\right)\right].$$

https://doi.org/10.4153/S0008439524000250 Published online by Cambridge University Press

As an analogue of Li coefficients for the Riemann zeta function, we have the following.

***Theorem 1.1*** *The function $Q(s,a)$ does not vanish when $\mathrm{Re}(s) > 1/2$ if and only if $\lambda_{n,a} \geq 0$ for all $n \in \mathbb{N}$.*

An arithmetic formula for $\lambda_{n,a}$ is stated in the following theorems.

***Theorem 1.2*** *We have*

$$\lambda_{n,a} = 1 - \frac{n}{2}(\log(4\pi) + \gamma) + \sum_{k=2}^{n}(-1)^k \binom{n}{k}\left(1 - 2^{-k}\right)\zeta(k) + \sum_{k=1}^{n}\binom{n}{k}\gamma_Q(k-1),$$

*where $\gamma_Q(n)$ are defined as follows:*

$$\frac{Q'}{Q}(s+1,a) + \frac{1}{s} = \sum_{n=0}^{\infty}\gamma_Q(n)s^n.$$

***Theorem 1.3*** *For $a = 1/2, 1/3, 1/4, 1/6$, under the RH, we have*

$$\lambda_{n,a} = \frac{n}{2}\log n + \frac{n}{2}(\gamma - 1 - \log 2\pi) + O(\sqrt{n}\log n).$$

For a fixed $l \in \mathbb{N}$, we have the following asymptotic formula of $\lambda_{l,a}$ when $a \to +0$. We can see that there exists $n \in \mathbb{N}$ such that $\lambda_{n,a} < 0$ by Theorem 1.1 and the fact that $Q(s,a)$ does not satisfy an analogue of the RH when $a \in \mathbb{Q} \cap (0,1/2)\backslash\{1/6, 1/4, 1/3\}$ (see [14, Proposition 1.4]). Clearly, this argument gives no information on the frequency of $n \in \mathbb{N}$, the smallest $n \in \mathbb{N}$ such that $\lambda_{n,a} < 0$ and so on. However, the next theorem implies that $\lambda_{2n,a} < 0$ if we fix any $n \in \mathbb{N}$ and then we take $a > 0$ sufficiently small.

***Theorem 1.4*** *Fix $l \in \mathbb{N}$. Then it holds that*

$$\lambda_{l,a} = \frac{(-1)^{l+1}}{(2a)^l} + O_l\left(a^{1-l}|\log a|\right), \qquad a \to +0.$$

*Especially, for any fixed $n \in \mathbb{N}$, there are $a > 0$ such that*

$$\lambda_{2n-1,a} > 0 \quad \text{and} \quad \lambda_{2n,a} < 0.$$

## 2 Proofs

### 2.1 Proof of Theorem 1.1

Since $\lambda_{-n,a} = \overline{\lambda_{n,a}} = \lambda_{n,a}$ for all $n \in \mathbb{N}$, then $\mathrm{Re}(\lambda_{-n,a}) = \mathrm{Re}(\lambda_{n,a}) = \lambda_{n,a}$. Using that $\xi_Q(s,a)$ is an entire function of order 1, and its zeros lie in the critical strip $0 < \mathrm{Re}(s) < 1$, we obtain that the series $\sum_{\rho \in Z_Q}\frac{1+|\mathrm{Re}(\rho)|}{(1+|\rho|)^2}$ is convergent. Application of [4, Theorem 1] to the multiset $Z_Q$ of zeros of $Q(s,a)$ gives that $\mathrm{Re}(\rho) \leq 1/2$ if and only if $\lambda_{n,a} \geq 0$ for all $n \in \mathbb{N}$. Now, the application of the same theorem to the multiset $1 - Z_Q = Z_Q$ gives $\mathrm{Re}(\rho) \geq 1/2$ if and only if $\lambda_{n,a} \geq 0$. This completes the proof.

https://doi.org/10.4153/S0008439524000250 Published online by Cambridge University Press

Theorem 1.1 can also be proved by the same argument used in [5, Theorem 1], which is due to Oesterlé.

## 2.2 Proof of Theorem 1.2

From the expression of $\xi_Q(s,a)$, one has

$$\frac{\xi_Q'}{\xi_Q}(s,a) = \frac{1}{s} + \frac{1}{s-1} - \frac{1}{2}\log\pi + \frac{1}{2}\frac{\Gamma'}{\Gamma}(s/2) + \frac{Q'}{Q}(s,a),$$

which is rewritten as

$$\frac{\xi_Q'}{\xi_Q}(s+1,a) = \frac{1}{s+1} + \frac{1}{s} - \frac{1}{2}\log\pi + \frac{1}{2}\frac{\Gamma'}{\Gamma}((s+1)/2) + \frac{Q'}{Q}(s+1,a). \tag{2.1}$$

Note that $Q(s,a)$ is a meromorphic function on the whole complex plane, which is holomorphic everywhere except for a simple pole at $s=1$ with residue 1 (see [13, Section 2.1]). Let us define the coefficients $\gamma_Q(n)$ and $\tau_Q(n)$ as follows:

$$\frac{Q'}{Q}(s+1,a) + \frac{1}{s} = \sum_{n=0}^{\infty}\gamma_Q(n)s^n \tag{2.2}$$

and

$$-\frac{1}{2}\log\pi + \frac{1}{2}\frac{\Gamma'}{\Gamma}((s+1)/2) = \sum_{n=0}^{\infty}\tau_Q(n)s^n. \tag{2.3}$$

By Equation (1.2), one has

$$\log\xi_Q(s,a) = \log\xi_Q(0,a) - \sum_{\rho_a\in Z_Q}\sum_{m=1}^{\infty}\frac{1}{m\rho^m}s^m.$$

From the functional equation for the function $\xi_Q(s,a)$, in the neighborhood of $s=0$, we have

$$\frac{\xi_Q'}{\xi_Q}(s+1,a) = -\frac{\xi_Q'}{\xi_Q}(-s,a) = \sum_{m=0}^{\infty}(-1)^m\sum_{\rho_a\in Z_Q}\frac{1}{\rho^{m+1}}s^m. \tag{2.4}$$

Comparing Equations (2.1)–(2.4), we get

$$(-1)^m\sum_{\rho_a\in Z_Q}\frac{1}{\rho^{m+1}} = (-1)^m + \gamma_Q(m) + \tau_Q(m),$$

for $m\geq 0$. Hence, the definition of $\lambda_{n,a}$ yields

$$\lambda_{n,a} = \sum_{k=1}^{n}(-1)^{k-1}\binom{n}{k}\sum_{\rho_a\in Z_Q}\frac{1}{\rho^k} = 1 + \sum_{k=1}^{n}\binom{n}{k}\gamma_Q(k-1) + \sum_{k=1}^{n}\binom{n}{k}\tau_Q(k-1),$$

where

$$\tau_Q(0) = -\frac{1}{2}\log\pi + \frac{1}{2}\psi(1/2) \text{ and } \tau_Q(k-1) = (-1)^k\sum_{m=0}^{\infty}\frac{1}{(2m+1)^k}$$

https://doi.org/10.4153/S0008439524000250 Published online by Cambridge University Press

using that $\psi(z) = -\gamma - \frac{1}{z} + \sum_{k=1}^{\infty} \frac{z}{k(k+z)}$. Here, $\psi(s) = \frac{\Gamma'}{\Gamma}(s)$ is the logarithmic derivative of the Gamma function. Since $\psi(1/2) = -\gamma - 2\log 2$, we obtain

$$\begin{aligned}\lambda_{n,a} &= 1 - \frac{n}{2}(\log(4\pi) + \gamma) + \sum_{k=2}^{n}(-1)^k\binom{n}{k}\sum_{m=0}^{\infty}\frac{1}{(2m+1)^k} + \sum_{k=1}^{n}\binom{n}{k}\gamma_Q(k-1)\\ &= 1 - \frac{n}{2}(\log(4\pi) + \gamma) + \sum_{k=2}^{n}(-1)^k\binom{n}{k}\left(1-2^{-k}\right)\zeta(k) + \sum_{k=1}^{n}\binom{n}{k}\gamma_Q(k-1).\end{aligned}$$

The equality above implies Theorem 1.2.

## 2.3 Proof of Theorem 1.3

Let us note that

$$\sum_{k=2}^{n}(-1)^k\binom{n}{k}\left(1-2^{-k}\right)\zeta(k) = \sum_{k=2}^{n}(-1)^l\binom{n}{k}\frac{\zeta(k,1/2)}{2^k},$$

where $\zeta(s,a)$ is the Hurwitz zeta function defined in Section 1.2. With the notation of Flajolet and Vespas [7, Lines 2–4, p. 70], this is $A_n(1,2)$ and which is equal to

$$\frac{n}{2}\psi(n) + n\left(\gamma - \frac{1}{2} + \frac{1}{2}\log 2\right) + o(1),$$

where the o(1) error term above is exponentially small and oscillating and equal to

$$\frac{1}{2}\left(\frac{n}{\pi}\right)^{1/4}\exp(-\sqrt{2\pi n})\cos\left(\sqrt{2\pi n} - \frac{5\pi}{8}\right) + O\left(n^{-1/4}e^{-\sqrt{2\pi n}}\right).$$

Then we have

$$\lambda_{n,a} = \frac{n}{2}\log n + \frac{n}{2}(\gamma - 1 - \log 2\pi) + \sum_{k=1}^{n}\binom{n}{k}\gamma_Q(k-1) + O(1).$$

It remains to prove that

$$\sum_{k=1}^{n}\binom{n}{k}\gamma_Q(k-1) = O(\sqrt{n}\log n). \tag{2.5}$$

To do so, we follow very closely the lines of the proof of the corresponding result in [8, Theorem 6.1] or [16, Lemma 3.3] and it will be shortened. We use the following kernel function:

$$k_n(s) := \left(1 + \frac{1}{s}\right)^n - 1 = \sum_{k=1}^{n}\binom{n}{k}\frac{1}{s^k}.$$

The residue theorem gives

$$\sum_{k=1}^{n}\binom{n}{k}\gamma_Q(k-1) = \frac{1}{2i\pi}\int_C k_n(s)\left(-\frac{Q'}{Q}(s+1,a)\right)ds,$$

where $C$ is a contour enclosing the point $s = 0$ counterclockwise on a circle of small enough positive radius. The residue comes entirely from the singularity at $s = 0$, as no other singularities lie inside the contour. Let $T = \sqrt{n} + \varepsilon_n$, for some $0 < \varepsilon_n < 1$.

https://doi.org/10.4153/S0008439524000250 Published online by Cambridge University Press

Now we follow very closely the lines in [16, pp. 1106–1107] using that the function $\frac{Q'}{Q}(s,a)$ satisfies the properties[1]

$$\frac{Q'}{Q}(s,a) = \sum_{\rho_a;\ |\mathrm{Im}(\rho_a - s)|<1} \frac{1}{s-\rho_a} + O(\log(1+|s|)),$$

for $-2 < \mathrm{Re}(s) < 2$ and

$$\left|\frac{Q'}{Q}(s+1,a)\right| = O(\log^2 T),$$

for $-2 \le \mathrm{Re}(s) \le 2$, and we get

$$\sum_{k=1}^{n} \binom{n}{k} \gamma_Q(k-1) = \lambda_{-n,a,T} + O(\sqrt{n}\log n),$$

where

$$\lambda_{-n,a,T} = \sum_{\rho_a \in Z_Q;\ |Im(\rho_a)| \le T}^{*} \left(1 - \left(1 - \frac{1}{\rho_a}\right)^n\right),$$

with $T = \sqrt{n} + \varepsilon_n$. For $a = 1/2, 1/3, 1/4, 1/6$, under the RH, since $\left|1 - \frac{1}{\rho_a}\right| = 1$ and using formula of $N_Q(T)$ given in Section 1.2, we obtain $\lambda_{n,a,T} = O(T\log T + 1)$. Therefore, Equation (2.5) follows from that $\lambda_{-n,a,\sqrt{n}} = \lambda_{-n,a,\sqrt{n}} = O(\sqrt{n}\log n)$.

***Remark*** Since $2Q(s,a) := Z(s,a) + P(s,a)$, from Corollary 2.3 below and [6, Equation (1.18)], we obtain

$$\gamma_Q(n) = \frac{1}{2}\left(\delta_n(a) + \frac{(-1)^n}{n!}(l_n(a) + l_n(1-a))\right),$$

where $\delta_n(a) = \frac{|\log a|^n}{an!} + O(1)$ and $l_n(a)$ are the coefficients in the expansion of $\mathrm{Li}_s(e^{2\pi i a})$ at $s = 1$; for $a \notin \mathbb{Z}$, one has

$$\mathrm{Li}_s(e^{2\pi i a}) = \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} l_n(a)(s-1)^n.$$

## 2.4 Proof of Theorem 1.4

To show Theorem 1.4, we quote the following lemmas from [2, 3].

***Lemma 2.1*** [3, Theorem 1] *We set*

$$(s-1)\zeta(s,a) = 1 + \sum_{n=0}^{\infty} \gamma_n(a)(s-1)^{n+1}, \qquad 0 < a \le 1.$$

[1]These properties are well known for the Riemann zeta function. The proof for the function $Q(s,a)$ is exactly the same since the Riemann–von Mangoldt formula holds for $Q(s,a)$ (see [14, Proposition 2.5] or [18, p. 217]).

https://doi.org/10.4153/S0008439524000250 Published online by Cambridge University Press

*Then it holds that*

$$\gamma_n(a) = \frac{(-1)^n}{n!} \lim_{m\to\infty} \left( \sum_{k=0}^{m} \frac{\log^n(k+a)}{k+a} - \frac{\log^{n+1}(m+a)}{n+1} \right).$$

***Lemma 2.2*** [2, Equation (26)] *Let $0 < a \le 1$, and let $n$ be a nonnegative integer. Then one has*

$$\begin{aligned}\zeta^{(n)}(0,a) = {} & \left(\frac{1}{2} - a\right) |\log a|^n - n! + n! a \sum_{m=n}^{\infty} \frac{|\log a|^m}{m!} \\ & + (-1)^n n \int_0^{\infty} \frac{\varphi(x) \log^{n-1}(x+a)}{(x+a)^2} dx \\ & - (-1)^n n(n-1) \int_0^{\infty} \frac{\varphi(x) \log^{n-2}(x+a)}{(x+a)^2} dx,\end{aligned}$$

*where $\varphi(x) = \int_0^x (y - \lfloor y \rfloor - 1/2) dy$ is periodic with period 1 and satisfies $2\varphi(x) = x(x-1)$ if $0 \le x \le 1$.*

By using the lemmas above, we immediately obtain the following.

***Corollary 2.3*** *When $a > 0$ is sufficiently small,*

$$(s-1)Z(s,a) = 2 + \sum_{n=0}^{\infty} \delta_n(a)(s-1)^{n+1}, \qquad \delta_n(a) = \frac{|\log a|^n}{an!} + O(1),$$

$$Z(s,a) = \sum_{n=1}^{\infty} \varepsilon_n(a) s^n, \qquad \varepsilon_n(a) = O(|\log a|^n).$$

**Proof** The first formula and estimation are easily proved by Lemma 2.1 (see also [3, Theorem 2]). For the first integral in Lemma 2.2, one has

$$\int_0^1 \frac{\varphi(x) \log^{n-1}(x+a)}{(x+a)^2} dx \ll \int_0^1 \frac{\log^{n-1}(x+a)}{x+a} dx = O(|\log a|^n),$$

$$\int_1^{\infty} \frac{\varphi(x) \log^{n-1}(x+a)}{(x+a)^2} dx \ll \int_1^{\infty} \frac{\log^{n-1}(x+a)}{(x+a)^2} dx = O(1)$$

from $x < x + a$ when $x, a > 0$. In addition, we have

$$a \sum_{m=n}^{\infty} \frac{|\log a|^m}{m!} \le a \sum_{m=0}^{\infty} \frac{|\log a|^m}{m!} = a e^{|\log a|} = a e^{-\log a} = 1, \qquad 0 < a < 1/2.$$

Hence, we obtain

$$\zeta(s,a) = \sum_{n=0}^{\infty} \frac{\zeta^{(n)}(0,a)}{n!} s^n, \qquad \zeta^{(n)}(0,a) = O(|\log a|^n).$$

Therefore, we have $\varepsilon_n(a) = O(|\log a|^n)$ and the second formula in this corollary by the definition of $Z(s,a)$ and $Z(0,a) = \zeta(0,a) + \zeta(0,1-a) = 0$ (see [15, Equation (4.11)]). ∎

https://doi.org/10.4153/S0008439524000250 Published online by Cambridge University Press

**Proof of Theorem 1.4** Recall the functional equation

$$Z(1-s,a) = \Gamma_{\cos}(s)P(s,a), \qquad \Gamma_{\cos}(s) := \frac{2\Gamma(s)}{(2\pi)^s}\cos\Big(\frac{\pi s}{2}\Big)$$

(see [15, Lemma 4.11]). By using $\Gamma_{\cos}(s)\Gamma_{\cos}(1-s) = 1$, we have

$$2Q(s,a) = Z(s,a) + P(s,a) = Z(s,a) + \Gamma_{\cos}(1-s)Z(1-s,a).$$

Let $|s-1|$ be sufficiently small. Then, by $\lim_{s\to 1}(s-1)Q(s,a) = 1$, the equation above, and the definitions of $Q(s,a)$ and $\xi_Q(s,a)$, we have

$$\begin{aligned}
\frac{d^l}{ds^l}\big[s^{l-1}\log\xi_Q(s,a)\big]_{s=1} &= \frac{d^l}{ds^l}\Big[s^{l-1}\log\big((s-1)Q(s,a)\big) + s^{l-1}\log\big(s\pi^{-s/2}\Gamma(s/2)\big)\Big]_{s=1} \\
&= \frac{d^l}{ds^l}\left[s^{l-1}\log\left(\frac{s-1}{2}\Big(Z(s,a) + \Gamma_{\cos}(1-s)Z(1-s,a)\Big)\right)\right]_{s=1} + O_l(1) \\
&= \frac{d^l}{ds^l}\left[s^{l-1}\log\left(1 + \sum_{n=0}^{\infty}\Big(\delta_n'(a) + \varepsilon_n'(a)\Big)(s-1)^{n+1}\right)\right]_{s=1} + O_l(1),
\end{aligned}$$

where $\delta_n'(a)$ and $\varepsilon_n'(a)$ are defined by

$$\delta_n'(a) := \frac{\delta_n(a)}{2}, \qquad (s-1)\Gamma_{\cos}(1-s)Z(1-s,a) = 2\sum_{n=0}^{\infty}\varepsilon_n'(a)(s-1)^{n+1}.$$

Clearly, the second estimation in Corollary 2.3 implies

$$Z(1-s,a) = \sum_{n=1}^{\infty}\varepsilon_n(a)(1-s)^n, \qquad \varepsilon_n(a) = O(|\log a|^n).$$

Thus, we can see that $\varepsilon_n'(a) = O(|\log a|^{n+1})$ from $\lim_{s\to 1}(s-1)\Gamma_{\cos}(1-s) = -2$ and the fact that the function $(s-1)\Gamma_{\cos}(1-s)$ does not depend on $a$. Put $\eta_n(a) := \delta_n'(a) + \varepsilon_n'(a)$. Then, for $n \geq 0$, we have

$$\eta_n(a) = \frac{1}{n!}\frac{|\log a|^n}{2a} + O(|\log a|^{n+1}), \qquad a \to +0 \tag{2.6}$$

by Corollary 2.3. By virtue of

$$\begin{aligned}
(a_0x + a_1x^2 + a_2x^3 + \cdots)^m &= a_0^m x^m + \binom{m}{1}a_0^{m-1}a_1x^{m+1} + \cdots \\
(a_0x + a_1x^2 + a_2x^3 + \cdots)^{m-1} &= a_0^m x^{m-1} + \binom{m-1}{1}a_0^{m-2}a_1x^m + \cdots \\
&\vdots \\
(a_0x + a_1x^2 + a_2x^3 + \cdots)^1 &= \cdots + a_mx^m + \cdots,
\end{aligned}$$

where $m \in \mathbb{N}$ and $a_m, x \in \mathbb{C}$, the coefficient of $(s-1)^l$ in the function

$$f(s,a) := \sum_{m=1}^{\infty}\frac{(-1)^{m+1}}{m}\left(\sum_{n=0}^{\infty}\eta_n(a)(s-1)^{n+1}\right)^m$$

https://doi.org/10.4153/S0008439524000250 Published online by Cambridge University Press

is expressed as

$$\frac{(-1)^{l+1}}{l}\big(\eta_0(a)\big)^l + \frac{(-1)^l}{l-1}\binom{l-1}{1}\eta_0(a)^{l-2}\eta_1(a) + \cdots + \frac{(-1)^{1+1}}{1}\eta_{l-1}(a). \tag{2.7}$$

Note that the function above is estimated by

$$\frac{(-1)^{l+1}}{l}\big(\eta_0(a)\big)^l + O_l\big(\eta_0(a)^{l-2}\eta_1(a)\big) = \frac{(-1)^{l+1}}{l}(2a)^{-l} + O_l\big(a^{1-l}|\log a|\big) \tag{2.8}$$

from (2.6) when $a \to +0$. We can find that

$$(s-1)\Big(Z(s,a) + \Gamma_{\cos}(1-s)Z(1-s,a)\Big) = 1 + \sum_{n=0}^{\infty}\eta_n(a)(s-1)^{n+1}$$

is analytic when $|s-1| < 1$ form the poles of $Z(s,a)$ and $\Gamma_{\cos}(1-s)$. So we can choose $|s-1| > 0$ such that

$$\sum_{n=0}^{\infty}\big|\eta_n(a)\big||s-1|^{n+1} < \frac{1}{2}.$$

Then, from (2.7), the Leibniz product rule, the definition of $\eta_n(a)$, and the Taylor expansion of $\log(1+x)$ with $|x| < 1$, one has

$$\begin{aligned}
&\frac{d^l}{ds^l}\Big[s^{l-1}\log\xi_Q(s,a)\Big]_{s=1} = \frac{d^l}{ds^l}\Big[s^{l-1}f(s,a)\Big]_{s=1} + O_l(1)\\
(\flat)\qquad &= \binom{l}{l}\frac{(-1)^{l+1}}{l}l!\big(\eta_0(a)\big)^l + O_l\big(\eta_0(a)^{l-2}\eta_1(a)\big)\\
(\natural)\qquad &\quad + \binom{l}{l-1}(l-1)\frac{(-1)^l}{l-1}(l-1)!\big(\eta_0(a)\big)^{l-1} + O_l\big(\eta_0(a)^{l-3}\eta_1(a)\big)\\
(\sharp)\qquad &\quad + \cdots + \binom{l}{1}(l-1)!\frac{(-1)^{1+1}}{1}\big(\eta_0(a)\big)^1 + O_l(1).
\end{aligned}$$

Note that $(\flat)$ comes from $f^{(l)}(s,a)$, $(\natural)$ is deduced by $f^{(l-1)}(s,a)$, and $(\sharp)$ derives from $f^{(1)}(s,a)$, $f^{(0)}(s,a)$, and $O_l(1)$ in the left-hand side of the formula above. Therefore, by (2.8), we obtain

$$\begin{aligned}
\frac{d^l}{ds^l}\Big[s^{l-1}\log\xi_Q(s,a)\Big]_{s=1} &= (-1)^{l+1}(l-1)!\big(\eta_0(a)\big)^l + O_l\big(\eta_0(a)^{l-2}\eta_1(a)\big)\\
&= (-1)^{l+1}\frac{(l-1)!}{(2a)^l} + O_l\big(a^{1-l}|\log a|\big),
\end{aligned}$$

which implies Theorem 1.4. ∎

At the end of the paper, we give numerical computation for $\lambda_{n,a}$ by Mathematica 13.0. Let

$$\lambda_{n,a}^{[k]} := \frac{1}{(n-1)!}\frac{d^n}{ds^n}\Big[s^{n-1}\log\xi_Q(s,a)\Big]_{s=1-10^{-k}}, \qquad \lambda_{n,a}^{*} := \frac{(-1)^{n+1}}{(2a)^n}.$$

Then, we have the following.

https://doi.org/10.4153/S0008439524000250 Published online by Cambridge University Press

For $n = 1$, we have

$$\begin{array}{lll}
a := 2^{-17} & \lambda_{1,a}^{[10]} = 65,537\ldots & \lambda_{1,a}^{[10]}/\lambda_{1,a}^{*} = 1.00001\ldots \\
a := 2^{-18} & \lambda_{1,a}^{[10]} = 131,074\ldots & \lambda_{1,a}^{[10]}/\lambda_{1,a}^{*} = 1.00002\ldots \\
a := 2^{-19} & \lambda_{1,a}^{[10]} = 262,151\ldots & \lambda_{1,a}^{[10]}/\lambda_{1,a}^{*} = 1.00003\ldots \\
a := 2^{-17} & \lambda_{1,a}^{[11]} = 65,536.6\ldots & \lambda_{1,a}^{[11]}/\lambda_{1,a}^{*} = 1.00001\ldots \\
a := 2^{-18} & \lambda_{1,a}^{[11]} = 131,073\ldots & \lambda_{1,a}^{[11]}/\lambda_{1,a}^{*} = 1.00001\ldots \\
a := 2^{-19} & \lambda_{1,a}^{[11]} = 262,145\ldots & \lambda_{1,a}^{[11]}/\lambda_{1,a}^{*} = 1.00000\ldots \\
a := 2^{-17} & \lambda_{1,a}^{[12]} = 655,365\ldots & \lambda_{1,a}^{[12]}/\lambda_{1,a}^{*} = 1.00001\ldots \\
a := 2^{-18} & \lambda_{1,a}^{[12]} = 131,073\ldots & \lambda_{1,a}^{[12]}/\lambda_{1,a}^{*} = 1.00000\ldots \\
a := 2^{-19} & \lambda_{1,a}^{[12]} = 262,145\ldots & \lambda_{1,a}^{[12]}/\lambda_{1,a}^{*} = 1.00000\ldots
\end{array}$$

For $n = 2$, we have

$$\begin{array}{lll}
a := 2^{-17} & \lambda_{2,a}^{[10]} = -4.29352\ldots \times 10^{9} & \lambda_{2,a}^{[10]}/\lambda_{2,a}^{*} = 0.999663\ldots \\
a := 2^{-18} & \lambda_{2,a}^{[10]} = -1.7177\ldots \times 10^{10} & \lambda_{2,a}^{[10]}/\lambda_{2,a}^{*} = 0.999836\ldots \\
a := 2^{-19} & \lambda_{2,a}^{[10]} = -6.87162\ldots \times 10^{10} & \lambda_{2,a}^{[10]}/\lambda_{2,a}^{*} = 0.999952\ldots \\
a := 2^{-17} & \lambda_{2,a}^{[11]} = -4.29478\ldots \times 10^{9} & \lambda_{2,a}^{[11]}/\lambda_{2,a}^{*} = 0.999956\ldots \\
a := 2^{-18} & \lambda_{2,a}^{[11]} = -1.71753\ldots \times 10^{10} & \lambda_{2,a}^{[11]}/\lambda_{2,a}^{*} = 0.999736\ldots \\
a := 2^{-19} & \lambda_{2,a}^{[11]} = -6.87149\ldots \times 10^{10} & \lambda_{2,a}^{[11]}/\lambda_{2,a}^{*} = 0.999933\ldots \\
a := 2^{-17} & \lambda_{2,a}^{[12]} = -4.29477\ldots \times 10^{9} & \lambda_{2,a}^{[12]}/\lambda_{2,a}^{*} = 0.999955\ldots \\
a := 2^{-18} & \lambda_{2,a}^{[12]} = -1.6911\ldots \times 10^{10} & \lambda_{2,a}^{[12]}/\lambda_{2,a}^{*} = 0.984353\ldots \\
a := 2^{-19} & \lambda_{2,a}^{[12]} = -6.87187\ldots \times 10^{10} & \lambda_{2,a}^{[12]}/\lambda_{2,a}^{*} = 0.999989\ldots
\end{array}$$

**Acknowledgements** The authors want to thank the anonymous referees for their many insightful comments and suggestions.

## References

[1] T. M. Apostol, *Introduction to analytic number theory*, Undergraduate Texts in Mathematics, Springer, New York, 1976.
[2] T. M. Apostol, *Formulas for higher derivatives of the Riemann zeta function*. Math. Comput. **44**(1985), no. 169, 223–232.
[3] B. C. Berndt, *On the Hurwitz zeta-function*. Rocky Mountain J. Math. **2**(1972), no. 1, 151–157.
[4] E. Bombieri and J. Lagarias, *Complements to Li's criterion for the Riemann hypothesis*. J. Number Theory 77(1999), 274–287.
[5] F. Brown, *Li's criterion and zero-free regions of L -functions*. J. Number Theory **111**(2005), 1–32.
[6] M. W. Coffey, *Series representations for the Stieltjes constants*. Rocky Mountain J. Math. **44**(2014), no. 2, 443–477.
[7] P. Flajolet and L. Vepstas, *On differences of zeta values*. J. Comput. Appl. Math. **220**(2008), nos. 1–2, 58–73.
[8] J. C. Lagarias, *Li coefficients for automorphic L-functions*. Ann. Inst. Fourier (Grenoble) **57**(2007), 1689–1740.
[9] A. Laurinčikas and R. Garunkštis, *The Lerch zeta-function*, Kluwer Academic Publishers, Dordrecht, 2002.
[10] X.-J. Li, *The positivity of a sequence of numbers and the Riemann hypothesis*. J. Number Theory **65**(1997), 325–333.
[11] K. Mazhouda and L. Smajlović, *Evaluation of the Li coefficients on function fields and applications*. Eur. J. Math. 5(2019), no. 2, 540–550.

https://doi.org/10.4153/S0008439524000250 Published online by Cambridge University Press

[12] K. Mazhouda and B. Sodaïgui, *The Li–Sekatskii coefficients for the Selberg class*. Int. J. Math. **33**(2022), no. 12, Article no. 2250075, 23 pp.
[13] T. Nakamura, *The functional equation and zeros on the critical line of the quadrilateral zeta function*. J. Number Theory **233**(2022), 432–455.
[14] T. Nakamura, *On Lerch's formula and zeros of the quadrilateral zeta function*. Preprint, 2022, arXiv:2001.01981.
[15] T. Nakamura, *On zeros of bilateral Hurwitz and periodic zeta and zeta star functions*. Rocky Mountain J. Math. **53**(2023), no. 1, 157–176.
[16] S. Omar and K. Mazhouda, *The Li criterion and the Riemann hypothesis for the Selberg class II*. J. Number Theory **130**(2010), no. 4, 1109–1114.
[17] L. Smajlović, *On Li's criterion for the Riemann hypothesis for the Selberg class*. J. Number Theory **130**(2010), no. 4, 828–851.
[18] E. C. Titchmarsh, *Theory of the Riemann zeta-function*. 2nd ed., Clarendon Press, Oxford, 1986.
[19] A. Voros, *Sharpenings of Li's criterion for the Riemann hypothesis*. Math. Phys. Anal. Geom. **9**(2006), 53–63.

*Department of Mathematics, University of Prishtina, Mother Theresa, No. 5, 10000 Prishtina, Kosovo*
*e-mail*: kajtaz.bllaca@uni-pr.edu

*Higher Institute of Applied Sciences and Technology, University of Sousse, Sousse 4003, Tunisia INSA Hauts-De-France, University Polytechnique Hauts-De-France, FR CNRS 2037, CERAMATHS, F-59313 Valenciennes, France*
*e-mail*: kamel.mazhouda@fsm.rnu.tn

*Department of Liberal Arts, Faculty of Science and Technology, Tokyo University of Science, 2641 Yamazaki, Noda-shi, Chiba-ken 278-8510, Japan*
*e-mail*: nakamuratakashi@rs.tus.ac.jp

https://doi.org/10.4153/S0008439524000250 Published online by Cambridge University Press