ADDING $\aleph_\omega$-MANY COHEN REALS

PEDRO MARUN, SAHARON SHELAH, AND COREY BACAL SWITZER

Abstract. Abstractly, the generic extensions after $\aleph_\omega$-many Cohen reals and $\aleph_{\omega+1}$-many Cohen reals must be different for reasons of uniform density of the relevant Boolean algebras. Nevertheless this is not satisfying and it would be nice to pin the difference between the two models down to some mathematical or combinatorial principle. In this article we provide such a principle.

§1. Introduction. Let $\mathrm{Add}(\omega, \aleph_\omega)$ be the poset for adding $\aleph_\omega$-many Cohen reals. In $V^{\mathrm{Add}(\omega,\aleph_\omega)}$, the continuum must be at least $\aleph_\omega$. By König's lemma, the continuum has uncountable cofinality, hence it must be at least $\aleph_{\omega+1}$. One can then ask whether this is different from adding $\aleph_{\omega+1}$-many Cohen reals, i.e., forcing with $\mathrm{Add}(\omega, \aleph_{\omega+1})$. More precisely, if G is $\mathrm{Add}(\omega, \aleph_\omega)$-generic over V, is there an H which is $\mathrm{Add}(\omega, \aleph_{\omega+1})$-generic over V such that $V[G] = V[H]$? The (folklore) answer to this is "No," and in fact $V[G]$ doesn't even contain a filter which is $\mathrm{Add}(\omega, \aleph_{\omega+1})$-generic over V, Proposition 2.5 below. However, the proof of this is somewhat unsatisfying, as it relies on abstract forcing considerations involving the (uniform) densities of the complete Boolean algebras. It is therefore desirable to exhibit a combinatorial or "mathematical" principle which distinguishes the two models. The purpose of this note is to provide one.

Definition 1.1. Let $\sigma \leq \theta < \mu < \lambda$ be cardinals. The statement $\mathsf{Pr}(\sigma, \theta, \mu, \lambda)$ states that there is an $\overline{\mathcal{F}}$ satisfying the following:

(A) $\overline{\mathcal{F}} = \langle \mathcal{F}_c : c \in [\lambda]^\sigma \rangle$ is a sequence so that for all $c \in [\lambda]^\sigma$ we have $\mathcal{F}_c \subseteq \mathcal{P}(c)$.
(B) For all $c \in [\lambda]^\sigma$ we have that $\mathcal{F}_c$ has cardinality at most θ.
(C) For every $A \in [\lambda]^\lambda$ there is a sequence $\langle B_i : i < \mu \rangle$ so that
  (a) $A = \bigcup_{i<\mu} B_i$ and
  (b) $B_i \cap c \in \mathcal{F}_c$ for all $c \in [\lambda]^\sigma$ and $i < \mu$.

We will show the following.

Theorem 1.2. *Assume* GCH. *Let* $\mathbb{P}_0 = \mathrm{Add}(\omega, \aleph_\omega)$ *and let* $\mathbb{P}_1 = \mathrm{Add}(\omega, \aleph_{\omega+1})$. *The following hold*:

(A) *For all* $n < \omega$ $\Vdash_{\mathbb{P}_0} \mathsf{Pr}(\aleph_n, \aleph_{n+1}, \aleph_\omega, \aleph_{\omega+1})$.
(B) *For all* $n < \omega$ $\Vdash_{\mathbb{P}_1} \neg\mathsf{Pr}(\aleph_n, \aleph_{n+1}, \aleph_\omega, \aleph_{\omega+1})$.

Received November 27, 2025 UTC.
2020 *Mathematics Subject Classification.* Primary 03E05, 03E35.
Key words and phrases. Cohen reals, forcing, singular cardinals.

© The Author(s), 2026. Published by Cambridge University Press on behalf of The Association for Symbolic Logic. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (https://creativecommons.org/licenses/by/4.0/), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

0022-4812/00/0000-0000
DOI:10.1017/jsl.2026.10241

https://doi.org/10.1017/jsl.2026.10241 Published online by Cambridge University Press

Thus the above says that in particular $\mathsf{Pr}(\aleph_0, \aleph_1, \aleph_\omega, \aleph_{\omega+1})$ is the desired principle. In fact Theorem 1.2 holds in more generality—namely we can take μ to be a strong limit cardinal of countable cofinality, and $\sigma < \mu$ infinite and we get that adding μ-many Cohen reals forces $\mathsf{Pr}(\sigma, 2^\sigma, \mu, \mu^+)$ while adding μ^+-many Cohen reals forces $\neg\mathsf{Pr}(\sigma, 2^\sigma, \mu, \mu^+)$. These two pieces are proved below as Lemmas 3.2 and 4.1, respectively.

The rest of this article is organized as follows. In Section 2, we make some preliminary observations about adding $\aleph_\omega$ versus adding $\aleph_{\omega+1}$ Cohen reals. In Section 3 we prove the generalization of Item (1) of Theorem 1.2 while in Section 4 we prove the generalization of Item (2) of Theorem 1.2. The article concludes with some questions for further research. Throughout our notation is mostly standard, conforming, e.g., to that of [7] or [9]. Those texts are referred to for any undefined terms.

§2. Some preliminary observations. If κ is an infinite cardinal, we let $\mathrm{Add}(\omega, \kappa)$ be the poset for adding κ-many Cohen reals. More precisely, $\mathrm{Add}(\omega, \kappa)$ is the set of finite partial functions $\kappa \rightharpoonup 2$, ordered by reverse inclusion. We will argue that, if $\kappa < \lambda$, then a generic extension of V by $\mathrm{Add}(\omega, \kappa)$ does not contain a filter that is $\mathrm{Add}(\omega, \lambda)$-generic over V. This result is folklore but a proof is hard to find in the published literature; we will follow the outline in [5].

We collect some facts from the elementary theory of Boolean algebras. If $\mathbb{B}$ is a Boolean algebra and $b \in \mathbb{B}$, then $\mathbb{B}_b = \{c \in \mathbb{B} : c \leq b\}$, which can be made into a Boolean algebra (with top element b) in an obvious way. We let $\mathbb{B}^+ = \mathbb{B} \setminus \{0\}$. If $\mathbb{A}$ is a subalgebra of $\mathbb{B}$, then $\mathbb{A}$ is said to be a *regular subalgebra of* $\mathbb{B}$ iff for all $X \subseteq \mathbb{A}$, if $\bigvee^{\mathbb{A}} X$ exists, then so does $\bigvee^{\mathbb{B}} X$ and moreover $\bigvee^{\mathbb{A}} X = \bigvee^{\mathbb{B}} A$. If $\mathbb{B}$ is complete, we will say that $\mathbb{A} \subseteq \mathbb{B}$ is a *complete subalgebra* if it is a regular subalgebra of $\mathbb{B}$ that is also complete as a Boolean algebra.

We will need the following cardinal function.

Definition 2.1. If $\mathbb{B}$ is a Boolean algebra, we define its *density* by

$$d(\mathbb{B}) := \min\{|D| : D \text{ is dense in } \mathbb{B}\},$$

where D is *dense* in $\mathbb{B}$ iff $D \subseteq \mathbb{B}^+$ and for all $b \in \mathbb{B}^+$ there exists $d \in D$ with $d \leq b$.

We say that $\mathbb{B}$ has *uniform density* iff $d(\mathbb{B}) = d(\mathbb{B}_b)$ for all $b \in \mathbb{B}^+$.

Unlike the topological notion, the Boolean algebraic density is nicely monotone.

Lemma 2.2 (Folklore). *Let $\mathbb{A}$ be a complete subalgebra of a complete Boolean algebra $\mathbb{B}$. Then $d(\mathbb{A}) \leq d(\mathbb{B})$.*

Proof. Let $D \subseteq \mathbb{B}$ be dense. For $d \in D$, let

$$u_d := \bigwedge\{a \in \mathbb{A} : d \leq a\}.$$

We will be done if we can argue that $\{u_d : d \in D\}$ is dense in $\mathbb{A}$. So, fix $a \in \mathbb{A}$. By the density of D, there exists some $d \in D$ with $d \leq a$, hence $u_d \leq a$. ⊣

Lemma 2.3 [6, Lemma 25.5(a)]. *Let $\mathbb{B}$ and $\mathbb{C}$ be complete Boolean algebras. Let G and H be $\mathbb{B}$ and $\mathbb{C}$ generic over V, respectively, and suppose that $V[G] = V[H]$. Then there exist $\pi \in V$, $b \in \mathbb{B}$, and $c \in \mathbb{C}$ such that $\pi : \mathbb{B}_b \cong \mathbb{C}_c$ and $\pi[\mathbb{B}_b \cap G] = \mathbb{C}_c \cap H$.*

https://doi.org/10.1017/jsl.2026.10241 Published online by Cambridge University Press

The *Cohen algebra on* κ is $\mathbb{C}_\kappa := \mathrm{ro}(\mathrm{Add}(\omega, \kappa))$, i.e., the Boolean completion of $\mathrm{Add}(\omega, \kappa)$.

LEMMA 2.4. *Let* κ *be an infinite cardinal. The algebra* $\mathbb{C}_\kappa$ *is homogeneous and has* (*uniform*) *density* κ.

PROOF. For the homogeneity, see [8, Corollary 12.5]. The uniform density assertion follows from homogeneity and the fact that $d(\mathbb{C}_\kappa)$ is also the smallest size of a dense subset of $\mathrm{Add}(\omega, \kappa)$, which is easily seen to be κ. ⊣

PROPOSITION 2.5 (Folklore). *Let* $\kappa < \lambda$ *be infinite cardinals. In* $V^{\mathrm{Add}(\omega,\kappa)}$ *there is no* $\mathrm{Add}(\omega, \lambda)$*-filter that is generic over* V.

PROOF. Suppose not. Then we can find G which is $\mathbb{C}_\kappa$-generic over V and H which is $\mathbb{C}_\lambda$-generic over V such that $H \in V[G]$. Obviously, $V \subseteq V[H] \subseteq V[G]$. By the intermediate model theorem (see [7, Lemma 15.43]), there exists $\mathbb{B}$ a complete regular subalgebra of $\mathbb{C}_\kappa$ such that $V[H] = V[G \cap \mathbb{B}]$. By Lemma 2.3, there exist $b \in \mathbb{B}$ and $c \in \mathbb{C}_\lambda$ such that $\mathbb{B}_b \cong (\mathbb{C}_\lambda)_c$ (in V). But then, by Lemmas 2.2 and 2.4,

$$\lambda = d((\mathbb{C}_\lambda)_c) = d(\mathbb{B}_b) \leq d(\mathbb{B}) = \kappa,$$

which is a contradiction. ⊣

REMARK 2.6. If we don't insist on fixing the ground model, the previous theorem can fail. In other words, it is consistent (modulo large cardinals) that there is a pair of models $V \subseteq W$ such that adding few Cohen reals to W adds a lot of Cohen reals to V (see [3, 4]).

Towards a slightly more mathematical distinction between the two models we also observe the following.

PROPOSITION 2.7. *Assume* CH. *Let* G_0 *be* $\mathrm{Add}(\omega, \aleph_\omega)$*-generic over* V *and let* G_1 *be* $\mathrm{Add}(\omega, \aleph_{\omega+1})$*-generic over* V.

(A) *In* $V[G_0]$ *there are* $\aleph_1$*-many Borel functions* $\{f_\alpha : \alpha \in \omega_1\}$ *and a set of* $\aleph_\omega$*-many reals* $A = \{a_\alpha : \alpha \in \aleph_\omega\}$, *so that the following hold*:
  (a) *For all* $\alpha < \omega_1$ *the domain of* f_α *is* $[\omega^\omega]^\omega$.
  (b) *For every* $x \in \omega^\omega$ *there is a* $c \in [\aleph_\omega]^\omega \cap V$ *and an* $\alpha < \omega_1$ *so that* $x = f_\alpha(\{a_\xi : \xi \in c\})$.

(B) *In* $V[G_1]$ *the following holds*: *for every* $\aleph_1$*-many Borel functions* $\{f_\alpha : \alpha \in \omega_1\}$ *and every set of* $\aleph_\omega$*-many reals* $A = \{a_\alpha : \alpha \in \aleph_\omega\}$, *there is an* $x \in \omega^\omega$ *so that* $x \notin \{f_\alpha(\{a_\xi : \xi \in c\}) : \alpha \in \omega_1\}$ *for any* $c \in [\aleph_\omega]^\omega \cap V$.

PROOF. Each part is a consequence of the ccc. For the first part let $\{f_\alpha : \alpha \in \omega_1\}$ be the ground model Borel functions and let $A = \{a_\alpha : \alpha \in \aleph_\omega\}$ be the set of Cohen generic reals coming from G_0. If $\dot{x}$ is a nice name for a real then it is well known, see, e.g., [12, Theorem 4.1.2], that there are $c \in [\aleph_\omega]^\omega \cap V$ so that $\dot{x}$ is forced to be in $V[a_\xi : \xi \in c]$ and in that model there is an $\alpha < \omega_1$ so that $\dot{x}^{G_0} = f_\alpha(\{a_\xi : \xi \in c\})$. For the second part, observe that any $\aleph_\omega$-many reals and any $\aleph_1$-many Borel functions were added by $\aleph_\omega$-many of the Cohen reals in G_1 and hence there is a Cohen real which is generic over an intermediate model containing all of them. ⊣

Arguably, Proposition 2.7 provides a more concrete distinction between the $\mathrm{Add}(\omega, \aleph_\omega)$- and $\mathrm{Add}(\omega, \aleph_{\omega+1})$-extensions than that offered by Proposition 2.5.

https://doi.org/10.1017/jsl.2026.10241 Published online by Cambridge University Press

However, from our perspective, this is still not fully satisfactory because (A)(b) and (B) in Proposition 2.7 make explicit reference to the ground model V, which is meta-mathematical in nature. This should be contrasted with Theorem 1.2, which is purely combinatorial.

We will now sketch why there is no concrete cardinal characteristic separating $V^{\mathrm{Add}(\omega,\aleph_\omega)}$ from $V^{\mathrm{Add}(\omega,\aleph_{\omega+1})}$. For any unproven assertions in this paragraph, refer to [2, Section 11.3]. To start, it's well known that the first $\aleph_1$-many Cohen reals added will form a (Cohen indestructible) non-meager family, hence $\mathrm{non}(\mathcal{M}) = \aleph_1$ in both models. Meanwhile, the covering of the meager ideal, $\mathrm{cov}(\mathcal{M})$, must be $2^{\aleph_0} = \aleph_{\omega+1}$ in both models. For the $\mathrm{Add}(\omega,\aleph_{\omega+1})$-extension this follows by the regularity of $\aleph_{\omega+1}$. For the $\mathrm{Add}(\omega,\aleph_\omega)$-extension, we note that by [1, Theorem 5.1], $\mathrm{cov}(\mathcal{M})$ has uncountable cofinality and hence cannot be $\aleph_\omega$. On the other hand, as $\mathrm{Add}(\omega,\aleph_\omega) \cong \mathrm{Add}(\omega,\aleph_\omega) \times \mathrm{Add}(\omega,\aleph_n)$ for any $n < \omega$, we obtain that $\mathrm{cov}(\mathcal{M}) > \aleph_n$ for all $n < \omega$. Putting everything together, we see that the cardinals in the Cichoń diagram are the same for both models. Applying the inequalities

$$\begin{aligned} \mathrm{cov}(\mathcal{M}) \leq \mathfrak{r} \leq \min\{\mathfrak{u},\mathfrak{i}\},\\ \mathfrak{m} \leq \mathfrak{p} \leq \mathfrak{t} \leq \mathfrak{h} \leq \mathfrak{b} \leq \mathrm{non}(\mathcal{M}),\\ \max\{\mathfrak{e},\mathfrak{s}\} \leq \mathrm{non}(\mathcal{M}), \end{aligned}$$

we see that all of the aforementioned characteristics agree in both models. By different methods, $\mathfrak{a} = \mathfrak{g} = \aleph_1$ in both models.

We conclude this section by making some simple observations about the principle $\mathsf{Pr}(\sigma,\theta,\mu,\lambda)$ in an attempt to clarify it and prepare the reader for the arguments in Lemmas 3.2 and 4.1. All of these are easy, as should be obvious to the reader who is clear on what the principle states.

PROPOSITION 2.8. *Let $\sigma \leq \theta \leq \mu \leq \lambda$ be infinite cardinals.*

(A) *If $\theta = 2^\sigma$, then $\mathsf{Pr}(\sigma,\theta,\mu,\lambda)$ is true for any choice of μ and λ.*
(B) *If $\lambda = \mu^+$, the statement $\mathsf{Pr}(\sigma,\theta,\mu,\lambda)$ is unchanged by allowing in clause (C) of the definition that A has size $\leq \lambda$.*
(C) *$\mathrm{Pr}(\sigma,\theta,\mu,\lambda)$ is monotone in θ and μ. More precisely, if $\theta_0 \leq \theta_1$ or $\mu_0 \leq \mu_1$ are infinite cardinals, then $\mathrm{Pr}(\sigma,\theta_0,\mu_0,\lambda)$ implies $\mathrm{Pr}(\sigma,\theta_1,\mu_1,\lambda)$.*

PROOF. For (A) note that if $\theta = 2^\sigma$ then letting $\mathcal{F}_c := \mathcal{P}(c)$ satisfies the requirements of the principle.

For (B), note that since $\sigma \leq \theta$ we have that for any $\overline{\mathcal{F}}$ as in the statement of Definition 1.1 we can define $\overline{\mathcal{F}}'$ by $\mathcal{F}'_c := \{\emptyset\} \cup \{\{\alpha\} \mid \alpha \in c\} \cup \mathcal{F}_c$ and $\mathcal{F}'_c$ will still have size at most θ. Now for each $A \subseteq \lambda$ of size $<\lambda$, by $\lambda = \mu^+$ we can enumerate A (possibly with repetitions) as $A = \{\alpha_\xi : \xi < \mu\}$. Then $\{\alpha_\xi\} \cap c \in \mathcal{F}'_c$ for any $c \in [\lambda]^\sigma$.

For (C), any witness to $\mathrm{Pr}(\sigma,\theta_0,\mu_0,\lambda)$ will witness $\mathrm{Pr}(\sigma,\theta_1,\mu_0,\lambda)$. For the monotonicity in μ, let $\mathcal{F}$ witness $\mathrm{Pr}(\sigma,\theta_0,\mu_0,\lambda)$ and note that we may assume without loss of generality that $\emptyset \in \mathcal{F}_c$ for every $c \in [\lambda]^\sigma$. Given $A \in [\lambda]^\lambda$, let $\langle B_\xi : \xi < \mu_0\rangle$ witness Definition 1.1(C) and set $B_\xi = \emptyset$ for $\xi \in [\mu_0,\mu_1)$. This works. ⊣

As a consequence of the above we have the following.

LEMMA 2.9. *If* GCH *holds then $\mathsf{Pr}(\sigma,\theta,\mu,\lambda)$ holds for every infinite $\sigma < \theta$.*

https://doi.org/10.1017/jsl.2026.10241 Published online by Cambridge University Press

§3. Adding few Cohens. In this section we prove the promised generalization of part 1 of Theorem 1.2. First, a definition is given as follows.

DEFINITION 3.1. Let $\mathbb{P}$ be a forcing poset and κ be a cardinal. We say that $\mathbb{P}$ has the *κ-covering property* iff for every $p \in \mathbb{P}$ and every $\mathbb{P}$-name $\dot{X}$ for a set of ordinals with $p \Vdash |\dot{X}| < \kappa$, there exists $q \leq p$ and Y with $|Y| < \kappa$ and $q \Vdash \dot{X} \subseteq \check{Y}$.

Clearly, if $\mathbb{P}$ has the κ-covering property, then forcing with $\mathbb{P}$ preserves κ. Also, if κ is regular, then every κ-cc poset has the κ-covering property.

In the next lemma, if κ is a cardinal and we're working in a generic extension $V^{\mathbb{P}}$, then the symbol $|\kappa|$ refers to the cardinality of κ as computed in $V^{\mathbb{P}}$.

LEMMA 3.2. *Assume $2^\sigma = \theta < \mu < \lambda$. Let $\mathbb{P}$ be a forcing notion that has the σ^+-covering property and such that $\Vdash |\dot{G}| \leq |\mu|$. Then $\Vdash_{\mathbb{P}} \mathsf{Pr}(|\sigma|, |\theta|, |\mu|, |\lambda|)$.*

Note that the first part of Theorem 1.2 follows from the above by assuming GCH and subbing in $(\aleph_n, \aleph_{n+1}, \aleph_\omega, \aleph_{\omega+1})$ for $(\sigma, \theta, \mu, \lambda)$.

PROOF. Fix $\sigma, \theta, \mu, \lambda$, and $\mathbb{P}$ as in the statement of the theorem. Since $\mathbb{P}$ has the σ^+-covering property, there is a name $\dot{\Phi}$ such that the following is forced by $\mathbb{P}$: "$\dot{\Phi} : [\lambda]^{|\sigma|} \to ([\lambda]^\sigma)^V$ and for all $c \in [\lambda]^{|\sigma|}$, $c \subseteq \dot{\Phi}(c)$."

Working in $V^{\mathbb{P}}$, for each $c \in [\lambda]^{|\sigma|}$ let $\mathcal{F}_c = \{a \cap c : a \in V \cap \mathcal{P}(\dot{\Phi}(c))\}$. Note that for every c we have that $\mathcal{F}_c$ has cardinality at most $(2^\sigma)^V = |\theta|$ and hence $\overline{\mathcal{F}} = \langle \mathcal{F}_c : c \in [\lambda]^\sigma \rangle$ satisfies the first two clauses of Definition 1.1. Back in V let $\dot{\overline{\mathcal{F}}}$ name this sequence and for each $\dot{c}$ let $\dot{\mathcal{F}}_{\dot{c}}$ be the name for the associated element. We need to see that the third clause of Definition 1.1 is forced.

Suppose that $\dot{A}$ is a $\mathbb{P}$-name such that $\Vdash \dot{A} \subseteq \lambda$. For each $q \in \mathbb{P}$, let $B_q = \{\xi < \lambda : q \Vdash \xi \in \dot{A}\}$. Note that $\Vdash \dot{A} = \bigcup_{q \in \dot{G}} B_q$. Now let $\dot{c}$ be a $\mathbb{P}$-name for a $|\sigma|$-sized subset of λ. If $q \in \mathbb{P}$, then $\Vdash B_q \cap \dot{c} = B_q \cap \dot{\Phi}(\dot{c}) \cap \dot{c} \in \dot{\mathcal{F}}_{\dot{c}}$ because $\Vdash \dot{\Phi}(\dot{c}) \in V$. $\dashv$

REMARK 3.3. The proof of the previous lemma shows something slightly more general, because it produces a decomposition for each subset of λ, not just those in $[|\lambda|]^{|\lambda|}$. Also, the decomposition is of size at most $|\mu|$, not exactly μ (to get exactly μ, simply pad with $\emptyset$'s).

§4. Adding many Cohens. Now we prove the failure of the principle after adding μ^+-many Cohen reals thus complementing Lemma 3.2.

LEMMA 4.1. *Assume μ is a strong limit cardinal of countable cofinality, $\sigma < \theta < \mu$, and $\lambda = \mu^+$. If $\mathbb{P} = \mathrm{Add}(\omega, \lambda)$, then $\Vdash_{\mathbb{P}} \neg \mathsf{Pr}(\sigma, \theta, \mu, \lambda)$.*

PROOF. We argue by contradiction. By homogeneity of $\mathbb{P}$, we may assume that $\Vdash_{\mathbb{P}} \mathrm{Pr}(\sigma, \theta, \mu, \lambda)$. Let $\dot{\overline{\mathcal{F}}}$ witness this. We aim to produce a name for a λ-sized subset of λ which cannot be decomposed as in Definition 1.1. Actually, we will produce θ^+ many sets and show by contradiction they cannot all be decomposed as we want.

Let $\dot{g}$ be a $\mathbb{P}$-name for the generic function $\lambda \to 2$. For $\xi < \theta^+$, let $\dot{A}_\xi$ be a name for the set $\{\alpha < \lambda : \dot{g}(\theta^+ \cdot \alpha + \xi) = 1\}$. An easy genericity argument confirms that each $\dot{A}_\xi$ is forced to have has size λ and that they are all distinct.

https://doi.org/10.1017/jsl.2026.10241 Published online by Cambridge University Press

Since we assume that it is forced that $\mathrm{Pr}(\sigma, \theta, \mu, \lambda)$ holds we have names $\dot{B}_{\xi,i}$ for $\xi < \theta^+$ and $i < \mu$ so that

$$\Vdash \dot{A}_\xi = \bigcup \{\dot{B}_{\xi,i} : i \in \mu\}.$$

Definition 1.1(C) holds. Our goal is to find a $c \in [\lambda]^\sigma$ such that $c \cap B_{\xi,i}$ are pairwise distinct for θ^+ many pairs (ξ, i). Since each of these must, by definition, be in $\dot{\mathcal{F}}_c$, we will contradict the assumption that this set has size at most θ.

To find this set of pairs (ξ, i) we will repeatedly thin out the sets we have constructed to get more and more homogeneity. To begin this process, for each $\xi < \theta^+$ and $\alpha < \lambda$ we can pick $(p_{\xi,\alpha}, i_{\xi,\alpha})$ so that $i_{\xi,\alpha} < \mu$ and $p_{\xi,\alpha} \Vdash \check{\alpha} \in B_{\xi,i_{\xi,\alpha}}$. Note that any such $p_{\xi,\alpha}$ forces in particular that $\check{\alpha} \in \dot{A}_\xi$ and therefore we have that

$$\langle \theta^+ \cdot \alpha + \xi, 1 \rangle \in p_{\xi,\alpha}. \tag{1}$$

Applying the pigeonhole principle judiciously to this situation and further thinning out we can find moreover for each $\alpha < \lambda$ a set $u_\alpha \in [\theta^+]^{\theta^+}$ so that:

- the sequence $\vec{i}_\alpha = \langle i_{\xi,\alpha} : \xi \in u_\alpha \rangle$ is bounded in μ;
- $\langle \mathrm{dom}(p_{\xi,\alpha}) : \xi \in u_\alpha \rangle$ forms a Δ-system;
- $\langle p_{\xi,\alpha} : \xi \in u_\alpha \rangle$ are pairwise compatible with common intersection r_α.

Fixing α, note that at most finitely many $\xi \in u_\alpha$ can have $\theta^+ \cdot \alpha + \xi \in \mathrm{dom}(r_\alpha)$. By throwing out these finitely many elements we can assume that

$$\forall \alpha < \lambda \forall \xi \in u_\alpha \, [\theta^+ \cdot \alpha + \xi \notin \mathrm{dom}(r_\alpha)]. \tag{2}$$

Claim 4.2. *There are $X_* \in [\lambda]^\lambda$, $u_* \in [\theta^+]^{\theta^+}$, and $\vec{i}_* \in {}^{\theta^+}\mu$ so that for every $\alpha \in X_*$ we have that $u_\alpha = u_*$ and $\vec{i}_\alpha = \vec{i}_*$.*

Proof of claim. This is another pigeonhole argument. There are $(\theta^+)^{\theta^+}$ possibilities for u_α and $(\theta^+)^{\theta^+} < \mu < \lambda$ because μ is strong limit, so there exist $X \in [\lambda]^\lambda$ and $u_* \in [\theta^+]^{\theta^+}$ such that $u_\alpha = u_*$ for all $\alpha \in X$. Now, if $\alpha \in X$, $\vec{i}_\alpha$ is bounded in μ, so we can find $v_\alpha < \mu$ such that $i_{\xi,\alpha} < v_\alpha$ for all $\xi \in u_*$. Shrinking X if necessary, we may find $v < \mu$ such that $v_\alpha = v$ for all $\alpha \in X$. But now there are $v^{\theta^+} < \mu$ many possibilities for $\vec{i}_\alpha$, so we can now find $X_* \in [X]^\lambda$ and $\vec{i}_*$ such that $\vec{i}_\alpha = \vec{i}$ for all $\alpha \in X_*$. ⊣

Next we aim for even more homogeneity. Let $S := \{\alpha \in [\mu, \lambda) : \mathrm{cf}(\alpha) = \theta^{++}\}$. For each $\delta \in S$, set $\beta_\delta := \min(X_* \setminus \delta)$.

Claim 4.3. *For each $\delta \in S$ there is a $\gamma_\delta \in [\mu, \delta)$ so that, if $\xi \in u_*$, then $p_{\xi,\beta_\delta} \restriction \delta = p_{\xi,\beta_\delta} \restriction \gamma_\delta$.*

Proof of claim. By the finiteness of the domains and the fact that δ is a limit ordinal there is always a $\gamma_{\delta,\xi} < \delta$ with the property that $p_{\xi,\beta_\delta} \restriction \delta = p_{\xi,\beta_\delta} \restriction \gamma_{\delta,\xi}$. The point is that now since δ has cofinality θ^{++} and the ξ's range over a subset of θ^+ there is a $\gamma_\delta = \sup_{\xi \in u_*} \gamma_{\delta,\xi} < \delta$. ⊣

Since $\delta \mapsto \gamma_\delta$ is a regressive function on S we can find a stationary set $S_* \subseteq S$ with the property that for all $\delta \in S_*$ we have $\gamma_\delta = \gamma_*$ for some fixed γ_*. Note that $|\gamma_*| = \mu$.

https://doi.org/10.1017/jsl.2026.10241 Published online by Cambridge University Press

Now for each $\alpha \in S_*$ consider the function $\alpha \mapsto \sup[\bigcup\{\mathrm{dom}(p_{\xi,\beta_\alpha}) : \xi \in u_*\}]$. We can find a club E bounding this function—i.e., for every $\alpha \in S_*$ we have

$$\sup\left[\bigcup\{\mathrm{dom}(p_{\xi,\beta_\alpha}) : \xi \in u_*\}\right] < \min[E \setminus (\alpha+1)]. \tag{3}$$

Let X_{**} be the set $\{\beta_\delta : \delta \in S_* \cap E\}$.

Claim 4.4. *Let $\delta, \varepsilon \in S_* \cap E$ with $\delta < \varepsilon$. If $\beta_\delta, \beta_\varepsilon$ are distinct elements of X_{**} then we have that*

$$\bigcup\{\mathrm{dom}(p_{\xi,\beta_\delta}) : \xi \in u_*\} \cap \bigcup\{\mathrm{dom}(p_{\xi,\beta_\varepsilon}) : \xi \in u_*\} \subseteq \gamma^*.$$

Proof. Fix $\xi, \eta \in u_*$. By (3), $\mathrm{dom}(p_{\xi\beta_\delta}) \subseteq \varepsilon$, hence

$$\begin{aligned}\mathrm{dom}(p_{\xi,\beta_\delta}) \cap \mathrm{dom}(p_{\eta,\beta_\varepsilon}) &= \mathrm{dom}(p_{\xi,\beta_\delta}) \cap \mathrm{dom}(p_{\eta,\beta_\varepsilon}) \cap \varepsilon \\ &= \mathrm{dom}(p_{\xi,\beta_\delta}) \cap \mathrm{dom}(p_{\xi,\beta_\varepsilon}) \cap \gamma_*, \\ &\subseteq \gamma^*,\end{aligned}$$

where the second equality is by the choice of γ_*. ⊣

Let $\langle W_n : n < \omega\rangle$ be a $\subseteq$-increasing sequence of sets of size $<\mu$ so that $\bigcup_{n<\omega} W_n = \gamma^*$. For $\delta \in E \cap S_*$ and $\xi \in u_*$ there is an $n_{\delta,\xi} \in \omega$ so that

$$\mathrm{dom}(p_{\xi,\beta_\delta}) \cap \gamma_* \subseteq W_{n_{\delta,\xi}} \tag{4}$$

simply by the finiteness of the domains. But this implies that for every $\delta \in S_* \cap E$ there is a $u_{*,\delta} \in [u_*]^{\theta^+}$ so that $n_{\delta,\xi} = n_\delta$ for some fixed $n_\delta \in \omega$ and every $\xi \in u_{*,\delta}$. By a pigeonhole argument, we can find a stationary set $S_{**} \subseteq S_* \cap E$, together with $n \in \omega$, $u_{**} \in [u_*]^{\theta^+}$ and a family $\langle p_\xi : \xi \in u_{**}\rangle$ so that, for every $\delta \in S_{**}$, the following hold:

- $n_\delta = n$;
- $u_{*,\delta} = u_{**}$;
- $\langle p_{\xi,\beta_\delta} \restriction W_n : \xi \in u_{**}\rangle = \langle p_\xi : \xi \in u_{**}\rangle$.

Reaching back to the beginning of the proof we can observe that $\langle p_\xi : \xi \in u_{**}\rangle$ forms a Δ-system by merit of being a family of restrictions of $\langle p_{\xi,\alpha} : \xi \in u_*\rangle$ which was constructed to be a Δ-system. Therefore, we can find a finite set $y \subseteq W_n$ and p^* so that for distinct $\xi_1, \xi_2 \in u_{**}$ we have $\mathrm{dom}(p_{\xi_1}) \cap \mathrm{dom}(p_{\xi_2}) = y$ and $p_{\xi_1} \restriction y = p_{\xi_2} \restriction y = p^*$. Let $c \subseteq \{\beta_\delta : \delta \in S_{**}\}$ be of size σ. Let $\dot{x}_\xi$ be a name for $\dot{B}_{\xi,i_\xi} \cap \check{c}$. By the assumption on the $\dot{B}_{\xi,i}$'s and the formulation of Definition 1.1 we must have that $\Vdash \dot{x}_\xi \in \overline{\dot{\mathcal{F}}}_{\check{c}}$ for every $\xi \in u_{**}$.

We will now contradict (B) by showing that the following claim holds.

Claim 4.5. *p^* forces that there is a set $u_\dagger \in [u_{**}]^{\theta^+}$ so that $\dot{x}_{\xi_1} \neq \dot{x}_{\xi_2}$ for every pair of distinct $\xi_1, \xi_2 \in u_\dagger$.*

Proof of claim. Let $\dot{u}_\dagger$ be the name for the set of $\xi \in u_{**}$ so that $p_\xi \in \dot{G}$. This set is forced by p^* to have size θ^+. To see this, observe that otherwise some $q \leq p^*$ forces it to have size at most θ so fix such a q and let $\delta < \theta^+$ be an ordinal so that $q \Vdash \dot{u}_\dagger \subseteq u_{**} \cap \delta$. Fix an $\xi \in u_{**}$. Since q strengthens p^*, if p_ξ and q are

https://doi.org/10.1017/jsl.2026.10241 Published online by Cambridge University Press

incompatible, there is an $\eta \notin y$ so that $p_\xi(\eta) \neq q(\eta)$ and in particular their domains intersect outside of y. Since the domains of the p_ξ's however form a Δ-system there is at most one such ξ for $\eta \in \mathrm{dom}(q) \setminus y$. As q has finite domain, there are cofinitely many $\xi \in u_{**}$ so that p_ξ and q are compatible, so in particular there is one such ξ with $\xi > \delta$. But now if $r \leq p_\xi, q$ then $r \Vdash \xi \in \dot{u}_\dagger \setminus \delta$, which is absurd.

Now, towards a contradiction suppose that $\xi_1 \neq \xi_2 \in u_{**}$ and let $p' \leq p^*$ force that $\xi_1, \xi_2 \in \dot{u}_\dagger$ but $\dot{x}_{\xi_1} = \dot{x}_{\xi_2}$. By Claim 4.4, for each $\zeta \in \mathrm{dom}(p') \setminus \gamma_*$, there exists at most one $\beta \in c$ such that $\zeta \in \mathrm{dom}(p_{\xi_1,\beta})$. Since c is infinite and $\mathrm{dom}(p')$ is finite, it follows that we can find some $\beta \in c$ such that

$$\forall \zeta \in \mathrm{dom}(p') \setminus \gamma_* [\zeta \notin \mathrm{dom}(p_{\xi_1,\beta})] \tag{5}$$

and

$$\theta^+ \cdot \beta + \xi_2 \notin \mathrm{dom}(p'). \tag{6}$$

By (1), recall that we have that $\theta^+ \cdot \beta + \xi_2 \in \mathrm{dom}(p_{\xi_2,\beta})$. On the other hand, by (2),

$$\theta^+ \cdot \beta + \xi_2 \notin \mathrm{dom}(r_\beta) = \mathrm{dom}(p_{\xi_1\beta}) \cap \mathrm{dom}(p_{\xi_2\beta}),$$

and therefore $\theta^+ \cdot \beta + \xi_2 \notin \mathrm{dom}(p_{\xi_1\beta})$. Define

$$r := p' \cup [p_{\xi_1,\beta} \restriction (\mathrm{dom}(p_{\xi_1,\beta}) \setminus \mathrm{dom}(p^*))] \cup \{\langle \theta^+ \cdot \beta + \xi_2, 0 \rangle\}.$$

We need to argue that r is a condition, the issue being the compatibility of p' and $p_{\xi_1,\beta}$. Fix $\zeta \in \mathrm{dom}(p') \cap \mathrm{dom}(p_{\xi_1,\beta})$. By (5), $\zeta < \gamma^*$, hence $\zeta \in W_n$ by (4). But $p_{\xi_1,\beta} \restriction W_n = p_{\xi_1}$ and $p' \Vdash \xi_1 \in u_\dagger$, so p' and p_{ξ_1} are compatible, therefore $p_{\xi_1,\beta}(\zeta) = p'(\zeta)$, as desired.

We have shown that r is a condition extending $p_{\xi_1,\beta}$, so $r \Vdash \beta \in B_{\xi_1,i_{\xi_1}}$ by the choice of $p_{\xi_1,\beta}$. On the other hand, $r \Vdash \beta \notin \dot{A}_{\xi_2}$ because $\langle \theta^+ \cdot \beta + \xi_2, 0 \rangle$, and therefore $r \Vdash \beta \notin \dot{B}_{\xi_2,i}$ for *any* $i < \mu$. This contradicts the assumption that p' forces $\dot{x}_{\xi_1} = \dot{x}_{\xi_2}$, thus proving the claim. ⊣

By the last claim, p^* forces that $\dot{\mathcal{F}}_{\check{c}}$ has size at least θ^+, so we have reached our final, promised, contradiction so we are done.

REMARK 4.6. In the above we really only needed that $\mathrm{cf}(\mu) < \theta^+$ to run the argument. Under this assumption, one writes $\gamma_* = \bigcup_{j<\nu} W_j$ for some regular $\nu \leq \theta$. The proof then proceeds the same, *mutatis mutandis*.

The principle $\mathsf{Pr}(\sigma, \theta, \mu, \lambda)$ is similar—though not evidently the same as that considered in [10] (see also [11] for a more complete treatment). In [10] principles of this form are shown to imply several mathematical statements involving, e.g., almost free groups. It would be interesting to know whether the same holds here.

QUESTION 4.7. *What applications does* $\mathsf{Pr}(\sigma, \theta, \mu, \lambda)$ *have to other types of structures? Does some mathematical consequence of* $\mathsf{Pr}(\sigma, \theta, \lambda, \mu)$, *for example, some statement about free groups along the lines of what is found in* [10], *serve to distinguish adding* $\aleph_\omega$ *from adding* $\aleph_{\omega+1}$ *-many Cohen reals*?

https://doi.org/10.1017/jsl.2026.10241 Published online by Cambridge University Press

More generally it would be nice to know more about the principle $\mathsf{Pr}(\sigma, \theta, \mu, \lambda)$. On this note we ask the following.

QUESTION 4.8. *What consequences more generally does* $\mathsf{Pr}(\sigma, \theta, \mu, \lambda)$ *have for the choices of* σ, θ, μ, *and* λ *considered in this article*? *For example, could it be equivalent to some partition principle or its negation*?

We note that our analysis above does not allow us to separate these principles when the final coordinate is different. Therefore we ask the following.

QUESTION 4.9. *Is it consistent that* $\mathrm{Pr}(\sigma, \theta, \mu, \lambda_0) \wedge \neg \mathrm{Pr}(\sigma, \theta, \mu, \lambda_1)$ *holds for some* $\lambda_0 < \lambda_1$?

Finally along these lines we also ask the following question.

QUESTION 4.10. *Can* Pr *exhibit any incompactness*? *For example, is it consistent that* $\mathrm{Pr}(\aleph_0, \aleph_1, \aleph_2, \aleph_n)$ *holds for every* $n > 2$ *but* $\mathrm{Pr}(\aleph_0, \aleph_1, \aleph_2, \aleph_\omega)$ *fails*?

Acknowledgments. References like [Sh:950, Th0.2=Ly5] mean that the internal label of Th0.2 is y5 in Sh:950. The reader should note that the version on S. Shelah's website is usually more up-to-date than the one in arXiv. This is publication number 1270 in Saharon Shelah's list.

Funding. Research of the second author was partially supported by the Israel Science Foundation (ISF) grant no: 2320/24 (2023-2027). The first author was supported by the Czech Academy of Sciences (RVO 67985840). The research of the third author was funded in whole or in part by the Austrian Science Fund (FWF) grant doi 10.55776/ESP548. For open access purposes, the author has applied a CC BY public copyright license to any author-accepted manuscript version arising from this submission.

REFERENCES

[1] T. BARTOSZYNSKI, *Invariants of measure and category*, ***Handbook of Set Theory***, vol. 1, 2, 3 (M. Foreman and A. Kanamori, editors), Springer, Dordrecht, 2010, pp. 491–555.

[2] A. BLASS, *Combinatorial cardinal characteristics of the continuum*, ***Handbook of Set Theory***, vol. 1 (M. Foreman and A. Kanamori, editors), Springer, Dordrecht, pp. 395–490.

[3] M. GITIK and M. GOLSHANI, *Adding a lot of Cohen reals by adding a few. I*. ***Transactions of the American Mathematical Society***, vol. 367 (2015), no. 1, pp. 209–229.

[4] ———, *Adding a lot of Cohen reals by adding a few. II*. ***Fundamenta Mathematicae***, vol. 231 (2015), no. 3, pp. 209–224.

[5] J. D. HAMKIN, *Reals added after Cohen forcing II*. MathOverflow. n.d. (Version: 2012-09-14). https://mathoverflow.net/q/107093

[6] T. JECH, ***Set Theory***, Pure and Applied Mathematics, Academic Press, Harcourt Brace Jovanovich, Publishers, New York, 1978.

[7] ———, ***Set Theory***, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2003.

[8] S. KOPPELBERG, ***Handbook of Boolean Algebras***, vol. 1, North-Holland, Amsterdam, 1989.

[9] K. KUNEN, ***Set Theory***, volume 34 of Studies in Logic (London), College Publications, London, 2011.

[10] S. SHELAH, *[Sh:108] On successors of singular cardinals*, ***Logic Colloquium '78 (Mons, 1978)***, volume 97 of Studies in Logic and the Foundations of Mathematics (M. Boffa, D. van Dalen and K. McAloon, editors), North-Holland, Amsterdam, 1979, pp. 357–380. MR567680.

https://doi.org/10.1017/jsl.2026.10241 Published online by Cambridge University Press

[11] ———, *[Sh:88a] Appendix*, 1985, pp. 483–495. Appendix of [Sh:88].

[12] J. Zapletal, ***Forcing Idealized***, volume 174 of Cambridge Tracts in Mathematics, Cambridge University Press, Cambridge, 2008.

INSTITUTE OF MATHEMATICS
CZECH ACADEMY OF SCIENCES
ŽITNÁ 25, PRAGUE 1, 115 67, CZECH REPUBLIC
E-mail: marun@math.cas.cz

EINSTEIN INSTITUTE OF MATHEMATICS
THE HEBREW UNIVERSITY OF JERUSALEM
9190401, JERUSALEM, ISRAEL
E-mail: shelah@math.huji.ac.il

and

DEPARTMENT OF MATHEMATICS
RUTGERS UNIVERSITY,
PISCATAWAY, NJ 08854-8019, USA

KURT GÖDEL RESEARCH CENTER, FACULTY OF MATHEMATICS
UNIVERSITY OF VIENNA
KOLINGASSE 14 – 16, 1090 WIEN, AUSTRIA
E-mail: corey.bacal.switzer@univie.ac.at

https://doi.org/10.1017/jsl.2026.10241 Published online by Cambridge University Press