References

[1] M. L. Mangano and S. J. Parke, "Multiparton amplitudes in gauge theories," *Phys. Rept.* **200**, 301 (1991) [hep-th/0509223].

[2] M. Srednicki, *Quantum Field Theory*, Cambridge, UK: Cambridge University Press (2007).

[3] L. J. Dixon, "Calculating scattering amplitudes efficiently," In *Boulder 1995, QCD and beyond*, 539–582 [hep-ph/9601359].

[4] L. J. Dixon, "Scattering amplitudes: the most perfect microscopic structures in the universe," *J. Phys.* A **44**, 454001 (2011) [arXiv:1105.0771 [hep-th]].

[5] S. J. Parke and T. R. Taylor, "An amplitude for *n* gluon scattering," *Phys. Rev. Lett.* **56**, 2459 (1986).

[6] K. Kampf, J. Novotny, and J. Trnka, "Tree-level amplitudes in the nonlinear sigma model," *JHEP* **1305**, 032 (2013) [arXiv:1304.3048 [hep-th]].

[7] R. Kleiss and H. Kuijf, "Multi-gluon cross-sections and five jet production at hadron colliders," *Nucl. Phys. B* **312**, 616 (1989).

[8] V. Del Duca, L. J. Dixon, and F. Maltoni, "New color decompositions for gauge amplitudes at tree and loop level," *Nucl. Phys. B* **571**, 51 (2000) [hep-ph/9910563].

[9] Z. Bern, J. J. M. Carrasco, and H. Johansson, "New relations for gauge-theory amplitudes," *Phys. Rev. D* **78**, 085011 (2008) [arXiv:0805.3993 [hep-ph]].

[10] C. F. Berger, V. Del Duca, and L. J. Dixon, "Recursive construction of Higgs-plus-multiparton loop amplitudes: The last of the phi-nite loop amplitudes," *Phys. Rev. D* **74**, 094021 (2006) [Erratum-*ibid.* D **76**, 099901 (2007)] [hep-ph/0608180];
S. D. Badger, E. W. N. Glover, and K. Risager, "One-loop phi-MHV amplitudes using the unitarity bootstrap," *JHEP* **0707**, 066 (2007) [arXiv:0704.3914 [hep-ph]];
L. J. Dixon and Y. Sofianatos, "Analytic one-loop amplitudes for a Higgs boson plus four partons," *JHEP* **0908**, 058 (2009) [arXiv:0906.0008 [hep-ph]];
S. Badger, E. W. Nigel Glover, P. Mastrolia, and C. Williams, "One-loop Higgs plus four gluon amplitudes: Full analytic results," *JHEP* **1001**, 036 (2010) [arXiv:0909.4475 [hep-ph]].

[11] L. J. Dixon, "A brief introduction to modern amplitude methods," [arXiv:1310.5353 [hep-ph]].

[12] F. A. Berends and W. T. Giele, "Recursive calculations for processes with n gluons," *Nucl. Phys. B* **306**, 759 (1988).
F. A. Berends, W. T. Giele, and H. Kuijf, "Exact and approximate expressions for multi-gluon scattering," *Nucl. Phys. B* **333**, 120 (1990).

[13] R. Britto, F. Cachazo, and B. Feng, "New recursion relations for tree amplitudes of gluons," *Nucl. Phys. B* **715**, 499 (2005) [hep-th/0412308].

Downloaded from https://www.cambridge.org/core. UNIVERSIDAD AUTONOMA METROPOLITANA, on 14 Nov 2026 at 00:33:59, subject to the Cambridge Core terms of use, available at https://www.cambridge.org/core/terms. https://doi.org/10.1017/CBO9781107706620.020

[14] R. Britto, F. Cachazo, B. Feng, and E. Witten, "Direct proof of tree-level recursion relation in Yang–Mills theory," *Phys. Rev. Lett.* **94**, 181602 (2005) [hep-th/0501052].
[15] F. Cachazo, P. Svrcek, and E. Witten, "MHV vertices and tree amplitudes in gauge theory," *JHEP* **0409**, 006 (2004) [hep-th/0403047].
[16] B. Feng, J. Wang, Y. Wang, and Z. Zhang, "BCFW recursion relation with nonzero boundary contribution," *JHEP* **1001**, 019 (2010) [arXiv:0911.0301 [hep-th]].
[17] E. Conde and S. Rajabi, "The twelve-graviton next-to-MHV amplitude from Risager's construction," *JHEP* **1209**, 120 (2012) [arXiv:1205.3500 [hep-th]].
[18] N. Arkani-Hamed and J. Kaplan, "On tree amplitudes in gauge theory and gravity," *JHEP* **0804**, 076 (2008) [arXiv:0801.2385 [hep-th]].
[19] C. Cheung, "On-shell recursion relations for generic theories," *JHEP* **1003**, 098 (2010) [arXiv:0808.0504 [hep-th]].
[20] P. Benincasa, C. Boucher-Veronneau, and F. Cachazo, "Taming tree amplitudes in general relativity," *JHEP* **0711**, 057 (2007) [hep-th/0702032 [HEP-TH]].
[21] H. Kawai, D. C. Lewellen, and S. H. H. Tye, "A relation between tree amplitudes of closed and open strings," *Nucl. Phys. B* **269**, 1 (1986).
[22] S. Sannan, "Gravity as the limit of the type II superstring theory," *Phys. Rev. D* **34**, 1749 (1986).
[23] N. Arkani-Hamed, F. Cachazo, C. Cheung, and J. Kaplan, "A duality for the S matrix," *JHEP* **1003**, 020 (2010) [arXiv:0907.5418 [hep-th]].
[24] A. Hodges, "Eliminating spurious poles from gauge-theoretic amplitudes," *JHEP* **1305**, 135 (2013) [arXiv:0905.1473 [hep-th]].
[25] M. Spradlin, A. Volovich, and C. Wen, "Three applications of a bonus relation for gravity amplitudes," *Phys. Lett. B* **674**, 69 (2009) [arXiv:0812.4767 [hep-th]].
[26] T. Cohen, H. Elvang, and M. Kiermaier, "On-shell constructibility of tree amplitudes in general field theories," *JHEP* **1104**, 053 (2011) [arXiv:1010.0257 [hep-th]].
[27] H. Elvang, D. Z. Freedman, and M. Kiermaier, "Recursion relations, generating functions, and unitarity sums in N = 4 SYM theory," *JHEP* **0904**, 009 (2009) [arXiv:0808.1720 [hep-th]].
[28] K. Risager, "A direct proof of the CSW rules," *JHEP* **0512**, 003 (2005) [hep-th/0508206].
[29] H. Elvang, D. Z. Freedman, and M. Kiermaier, "Proof of the MHV vertex expansion for all tree amplitudes in N = 4 SYM theory," *JHEP* **0906**, 068 (2009) [arXiv:0811.3624 [hep-th]].
[30] L. J. Dixon, E. W. N. Glover, and V. V. Khoze, "MHV rules for Higgs plus multi-gluon amplitudes," *JHEP* **0412**, 015 (2004) [hep-th/0411092].
[31] S. D. Badger, E. W. N. Glover, and V. V. Khoze, "MHV rules for Higgs plus multi-parton amplitudes," *JHEP* **0503**, 023 (2005) [hep-th/0412275].
[32] A. Brandhuber, B. Spence, and G. Travaglini, "Tree-level formalism," *J. Phys. A* **44**, 454002 (2011) [arXiv:1103.3477 [hep-th]].
[33] A. Gorsky and A. Rosly, "From Yang–Mills Lagrangian to MHV diagrams," *JHEP* **0601**, 101 (2006) [hep-th/0510111];
P. Mansfield, "The Lagrangian origin of MHV rules," *JHEP* **0603**, 037 (2006) [hep-th/0511264];

Downloaded from https://www.cambridge.org/core. UNIVERSIDAD AUTONOMA METROPOLITANA, on 14 Nov 2026 at 00:33:59, subject to the Cambridge Core terms of use, available at https://www.cambridge.org/core/terms. https://doi.org/10.1017/CBO9781107706620.020

J. H. Ettle and T. R. Morris, "Structure of the MHV-rules Lagrangian," *JHEP* **0608**, 003 (2006) [hep-th/0605121];
J. H. Ettle, C.-H. Fu, J. P. Fudger, P. R. W. Mansfield, and T. R. Morris, "S-matrix equivalence theorem evasion and dimensional regularisation with the canonical MHV Lagrangian," *JHEP* **0705**, 011 (2007) [hep-th/0703286];
H. Feng and Y.-t. Huang, "MHV Lagrangian for N = 4 super Yang–Mills," *JHEP* **0904**, 047 (2009) [hep-th/0611164].

[34] R. Boels, L. J. Mason, and D. Skinner, "From twistor actions to MHV diagrams," *Phys. Lett. B* **648**, 90 (2007) [hep-th/0702035].

[35] N. E. J. Bjerrum-Bohr, D. C. Dunbar, H. Ita, W. B. Perkins, and K. Risager, "MHV-vertices for gravity amplitudes," *JHEP* **0601**, 009 (2006) [hep-th/0509016].

[36] M. Bianchi, H. Elvang, and D. Z. Freedman, "Generating tree amplitudes in N = 4 SYM and N = 8 SG," *JHEP* **0809**, 063 (2008) [arXiv:0805.0757 [hep-th]].

[37] J. Wess and J. Bagger, *Supersymmetry and Supergravity*, Princeton, USA: University Press (1992).

[38] M. T. Grisaru and H. N. Pendleton, "Some properties of scattering amplitudes in supersymmetric theories," *Nucl. Phys. B* **124**, 81 (1977);
M. T. Grisaru, H. N. Pendleton, and P. van Nieuwenhuizen, "Supergravity and the S matrix," *Phys. Rev. D* **15**, 996 (1977).

[39] D. Z. Freedman and A. Van Proeyen, Supergravity, Cambridge, UK: Cambridge University Press (2012).

[40] L. Brink, J. H. Schwarz, and J. Scherk, "Supersymmetric Yang–Mills theories," *Nucl. Phys. B* **121**, 77 (1977).

[41] A. Ferber, "Supertwistors and conformal supersymmetry," *Nucl. Phys. B* **132**, 55 (1978).

[42] H. Elvang, D. Z. Freedman, and M. Kiermaier, "Solution to the Ward identities for superamplitudes," *JHEP* **1010**, 103 (2010) [arXiv:0911.3169 [hep-th]].

[43] H. Elvang, D. Z. Freedman, and M. Kiermaier, "SUSY Ward identities, superamplitudes, and counterterms," *J. Phys. A* **44**, 454009 (2011) [arXiv:1012.3401 [hep-th]].

[44] M. Kiermaier and S. G. Naculich, "A super MHV vertex expansion for N = 4 SYM theory," *JHEP* **0905**, 072 (2009) [arXiv:0903.0377 [hep-th]].

[45] N. Arkani-Hamed, "What is the simplest QFT?," talk given at the Paris Workshop *Wonders of Gauge Theory and Supergravity*, 24 June 2008.

[46] A. Brandhuber, P. Heslop, and G. Travaglini, "A note on dual superconformal symmetry of the N = 4 super Yang–Mills S-matrix," *Phys. Rev. D* **78**, 125005 (2008) [arXiv:0807.4097 [hep-th]].

[47] N. Arkani-Hamed, F. Cachazo, and J. Kaplan, "What is the simplest quantum field theory?," *JHEP* **1009**, 016 (2010) [arXiv:0808.1446 [hep-th]].

[48] Z. Bern, J. J. M. Carrasco, H. Ita, H. Johansson, and R. Roiban, "On the structure of supersymmetric sums in multi-loop unitarity cuts," *Phys. Rev. D* **80**, 065029 (2009) [arXiv:0903.5348 [hep-th]].

[49] J. M. Drummond and J. M. Henn, "All tree-level amplitudes in N = 4 SYM," *JHEP* **0904**, 018 (2009) [arXiv:0808.2475 [hep-th]].

Downloaded from https://www.cambridge.org/core. UNIVERSIDAD AUTONOMA METROPOLITANA, on 14 Nov 2026 at 00:33:59, subject to the Cambridge Core terms of use, available at https://www.cambridge.org/core/terms. https://doi.org/10.1017/CBO9781107706620.020

[50] J. M. Drummond, J. Henn, G. P. Korchemsky, and E. Sokatchev, "Dual superconformal symmetry of scattering amplitudes in N = 4 super-Yang–Mills theory," *Nucl. Phys. B* **828**, 317 (2010) [arXiv:0807.1095 [hep-th]].
[51] J. M. Drummond, M. Spradlin, A. Volovich, and C. Wen, "Tree-level amplitudes in N = 8 supergravity," *Phys. Rev. D* **79**, 105018 (2009) [arXiv:0901.2363 [hep-th]].
[52] J. M. Drummond, "Hidden simplicity of gauge theory amplitudes," *Class. Quant. Grav.* **27**, 214001 (2010) [arXiv:1010.2418 [hep-th]].
[53] R. Penrose, "Twistor algebra," *J. Math. Phys.* **8**, 345 (1967).
[54] E. Witten, "Perturbative gauge theory as a string theory in twistor space," *Commun. Math. Phys.* **252**, 189 (2004) [hep-th/0312171].
[55] P. A. M. Dirac, "Wave equations in conformal space," *Annals Math.* **37**, 429 (1936).
[56] W. Siegel, "Embedding versus 6D twistors," [arXiv:1204.5679 [hep-th]].
[57] W. Siegel, "Fields," [hep-th/9912205].
[58] J. M. Drummond, J. Henn, V. A. Smirnov, and E. Sokatchev, "Magic identities for conformal four-point integrals," *JHEP* **0701**, 064 (2007) [hep-th/0607160].
[59] J. M. Drummond, J. M. Henn, and J. Plefka, "Yangian symmetry of scattering amplitudes in N = 4 super Yang–Mills theory," *JHEP* **0905**, 046 (2009) [arXiv:0902.2987 [hep-th]].
[60] L. J. Mason and D. Skinner, "Dual superconformal invariance, momentum twistors and grassmannians," *JHEP* **0911**, 045 (2009) [arXiv:0909.0250 [hep-th]].
N. Arkani-Hamed, F. Cachazo, and C. Cheung, "The Grassmannian origin of dual superconformal invariance," *JHEP* **10033**, 036 (2010) [arXiv: 0909.0483 [hep-th]].
H. Elvang, Y.-t. Huang, C. Keeler, *et al.*, "Grassmannians for scattering amplitudes in 4d N =4 SYM and 3d ABJM," 2014 [arXiv: 1410.0621 [hep.th]].
[61] R. Roiban, "Review of AdS/CFT integrability, Chapter V.1: Scattering amplitudes – a brief introduction," *Lett. Math. Phys.* **99**, 455 (2012) [arXiv:1012.4001 [hep-th]].
[62] Lorenzo Magnea, *Lecture notes on Perturbative QCD* at the National School of Theoretical Physics of the University of Parma (2008). http://personalpages.to.infn.it/∼magnea/QCD.pdf
[63] L. V. Bork, D. I. Kazakov, G. S. Vartanov, and A. V. Zhiboedov, "Construction of infrared finite observables in N = 4 super Yang–Mills theory," *Phys. Rev. D* **81**, 105028 (2010) [arXiv:0911.1617 [hep-th]].
[64] Z. Bern, G. Chalmers, L. J. Dixon, and D. A. Kosower, "One loop N gluon amplitudes with maximal helicity violation via collinear limits," *Phys. Rev. Lett.* **72**, 2134 (1994) [hep-ph/9312333].
[65] G. Mahlon, "Multi-gluon helicity amplitudes involving a quark loop," *Phys. Rev. D* **49**, 4438 (1994) [hep-ph/9312276].
[66] Z. Bern, L. J. Dixon, and D. A. Kosower, "Dimensionally regulated pentagon integrals," *Nucl. Phys. B* **412**, 751 (1994) [hep-ph/9306240].
[67] Z. Bern, L. J. Dixon, D. C. Dunbar, and D. A. Kosower, "One loop n point gauge theory amplitudes, unitarity and collinear limits," *Nucl. Phys. B* **425**, 217 (1994) [hep-ph/9403226];
Z. Bern, L. J. Dixon, D. C. Dunbar, and D. A. Kosower, "Fusing gauge theory tree amplitudes into loop amplitudes," *Nucl. Phys. B* **435**, 59 (1995) [hep-ph/9409265].

Downloaded from https://www.cambridge.org/core. UNIVERSIDAD AUTONOMA METROPOLITANA, on 14 Nov 2026 at 00:33:59, subject to the Cambridge Core terms of use, available at https://www.cambridge.org/core/terms. https://doi.org/10.1017/CBO9781107706620.020

[68] J. J. M. Carrasco and H. Johansson, "Generic multiloop methods and application to N = 4 super-Yang–Mills," *J. Phys. A* **44**, 454004 (2011) [arXiv:1103.3298 [hep-th]].
[69] Z. Bern and Y.-t. Huang, "Basics of generalized unitarity," *J. Phys.* A **44**, 454003 (2011) [arXiv:1103.1869 [hep-th]].
[70] H. Ita, "Susy theories and QCD: Numerical approaches," *J. Phys. A* **44**, 454005 (2011) [arXiv:1109.6527 [hep-th]].
[71] R. Britto, "Loop amplitudes in gauge theories: Modern analytic approaches," *J. Phys. A* **44**, 454006 (2011) [arXiv:1012.4493 [hep-th]].
[72] W. L. van Neerven and J. A. M. Vermaseren, "Large loop integrals," *Phys. Lett. B* **137**, 241 (1984).
[73] Z. Bern, L. J. Dixon, and D. A. Kosower, "Dimensionally regulated one loop integrals," *Phys. Lett. B* **302**, 299 (1993) [Erratum-*ibid. B* **318**, 649 (1993)] [hep-ph/9212308].
[74] Z. Bern, L. J. Dixon, and D. A. Kosower, "Dimensionally regulated pentagon integrals," *Nucl. Phys. B* **412**, 751 (1994) [hep-ph/9306240].
[75] L. M. Brown and R. P. Feynman, "Radiative corrections to Compton scattering," *Phys. Rev.* **85**, 231 (1952);
G. 't Hooft and M. J. G. Veltman, "Scalar one loop integrals," *Nucl. Phys. B* **153**, 365 (1979).
[76] G. Passarino and M. J. G. Veltman, "One loop corrections for e+ e− annihilation into mu+ mu− in the Weinberg model," *Nucl. Phys. B* **160**, 151 (1979).
[77] R. K. Ellis, Z. Kunszt, K. Melnikov, and G. Zanderighi, "One-loop calculations in quantum field theory: from Feynman diagrams to unitarity cuts," *Phys. Rept.* **518**, 141 (2012) [arXiv:1105.4319 [hep-ph]].
[78] H. Johansson, D. A. Kosower, and K. J. Larsen, "An overview of maximal unitarity at two loops," *PoS LL* **2012**, 066 (2012) [arXiv:1212.2132 [hep-th]].
[79] C. Anastasiou, R. Britto, B. Feng, Z. Kunszt, and P. Mastrolia, "D-dimensional unitarity cut method," *Phys. Lett. B* **645**, 213 (2007) [hep-ph/0609191];
R. Britto and B. Feng, "Integral coefficients for one-loop amplitudes," *JHEP* **0802**, 095 (2008) [0711.4284 [hep-ph]];
R. Britto and B. Feng, "Unitarity cuts with massive propagators and algebraic expressions for coefficients," *Phys. Rev. D* **75**, 105006 (2007) [hep-ph/0612089];
G. Ossola, C. G. Papadopoulos, and R. Pittau, "Reducing full one-loop amplitudes to scalar integrals at the integrand level," *Nucl. Phys. B* **763**, 147 (2007) [hep-ph/0609007];
R. Britto, B. Feng, and P. Mastrolia, "Closed-form decomposition of one-loop massive amplitudes," *Phys. Rev. D* **78**, 025031 (2008) [0803.1989 [hep-ph]];
D. Forde, "Direct extraction of one-loop integral coefficients," *Phys. Rev. D* **75**, 125019 (2007) [0704.1835 [hep-ph]];
[80] Z. Bern and A. G. Morgan, "Massive loop amplitudes from unitarity," *Nucl. Phys. B* **467**, 479 (1996) [hep-ph/9511336];
Z. Bern, L. J. Dixon, and D. A. Kosower, "Progress in one-loop QCD computations," *Ann. Rev. Nucl. Part. Sci.* **46**, 109 (1996) [hep-ph/9602280].

Downloaded from https://www.cambridge.org/core. UNIVERSIDAD AUTONOMA METROPOLITANA, on 14 Nov 2026 at 00:33:59, subject to the Cambridge Core terms of use, available at https://www.cambridge.org/core/terms. https://doi.org/10.1017/CBO9781107706620.020

[81] S. D. Badger, "Direct extraction of one loop rational terms," *JHEP* **0901**, 049 (2009) [0806.4600 [hep-ph]].
[82] M. B. Green, J. H. Schwarz, and L. Brink, "N = 4 Yang–Mills and N = 8 supergravity as limits of string theories," *Nucl. Phys. B* **198**, 474 (1982).
[83] Z. Bern, N. E. J. Bjerrum-Bohr, and D. C. Dunbar, "Inherited twistor-space structure of gravity loop amplitudes," *JHEP* **0505**, 056 (2005) [hep-th/0501137];
N. E. J. Bjerrum-Bohr, D. C. Dunbar, H. Ita, W. B. Perkins, and K. Risager, "The no-triangle hypothesis for N = 8 supergravity," *JHEP* **0612**, 072 (2006) [hep-th/0610043];
N. E. J. Bjerrum-Bohr and P. Vanhove, "Absence of triangles in maximal supergravity amplitudes," *JHEP* **0810**, 006 (2008) [arXiv:0805.3682 [hep-th]].
[84] Z. Bern, J. J. Carrasco, D. Forde, H. Ita, and H. Johansson, "Unexpected cancellations in gravity theories," *Phys. Rev. D* **77**, 025010 (2008) [arXiv:0707.1035 [hep-th]].
[85] S. Lal and S. Raju, "The next-to-simplest quantum field theories," *Phys. Rev. D* **81**, 105002 (2010) [arXiv:0910.0930 [hep-th]].
[86] D. C. Dunbar, J. H. Ettle, and W. B. Perkins, "Perturbative expansion of $N < 8$ supergravity," *Phys. Rev. D* **83**, 065015 (2011) [arXiv:1011.5378 [hep-th]].
[87] H. Elvang, Y.-t. Huang, and C. Peng, "On-shell superamplitudes in $N < 4$ SYM," *JHEP* **1109**, 031 (2011) [arXiv:1102.4843 [hep-th]].
[88] Y.-t. Huang, D. A. McGady, and C. Peng, "One-loop renormalization and the S-matrix," [arXiv:1205.5606 [hep-th]].
[89] N. Marcus, "Composite anomalies in supergravity," *Phys. Lett. B* **157**, 383 (1985).
[90] P. di Vecchia, S. Ferrara, and L. Girardello, "Anomalies of hidden local chiral symmetries in sigma models and extended supergravities," *Phys. Lett. B* **151**, 199 (1985).
[91] J. M. Drummond, J. Henn, G. P. Korchemsky, and E. Sokatchev, "Generalized unitarity for N = 4 super-amplitudes," *Nucl. Phys. B* **869**, 452 (2013) [arXiv:0808.0491 [hep-th]].
[92] A. Brandhuber, P. Heslop, and G. Travaglini, "One-loop amplitudes in N = 4 super Yang–Mills and anomalous dual conformal symmetry," *JHEP* **0908**, 095 (2009) [arXiv:0905.4377 [hep-th]].
[93] H. Elvang, D. Z. Freedman, and M. Kiermaier, "Dual conformal symmetry of 1-loop NMHV amplitudes in N = 4 SYM theory," *JHEP* **1003**, 075 (2010) [arXiv:0905.4379 [hep-th]].
[94] G. P. Korchemsky and E. Sokatchev, "Symmetries and analytic properties of scattering amplitudes in N = 4 SYM theory," *Nucl. Phys. B* **832**, 1 (2010) [arXiv:0906.1737 [hep-th]].
[95] J. Gluza, K. Kajda, and D. A. Kosower, "Towards a basis for planar two-loop integrals," *Phys. Rev. D* **83**, 045012 (2011) [arXiv:1009.0472 [hep-th]].
[96] D. A. Kosower and K. J. Larsen, "Maximal unitarity at two loops," *Phys. Rev. D* **85**, 045017 (2012) [arXiv:1108.1180 [hep-th]];
H. Johansson, D. A. Kosower, and K. J. Larsen, "Two-loop maximal unitarity with external masses," *Phys. Rev. D* **87**, 025030 (2013) [arXiv:1208.1754 [hep-th]].
[97] S. Badger, H. Frellesvig, and Y. Zhang, "Hepta-cuts of two-loop scattering amplitudes," *JHEP* **1204**, 055 (2012) [arXiv:1202.2019 [hep-ph]].

Downloaded from https://www.cambridge.org/core. UNIVERSIDAD AUTONOMA METROPOLITANA, on 14 Nov 2026 at 00:33:59, subject to the Cambridge Core terms of use, available at https://www.cambridge.org/core/terms. https://doi.org/10.1017/CBO9781107706620.020

[98] Y. Zhang, "Integrand-level reduction of loop amplitudes by computational algebraic geometry methods," *JHEP* **1209**, 042 (2012) [arXiv:1205.5707 [hep-ph]].
[99] M. Sgaard, "Global residues and two-loop hepta-cuts," *JHEP* **1309**, 116 (2013) [arXiv:1306.1496 [hep-th]].
[100] A. V. Smirnov and A. V. Petukhov, "The number of master integrals is finite," *Lett. Math. Phys.* **97**, 37 (2011) [arXiv:1004.4199 [hep-th]].
[101] C. Anastasiou, Z. Bern, L. J. Dixon, and D. A. Kosower, "Planar amplitudes in maximally supersymmetric Yang–Mills theory," *Phys. Rev. Lett.* **91**, 251602 (2003) [hep-th/0309040].
[102] Z. Bern, L. J. Dixon, and V. A. Smirnov, "Iteration of planar amplitudes in maximally supersymmetric Yang–Mills theory at three loops and beyond," *Phys. Rev. D* **72**, 085001 (2005) [hep-th/0505205].
[103] Z. Bern, J. S. Rozowsky, and B. Yan, "Two loop four gluon amplitudes in N = 4 superYang–Mills," *Phys. Lett. B* **401**, 273 (1997) [hep-ph/9702424].
[104] Z. Bern, M. Czakon, D. A. Kosower, R. Roiban, and V. A. Smirnov, "Two-loop iteration of five-point N = 4 super-Yang–Mills amplitudes," *Phys. Rev. Lett.* **97**, 181601 (2006) [hep-th/0604074].
[105] F. Cachazo, M. Spradlin, and A. Volovich, "Iterative structure within the five-particle two-loop amplitude," *Phys. Rev. D* **74**, 045020 (2006) [hep-th/0602228].
[106] L. F. Alday and J. Maldacena, "Comments on gluon scattering amplitudes via AdS/CFT," *JHEP* **0711**, 068 (2007) [arXiv:0710.1060 [hep-th]].
[107] L. F. Alday and J. M. Maldacena, "Gluon scattering amplitudes at strong coupling," *JHEP* **0706**, 064 (2007) [arXiv:0705.0303 [hep-th]].
[108] Z. Bern, L. J. Dixon, D. A. Kosower, *et al.*, "The two-loop six-gluon MHV amplitude in maximally supersymmetric Yang–Mills theory," *Phys. Rev. D* **78** (2008) 045007 [arXiv:0803.1465 [hep-th]].
[109] F. Cachazo, M. Spradlin, and A. Volovich, "Leading singularities of the two-loop six-particle MHV amplitude," *Phys. Rev. D* **78**, 105022 (2008) [arXiv:0805.4832 [hep-th]].
[110] V. Del Duca, C. Duhr, and V. A. Smirnov, "An analytic result for the two-loop hexagon Wilson loop in N = 4 SYM," *JHEP* **1003**, 099 (2010) [arXiv:0911.5332 [hep-ph]].
[111] V. Del Duca, C. Duhr, and V. A. Smirnov, "The two-loop hexagon Wilson loop in N = 4 SYM," *JHEP* **1005**, 084 (2010) [arXiv:1003.1702 [hep-th]].
[112] A. B. Goncharov, M. Spradlin, C. Vergu, and A. Volovich, "Classical polylogarithms for amplitudes and Wilson loops," *Phys. Rev. Lett.* **105**, 151605 (2010) [arXiv:1006.5703 [hep-th]].
[113] E. W. Nigel Glover and C. Williams, "One-loop gluonic amplitudes from single unitarity cuts," *JHEP* **0812**, 067 (2008) [0810.2964 [hep-th]];
I. Bierenbaum, S. Catani, P. Draggiotis, and G. Rodrigo, "A tree-loop duality relation at two loops and beyond," *JHEP* **1010**, 073 (2010) [arXiv:1007.0194 [hep-ph]];
H. Elvang, D. Z. Freedman, and M. Kiermaier, "Integrands for QCD rational terms and N = 4 SYM from massive CSW rules," *JHEP* **1206**, 015 (2012) [arXiv:1111.0635 [hep-th]].
[114] S. Caron-Huot, "Loops and trees," *JHEP* **1105**, 080 (2011) [arXiv:1007.3224 [hep-ph]].

Downloaded from https://www.cambridge.org/core. UNIVERSIDAD AUTONOMA METROPOLITANA, on 14 Nov 2026 at 00:33:59, subject to the Cambridge Core terms of use, available at https://www.cambridge.org/core/terms. https://doi.org/10.1017/CBO9781107706620.020

[115] N. Arkani-Hamed, J. L. Bourjaily, F. Cachazo, S. Caron-Huot, and J. Trnka, "The all-loop integrand for scattering amplitudes in planar N = 4 SYM," *JHEP* **1101**, 041 (2011) [arXiv:1008.2958 [hep-th]].

[116] R. H. Boels, "On BCFW shifts of integrands and integrals," *JHEP* **1011**, 113 (2010) [arXiv:1008.3101 [hep-th]].

[117] Z. Bern, J. J. M. Carrasco, H. Johansson, and D. A. Kosower, "Maximally supersymmetric planar Yang–Mills amplitudes at five loops," *Phys. Rev. D* **76**, 125020 (2007) [0705.1864 [hep-th]].

[118] R. Britto, F. Cachazo, and B. Feng, "Generalized unitarity and one-loop amplitudes in N = 4 super-Yang–Mills," *Nucl. Phys. B* **725**, 275 (2005) [hep-th/0412103];
E. I. Buchbinder and F. Cachazo, "Two-loop amplitudes of gluons and octa-cuts in N = 4 super Yang–Mills," *JHEP* **0511**, 036 (2005) [hep-th/0506126].

[119] N. Arkani-Hamed, J. L. Bourjaily, F. Cachazo, and J. Trnka, "Local integrals for planar scattering amplitudes," *JHEP* **1206**, 125 (2012) [arXiv:1012.6032 [hep-th]].
For extensions, see J. L. Bourjaily, S. Caron-Huot, and J. Trnka, "Dual-conformal regularization of infrared loop divergences and the chiral box expansion," [arXiv:1303.4734 [hep-th]].

[120] N. Arkani-Hamed, J. L. Bourjaily, F. Cachazo, *et al.*, "Scattering amplitudes and the positive Grassmannian," [arXiv:1212.5605 [hep-th]].

[121] F. Cachazo, "Sharpening the leading singularity," [arXiv:0803.1988 [hep-th]].

[122] J. M. Drummond and L. Ferro, "Yangians, Grassmannians and T-duality," *JHEP* **1007**, 027 (2010) [arXiv:1001.3348 [hep-th]].

[123] J. M. Drummond and L. Ferro, "The Yangian origin of the Grassmannian integral," *JHEP* **1012**, 010 (2010) [arXiv:1002.4622 [hep-th]];
G. P. Korchemsky and E. Sokatchev, "Superconformal invariants for scattering amplitudes in N = 4 SYM theory," *Nucl. Phys. B* **839**, 377 (2010) [arXiv:1002.4625 [hep-th]].

[124] R. Roiban, M. Spradlin, and A. Volovich, "On the tree level S matrix of Yang–Mills theory," *Phys. Rev. D* **70**, 026009 (2004) [hep-th/0403190].

[125] M. Spradlin and A. Volovich, "From twistor string theory to recursion relations," *Phys. Rev. D* **80**, 085022 (2009) [arXiv:0909.0229 [hep-th]].

[126] N. Arkani-Hamed, J. Bourjaily, F. Cachazo, and J. Trnka, "Unification of residues and Grassmannian dualities," *JHEP* **1101**, 049 (2011) [arXiv:0912.4912 [hep-th]].

[127] J. L. Bourjaily, J. Trnka, A. Volovich, and C. Wen, "The Grassmannian and the twistor string: Connecting all trees in N = 4 SYM," *JHEP* **1101**, 038 (2011) [arXiv:1006.1899 [hep-th]].

[128] M. Bullimore, L. J. Mason, and D. Skinner, "Twistor-strings, Grassmannians and leading singularities," *JHEP* **1003**, 070 (2010) [arXiv:0912.0539 [hep-th]].

[129] L. Dolan and P. Goddard, "Complete equivalence between gluon tree amplitudes in twistor string theory and in gauge theory," *JHEP* **1206**, 030 (2012) [arXiv:1111.0950 [hep-th]].

[130] N. Arkani-Hamed, J. L. Bourjaily, F. Cachazo, A. Hodges, and J. Trnka, "A note on polytopes for scattering amplitudes," *JHEP* **1204**, 081 (2012) [arXiv:1012.6030 [hep-th]].

Downloaded from https://www.cambridge.org/core. UNIVERSIDAD AUTONOMA METROPOLITANA, on 14 Nov 2026 at 00:33:59, subject to the Cambridge Core terms of use, available at https://www.cambridge.org/core/terms. https://doi.org/10.1017/CBO9781107706620.020

[131] M. Bullimore, L. J. Mason, and D. Skinner, "MHV diagrams in momentum twistor space," *JHEP* **1012**, 032 (2010) [arXiv:1009.1854 [hep-th]].
[132] N. Arkani-Hamed and J. Trnka, "The amplituhedron," [arXiv:1312.2007 [hep-th]]; N. Arkani-Hamed and J. Trnka, "Into the amplituhedron," [arXiv:1312.7878 [hep-th]].
[133] L. Mason and D. Skinner, "Amplitudes at weak coupling as polytopes in AdS_5," *J. Phys. A* **44**, 135401 (2011) [arXiv:1004.3498 [hep-th]].
[134] H. Nastase and H. J. Schnitzer, "Twistor and polytope interpretations for subleading color one-loop amplitudes," *Nucl. Phys. B* **855**, 901 (2012) [arXiv:1104.2752 [hep-th]].
[135] R. Boels, "Covariant representation theory of the Poincaré algebra and some of its extensions," *JHEP* **1001**, 010 (2010) [arXiv:0908.0738 [hep-th]].
[136] S. Caron-Huot and D. O'Connell, "Spinor helicity and dual conformal symmetry in ten dimensions," *JHEP* **1108**, 014 (2011) [arXiv:1010.5487 [hep-th]].
[137] R. H. Boels and D. O'Connell, "Simple superamplitudes in higher dimensions," *JHEP* **1206**, 163 (2012) [arXiv:1201.2653 [hep-th]].
[138] S. Davies, "One-loop QCD and Higgs to partons processes using six-dimensional helicity and generalized unitarity," *Phys. Rev. D* **84**, 094016 (2011) [arXiv:1108.0398 [hep-ph]].
[139] Z. Bern, J. J. Carrasco, T. Dennen, Y.-t. Huang, and H. Ita, "Generalized unitarity and six-dimensional helicity," *Phys. Rev. D* **83**, 085022 (2011) [arXiv:1010.0494 [hep-th]].
[140] N. Craig, H. Elvang, M. Kiermaier, and T. Slatyer, "Massive amplitudes on the Coulomb branch of N = 4 SYM," *JHEP* **1112**, 097 (2011) [arXiv:1104.2050 [hep-th]].
[141] C. Cheung and D. O'Connell, "Amplitudes and spinor-helicity in six dimensions," *JHEP* **0907**, 075 (2009) [arXiv:0902.0981 [hep-th]].
[142] T. Dennen, Y.-t. Huang, and W. Siegel, "Supertwistor space for 6D maximal super Yang–Mills," *JHEP* **1004**, 127 (2010) [arXiv:0910.2688 [hep-th]].
[143] A. Brandhuber, D. Korres, D. Koschade, and G. Travaglini, "One-loop amplitudes in six-dimensional (1,1) theories from generalised unitarity," *JHEP* **1102**, 077 (2011) [arXiv:1010.1515 [hep-th]];
C. Saemann, R. Wimmer, and M. Wolf, "A twistor description of six-dimensional N = (1,1) super Yang–Mills theory," *JHEP* **1205**, 020 (2012) [arXiv:1201.6285 [hep-th]].
[144] T. Dennen and Y.-t. Huang, "Dual conformal properties of six-dimensional maximal super Yang–Mills amplitudes," *JHEP* **1101**, 140 (2011) [arXiv:1010.5874 [hep-th]].
[145] T. Chern, "Superconformal field theory in six dimensions and supertwistor," [arXiv:0906.0657 [hep-th]];
M. Chiodaroli, M. Gunaydin, and R. Roiban, "Superconformal symmetry, and maximal supergravity in various dimensions," *JHEP* **1203**, 093 (2012) [arXiv:1108.3085 [hep-th]];
L. J. Mason, R. A. Reid-Edwards, and A. Taghavi-Chabert, "Conformal field theories in six-dimensional twistor space," *J. Geom. Phys.* **62**, 2353 (2012) [arXiv:1111.2585 [hep-th]];

Downloaded from https://www.cambridge.org/core. UNIVERSIDAD AUTONOMA METROPOLITANA, on 14 Nov 2026 at 00:33:59, subject to the Cambridge Core terms of use, available at https://www.cambridge.org/core/terms. https://doi.org/10.1017/CBO9781107706620.020

C. Saemann and M. Wolf, "On twistors and conformal field theories from six dimensions," *J. Math. Phys.* **54**, 013507 (2013) [arXiv:1111.2539 [hep-th]].

[146] B. Czech, Y.-t. Huang, and M. Rozali, "Amplitudes for multiple M5 branes," *JHEP* **1210**, 143 (2012) [arXiv:1110.2791 [hep-th]].

[147] C. Saemann and M. Wolf, "Non-Abelian tensor multiplet equations from twistor space," [arXiv:1205.3108 [hep-th]].

[148] Y.-t. Huang and A. E. Lipstein, "Amplitudes of 3D and 6D maximal superconformal theories in supertwistor space," *JHEP* **1010**, 007 (2010) [arXiv:1004.4735 [hep-th]].

[149] L. F. Alday, J. M. Henn, J. Plefka, and T. Schuster, "Scattering into the fifth dimension of N = 4 super Yang–Mills," *JHEP* **1001**, 077 (2010) [arXiv:0908.0684 [hep-th]].

[150] A. Agarwal, N. Beisert, and T. McLoughlin, "Scattering in mass-deformed $N \geq 4$ Chern–Simons models," *JHEP* **0906**, 045 (2009) [arXiv:0812.3367 [hep-th]].

[151] A. Gustavsson, "Algebraic structures on parallel M2-branes," *Nucl. Phys. B* **811**, 66 (2009) [arXiv:0709.1260 [hep-th]].

[152] J. Bagger and N. Lambert, "Gauge symmetry and supersymmetry of multiple M2-branes," *Phys. Rev. D* **77**, 065008 (2008) [arXiv:0711.0955 [hep-th]].

[153] M. A. Bandres, A. E. Lipstein, and J. H. Schwarz, "N = 8 superconformal Chern–Simons theories," *JHEP* **0805**, 025 (2008) [arXiv:0803.3242 [hep-th]].

[154] J. Gomis, G. Milanesi, and J. G. Russo, "Bagger-Lambert theory for general Lie algebras," *JHEP* **0806**, 075 (2008) [arXiv:0805.1012 [hep-th]].

[155] S. Benvenuti, D. Rodriguez-Gomez, E. Tonni, and H. Verlinde, "N = 8 superconformal gauge theories and M2 branes," *JHEP* **0901**, 078 (2009) [arXiv:0805.1087 [hep-th]].

[156] P. -M. Ho, Y. Imamura, and Y. Matsuo, "M2 to D2 revisited," *JHEP* **0807**, 003 (2008) [arXiv:0805.1202 [hep-th]].

[157] T. Bargheer, F. Loebbert, and C. Meneghelli, "Symmetries of tree-level scattering amplitudes in N = 6 superconformal Chern–Simons theory," *Phys. Rev. D* **82**, 045016 (2010) [arXiv:1003.6120 [hep-th]].

[158] O. Aharony, O. Bergman, D. L. Jafferis, and J. Maldacena, "N = 6 superconformal Chern–Simons-matter theories, M2-branes and their gravity duals," *JHEP* **0810**, 091 (2008) [arXiv:0806.1218 [hep-th]].

[159] M. Benna, I. Klebanov, T. Klose, and M. Smedback, "Superconformal Chern–Simons theories and AdS(4)/CFT(3) correspondence," *JHEP* **0809**, 072 (2008) [arXiv:0806.1519 [hep-th]].

[160] M. A. Bandres, A. E. Lipstein, and J. H. Schwarz, "Studies of the ABJM theory in a formulation with manifest SU(4) R-symmetry," *JHEP* **0809**, 027 (2008) [arXiv:0807.0880 [hep-th]].

[161] A. Gustavsson, "Selfdual strings and loop space Nahm equations," *JHEP* **0804**, 083 (2008) [arXiv:0802.3456 [hep-th]];
J. Bagger and N. Lambert, "Three-algebras and N = 6 Chern–Simons gauge theories," *Phys. Rev. D* **79**, 025002 (2009) [arXiv:0807.0163 [hep-th]].

[162] D. Gang, Y.-t. Huang, E. Koh, S. Lee, and A. E. Lipstein, "Tree-level recursion relation and dual superconformal symmetry of the ABJM theory," *JHEP* **1103**, 116 (2011) [arXiv:1012.5032 [hep-th]].

Downloaded from https://www.cambridge.org/core. UNIVERSIDAD AUTONOMA METROPOLITANA, on 14 Nov 2026 at 00:33:59, subject to the Cambridge Core terms of use, available at https://www.cambridge.org/core/terms. https://doi.org/10.1017/CBO9781107706620.020

[163] Y.-t. Huang and A. E. Lipstein, "Dual superconformal symmetry of N = 6 Chern–Simons theory," *JHEP* **1011**, 076 (2010) [arXiv:1008.0041 [hep-th]].
[164] T. Bargheer, N. Beisert, F. Loebbert, and T. McLoughlin, "Conformal anomaly for amplitudes in $\mathcal{N} = 6$ superconformal Chern–Simons theory," *J. Phys. A* **45**, 475402 (2012) [arXiv:1204.4406 [hep-th]].
[165] M. S. Bianchi, M. Leoni, A. Mauri, S. Penati, and A. Santambrogio, "One loop amplitudes In ABJM," *JHEP* **1207**, 029 (2012) [arXiv:1204.4407 [hep-th]].
[166] A. Brandhuber, G. Travaglini, and C. Wen, "All one-loop amplitudes in N = 6 superconformal Chern–Simons theory," *JHEP* **1210**, 145 (2012) [arXiv:1207.6908 [hep-th]].
[167] W. -M. Chen and Y.-t. Huang, "Dualities for loop amplitudes of N = 6 Chern–Simons matter theory," *JHEP* **1111**, 057 (2011) [arXiv:1107.2710 [hep-th]];
[168] A. Brandhuber, G. Travaglini, and C. Wen, "A note on amplitudes in N = 6 superconformal Chern–Simons theory," *JHEP* **1207**, 160 (2012) [arXiv:1205.6705 [hep-th]].
[169] M. S. Bianchi, M. Leoni, A. Mauri, S. Penati, and A. Santambrogio, "Scattering amplitudes/Wilson loop duality in ABJM theory," *JHEP* **1201**, 056 (2012) [arXiv:1107.3139 [hep-th]].
[170] S. Caron-Huot and Y.-t. Huang, "The two-loop six-point amplitude in ABJM theory," *JHEP* **1303**, 075 (2013) [arXiv:1210.4226 [hep-th]].
[171] S. Lee, "Yangian invariant scattering amplitudes in supersymmetric Chern–Simons theory," *Phys. Rev. Lett.* **105**, 151603 (2010) [arXiv:1007.4772 [hep-th]].
[172] Y. -t. Huang, C. Wen, and D. Xie, "The positive orthogonal Grassmannian and loop amplitudes of ABJM," [arXiv:1402.1479 [hep-th]].
[173] Y.-t. Huang and S. Lee, "A new integral formula for supersymmetric scattering amplitudes in three dimensions," *Phys. Rev. Lett.* **109**, 191601 (2012) [arXiv:1207.4851 [hep-th]].
[174] O. T. Engelund and R. Roiban, "A twistor string for the ABJ(M) theory," [arXiv:1401.6242 [hep-th]].
[175] S. Weinberg, *Gravitation and Cosmology: Principles and Applications of the General Theory of Relativity*, USA: John Wiley & Sons (1972).
[176] R. M. Wald, *General Relativity*, Chicago, USA: University Press (1984).
[177] S. M. Carroll, *Spacetime and Geometry: An Introduction to General Relativity*, San Francisco, USA: Addison-Wesley (2004).
[178] H. Elvang and D. Z. Freedman, unpublished notes (2007).
[179] B. S. DeWitt, "Quantum theory of gravity. 2. The manifestly covariant theory," *Phys. Rev.* **162**, 1195 (1967);
B. S. DeWitt, "Quantum theory of gravity. 3. Applications of the covariant theory," *Phys. Rev.* **162**, 1239 (1967);
M. J. G. Veltman, "Quantum theory of gravitation," *Conf. Proc. C* **7507281**, 265 (1975).
[180] F. A. Berends, W. T. Giele, and H. Kuijf, "On relations between multi-gluon and multigraviton scattering," *Phys. Lett. B* **211**, 91 (1988).
[181] J. Bedford, A. Brandhuber, B. J. Spence, and G. Travaglini, "A recursion relation for gravity amplitudes," *Nucl. Phys. B* **721**, 98 (2005) [hep-th/0502146].

Downloaded from https://www.cambridge.org/core. UNIVERSIDAD AUTONOMA METROPOLITANA, on 14 Nov 2026 at 00:33:59, subject to the Cambridge Core terms of use, available at https://www.cambridge.org/core/terms. https://doi.org/10.1017/CBO9781107706620.020

[182] H. Elvang and D. Z. Freedman, "Note on graviton MHV amplitudes," *JHEP* **0805**, 096 (2008) [arXiv:0710.1270 [hep-th]].
[183] D. Nguyen, M. Spradlin, A. Volovich, and C. Wen, "The tree formula for MHV graviton amplitudes," *JHEP* **1007**, 045 (2010) [arXiv:0907.2276 [hep-th]].
[184] Z. Bern, L. J. Dixon, M. Perelstein, and J. S. Rozowsky, "Multileg one loop gravity amplitudes from gauge theory," *Nucl. Phys. B* **546**, 423 (1999) [hep-th/9811140].
[185] Z. Bern and A. K. Grant, "Perturbative gravity from QCD amplitudes," *Phys. Lett. B* **457**, 23 (1999) [hep-th/9904026].
[186] Z. Bern, L. J. Dixon, D. C. Dunbar, *et al.*, "On perturbative gravity and gauge theory," *Nucl. Phys. Proc. Suppl.* **88**, 194 (2000) [hep-th/0002078].
[187] W. Siegel, "Two vierbein formalism for string inspired axionic gravity," *Phys. Rev. D* **47**, 5453 (1993) [hep-th/9302036].
[188] Z. Bern, "Perturbative quantum gravity and its relation to gauge theory," *Living Rev. Rel.* **5**, 5 (2002) [gr-qc/0206071].
[189] D. Z. Freedman, "Some beautiful equations of mathematical physics," In *ICTP (ed.): The Dirac Medals of the ICTP 1993* 25–53, and CERN Geneva - TH.-7367 (94/07,rec.Sep.) [hep-th/9408175].
[190] S. J. Gates, M. T. Grisaru, M. Rocek, and W. Siegel, "Superspace or one thousand and one lessons in supersymmetry," *Front. Phys.* **58**, 1 (1983) [hep-th/0108200].
[191] B. de Wit and D. Z. Freedman, "On SO(8) extended supergravity," *Nucl. Phys. B* **130**, 105 (1977).
[192] E. Cremmer and B. Julia, "The N = 8 supergravity theory. 1. The Lagrangian," *Phys. Lett. B* **80**, 48 (1978);
E. Cremmer and B. Julia, "The SO(8) supergravity," *Nucl. Phys. B* **159**, 141 (1979).
[193] B. de Wit and H. Nicolai, "N = 8 supergravity," *Nucl. Phys. B* **208**, 323 (1982).
[194] A. Hodges, "A simple formula for gravitational MHV amplitudes," [arXiv:1204.1930 [hep-th]].
[195] F. Cachazo, L. Mason, and D. Skinner, "Gravity in twistor space and its Grassmannian formulation," [arXiv:1207.4712 [hep-th]].
[196] S. He, "A link representation for gravity amplitudes," [arXiv:1207.4064 [hep-th]].
[197] F. Cachazo and Y. Geyer, "A 'twistor string' inspired formula for tree-level scattering amplitudes in N = 8 SUGRA," [arXiv:1206.6511 [hep-th]].
[198] D. Skinner, "Twistor strings for N = 8 supergravity," [arXiv:1301.0868 [hep-th]].
[199] F. Cachazo, S. He, and E. Y. Yuan, "Scattering equations and KLT orthogonality," [arXiv:1306.6575 [hep-th]].
[200] F. Cachazo, S. He, and E. Y. Yuan, "Scattering of massless particles in arbitrary dimension," [arXiv:1307.2199 [hep-th]].
[201] S. L. Adler, "Consistency conditions on the strong interactions implied by a partially conserved axial vector current," *Phys. Rev.* **137**, B1022 (1965).
[202] S. R. Coleman, "Secret symmetry: An introduction to spontaneous symmetry breakdown and gauge fields," *Subnucl. Ser.* **11**, 139 (1975).
[203] M. Kiermaier, "The Coulomb-branch S-matrix from massless amplitudes," [arXiv:1105.5385 [hep-th]].

Downloaded from https://www.cambridge.org/core. UNIVERSIDAD AUTONOMA METROPOLITANA, on 14 Nov 2026 at 00:33:59, subject to the Cambridge Core terms of use, available at https://www.cambridge.org/core/terms. https://doi.org/10.1017/CBO9781107706620.020

[204] G. 't Hooft and M. J. G. Veltman, "One loop divergencies in the theory of gravitation," *Annales Poincaré Phys. Theor. A* **20**, 69 (1974).
[205] M. H. Goroff and A. Sagnotti, "Quantum gravity at two loops," *Phys. Lett. B* **160**, 81 (1985).
[206] A. E. M. van de Ven, "Two loop quantum gravity," *Nucl. Phys. B* **378**, 309 (1992).
[207] S. Deser and P. van Nieuwenhuizen, "One loop divergences of quantized Einstein–Maxwell fields," *Phys. Rev. D* **10**, 401 (1974).
[208] M. T. Grisaru, P. van Nieuwenhuizen, and J. A. M. Vermaseren, "One loop renormalizability of pure supergravity and of Maxwell–Einstein theory in extended supergravity," *Phys. Rev. Lett.* **37**, 1662 (1976).
[209] M. T. Grisaru, "Two loop renormalizability of supergravity," *Phys. Lett. B* **66**, 75 (1977).
[210] E. Tomboulis, "On the two loop divergences of supersymmetric gravitation," *Phys. Lett. B* **67**, 417 (1977).
[211] S. Deser, J. H. Kay, and K. S. Stelle, "Renormalizability properties of supergravity," *Phys. Rev. Lett.* **38**, 527 (1977).
[212] Z. Bern, L. J. Dixon, and R. Roiban, "Is N = 8 supergravity ultraviolet finite?," *Phys. Lett. B* **644**, 265 (2007) [hep-th/0611086].
[213] Z. Bern, J. J. Carrasco, L. J. Dixon, *et al.*, "Three-loop superfiniteness of N = 8 supergravity," *Phys. Rev. Lett.* **98**, 161303 (2007) [hep-th/0702112].
[214] Z. Bern, J. J. M. Carrasco, L. J. Dixon, H. Johansson, and R. Roiban, "Manifest ultraviolet behavior for the three-loop four-point amplitude of N = 8 supergravity," *Phys. Rev. D* **78**, 105019 (2008) [arXiv:0808.4112 [hep-th]].
[215] P. S. Howe and K. S. Stelle, "Supersymmetry counterterms revisited," *Phys. Lett. B* **554**, 190 (2003) [hep-th/0211279].
[216] Z. Bern, J. J. Carrasco, L. J. Dixon, H. Johansson, and R. Roiban, "The ultraviolet behavior of N = 8 supergravity at four loops," *Phys. Rev. Lett.* **103**, 081301 (2009) [arXiv:0905.2326 [hep-th]].
[217] Z. Bern, J. J. M. Carrasco, L. J. Dixon, H. Johansson, and R. Roiban, "The complete four-loop four-point amplitude in N = 4 super-Yang–Mills theory," *Phys. Rev. D* **82**, 125040 (2010) [arXiv:1008.3327 [hep-th]].
[218] J. Bjornsson and M. B. Green, "5 loops in 24/5 dimensions," *JHEP* **1008**, 132 (2010) [arXiv:1004.2692 [hep-th]].
[219] H. Elvang, D. Z. Freedman, and M. Kiermaier, "A simple approach to counterterms in N = 8 supergravity," *JHEP* **1011**, 016 (2010) [arXiv:1003.5018 [hep-th]].
[220] H. Elvang and M. Kiermaier, "Stringy KLT relations, global symmetries, and $E_{7(7)}$ violation," *JHEP* **1010**, 108 (2010) [arXiv:1007.4813 [hep-th]].
[221] N. Beisert, H. Elvang, D. Z. Freedman, *et al.*, "E7(7) constraints on counterterms in N = 8 supergravity," *Phys. Lett. B* **694**, 265 (2010) [arXiv:1009.1643 [hep-th]].
[222] P. van Nieuwenhuizen and C. C. Wu, "On integral relations for invariants constructed from three riemann tensors and their applications in quantum gravity," *J. Math. Phys.* **18**, 182 (1977).
[223] G. Bossard, C. Hillmann, and H. Nicolai, "E7(7) symmetry in perturbatively quantised N = 8 supergravity," *JHEP* **1012**, 052 (2010) [arXiv:1007.5472 [hep-th]].

Downloaded from https://www.cambridge.org/core. UNIVERSIDAD AUTONOMA METROPOLITANA, on 14 Nov 2026 at 00:33:59, subject to the Cambridge Core terms of use, available at https://www.cambridge.org/core/terms. https://doi.org/10.1017/CBO9781107706620.020

[224] D. Z. Freedman and E. Tonni, "The $D^{2k}R^4$ invariants of $N = 8$ supergravity," *JHEP* **1104**, 006 (2011) [arXiv:1101.1672 [hep-th]].

[225] S. Deser and J. H. Kay, "Three loop counterterms for extended supergravity," *Phys. Lett. B* **76**, 400 (1978).

[226] J. M. Drummond, P. J. Heslop, and P. S. Howe, "A note on N = 8 counterterms," [arXiv:1008.4939 [hep-th]].

[227] G. Bossard and H. Nicolai, "Counterterms vs. dualities," *JHEP* **1108**, 074 (2011) [arXiv:1105.1273 [hep-th]].

[228] R. Kallosh and T. Kugo, "The footprint of E(7(7)) amplitudes of N = 8 supergravity," *JHEP* **0901**, 072 (2009) [arXiv:0811.3414 [hep-th]];
R. Kallosh, "$E_{7(7)}$ symmetry and finiteness of N = 8 supergravity," *JHEP* **1203**, 083 (2012) [arXiv:1103.4115 [hep-th]];
R. Kallosh, "N = 8 counterterms and $E_{7(7)}$ current conservation," *JHEP* **1106**, 073 (2011) [arXiv:1104.5480 [hep-th]];
R. Kallosh and T. Ortin, "New E77 invariants and amplitudes," *JHEP* **1209**, 137 (2012) [arXiv:1205.4437 [hep-th]];
M. Gunaydin and R. Kallosh, "Obstruction to $E_{7(7)}$ deformation in N = 8 supergravity," [arXiv:1303.3540 [hep-th]];
J. J. M. Carrasco and R. Kallosh, "Hidden supersymmetry may imply duality invariance," [arXiv:1303.5663 [hep-th]].

[229] S. Stieberger and T. R. Taylor, "Complete six-gluon disk amplitude in superstring theory," *Nucl. Phys. B* **801**, 128 (2008) [arXiv:0711.4354 [hep-th]].

[230] J. Broedel and L. J. Dixon, "R**4 counterterm and E(7)(7) symmetry in maximal supergravity," *JHEP* **1005**, 003 (2010) [arXiv:0911.5704 [hep-th]].

[231] N. Berkovits, "New higher-derivative R**4 theorems," *Phys. Rev. Lett.* **98**, 211601 (2007) [arXiv:hep-th/0609006];
M. B. Green, J. G. Russo, and P. Vanhove, "Non-renormalisation conditions in type II string theory and maximal supergravity," *JHEP* **0702**, 099 (2007) [arXiv:hep-th/0610299];
M. B. Green, J. G. Russo, and P. Vanhove, "Ultraviolet properties of maximal supergravity," *Phys. Rev. Lett.* **98**, 131602 (2007) [arXiv:hep-th/0611273];
M. B. Green, J. G. Russo, and P. Vanhove, "Modular properties of two-loop maximal supergravity and connections with string theory," *JHEP* **0807**, 126 (2008) [arXiv:0807.0389 [hep-th]];
N. Berkovits, M. B. Green, J. G. Russo, and P. Vanhove, "Non-renormalization conditions for four-gluon scattering in supersymmetric string and field theory," *JHEP* **0911**, 063 (2009) [arXiv:0908.1923 [hep-th]];
P. Vanhove, "The critical ultraviolet behaviour of N = 8 supergravity amplitudes," [arXiv:1004.1392 [hep-th]].

[232] Z. Bern, J. J. M. Carrasco, L. J. Dixon, H. Johansson, and R. Roiban, "Simplifying multiloop integrands and ultraviolet divergences of gauge theory and gravity amplitudes," *Phys. Rev. D* **85**, 105014 (2012) [arXiv:1201.5366 [hep-th]].

[233] P. S. Howe and U. Lindstrom, "Higher order invariants in extended supergravity," *Nucl. Phys. B* **181**, 487 (1981).

Downloaded from https://www.cambridge.org/core. UNIVERSIDAD AUTONOMA METROPOLITANA, on 14 Nov 2026 at 00:33:59, subject to the Cambridge Core terms of use, available at https://www.cambridge.org/core/terms. https://doi.org/10.1017/CBO9781107706620.020

[234] R. E. Kallosh, "Counterterms in extended supergravities," *Phys. Lett. B* **99**, 122 (1981).
[235] G. Bossard, P. S. Howe, K. S. Stelle, and P. Vanhove, "The vanishing volume of D = 4 superspace," *Class. Quant. Grav.* **28**, 215005 (2011) [arXiv:1105.6087 [hep-th]].
[236] N. Berkovits, "Super Poincaré covariant quantization of the superstring," *JHEP* **0004**, 018 (2000) [hep-th/0001035].
[237] M. B. Green, H. Ooguri, and J. H. Schwarz, "Nondecoupling of maximal supergravity from the superstring," *Phys. Rev. Lett.* **99**, 041601 (2007) [arXiv:0704.0777 [hep-th]].
[238] T. Banks, "Arguments against a finite N = 8 supergravity," [arXiv:1205.5768 [hep-th]].
[239] M. Bianchi, S. Ferrara, and R. Kallosh, "Perturbative and non-perturbative N = 8 supergravity," *Phys. Lett. B* **690**, 328 (2010) [arXiv:0910.3674 [hep-th]].
[240] Z. Bern, S. Davies, T. Dennen, Y.-t. Huang, and J. Nohle, "Color-kinematics duality for pure Yang–Mills and gravity at one and two loops," [arXiv:1303.6605 [hep-th]].
[241] Z. Bern, S. Davies, T. Dennen, A. V. Smirnov, and V. A. Smirnov, "The ultraviolet properties of N = 4 supergravity at four loops," *Phys. Rev. Lett.* **111**, 231302 (2013) [arXiv:1309.2498 [hep-th]].
[242] Z. Bern, L. J. Dixon, D. C. Dunbar, M. Perelstein, and J. S. Rozowsky, "On the relationship between Yang–Mills theory and gravity and its implication for ultraviolet divergences," *Nucl. Phys. B* **530**, 401 (1998) [hep-th/9802162].
[243] D. C. Dunbar, B. Julia, D. Seminara, and M. Trigiante, "Counterterms in type I supergravities," *JHEP* **0001**, 046 (2000) [hep-th/9911158].
[244] Z. Bern, S. Davies, T. Dennen, and Y.-t. Huang, "Ultraviolet cancellations in half-maximal supergravity as a consequence of the double-copy structure," *Phys. Rev. D* **86**, 105014 (2012) [arXiv:1209.2472 [hep-th]].
[245] Z. Bern, S. Davies, T. Dennen, and Y.-t. Huang, "Absence of three-loop four-point divergences in N = 4 supergravity," *Phys. Rev. Lett.* **108**, 201301 (2012) [arXiv:1202.3423 [hep-th]];
[246] M. Fischler, "Finiteness calculations for O(4) through O(8) extended supergravity and O(4) supergravity coupled to selfdual O(4) matter," *Phys. Rev. D* **20**, 396 (1979).
[247] Z. Bern, S. Davies, and T. Dennen, "The ultraviolet structure of half-maximal supergravity with matter multiplets at two and three loops," [arXiv:1305.4876 [hep-th]].
[248] G. Bossard, P. S. Howe, and K. S. Stelle, "Invariants and divergences in half-maximal supergravity theories," [arXiv:1304.7753 [hep-th]].
[249] D. Vaman and Y.-P. Yao, "Constraints and generalized gauge transformations on tree-level gluon and graviton amplitudes," *JHEP* **1011**, 028 (2010) [arXiv:1007.3475 [hep-th]].
[250] R. H. Boels and R. S. Isermann, "On powercounting in perturbative quantum gravity theories through color-kinematic duality," *JHEP* **1306**, 017 (2013) [arXiv:1212.3473].

Downloaded from https://www.cambridge.org/core. UNIVERSIDAD AUTONOMA METROPOLITANA, on 14 Nov 2026 at 00:33:59, subject to the Cambridge Core terms of use, available at https://www.cambridge.org/core/terms. https://doi.org/10.1017/CBO9781107706620.020

[251] Z. Bern, T. Dennen, Y.-t. Huang, and M. Kiermaier, "Gravity as the square of gauge theory," *Phys. Rev. D* **82**, 065003 (2010) [arXiv:1004.0693 [hep-th]].

[252] M. Kiermaier, Talk at *Amplitudes 2010*, May 2010 at QMUL, London, UK. http://www.strings.ph.qmul.ac.uk/∼theory/Amplitudes2010/

[253] N. E. J. Bjerrum-Bohr, P. H. Damgaard, T. Sondergaard, and P. Vanhove, "The momentum kernel of gauge and gravity theories," *JHEP* **1101**, 001 (2011) [arXiv:1010.3933 [hep-th]].

[254] C. R. Mafra, O. Schlotterer, and S. Stieberger, "Explicit BCJ numerators from pure spinors," *JHEP* **1107**, 092 (2011) [arXiv:1104.5224 [hep-th]];
C.-H. Fu, Y.-J. Du, and B. Feng, "An algebraic approach to BCJ numerators," *JHEP* **1303**, 050 (2013) [arXiv:1212.6168 [hep-th]].

[255] N. E. J. Bjerrum-Bohr, P. H. Damgaard, and P. Vanhove, "Minimal basis for gauge theory amplitudes," *Phys. Rev. Lett.* **103**, 161602 (2009) [0907.1425 [hep-th]];
S. Stieberger, "Open & closed vs. pure open string disk amplitudes," [arXiv:0907.2211 [hep-th]];
C. R. Mafra and O. Schlotterer, "The structure of n-point one-loop open superstring amplitudes," [arXiv:1203.6215 [hep-th]];
O. Schlotterer and S. Stieberger, "Motivic multiple zeta values and superstring amplitudes," [arXiv:1205.1516 [hep-th]];
J. Broedel, O. Schlotterer, and S. Stieberger, "Polylogarithms, multiple zeta values and superstring amplitudes," [arXiv:1304.7267 [hep-th]].

[256] S. H. Henry Tye and Y. Zhang, "Dual identities inside the gluon and the graviton scattering amplitudes," *JHEP* **1006**, 071 (2010) [Erratum-*ibid.* **1104**, 114 (2011)] [arXiv:1003.1732 [hep-th]].

[257] B. Feng, R. Huang, and Y. Jia, "Gauge amplitude identities by on-shell recursion relation in s-matrix program," *Phys. Lett. B* **695**, 350 (2011) [arXiv:1004.3417 [hep-th]].

[258] F. Cachazo, "Fundamental BCJ relation in N = 4 SYM from the connected formulation," [arXiv:1206.5970 [hep-th]].

[259] N. E. J. Bjerrum-Bohr, P. H. Damgaard, R. Monteiro, and D. O'Connell, "Algebras for amplitudes," *JHEP* **1206**, 061 (2012) [arXiv:1203.0944 [hep-th]].

[260] R. Monteiro and D. O'Connell, "The kinematic algebra from the self-dual sector," *JHEP* **1107**, 007 (2011) [arXiv:1105.2565 [hep-th]].

[261] M. Tolotti and S. Weinzierl, "Construction of an effective Yang–Mills Lagrangian with manifest BCJ duality," [arXiv:1306.2975 [hep-th]].

[262] Z. Bern, J. J. M. Carrasco, and H. Johansson, "Perturbative quantum gravity as a double copy of gauge theory," *Phys. Rev. Lett.* **105**, 061602 (2010) [arXiv:1004.0476 [hep-th]].

[263] J. J. Carrasco and H. Johansson, "Five-point amplitudes in N = 4 super-Yang–Mills theory and N = 8 supergravity," *Phys. Rev. D* **85**, 025006 (2012) [arXiv:1106.4711 [hep-th]].

[264] N. E. J. Bjerrum-Bohr, T. Dennen, R. Monteiro, and D. O'Connell, "Integrand oxidation and one-loop colour-dual numerators in N = 4 gauge theory," [arXiv:1303.2913 [hep-th]].

Downloaded from https://www.cambridge.org/core. UNIVERSIDAD AUTONOMA METROPOLITANA, on 14 Nov 2026 at 00:33:59, subject to the Cambridge Core terms of use, available at https://www.cambridge.org/core/terms. https://doi.org/10.1017/CBO9781107706620.020

[265] R. H. Boels, R. S. Isermann, R. Monteiro, and D. O'Connell, "Colour-kinematics duality for one-loop rational amplitudes," *JHEP* **1304**, 107 (2013) [arXiv:1301.4165 [hep-th]].

[266] J. J. M. Carrasco, M. Chiodaroli, M. Günaydin, and R. Roiban, "One-loop four-point amplitudes in pure and matter-coupled $N \leq 4$ supergravity," *JHEP* **1303**, 056 (2013) [arXiv:1212.1146 [hep-th]].

[267] M. Chiodaroli, Q. Jin, and R. Roiban, "Color/kinematics duality for general abelian orbifolds of N = 4 super Yang–Mills theory," *JHEP* **1401**, 152 (2014) [arXiv:1311.3600 [hep-th]].

[268] J. Nohle, "Color-kinematics duality in one-loop four-gluon amplitudes with matter," [arXiv:1309.7416 [hep-th]].

[269] Z. Bern, C. Boucher-Veronneau, and H. Johansson, "$N \geq 4$ supergravity amplitudes from gauge theory at one loop," *Phys. Rev. D* **84**, 105035 (2011) [arXiv:1107.1935 [hep-th]];
C. Boucher-Veronneau and L. J. Dixon, "$N \geq 4$ supergravity amplitudes from gauge theory at two loops," *JHEP* **1112**, 046 (2011) [arXiv:1110.1132 [hep-th]].

[270] M. T. Grisaru and W. Siegel, "Supergraphity. 2. Manifestly covariant rules and higher loop finiteness," *Nucl. Phys. B* **201**, 292 (1982) [Erratum-*ibid. B* **206**, 496 (1982)].

[271] S. Ferrara, R. Kallosh, and A. Van Proeyen, "Conjecture on hidden superconformal symmetry of N = 4 supergravity," *Phys. Rev. D* **87**, 025004 (2013) [arXiv:1209.0418 [hep-th]].

[272] J. Broedel and L. J. Dixon, "Color-kinematics duality and double-copy construction for amplitudes from higher-dimension operators," *JHEP* **1210**, 091 (2012) [arXiv:1208.0876 [hep-th]].

[273] R. H. Boels, B. A. Kniehl, O. V. Tarasov, and G. Yang, "Color-kinematic duality for form factors," *JHEP* **1302**, 063 (2013) [arXiv:1211.7028 [hep-th]];

[274] T. Bargheer, S. He, and T. McLoughlin, "New relations for three-dimensional supersymmetric scattering amplitudes," *Phys. Rev. Lett.* **108**, 231601 (2012) [arXiv:1203.0562 [hep-th]].

[275] Y.-t. Huang and H. Johansson, "Equivalent D = 3 supergravity amplitudes from double copies of three-algebra and two-algebra gauge theories," [arXiv:1210.2255 [hep-th]].

[276] A. Zee, "Quantum field theory in a nutshell," Princeton, USA: Princeton University Press (2010).

[277] M. D. Schwartz, *Quantum Field Theory and the Standard Model*, Cambridge, UK: Cambridge University Press (2013).

[278] J. M. Henn and J. C. Plefka, "Scattering amplitudes in gauge theories," *Lecture Notes in Physics 883* Heidelbera: Springer (2014).

[279] M. Wolf, "A first course on twistors, integrability and gluon scattering amplitudes," *J. Phys. A* **43**, 393001 (2010) [arXiv:1001.3871 [hep-th]].

[280] Z. Bern, L. J. Dixon, and D. A. Kosower, "On-shell methods in perturbative QCD," *Annals Phys.* **322**, 1587 (2007) [arXiv:0704.2798 [hep-ph]].

[281] M. E. Peskin, "Simplifying multi-jet QCD computation," [arXiv:1101.2414 [hep-ph]].

Downloaded from https://www.cambridge.org/core. UNIVERSIDAD AUTONOMA METROPOLITANA, on 14 Nov 2026 at 00:33:59, subject to the Cambridge Core terms of use, available at https://www.cambridge.org/core/terms. https://doi.org/10.1017/CBO9781107706620.020

[282] J. F. Donoghue, "Introduction to the effective field theory description of gravity," gr-qc/9512024.
[283] L. J. Dixon, "Ultraviolet behavior of N = 8 supergravity," [arXiv:1005.2703 [hep-th]].
[284] Z. Bern, P. Gondolo, and M. Perelstein, "Neutralino annihilation into two photons," *Phys. Lett. B* **411**, 86 (1997) [hep-ph/9706538];
Z. Bern, A. De Freitas, and L. J. Dixon, "Two loop helicity amplitudes for gluon–gluon scattering in QCD and supersymmetric Yang–Mills theory," *JHEP* **0203**, 018 (2002) [hep-ph/0201161];
Z. Bern, A. De Freitas, and L. J. Dixon, "Two loop helicity amplitudes for quark gluon scattering in QCD and gluino gluon scattering in supersymmetric Yang–Mills theory," *JHEP* **0306**, 028 (2003) [hep-ph/0304168].
[285] S. J. Bidder, N. E. J. Bjerrum-Bohr, D. C. Dunbar, and W. B. Perkins, "One-loop gluon scattering amplitudes in theories with N < 4 supersymmetries," *Phys. Lett. B* **612**, 75 (2005) [hep-th/0502028].
[286] R. Britto, E. Buchbinder, F. Cachazo, and B. Feng, "One-loop amplitudes of gluons in SQCD," *Phys. Rev. D* **72**, 065012 (2005) [hep-ph/0503132].
[287] S. Lal and S. Raju, "Rational terms in theories with matter," *JHEP* **1008**, 022 (2010) [arXiv:1003.5264 [hep-th]].
[288] S. Dittmaier, "Weyl-van der Waerden formalism for helicity amplitudes of massive particles," *Phys. Rev. D* **59**, 016007 (1998) [hep-ph/9805445].
[289] R. Boels and C. Schwinn, "CSW rules for massive matter legs and glue loops," *Nucl. Phys. Proc. Suppl.* **183**, 137 (2008) [arXiv:0805.4577 [hep-th]].
[290] R. H. Boels, "No triangles on the moduli space of maximally supersymmetric gauge theory," *JHEP* **1005**, 046 (2010) [arXiv:1003.2989 [hep-th]].
[291] P. Ferrario, G. Rodrigo, and P. Talavera, "Compact multigluonic scattering amplitudes with heavy scalars and fermions," *Phys. Rev. Lett.* **96**, 182001 (2006) [hep-th/0602043].
[292] D. Forde and D. A. Kosower, "All-multiplicity amplitudes with massive scalars," *Phys. Rev. D* **73**, 065007 (2006) [hep-th/0507292].
[293] G. Rodrigo, "Multigluonic scattering amplitudes of heavy quarks," *JHEP* **0509**, 079 (2005) [hep-ph/0508138].
[294] C. Cheung, D. O'Connell, and B. Wecht, "BCFW recursion relations and string theory," *JHEP* **1009**, 052 (2010) [arXiv:1002.4674 [hep-th]].
[295] R. H. Boels, D. Marmiroli, and N. A. Obers, "On-shell recursion in string theory," *JHEP* **1010**, 034 (2010) [arXiv:1002.5029 [hep-th]].
[296] K. Kampf, J. Novotny, and J. Trnka, "Recursion relations for tree-level amplitudes in the SU(N) non-linear sigma model," *Phys. Rev. D* **87**, 081701 (2013) [arXiv:1212.5224 [hep-th]].
[297] Z. Bern, L. J. Dixon, and D. A. Kosower, "The last of the finite loop amplitudes in QCD," *Phys. Rev. D* **72**, 125003 (2005) [hep-ph/0505055].
[298] B. Feng and M. Luo, "An introduction to on-shell recursion relations," [arXiv:1111.5759 [hep-th]].

Downloaded from https://www.cambridge.org/core. UNIVERSIDAD AUTONOMA METROPOLITANA, on 14 Nov 2026 at 00:33:59, subject to the Cambridge Core terms of use, available at https://www.cambridge.org/core/terms. https://doi.org/10.1017/CBO9781107706620.020

[299] J. M. Drummond, G. P. Korchemsky, and E. Sokatchev, "Conformal properties of four-gluon planar amplitudes and Wilson loops," *Nucl. Phys. B* **795**, 385 (2008) [arXiv:0707.0243 [hep-th]].
[300] N. Berkovits and J. Maldacena, "Fermionic T-duality, dual superconformal symmetry, and the amplitude/Wilson loop connection," *JHEP* **0809**, 062 (2008) [arXiv:0807.3196 [hep-th]].
[301] N. Beisert, R. Ricci, A. A. Tseytlin, and M. Wolf, "Dual superconformal symmetry from AdS(5) x S**5 superstring integrability," *Phys. Rev. D* **78**, 126004 (2008) [arXiv:0807.3228 [hep-th]].
[302] A. Brandhuber, P. Heslop, and G. Travaglini, "MHV amplitudes in N = 4 super Yang–Mills and Wilson loops," *Nucl. Phys. B* **794**, 231 (2008) [arXiv:0707.1153 [hep-th]].
[303] L. J. Mason and D. Skinner, "The complete planar S-matrix of N = 4 SYM as a Wilson loop in twistor space," *JHEP* **1012**, 018 (2010) [arXiv:1009.2225 [hep-th]].
[304] S. Caron-Huot, "Notes on the scattering amplitude / Wilson loop duality," *JHEP* **1107**, 058 (2011) [arXiv:1010.1167 [hep-th]].
[305] B. Eden, P. Heslop, G. P. Korchemsky, and E. Sokatchev, "The super-correlator/super-amplitude duality: Part I," *Nucl. Phys. B* **869**, 329 (2013) [arXiv:1103.3714 [hep-th]]; Part II, *Nucl. Phys. B* **869**, 378 (2013) [arXiv:1103.4353 [hep-th]].
[306] L. F. Alday, B. Eden, G. P. Korchemsky, J. Maldacena, and E. Sokatchev, "From correlation functions to Wilson loops," *JHEP* **1109**, 123 (2011) [arXiv:1007.3243 [hep-th]].
[307] T. Adamo, M. Bullimore, L. Mason, and D. Skinner, "A proof of the supersymmetric correlation function / Wilson loop correspondence," *JHEP* **1108**, 076 (2011) [arXiv:1103.4119 [hep-th]].
[308] L. F. Alday and R. Roiban, "Scattering amplitudes, Wilson loops and the string/gauge theory correspondence," *Phys. Rept.* **468**, 153 (2008) [arXiv:0807.1889 [hep-th]].
[309] R. M. Schabinger, "One-loop N = 4 super Yang–Mills scattering amplitudes in d dimensions, relation to open strings and polygonal Wilson loops," *J. Phys. A* **44**, 454007 (2011) [arXiv:1104.3873 [hep-th]].
[310] J. M. Henn, "Duality between Wilson loops and gluon amplitudes," *Fortsch. Phys.* **57**, 729 (2009) [arXiv:0903.0522 [hep-th]].
[311] T. Adamo, M. Bullimore, L. Mason, and D. Skinner, "Scattering amplitudes and Wilson loops in twistor space," *J. Phys. A* **44**, 454008 (2011) [arXiv:1104.2890 [hep-th]].
[312] L. F. Alday, D. Gaiotto, J. Maldacena, A. Sever, and P. Vieira, "An operator product expansion for polygonal null Wilson loops," *JHEP* **1104**, 088 (2011) [arXiv:1006.2788 [hep-th]].
[313] B. Basso, A. Sever, and P. Vieira, "Space-time S-matrix and flux-tube S-matrix at finite coupling," *Phys. Rev. Lett.* **111**, 091602 (2013) [arXiv:1303.1396 [hep-th]].
[314] B. Basso, A. Sever, and P. Vieira, "Space-time S-matrix and flux tube S-matrix II. Extracting and matching data," *JHEP* **1401**, 008 (2014) [arXiv:1306.2058 [hep-th]].

Downloaded from https://www.cambridge.org/core. UNIVERSIDAD AUTONOMA METROPOLITANA, on 14 Nov 2026 at 00:33:59, subject to the Cambridge Core terms of use, available at https://www.cambridge.org/core/terms. https://doi.org/10.1017/CBO9781107706620.020

[315] B. Basso, A. Sever, and P. Vieira, "Space-time S-matrix and flux-tube S-matrix III. The two-particle contributions," [arXiv:1402.3307 [hep-th]].
[316] L. J. Dixon, J. M. Drummond, M. von Hippel, and J. Pennington, "Hexagon functions and the three-loop remainder function," *JHEP* **1312**, 049 (2013) [arXiv:1308.2276 [hep-th]].
[317] L. J. Dixon, J. M. Drummond, C. Duhr, and J. Pennington, "The four-loop remainder function and multi-Regge behavior at NNLLA in planar N = 4 super-Yang–Mills theory," [arXiv:1402.3300 [hep-th]].
[318] R. Penrose and W. Rindler, *Spinors and Space-Time*, vol. 2, Cambridge: Cambridge University Press (1986).
[319] R. Ward and R. Wells, *Twistor Geometry and Field Theory*, Cambridge: Cambridge University Press (1990).
[320] S. Huggett and P. Tod, *An Introduction to Twistor Theory*, Student Texts 4, London: London Mathematical Society (1985).
[321] F. Cachazo and P. Svrcek, "Lectures on twistor strings and perturbative Yang–Mills theory," *PoS RTN* **2005**, 004 (2005) [hep-th/0504194].

Downloaded from https://www.cambridge.org/core. UNIVERSIDAD AUTONOMA METROPOLITANA, on 14 Nov 2026 at 00:33:59, subject to the Cambridge Core terms of use, available at https://www.cambridge.org/core/terms. https://doi.org/10.1017/CBO9781107706620.020