doi:10.1017/S0043933919000060

Fertility and hatchability in duck eggs

M.E. ABD EL-HACK[1]*, C.B. HURTADO[2], D.M. TORO[3], M. ALAGAWANY[1]*, E.M. ABDELFATTAH[4] and S.S. ELNESR[5]

[1]Poultry Department, Faculty of Agriculture, Zagazig University, Zagazig 44511, Egypt; [2]Department of Livestock Sciences, Faculty of Veterinary Medicine and Zootechnic, University of Córdoba, Monteria 230002, Colombia; [3]Laboratory of Animal Nutrition, Faculty of Natural Sciences, Autonomous University of Queretaro, Queretaro 76230, Mexico; [4]Department of Animal Science, University of California Davis, Meyer Hall, One Shields Avenue, Davis, CA 95616. USA; [5]Poultry Production Department, Faculty of Agriculture, Fayoum University, Fayoum, 63514, Egypt
***Corresponding author: m.ezzat@zu.edu.eg; dr.mahmoud.alagwany@gmail.com**

Ducks are waterfowl belonging to the *Anatidae* family of cosmopolitan distribution. In duck production systems, obtaining ducklings at one-day-old is determinant for the productive chain. The egg production in some species of ducks reaches about 250 to 300 eggs per year. Obtaining one-day-old ducklings can be done by natural incubation with a broody female duck or artificially in an incubator. During artificial incubation, fertility and hatchability are the most important indicators that must be controlled, because they influence the supply of ducklings to the farm. Many factors are related to fertility and hatching, such as environmental conditions, production system, season, nutrition, management of broodstock, storage time of egg and cleaning of eggs before the incubation. According to some reports, Pekin eggs have greater hatchability than Muscovy eggs. The eggs of Muscovy have presented values lower than 22.7% of hatchability. The hatchability of Pekin duck eggs was 78.0% in the spring, while in summer it was around 46.5%. The best hatchability is observed during the winter (57.68%), as in the summer it decreases to 54.14%. The reproductive characteristics of flocks, age, external and internal quality of the egg, male female relation, and presence of lethal genes are factors that directly involve breeders. Larger sexual ratios between males and females of 1:4.3 to 1: 10 cause reduced egg fertility from 75.9% down to 49.6%. Successful production of day-old ducklings starts with the proper selection and management of breeding stock, proper post-lay handling of fertile eggs and the correct incubation process. There are different methods used to improve the hatchability such as dipping eggs in nutrients during the incubation period.

Keywords: ducks; fertility; hatchability; eggs; incubation

© World's Poultry Science Association 2019

Received for publication November 3, 2018
Accepted for publication September 4, 2019

https://doi.org/10.1017/S0043933919000060 Published online by Cambridge University Press

Introduction

Ducks are waterfowl that belong to the *Anatidae* family. Other waterfowl such as loons, grebes, gallinules and coots may be confused with this species (Idahor *et al.,* 2015; Ericson *et al.,* 2017; Farghly *et al.,* 2018a) but do not represent a monophyletic group; therefore, swans and geese are not considered ducks. They are distributed throughout the world, except in Antarctica (Idahor *et al.,* 2015), due to their adaptability to different environments. Because of the variable market demand for a one-day-old chick by the duck industry, fertility and hatchability are important economic factors which represent the major components of reproductive performance and are sensitive to environmental and genetic factors (Widiyaningrum *et al.,* 2016).

Fertility is defined as the percentage of embryonated eggs after three days placement in the incubator; meanwhile, hatchability is the percentage of fertile eggs that hatch (King'ori, 2011; Taplah *et al.,* 2018). Fertility and hatchability are affected by genetic and non-genetic factors. For a particular variety, non-genetic factors have a greater influence on fertility and hatchability in production (King'ori, 2011), Including management of the breeder, egg quality, and incubation processes. The management of breeding stock includes genetic selection (Drouilhet *et al.,* 2014), age of the breeders (King'ori, 2011), season and feeding (King'ori, 2011), breeding system and breeding technology (Weis *et al.,* 2011) and egg quality (Hester, 2017). The objective of the following review was to provide advanced information about the factors that affect fertility and hatchability of duck eggs and ways to improve them.

Fertility

Fertility can be affected by the quality of broodstock, male:female ratio, environmental temperature, storage time and housing systems. Brillard (2003) stated that fertility depends on the ability of females to ovulate, store sperm and provide an appropriate environment for the formation and development of the egg. Likewise, Brillard (2003) and Mohan *et al.* (2018) reported that the quality and quantity of semen deposited by the male were important to obtain good fertility. However, there are differences between duck breeds.

Muscovy ducks age influences the reproductive characteristics of both females and males (Yakubu, 2013). For flocks from the same breed, differences in fertility have been reported for different batches of eggs. In Muscovy ducks, the highest fertility is obtained at the peak of posture, compared to the moments before or after the peak of posture (Nickolova, 2005). Genitalia of waterfowl is more complicated than chickens, making the fertility issue more prevalent in ducks than in chickens, as well as the issue of sexual dimorphism in body size in some breeds of ducks and crossbreeding (Yakubu *et al.,* 2015).

The male:female ratio in all poultry species plays an important role in achieving greater fertility. For Muscovy ducks a ratio of one drake to five ducks is used according to Idahor *et al.* (2015) and one to four according to Nickolova (2005) and Alonso-Alvarez (2006). In Muscovy ducks, a ratio of one to six has been used satisfactorily where there is a shortage of males (Banerjee, 2013). However, larger sexual ratios of 1:4.3 to 1:10 cause a reduced egg fertility from 75.9% down to 49.6% (Nickolova, 2005), which corresponded with a loss in fertility from 97.09% to 93.41% when the sex ratio was increased from 1:5 to 1:8. In Khaki Campbell ducks, Giri *et al.* (2014) studies the ideal mating sex ratio for production of fertile eggs under intensive rearing and concluded that a 1:5 sex mating ratio was ideal for better fertility and hatchability.

https://doi.org/10.1017/S0043933919000060 Published online by Cambridge University Press

An increase of the environmental temperature above the optimal ranges of thermal comfort in poultry affects productive performance of breeders. According to Chowdhury *et al.* (2004), heat stress causes a decrease in the number of germ cells, ovular release, fertilisation and survival capacity of the embryo and abnormalities in sperm of the Northern Pintail duck (Penfold *et al.,* 2000). The latter decreases deposition of sperm in the host gland in the reproductive system of female (Brillard, 2003; Abd El-Hack *et al.,* 2018). Fertility was significantly higher at months from January to May as compared to June to August months and was significantly lowered by 9.70%, 12.72% and 14.29% in Summer as compared to Autumn, Winter and Spring season, respectively (Awad, 2013)

The storage time before the incubation of Pekin duck eggs has a significant effect on their fertility, hatchability and early embryo mortality (Waehner *et al.,* 2015). Reyna and Burggren (2017) reported a decrease in the fertility of duck eggs stored for more than six days from laying to incubation.

Housing system influences the fertility of ducks, allows mass mating and provides access to swimming, which significantly increases the fertility rate as ducks are waterfowl by nature and prefer to mate in water (Ojewola, 2006). However, egg weight linearly improves fertility; in Khaki Campbell ducks, the egg weight increased from 60 g to 75 g when they had water access, which positively affected egg fertility (Giri *et al.,* 2014).

Hatchability

Factors attributed to the breeding birds, such as genetic selection, management and feeding, handling and storage of eggs and conditions inside the incubator can influence the hatchability of duck eggs. Archer *et al.* (2017) and Ramli *et al.* (2017) stated that the temperature, relative humidity, ventilation and turning of the eggs throughout incubation and hatching were environmental factors that can modify hatchability.

BREEDER MANAGEMENT

Breeders must use breeds with good genetic characteristics at optimal ages to obtain high fertility, hatchability and growth performance of progeny. The time of year and the feeding conditions play an important role in the development of the embryo, before and after incubation.

Genetic characteristics can influence the productive indicators of different breeds of ducks in varying ways. The Muscovy lineages, compared to the Pekin duck, show higher mortality due to adhesion to the egg shell and for normal ducklings, however, Pekin duck eggs have better hatchability than Muscovy eggs (Rashid *et al.,* 2009). The eggs of the Muscovy such have shown less than 22.7% of hatchability (Ali *et al.,* 1989). The fertility and hatchability of laying Brown Tsaiya ducks were increased when a genetic selection was made and one artificial insemination per week was combined with Muscovy semen (Cheng *et al.,* 2002).

AGE

The age of female ducks has more influence on fertility than age of males (Brommer and Rattiste, 2008). After 24 days of incubation, the relative weight of the yolk sac, with respect to total egg weight, increased in hens at 36 weeks compared with Pekin ducks at 26 and 31 weeks of age (Applegate *et al.,* 1998). On the other hand, increasing the age of the female decreases the functioning of the sperm storage tubules, leading to problems in the reproductive system which influences egg quality.

https://doi.org/10.1017/S0043933919000060 Published online by Cambridge University Press

In addition, the age of ducks affects the external and internal quality of eggs, influencing both hatchability and fertility. The ratio between yolk and albumin decreased with increasing age of the birds (Peebles *et al.,* 2001). Larger eggs have comparatively less shell area per unit of interior weight than smaller ones, therefore, those from older broodstock have less weight when calculated in grams and more in percentage than the eggs of younger broodstock (Onasanya *et al.,* 2013). Age affects the deposition of calcium and minerals in the shell (Onbaşılar *et al.,* 2014). The shell allows gaseous exchange and loss of moisture from the egg, and poor shell quality results in excessive loss of relative humidity during incubation (Peebles *et al.,* 2001).

SEASON

The season when laying takes place affects fertility of ages and hatchability of ducklings, due to differences in temperature, wind speed, rainfall and relative humidity (Buhr, 1995; Farghly *et al.,* 2018b; Li *et al.,* 2018). Chowdhury *et al.* (2004) observed a high significance between the month and hatchability of Pekin duck eggs, where the hatchability in January was 59.54 ± 0.79%, but in July it decreased to 48.27 ± 0.79%. The best hatchability of Pekin duck eggs is observed during the winter (57.68 ± 0.59%), but during the summer it decreases to 54.14 ± 0.59%, and the lowest values are found in the monsoon or rainy season (49.13 ± 0.59%) (Chowdhury *et al.*, 2004). Hatchability of duck eggs showed the highest value in March (65.21%), but the lowest value (34.96%) was observed in August (Awad, 2013). Fertility and hatchability were lower in the summer months compared to other seasons due to high environmental temperature that causes a decrease in the reproductive efficiency in both male and female ducks.

NUTRITION

During egg formation, essential nutrients are deposited for the development of embryo during the incubation period and form reserves for the developing embryo until the first meal becomes available to the young birds (El-Kholy *et al.,* 2019; Saeed *et al.,* 2019). The nutrients are used for tissue formation, heat generation and muscle activity, and are stored in both albumin and yolk (Uni *et al.,* 2012; Onbaşılar *et al.,* 2014). Egg albumin from eggs from Pekin ducks is mainly water (85.7 to 88.1%) containing 10.9% to 13.1% water-soluble proteins and represents from 53.0% to 55.6% of the total egg content. Although the yolk only represents 31.3% to 33.9% of the egg content and 41.7% to 44.1% is water, it has higher protein reserves (17.3% to 17.8%), lipids (35% to 38.4%) and minerals (Yair and Uni, 2011).

Nutrient deficiencies impede the proper development of embryos, reduce hatchability and increase embryonic death, in addition to causing disorders of the musculoskeletal system, immune system and cardiovascular system (Uni *et al.,* 2012). The nutrients in the egg determine the weight of the duckling at birth, body size and hatchability. Therefore, feeding ducks correctly can increase deposition of nutrients in the egg, without causing any change in weight of the broodstock, which are important factors to consider to avoid low quality ejaculation and ovulation and, at an extreme, early ovarian and testicular regression (Brillard, 2003).

Certain nutrients and food ingredients have adverse effects on the quality and quantity of eggs produced by ducks. Anti-nutritional factors, such as gossypol from cotton seed meal, cause infertility in males manifested as immobility of sperm caused by damage to the mitochondria located in the tail of sperm, and extensive damage to the germinal epithelium (Randel *et al.,* 1992). Increased consumption of gossypol causes increased permeability of the yolk sac membrane which leads to excessive pigmentation of the albumin, known as 'pink disease' or white albumen which is mottled or gummy. It is

https://doi.org/10.1017/S0043933919000060 Published online by Cambridge University Press

necessary for producers to constantly update feeding standards and include new recommendations to ensure the quantity and quality of food necessary to the layer ducks.

Egg factors

Eggs are normally fertile four days after the deposition of the semen inside the female, and in this period all necessary components for the development of embryo until the birth should be present in the egg. Under normal circumstances, infertility is due to different factors related to the management of the broodstock and environmental conditions, since these are related to modifications in the physical and chemical characteristics that reduce embryonic development and hatching of good quality ducklings (Narushin and Romanov, 2002).

The management of broodstock plays an important role to ensure a good hatchability, breed used, season, health, and nutrition, also size, weight and quality of egg, in addition to the duration and storage conditions (Wilson, 1991; Onasanya *et al.,* 2013). The quality of eggs for incubation is determined by both external and internal characteristics, however, there are other factors that prevent hatching, such as the inheritance of lethal genes, lack of nutrients and sudden variations in incubation conditions (King'ori, 2011).

EGG QUALITY

Egg quality has a significant effect on hatchability, as conditions of the microenvironment during the storage process and early incubation modify both the external and internal quality of eggs (Narushin and Romanov, 2002). External characteristics such as weight, index form (described as maximum breadth to length ratio), shell thickness, porosity, surface quality of the shell and resistance to breakage have been measured (Toro *et al.,* 2015). Internal quality is determined by the height of the dense albumen and the yolk, in addition to Haugh Units. Haugh units are calculated as:

$$HU=100 \log (H+7.57-1.7W^{0.37})$$

where HU=Haugh units, H=albumin height and W=egg weight (Toro *et al.,* 2015).

Heavier eggs are more likely to hatch than those of low weight; however, there are contradictions about the weight of eggs that are not within the average values for incubation (King'ori, 2011). An increase in weight, shell thickness and internal contents of eggs leads to a higher total weight, which in turn reflects more reserves of nutrients and energy (Toro *et al.,* 2015). Light weight eggs have relatively shorter incubation times (Weis *et al.,* 2011) and the weight of hatchlings is higher from large eggs. It is important to note that, during incubation, larger embryos produce a greater amount of heat, which requires good ventilation to maintain the correct temperature (Jibrin *et al.,* 2011).

Likewise, egg size affects hatchability. Wilson (1991) and Weis *et al.* (2011) found that medium-sized eggs from Muscovy ducks had better hatchability than small eggs. In this sense Demirel and Kırıkçi (2009), found a greater increase in yolk compared to albumin as the egg size increased, which could be a major influence.

With longer storage time, HU decreases and pH of the albumin increases, apparently due to loss of water (Martínez *et al.,* 2014). Elibol *et al.* (2002) stated that the low pH in fresh eggs did not affect fertility and hatchability, however, other authors indicated the opposite, and perhaps this difference may be due to different flock age (Benton and Brake, 1994).

https://doi.org/10.1017/S0043933919000060 Published online by Cambridge University Press

EMBRYO SURVIVAL

According to Kuurman *et al.* (2003), mortality describes a curved, diphasic Weibull distribution, due to the peaks of mortality in the first and last third of incubation. The survival of the embryo does not only depend on environmental conditions before and during incubation, factors related to the genotype of the dam and the sire, mainly the dam, affects the survival of the duckling within the egg. Chromosomal aberrations and lethal genes acquired from the sire and dam can cause high percentages of early embryonic mortality (Liptoi and Hidas, 2006). The stage of development decreases the heritability of the susceptibility to embryonic death, from 0.09 for early mortality to 0.05 for late mortality, based on the parent component and from 0.25 to 0.18 based on the dam component (Beaumont *et al.*, 1997). In this sense, Bennewitz *et al.* (2007) reported low direct heritability of hatchability, based on linear and threshold models respectively, with hatching capacity treated as a trait of the dam.

EGG STORAGE

The microenvironment conditions during storage, storage time and position of the eggs on the incubation tray are important to guarantee a profitable production of ducklings, especially for production in the tropics and subtropics. To avoid the effect of heat stress on the embryo during storage, temperature should be 13°C (55°F) and never higher than 24°C, and relative humidity should be less than 75% but higher than 40% (Buhr, 1995). Eggs stored under refrigeration should be placed for 4 h at room temperature before incubating (Onbaşılar *et al.,* 2007). Onbaşılar *et al.* (2007) reported that hatchability was lower and early embryonic death rate was the highest in eggs stored for 11 d. Saha *et al.* (1992) reported that hatchability of Khaki Campbell duck eggs was greatest in eggs stored for 3 d compared to 7 d pre-incubation.

It is advisable not to store eggs for more than two weeks, as hatchability decreases significantly after 14 d. Researcher have shown that storage period is related to both early and late embryonic deaths (Onbaşılar *et al.,* 2007). The hatchability of total and fertile eggs decreased with increasing storage period, since early and late embryonic death was higher due to loss of water and degradation of albumin during storage.

However, the optimal storage period is not fixed. It varies according to the age of the batch and the strain and species, due to the differences in the quality of the albumin. Keeping eggs with the small end down and the large end upwards can result in more successful hatching than when doing the opposite, perhaps because the incorrect orientation of the head of the embryo towards the larger pole prevents correct elimination of water. In this sense, Bauer *et al.* (1990) reported a 17% decrease in the hatchability of embryos of broilers located with the small pole up. Apparently, this happens because the embryo cannot find the air chamber that is located in the round pole, especially at the end of incubation (Bauer *et al.,* 1990).

Conclusions

Success of the duck sector at small or large scale depends on a regular supply of one-day-old chicks. The production of ducklings is influenced by the fertility and hatchability of the eggs, which are important economic factors representing the major components of reproductive performance. Fertility and hatchability are most sensitive to the environmental and genetic influences. There are many factors related to producers, breeders and environmental conditions during incubation that can influence these parameters, including managerial, nutritional and genetic factors as well as eggs quality and incubator factors. There are different methods used to improve the

https://doi.org/10.1017/S0043933919000060 Published online by Cambridge University Press

hatchability percentage of waterfowl eggs - cooling and periodic spraying with water are some of the most commonly used. Dipping eggs and *in ovo* injection with nutrients such as ascorbic acid and vitamin E during the incubation period is one of the tools used to improve hatchability percentage.

References

ABD EL-HACK, M.E., ALAGAWANY, M. and NORELDIN, A.E. (2018) Managerial and nutritional trends to mitigate heat stress risks in poultry farms, in: *The Handbook of Environmental Chemistry*, pp 1-14 (Springer, Berlin, Heidelberg. Germany).

ALI, M.A., RAHAMAN, M.A., WAHID, M.A. and SINGH, T.K. (1989) The hatchability of Muscovy duck eggs under natural and artificial incubation. *Bangladesh Journal Animal Science* **8 (1)**: 13.

ALONSO-ALVAREZ, C. (2006) Manipulation of primary sex-ratio: an updated review. *Avian and Poultry Biology Reviews* **17 (1)**: 1-20.

APPLEGATE, T.J., HARPER, D. and LILBURN, M.S. (1998) Effect of hen production age on egg composition and embryo development in commercial Pekin ducks. *Poultry Science* **77 (11)**: 1608-1612.

ARCHER, G.S., JEFFREY, D. and TUCKER, Z. (2017) Effect of the combination of white and red LED lighting during incubation on layer, broiler, and Pekin duck hatchability. *Poultry Science* **96 (8)**: 2670-2675.

AWAD, A.L. (2013) Field study on hatching traits of duck eggs under Egyptian environmental conditions. *Egyptian Poultry Science Journal* **33**: 849-863.

BANERJEE, S. (2013) Morphological traits of duck and geese breeds of West Bengal, India. *Animal Genetic Resources/Resources Génétiques Animales/Recursos Genéticos Animales* **52**: 1-16.

BAUER, F., TULLETT, S.G. and WILSON, H.R. (1990) Effects of setting eggs small end up on hatchability and posthatching performance of broilers. *British Poultry Science* **31 (4)**: 715-724.

BEAUMONT, C., MILLET, N., LE BIHAN-DUVAL, E., KIPI, A. and DUPUY, V. (1997) Genetic parameters of survival to the different stages of embryonic death in laying hens. *Poultry Science* **76 (9)**: 1193-1196.

BENNEWITZ, J., MORGADES, O., PREISINGER, R., THALLER, G. and KALM, E. (2007) Variance component and breeding value estimation for reproductive traits in laying hens using a Bayesian threshold model. *Poultry Science* **86 (5)**: 823-828.

BENTON, C.E. and BRAKE, J. (1994) The effect of the presence of an embryo on albumen height and pH during preincubation storage and incubation. *Poultry Science* **73 (Suppl 1)**: 38.

BRILLARD, J.P. (2003) Practical aspects of fertility in poultry. *World's Poultry Science Journal* **59 (4)**: 441-446.

BROMMER, J.E. and RATTISTE, K. (2008) "Hidden" reproductive conflict between mates in a wild bird population. *Evolution: International Journal of Organic Evolution* **62 (9)**: 2326-2333.

BUHR, R.J. (1995) Incubation relative humidity effects on allantoic fluid volume and hatchability. *Poultry Science* **74 (5)**: 874-884.

CHENG, Y.S., ROUVIER, R., POIVEY, J.P., TAI, J.J.L., TAI, C. and HUANG, S.C. (2002) Selection responses for the number of fertile eggs of the Brown Tsaiya duck (*Anas platyrhynchos*) after a single artificial insemination with pooled Muscovy (*Cairina moschata*) semen. *Genetics Selection Evolution* **34 (5)**: 597.

CHOWDHURY, M.M.I., ASHRAF, A., MONDAL, S.P., MONDOL, N.M.A.A.M. and HASAN, M.M. (2004) Effect of season on the hatchability of duck eggs. *International Journal of Poultry Science* **3 (6)**: 419-421.

DEMIREL, S. and KIRIKÇI, K. (2009) Effect of different egg storage times on some egg quality characteristics and hatchability of pheasants (*Phasianus colchicus*). *Poultry Science* **88 (2)**: 440-444.

DROUILHET, L., BASSO, B., BERNADET, M.D., CORNUEZ, A., BODIN, L., DAVID, I. and MARIE-ETANCELIN, C. (2014) Improving residual feed intake of mule progeny of Muscovy ducks: genetic parameters and responses to selection with emphasis on carcass composition and fatty liver quality. *Journal of Animal Science* **92 (10)**: 4287-4296.

ELIBOL, O., PEAK, S.D. and BRAKE, J. (2002) Effect of flock age, length of egg storage, and frequency of turning during storage on hatchability of broiler hatching eggs. *Poultry Science* **81 (7)**: 945-950.

EL-KHOLY, M.S., IBRAHIM, Z.A., EL-MEKKAWY, M.M. and ALAGAWANY, M. (2019) Influence of in-ovo administration of some water-soluble vitamins on hatchability traits, growth, carcass traits and blood chemistry of Japanese quails. *Annals of Animal Science* **19**: 97-111.

ERICSON, P.G., QU, Y., BLOM, M.P., JOHANSSON, U.S. and IRESTEDT, M. (2017) A genomic perspective of the pink-headed duck *Rhodonessa caryophyllacea* suggests a long history of low effective population size. *Scientific Reports* **7 (1)**: 16853.

https://doi.org/10.1017/S0043933919000060 Published online by Cambridge University Press

FARGHLY, M.F.A., ABD EL-HACK, M.E., ALAGAWANY, M., SAADELDIN, I.M. and SWELUM, A. A. (2018a) Ameliorating deleterious effects of heat stress on growing Muscovy ducklings using feed withdrawal and cold water. *Poultry Science* **98**: 251-259.

FARGHLY, M.F.A., ABD EL-HACK, M.E., ALAGAWANY, M., SAADELDIN, I.M. and SWELUM, A. A. (2018b) Wet feed and cold water as heat stress modulators in growing Muscovy ducklings. *Poultry Science* **97**: 1588-1594.

GIRI, S.C., KUMAR, P., JAISWARA, R., BAIS, R.K.S., SAHOO, S.K., SASTRY, K.V.H. and SARAN, S. (2014) Ideal mating sex ratio in duck houses for optimum production of fertile eggs. *Indian Journal of Poultry Science* **49 (1)**: 106-107.

HESTER, P.Y. (2017) Effects of temperature and storage conditions on eggs, in: *Egg innovations and strategies for improvements*, Section III Food Safety, Chapter 12, pp. 125-134.

IDAHOR, K.O., AKINOLA, L.A.F. and CHIA, S.S. (2015) Egg colour, weight and shape: possible indices in the predetermination of duckling sex. *Journal of Recent Advances in Agriculture* **3 (1)**: 337-344.

JIBRIN, M.M., IDIKE, F.I., AHMAD, K. and IBRAHIM, U. (2011) Modelling incubation temperature: the effects of incubator design, embryonic development and egg size. *Journal of Agricultural Engineering and Technology* **19 (1)**: 46-59.

KING'ORI, A.M. (2011) Review of the factors that influence egg fertility and hatchability in poultry. *International Journal of Poultry Science* **10 (6)**: 483-492.

KUURMAN, W.W., BAILEY, B.A., KOOPS, W.J. and GROSSMAN, M. (2003) A model for failure of a chicken embryo to survive incubation. *Poultry Science* **82 (2)**: 214-222.

LI, X., WEN, Y., ZHANG, J., LIU, L., JIN, L., YAN, T. and WANG, Y. (2018) The effect of low-temperature event on the survival and growth of *Juglans mandshurica* seedlings within forest gaps. *Journal of Forestry Research* **29 (4)**: 943-951.

LIPTOI, K. and HIDAS, A. (2006) Investigation of possible genetic background of early embryonic mortality in poultry. *World's Poultry Science Journal* **62 (2)**: 326-337.

MARTÍNEZ, Y., GUERRA, L.D., RODRÍGUEZ, R. and BETANCUR, C.A. (2014) Effect of pre-incubation storage conditions on embryonic development and chick quality of Camperos breeders. *International Journal of Animal and Veterinary Advances* **6 (3)**: 108-111.

MOHAN, J., SHARMA, S., KOLLURI, G. and DHAMA, K. (2018) History of artificial insemination in poultry, its components and significance. *World's Poultry Science Journal* **74 (3)**: 1-14.

NARUSHIN, V.A. and ROMANOV, M.N. (2002) Egg physical characteristics and hatchability. *World's Poultry Science Journal* **58 (3)**: 297-303.

NICKOLOVA, M. (2005) Effect of the sex ratio on the egg fertility of Muscovy duck (*Cairina moshcata*). *Journal of Central European Agriculture* **5 (4)**: 367-372.

OJEWOLA, F.E.G. (2006) Effect of Management Systems on Semen Quality of Muscovy Drakes. *International Journal of Poultry Science* **5 (5)**: 482-484.

ONASANYA, G.O. and IKEOBI, C.O.N. (2013) Egg physical traits, performance, fertility and hatchability in exotic and Nigerian indigenous chickens. *Standard Research Journal of Agricultural Sciences* **1 (1)**: 1-8.

ONBAŞILAR, E., ERDEM, E., HACAN, Ö. and YALÇIN, S. (2014) Effects of breeder age on mineral contents and weight of yolk sac, embryo development, and hatchability in Pekin ducks. *Poultry Science* **93 (2)**: 473-478.

ONBAŞILAR, E., POYRAZ, Ö. and ERDEM, E. (2007) Effects of egg storage period on hatching egg quality, hatchability, chick quality and relative growth in Pekin ducks. *Archiv für Geflügelkunde* **71 (4)**: 187-191.

PEEBLES, E.D., DOYLE, S.M., ZUMWALT, C.D., GERARD, P.D., LATOUR, M.A., BOYLE, C.R. and SMITH, T.W. (2001) Breeder age influences embryogenesis in broiler hatching eggs. *Poultry Science* **80 (3)**: 272-277.

PENFOLD, L.M., WILDT, D.E., HERZOG, T.L., LYNCH, W., WARE, L., DERRICKSON, S.E. and MONFORT, S.L. (2000) Seasonal patterns of LH, testosterone and semen quality in the Northern pintail duck (*Anas acuta*). *Reproduction, Fertility and Development* **12 (4)**: 229-235.

RAMLI, M.B., WAHAB, M.S., ZAIN, B.A., RAUS, A.A., RAJA, P. and PAHAT, B. (2017) Effect of incubation temperature on ikta's quail breed with new rolling mechanism system. *Journal of Mechanical Engineering* **SI 4 (3)**: 78-88.

RANDEL, R.D., CHASE, C.C. (Jr) and WYSE, S.J. (1992) Effects of gossypol and cottonseed products on reproduction of mammals. *Journal of Animal Science* **70 (5)**: 1628-1638.

RASHID, M., KAWSAR, M., MIAH, M. and HOWLIDER, M. (2009) Fertility and hatchability of Pekin and Muscovy duck eggs and performance of their ducklings. *Progressive Agriculture* **20 (1-2)**: 93-98.

REYNA, K.S. and BURGGREN, W.W. (2017) Altered embryonic development in northern bobwhite quail (*Colinus virginianus*) induced by pre-incubation oscillatory thermal stresses mimicking global warming predictions. *PloS One* **12 (9)**: e0184670.

https://doi.org/10.1017/S0043933919000060 Published online by Cambridge University Press

SAEED, M., BABAZADEH, D., NAVEED, M., ALAGAWANY, M., ABD EL-HACK, M.E., ARAIN, M. A., TIWARI, R., SACHAN, S., KARTHIK, K., DHAMA, K., ELNESR, S.S. and CHAO, S. (2019) In ovo delivery of various biological supplements, vaccines and drugs in poultry: current knowledge. *Journal of Science of Food and Agriculture* **99**: 3727-3739.

SAHA, S.K., CHOWDHURY, S.D. and HAMID, M.A. (1992) A study on the incubation of indigenous (Desi), Khaki Campbell and crossbred (Indian Runner x Khaki Campbell, F1) duck eggs under two pre-incubation holding periods. *Asian-Australasian Journal of Animal Sciences* **5 (3)**: 541.

TAPLAH JR, A.J., SUMINISTRADO, D.C., AMONGO, R.M.C., PARAS, F.O. (Jr), ELAURIA, J.C. and TORKPAH, D.P. (2018) Economic analysis of duck eggs incubation using hot spring as heat source. *Journal of Development and Agricultural Economics* **10 (2)**: 38-44.

TORO, D.M., AGUILAR, Y.M., BERTOT, R.R., HURTADO, C.B. and NAVA, O.R. (2015) Effect of dietary supplementation with *Morinda citrifolia* on productivity and egg quality of laying hens. *Revista Ciencia y Agricultura* **12 (2)**: 7-12.

UNI, Z., YADGARY, L. and YAIR, R. (2012) Nutritional limitations during poultry embryonic development. *Journal of Applied Poultry Research* **21 (1)**: 175-184.

WAEHNER, M., PINGEL, H. and HAIDONG, S. (2015) Effect of prolonged storage of eggs of Pekin ducks with periodical warming on internal egg quality and hatchability. Proceedings of the 4th International Congress on New Perspectives and Challenges of Sustainable Livestock Production, October 7-9, 2015, Belgrade, Serbia, pp: 140-144.

WEIS, J., HRNČÁR, C., PÁL, G., BARAŇSKA, B., BUJKO, J. and MALÍKOVÁ, L. (2011) Effect of the egg size on egg loses and hatchability of the Muscovy duck. *Scientific Papers Animal Science and Biotechnologies* **44 (1)**: 354-356.

WIDIYANINGRUM, P., LISDIANA, L. and UTAMI, N. (2016) Egg production and hatchability of local ducks under semi intensive vs extensive managements. *Journal of the Indonesian Tropical Animal Agriculture* **41 (2)**: 77-82.

WILSON, H.R. (1991) Interrelationships of egg size, chick size, post hatching growth and hatchability. *World's Poultry Science Journal* **47 (1)**: 5-20.

YAIR, R. and UNI, Z. (2011) Content and uptake of minerals in the yolk of broiler embryos during incubation and effect of nutrient enrichment. *Poultry Science* **90 (7)**: 1523-1531.

YAKUBU, A. (2013) Characterization of the local Muscovy duck in Nigeria and its potential for egg and meat production. *World's Poultry Science Journal* **69 (4)**: 931-938.

YAKUBU, A., MUHAMMED, M.M., ARI, M.M., MUSA-AZARA, I.S. and OMEJE, J.N. (2015) Correlation and path coefficient analysis of body weight and morphometric traits of two exotic genetic groups of ducks in Nigeria. *Bangladesh Journal of Animal Science* **44 (1)**: 1-9.

https://doi.org/10.1017/S0043933919000060 Published online by Cambridge University Press

https://doi.org/10.1017/S0043933919000060 Published online by Cambridge University Press