Proceedings of the Royal Society of Edinburgh, **155**, 1051–1060, *2025*
DOI:10.1017/prm.2023.126

The lower bounds of non-real eigenvalues for singular indefinite Sturm–Liouville problems

Fu Sun
School of Statistics and Data Science, Qufu Normal University, Qufu 273165, People's Republic of China (sfmath@163.com)

(Received 24 February 2023; accepted 17 November 2023)

The present paper deals with the non-real eigenvalues for singular indefinite Sturm–Liouville problems. The lower bounds on non-real eigenvalues for this singular problem associated with a special separated boundary condition are obtained.

Keywords: Sturm–Liouville problem; non-real eigenvalue; indefinite; lower bounds

2020 *Mathematics Subject Classification:* 34B24; 34L15; 47B50

1. Introduction

In the present paper, we consider the eigenvalue problem

$$-(py')' + qy = \lambda wy, \quad [y, u_{-1}](-1) = 0 = [y, u_1](1), \tag{1.1}$$

where the functions p, q, w are real-valued and w changes sign on $(-1, 1)$ and u_{-1}, u_1 are the principal solutions of differential expression

$$-(py')' + qy = \lambda wy \tag{1.2}$$

at -1, 1 for $\lambda = 0$, respectively. Such a problem is called *indefinite* and the indefinite nature, that non-real spectral points may appear, was noticed by Haupt [**11**] and Richardson [**19**] at the beginning of the last century and has attracted a lot of attention in the recent years, see [**1**, **7**, **8**, **10**, **14**].

A priori bounds on non-real eigenvalues for indefinite Sturm–Liouville problems were raised in [**15**] by Mingarelli and stressed by Kong *et al.* [**13**]. Recently, the regular indefinite case of this problem was solved by Qi *et al.* in [**2**, **17**, **18**, **21**]. For the singular indefinite Sturm–Liouville problems with limit-point type endpoints

$$(Af)(x) := \operatorname{sgn}(x)(-f''(x) + V(x)f(x)) = \lambda f(x), \quad x \in \mathbb{R}, \tag{1.3}$$

the authors in [**3**] provided sufficient conditions for the existence of non-real eigenvalues. The explicit bounds on the non-real eigenvalues of (1.3) were obtained in [**4**]. In [**5**, **6**], the authors not only estimated the absolute values of the non-real eigenvalues in terms of the L^1-norm of the continuous potential, but also obtained the bounds on the imaginary parts and absolute values of these eigenvalues in terms

© The Author(s), 2023. Published by Cambridge University Press on behalf of The Royal Society of Edinburgh

https://doi.org/10.1017/prm.2023.126 Published online by Cambridge University Press

of the L^1-norm of the potential and its negative part. Recently in [**20**], the authors solved the estimates of absolute values on the non-real eigenvalues for the singular indefinite Sturm–Liouville eigenvalue problems with limit-circle type non-oscillation endpoints associated with a special self-adjoint boundary condition.

In the present paper, we will focus on the singular indefinite Sturm–Liouville eigenvalue problems with self-adjoint boundary conditions associated with principal solution at endpoints. The lower bounds of this eigenvalue problem are obtained under the some conditions. This paper is organized as follows: the preliminary knowledge and the main results, theorems 2.2 and 2.3, are stated in § 2 and their proofs are given in § 3.

2. Preliminary knowledge and main results

In this section, we give some basic knowledge for the singular differential equation (1.2) under the standard conditions that p, q, w are real-valued functions satisfying

$$p > 0, |w| > 0 \text{ a.e. on } (-1,1),\ \frac{1}{p}, w, q \in L^1_{\mathrm{loc}}(-1,1), \int_{-1}^{1} \left(\left| \frac{1}{p} \right| + |q| + |w| \right) = \infty \tag{2.1}$$

and w changes its sign on $(-1, 1)$ in the meaning that

$$\mathrm{mes}\{x : w(x) > 0\} > 0, \quad \mathrm{mes}\{x : w(x) < 0\} > 0.$$

We first introduce some concepts. For fixed $\lambda \in \mathbb{R}$, a real solution $u(x)$ of (1.2) is called a *principal solution* at 1 if there exists $c \in (-1, 1)$ such that $u(x) \neq 0$, $x \in (c, 1)$, $\int_c^1 1/(pu^2) = \infty$. A real solution $v(x)$ of (1.2) is called a *non-principal solution* at 1 if there exists $c \in (-1, 1)$ such that $v(x) \neq 0$, $x \in (c, 1)$, $\int_c^1 1/(pv^2) < \infty$. If u and v are principal and non-principal solutions at 1, respectively, then $u(x)/v(x) \to 0$ as $x \to 1$. (cf. [**16**] and [**22**, Theorem 6.2.2]). In order to give the asymptotic behaviours of eigenfunctions at the endpoints, we assume that

$$\Upsilon(t) := \sup_{-1<x<1} \left| \frac{1}{p(t)} \int_t^x q(s)\mathrm{d}s \right| \in L^1(-1,1), \quad \int_c^x \frac{1}{p(t)}\mathrm{d}t \in L^2_{|w|}(-1,1) \tag{2.2}$$

for some (and hence for all) $c \in (-1, 1)$. Throughout this paper, the functions p, q, w always satisfy (2.1) and (2.2). Then

LEMMA 2.1. *[**20**, Lemma 2.3] Assume that (2.2) holds, $u_1(\cdot)$ is a principal solution of (1.2) at 1 for $\lambda = 0$. Let y be an eigenfunction of (1.1) corresponding to the eigenvalue λ. If either $\int_c^1 1/p(t)dt = \infty$ or $\int_c^1 1/p(t)dt < \infty$, $q \in L^1(-1, 1)$, then y is bounded and*

$$[y, u_1](1) = 0 \Leftrightarrow (py')(x)y(x) \to 0 \text{ as } \quad x \to 1. \tag{2.3}$$

The similar conclusion holds for $x \to -1$.

https://doi.org/10.1017/prm.2023.126 Published online by Cambridge University Press

The operator S associated with the right-definite problem

$$\begin{cases} -(py')'(x)+q(x)y(x)=\lambda|w(x)|y(x), \\ [y,u_{-1}](-1)=0,\ [y,u_1](1)=0 \end{cases} \tag{2.4}$$

is defined as $Sy=\frac{1}{|w|}\tau y$ for $y\in D(S)$, where $\tau y:=-(py')'(x)+q(x)y(x)$,

$$D(S)=\{y\in L^2_{|w|}(-1,1):\ y,py'\in AC_{loc}(-1,1),\tau y/|w|\in L^2_{|w|}(-1,1),\ \mathcal{B}y=0\}$$

and $\mathcal{B}y=0:=\ [y,\,u_{-1}](-1)=0=[y,\,u_1](1)$. It follows from [**12**] and [**22**, Theorem 10.6.2, p.195] that the operator S is self-adjoint in the Hilbert space $(L^2_{|w|},\,(\cdot,\,\cdot)_{|w|})$ and it has discrete spectrum consisting of an infinite number of eigenvalues $\{\mu_n:n\in\mathbb{N}:=\{1,\,2,\,\cdots\}\}$, which are all real, unbounded from above and bounded from below, i.e. $-\infty<\mu_1<\mu_2<\mu_3<\cdots\to+\infty$.

Let $K=(L^2_{|w|}(-1,\,1),\,[\cdot,\,\cdot]_w)$ be the Krein space with the indefinite inner product $[f,\,g]_w=\int_{-1}^{1}w(x)f(x)\overline{g(x)}\,\mathrm{d}x$, $f,\,g\in L^2_{|w|}(-1,\,1)$ and $J=\operatorname{sgn}w$ the fundamental symmetry operator. The operator T in K is defined as

$$Ty=\frac{1}{w}\tau y,\quad y\in D(T)=D(S).$$

Then $S=JT$, $[Tf,\,g]_w=(Sf,\,g)_{|w|}$, $f,\,g\in D(T)$ and T is a self-adjoint operator in K [**3**, **7**, **9**]. In the following, we denote the resolvent set of S by $\rho(S)$.

Now, we state the lower bound result on T.

THEOREM 2.2. *Let T and S be defined as above. Suppose that $0\in\rho(S)$ and S^{-1} is compact. Let $\mu^+:=\min\ \sigma(S)\cap(0,\,\infty)$, $\mu^-:=\min\ \{|\lambda|:\lambda\in\sigma(S)\cap(-\infty,\,0)\}$, where $\min\ \emptyset:=\infty$. Then for each eigenvalue λ of T we have $|\lambda|\geqslant\min\ \{\mu^+,\,\mu^-\}$.*

Moreover, if λ corresponds to an eigenvector ϕ of T with $[\phi,\,\phi]_w=0$, then the following, in general stronger, estimate holds $|\lambda|^2\geqslant-\mu^+\mu^-$.

In order to give another result of the lower bound on T, we assume that $q_-(x)=\max\{0,\,-q(x)\}$ and for some $C,\ C_0,\ C_1,\ C_2>0,\ x\in(-1,\,1)$

$$\left|\frac{1-x}{\sqrt{p(x)}}\int_{-1}^{x}q_-(t)\mathrm{d}t\right|\leqslant C_0,\quad \left|\frac{x+1}{\sqrt{p(x)}}\int_{x}^{1}q_-(t)\mathrm{d}t\right|\leqslant C_1,$$

$$W(x)=\int_{-1}^{x}w(t)\mathrm{d}t,\quad \frac{W(x)}{\sqrt{p(x)}}\leqslant C_2,\quad C=\sqrt{2}C_2. \tag{2.5}$$

It is easy to verify that if $q\in L^2(-1,\,1)$, $p(x)=1-x^2$, $w(x)=x$, then (2.5) holds. Let

$$\gamma_t:=\min\left\{\int_{t_0}^{t_0+t}|w(x)|\mathrm{d}x:t\in(-1,1),\ t_0\in(-1,1)\right\},$$

$$\Delta_{w,1,n}=\|w\|_1+C\left(\sqrt{\Delta}+\sqrt{\Delta_n}\right),\quad \Delta=2\int_{-1}^{1}q_-(t)\mathrm{d}t+8\alpha^2,$$

$$\Delta_n=2\int_{-1}^{1}q_-(t)\mathrm{d}t+8\alpha^2+2|\mu_n|\|w\|_1,\quad \alpha=\frac{C_0+C_1}{2}. \tag{2.6}$$

https://doi.org/10.1017/prm.2023.126 Published online by Cambridge University Press

With this notation, we give the following result.

THEOREM 2.3. *Assume that λ and μ_n are the non-real eigenvalue of (1.1) and the nth eigenvalue of right-definite problem (2.4), respectively. Let $\mu_{h-1} < 0 < \mu_h$ for some positive integer $h \geqslant 2$, then the eigenvalue λ satisfies*

$$|\lambda|^2 \geqslant \frac{\mu_h \mu_{h-1}^2 \gamma_\delta}{16(\mu_h - \mu_{h-1})} \left(\sum_{n=1}^{h-1} \frac{\Delta_{w,1,n}^2}{\gamma_{\delta_n}} \right)^{-1}.$$

3. The proofs of theorems 2.2 and 2.3

In order to prove theorems 2.2 and 2.3, we prepare some lemmas. The following lemma is the estimates of $\|\sqrt{p}\phi'\|_2$, where ϕ is an eigenfunction of (1.1) corresponding to a non-real eigenvalue λ. That is $\mathcal{B}\phi = 0$ and

$$-(p\phi')' + q\phi = \lambda w\phi. \tag{3.1}$$

Since problem (1.1) is a linear system and ϕ is continuous, we can choose ϕ satisfies $\max\{|\phi(x)| : x \in [-1, 1]\} = 1$ in the following discussion.

LEMMA 3.1. *Let λ and ϕ be defined as above. Then*

$$\int_{-1}^{1} w|\phi|^2 = 0, \quad \int_{-1}^{1} p|\phi'|^2 \leqslant \Delta, \tag{3.2}$$

where Δ is given by (2.6).

Proof. It follows from $\mathcal{B}\phi = 0$ and lemma 2.1 that ϕ is bounded and satisfies $(p\phi'\overline{\phi})(x) \to 0$ as $x \to -1$ or 1. Multiplying both sides of (3.1) by $\overline{\phi}$ and integrating over the interval $[a, b]$, then

$$\int_{-1}^{1} p|\phi'|^2 + \int_{-1}^{1} q|\phi|^2 = \lambda \int_{-1}^{1} w|\phi|^2. \tag{3.3}$$

From $\operatorname{Im}\lambda \neq 0$ and (3.3) one sees that $\int_{-1}^{1} w|\phi|^2 = 0$, and hence

$$\int_{-1}^{1} p|\phi'|^2 + \int_{-1}^{1} q|\phi|^2 = 0. \tag{3.4}$$

Set

$$Q(x) = \int_{-1}^{x} q_-(t)\mathrm{d}t - \frac{x+1}{2} \int_{-1}^{1} q_-(t)\mathrm{d}t$$

Then one can verify that

$$Q(-1) = 0 = Q(1),\ Q'(x) = q_-(x) - \frac{1}{2}\int_{-1}^{1} q_-(t)\mathrm{d}t \text{ a.e. } x \in (-1, 1) \quad \text{and}$$

$$|Q(x)| \leqslant \left| \frac{1-x}{2} \int_{-1}^{x} q_-(t)\mathrm{d}t \right| + \left| \frac{x+1}{2} \int_{x}^{1} q_-(t)\mathrm{d}t \right| \leqslant \frac{\sqrt{p(x)}}{2}(C_0 + C_1) = \alpha\sqrt{p(x)}.$$

https://doi.org/10.1017/prm.2023.126 Published online by Cambridge University Press

As a result, this together with (3.4) and Cauchy–Schwarz inequality yields that

$$\begin{aligned}\int_{-1}^{1} q_-|\phi|^2 &= \int_{-1}^{1}\left(Q' + \frac{1}{2}\int_{-1}^{1} q_-\right)|\phi|^2 \\ &\leqslant \int_{-1}^{1} q_- - 2\,\mathrm{Re}\left(\int_{-1}^{1} Q\phi'\overline{\phi}\right) \\ &\leqslant \int_{-1}^{1} q_- + \frac{1}{2}\int_{-1}^{1} p|\phi'|^2 + 4\alpha^2. \end{aligned} \tag{3.5}$$

It follows from (3.4), (3.5) and $q = q_+ - q_-$, $q_\pm = \max\{0, \pm q\}$ that

$$\int_{-1}^{1} p|\phi'|^2 = -\int_{-1}^{1} q|\phi|^2 \leqslant \int_{-1}^{1} q_-|\phi|^2 \leqslant \int_{-1}^{1} q_- + \frac{1}{2}\int_{-1}^{1} p|\phi'|^2 + 4\alpha^2.$$

So the inequalities in (3.2) holds immediately. □

Similarly with the argument of lemma 3.1, we give the estimates of $\|\sqrt{p}\psi_n'\|_2$, where ψ_n is the eigenfunction that satisfies $\max\{|\psi_n(x)| : x \in [-1, 1]\} = 1$ corresponding to the nth eigenvalue μ_n of (2.4).

LEMMA 3.2. *Suppose that μ_n and ψ_n are defined as above. Then $\int_{-1}^{1} p|\psi_n'|^2 \leqslant \Delta_n$, where Δ_n is given by (2.6).*

Proof. Let μ_n and ψ_n be defined as above, then

$$-(p\psi_n')' + q\psi_n = \mu_n|w|\psi_n, \quad \mathcal{B}\psi_n = 0. \tag{3.6}$$

From $\mathcal{B}\psi_n = 0$ and lemma 2.1 that ψ_n is bounded and satisfies $(p\psi_n'\psi_n)(x) \to 0$, $x \to -1$ or 1. Multiplying both sides of (3.6) by ψ_n and integrating over the interval $(-1, 1)$, we have

$$\int_{-1}^{1} p|\psi_n'|^2 + \int_{-1}^{1} q|\psi_n|^2 = \mu_n \int_{-1}^{1} |w||\psi_n|^2. \tag{3.7}$$

With the similar argument in (3.5), one can prove that

$$\int_{-1}^{1} q_-|\psi_n|^2 \leqslant \int_{-1}^{1} q_- + \frac{1}{2}\int_{-1}^{1} p|\psi_n'|^2 + 4\alpha^2.$$

This together with (3.7) and $q = q_+ - q_-$ yields that

$$\begin{aligned}\int_{-1}^{1} p|\psi_n'|^2 &\leqslant \int_{-1}^{1} q_-|\psi_n|^2 + \mu_n \int_{-1}^{1} |w||\psi_n|^2 \\ &\leqslant \int_{-1}^{1} q_- + \frac{1}{2}\int_{-1}^{1} p|\psi_n'|^2 + 4\alpha^2 + |\mu_n|\|w\|_1. \end{aligned}$$

And hence

$$\int_{-1}^{1} p|\psi_n'|^2 \leqslant 2\int_{-1}^{1} q_- + 8\alpha^2 + 2|\mu_n|\|w\|_1.$$

This completes the proof of lemma 3.2. □

https://doi.org/10.1017/prm.2023.126 Published online by Cambridge University Press

For any $\varepsilon > 0$, let

$$\delta = \sup\left\{\min\left\{\tilde{\delta}, \frac{1}{2}\right\} : \int_I \frac{1}{p} \leqslant \frac{1}{4\Delta}, \text{for any } I \subset [-1/2, 1/2] \text{ with length } \tilde{\delta}\right\} \quad (3.8)$$

$$\delta_n = \sup\left\{\min\left\{\tilde{\delta}, \frac{1}{2}\right\} : \int_I \frac{1}{p} \leqslant \frac{1}{4\Delta_n}, \text{for any } I \subset [-1/2, 1/2] \text{ with length } \tilde{\delta}\right\} \quad (3.9)$$

From the definition of δ in (3.8), one sees that $\delta \in (0,\, 1/2]$ and $\int_I \frac{1}{p} \leqslant \frac{1}{4\Delta}$ for any interval $I \subset [-1/2,\, 1/2]$ with length δ. The conclusion holds for δ_n.

LEMMA 3.3. *Let λ and ϕ be defined as above. Then there exists an interval $\tilde{I} \subset (-1,\, 1)$ with δ in length, such that $|\phi(\cdot)| \geqslant 1/2$ on $\tilde{I}$.*

Proof. For any interval $I \subset [-1/2,\, 1/2]$ with length δ, it follows from Cauchy–Schwarz inequality and lemma 3.1 that

$$\left(\int_I |\phi'|\right)^2 \leqslant \int_I \frac{1}{p} \int_{-1}^{1} p|\phi'|^2 \leqslant \frac{1}{4}. \quad (3.10)$$

Since $\max\{|\phi(x)| : x \in [-1,\, 1]\} = 1$, there exists $x_0 \in [-1/2,\, 1/2]$ such that $|\phi(x_0)| \leqslant 1$. Hence, for $x \in (-1,\, 1)$ and $|x - x_0| \leqslant \delta$,

$$\Big||\phi(x)| - 1\Big| \leqslant \Big||\phi(x)| - |\phi(x_0)|\Big| \leqslant |\phi(x) - \phi(x_0)| = \left|\int_{x_0}^{x} \phi'(t)\,\mathrm{d}t\right| \leqslant \frac{1}{2}$$

by (3.10), and hence

$$|\phi(x)| \geqslant \frac{1}{2} \quad \text{on} \quad \tilde{I} = [-\delta + x_0, x_0] \quad \text{or} \quad [x_0, x_0 + \delta].$$

From $\delta \in (0,\, 1/2]$ one sees that $(-1,\, 1)$ contains at least one such interval $\tilde{I}$. □

Similar with lemma 3.3 we have

LEMMA 3.4. *Assume that μ_n is an eigenvalue of (2.4) and ψ_n is the corresponding eigenfunction. Then there exists an interval $\tilde{I}_n \subset (-1,\, 1)$ with δ_n in length, such that $|\psi_n(\cdot)| \geqslant 1/2$ on $\tilde{I}_n$.*

Applying the above lemmas we now prove the main results of theorems 2.2 and 2.3.

The proof of theorem 2.2. Let μ_n be the nth eigenvalue of right-definite problem (2.4) and ψ_n the corresponding eigenfunction. From $\psi_n \in D(S)$ is linearly independent, one sees that $\{\psi_n : n \geqslant 1\}$ forms an orthonormal system. Let ϕ be an eigenfunction of T associated with eigenvalue λ such that $\int_{-1}^{1} |w||\phi|^2 = 1$. Since S

https://doi.org/10.1017/prm.2023.126 Published online by Cambridge University Press

is a self-adjoint operator and S^{-1} is compact, we can expand ϕ via the orthonormal system ψ_n, i.e. $\phi = \sum_{n=1}^{\infty} (\phi, \psi_n)_{|w|}\psi_n$. Then from $\int_{-1}^{1} |w||\phi|^2 = 1$, we have

$$\sum_{n=1}^{\infty} |(\phi, \psi_n)_{|w|}|^2 = 1, \quad \sum_{n=1}^{\infty} |(\phi, \psi_n)_{|w|}|^2 \mu_n^2 = |\lambda|^2. \tag{3.11}$$

It follows from $\mu^+ = \min\ \sigma(S) \cap (0, \infty)$ and $\mu^- = \min\ \{|\lambda| : \lambda \in \sigma(S) \cap (-\infty, 0)\}$ that $|\mu_n| \geqslant \min\ \{\mu^+, \mu^-\}$ for $n \geqslant 1$. This together with (3.11) yields that

$$|\lambda| \geqslant \min\ \{\mu^+, \mu^-\}.$$

If λ corresponds to an eigenvector ϕ of T with $[\phi, \phi]_w = 0$, then it follows from (3.11) that

$$\sum_{n=1}^{\infty} \left|(\phi, \psi_n)_{|w|}\right|^2 \left(\mu_n - \frac{1}{2}(\mu^+ + \mu^-)\right)^2 = |\lambda|^2 + \frac{1}{4}(\mu^+ + \mu^-)^2. \tag{3.12}$$

From $|\mu_n| \geqslant \min\ \{\mu^+, \mu^-\}$, one sees that

$$\left(\mu_n - \frac{1}{2}(\mu^+ + \mu^-)\right)^2 - \frac{1}{4}(\mu^+ - \mu^-)^2 = (\mu_n - \mu^+)(\mu_n - \mu^-) \geqslant 0. \tag{3.13}$$

From (3.11), (3.12) and (3.13), we have that

$$\begin{aligned} |\lambda|^2 &= \left(\mu_n - \frac{1}{2}(\mu^+ + \mu^-)\right)^2 - \frac{1}{4}(\mu^+ + \mu^-)^2 \\ &= \left(\mu_n - \frac{1}{2}(\mu^+ + \mu^-)\right)^2 - \frac{1}{4}(\mu^+ - \mu^-)^2 - \mu^+\mu^- \geqslant -\mu^+\mu^-, \end{aligned}$$

which completes the proof of theorem 2.2. □

The proof of theorem 2.3. Let μ_n be the nth eigenvalue of (2.4) and ψ_n the corresponding eigenfunction such that $\max\{|\psi_n(x)| : x \in [-1, 1]\} = 1$, $n \geqslant 1$. It follows from lemmas 3.3, 3.4 and the definition of γ_t in (2.6) that

$$\|\phi\|_{|w|}^2 = \int_{-1}^{1} |w||\phi|^2 \geqslant \int_I |w||\phi|^2 \geqslant \int_I \frac{|w|}{4} \geqslant \frac{\gamma_\delta}{4}, \tag{3.14}$$

$$\|\psi_n\|_{|w|}^2 = \int_{-1}^{1} |w||\psi_n|^2 \geqslant \int_{I_n} |w||\psi_n|^2 \geqslant \int_{I_n} \frac{|w|}{4} \geqslant \frac{\gamma_{\delta_n}}{4},\ n \geqslant 1. \tag{3.15}$$

From the definition of $W(x) = \int_{-1}^{x} w(t)\mathrm{d}t$, one sees that

$$[\phi, \psi_n]_w = \int_{-1}^{1} w\phi\overline{\psi_n} = \int_{-1}^{1} W'\phi\overline{\psi_n} = \phi(1)\overline{\psi_n}(1)\int_{-1}^{1} w - \int_{-1}^{1} W(\phi'\overline{\psi_n} + \phi\overline{\psi_n'}).$$

https://doi.org/10.1017/prm.2023.126 Published online by Cambridge University Press

This together with (2.5) and lemmas 3.1 and 3.2 that

$$\begin{aligned}\left|[\phi,\psi_n]_w\right| &\leqslant \int_{-1}^{1}|w|+\left|\int_{-1}^{1}W(\phi'\overline{\psi_n}+\phi\overline{\psi_n'})\right|\\ &\leqslant \|w\|_1+\left|\int_{-1}^{1}\frac{W}{\sqrt{p}}\sqrt{p}\phi'\overline{\psi_n}+\int_{-1}^{1}\frac{W}{\sqrt{p}}\phi\sqrt{p}\overline{\psi_n'}\right|\\ &\leqslant \|w\|_1+C_2\left(\int_{-1}^{1}|\sqrt{p}\phi'\overline{\psi_n}|+\int_{-1}^{1}|\phi\sqrt{p}\overline{\psi_n'}|\right)\\ &\leqslant \|w\|_1+C\left(\sqrt{\Delta}+\sqrt{\Delta_n}\right)=\Delta_{w,1,n}.\end{aligned}$$

Furthermore, for $n\geqslant 1$ we get

$$\lambda[\phi,\psi_n]_w=[T\phi,\psi_n]_w=(S\phi,\psi_n)_{|w|}=(\phi,S\psi_n)_{|w|}=\mu_n(\phi,\psi_n)_{|w|}. \tag{3.16}$$

If we set

$$\Lambda_n=(\phi,\psi_n)_{|w|}=\frac{(\phi,\psi_n)_{|w|}}{\|\phi\|_{|w|}\|\psi_n\|_{|w|}},$$

then (3.14)–(3.16) give that

$$|\Lambda_n|\leqslant\frac{4|\lambda|\left|[\phi,\psi_n]_w\right|}{|\mu_n|\sqrt{\gamma_\delta\gamma_{\delta_n}}}\leqslant\frac{4|\lambda|\Delta_{w,1,n}}{|\mu_n|\sqrt{\gamma_\delta\gamma_{\delta_n}}}. \tag{3.17}$$

From $\sum\limits_{n=1}^{\infty}|(\phi,\,\psi_n)_{|w|}|^2\mu_n=\lambda[\phi,\,\phi]_w$ and $[\phi,\,\phi]_w=\int_{-1}^{1}w|\phi|^2=0$ in lemma 3.1, we have

$$0=\lambda[\phi,\phi]_w=\sum_{n=1}^{\infty}\left|(\phi,\psi_n)_{|w|}\right|^2\mu_n=\sum_{n=1}^{\infty}|\Lambda_n|^2\mu_n=\sum_{n=1}^{h-1}|\Lambda_n|^2\mu_n+\sum_{n=h}^{\infty}|\Lambda_n|^2\mu_n,$$

and hence $-\sum\limits_{n=1}^{h-1}|\Lambda_n|^2\mu_n=\sum\limits_{n=h}^{\infty}|\Lambda_n|^2\mu_n$. Thus, by (3.11) and $\mu_{h-1}<0<\mu_h$, $h\geqslant 2$,

$$\begin{aligned}\sum_{n=1}^{h-1}|\Lambda_n|^2(\mu_h-\mu_n)&=\sum_{n=1}^{h-1}|\Lambda_n|^2\mu_h+\sum_{n=h}^{\infty}|\Lambda_n|^2\mu_n\\ &=\sum_{n=1}^{h}|\Lambda_n|^2\mu_h+\sum_{n=h+1}^{\infty}|\Lambda_n|^2\mu_n\geqslant\sum_{n=1}^{h}|\Lambda_n|^2\mu_h+\sum_{n=h+1}^{\infty}|\Lambda_n|^2\mu_h\\ &=\sum_{n=1}^{\infty}|\Lambda_n|^2\mu_h=\mu_h.\end{aligned} \tag{3.18}$$

By $\mu_h>0$, $\mu_n\leqslant\mu_{h-1}<0$, $1\leqslant n\leqslant h-1$, we have

$$\frac{1}{\mu_n^2}\leqslant\frac{1}{\mu_{h-1}^2},\quad\frac{\mu_h}{\mu_n^2}\leqslant\frac{\mu_h}{\mu_{h-1}^2},\quad\frac{-1}{\mu_n}\leqslant\frac{-1}{\mu_{h-1}}=\frac{-\mu_{h-1}}{\mu_{h-1}^2}.$$

https://doi.org/10.1017/prm.2023.126 Published online by Cambridge University Press

This together with the assumption that $\mu_1 \leqslant \cdots \leqslant \mu_{h-1} < 0 < \mu_h$, $h \geqslant 2$ yields that

$$\frac{\mu_h - \mu_n}{\mu_n^2} = \frac{\mu_h}{\mu_n^2} - \frac{1}{\mu_n} \leqslant \frac{\mu_h}{\mu_{h-1}^2} - \frac{\mu_{h-1}}{\mu_{h-1}^2} = \frac{\mu_h - \mu_{h-1}}{\mu_{h-1}^2}, \quad 1 \leqslant n \leqslant h-1. \tag{3.19}$$

Hence, by (3.17)–(3.19) we have

$$\begin{aligned}\mu_h &= \sum_{n=1}^{h-1} |\Lambda_n|^2(\mu_h - \mu_n) \leqslant \sum_{n=1}^{h-1} \frac{16|\lambda|^2\Delta_{w,1,n}^2(\mu_h - \mu_n)}{|\mu_n|^2\gamma_\delta\gamma_{\delta_n}} = \frac{16|\lambda|^2}{\gamma_\delta}\sum_{n=1}^{h-1}\frac{\mu_h - \mu_n}{|\mu_n|^2\gamma_{\delta_n}}\Delta_{w,1,n}^2 \\ &\leqslant \frac{16|\lambda|^2}{\gamma_\delta}\sum_{n=1}^{h-1}\frac{\mu_h - \mu_{h-1}}{\mu_{h-1}^2\gamma_{\delta_n}}\Delta_{w,1,n}^2 = \frac{16|\lambda|^2(\mu_h - \mu_{h-1})}{\mu_{h-1}^2\gamma_\delta}\sum_{n=1}^{h-1}\frac{\Delta_{w,1,n}^2}{\gamma_{\delta_n}}.\end{aligned}$$

Therefore,

$$|\lambda|^2 \geqslant \frac{\mu_h\mu_{h-1}^2\gamma_\delta}{16(\mu_h - \mu_{h-1})}\left(\sum_{n=1}^{h-1}\frac{\Delta_{w,1,n}^2}{\gamma_{\delta_n}}\right)^{-1}.$$

This completes the proof of theorem 2.3. □

Acknowledgements

The author is deeply grateful to the editors and reviewers for a careful reading and very helpful suggestions which improved and strengthened the presentation of this manuscript. The author would like to thank Professor Jiangang Qi (Shandong University) and Professor Zhaowen Zheng (Guangdong Polytech Normal University) for their generous help. This research was partially supported by the National Natural Science Foundation of China (grant: 12101356), Natural Science Foundation of Shandong Province (grant: ZR2021QA065) and China Postdoctoral Science Foundation (grant: 2023M732023).

References

1 F. Atkinson and D. Jabon. Indefinite Sturm-Liouville problems. In *'Proceedings of the Focused Research Program on Spectral Theory and Boundary Value Problems* Vol.I', (ed. H. Kaper, M. K. Kwong, and A. Zettle) (Argonne National Laboratory, Argonne, IL, 1984), pp. 31–45.

2 J. Behrndt, S. Chen, F. Philipp and J. Qi. Estimates on the non-real eigenvalues of regular indefinite Sturm-Liouvilleproblems. *Proc. Roy. Soc. Edinburgh. A.* **144** (2014), 1113–1126.

3 J. Behrndt, Q. Katatbeh and C. Trunk. Non-real eigenvalues of singular indefinite Sturm-Liouville operators. *Proc. Amer. Math. Soc.* **137** (2009), 3797–3806.

4 J. Behrndt, F. Philipp and C. Trunk. Bounds on the non-real spectrum of differential operators with indefinite weights. *Math. Ann.* **357** (2013), 185–213.

5 J. Behrndt, P. Schmitz and C. Trunk. Bounds on the non-real spectrum of a singular indefinite Sturm-Liouville operator on $\mathbb{R}$. *Proc. Appl. Math. Mech.* **16** (2016), 881–882.

6 J. Behrndt, P. Schmitz and C. Trunk. Spectral bounds for singular indefinite Sturm-Liouville operators with L^1-potentials, arXiv:1709.04994v2 [math.SP] 18 Dec 2017.

7 J. Behrndt and C. Trunk. On the negative squares of indefinite Sturm-Liouville operators. *J. Differ. Equ.* **238** (2007), 491–519.

8 P. Binding and H. Volkmer. Eigencurves for two-parameter Sturm-Liouville equations. *SIAM Rev.* **38** (1996), 27–48.

https://doi.org/10.1017/prm.2023.126 Published online by Cambridge University Press

9 B. Čurgus and H. Langer. A Krein space approach to symmetric ordinary differential operators with an indefinite weight functions. *J. Differ. Equ.* **79** (1989), 31–61.

10 J. Fleckinger and A. B. Mingarelli, On the eigenfunctions of non-definite elliptic operators. In *Differential Equations* Vol. 92, (ed. I. W. Knowles and R. T. Lewis) (North-Holland, The Netherlands, 1984), pp. 219–227.

11 O. Haupt. Über eine Methode zum Beweis von Oszillationstheoremen. *Math. Ann.* **76** (1915), 67–104.

12 L. Kong, Q. Kong, H. Wu and A. Zettl. Regular approximations of singular Sturm-Liouville problems with limit-circle endpoints. *Result. Math.* **45** (2004), 274–292.

13 Q. Kong, M. Möller, H. Wu and A. Zettl. Indefinite Sturm-Liouville problems. *Proc. Roy. Soc. Edinburgh. Sect. A.* **133** (2003), 639–652.

14 A. B. Mingarelli, *Indefinite Sturm-Liouville problems*, Lecture Notes in Math., Vol. 964, (Springer, Berlin, New York, 1982, pp. 519–528).

15 A. B. Mingarelli, A survey of the regular weighted Sturm-Liouville problem – the non-definite case, e-print arXiv:1106.6013v1 [math.CA], 2011.

16 H. D. Niessen and A. Zettl. Singular Sturm-Liouville problem: the Friedrichs extension and comparison of eigenvalues. *Proc. London Math. Soc.* **64** (1992), 545–578.

17 J. Qi and S. Chen. A priori bounds and existence of non-real eigenvalues of indefinite Sturm-Liouville problems. *J. Spectr. Theory.* **4** (2014), 53–63.

18 J. Qi, B. Xie and S. Chen. The upper and lower bounds on non-real eigenvalues of indefinite Strum-Liouville problems. *Pro. Amer. Math. Soc.* **144** (2016), 547–559.

19 R. G. D. Richardson. Theorems of oscillation for two linear differential equations of second order with two parameters. *Trans. Amer. Math. Soc.* **13** (1912), 22–34.

20 F. Sun and J. Qi. A priori bounds and existence of non-real eigenvalues for singular indefinite Sturm-Liouville problems with limit-circle type endpoints. *Proc. Roy. Soc. Edinburgh A.* **150** (2020), 2607–2619.

21 B. Xie and J. Qi. Non-real eigenvalues of indefinite Sturm-Liouville problems. *J. Differ. Equ.* **8** (2013), 2291–2301.

22 A. Zettl, *Sturm-Liouville Theory*, Math. Surveys Monogr., Vol. 121, (Amer. Math. Soc., Providence, RI, 2005).

https://doi.org/10.1017/prm.2023.126 Published online by Cambridge University Press