The Journal of Symbolic Logic
Volume 91, Number 2, June 2026

A HIERARCHY ON NON-ARCHIMEDEAN POLISH GROUPS ADMITTING A COMPATIBLE COMPLETE LEFT-INVARIANT METRIC

LONGYUN DING AND XU WANG

Abstract. In this article, we introduce a hierarchy on the class of non-archimedean Polish groups that admit a compatible complete left-invariant metric. We denote this hierarchy by α-CLI and L-α-CLI where α is a countable ordinal. We establish three results:

(1) G is 0-CLI iff $G = \{1_G\}$;
(2) G is 1-CLI iff G admits a compatible complete two-sided invariant metric; and
(3) G is L-α-CLI iff G is locally α-CLI, i.e., G contains an open subgroup that is α-CLI.

Subsequently, we show this hierarchy is proper by constructing non-archimedean CLI Polish groups G_α and H_α for $\alpha < \omega_1$, such that:

(1) H_α is α-CLI but not L-β-CLI for $\beta < \alpha$; and
(2) G_α is $(\alpha + 1)$-CLI but not L-α-CLI.

§1. Introduction. A Polish group is *non-archimedean* if it has a neighborhood basis of its identity element consisting of open subgroups. By a theorem of Becker and Kechris (cf. [1, Theorem 1.5.1]), a Polish group is non-archimedean iff it is homeomorphic to a closed subgroup of S_∞, the group of all permutations of $\mathbb{N}$ equipped with the pointwise convergence topology. A metric d on a group G is *left-invariant* if $d(gh, gk) = d(h, k)$ for all $g, h, k \in G$. A Polish group is *CLI* if it admits a compatible complete left-invariant metric.

Malicki [5] defined a notion of orbit tree T_G for each closed subgroup G of S_∞, and showed that G is CLI iff T_G is well-founded. Moreover, he proved that the heights of orbit trees of all CLI closed subgroups of S_∞ are cofinal in ω_1. Malicki proved that the family of all CLI groups is coanalytic non-Borel based on this cofinality. After that, Xuan defined a different kind of orbit trees and showed that, a closed subgroup of S_∞ is locally compact iff its orbit tree has finite height (cf. [6, Theorem 3.7]). It is worth noting that both kinds of orbit trees defined by Malicki and Xuan are all defined on closed subgroups of S_∞ rather than directly on non-archimedean Polish groups. As a result, two topologically isomorphic closed subgroups of S_∞ can have completely different orbit trees, and even the rank of their orbit trees can be different. This suggests that one cannot use ranks of orbit trees directly to define a hierarchy.

In this article, for a given non-archimedean CLI Polish group G, we use a neighborhood basis of the identity 1_G to define a new type of orbit trees. This differs from the approach which employs a closed subgroup of S_∞ topologically

Received September 13, 2023.
2020 *Mathematics Subject Classification.* Primary 03E15, Secondary 22A05.
Key words and phrases. hierarchy, non-archimedean Polish group, tree.

© The Author(s), 2024. Published by Cambridge University Press on behalf of The Association for Symbolic Logic.
0022-4812/26/9102-0010
DOI:10.1017/jsl.2024.7

https://doi.org/10.1017/jsl.2024.7 Published online by Cambridge University Press

isomorphic to G. More specifically, let $\mathcal{G} = (G_n)$ be a decreasing sequence of open subgroups of G with $G_0 = G$, such that (G_n) forms a neighborhood basis of 1_G. We will define a well-founded tree $T_{\mathcal{G}}^{X(\mathcal{G})}$ and denote its rank by $\rho(\mathcal{G})$. We let $\mathrm{rank}(G)$ be the ordinal given by $\max\{\beta : \rho(\mathcal{G}) \geq \omega \cdot \beta\}$. Then we shall prove that the following are independent from the choice of $\mathcal{G}$: (a) the value of the ordinal $\mathrm{rank}(G)$; and (b) whether $\rho(\mathcal{G})$ is a limit ordinal or not. These facts allow us to form a well-defined hierarchy on the class of non-archimedean CLI Polish groups: given an ordinal $\alpha < \omega_1$,

(1) if $\rho(\mathcal{G}) \leq \omega \cdot \alpha$, we say G is α-CLI;
(2) if $\rho(\mathcal{G}) \leq \omega \cdot \alpha + m$ for some $m < \omega$, i.e., $\mathrm{rank}(G) \leq \alpha$, we say G is L-α-CLI.

It is clear that, if G is L-α-CLI, then it is also $(\alpha + 1)$-CLI.

The following theorem shows that the hierarchy classifies non-archimedean CLI Polish groups in a good manner:

THEOREM 1.1. *Let G be a non-archimedean CLI Polish group and α be a countable ordinal. Then*:

(1) *G is 0-CLI iff $G = \{1_G\}$*;
(2) *G is L-0-CLI iff G is discrete*;
(3) *G is 1-CLI iff G is TSI, i.e., G admits a compatible complete two-sided invariant metric*;
(4) *G is L-α-CLI iff G is locally α-CLI, i.e., G has an open subgroup which is α-CLI.*

It is well-known that all compact Polish groups are TSI (cf. [2, Theorem 2.1.5]), and all locally compact Polish groups are CLI (cf. [2, Theorem 2.2.5]). Now we know that all compact non-archimedean Polish groups are 1-CLI, and all locally compact non-archimedean Polish groups are L-1-CLI.

THEOREM 1.2. *Let G be a non-archimedean CLI Polish group and α be a countable ordinal. Assume H and N are closed subgroups of G, and that N is normal in G. If G is α-CLI (or L-α-CLI), so are H and G/N. In particular, we have* $\mathrm{rank}(H) \leq \mathrm{rank}(G)$ *and* $\mathrm{rank}(G/N) \leq \mathrm{rank}(G)$.

THEOREM 1.3. *Let (G^i) be a sequence of non-archimedean CLI Polish groups, $\alpha < \omega_1$, and let $G = \prod_i G^i$. Then we have*:

(1) *G is α-CLI iff all G^i are α-CLI; and*
(2) *G is L-α-CLI iff all G^i are L-α-CLI and for all but finitely many i, G^i is α-CLI.*

Finally, we prove the following theorem, which indicates that this hierarchy is proper:

THEOREM 1.4. *For any $\alpha < \omega_1$, there exist two non-archimedean CLI Polish groups G_α and H_α with* $\mathrm{rank}(G_\alpha) = \mathrm{rank}(H_\alpha) = \alpha$ *such that H_α is α-CLI and G_α is L-α-CLI but not α-CLI.*

§2. Preliminaries. We denote the class of all ordinals by Ord. For any $\alpha \in \mathrm{Ord}$, we define

$$\omega(\alpha) = \max\{0, \lambda : \lambda \leq \alpha \text{ is a limit ordinal}\}.$$

Then $\alpha = \omega(\alpha) + m$ for some $m < \omega$.

https://doi.org/10.1017/jsl.2024.7 Published online by Cambridge University Press

Let E be an equivalence relation on a set X, $x \in X$, and $A \subseteq X$. The E-equivalence class of x is $[x]_E = \{y \in X : xEy\}$. Similarly, the E-saturation of A is $[A]_E = \{y \in X : \exists z \in A\,(yEz)\}$.

The identity element of a group G is denoted by 1_G. Let H be a subgroup of G, we denote the set of all left-cosets of H by G/H.

A topological space is *Polish* if it is separable and completely metrizable. A topological group is *Polish* if its underlying topology is Polish. Let G be a Polish group and X a Polish space, an action of G on X, denoted by $G \curvearrowright X$, is a map $a : G \times X \to X$ that satisfies $a(1_G, x) = x$ and $a(gh, x) = a(g, a(h, x))$ for $g, h \in G$ and $x \in X$. The pair (X, a) is called a *Polish G-space* if a is continuous. For brevity, we write $g \cdot x$ in place of $a(g, x)$. The *orbit equivalence relation* E_G^X is defined as $xE_G^X y \iff \exists g \in G\,(g \cdot x = y)$. Note that the E_G^X-equivalence class of x is $G \cdot x = \{g \cdot x : g \in G\}$, which is also called the *G-orbit* of x. Similarly, for $A \subseteq X$, the E_G^X-saturation of A is $G \cdot A = \{g \cdot x : g \in G \wedge x \in A\}$.

Let $<$ be a binary relation on a set T. We say that $(T, <)$ is a *tree* if:

(1) $\forall s \in T\,(s \not< s)$,
(2) $\forall s, t, u \in T\,((s < t \wedge t < u) \Rightarrow s < u)$,
(3) $\forall s \in T\,(|\{t \in T : t < s\}| < \omega \wedge \forall t, u < s\,(t = u \vee t < u \vee u < t))$.

For $s \in T$, we define $\mathrm{lh}(s) = |\{t \in T : t < s\}|$, which is called the length of s. It is clear that $s < t$ implies $\mathrm{lh}(s) < \mathrm{lh}(t)$. For $n < \omega$, we denote the nth level of T by

$$L_n(T) = \{s \in T : \mathrm{lh}(s) = n\}.$$

Each element in $L_0(T)$ is called a *root* of T.

Let $(T, <)$ be a tree. We say that T is *well-founded* if any non-empty subset of T contains at least a maximal element, or equivalently (under AC), if T contains no infinite strictly increasing sequence. Let T be a well-founded tree. We define the rank function $\rho_T : T \to \mathrm{Ord}$ by transfinite induction as

$$\rho_T(s) = \sup\{\rho_T(t) + 1 : s < t \wedge t \in T\}.$$

If $\rho_T(s) = 0$, we say that s is a *terminal* of T. Then we define

$$\rho(T) = \sup\{\rho_T(s) + 1 : s \in T\}.$$

So $\rho(T) = 0$ iff $T = \emptyset$. It is clear that $\rho(T) = \sup\{\rho_T(s) + 1 : s \in L_0(T)\}$. If $L_0(T) = \{s_0\}$ is a singleton, then $\rho(T) = \rho_T(s_0) + 1$ is a successor ordinal.

For $s \in T$, we define

$$T_s = \{t \in T : s = t \vee s < t\}.$$

Since $L_0(T_s) = \{s\}$, we have $\rho(T_s) = \rho_T(s) + 1$. For the convenience of discussion, we let $T_s = \emptyset$ whenever $s \notin T$; in other words, $\rho(T_s) = 0$. This convention will be useful in some proofs (see Lemma 4.3). Note that $\rho(T_s)$ is always a non-limit ordinal regardless of $s \in T$ or not.

First, we note the following facts:

PROPOSITION 2.1. *Let T be a well-founded tree, then*

$$\rho(T) = \sup\{\rho(T_s) : s \in T\} = \sup\{\rho(T_s) : s \in T \wedge s \in L_0(T)\},$$

https://doi.org/10.1017/jsl.2024.7 Published online by Cambridge University Press

and for all $s \in T$,

$$\begin{aligned}\rho(T_s) &= \sup\{\rho(T_t) : s < t \wedge t \in T\} + 1\\ &= \sup\{\rho(T_t) : s < t \wedge t \in T \wedge \mathrm{lh}(t) = \mathrm{lh}(s) + 1\} + 1.\end{aligned}$$

PROPOSITION 2.2. *Let* $(T, <)$ *be a well-founded tree,* $k < \omega$. *Then we have*:

(1) $\sup\{\rho(T_s) : s \in L_k(T)\} \leq \rho(T) \leq \sup\{\rho(T_s) : s \in L_k(T)\} + k$;
(2) $\omega(\rho(T)) = \omega(\sup\{\rho(T_s) : s \in L_k(T)\})$;
(3) *if* $\rho(T_s) \geq \alpha$ *for some* $s \in L_k(T)$, *then* $\rho(T) \geq \alpha + k$.

PROOF. It is routine to prove clause (1) by induction on k based on Proposition 2.1. Clause (2) is an easy corollary of (1). And clause (3) is trivial. ⊣

Let $(S, <)$ and $(T, <)$ be two trees. A map $\phi : S \to T$ is said to be an *order-preserving map* if

$$\forall s, t \in S\, (s < t \Rightarrow \phi(s) < \phi(t)).$$

It is said to be an *order-preserving embedding (isomorphism)* if it is injective (bijective) and

$$\forall s, t \in S\, (s < t \iff \phi(s) < \phi(t)).$$

In particular, an order-preserving map ϕ is said to be *Lipschitz* if $\mathrm{lh}(\phi(s)) = \mathrm{lh}(s)$ for all $s \in S$.

PROPOSITION 2.3. *Let* $(S, <)$ *and* $(T, <)$ *be trees, and assume that* $(T, <)$ *is well-founded. If there exists an order-preserving map* $\phi : S \to T$, *then* $(S, <)$ *is also well-founded, and* $\rho(S) \leq \rho(T)$ *holds.*

PROOF. If S contains an infinite strictly increasing sequence (s_n), then $(\phi(s_n))$ is an infinite strictly increasing sequence in T, contradicting that $(T, <)$ is well-founded.

Now we prove that $\rho_S(s) \leq \rho_T(\phi(s))$ for $s \in S$ by induction on $\rho_S(s)$. For the basis of induction, note that $\rho_S(s) = 0 \leq \rho_T(\phi(s))$ clearly holds; and for the inductive step, we have the following inequality:

$$\begin{aligned}\rho_S(s) &= \sup\{\rho_S(u) + 1 : s < u \wedge u \in S\}\\ &\leq \sup\{\rho_T(\phi(u)) + 1 : s < u \wedge u \in S\}\\ &\leq \sup\{\rho_T(t) + 1 : \phi(s) < t \wedge t \in T\} = \rho_T(\phi(s)).\end{aligned}$$

This implies $\rho(S) \leq \rho(T)$. ⊣

§3. Definition of the hierarchy

DEFINITION 3.1. Let X be a set, and $\mathcal{E} = (E_n)$ be a decreasing sequence of equivalence relations on X, i.e., $E_n \supseteq E_{n+1}$ for each $n < \omega$. We define

$$T^X_{\mathcal{E}} = \{(n, C) : \exists x \in X\, (C = [x]_{E_n} \neq \{x\})\}.$$

For $(n, C), (m, D) \in T^X_{\mathcal{E}}$, we define

$$(n, C) < (m, D) \iff n < m \wedge C \supseteq D.$$

https://doi.org/10.1017/jsl.2024.7 Published online by Cambridge University Press

It is straightforward to check that $(T_{\mathcal{E}}^X, <)$ is a tree. Note that $(n, C) \in T_{\mathcal{E}}^X$ iff C is a non-singleton equivalence class of E_n. Here we omit all singleton classes in our definition. This will be crucial in the proof of Lemma 3.3.

DEFINITION 3.2. Let G be a non-archimedean Polish group. We denote by $\mathrm{dgnb}(G)$ the set of all decreasing sequences $\mathcal{G} = (G_n)$ of open subgroups of G, such that $G_0 = G$ and (G_n) forms a neighborhood basis of 1_G. Here 'dgnb' stands for 'decreasing group neighborhood basis'.

Let X be a countable discrete Polish G-space, $\mathcal{G} = (G_n) \in \mathrm{dgnb}(G)$. We define $E_n = E_{G_n}^X$, i.e., $xE_n y \iff \exists g \in G_n\, (g \cdot x = y)$, and hence $[x]_{E_n} = G_n \cdot x$. Then we write $\mathcal{E} = (E_n)$ and

$$T_{\mathcal{G}}^X = T_{\mathcal{E}}^X.$$

Therefore, $(n, C) \in T_{\mathcal{G}}^X$ iff C is a non-singleton G_n-orbit. We point out that our definition is different from Malicki's and Xuan's, which are based on infinite orbits.

The following lemma builds a connection between non-archimedean CLI Polish groups and well-founded trees. Similar results also appear in [5, Theorem 3] and [6, Theorem 3.9].

LEMMA 3.3. *Let G be a non-archimedean CLI Polish group, X a countable discrete Polish G-space, and let $\mathcal{G} = (G_n) \in \mathrm{dgnb}(G)$. Then $T_{\mathcal{G}}^X$ is well-founded.*

PROOF. Assume for contradiction that $T_{\mathcal{G}}^X$ is ill-founded, then there exists an infinite sequence (n, C_n), $n < \omega$ in $T_{\mathcal{G}}^X$ with $C_n \supseteq C_{n+1}$ for each $n < \omega$.

Let d be a compatible complete left-invariant metric on G.

Fix an $x_0 \in C_0$. Then we have $G_0 \cdot x_0 = C_0 \supseteq C_1$, so we can find a $g_0 \in G_0$ such that $g_0 \cdot x_0 \in C_1$. Inductively, we can find a $g_n \in G_n$ for each $n < \omega$ such that $g_n g_{n-1} \dots g_0 \cdot x_0 \in C_{n+1}$. Put $h_n = g_0^{-1} \dots g_n^{-1}$ for each $n < \omega$. Then, for any $n, p < \omega$, we have $d(h_{n+p}, h_n) = d(h_n^{-1} h_{n+p}, 1_G) = d(g_{n+1}^{-1} \dots g_{n+p}^{-1}, 1_G) \leq \mathrm{diam}(G_{n+1}) \to 0$ as $n \to \infty$. It follows that (h_n) is a d-Cauchy sequence in G, so it converges to some $h \in G$.

Let $x_\infty = h^{-1} \cdot x_0$. Since $h_n \to h$, we have $h_n^{-1} \to h^{-1}$, and hence $h_n^{-1} \cdot x_0 \to h^{-1} \cdot x_0 = x_\infty$. Note that X is discrete, so there exists an integer N such that $h_n^{-1} \cdot x_0 = x_\infty$ for any $n > N$, thus $x_\infty = g_n g_{n-1} \dots g_0 \cdot x_0 \in C_{n+1}$. This implies that $x_\infty \in \bigcap_n C_n$ and $G_n \cdot x_\infty = C_n$ for each $n < \omega$.

Finally, define $G_{x_\infty} = \{g \in G : g \cdot x_\infty = x_\infty\}$ and put $f : G \to X$ as $f(g) = g \cdot x_\infty$. Since f is continuous and $\{x_\infty\}$ is clopen in X, it follows that $G_{x_\infty} = f^{-1}(x_\infty)$ is a clopen subgroup of G. So there exists an $m < \omega$ such that $G_m \subseteq G_{x_\infty}$. We now have $C_m = G_m \cdot x_\infty = \{x_\infty\}$, which is a singleton G_m-orbit, contradicting that $(m, C_m) \in T_{\mathcal{G}}^X$. ⊣

Given two sets X and Y. Let $\mathcal{E} = (E_n)$ and $\mathcal{F} = (F_n)$ be two decreasing sequences of equivalence relations on X and Y, respectively. Let $\theta : X \to Y$ be an injection. For $(n, C) \in T_{\mathcal{E}}^X$, define $\phi(n, C) = (n, [\theta(C)]_{F_n})$.

PROPOSITION 3.4. (1) *If θ is an (E_n, F_n)-homomorphism for each $n < \omega$, i.e.,*

$$\forall n < \omega\, \forall x, x' \in X\, (xE_n x' \Rightarrow \theta(x) F_n \theta(x')),$$

then ϕ is an order-preserving map from $T_{\mathcal{E}}^X$ to $T_{\mathcal{F}}^Y$.

https://doi.org/10.1017/jsl.2024.7 Published online by Cambridge University Press

(2) *If θ is an reduction of E_n to F_n for each $n < \omega$, i.e.,*

$$\forall n < \omega \, \forall x, x' \in X \, (xE_n x' \Leftrightarrow \theta(x) F_n \theta(x')),$$

then ϕ is an Lipschitz embedding from $T^X_{\mathcal{E}}$ to $T^Y_{\mathcal{F}}$. In particular, if θ is a bijection, then ϕ is an order-preserving isomorphism.

PROOF. Note that θ is injective. Let us prove that $\phi(n, C) \in T^Y_{\mathcal{F}}$ holds for an arbitrary $(n, C) \in T^X_{\mathcal{E}}$. Indeed, it suffices to note that since C is not a singleton, neither is $[\phi(C)]_{F_n}$. This proves $\phi(n, C) \in T^Y_{\mathcal{F}}$. The rest of the proof is trivial. ⊣

DEFINITION 3.5. Let $(T, <)$ be a tree, (n_i) a strictly increasing sequence of natural numbers. We define

$$T|(n_i) = \bigcup_i L_{n_i}(T).$$

Note that $(T|(n_i), <)$ is also a tree, and that $L_j(T|(n_i)) = L_{n_j}(T)$ holds for each $j < \omega$. We call $T|(n_i)$ a *level-subtree* of T.

LEMMA 3.6. *Let $(T, <)$ be a well-founded tree, (n_i) a strictly increasing sequence of natural numbers. Then we have*:

$$\omega(\rho(T)) \leq \rho(T|(n_i)) \leq \rho(T).$$

In particular, if $\rho(T)$ is a limit ordinal, then $\rho(T|(n_i)) = \rho(T)$.

PROOF. $\rho(T|(n_i)) \leq \rho(T)$ follows from Proposition 2.3. We prove $\omega(\rho(T)) \leq \rho(T|(n_i))$ by induction on $\rho(T)$.

First, if $\rho(T) < \omega$, then $\rho(T) = \min\{n : L_n(T) = \emptyset\}$, and hence $\rho(T|(n_i)) = \min\{i : L_{n_i}(T) = \emptyset\}$. So we have $\omega(\rho(T)) = 0 \leq \rho(T|(n_i))$.

For $t \in T|(n_i)$, note that $(T|(n_i))_t = \{u \in T|(n_i) : t = u \vee t < u\}$ is a level-subtree of T_t as well.

Case 1: If $\rho(T)$ is a limit ordinal, then $\omega(\rho(T)) = \rho(T)$. Proposition 2.1 implies that $\rho(T) = \sup\{\rho(T_t) : t \in T\}$. Since $\rho(T)$ is a limit ordinal and $\rho(T_t)$ is a successor ordinal, we have $\rho(T_t) < \rho(T)$ for $t \in T$.

Subcase 1.1: If there is no maximum in $\{\omega(\rho(T_t)) : t \in T\}$, we have

$$\rho(T) = \sup\{\rho(T_t) : t \in T\} = \sup\{\omega(\rho(T_t)) : t \in T\}.$$

By the inductive hypothesis, we have $\omega(\rho(T_t)) \leq \rho((T|(n_i))_t)$. Proposition 2.3 gives $\rho((T|(n_i))_t) \leq \rho(T|(n_i))$ for each $t \in T$, so we have $\rho(T|(n_i)) = \rho(T)$.

Subcase 1.2: Otherwise, let $\alpha = \max\{\omega(\rho(T_t)) : t \in T\}$. Since $\rho(T_t) < \rho(T)$ for $t \in T$, we have $\rho(T) = \alpha + \omega$. We can find a sequence t_m, $m < \omega$ in L_0 such that $\rho(T_{t_m}) = \alpha + k_m$ with $\sup\{k_m : m < \omega\} = \omega$. By Proposition 2.1, for each $m < \omega$ and $n < k_m$ we can find $t^n_m \in L_n(T)$ such that $t_m = t^0_m < t^1_m < \cdots < t^{k_m - 1}_m$ and $\rho(T_{t^n_m}) = \alpha + (k_m - n)$. For $k_m > n_0$, let i_m be the largest i such that $n_i < k_m$, then $t^{n_j}_m \in L_j(T|(n_i)) = L_{n_j}(T)$ for $j \leq i_m$. By the inductive hypothesis, $\alpha = \omega(\rho(T_{t^{n_j}_m})) \leq \rho((T|(n_i))_{t^{n_j}_m})$. Since $\rho((T|(n_i))_{t^{n_j}_m}) \geq \rho((T|(n_i))_{t^{n_{j+1}}_m}) + 1$ for each $j < i_m$, we have $\rho((T|(n_i))_{t^{n_0}_m}) \geq \alpha + i_m$. By the definition of i_m, we have $\sup\{i_m : m < \omega\} = \omega$. This gives $\rho(T|(n_i)) = \alpha + \omega = \rho(T)$.

https://doi.org/10.1017/jsl.2024.7 Published online by Cambridge University Press

Case 2: If $\rho(T) = \omega(\rho(T)) + n$ with $1 \leq n < \omega$, then there exists some $t_0 \in L_0(T)$ such that $\rho(T) = \rho_T(t_0) + 1$. Since $\{u \in T|(n_i) : t_0 < u\}$ is a level-subtree of $\{u \in T : t_0 < u\}$ and $\rho(\{u \in T : t_0 < u\}) = \rho_T(t_0) < \rho(T)$, by the inductive hypothesis and Proposition 2.3, we have

$$\omega(\rho_T(t_0)) = \omega(\rho(\{u \in T : t_0 < u\})) \leq \rho(\{u \in T|(n_i) : t_0 < u\}) \leq \rho(T|(n_i)).$$

It follows that $\omega(\rho(T)) = \omega(\rho_T(t_0)) \leq \rho(T|(n_i))$. ⊣

In general, the tree $T^X_{\mathcal{G}}$ and the ordinal $\rho(T^X_{\mathcal{G}})$ depend on $\mathcal{G}$, and not only on the action $G \curvearrowright X$. The following key lemma shows that $\omega(\rho(T^X_{\mathcal{G}}))$ is independent from the choice of $\mathcal{G}$.

LEMMA 3.7. *Let G be a non-archimedean CLI Polish group, X be a countable discrete Polish G-space, and let $\mathcal{G} = (G_n), \mathcal{G}' = (G'_n) \in \mathrm{dgnb}(G)$. Then*

$$\omega(\rho(T^X_{\mathcal{G}})) = \omega(\rho(T^X_{\mathcal{G}'})).$$

PROOF. (1) First, we consider the case where (G'_n) is a subsequence of (G_n), i.e., there is a strictly increasing sequence (n_i) of natural numbers such that $G'_i = G_{n_i}$ for each $i < \omega$.

We define $\psi : T^X_{\mathcal{G}'} \to T^X_{\mathcal{G}}$ as $\psi(i, C) = (n_i, C)$. It is clear that ψ is an order-preserving isomorphism from $T^X_{\mathcal{G}'}$ onto $T^X_{\mathcal{G}}|(n_i)$. It follows that

$$\omega(\rho(T^X_{\mathcal{G}})) \leq \rho(T^X_{\mathcal{G}'}) = \rho(T^X_{\mathcal{G}}|(n_i)) \leq \rho(T^X_{\mathcal{G}}).$$

So we have $\omega(\rho(T^X_{\mathcal{G}})) = \omega(\rho(T^X_{\mathcal{G}'}))$.

(2) Since $(G_n), (G'_n) \in \mathrm{dgnb}(G)$, we can find two strictly increasing natural numbers (n_i) and (m_j) such that $n_0 = 0, m_0 = 0$, and

$$G_0 \supseteq G'_{m_0} \supseteq G_{n_1} \supseteq G'_{m_1} \supseteq G_{n_2} \supseteq \ldots.$$

Define $H_{2i} = G_{n_i}$ and $H_{2i+1} = G'_{m_i}$ for each $i < \omega$. Then $(H_k) \in \mathrm{dgnb}(G)$. Put $\mathcal{H} = (H_k), \mathcal{K} = (G_{n_i})$, and $\mathcal{K}' = (G'_{m_i})$.

Note that (G_{n_i}) is a subsequence of (G_n) and also a subsequence of (H_k). From (1), we obtain

$$\omega(\rho(T^X_{\mathcal{G}})) = \omega(\rho(T^X_{\mathcal{K}})) = \omega(\rho(T^X_{\mathcal{H}})).$$

Similarly, we obtain

$$\omega(\rho(T^X_{\mathcal{G}'})) = \omega(\rho(T^X_{\mathcal{K}'})) = \omega(\rho(T^X_{\mathcal{H}})).$$

So we have $\omega(\rho(T^X_{\mathcal{G}})) = \omega(\rho(T^X_{\mathcal{G}'}))$. ⊣

Now we are going to find a special G-space $X(\mathcal{G})$ such that $\omega(\rho(T^{X(\mathcal{G})}_{\mathcal{G}}))$ attains the maximum value among all $\omega(\rho(T^X_{\mathcal{G}}))$. This leads to the conclusion that the value of $\omega(\rho(T^{X(\mathcal{G})}_{\mathcal{G}}))$ is determined by G itself.

LEMMA 3.8. *Given two sets X and Y. Let $\mathcal{E} = (E_n)$ and $\mathcal{F} = (F_n)$ be two decreasing sequences of equivalence relations on X and Y, respectively. Let $\theta : X \to Y$ be a surjection such that*

$$\forall n < \omega \, \forall x \in X \, (\theta([x]_{E_n}) = [\theta(x)]_{F_n}).$$

https://doi.org/10.1017/jsl.2024.7 Published online by Cambridge University Press

Then there exists a Lipschitz embedding $\psi : T^Y_{\mathcal{F}} \to T^X_{\mathcal{E}}$. In particular, if $T^X_{\mathcal{E}}$ is well-founded, so is $T^Y_{\mathcal{F}}$, and $\rho(T^Y_{\mathcal{F}}) \leq \rho(T^X_{\mathcal{E}})$ holds.

PROOF. For any $(n, C) \in T^Y_{\mathcal{F}}$, we construct $\psi(n, C)$ by induction on n such that $\psi(n, C) = (n, [x]_{E_n})$ for some $x \in X$ with $[\theta(x)]_{F_n} = C$.

For $n = 0$, since θ is a surjection, we can find an $x \in X$ with $\theta(x) \in C$. Then we let $\psi(0, C) = (0, [x]_{E_0})$.

For $n > 0$, since C is an F_n-equivalence class, there exists a unique F_{n-1}-equivalence class $D \supseteq C$. By the inductive hypothesis, we can find an $x' \in X$ such that $\psi(n-1, D) = (n-1, [x']_{E_{n-1}})$ with $[\theta(x')]_{F_{n-1}} = D$. Since $\theta([x']_{E_{n-1}}) = [\theta(x')]_{F_{n-1}} = D \supseteq C$, we can find an $x \in [x']_{E_{n-1}}$ such that $\theta(x) \in C$. Then we put $\psi(n, C) = (n, [x]_{E_n})$.

Since C is not a singleton, the equality $\theta([x]_{E_n}) = [\theta(x)]_{F_n} = C$ implies that $[x]_{E_n}$ is not a singleton. So $\psi(n, C) \in T^X_{\mathcal{E}}$. From the construction, it is routine to check that $\psi : T^Y_{\mathcal{F}} \to T^X_{\mathcal{E}}$ is a Lipschitz embedding.

Finally, if $T^X_{\mathcal{E}}$ is well-founded, then by Proposition 2.3, $T^Y_{\mathcal{F}}$ is well-founded as well, and $\rho(T^Y_{\mathcal{F}}) \leq \rho(T^X_{\mathcal{E}})$. ⊣

DEFINITION 3.9. Let G be a non-archimedean CLI Polish group, and let $\mathcal{G} = (G_n) \in \mathrm{dgnb}(G)$. For each $k < \omega$, we define an action $G \curvearrowright G/G_k$ as $g \cdot hG_k = ghG_k$ for $g, h \in G$, and let $\rho^k(\mathcal{G})$ denote $\rho(T^{G/G_k}_{\mathcal{G}})$. Afterwards, we let $X(\mathcal{G}) = \bigcup_k G/G_k$ and $\rho(\mathcal{G}) = \rho(T^{X(\mathcal{G})}_{\mathcal{G}})$.

Note that $G \cdot gG_k = \{hgG_k : h \in G\} = G/G_k$ for any $g \in G$.

It is clear that $\rho^0(\mathcal{G}) = 0$, since $G = G_0$.

LEMMA 3.10. (1) $\rho(\mathcal{G}) = \sup\{\rho^k(\mathcal{G}) : k < \omega\}$.

(2) $(\rho^k(\mathcal{G}))$ *is an increasing sequence of countable non-limit ordinals.*

PROOF. (1) Note that $L_0(T^{X(\mathcal{G})}_{\mathcal{G}}) = \{(0, G/G_k) : G \neq G_k\}$. We point out that $(T^{X(\mathcal{G})}_{\mathcal{G}})_{(0,G/G_k)} \cong T^{G/G_k}_{\mathcal{G}}$ for $G \neq G_k$, and $(T^{X(\mathcal{G})}_{\mathcal{G}})_{(0,G/G_k)} = T^{G/G_k}_{\mathcal{G}} = \emptyset$ for $G = G_k$. So

$$\begin{aligned}\rho(\mathcal{G}) &= \rho(T^{X(\mathcal{G})}_{\mathcal{G}}) = \sup\{\rho((T^{X(\mathcal{G})}_{\mathcal{G}})_{(0,G/G_k)}) : k < \omega\} \\ &= \sup\{\rho(T^{G/G_k}_{\mathcal{G}}) : k < \omega\} = \sup\{\rho^k(\mathcal{G}) : k < \omega\}.\end{aligned}$$

(2) Given $k < \omega$, we define $\theta : G/G_{k+1} \to G/G_k$ as $\theta(gG_{k+1}) = gG_k$ for $g \in G$. It is clear that θ is well-defined and is a surjection. Furthermore, for $n < \omega$ and $g \in G$, we have

$$\begin{aligned}\theta(G_n \cdot gG_{k+1}) &= \{\theta(hgG_{k+1}) : h \in G_n\} \\ &= \{hgG_k : h \in G_n\} = G_n \cdot gG_k = G_n \cdot \theta(gG_{k+1}).\end{aligned}$$

From Lemma 3.8, we have

$$\rho(T^{G/G_k}_{\mathcal{G}}) \leq \rho(T^{G/G_{k+1}}_{\mathcal{G}}),$$

i.e., $(\rho^k(\mathcal{G}))$ is increasing.

https://doi.org/10.1017/jsl.2024.7 Published online by Cambridge University Press

For each $k < \omega$, since $T_{\mathcal{G}}^{G/G_k}$ is countable, $\rho^k(\mathcal{G})$ is countable as well. If $G = G_k$, then $T_{\mathcal{G}}^{G/G_k} = \emptyset$; otherwise if $G \neq G_k$, then $L_0(T_{\mathcal{G}}^{G/G_k}) = \{(0, G/G_k)\}$ is a singleton. So $\rho^k(\mathcal{G}) = \rho(T_{\mathcal{G}}^{G/G_k})$ is either 0 or a successor ordinal. ⊣

Recall that a G-space X is said to be *transitive* if X itself is an orbit.

LEMMA 3.11. *Let G be a non-archimedean CLI Polish group, X a countable discrete transitive Polish G-space, and let $\mathcal{G} = (G_n) \in \mathrm{dgnb}(G)$. Then we can find some $k < \omega$ such that $\rho(T_{\mathcal{G}}^X) \leq \rho^k(\mathcal{G})$.*

PROOF. Fix an $x \in X$. Since $\{x\}$ is clopen in X, by the continuity of the group action of G on X, we know G_x is a clopen subgroup of G. So there is some $k < \omega$ such that $G_k \subseteq G_x$. Then we can define $\theta : G/G_k \to X$ as $\theta(gG_k) = g \cdot x$ for $g \in G$. Since X is a transitive G-space, θ is surjective and $\theta(G_n \cdot gG_k) = G_n \cdot \theta(gG_k)$ for each $n < \omega$ and $g \in G$. By Lemma 3.8, we have $\rho(T_{\mathcal{G}}^X) \leq \rho^k(\mathcal{G})$. ⊣

COROLLARY 3.12. *Let G be a non-archimedean CLI Polish group, X a countable discrete Polish G-space, and let $\mathcal{G} = (G_n) \in \mathrm{dgnb}(G)$. Then we have*

$$\rho(T_{\mathcal{G}}^X) \leq \rho(\mathcal{G}).$$

PROOF. Note that $L_0(T_{\mathcal{G}}^X) = \{(0, G \cdot x) : x \in X \wedge G \cdot x \neq \{x\}\}$ and $T_{\mathcal{G}}^{G \cdot x} \cong (T_{\mathcal{G}}^X)_{(0, G \cdot x)}$ for $G \cdot x \neq \{x\}$, so we have

$$\rho(T_{\mathcal{G}}^X) = \sup\{\rho((T_{\mathcal{G}}^X)_{(0,G\cdot x)}) : x \in X\} = \sup\{\rho(T_{\mathcal{G}}^{G\cdot x}) : x \in X\}.$$

By Lemma 3.11, we have

$$\rho(T_{\mathcal{G}}^X) \leq \sup\{\rho^k(\mathcal{G}) : k < \omega\} = \rho(\mathcal{G}).$$ ⊣

COROLLARY 3.13. *Let G be a non-archimedean Polish group, $\mathcal{G} = (G_n) \in \mathrm{dgnb}(G)$. Then G is CLI iff $T_{\mathcal{G}}^{G/G_k}$ is well-founded for any $k < \omega$.*

PROOF. The ($\Rightarrow$) part follows from Lemma 3.3.

($\Leftarrow$). Given a countable Polish G-space X. Following the arguments in the proof of Lemma 3.11, we can see that, for any $x \in X$, there is a $k < \omega$ and a Lipschitz embedding from $T_{\mathcal{G}}^{G\cdot x}$ to $T_{\mathcal{G}}^{G/G_k}$. Since $T_{\mathcal{G}}^{G/G_k}$ is well-founded, so is $T_{\mathcal{G}}^{G\cdot x}$. By the arbitrariness of $x \in X$, we conclude that $T_{\mathcal{G}}^X$ is also well-founded. Combining this with [5, Theorem 6], it follows that G is CLI. ⊣

THEOREM 3.14. *Let G be a non-archimedean CLI Polish group, and $\mathcal{G} = (G_n), \mathcal{G}' = (G'_n) \in \mathrm{dgnb}(G)$. Then $\omega(\rho(\mathcal{G})) = \omega(\rho(\mathcal{G}'))$ holds.*

PROOF. By Corollary 3.12, we have $\rho(T_{\mathcal{G}'}^{X(\mathcal{G})}) \leq \rho(\mathcal{G}')$. Thus Lemma 3.7 implies

$$\omega(\rho(\mathcal{G})) = \omega(\rho(T_{\mathcal{G}}^{X(\mathcal{G})})) = \omega(\rho(T_{\mathcal{G}'}^{X(\mathcal{G})})) \leq \omega(\rho(\mathcal{G}')),$$

and vice versa. ⊣

By the preceding theorem, there is a unique ordinal $\beta < \omega_1$, which is independent from the choice of $\mathcal{G} = (G_n) \in \mathrm{dgnb}(G)$, such that

$$\omega(\rho(\mathcal{G})) = \omega \cdot \beta.$$

https://doi.org/10.1017/jsl.2024.7 Published online by Cambridge University Press

Consequently, we can define a rank for a given non-archimedean CLI Polish group as follows:

DEFINITION 3.15. Suppose G is a non-archimedean CLI Polish group. We define

$$\mathrm{rank}(G) = \text{the unique } \beta \text{ such that } \omega(\rho(\mathcal{G})) = \omega \cdot \beta$$

for some $\mathcal{G} \in \mathrm{dgnb}(G)$.

LEMMA 3.16. (1) *If* $\rho(\mathcal{G}) = \omega \cdot \mathrm{rank}(G)$, *then either* $\mathrm{rank}(G) = 0$ *or* $\rho^k(\mathcal{G}) < \omega \cdot \mathrm{rank}(G)$ *for any* $k < \omega$.
(2) *If* $\rho(\mathcal{G}) > \omega \cdot \mathrm{rank}(G)$, *then there exists an* $m > 0$ *such that* $\rho^k(\mathcal{G}) = \omega \cdot \mathrm{rank}(G) + m$ *for large enough* $k < \omega$.

PROOF. (1) If $\mathrm{rank}(G) > 0$, then $\omega \cdot \mathrm{rank}(G)$ is a limit ordinal. By Lemma 3.10, $\rho^k(\mathcal{G})$ is either 0 or a successor ordinal, so $\rho^k(\mathcal{G}) < \omega \cdot \mathrm{rank}(G)$ for any $k < \omega$.
(2) Clearly, $\rho(\mathcal{G}) = \omega \cdot \mathrm{rank}(G) + m$ for some $0 < m < \omega$. Again by Lemma 3.10, $(\rho^k(\mathcal{G}))$ is increasing, so $\rho^k(\mathcal{G}) = \omega \cdot \mathrm{rank}(G) + m$ for large enough $k < \omega$. ⊣

THEOREM 3.17. *Let* G *be a non-archimedean CLI Polish group,* $\mathcal{G} = (G_n), \mathcal{G}' = (G'_n) \in \mathrm{dgnb}(G)$. *Then* $\rho(\mathcal{G}) = \omega \cdot \mathrm{rank}(G)$ *iff* $\rho(\mathcal{G}') = \omega \cdot \mathrm{rank}(G)$.

PROOF. If $\mathrm{rank}(G) = 0$, then $\rho(\mathcal{G}) = \omega \cdot \mathrm{rank}(G)$ implies that $\rho^k(\mathcal{G}) = 0$ for any $k < \omega$. So $T_{\mathcal{G}}^{G/G_k} = \emptyset$, and hence $G_0 \cdot G_k \notin T_{\mathcal{G}}^{G/G_k}$. It follows that $G_0 \cdot G_k = \{G_k\}$, i.e., $G = G_0 = G_k$ for any $k < \omega$. This implies that $G = \{1_G\}$. Then we can easily see that $T_{\mathcal{G}'}^{G/G'_k} = \emptyset$ for $k < \omega$. So $\rho(\mathcal{G}') = \omega \cdot \mathrm{rank}(G)$ holds. And vice versa.

If $\mathrm{rank}(G) > 0$, assume for contradiction that $\rho(\mathcal{G}) = \omega \cdot \mathrm{rank}(G)$, but $\rho(\mathcal{G}') > \omega \cdot \mathrm{rank}(G)$. From Lemma 3.16, we have $\rho^k(\mathcal{G}) < \omega \cdot \mathrm{rank}(G)$ for any $k < \omega$, but $\rho^l(\mathcal{G}') = \omega \cdot \mathrm{rank}(G) + m$ for some $0 < m < \omega$ and large enough $l < \omega$. From Lemmas 3.7 and 3.11, we can conclude that for any $l < \omega$, there is some $k < \omega$ such that

$$\omega(\rho^l(\mathcal{G}')) = \omega(\rho(T_{\mathcal{G}'}^{G/G'_l})) = \omega(\rho(T_{\mathcal{G}}^{G/G'_l})) \leq \rho(T_{\mathcal{G}}^{G/G'_l}) \leq \rho^k(\mathcal{G}).$$

A contradiction! ⊣

Now we are ready to define a hierarchy on the class of non-archimedean CLI Polish groups.

DEFINITION 3.18. Let G be a non-archimedean CLI Polish group, $\mathcal{G} = (G_n) \in \mathrm{dgnb}(G)$, and let $\alpha < \omega_1$ be an ordinal.

(1) If $\rho(\mathcal{G}) \leq \omega \cdot \alpha$, we say that G is *α-CLI*.
(2) If $\omega(\rho(\mathcal{G})) \leq \omega \cdot \alpha$, i.e., $\mathrm{rank}(G) \leq \alpha$, we say that G is *L-α-CLI*.

It is clear that, if G is L-α-CLI, then it is also $(\alpha + 1)$-CLI.

From Theorems 3.14 and 3.17, we see that the definitions of α-CLI and L-α-CLI are independent from the choice of $\mathcal{G} \in \mathrm{dgnb}(G)$.

Recall that a metric d on a group G is *two-sided invariant* if $d(gh, gk) = d(h, k) = d(hg, kg)$ for all $g, h, k \in G$. A Polish group is *TSI* if it admits a compatible complete two-sided invariant metric.

https://doi.org/10.1017/jsl.2024.7 Published online by Cambridge University Press

THEOREM 3.19. *Let G be a non-archimedean CLI Polish group. The following hold*:
(1) *G is 0-CLI iff $G = \{1_G\}$*;
(2) *G is L-0-CLI iff G is discrete*; *and*
(3) *G is 1-CLI iff G is TSI.*

PROOF. Fix a sequence $\mathcal{G} = (G_n) \in \mathrm{dgnb}(G)$.
(1) It follows from the first paragraph of the proof of Theorem 3.17.
(2) If G is L-0-CLI, then $\mathrm{rank}(G) = 0$. So there is an $m < \omega$ such that $\rho(T_{\mathcal{G}}^{G/G_k}) = \rho^k(\mathcal{G}) = m$ for large enough $k < \omega$. Then we have $L_m(T_{\mathcal{G}}^{G/G_k}) = \emptyset$. This implies that $G_m \cdot G_k = \{G_k\}$. So $G_m \subseteq G_k$ for large enough $k < \omega$, and thus $G_m = \{1_G\}$. It follows that G is discrete.
On the other hand, if G is discrete, then there is an $m < \omega$ such that $G_m = \{1_G\}$. Therefore, for any $k < \omega$, we have $L_m(T_{\mathcal{G}}^{G/G_k}) = \emptyset$, and hence $\rho^k(\mathcal{G}) \leq m$. This implies that $\mathrm{rank}(G) = 0$, i.e., G is L-0-CLI.
(3) If G is 1-CLI, then Lemma 3.16 implies that $\rho^k(\mathcal{G}) < \omega$ for any $k < \omega$. So there is an $m_k < \omega$ such that $L_{m_k}(T_{\mathcal{G}}^{G/G_k}) = \emptyset$. It follows that, for $g \in G$, $G_{m_k} \cdot gG_k = \{gG_k\}$, so $g^{-1}G_{m_k}g \subseteq G_k$. Put $U_k = \bigcup\{g^{-1}G_{m_k}g : g \in G\} \subseteq G_k$. We can see that (U_k) is a neighborhood basis of 1_G with $g^{-1}U_kg = U_k$ for all $g \in G$. By Klee's theorem (cf. [4] or [2, Exercise 2.1.4]), G is TSI.
On the other hand, if G is TSI, again by Klee's theorem, we can find a neighborhood basis (U_m) of 1_G with $g^{-1}U_mg = U_m$ for all $g \in G$. For any $n < \omega$, there is an $m_n < \omega$ such that $U_{m_n} \subseteq G_n$. Let $V_n = U_{m_n} \cap U_{m_n}^{-1}$ and $G'_n = \bigcup_i V_n^i$. Then G'_n is an open normal subgroup of G with $G'_n \subseteq G_n$. So $(G'_n) \in \mathrm{dgnb}(G)$. Let $\mathcal{G}' = (G'_n)$. Then $G'_k \cdot gG'_k = \{gG'_k\}$ for all $g \in G$ and $k < \omega$, thus $L_k(T_{\mathcal{G}'}^{G/G'_k}) = \emptyset$. So $\rho^k(\mathcal{G}') \leq k < \omega$, and hence G is 1-CLI. ⊣

Clause (2) in the preceding theorem can be generalized to all $\alpha < \omega_1$.

DEFINITION 3.20. Let G be a non-archimedean CLI Polish group, and $\alpha < \omega_1$. We say that G is *locally α-CLI* if G has an open subgroup which is α-CLI.

THEOREM 3.21. *Let G be a non-archimedean CLI Polish group, and $\alpha < \omega_1$. Then G is L-α-CLI iff G is locally α-CLI.*

PROOF. ($\Rightarrow$). Suppose G is L-α-CLI. Without loss of generality, we may assume that G is not α-CLI. Fix a sequence $\mathcal{G} = (G_n) \in \mathrm{dgnb}(G)$. There exists an $m \geq 1$ such that $\rho(\mathcal{G}) = \omega \cdot \alpha + m$, and thus we can pick a $k_0 > m$ such that $\rho^k(\mathcal{G}) = \omega \cdot \alpha + m$ for any $k \geq k_0$. We will show that G_{k_0} is α-CLI.
Let $H = G_{k_0}$ and $H_n = G_{n+k_0}$ for $n < \omega$. Then $(H_n) \in \mathrm{dgnb}(H)$. Put $\mathcal{H} = (H_n)$. Given $k < \omega$, define $\phi : T_{\mathcal{H}}^{H/H_k} \to (T_{\mathcal{G}}^{G/G_{k+k_0}})_{(k_0, G_{k_0}/G_{k+k_0})}$ as $\phi(n, C) = (n + k_0, C)$. It is trivial to see that ϕ is an order-preserving isomorphism. From Proposition 2.2, since $k_0 > m$, we have

$$\rho^k(\mathcal{H}) = \rho(T_{\mathcal{H}}^{H/H_k}) = \rho((T_{\mathcal{G}}^{G/G_{k+k_0}})_{(k_0, G_{k_0}/G_{k+k_0})}) \leq \omega \cdot \alpha.$$

So $\rho(\mathcal{H}) \leq \omega \cdot \alpha$, and hence $H = G_{k_0}$ is α-CLI.
($\Leftarrow$). Suppose G is locally α-CLI. Let H be an open subgroup of G which is α-CLI, and let $\mathcal{H} = (H_n) \in \mathrm{dgnb}(H)$. Then $\rho^k(\mathcal{H}) \leq \omega \cdot \alpha$ for $k < \omega$.

https://doi.org/10.1017/jsl.2024.7 Published online by Cambridge University Press

Put $G_0 = G$ and $G_n = H_{n-1}$ for $n \geq 1$. Then $(G_n) \in \mathrm{dgnb}(G)$. Put $\mathcal{G} = (G_n)$. Given $g \in G$ and $k < \omega$, by similar arguments in the $(\Rightarrow)$ part, we have $T_{\mathcal{H}}^{H \cdot gH_k} \cong (T_{\mathcal{G}}^{G/G_{k+1}})_{(1, G_1 \cdot gG_{k+1})}$. By Lemma 3.11, there exists an $l < \omega$ such that $\rho(T_{\mathcal{H}}^{H \cdot gH_k}) \leq \rho^l(\mathcal{H})$. Therefore, by Proposition 2.1,

$$\begin{aligned}
\rho^{k+1}(\mathcal{G}) &= \rho(T_{\mathcal{G}}^{G/G_{k+1}}) \\
&\leq \sup\{\rho((T_{\mathcal{G}}^{G/G_{k+1}})_{(1, G_1 \cdot gG_{k+1})}) : g \in G\} + 1 \\
&= \sup\{\rho(T_{\mathcal{H}}^{H \cdot gH_k}) : g \in G\} + 1 \\
&\leq \sup\{\rho^l(\mathcal{H}) : l < \omega\} + 1 \\
&\leq \omega \cdot \alpha + 1.
\end{aligned}$$

So $\rho(\mathcal{G}) \leq \omega \cdot \alpha + 1$, and hence G is L-α-CLI. ⊣

§4. Properties of the hierarchy

THEOREM 4.1. *Let G be a non-archimedean CLI Polish group, H a closed subgroup of G, and $\alpha < \omega_1$. If G is α-CLI (or L-α-CLI), so is H. In particular, we have* $\mathrm{rank}(H) \leq \mathrm{rank}(G)$.

PROOF. Let $\mathcal{G} = (G_n) \in \mathrm{dgnb}(G)$, and put $H_n = H \cap G_n$ for $n < \omega$. It is clear that $(H_n) \in \mathrm{dgnb}(H)$. Put $\mathcal{H} = (H_n)$. We only need to show that $\rho(\mathcal{H}) \leq \rho(\mathcal{G})$.

Given $k < \omega$, define $\theta : H/H_k \to G/G_k$ as $\theta(hH_k) = hG_k$ for $h \in H$. By Proposition 3.4, there is a Lipschitz order-preserving map from $T_{\mathcal{H}}^{H/H_k}$ to $T_{\mathcal{G}}^{G/G_k}$. So Proposition 2.3 implies

$$\rho^k(\mathcal{H}) = \rho(T_{\mathcal{H}}^{H/H_k}) \leq \rho(T_{\mathcal{G}}^{G/G_k}) = \rho^k(\mathcal{G}).$$

Then we have $\rho(\mathcal{H}) \leq \rho(\mathcal{G})$ as desired. ⊣

THEOREM 4.2. *Let G be a non-archimedean CLI Polish group, N a closed normal subgroup of G, and $\alpha < \omega_1$. If G is α-CLI (or L-α-CLI), so is G/N. In particular, we have* $\mathrm{rank}(G/N) \leq \mathrm{rank}(G)$.

PROOF. Let $\mathcal{G} = (G_n) \in \mathrm{dgnb}(G)$, and put $H_n = G_n \cdot N = \{\hat{g}N : \hat{g} \in G_n\}$ for $n < \omega$. It is clear that $H_0 = G/N$ and $(H_n) \in \mathrm{dgnb}(G/N)$. Put $\mathcal{H} = (H_n)$. We only need to show that $\rho(\mathcal{H}) \leq \rho(\mathcal{G})$.

Given $k < \omega$, define $\theta : G/G_k \to (G/N)/H_k$ as $\theta(gG_k) = (gN)H_k$ for $g \in G$. Note that for $n < \omega$,

$$\begin{aligned}
\theta(G_n \cdot gG_k) &= \theta(\{\hat{g}gG_k : \hat{g} \in G_n\}) \\
&= \{(\hat{g}gN)H_k : \hat{g} \in G_n\} \\
&= \{(\hat{g}N)(gN)H_k : \hat{g} \in G_n\} \\
&= \{(\hat{g}N)\theta(gG_k) : \hat{g} \in G_n\} \\
&= \{\hat{g}N : \hat{g} \in G_n\}\theta(gG_k) = H_n \cdot \theta(gG_k).
\end{aligned}$$

By Lemma 3.8, we have

$$\rho^k(\mathcal{H}) = \rho(T_{\mathcal{H}}^{(G/N)/H_k}) \leq \rho(T_{\mathcal{G}}^{G/G_k}) = \rho^k(\mathcal{G}).$$

So $\rho(\mathcal{H}) \leq \rho(\mathcal{G})$ holds as desired. ⊣

https://doi.org/10.1017/jsl.2024.7 Published online by Cambridge University Press

The above two theorems involve closed subgroups and quotient groups. Now we turn to product groups, which are more complicated. We discuss finite product groups first.

LEMMA 4.3. *Let X, Y be two sets, $\mathcal{E} = (E_n)$ and $\mathcal{F} = (F_n)$ be two decreasing sequences of equivalence relations on X and Y, respectively. Let $\mathcal{E} \times \mathcal{F} = (E_n \times F_n)$.*

(1) *$T^{X\times Y}_{\mathcal{E}\times\mathcal{F}}$ is well-founded iff $T^X_{\mathcal{E}}$ and $T^Y_{\mathcal{F}}$ are well-founded.*

(2) *If $T^{X\times Y}_{\mathcal{E}\times\mathcal{F}}$ is well-founded, then we have*

$$\rho(T^{X\times Y}_{\mathcal{E}\times\mathcal{F}}) = \max\{\rho(T^X_{\mathcal{E}}), \rho(T^Y_{\mathcal{F}})\}.$$

PROOF. First, note that $[(x,y)]_{E_n\times F_n} = [x]_{E_n} \times [y]_{F_n}$ for all $(x,y) \in X \times Y$ and $n < \omega$.

(1) For any sequences $(x_n), (y_n)$ in X, Y, respectively, $((n, [(x_n, y_n)]_{E_n\times F_n}))$ is an infinite branch of $T^{X\times Y}_{\mathcal{E}\times\mathcal{F}}$ iff either $((n, [x_n]_{E_n}))$ or $((n, [y_n]_{F_n}))$ is an infinite branch of $T^X_{\mathcal{E}}$ or $T^Y_{\mathcal{F}}$, respectively.

(2) If $T^{X\times Y}_{\mathcal{E}\times\mathcal{F}}$ is well-founded, by (1), $T^X_{\mathcal{E}}$ and $T^Y_{\mathcal{F}}$ are also well-founded. For all $(x,y) \in X \times Y$ and $n < \omega$, note that

$$\begin{aligned} & (n, [(x,y)]_{E_n\times F_n}) \in T^{X\times Y}_{\mathcal{E}\times\mathcal{F}} \\ \iff\ & [(x,y)]_{E_n\times F_n} \neq \{(x,y)\} \\ \iff\ & [x]_{E_n} \neq \{x\} \vee [y]_{F_n} \neq \{y\} \\ \iff\ & (n, [x]_{E_n}) \in T^X_{\mathcal{E}} \vee (n, [y]_{F_n}) \in T^Y_{\mathcal{F}}. \end{aligned}$$

By Proposition 2.1, it is routine to prove

$$\rho((T^{X\times Y}_{\mathcal{E}\times\mathcal{F}})_{(n,[(x,y)]_{E_n\times F_n})}) = \max\{\rho((T^X_{\mathcal{E}})_{(n,[x]_{E_n})}), \rho((T^Y_{\mathcal{F}})_{(n,[y]_{F_n})})\}$$

by induction on $\rho((T^{X\times Y}_{\mathcal{E}\times\mathcal{F}})_{(n,[(x,y)]_{E_n\times F_n})})$. Taking supremum on both sides of the above formula, we get

$$\rho(T^{X\times Y}_{\mathcal{E}\times\mathcal{F}}) = \max\{\rho(T^X_{\mathcal{E}}), \rho(T^Y_{\mathcal{F}})\}. \qquad \dashv$$

COROLLARY 4.4. *Let G and H be two non-archimedean CLI Polish groups, $\mathcal{G} = (G_n) \in \mathrm{dgnb}(G)$ and $\mathcal{H} = (H_n) \in \mathrm{dgnb}(H)$. Then we have $\mathcal{G} \times \mathcal{H} = (G_n \times H_n) \in \mathrm{dgnb}(G \times H)$ and*

$$\rho^k(\mathcal{G} \times \mathcal{H}) = \max\{\rho^k(\mathcal{G}), \rho^k(\mathcal{H})\} \quad (\forall k < \omega),$$

$$\rho(\mathcal{G} \times \mathcal{H}) = \max\{\rho(\mathcal{G}), \rho(\mathcal{H})\},$$

$$\mathrm{rank}(G \times H) = \max\{\mathrm{rank}(G), \mathrm{rank}(H)\}.$$

PROOF. For any $k < \omega$, put $X = G/G_k$, $Y = H/H_k$, and define E_n, F_n on X and Y for each $n < \omega$, respectively, by

$$(\hat{g}G_k, \tilde{g}G_k) \in E_n \iff \exists g \in G_n\, (g\hat{g}G_k = \tilde{g}G_k) \quad (\forall \hat{g}, \tilde{g} \in G),$$

$$(\hat{h}H_k, \tilde{h}H_k) \in F_n \iff \exists h \in H_n\, (h\hat{h}H_k = \tilde{h}H_k) \quad (\forall \hat{h}, \tilde{h} \in H).$$

https://doi.org/10.1017/jsl.2024.7 Published online by Cambridge University Press

Then Lemma 4.3 gives $\rho^k(\mathcal{G}\times\mathcal{H}) = \max\{\rho^k(\mathcal{G}), \rho^k(\mathcal{H})\}$. The rest follows trivially. ⊣

Now we are ready to discuss countably infinite product groups.

LEMMA 4.5. *Let (G^i) be a sequence of non-archimedean CLI Polish groups and $\mathcal{G}^i = (G^i_n) \in \mathrm{dgnb}(G^i)$ for each $i<\omega$. We define $G = \prod_i G^i$ and*

$$G_n = \prod_{i<n} G^i_n \times \prod_{i\geq n} G^i \quad (\forall n<\omega).$$

Then $\mathcal{G} = (G_n) \in \mathrm{dgnb}(G)$ and for $k<\omega$, we have

$$\rho^k(\mathcal{G}) \leq \max\{\rho^k(\mathcal{G}^i) : i<k\} + k.$$

PROOF. Given $k<\omega$, we put $Y = G^0/G^0_k \times \cdots \times G^{k-1}/G^{k-1}_k$ and define $\theta : G/G_k \to Y$ as

$$\theta((g_i)G_k) = (g_0G^0_k, \dots, g_{k-1}G^{k-1}_k)$$

for $(g_i)\in G = \prod_i G^i$. By the definition of G_k, it is easy to see that θ is a bijection. Moreover, we put

$$H_n = \begin{cases} \prod_{i<n} G^i_n \times \prod_{n\leq i<k} G^i, & n<k,\\ \prod_{i<k} G^i_n, & n\geq k,\end{cases}$$

then θ is a reduction of $E^{G/G_k}_{G_n}$ to $E^Y_{H_n}$ for each $n<\omega$. Put $\mathcal{H} = (H_n)$. By Proposition 3.4, $(n, G_n\cdot(g_i)G_k) \mapsto (n, H_n\cdot\theta((g_i)G_k))$ is an order-preserving isomorphism from $T^{G/G_k}_{\mathcal{G}}$ to $T^Y_{\mathcal{H}}$. So $\rho^k(\mathcal{G}) = \rho(T^{G/G_k}_{\mathcal{G}}) = \rho(T^Y_{\mathcal{H}})$.

For $n\geq k$ and $(g_i)\in G$, we have

$$H_n\cdot\theta((g_i)G_k) = (G^0_n\cdot g_0G^0_k)\times\cdots\times(G^{k-1}_n\cdot g_{k-1}G^{k-1}_k).$$

Lemma 4.3 gives

$$\rho((T^Y_{\mathcal{H}})_{(k,H_k\cdot\theta((g_i)G_k))}) = \max\{\rho((T^{G^i/G^i_k}_{\mathcal{G}^i})_{(k,G^i_k\cdot g_iG^i_k)}) : i<k\}.$$

Hence, by Proposition 2.2(1),

$$\begin{aligned}\rho^k(\mathcal{G}) &= \rho(T^{G/G_k}_{\mathcal{G}}) = \rho(T^Y_{\mathcal{H}})\\ &\leq \sup\{\rho((T^Y_{\mathcal{H}})_{(k,H_k\cdot\theta((g_i)G_k))}) : (g_i)\in G\} + k\\ &= \sup\{\max\{\rho((T^{G^i/G^i_k}_{\mathcal{G}^i})_{(k,G^i_k\cdot g_iG^i_k)}) : i<k\} : (g_i)\in G\} + k\\ &= \max\{\sup\{\rho((T^{G^i/G^i_k}_{\mathcal{G}^i})_{(k,G^i_k\cdot gG^i_k)}) : g\in G^i\} : i<k\} + k\\ &\leq \max\{\rho(T^{G^i/G^i_k}_{\mathcal{G}^i}) : i<k\} + k\\ &= \max\{\rho^k(\mathcal{G}^i) : i<k\} + k.\end{aligned}$$ ⊣

THEOREM 4.6. *Let (G^i) be a sequence of non-archimedean CLI Polish groups, $\alpha<\omega_1$, and let $G = \prod_i G^i$. Then we have*:

(1) *G is α-CLI iff all G^i are α-CLI; and*
(2) *G is L-α-CLI iff all G^i are L-α-CLI and for all but finitely many i, G^i is α-CLI.*

https://doi.org/10.1017/jsl.2024.7 Published online by Cambridge University Press

PROOF. Fix a $\mathcal{G}^i = (G^i_n) \in \mathrm{dgnb}(G^i)$ for each $i < \omega$. Put

$$G_n = \prod_{i<n} G^i_n \times \prod_{i\geq n} G^i \quad (\forall n < \omega).$$

(1) Assume G is α-CLI. Since each G^i is topologically isomorphic to a closed subgroup of G, by Theorem 4.1, G^i is also α-CLI.

On the other hand, assume all G^i are α-CLI. Now by Lemma 3.16(1), $\rho^k(\mathcal{G}^i) < \omega \cdot \alpha$ holds for all $i, k < \omega$. Then Lemma 4.5 implies

$$\rho^k(\mathcal{G}) \leq \max\{\rho^k(\mathcal{G}^i) : i < k\} + k < \omega \cdot \alpha$$

holds for each $k < \omega$. Consequently, G is α-CLI.

(2) Assume G is L-α-CLI. Since each G^i is topologically isomorphic to a closed subgroup of G, we can see G^i is L-α-CLI too. Moreover, by Theorem 3.21, there is an open subgroup H of G which is α-CLI. Hence, G_n is a clopen subgroup of H for some $n < \omega$, from which we can see this G_n is also α-CLI. Now by (1), we conclude that for all $i \geq n$, G^i is α-CLI.

On the other hand, assume all G^i are L-α-CLI, and there is an $m < \omega$ such that G^i is α-CLI for each $i \geq m$. Then (1) implies that $\prod_{i\geq m} G^i$ is α-CLI. Note that $G = G^0 \times \cdots \times G^{m-1} \times \prod_{i\geq m} G^i$. By Corollary 4.4, we have

$$\mathrm{rank}(G) = \max\{\mathrm{rank}(G^0), \ldots, \mathrm{rank}(G^{m-1}), \mathrm{rank}(\prod_{i\geq m} G^i)\} \leq \alpha,$$

i.e., G is L-α-CLI. ⊣

In the rest of this article, we will show that the notions of α-CLI, L-α-CLI, together with $\mathrm{rank}(G)$, form a proper hierarchy on the class of non-archimedean CLI Polish groups. For this purpose, we will construct groups which are α-CLI but not L-β-CLI for all $\beta < \alpha$, and groups which are L-α-CLI but not α-CLI, by induction on $\alpha < \omega_1$. We consider the case concerning successor ordinals first.

COROLLARY 4.7. *Let (G^i) be a sequence of non-archimedean CLI Polish groups, $\alpha < \omega_1$, and let $G = \prod_i G^i$. If all G^i are L-α-CLI but not α-CLI, then G is $(\alpha+1)$-CLI but not L-α-CLI.*

PROOF. Note that all G^i are $(\alpha+1)$-CLI but not α-CLI. ⊣

From Theorem 3.19 and [3, Theorem 1.1], a non-archimedean CLI Polish group G is 1-CLI iff G is isomorphic to a closed subgroup of a product $\prod_i G^i$, where each G^i is L-0-CLI. However, we do not know whether the following generalization of Corollary 4.7 is true:

QUESTION 4.8. *Let $0 < \alpha < \omega_1$, and let G be an $(\alpha+1)$-CLI group. Can we find a sequence of L-α-CLI groups G^i such that G is isomorphic to a closed subgroup of $\prod_i G^i$?*

Let G and Λ be two groups. Recall that the *wreath product* $\Lambda \wr G$ is the set $\Lambda \times G^\Lambda$ with the following group operation: given $(\hat{\lambda}, \hat{\chi}), (\tilde{\lambda}, \tilde{\chi}) \in \Lambda \times G^\Lambda$, we have

$$(\hat{\lambda}, \hat{\chi})(\tilde{\lambda}, \tilde{\chi}) = (\hat{\lambda}\tilde{\lambda}, \chi),$$

https://doi.org/10.1017/jsl.2024.7 Published online by Cambridge University Press

where $\chi(\lambda) = \hat{\chi}(\lambda)\tilde{\chi}(\hat{\lambda}^{-1}\lambda)$ for each $\lambda \in \Lambda$. If Λ is countable discrete and G is Polish, then $\Lambda \wr G$ equipped with the product topology of $\Lambda \times G^{\Lambda}$ is also a Polish group.

THEOREM 4.9. *Let G be a non-archimedean CLI Polish group, Λ an infinite countable discrete group, and $\alpha < \omega_1$. If G is $(\alpha+1)$-CLI but not α-CLI, then $\Lambda \wr G$ is L-$(\alpha+1)$-CLI but not $(\alpha+1)$-CLI.*

PROOF. For brevity, we denote $\Lambda \wr G$ by H. Let λ_i, $i < \omega$ be an enumeration of Λ without repetition. Note that the underlying space of $\Lambda \wr G$ is $\Lambda \times G^{\Lambda}$. For $(\lambda, \chi) \in \Lambda \wr G$, put $\pi_{\Lambda}(\lambda, \chi) = \lambda$ and $\pi^i_G(\lambda, \chi) = \chi(\lambda_i)$.

Let $\mathcal{G} = (G_n) \in \mathrm{dgnb}(G)$. Since G is not α-CLI, $G \neq \{1_G\}$. Without loss of generality, we can assume that $G \neq G_1$. Put $H_0 = H$ and

$$H_{n+1} = \{(1_{\Lambda}, \chi) : \chi \in G^{\Lambda} \wedge \forall i < n\, (\chi(\lambda_i) \in G_n)\}$$

for $n < \omega$. Then $\mathcal{H} = (H_n) \in \mathrm{dgnb}(H)$. Note that the open subgroup $H_1 = \{1_{\Lambda}\} \times G^{\Lambda}$ is topologically isomorphic to G^{ω}. By Theorem 4.6, H_1 is $(\alpha+1)$-CLI. So we can see that H is L-$(\alpha+1)$-CLI from Theorem 3.21.

Since G is $(\alpha+1)$-CLI but not α-CLI, $\omega \cdot \alpha < \rho(\mathcal{G}) \leq \omega \cdot (\alpha+1)$. By Lemma 3.16, there exist $1 \leq m, k < \omega$ such that $\rho^k(\mathcal{G}) = \omega \cdot \alpha + m$. To see that H is not $(\alpha+1)$-CLI, we will show that $\rho^{k+1}(\mathcal{H}) > \omega \cdot (\alpha+1)$ as follows.

For any $(\lambda_l, \hat{\chi}) \in H$ and $(1_{\Lambda}, \tilde{\chi}) \in H_{k+1}$, note that $(\lambda_l, \hat{\chi})(1_{\Lambda}, \tilde{\chi}) = (\lambda_l, \hat{\chi}\tilde{\chi}_l)$, where $\tilde{\chi}_l(\lambda) = \tilde{\chi}(\lambda_l^{-1}\lambda)$ for $\lambda \in \Lambda$. It follows that

$$(\lambda_l, \hat{\chi})H_{k+1} = \{(\lambda_l, \chi) : \chi \in G^{\Lambda} \wedge \forall i < k\, (\chi(\lambda_l\lambda_i) \in \hat{\chi}(\lambda_l\lambda_i)G_k)\}.$$

There is a unique $l_i < \omega$ such that $\lambda_{l_i} = \lambda_l\lambda_i$ for $i < k$. It is clear that $\pi_{\Lambda}((\lambda_l, \hat{\chi})H_{k+1}) = \{\lambda_l\}$ and for $j < \omega$,

$$\pi^j_G((\lambda_l, \hat{\chi})H_{k+1}) = \begin{cases} \hat{\chi}(\lambda_{l_i})G_k, & j = l_i, i < k, \\ G, & \text{otherwise.} \end{cases}$$

Let $m_l = \max\{l_i : i < k\}$. Then it is clear that $m_l \geq k-1$ for $l < \omega$ and $\sup\{m_l : l < \omega\} = \omega$. Note that $\pi^{m_l}_G(H_{m_l+1} \cdot (\lambda_l, \hat{\chi})H_{k+1}) = G/G_k$ is not a singleton, and so $(m_l+1, H_{m_l+1} \cdot (\lambda_l, \hat{\chi})H_{k+1}) \in T^{H/H_{k+1}}_{\mathcal{H}}$. Also note that

$$(T^{H/H_{k+1}}_{\mathcal{H}})_{(m_l+2, H_{m_l+2} \cdot (\lambda_l, \hat{\chi})H_{k+1})} \cong T^{H_{m_l+2} \cdot (\lambda_l, \hat{\chi})H_{k+1}}_{\mathcal{H}'}, \text{ and}$$

$$(T^{G/G_k}_{\mathcal{G}})_{(m_l+1, G_{m_l+1} \cdot gG_k)} \cong T^{G_{m_l+1} \cdot gG_k}_{\mathcal{G}'},$$

where $\mathcal{H}' = (H_{n+m_l+2})$ and $\mathcal{G}' = (G_{n+m_l+1})$ for $n < \omega$. Define $\theta : H/H_{k+1} \to G/G_k$ as $\theta(C) = \pi^{m_l}_G(C)$. Now we restrict θ as a function that maps $H_{m_l+2} \cdot (\lambda_l, \hat{\chi})H_{k+1}$ onto $G_{m_l+1} \cdot \hat{\chi}(\lambda_{m_l})G_k$. Then apply Lemma 3.8 and Proposition 2.2 to obtain the following:

https://doi.org/10.1017/jsl.2024.7 Published online by Cambridge University Press

$$\begin{aligned}
& \rho((T_{\mathcal{H}}^{H/H_{k+1}})_{(m_l+1, H_{m_l+1}\cdot(\bar{\lambda}_l,\bar{\chi})H_{k+1})}) \\
\geq\ & \sup\{\rho((T_{\mathcal{H}}^{H/H_{k+1}})_{(m_l+2, H_{m_l+2}\cdot(\bar{\lambda}_l,\hat{\chi})H_{k+1})}) : \forall i < m_l\ (\bar{\chi}(\lambda_i) = \hat{\chi}(\lambda_i))\} + 1 \\
\geq\ & \sup\{\rho((T_{\mathcal{G}}^{G/G_k})_{(m_l+1, G_{m_l+1}\cdot gG_k)}) : g \in G\} + 1 \\
\geq\ & \omega(\rho(T_{\mathcal{G}}^{G/G_k})) + 1 \\
=\ & \omega \cdot \alpha + 1.
\end{aligned}$$

This implies $\rho(T_{\mathcal{H}}^{H/H_{k+1}}) \geq \omega \cdot \alpha + m_l + 2$ for all $l < \omega$, and hence

$$\rho^{k+1}(\mathcal{H}) = \rho(T_{\mathcal{H}}^{H/H_{k+1}}) \geq \omega \cdot \alpha + \omega = \omega \cdot (\alpha + 1).$$

Since $\rho^{k+1}(\mathcal{H})$ is a successor ordinal, we have $\rho^{k+1}(\mathcal{H}) > \omega \cdot (\alpha + 1)$. ⊣

Now we turn to the case concerning limit ordinals. To do this, we need two lemmas.

LEMMA 4.10. *Let (G^i) be a sequence of Polish groups, and let H^i be an open subgroup of G^i for each $i < \omega$. Suppose $H = \prod_i H^i$ and*

$$G = \{(g_i) \in \prod_i G^i : \forall^\infty i\ (g_i \in H^i)\}$$

is equipped with the topology τ generated by the sets of the form $(g_i)U$ for $(g_i) \in G$ and U open in H. Then (G, τ) is a Polish group and τ is the unique group topology on G making H an open subgroup of G.

PROOF. For each $(g_i) \in G$, the subspace $(g_i)H$ of (G, τ) is homeomorphic to H, so it is Polish. Let D^i be a subset of G^i which meets every coset of H^i at exactly one point. Note that D^i is countable for all $i < \omega$. We define

$$D = \{(g_i) \in \prod_i D^i : \forall^\infty i\ (g_i = 1_{G^i})\}.$$

It is clear that $G/H = \{(g_i)H : (g_i) \in D\}$ is countable. So (G, τ) is a sum of countably many Polish spaces, thus is a Polish space.

For $(g_i), (h_i) \in G$ and U open in H with $1_H \in H$, there exists an $m < \omega$ such that $g_i, h_i \in H^i$ for $i > m$ and $U^0 \times \cdots \times U^m \times \prod_{i>m} H^i \subseteq U$, where U^i is an open subset of H^i with $1_{H^i} \in U^i$ for each $i \leq m$. We can find open neighborhoods V^i and W^i of 1_{H^i} with $(g_i V^i)(h_i W^i)^{-1} \subseteq g_i h_i^{-1} U^i$ for $i \leq m$. Now let $V = V^0 \times \cdots \times V^m \times \prod_{i \geq m} H^i$ and $W = W^0 \times \cdots \times W^m \times \prod_{i \geq m} H^i$, then V and W are open neighborhoods of 1_H, and $((g_i)V)((h_i)W)^{-1} \subseteq (g_i h_i^{-1})U$. So (G, τ) is a Polish group.

Finally, suppose τ' is another group topology on G such that H is an open subgroup of G. Then for each $(g_i) \in G$, the subspace $(g_i)H$ of (G, τ') is homeomorphic to H, so the restrictions of τ and τ' on $(g_i)H$ are the same. Hence, $\tau = \tau'$. ⊣

LEMMA 4.11. *Let (G^i) be a sequence of non-archimedean CLI Polish groups, $\mathcal{G}^i = (G^i_n) \in \mathrm{dgnb}(G^i)$ for each $i < \omega$, and let $0 < \alpha < \omega_1$. Suppose*

https://doi.org/10.1017/jsl.2024.7 Published online by Cambridge University Press

$$\sup\{\rho^1(\mathcal{G}^i) : i < \omega\} = \omega \cdot \alpha, \text{ and}$$

$$G = \{(g_i) \in \prod_i G^i : \forall^\infty i\, (g_i \in G_1^i)\}$$

is equipped with the unique group topology making $\prod_i G_1^i$ *an open subgroup of* G. *Then* G *is not* α*-CLI.*

PROOF. Put $G_0 = G$ and for $n \geq 1$, let

$$G_n = \prod_{i<n-1} G_n^i \times \prod_{i \geq n-1} G_1^i.$$

It is clear that $G_1 = \prod_i G_1^i$ and $\mathcal{G} = (G_n) \in \mathrm{dgnb}(G)$.

Given $j < \omega$, define $\theta : G/G_1 \to G^j/G_1^j$ as $\theta((g_i)G_1) = g_j G_1^j$ for $(g_i) \in G$. Applying Lemma 3.8 to the restriction of θ as in the proof of Theorem 4.9, we have

$$\begin{aligned}\rho(T_{\mathcal{G}}^{G/G_1}) &= \sup\{\rho((T_{\mathcal{G}}^{G/G_1})_{(1,G_1\cdot(g_i)G_1)}) : (g_i) \in G\} + 1\\ &\geq \sup\{\rho((T_{\mathcal{G}^j}^{G^j/G_1^j})_{(1,G_1^j\cdot gG_1^j)}) : g \in G^j\} + 1\\ &= \rho(T_{\mathcal{G}^j}^{G^j/G_1^j}) = \rho^1(\mathcal{G}^j).\end{aligned}$$

Therefore,

$$\rho^1(\mathcal{G}) = \rho(T_{\mathcal{G}}^{G/G_1}) \geq \sup\{\rho^1(\mathcal{G}^j) : j < \omega\} = \omega \cdot \alpha.$$

Since $\rho^1(\mathcal{G})$ is a non-limit ordinal and $\alpha > 0$, we have $\rho^1(\mathcal{G}) > \omega \cdot \alpha$. So G is not α-CLI. ⊣

Finally, we complete the construction in the following theorem.

THEOREM 4.12. *For any* $\alpha < \omega_1$, *there exist two non-archimedean CLI Polish groups* G_α *and* H_α *with* $\mathrm{rank}(G_\alpha) = \mathrm{rank}(H_\alpha) = \alpha$ *such that* H_α *is* α*-CLI and* G_α *is L-*α*-CLI but not* α*-CLI.*

PROOF. We construct G_α and H_α by induction on α. From Corollary 4.7 and Theorem 4.9, we only need to consider the case where α is a limit ordinal.

Let (α_i) be a sequence of ordinals less than α with $\sup\{\alpha_i : i < \omega\} = \alpha$. By the inductive hypothesis, we can find a non-archimedean CLI Polish group G^i for each $i < \omega$ such that G^i is L-α_i-CLI but not α_i-CLI. It is clear that $\mathrm{rank}(G^i) = \alpha_i < \alpha$.

Put $H_\alpha = \prod_i G^i$. Theorem 4.6(1) implies that H_α is α-CLI. By Theorem 4.1, $\mathrm{rank}(H_\alpha) \geq \mathrm{rank}(G^i)$ for each $i < \omega$. So $\mathrm{rank}(H_\alpha) = \alpha$.

For $i < \omega$, let $\mathcal{G}^i = (G_n^i) \in \mathrm{dgnb}(G^i)$. By Lemma 3.16, there exist $0 < k_i < \omega$ such that $\omega(\rho^{k_i}(\mathcal{G}^i)) = \omega \cdot \alpha_i$. Put $H_0^i = G^i$ and $H_{n+1}^i = G_{n+k_i}^i$ for $n < \omega$. Then $\mathcal{H}^i = (H_n^i) \in \mathrm{dgnb}(G^i)$ and $H_1^i = G_{k_i}^i$. By Lemma 3.7, we have

$$\omega(\rho^1(\mathcal{H}^i)) = \omega(\rho(T_{\mathcal{H}^i}^{G^i/H_1^i})) = \omega(\rho(T_{\mathcal{G}^i}^{G^i/G_{k_i}^i})) = \omega(\rho^{k_i}(\mathcal{G}^i)) = \omega \cdot \alpha_i.$$

https://doi.org/10.1017/jsl.2024.7 Published online by Cambridge University Press

So $\sup\{\rho^1(\mathcal{H}^i) : i < \omega\} = \omega \cdot \alpha$. Now we let

$$G_\alpha = \{(g_i) \in \prod_i G^i : \forall^\infty i\, (g_i \in G^i_{k_i})\}$$

be equipped with the unique group topology making $\prod_i G^i_{k_i}$ an open subgroup of G_α. By Lemma 4.11, G_α is not α-CLI. It is clear that the open subgroup $\prod_i G^i_{k_i}$ is α-CLI, so G_α is L-α-CLI, and hence $\operatorname{rank}(G_\alpha) = \alpha$. ⊣

Acknowledgements. We are grateful to Su Gao and Victor Hugo Yañez for carefully reading the manuscript, and giving much useful advice on grammar and language. We also thank the anonymous referee for helpful suggestions in both the context of definitions and language.

Funding. Research is partially supported by the National Natural Science Foundation of China (Grant No. 11725103).

REFERENCES

[1] H. Becker and A. S. Kechris, ***The Descriptive Set Theory of Polish Group Actions***, London Mathematical Society Lecture Note Series, 232, Cambridge University Press, Cambridge, 1996.

[2] S. Gao, ***Invariant Descriptive Set Theory***, Monographs and Textbooks in Pure and Applied Mathematics, 293, CRC Press, Boca Raton, 2009.

[3] S. Gao and M. Xuan, *On non-Archimedean Polish groups with two-sided invariant metrics*. ***Topology and its Applications***, vol. 161 (2014), pp. 343–353.

[4] V. L. Klee, *Invariant metrics in groups (solution of a problem of Banach)*. ***Proceedings of the American Mathematical Society***, vol. 3 (1952), pp. 484–487.

[5] M. Malicki, *On Polish groups admitting a compatible complete left-invariant metric*, this Journal, vol. 76 (2011), pp. 437–447.

[6] M. Xuan, ***On Steinhaus sets, orbit trees and universal properties of various subgroups in the permutation group of natural numbers***, Ph.D. thesis, University of North Texas, 2012.

SCHOOL OF MATHEMATICAL SCIENCES AND LPMC
NANKAI UNIVERSITY
TIANJIN 300071
P.R.CHINA
E-mail: dingly@nankai.edu.cn
E-mail: 1120210025@mail.nankai.edu.cn

https://doi.org/10.1017/jsl.2024.7 Published online by Cambridge University Press