mood-stabilising properties of lithium in patients.

Enhanced 5-HT_{1A} neurotransmission appears to dissipate within a few days of lithium withdrawal, revealing and leaving unopposed the pro-manic behavioural effects of increased 5-HT_2 activity. These findings in animals support a hypothesis of 5-HT_2 receptor over-activity in mania and hypomania, with clear therapeutic indications. Indirectly, these findings also suggest that claims for lithium's therapeutic effects should not 'be taken with a pinch of salt'.

Cookson, J. (1997) Lithium: balancing risks and benefits. *British Journal of Psychiatry*, **117**, 120–124.

Harrison-Read, P. E. (1988) Lithium differentially alters selective attention to novelty in rats depending on length of treatment and state of withdrawal: a possible role for 5-HT receptor subtypes. In *Lithium: Organic Pharmacology and Psychiatric Use* (ed. N. J. Burch), pp. 319–323. Oxford: IRL Press.

Moncrieff, J. (1997) Lithium: evidence reconsidered. *British Journal of Psychiatry*, **171**, 113–119

P. E. Harrison-Read Park Royal Centre for Mental Health, Central Middlesex Hospital, Acton Lane, London NW10 7NS

Ethnicity and clozapine metabolism

Sir: Taylor's review (1997) focused on the variability of clozapine metabolism from the effect of compounds on the cytochrome P450 system. We would like to highlight the possible influence of ethnicity on clozapine metabolism. To date, there are only three studies published in English of plasma clozapine levels in Asians. Chang *et al* (1993) found the daily mean plasma clozapine level of 77 of their Chinese patients with schizophrenia was significantly higher than those reported in American and European studies. Similarly, Chong *et al* (1997) found the mean plasma clozapine concentration attained in Chinese patients was higher than that reported in American studies, even though the Chinese patients had a lower mean daily dose of clozapine. However, it is difficult to interpret the findings from these two studies as comparisons were made with studies done elsewhere, by different workers and where many confounding variables were not controlled for. In a comparison study of 17 Korean–Americans and 17 Caucasians within the same treatment centre, Matsuda *et al* (1996) found a lower mean clozapine concentration in their Korean patients, which remained significant even after controlling for differences in daily clozapine doses. However, their study was limited by the small sample size and the failure to control for body weight. At present, there is only a suggestion of cross-ethnic differences in clozapine metabolism between Asians and Caucasians and perhaps even among Asian subgroups. More comparative studies need to be done before this can be resolved. These studies would need to control for variables like body weight, dose per kilogram, gender, and those compounds listed by Taylor (1997). Such studies should also guard against lumping the different Asian groups (Matsuda *et al*, 1996). We wish to emphasise that ethnicity is a confounding variable in psychopharmacology research and an important consideration in clinical practice.

Chang, W. H., Chein, C. P., Lin, S. K., *et al* (1993) Elevated clozapine concentrations in Chinese patients. *Neuropsychopharmacology* **9**, 117S–118S.

Chong, S. A., Tan, C. H., Khoo, Y. H., *et al* (1997) Clinical evaluation and plasma clozapine concentrations in Chinese patients with schizophrenia. *Therapeutic Drug Monitoring*, **19**, 219–223.

Matsuda, K. T., Cho, M. C., Lin, K. M., *et al* (1996) Clozapine dosage, serum levels, efficacy, and side-effect profiles: A comparison of Korean–American and Caucasian patients. *Psychopharmacology Bulletin*, **32**, 253–257.

Taylor, D. (1997) Pharmacokinetic interactions involving clozapine. *British Journal of Psychiatry*, **171**, 109–112.

S.-A. Chong Woodbridge Hospital and Institute of Mental Health, 10 Buangkok Green, Singapore 539747

G. Remington Medication Assessment Program, Schizophrenia Division, Clarke Institute of Psychiatry, 250 College Street, Toronto, Ontario M5T 1R8

One hundred years ago

County Armagh Asylum

At a meeting of the governors held on Jan. 10th the new superintendent, Mr. Lawless, in his report said that the time had now come when it became necessary to consider the appointment of an assistant medical officer. He drew attention also to the fact that there was no supply of drinking water to the new hospital, the existing supply coming from the river—in his opinion a dangerous source. He suggested that they should apply to the town commissioners for a supply. The board decided that the matter of the appointment of an assistant medical officer should be postponed for the present.

Lancet, 15 January 1898, 187.

Researched by Henry Rollin, Emeritus Consultant Psychiatrist, Horton Hospital, Epsom, Surrey

Erratum

Fenton, W. S., Blyler, C. R., Wyatt, R. J., *et al*, *BJP*, 171, 265–268. In Table 1 (p. 266) the prevalence of dyskinesia among 94 patients in the Fenton *et al* (1994) study should read 23% (not 28%). The authors apologise for any inconvenience caused by this error.

https://doi.org/10.1192/S0007125000149311 Published online by Cambridge University Press