Psychological Medicine

cambridge.org/psm

Interoceptive deficits, non-suicidal self-injury, and suicide risk: a multi-sample study of indirect effects

Christopher R. Hagan[1], Megan L. Rogers[2], Amy M. Brausch[3], Jennifer J. Muehlenkamp[1] and Thomas E. Joiner Jr[2]

[1]University of Wisconsin Eau Claire; [2]Florida State University and [3]Western Kentucky University

Original Article

Cite this article: Hagan CR, Rogers ML, Brausch AM, Muehlenkamp JJ, Joiner Jr TE (2019). Interoceptive deficits, non-suicidal self-injury, and suicide risk: a multi-sample study of indirect effects. *Psychological Medicine* **49**, 2789–2800. https://doi.org/10.1017/S0033291718003872

Received: 21 April 2018
Revised: 20 November 2018
Accepted: 27 November 2018
First published online: 3 January 2019

Key words:
Interoceptive deficits; non-suicidal self-injury; pain tolerance; suicide

Author for correspondence:
Christopher Hagan, E-mail: christopher.ryan.hagan@gmail.com

Abstract

Background. Interoceptive deficits (ID) have been associated with non-suicidal self-injury (NSSI) and suicidal behavior in multiple studies. Many of these studies are limited in scope, and have not fully examined possible mechanisms explaining how ID affect suicidal behavior.

Methods. This study assesses how self-reported ID relate to suicide ideation and attempts in six distinct and geographically diverse samples of adults ($n = 2706$) and one sample of adolescents ($n = 436$). Participants responded to a variety of self-report questionnaires and interviews.

Results. Contrary to our hypothesis, self-reported ID were only associated with suicidal ideation in two samples, one of which was the adolescent sample. Largely consistent with our predictions, self-reported ID exhibited an indirect effect on suicide attempts through versatility of NSSI in four of the five adult samples tested. Finally, the indirect effects of self-reported ID on suicide attempts through NSSI versatility did not act indirectly through behaviorally assessed pain tolerance.

Conclusions. We found that, in adults, self-reported ID are not associated with suicidal ideation, but are connected with a history of suicide attempts, through an indirect effect via NSSI. Our findings also indicate that the mechanism of action leading from self-reported ID to suicidal behavior may differ between adolescents and adults, and relate to suicidal behavior independent of pain tolerance. Clinical implications and future directions are discussed.

Over 800 000 people die by suicide annually (World Health Organization, 2014) and suicide is now the second leading cause of death in adolescents and young adults in America (Hoyert and Xu, 2012). Although suicide risk assessment and prevention have been researched extensively, recent meta-analyses (May and Klonsky, 2016; Franklin *et al.*, 2017) have demonstrated that decades of research have only minimally improved the ability to predict and prevent suicide in specific individuals. A key difficulty is that research has primarily focused on risk factors that predict suicidal ideation, which are generally poor predictors of suicidal behavior (May and Klonsky, 2016). Recent theoretical models have attempted to address this concern by focusing on different factors that lead to suicidal ideation and that contribute to the capacity to make a lethal or potentially lethal suicide attempt.

Drawing from several theoretical models, such as the Interpersonal Theory of Suicide (ITS; Van Orden *et al.*, 2010), the Three Step Theory (3ST; Klonsky and May, 2015), and the Integrated Motivational-Volitional Model (IMV; O'Connor *et al.*, 2016), capacity for suicidal behavior can be viewed as a combination of dispositional (e.g. genetically derived pain tolerance or reactivity) and acquired (e.g. learned fearlessness of pain and death and familiarity with the use of lethal means) components. One way in which capacity for suicidal behavior can be strengthened is through increased pain tolerance, which can occur through exposure to painful events including accidental injuries and non-suicidal self-injury (NSSI; Hooley *et al.*, 2010; Franklin *et al.*, 2011). A variety of factors contribute to these components of capacity and are in need of additional research regarding how they specifically lead to risk for suicidal behavior.

Deficits in interoception, defined as the ability to effectively and accurately perceive the physiological condition of one's body (Craig, 2002), may be a factor that contributes to capacity for suicidal behavior. Interoception is posited to encompass numerous domains, including awareness and trusting of bodily states, emotional awareness and reactions, attentional and self-regulatory responses, and listening to and noticing one's body sensations (Mehling *et al.*, 2012). The mental health benefits of interoceptive awareness are clear: converging evidence suggests that interoceptive awareness is central to emotional experience and shapes affective functioning (Blascovich and Mendes, 2010; MacCormack and Lindquist, 2017), is linked to greater mindfulness (Carruthers, 2008; Hölzel *et al.*, 2011; Mehling *et al.*, 2012), and promotes

© Cambridge University Press 2019

CAMBRIDGE UNIVERSITY PRESS

stronger emotion regulation (Füstös *et al.*, 2013) and empathy (Fukushima *et al.*, 2011; Ernst *et al.*, 2013).

However, interoceptive deficits (ID) have been identified in various forms of psychopathology, including those with disordered eating (Sim and Zeman, 2004; Peat and Muehlenkamp, 2011) and those who engage NSSI (Favaro and Santonastaso, 1998; Smith *et al.*, 2018). Those with both disordered eating and who engage in NSSI exhibit the greatest levels of ID (Ross *et al.*, 2009; Claes *et al.*, 2012; Muehlenkamp *et al.*, 2012; Noma *et al.*, 2015). Furthermore, ID significantly predicted suicide attempts, both cross-sectionally and longitudinally, in women with bulimia nervosa (Franko *et al.*, 2004; Forcano *et al.*, 2009). A key limitation to much of the existing literature on this topic is the frequent use of samples of women with eating disorders, limiting the potential generalizability of these results (Franko *et al.*, 2004; Forcano *et al.*, 2009; Dodd *et al.*, 2018; Smith *et al.*, 2018). Importantly, ID have been repeatedly linked to both suicidal ideation (Rogers *et al.*, 2018) and suicide attempts (Forrest *et al.*, 2015; Dodd *et al.*, 2018; Rogers *et al.*, 2018). The lack of interoceptive awareness related to these ID could contribute to a reduction in the barriers to NSSI, such as an awareness of one's bodily sensations (Rogers *et al.*, 2018). Properly formed, this awareness may curtail or prevent self-inflicted injuries through an increased initial pain threshold (Pollatos *et al.*, 2012; Weiss *et al.*, 2014). Indirectly, this contribution to NSSI engagement may contribute to changes in pain tolerance through engagement in NSSI as supported by Dodd *et al.* (2018).

NSSI has been repeatedly found to be a robust predictor of suicidal behavior in a variety of samples (Muehlenkamp and Gutierrez, 2007; Guan *et al.*, 2012; Klonsky *et al.*, 2013). Specifically, NSSI versatility (i.e. the number of different methods an individual has utilized to engage in NSSI) has been found to be a stronger longitudinal predictor of risk than frequency of NSSI (Turner *et al.*, 2013). Additionally, a meta-analysis revealed that after suicidal ideation, NSSI versatility had the strongest association with suicide attempts out of 20 relevant variables (Victor and Klonsky, 2014). Bringing this all together, one recent study found that ID were related to suicide attempts indirectly through NSSI and then through self-reported pain tolerance (Dodd *et al.*, 2018); although this study consisted exclusively of women with eating disorders.

In the current study, we tested three hypotheses across seven independent samples. First, we predicted that, consistent with past research, self-reported ID would be directly and independently associated with suicidal ideation beyond the effects of gender, age, NSSI versatility, and depression. Second, we predicted that self-reported ID would play an indirect role in suicidal actions, through direct effects on NSSI versatility. Finally, we sought to replicate and extend the recent findings of Dodd *et al.* (2018), hypothesizing that the full indirect pathway from ID to suicide attempts would progress through NSSI versatility and then pain tolerance. This specific pathway was predicted due to the likelihood that pain tolerance may increase due to repeated exposure to painful events (i.e. NSSI) and support for this model from relevant theories of suicide (Van Orden *et al.*, 2010; Klonsky and May, 2015; O'Connor *et al.*, 2016) and in previous research (Dodd *et al.*, 2018). Notably, the current study expanded on previous findings by assessing pain tolerance behaviorally rather than relying on self-report. Furthermore, this study adds to the current literature by testing these hypotheses in multiple independent samples, not limited to women with eating disorders.

Method

Participants and procedures

This study analyzed data from seven independent samples to conceptually replicate findings across samples. These samples were selected for secondary data analysis based on their inclusion of relevant measures; however, it is worth noting that none of these studies were originally designed to specifically test the hypotheses from this study. As described in the Measures section and summarized in Table 1, some variables were assessed using different measures and some variables (i.e. pain tolerance and suicide attempts) were not available in every sample. Each sample is briefly described below with demographic information summarized in Table 2. All studies were approved by the appropriate Institutional Review Boards prior to data collection.

Sample 1

Sample 1 includes 419 undergraduate students from a large Midwestern public university. Participants were recruited from all psychology department courses and invited to complete self-report questionnaires in a research laboratory in small groups as part of a larger study. Approximately 7.2% of data were missing at random in this sample, this reduced the *n* used to test the first hypothesis to 313 for this sample. Participants were compensated with course credit and local mental health resources were provided to all participants. Of these students, 7.2% reported suicidal ideation and 19.8% reported engaging in NSSI. Suicide attempt data were not available in this dataset, so this sample is not included in analyses testing our second and third hypotheses.

Sample 2

Sample 2 includes 1371 undergraduate students from a large Southwestern public university. Of note, this sample was primarily made up of Asian-American students (50.6%). Approximately 1.6% of data were missing in this sample. Participants answered questionnaires online as part of a larger study and were compensated with course credit. Of these students, 17.7% reported suicidal ideation and 24.9% reported engaging in NSSI. Among participants who reported a previous suicide attempt ($n = 46$), the mean number of attempts was 1.57 (S.D. = 0.89). Local and national mental health resource information was provided to all participants.

Sample 3

Sample 3 includes 193 undergraduate students from a large public university in the Southeastern USA. Participants completed questionnaires and a pressure pain assessment individually in a university laboratory for course credit. There were no missing data in this sample. Of these students, 17.1% reported suicidal ideation and 23.3% reported engaging in NSSI. Among participants who reported a previous suicide attempt ($n = 18$), the mean number of attempts was 1.33 (S.D. = 0.69). Local and national mental health resource information was provided to all participants.

Sample 4

Sample 4 includes 102 community members (exclusive of undergraduate students) from a mid-sized city in the Southeastern USA. Research procedures for this sample were identical to those of Sample 3, except that participants were compensated with a small gift card and a raffle entry for a tablet. There were no missing data in this sample. Of these community members, 15.7% reported suicidal ideation and 26.5% reported engaging

https://doi.org/10.1017/S0033291718003872 Published online by Cambridge University Press

https://doi.org/10.1017/S0033291718003872 Published online by Cambridge University Press

Table 1. Descriptive data

	Sample 1 ($n = 419$)		Sample 2 ($n = 1371$)		Sample 3 ($n = 193$)		Sample 4 ($n = 102$)		Sample 5 ($n = 204$)		Sample 6 ($n = 417$)		Sample 7 ($n = 436$)	
	M	S.D.	M	S.D.	M	S.D.	M	S.D.	M	S.D.	M	S.D.	M	S.D.
Self-reported ID	2.14	0.66	2.60	0.84	2.75	0.98	2.63	0.89	2.22	0.92	2.53	0.97	2.20	0.96
Measure	EDI		EDI		EDI		EDI		EDI		EDI		EDI	
Range	1–5.00		1–6.00		1–5.75		1–4.75		1–5.00		1–5.56		1–5.50	
Suicide attempts	NA	NA	0.05	0.33	0.12	0.44	0.26	0.83	0.30	1.22	0.13	0.59	0.06	0.36
Attempts if >0			1.57	0.89	1.33	0.69	1.93	1.38	2.70	2.65	2.07	1.14	1.79	0.98
Measure			Single item		SITBI		SITBI		SITBI		SHBQ		SHBQ	
Range			0–4		0–3		0–5		0–10		0–4		0–4	
Suicidal ideation	0.07	0.26	0.18	0.38	0.76	2.42	1.12	3.97	2.14	5.28	0.32	0.47	4.60	9.66
Ideation if >0	1.00	0.00	–	–	4.45	4.27	7.13	7.78	10.90	6.85	–	–	8.36	11.76
Measure	BDI-II item 9		Single item		BSS		BSS		BSS		SHBQ		SIQ-JR	
Range	0–1		0 or 1		0–20		0–26		0–30		0 or 1		0–78	
NSSI versatility	0.36	0.85	0.48	1.06	0.56	1.17	0.73	1.44	0.53	1.28	1.18	1.97	0.66	1.32
NSSI if >0	1.61	1.12	1.91	1.33	2.40	1.20	2.74	1.51	2.84	1.48	3.09	2.05	2.36	1.49
Measure	DSHI		DSHI		SITBI		SITBI		SITBI		ISAS		ISAS	
Range	0–7		0–8		0–6		0–6		0–7		0–10		0–7	
Depression	6.98	7.11	14.90	10.02	3.18	3.39	4.47	4.34	4.12	5.62	17.11	12.30	52.15	15.84
Measure	BDI-II w/o item 9		CES-D		DASS		DASS		DASS		CES-D		RADS-2	
Range	0–44		0–52		0–20		0–18		0–21		0–55		30–106	
Pain tolerance	NA	NA	NA	NA	121.00	125.13	161.30	152.69	NA	NA	NA	NA	NA	NA
Range					13.49–480		10.23–480							

For full information on each measure used, including the full title of the measure, please see the Method section.

M, mean; S.D., standard deviation; ID, interoceptive deficits; NA, measure not available in this study; Suicide attempts, number of previous suicide attempts; Attempts/Ideation/NSSI if >0 includes the mean and standard deviation when limiting the sample only to those with a score >0; 'Single item' indicates that the construct was assessed using a single item that is not part of a larger scale (full text of these items is available in the Method section); NSSI versatility, number of self-injury methods endorsed; Range, actual reported range of scores.

https://doi.org/10.1017/S0033291718003872 Published online by Cambridge University Press

Table 2. Demographics

	Sample 1 ($n = 419$)		Sample 2 ($n = 1371$)		Sample 3 ($n = 193$)		Sample 4 ($n = 102$)		Sample 5 ($n = 204$)		Sample 6 ($n = 417$)		Sample 7 ($n = 436$)	
	n	% of group	*n*	% of group	*n*	% of group	*n*	% of group	*n*	% of group	*n*	% of group	*n*	% of group
Gender														
Male	69	16.5	351	25.6	45	23.3	28	27.5	89	43.6	112	26.9	201	46.3
Female	326	77.8	1018	74.3	148	76.7	74	72.5	115	56.4	302	72.4	229	52.8
Other	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	3	0.6	4	0.97
Race/ethnicity														
White	362	86.4	235	17.1	122	63.2	78	76.5	169	82.8	326	78.2	368	85.4
African-Am	4	1.0	25	1.8	29	15.0	12	11.8	12	5.9	36	8.6	10	2.3
Hispanic/Latino	1	0.2	194	14.2	28	14.5	5	4.9	6	2.9	17	4.1	12	2.8
Asian	9	2.1	694	50.6	8	4.1	2	2.0	13	6.4	15	3.6	8	1.9
Middle Eastern	NA	NA	173	12.6	4	2.1	2	2.0	0	0.0	NA	NA	NA	NA
AK Native/AI	14	3.3	6	0.4	1	0.5	1	1.0	3	1.5	2	0.5	4	0.9
HI Native/Pac Is	NA	NA	NA	NA	1	0.5	2	2.0	0	0.0	0	0	NA	NA
Multiracial	3	0.7	35	2.6	NA	NA	NA	NA	NA	NA	13	3.1	20	4.6
Other	2	0.5	5	0.4	0	0.0	0	0.0	0	0.0	6	1.4	9	2.1
	M	S.D.	M	S.D.	M	S.D.	M	S.D.	M	S.D.	M	S.D.	M	S.D.
Age	20.5	3.76	20.1	2.10	19.2	2.39	38.0	16.4	34.8	10.53	19.8	3.89	13.2	1.09

African-Am, African-American/Black; Ak Native/AI, Alaska Native/American Indian; HI Native/Pac Is, Hawaiian Native/Pacific Islander; M, mean; S.D., standard deviation; NA, response option not available in this study. In Sample 1, 5.7% of participants did not report gender or race. In Sample 2, 0.1% did not report gender, and 0.3% did not report race. In Sample 5, 0.5% did not report race.

in NSSI. Among participants who reported a previous suicide attempt ($n = 14$), the mean number of attempts was 1.93 (S.D. = 1.38).

Sample 5

Sample 5 includes 204 American participants from 40 of the 50 states recruited through Amazon's Mechanical Turk (MTurk) service. Many recent studies have utilized this recruitment system, and it has provided high-quality data (Buhrmester *et al.*, 2011; Mason and Suri, 2012). Participants completed this study online and were provided with a small financial incentive. Approximately 0.4% of data were missing in this sample. Of these adults, 19.6% reported suicidal ideation and 18.2% reported engaging in NSSI. Among participants who reported a previous suicide attempt ($n = 23$), the mean number of attempts was 2.70 (S.D. = 2.65). National mental health resource information was provided to all participants.

Sample 6

Sample 6 includes 417 university students enrolled in undergraduate psychology courses at a Southern public university. Participants were recruited from all psychology department courses and invited to come into the research laboratory to complete a packet of self-report questionnaires in small groups as part of a larger study. Participants received course credit for participation. Mental health services referral information and resources were provided to participants. Approximately 5.0% of data were missing in this sample. Of these students, 31.5% reported suicidal ideation and 38.0% reported engaging in NSSI. Among individuals who reported a suicide attempt ($n = 28$), the mean number of attempts was 2.07 (S.D. = 1.14)

Sample 7

Sample 7 includes 436 adolescents from public middle and high schools in the south-central region of the USA. Participants were recruited from seventh and ninth grades and completed self-report questionnaires at school as part of a larger study after parental consent was obtained. Adolescents received $10 for their participation. Participants who endorsed critical items on measures of suicide ideation and attempts were referred to school counselors for further assessment. Approximately 5.0% of data were missing in this sample. Of these adolescents, 55.0% reported suicidal ideation and 27.8% reported engaging in NSSI. Among adolescents who reported a suicide attempt ($n = 14$), the mean number of attempts was 1.79 (S.D. = 0.98).

Measures

Demographics

Data on gender (males coded as 1, females coded as 0), age, and racial/ethnic identity were collected through demographic questionnaires.

Interoceptive deficits

Self-reported ID were assessed through the nine-item Interoceptive Awareness subscale of the Eating Disorders Inventory-3 (EDI; Garner, 2004). The research scoring procedures, in which participants provide responses on a six-point scale, were utilized in all samples to allow for a wider variety of responses compared with the truncated clinical scoring procedures. Samples 2–5 included an older version of the EDI (Garner *et al.*, 1983). This version includes identical items to the EDI-3, with the exception of one missing item. Due to the discrepancy in item numbers, mean Interoceptive Awareness subscale scores were calculated (instead of a summed score) in each sample and used in analyses (possible range: 1–6). Cronbach's α values were good to excellent in all samples ($\alpha = 0.84$–0.92).

Suicide attempts

Sample 1 did not assess suicide attempt history. Sample 2 utilized a single item question with the same wording as the item used in samples 6 and 7 from the Self-Harm Behaviors Questionnaire (SHBQ) to assess suicide attempt history. Lifetime number of suicide attempts were assessed using the Self-Injurious Thoughts & Behaviors Interview-Short Self Report Form (SITBI; Nock *et al.*, 2007) in Samples 3–5. The SITBI asks, 'How many suicide attempts have you made in your lifetime?' Samples 6–7 used the suicide attempt frequency item from the SHBQ (Gutierrez *et al.*, 2001), which asks, 'How many times have you attempted suicide?' and has response options of '0', 'once', 'twice', 'three times', and 'four or more times'. The upper response was coded as four attempts in this study.

Suicidal ideation

Sample 1 utilized item 9 of the Beck Depression Inventory Second Edition (BDI-II; described below; Beck *et al.*, 1996) to assess suicidal ideation (possible range: 0–3). Sample 2 assessed suicidal ideation through a single item with a yes or no response option, coded 0 (no) or 1 (yes), 'Have you ever seriously considered suicide?' Samples 3–5 assessed suicidal ideation with the Beck Scale for Suicidal Ideation (BSS; Beck *et al.*, 1988). Nineteen items from this 21-item measure were included. Two items assessing suicidal behavior were excluded. Cronbach's α values were good to excellent in all three samples: $\alpha_{\text{Sample 3}} = 0.88$; $\alpha_{\text{Sample 4}} = 0.88$; $\alpha_{\text{Sample 5}} = 0.96$ and the possible range of scores is 0–38. Sample 6 utilized the suicidal ideation item from the SHBQ (Gutierrez *et al.*, 2001) which asks 'Have you ever talked or thought about wanting to die? Killing yourself?' A yes response to either question is coded as positive for suicidal ideation (i.e. '1'). Sample 7 utilized the Suicide Ideation Questionnaire-JR (SIQ-JR; Reynolds, 1988) as a continuous measure of ideation. The SIQ-JR consists of 15 items that ask about thoughts related to suicide, death, dying, and making plans and preparations, rated on a 0–6 Likert scale ranging from 'I never have this thought' to 'I have this thought every day'. The internal consistency of the SIQ-JR was high: $\alpha = 0.93$ and the possible range of scores is 0–90.

Non-suicidal self-injury

NSSI was assessed using a measure of NSSI versatility, a count of the number of different methods an individual has used throughout their lifetime to engage in NSSI. This metric has been strongly linked to suicidal behavior in a number of studies (e.g. Muehlenkamp and Gutierrez, 2007; Turner *et al.*, 2013). Samples 1 and 2 obtained these data from the Deliberate Self-Harm Inventory (DSHI; Gratz, 2001), which assesses NSSI history (possible range: 0–15). Samples 3–5 utilized the SITBI (Nock *et al.*, 2007) to determine this information (possible range: 0–7). Samples 6–7 utilized responses from the Inventory of Statements About Self-Injury (ISAS; Klonsky and Glenn, 2009; possible range: 0–13). All NSSI methods endorsed were summed to create a total number of NSSI methods. Most of the methods described in these three measures are present in at least two of the three measures. All three measures include the most common forms of NSSI (e.g. cutting, burning, hitting). Some of the SITBI

https://doi.org/10.1017/S0033291718003872 Published online by Cambridge University Press

methods are combined (e.g. cutting and carving one's skin is a single option). The DSHI also includes a few rarely or never endorsed methods such as dripping acid onto one's skin. Each of these widely used assessment measures assess the same phenomenon of NSSI versatility, but break it into more or fewer specific categories.

Depression

In Sample 1, depression was measured using the BDI-II (Beck *et al.*, 1996) total score, excluding item 9, as item 9 was utilized as a dependent variable assessing suicidal ideation. The BDI-II is a well-validated, 21-item measure of depression symptom severity with a possible range of scores in this study of 0–60, and had a strong Cronbach's α value in this study ($\alpha = 0.90$). In Sample 1, 77.1% scored in the minimal range (0–13), 7.4% in the mild range (14–19), 4.5% in the moderate range (20–28), and 1.7% in the severe (29–63) range.

Samples 2 and 6 used the Center for Epidemiological Studies Depression scale (CES-D; Radloff, 1977). The CES-D is a widely used, well-validated 20-item measure of depressive symptom severity, which demonstrated a strong Cronbach α value in these samples ($\alpha = 0.90$, 0.93, respectively). It has a possible score range of 0–60 with scores at or above 16 indicating those at risk for clinical depression (Lewinsohn *et al.*, 1997). In Sample 2, 37.9% of participants scored in the at-risk range, and in Sample 6, 40.0% were at risk.

Samples 3–5 used the depression subscale of the Depression Anxiety Stress Scale (DASS; Henry and Crawford, 2005), a 21-item measure of depression, anxiety, and stress. This subscale has a possible range of scores from 0 to 21, and demonstrated good-to-excellent Cronbach's α values in all three samples ($\alpha_{\text{Sample 3}} = 0.85$; $\alpha_{\text{Sample 4}} = 0.89$; $\alpha_{\text{Sample 5}} = 0.96$). Severity ranges are based on the doubled score for the 21-item version of the DASS used in these samples. In Sample 3, 75.1% scored in the 'normal' range (0–9), 9.3% in the mild range (10–13), 9.3% in the moderate range (14–20), 5.7% in the severe range (21–27), and 0.5% in the extremely severe range (28–42). In Sample 4, 61.8% scored in the 'normal' range, 12.7% in the mild range, 14.7% as moderate, 4.9% severe, and 5.9% as extremely severe. In Sample 5, 69.1% scored in the 'normal' range, 6.4% as mild, 10.3% moderate, 4.4% severe, and 9.8% as extremely severe.

Sample 7 assessed depression with the Reynolds Adolescent Depression Scale – Second Edition (RADS-2; Reynolds, 2002). The RADS-2 is a widely used 30-item measure of depressive symptoms that has been validated for use with adolescents, with excellent reliability in the current sample ($\alpha = 0.94$), and a possible range of 30–120, with scores at or above 76 representing those at risk for clinical depression. In Sample 6, 9.4% of adolescents scored above the cut-off.

Pain tolerance

Pain tolerance was assessed with a pressure algometer that applies pressure to the back of a participant's middle finger. This algometer's 755 g mass at a blunted tip was applied to the non-dominant middle finger between the central knuckles. Given the design of this tool (i.e. a uniform amount of pressure is constantly applied), tolerance was measured as the amount of time that passed until the participant reached their maximum pain tolerance and lifted the weight off of their finger, with a maximum time of 8 min. The algometer used in these studies was built to match the specifications of algometers that have been previously used in psychological research (Hooley and St. Germain, 2014). These data were only available from Samples 3 and 4.

Data analytic strategy

Linear multiple regression analyses were conducted to test the first hypothesis assessing the relationship between self-reported ID and suicidal ideation using SPSS version 24. Analyses for Samples 2 and 6 utilized logistic regression analyses as the suicidal ideation variable was a dichotomous question. In all models, suicidal ideation was entered as the dependent variable with self-reported ID, NSSI versatility, depression scores, gender, and age included as independent variables. NSSI versatility, gender, age, and depression scores, which were assessed in all samples, were entered as covariates to control for possible differences in suicidal ideation based on gender and age (as these variables differed notably between some samples) and for known differences in suicidal ideation among those with and without depressive symptoms and a history of NSSI engagement (May and Klonsky, 2016). Carlson and Wu (2012) recommend including only covariates that have significant associations with predictor variables. NSSI versatility and depression scores were correlated with self-reported ID in all samples, and gender and age were each correlated with self-reported ID in four out of the seven samples. Given our primary goal of evaluating the role of self-reported ID in suicidal ideation and providing clearly comparable analyses across these seven samples, all four covariates were included in each test of this hypothesis. Missing data in each sample were handled with listwise deletion. The model effect size for each sample is reported as well.

To test the second hypothesis assessing the possible indirect role of self-reported ID in suicidal actions through NSSI, NSSI versatility was entered as a mediator between self-reported ID and suicide attempts using zero inflated negative binomial (ZINB) analyses in MPlus Version 6.11. Finally, to test the third hypothesis that the indirect effect of self-reported ID operates through NSSI versatility and then through pain tolerance on suicide attempts, we conducted a ZINB indirect effects model in MPlus version 6.11. We utilized maximum likelihood estimation with robust standard errors when testing both the second and third hypotheses.

Results

Data summarizing means and standard deviations of data from all samples are included in Table 1. Correlations between these variables are presented for each sample in online Supplementary Tables S1–S4[†1].

The first hypothesis predicted that self-reported ID would be directly associated with suicidal ideation beyond the effects of gender, age, NSSI versatility, and depression. Sample 1, composed of Midwestern public university students supported this hypothesis, finding that only self-reported ID and depression scores significantly accounted for variation in suicidal ideation

†The notes appear after the main text.

[1]As relatively little is known about the simple relationship between self-reported ID and suicidal ideation and attempts outside of eating disorder samples, we also included the results of simple regression analyses using self-reported ID scores as the independent variable and suicidal ideation and attempts as the outcome variables in online Supplementary Table S5.

https://doi.org/10.1017/S0033291718003872 Published online by Cambridge University Press

[$F_{(5,308)} = 22.99$, $p < 0.001$; $R^2 = 0.27$[2]; see Table 3]. Sample 7 was the only other sample that supported this hypothesis [$F_{(5,342)} = 76.86$, $p < 0.001$; $R^2 = 0.52$; see Table 3]. In Sample 7, an adolescent sample, depression, and NSSI versatility were significant indicators of ideation, and self-reported ID accounted for unique variance in ideation above and beyond the covariates.

In Sample 2 (Southwestern university students), Sample 3 (Southeastern university students), and Sample 5 (MTurk workers), only depression and NSSI versatility accounted for significant variance in suicidal ideation. Sample 2 produced a full model accounting for approximately 20% of variation in suicidal ideation [Nagelkerke Pseudo $R^2 = 0.20$; $\chi^2(5) = 164.56$, $p < 0.001$]. The full model from Sample 3 accounted for only 10% of suicidal ideation variance [$R^2 = 0.10$; $F_{(5,187)} = 4.05$, $p = 0.002$] but Sample 5 accounted for nearly 58% of suicidal ideation variance [$R^2 = 0.58$; $F_{(5,192)} = 51.97$, $p < 0.001$]. The model for Sample 4 (Southeastern community residents) accounted for 40% of the variation in suicidal ideation, but this effect was driven by depression scores and gender; neither NSSI versatility nor self-reported ID were significant indicators of suicidal ideation in this sample [$R^2 = 0.40$; $F_{(5,96)} = 12.65$, $p < 0.001$]. In Study 6, gender, depression, and NSSI versatility accounted for significant variance in ideation [$\chi^2(5) = 98.85$, $p < 0.001$; Nagelkerke Pseudo $R^2 = 0.37$; see Table 3].

The second hypothesis states that self-reported ID play an indirect role in suicidal actions (i.e. number of suicide attempts) through NSSI versatility. Table 4 includes data from each step of the indirect effects analyses for each of the six samples used to test this hypothesis (Sample 1 was excluded because it did not include an evaluation of suicide attempt history). Figure 1 illustrates the results from Sample 5 as an example of the visual representation of these analyses. Results from most samples found statistically significant indirect effects of self-reported ID on suicide attempts through NSSI, supporting our hypothesis with the exceptions of Sample 4, the Southeastern community sample, and Sample 7, the adolescent sample.

The final hypothesis states that in addition to self-reported ID exerting an effect on suicide attempts through an effect on NSSI, NSSI additionally affects suicide attempts indirectly through elevated pain tolerance. Only Samples 3 and 4 included an assessment of pain tolerance, so this hypothesis was only tested in those two samples. First, in Sample 3, self-reported ID were significantly associated with NSSI versatility ($\beta = 0.21$, $p = 0.003$, IRR = 1.23). However, self-reported ID and NSSI were each unrelated to pain tolerance ($\beta = 0.07$ and 0.10, $p = 0.43$ and 0.22, IRR = 1.07 and 1.11, respectively), and self-reported ID ($\beta = 0.15$, $p = 0.646$, IRR = 1.16), NSSI versatility ($\beta = 0.74$, $p = 0.507$, IRR = 2.10), and pain tolerance ($\beta = 0.53$, $p = 0.486$, IRR = 1.70) were unrelated to the presence of a lifetime suicide attempt after taking into account excess zeros using the ZINB analysis. The indirect effect of self-reported ID on pain tolerance through NSSI versatility ($B = 2.56$, S.E. = 2.36, $p = 0.28$, IRR = 12.94) and of NSSI versatility on suicide attempts through pain tolerance ($B = 0.03$, S.E. = 0.04, $p = 0.46$, IRR = 1.03) were non-significant. Similarly, in Sample 4, self-reported ID were positively related to NSSI versatility ($\beta = 0.31$, $p = 0.004$, IRR = 1.36), and self-reported ID ($\beta = -0.05$, $p = 0.60$, IRR = 0.95) and NSSI ($\beta = 0.05$, $p = 0.63$, IRR = 1.05) were each unrelated to pain tolerance. Furthermore, self-reported ID ($\beta = -0.11$, $p = 0.948$, IRR = 0.90), NSSI versatility ($\beta = 1.01$, $p = 0.935$, IRR = 2.75), and pain tolerance ($\beta = -0.19$, $p = 0.935$, IRR = 0.83) were not related to the presence of a lifetime suicide attempt after taking into account excess zeros using the ZINB analysis. As in Sample 3, the indirect effect of self-reported ID on pain tolerance through NSSI versatility ($B = 2.58$, S.E. = 5.52, $p = 0.64$, IRR = 13.20), and of NSSI versatility on suicide attempts through pain tolerance ($B = 0.01$, S.E. = 0.02, $p = 0.69$, IRR = 1.01), were non-significant.

In summary, results from adult samples generally found that self-reported ID are not directly associated with suicidal ideation, but that they are indirectly connected to suicide attempts through NSSI versatility. Conversely, in our one adolescent sample, self-reported ID were associated with suicidal ideation beyond the effect of multiple covariates but were not indirectly associated with a history of suicide attempts. Finally, the indirect path from self-reported ID to suicide attempts does not appear to flow through NSSI and then through pain tolerance in the two adult samples we tested, contrary to previous, similar evaluations of this pathway.

Discussion

This study sought to understand the role of self-reported ID on suicidal ideation and behavior in a variety of samples, distinct from the severely eating disordered samples in which the relationship between ID and suicide ideation and behavior is often studied. We tested three hypotheses, first that self-reported ID would be directly and independently associated with suicidal ideation beyond the effects of gender, age, NSSI versatility, and depression. Second, we predicted that self-reported ID would play an indirect role in suicidal actions, through direct effects on NSSI versatility. Finally, we tested an alternative indirect pathway from self-reported ID to suicide attempts through NSSI versatility and then through behaviorally assessed pain tolerance.

Overall, Hypothesis 1, which predicted a direct association between self-reported ID and suicidal ideation, received mixed results. Only data from Samples 1 and 7 supported this hypothesis when including gender, age, NSSI versatility, and depression as covariates. Additionally, the strength of the effect of self-reported ID on suicidal ideation was the weakest of all statistically significant variables in both samples. Sample 1 utilized a single-item measure of suicidal ideation that had limited variability (i.e. item 9 from the BDI-II), which may have accounted for the divergent results from the other adults samples. However, this single item has previously been used to assess suicidal ideation (Anestis *et al.*, 2011; Venta *et al.*, 2014), and Sample 2 also used a single-item measure of ideation but did not provide support for this hypothesis. Sample 7 was a sample of adolescents, so the divergent findings of this sample may indicate that self-reported ID have a different relationship with suicidal ideation in adolescents compared with adults. Based on the five study samples that did not support this hypothesis, which include over 2200 unique individuals, self-reported ID do not appear to be directly associated with suicidal ideation beyond the effects of gender, age, depressive symptoms, and NSSI versatility for adults. The significant finding from Sample 7 may indicate that there is a direct association in adolescents, but replication and additional research are needed to confirm this finding, and to compare possible age-related differences in the relationship between self-reported ID and suicidal ideation between adolescents and adults.

Four of five tested adult samples (three geographically diverse undergraduate samples and one diverse community sample)

[2]All R^2 effect sizes described here are for the full model.

https://doi.org/10.1017/S0033291718003872 Published online by Cambridge University Press

https://doi.org/10.1017/S0033291718003872 Published online by Cambridge University Press

Table 3. Regression analyses predicting suicidal ideation

	Sample 1		Sample 2			Sample 3		Sample 4		Sample 5		Sample 6			Sample 7	
Predictor	β	S.E.	B	S.E.	OR	β	S.E.	β	S.E.	β	S.E.	*B*	S.E.	OR	*B*	S.E.
Self-reported ID	0.18**	0.021	1.15	0.113	1.15	−0.05	0.210	−0.15	0.419	−0.10	0.339	0.35	0.22	1.42	0.21***	0.50
NSSI	−0.03	0.015	1.81***	0.070	1.81	0.19**	0.149	0.02	0.235	0.25***	0.210	0.32***	0.09	1.37	0.22***	0.32
Depression	0.43***	0.002	1.05***	0.009	1.05	0.26**	0.058	0.71***	0.085	0.68***	0.053	0.06***	0.02	1.06	0.43***	0.03
Gender	0.08	0.034	1.05	0.187	1.05	0.02	0.410	−0.27**	0.763	0.04	0.489	0.74*	0.35	2.09	−0.01	0.76
Age	−0.08	0.003	1.03	0.038	1.03	−0.07	0.071	0.09	0.021	−0.04	0.023	0.03	0.03	1.03	0.01	0.36

*$p < 0.05$; **$p < 0.01$; ***$p \leqslant 0.001$.
ID, interoceptive deficits; NSSI, non-suicidal self-injury versatility; Gender, male coded as 1, female as 0; β, standardized β; S.E., standard error; OR, odds ratio.

Table 4. Indirect effects of self-reported interoceptive deficits on suicide attempts through NSSI versatility

	a path ID→NSSI				*b* path NSSI→SA				*c* path ID→SA total effect				*c*′ path ID→SA direct effect				Indirect effect			
	B	SE	*p*	IRR	*B*	SE	*p*	IRR	*B*	SE	*p*	IRR	*B*	SE	*p*	IRR	*B*	SE	*p*	IRR
Sample 2	0.24	0.04	<0.001	1.27	0.99	0.12	<0.001	2.70	0.49	0.18	0.008	1.63	−0.04	0.21	0.85	0.96	0.24	0.05	<0.001	1.27
Sample 3	0.25	0.09	0.005	1.28	0.45	0.13	<0.001	1.57	0.35	0.24	0.14	1.42	0.29	0.27	0.29	1.33	0.11	0.05	0.017	1.12
Sample 4	0.44	0.18	0.012	1.55	0.42	0.33	0.21	1.51	0.21	0.49	0.67	1.23	−0.17	0.63	0.79	0.84	0.18	0.14	0.20	1.20
Sample 5	0.65	0.14	<0.001	1.91	0.61	0.18	0.001	1.83	0.90	0.27	0.001	2.45	0.43	0.32	0.17	1.54	0.39	0.14	0.004	1.48
Sample 6	0.85	0.11	<0.001	2.34	0.28	0.09	0.002	1.32	1.06	0.20	<0.001	2.90	0.82	0.24	0.001	2.28	0.23	0.08	0.002	1.26
Sample 7	0.59	0.08	<0.001	1.80	0.23	0.13	0.07	1.26	1.49	0.39	<0.001	4.45	1.29	0.40	0.001	3.62	0.14	0.08	0.09	1.15

NSSI, non-suicidal self-injury versatility; SA, suicide attempt; *B*, unstandardized β; IRR, incidence rate ratios.




Fig. 1. Visual representation of the indirect effects analyses. Data are reported from Sample 5.

supported the second hypothesis, which predicted an indirect effect between self-reported ID and suicide attempts through NSSI versatility. While the adolescent sample (i.e. Sample 7) did not show a significant indirect effect, it should be noted that the indirect effect of self-reported ID on suicide attempts through NSSI versatility approached significance in this adolescent sample ($p = 0.09$). A more clinically based sample of adolescents may result in a significant indirect relationship between these variables. This topic merits more attention in future research endeavors.

The hypothesis that these indirect effects would act on suicidal behavior through behaviorally assessed pain tolerance, consistent with Dodd *et al.*'s (2018) finding of an indirect effect through self-reported pain tolerance, was not supported in either the Southeastern undergraduate or the community-based sample, which was the one adult sample to not find a significant indirect effect in the second hypothesis. The use of a behavioral measure of pain tolerance (i.e. blunt pressure pain tolerance) in these two samples did not support an indirect role of pain tolerance on suicide attempts. This indicates that although self-reported ID appear to act on suicide attempts through NSSI versatility, that link does not appear to be due to increased levels of pain tolerance. Self-reported ID may be another component of the breadth of ways in which individuals are predisposed to have an elevated capacity for suicidal behavior. Phenomena including having an outsider's perspective of one's own body and having low body regard, which have been shown to moderate risk for NSSI (Muehlenkamp *et al.*, 2005, 2013), may also influence this relationship.

Our findings that self-reported ID were not associated directly with pain tolerance also diverges from previous research, which support a link between ID and pain tolerance (Pollatos *et al.*, 2012; Weiss *et al.*, 2014). These differences may be due to differences in the assessment of ID (i.e. this study used a self-report measure that primarily assesses awareness of emotional experiences, rather than an assessment of accuracy of heartbeat detection) and the use of samples of relatively healthy undergraduates and adults rather than a study of somatoform patients and healthy controls.

Overall, results from six adult samples largely (with the exception of one undergraduate sample) found that self-reported ID are not directly associated with suicidal ideation, but that they are indirectly connected to suicide attempts through NSSI versatility, with the exception of one community-based sample. These findings that self-reported ID appear to be specifically associated with suicidal behavior and not suicidal ideation in adults indicates that self-reported ID may be one of few identified indicators of risk for suicidal behavior. Future studies should seek to determine how self-reported ID best fits into theoretical conceptualizations of capacity for suicide (e.g. as an acquired and/or dispositional factor). Although self-reported ID appear to primarily increase risk through greater NSSI versatility, it does not appear that this occurs due to increases in pain tolerance. One way in which this may occur could be through an effect on diminished body regard and a feeling of disconnection from one's body which then reduces psychological barriers against harming oneself, making NSSI more likely. These possibilities all require testing in future studies.

Assessments of ID may be able to provide a non-face valid way of assessing suicide risk (within the context of a broader suicide risk assessment) to which individuals who want to conceal suicidal thoughts or behaviors may be willing to respond. Additional research is needed to explore potential mechanisms connecting ID to suicidal behavior and the potential utility of using ID assessments as a non-face valid measure of increased suicide risk.

Although these results will need replication, especially within broad clinical samples, preliminary clinical implications can be proposed. Consistent with other studies (Nock *et al.*, 2006; Muehlenkamp and Gutierrez, 2007), NSSI versatility was correlated with suicidal behavior in all our samples. Clients with a significant or ongoing NSSI history should be regularly assessed for suicide risk with a thorough suicide risk evaluation (see Chu *et al.*, 2015). Furthermore, among these clients, an assessment of their awareness of and the accuracy of their interoceptive experiences may be useful. For those with ID, training to be more aware and mindful of their bodily signals, such as is used in treatments for anxiety (Barlow *et al.*, 2016; Blakey and Abramowitz, 2018) and in Dialectical Behavior Therapy (Linehan, 1993), may be clinically useful for reducing suicide risk.

This study's primary strength is that it includes seven independent and diverse samples from different regions of the USA, providing improved generalizability of the results to the population at large. This is also one of a few studies examining the role of self-reported ID on suicide to include samples that are not exclusively limited to women with eating disorders. As these studies were each implemented independently as parts of other research endeavors, different measures were used to assess the variables of interest. The consistency of the findings across different measures shows that these findings may reflect true effects.

Although this study used a wide sample of over 3000 individuals, it is limited in that none of the included samples were from a clinical population; further research in clinical samples is needed. This study was also limited by the self-report (with the exception of behaviorally assessed pain tolerance), cross-sectional design used in all samples. This prevents our ability to determine temporal precedence and causality of the variables assessed in this study. Additionally, these studies were designed to test other hypotheses and were not originally designed to test the hypotheses included in this article. Future studies would benefit from including behavioral measures of interoceptive abilities such as a heartbeat detection test. Furthermore, changes in ID (and any related change in pain tolerance) over time, especially during the transition from adolescence to adulthood should be investigated. Additionally, although gender was included as a covariate, these analyses were not conducted independently for men and women. Future studies would benefit from explicitly testing

https://doi.org/10.1017/S0033291718003872 Published online by Cambridge University Press

these relationships to determine if they function differently in men and women.

Finally, our study was limited in that each sample utilized a self-report measure of ID that was designed for use in relation to disordered eating (i.e. the Eating Disorder Inventory) and which only assesses one facet of interoceptive abilities (i.e. awareness of emotional experiences; Khalsa *et al.*, 2018). However, as noted by previous research, interoceptive awareness is multidimensional (Mehling *et al.*, 2012) and perceived interoceptive awareness does not always correspond to actual interoceptive awareness (Garfinkel *et al.*, 2015). Facets of interoceptive awareness likely relate to suicide risk differentially, as demonstrated by Rogers *et al.* (2018), who found that suicide attempters, but not suicide ideators, had a greater tendency to ignore or distract themselves from bodily sensations, less self-regulation using body sensations, and less trust of their body sensations compared with those without a history of suicidal ideation or attempts. These facets may similarly relate more strongly to NSSI and pain tolerance than the type of interoceptive awareness assessed by the Interoceptive Awareness subscale of the EDI, which primarily contains items pertaining to interoceptive emotional awareness. Thus, future work should integrate multidimensional assessments [e.g. heartbeat perception tests and the Multidimensional Assessment of Interoceptive Awareness (Mehling *et al.*, 2012), a self-report measure that assesses a variety of interoceptive components] to further elucidate the nature of the associations between ID, NSSI, pain tolerance, and suicidal behavior across the range of interoceptive facets including attentional biases, distortions of physiological sensitivity, cognitive biases, and insight impairments. As interoception is currently understood to be a multi-faceted, complex way in which the nervous system processes internal stimuli, with numerous potential overlapping elements with how emotion and pain are processed (Khalsa *et al.*, 2018), many aspects of interoception remain to be assessed in the context of suicide risk. Specific interoceptive symptoms have been implicated in numerous psychiatric disorders including panic disorder, somatic symptoms disorders, eating disorders, post-traumatic stress disorder, and autism spectrum disorders. While some specific symptoms overlap between disorders (e.g. failure to anticipate changes to interoceptive states in mood and anxiety disorders), other situations may involve a unique constellation of interoceptive symptoms specific to the individual and disorder (see Khalsa *et al.*, 2018 for a brief review). Research remains to be conducted on the specific interoceptive symptoms that define the relationship between ID and suicide based on an expanded conceptualization of interoception that includes multifaceted assessments that go beyond self-reported ID and a focus on awareness of emotional experiences.

Conclusion

This study, incorporating seven independent samples, largely found that among adults, self-reported ID are not uniquely associated with suicidal ideation beyond other relevant factors, but are associated with suicide attempt history indirectly through NSSI versatility and not through behaviorally assessed physical pain tolerance. Furthermore, self-reported ID are associated with suicidal ideation in adolescents. Awareness of and treatments addressing ID in the context of NSSI and suicide risk may be beneficial, even outside of the eating disordered patient populations in which ID have typically been studied.

Author ORCIDs. Christopher R. Hagan 0000-0001-5603-7433

Financial support. Support from NIGMS Grant #8P20GM103436-14 was used in the collection of data for Sample 7.

Supplementary material. The supplementary material for this article can be found at https://doi.org/10.1017/S0033291718003872.

References

Anestis MD, Bagge CL, Tull MT and Joiner TE (2011) Clarifying the role of emotion dysregulation in the interpersonal-psychological theory of suicidal behavior in an undergraduate sample. *Journal of Psychiatric Research* **45**, 603–611.

Barlow DH, Allen LB and Choate ML (2016) Toward a unified treatment for emotional disorders – republished article. *Behavior Therapy* **47**, 838–853.

Beck AT, Steer RA and Ranieri WF (1988) Scale for suicide ideation: psychometric properties of a self-report version. *Journal of Clinical Psychology* **44**, 499–505.

Beck AT, Steer RA and Brown GK (1996) *Manual for the BDI-II*. San Antonio, TX: Psychological Corporation.

Blakey SM and Abramowitz JS (2018) Interoceptive exposure: an overlooked modality in the cognitive-behavioral treatment of OCD. *Cognitive and Behavioral Practice* **25**, 145–155.

Blascovich J and Mendes WB (2010) Social psychophysiology and embodiment. In Fiske ST, Gilbert DT and Lindzey G (eds), *Handbook of Social Psychology*, 5th Edn. Hoboken, NJ: John Wiley & Sons, Inc, pp. 194–227.

Buhrmester M, Kwang T and Gosling SD (2011) Amazon's Mechanical Turk: a new source of inexpensive, yet high-quality data? *Perspectives on Psychological Science* **6**, 3–5.

Carlson KD and Wu J (2012) The illusion of statistical control: control variable practice in management research. *Organizational Research Methods* **15**, 413–435.

Carruthers G (2008) Types of body representation and the sense of embodiment. *Consciousness and Cognition* **17**, 1302–1316.

Chu C, Klein KM, Buchman-Schmitt J, Hom MA, Hagan CR and Joiner TE (2015) Routinized assessment of suicide risk in clinical practice: an empirically-informed update. *Journal of Clinical Psychology* **71**, 1186–1200.

Claes L, Jiménez-Murcia S, Agüera Z, Castro R, Sánchez I, Menchón JM and Fernández-Aranda F (2012) Male eating disorder patients with and without non-suicidal self-injury: a comparison of psychopathological and personality features. *European Eating Disorders Review* **20**, 335–338.

Craig AD (2002) How do you feel? Interoception: the sense of the physiological condition of the body. *Nature Reviews Neuroscience* **3**, 655–666.

Dodd DR, Smith AR, Forrest LN, Witte TK, Bodell L, Bartlett M, Siegfried N and Goodwin N (2018) Interoceptive deficits, nonsuicidal self-injury, and suicide attempts among women with eating disorders. *Suicide & Life Threatening Behavior* **48**, 438–448.

Ernst J, Northoff G, Böker H, Seifritz E and Grimm S (2013) Interoceptive awareness enhances neural activity during empathy. *Human Brain Mapping* **34**, 1615–1624.

Favaro A and Santonastaso P (1998) Impulsive and compulsive self-injurious behavior in bulimia nervosa: prevalence and psychological correlates. *The Journal of Nervous and Mental Disease* **186**, 157–165.

Forcano L, Fernández-Aranda F, Álvarez-Moya E, Bulik C, Granero R, Gratacòs M, Jiménez-Murcia S, Krug I, Mercader JM, Riesco N, Saus E, Santamaría JJ and Estivill X (2009) Suicide attempts in bulimia nervosa: personality and psychopathological correlates. *European Psychiatry* **24**, 91–97.

Forrest LN, Smith AR, White RD and Joiner TE (2015) (Dis)connected: an examination of interoception in individuals with suicidality. *Journal of Abnormal Psychology* **124**, 754–764.

Franklin JC, Hessel ET and Prinstein MJ (2011) Clarifying the role of pain tolerance in suicidal capability. *Psychiatry Research* **189**, 362–367.

Franklin JC, Ribeiro JD, Fox KR, Bentley KH, Kleiman EM, Huang X, Musacchio KM, Jaroszewski AC, Chang BP and Nock MK (2017) Risk factors for suicidal thoughts and behaviors: a meta-analysis of 50 years of research. *Psychological Bulletin* **143**, 187–232.

https://doi.org/10.1017/S0033291718003872 Published online by Cambridge University Press

Franko DL, Keel PK, Dorer DJ, Blais MA, Delinsky SS, Eddy KT, Charat V, Renn R and Herzog DB (2004) What predicts suicide attempts in women with eating disorders? *Psychological Medicine* **34**, 843–853.

Fukushima H, Terasawa Y and Umeda S (2011) Association between interoception and empathy: evidence from heartbeat-evoked brain potential. *International Journal of Psychophysiology* **79**, 259–265.

Füstös J, Gramann K, Herbert BM and Pollatos O (2013) On the embodiment of emotion regulation: interoceptive awareness facilitates reappraisal. *Social Cognitive and Affective Neuroscience* **8**, 911–917.

Garfinkel SN, Seth AK, Barrett AB, Suzuki K and Critchley HD (2015) Knowing your own heart: distinguishing interoceptive accuracy from interoceptive awareness. *Biological Psychology* **104**, 65–74.

Garner DM (2004) *EDI 3: Eating Disorder Inventory-3: Professional Manual*. Lutz, FL: Psychological Assessment Resources.

Garner DM, Olmstead MP and Polivy J (1983) Development and validation of a multidimensional eating disorder inventory for anorexia nervosa and bulimia. *International Journal of Eating Disorders* **2**,, 15–34.

Gratz KL (2001) Measurement of deliberate self-harm: preliminary data on the deliberate self-harm inventory. *Journal of Psychopathology and Behavioral Assessment* **23**, 253–263.

Guan K, Fox KR and Prinstein MJ (2012) Nonsuicidal self-injury as a time-invariant predictor of adolescent suicide ideation and attempts in a diverse community sample. *Journal of Consulting and Clinical Psychology* **80**, 842–849.

Gutierrez PM, Osman A, Barrios FX and Kopper BA (2001) Development and initial validation of the Self-Harm Behavior Questionnaire. *Journal of Personality Assessment* **77**, 475–490.

Henry JD and Crawford JR (2005) The short-form version of the Depression Anxiety Stress Scales (DASS-21): construct validity and normative data in a large non-clinical sample. *British Journal of Clinical Psychology* **44**, 227–239.

Hölzel BK, Lazar SW, Gard T, Schuman-Olivier Z, Vago DR and Ott U (2011) How does mindfulness meditation work? Proposing mechanisms of action from a conceptual and neural perspective. *Perspectives on Psychological Science* **6**, 537–559.

Hooley JM and St Germain SA (2014) Should we expand the conceptualization of self-injurious behavior? Rationale, review, and recommendations. In Nock MK (ed.), *The Oxford Handbook of Suicide and Self-Injury*. Oxford, UK: Oxford University Press, pp. 47–60.

Hooley JM, Ho DT, Slater J and Lockshin A (2010) Pain perception and non-suicidal self-injury: a laboratory investigation. *Personality Disorders: Theory, Research, and Treatment* **1**, 170–179.

Hoyert DL and Xu J (2012) Deaths: preliminary data for 2011. *National Vital Statistics Reports* **61**, 1–64.

Khalsa SS, Adolphs R, Cameron OG, Critchley HD, Davenport PW, Feinstein JS, Feusner JD, Garfinkel SN, Lane RD, Mehling WE, Meuret AE, Nemeroff CB, Oppenheimer S, Petzschner FH, Pollatos O, Rhudy JL, Schramm LP, Simmons WK, Stein MB, Stephan KE, Van den Bergh O, Van Diest I, von Leupoldt A and Paulus MP, Interoception Summit 2016 participants (2018) Interoception and mental health: a roadmap. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging* **3**, 501–513.

Klonsky ED and Glenn CR (2009) Assessing the functions of non-suicidal self-injury: psychometric properties of the inventory of statements about self-injury (ISAS). *Journal of Psychopathology and Behavioral Assessment* **31**, 215–219.

Klonsky ED and May AM (2015) The three-step theory (3ST): a new theory of suicide rooted in the 'ideation-to-action' framework. *International Journal of Cognitive Therapy* **8**, 114–129.

Klonsky ED, May AM and Glenn CR (2013) The relationship between non-suicidal self-injury and attempted suicide: converging evidence from four samples. *Journal of Abnormal Psychology* **122**, 231–237.

Lewinsohn PM, Seeley JR, Roberts RE, Allen NB (1997) Center for Epidemiologic Studies Depression Scale (CES-D) as a screening instrument for depression among community-residing older adults. *Psychology and Aging* **12**, 277–287.

Linehan MM (1993) *Cognitive Behavioral Therapy of Borderline Personality Disorder*. New York: Guilford Press.

MacCormack JK and Lindquist KA (2017) Bodily contributions to emotion: Schachter's legacy for a psychological constructionist view on emotion. *Emotion Review* **9**, 36–45.

Mason W and Suri S (2012) Conducting behavioral research on Amazon's Mechanical Turk. *Behavior Research* **44**, 1–23.

May AM and Klonsky ED (2016) What distinguishes suicide attempters from suicide ideators? A meta-analysis of potential factors. *Clinical Psychology: Science and Practice* **23**, 5–20.

Mehling WE, Price C, Daubenmier JJ, Acree M, Bartmess E and Stewart A (2012) The multidimensional assessment of interoceptive awareness (MAIA). *PLoS ONE* **7**, e48230.

Muehlenkamp JJ and Gutierrez PM (2007) Risk for suicide attempts among adolescents who engage in non-suicidal self-injury. *Archives of Suicide Research* **11**, 69–82.

Muehlenkamp JJ, Swanson JD and Brausch AM (2005) Self-objectification, risk taking, and self-harm in college women. *Psychology of Women Quarterly* **29**, 24–32.

Muehlenkamp JJ, Peat CM, Claes L and Smits D (2012) Self-injury and disordered eating: expressing emotion dysregulation through the body. *Suicide and Life-Threatening Behavior* **42**, 416–425.

Muehlenkamp JJ, Bagge CL, Tull MT and Gratz KL (2013) Body regard as a moderator of the relation between emotion dysregulation and nonsuicidal self-injury. *Suicide and Life-Threatening Behavior* **43**, 479–493.

Nock MK, Joiner TE, Gordon KH, Lloyd-Richardson E and Prinstein MJ (2006) Non-suicidal self-injury among adolescents: diagnostic correlates and relation to suicide attempts. *Psychiatry Research* **144**, 65–72.

Nock MK, Holmberg EB, Photos VI and Michel BD (2007) Self-injurious thoughts and behaviors interview: development, reliability, and validity in an adolescent sample. *Psychological Assessment* **19**, 309–317.

Noma S, 'ichi UT, Ono M, Miyagi T and Murai T (2015) Differences between nonsuicidal self-injury and suicidal behavior in patients with eating disorders. *Journal of Psychiatric Practice* **21**, 198–207.

O'Connor RC, Cleare S, Eschle S, Wetherall K and Kirtley OJ (2016) The integrated motivational-volitional model of suicidal behaviour: an update. In O'Connor RC and Pirkis J (eds), *The International Handbook of Suicide Prevention*, 2nd Edn. Chichester: Wiley-Blackwell, pp. 220–240.

Peat CM and Muehlenkamp JJ (2011) Self-objectification, disordered eating, and depression: a test of mediational pathways. *Psychology of Women Quarterly* **35**, 441–450.

Pollatos O, Füstös J and Critchley HD (2012) On the generalised embodiment of pain: how interoceptive sensitivity modulates cutaneous pain perception. *Pain* **153**, 1680–1686.

Radloff LS (1977) The CES-D scale: a self-report depression scale for research in the general population. *Applied Psychological Measurement* **1**, 385–401.

Reynolds WM (1988) *SIQ, Suicidal Ideation Questionnaire: Professional Manual*. Lutz, FL: Psychological Assessment Resources.

Reynolds WM (2002) *RADS-2: Reynolds Adolescent Depression Scale*. Lutz, FL: Psychological Assessment Resources.

Rogers ML, Hagan CR and Joiner TE (2018) Examination of interoception along the suicidality continuum. *Journal of Clinical Psychology* **74**, 1004–1016.

Ross S, Heath NL and Toste JR (2009) Non-suicidal self-injury and eating pathology in high school students. *American Journal of Orthopsychiatry* **79**, 83–92.

Sim L and Zeman J (2004) Emotion awareness and identification skills in adolescent girls with bulimia nervosa. *Journal of Clinical Child and Adolescent Psychology* **33**, 760–771.

Smith A, Forrest L and Velkoff E (2018) Out of touch: interoceptive deficits are elevated in suicide attempters with eating disorders. *Eating Disorders* **26**, 52–65.

Turner BJ, Layden BK, Butler SM and Chapman AL (2013) How often, or how many ways: clarifying the relationship between non-suicidal self-injury and suicidality. *Archives of Suicide Research* **17**, 397–415.

Van Orden KA, Witte TK, Cukrowicz KC, Braithwaite SR, Selby EA and Joiner Jr TE (2010) The interpersonal theory of suicide. *Psychological Review* **117**, 575–600.

Venta A, Mellick W, Schatte D and Sharp D (2014) Preliminary evidence that thoughts of thwarted belongingness mediate the relations between

https://doi.org/10.1017/S0033291718003872 Published online by Cambridge University Press

level of attachment insecurity and depression and suicide-related thoughts in inpatient adolescents. *Journal of Social and Clinical Psychology* **33**, 428–447.

Victor SE and Klonsky ED (2014) Correlates of suicide attempts among self-injurers: a meta-analysis. *Clinical Psychology Review* **34**, 282–297.

Weiss S, Sack M, Henningsen P and Pollatos O (2014) On the interaction of self-regulation, interoception and pain perception. *Psychopathology* **47**, 377–382.

World Health Organization – Preventing suicide: A global imperative. Available at http://www.who.int/mental_health/suicide-prevention/world_report_2014/en/) (Accessed 8 September 2014).

https://doi.org/10.1017/S0033291718003872 Published online by Cambridge University Press