Closed form expressions for two harmonic continued fractions

MARTIN BUNDER, JOSEPH TONIEN

Introduction

A continued fraction is an expression of the form

$$a_0 + \cfrac{b_0}{a_1 + \cfrac{b_1}{a_2 + \cfrac{b_2}{\ddots}}}.$$

The expression can continue for ever, in which case it is called an *infinite* continued fraction, or it can stop after some term, when we call it a *finite* continued fraction. For irrational numbers, a continued fraction expansion often reveals beautiful number patterns which remain obscured in their decimal expansion. The interested reader is referred to [1] for a collection of many interesting continued fractions for famous mathematical constants.

When we truncate a continued fraction after some number of terms, we get what is called a *convergent*. In particular, the following finite continued fraction

$$a_0 + \cfrac{b_0}{a_1 + \cfrac{b_1}{\ddots + \cfrac{b_{n-1}}{a_n}}}$$

is called the nth convergent of the above mentioned continued fraction. If the limit, as n approaches ∞, of the nth convergent exists, we say that the infinite continued fraction converges and the limit is its value.

If in the above continued fraction all numerators b_i are 1, we denote it by $[a_0, a_1, a_2, a_3, \dots]$, and its nth convergent by $[a_0, a_1, a_2, a_3, \dots, a_n]$. If in addition the coefficients a_n are positive integers, it is not hard to prove that the infinite continued fraction always converges. Continued fractions of this type have applications in cryptography [2, 3]. When the coefficients a_n are positive real numbers, there is a classical theorem due to Seidel and Stern [4, 5], dating back to the 1840s:

The Seidel-Stern Theorem [6, 7]: If $a_n > 0$ then $[a_0, a_1, a_2, a_3, \dots]$ converges if, and only if, $\sum a_n$ diverges.

Since the harmonic series $\sum \frac{1}{n}$ diverges, by the Seidel-Stern Theorem, the infinite continued fraction $[\frac{t}{1}, \frac{t}{2}, \frac{t}{3}, \dots]$ converges for any positive real number t. Let us call these continued fractions the *harmonic continued fractions* and denote them by

https://doi.org/10.1017/mag.2017.125 Published online by Cambridge University Press

$$HCF(t) = \frac{t}{1} + \cfrac{1}{\frac{t}{2} + \cfrac{1}{\frac{t}{3} + \cfrac{1}{\ddots}}}.$$

When $t = 1$ and $t = 2$, the values of the harmonic continued fractions are known.

Theorem 1:

$$HCF(1) = \frac{1}{1} + \cfrac{1}{\frac{1}{2} + \cfrac{1}{\frac{1}{3} + \cfrac{1}{\frac{1}{4} + \cfrac{1}{\ddots}}}} = \frac{2}{\pi - 2}.$$

Theorem 2:

$$HCF(2) = \frac{2}{1} + \cfrac{1}{\frac{2}{2} + \cfrac{1}{\frac{2}{3} + \cfrac{1}{\frac{2}{4} + \cfrac{1}{\ddots}}}} = \frac{1}{2\ln 2 - 1}.$$

The two harmonic continued fractions in Theorem 1 and Theorem 2 are derived from the following continued fractions of Euler [8]:

$$1 + \cfrac{1}{1 + \cfrac{1 \times 2}{1 + \cfrac{2 \times 3}{1 + \cfrac{3 \times 4}{\ddots}}}} = \frac{\pi}{2}, \qquad 2 + \cfrac{1 \times 2}{2 + \cfrac{2 \times 3}{2 + \cfrac{3 \times 4}{2 + \cfrac{4 \times 5}{\ddots}}}} = \frac{1}{2\ln 2 - 1}.$$

Euler's proof involved the Wallis product and differential equations. In this paper, we offer an elementary and direct proof of Theorem 1 and Theorem 2. Our proof uses the Euler-Wallis recurrence formula to establish a *closed form formula* for the convergents of the continued fractions $HCF(1)$ and $HCF(2)$.

Theorem 3: For any natural number $n \geqslant 1$,

$$\frac{1}{1} + \cfrac{1}{\frac{1}{2} + \cfrac{1}{\frac{1}{3} + \cfrac{1}{\ddots + \cfrac{1}{\frac{1}{2n-1}}}}} = \frac{1}{\frac{1}{2n}\frac{2^2 4^2 \ldots (2n)^2}{1^2 3^2 \ldots (2n-1)^2} - 1},$$

https://doi.org/10.1017/mag.2017.125 Published online by Cambridge University Press

$$\frac{1}{1}+\cfrac{1}{\frac{1}{2}+\cfrac{1}{\frac{1}{3}+\cfrac{1}{\ddots+\cfrac{1}{\frac{1}{2n}}}}}=\cfrac{1}{\frac{1}{2n+1}\cfrac{2^2 4^2 \ldots (2n)^2}{1^2 3^2 \ldots (2n-1)^2}-1}.$$

Theorem 4: For any natural number $n \geqslant 1$,

$$\frac{2}{1}+\cfrac{1}{\frac{2}{2}+\cfrac{1}{\frac{2}{3}+\cfrac{1}{\ddots+\cfrac{1}{\frac{2}{n}}}}}=\cfrac{1}{\sum_{i=0}^{n-1}\frac{(-1)^i}{(i+1)(i+2)}}.$$

Euler-Wallis recurrence formulas

The following theorem due to William Brouncker [1620-1684], the first President of the Royal Society – is called *the fundamental theorem of continued fractions*. It gives us recursive formulas to calculate the numerator p_n and the denominator q_n of the convergents. John Wallis [1616-1703] and Leonhard Euler [1707-1783] in [8] made extensive use of these formulas which are now called the Euler-Wallis formulas.

Theorem 5: For any $n \geqslant 0$, the nth convergent can be determined as

$$\left[a_0, a_1, a_2, \ldots, a_n\right] = \frac{p_n}{q_n},$$

where the sequences $\{p_n\}_{n \geqslant -2}$ and $\{q_n\}_{n \geqslant -2}$ are specified as follows:

$$p_{-2} = 0, p_{-1} = 1, p_n = a_n p_{n-1} + p_{n-2}, \text{ for all } n \geqslant 0,$$

$$q_{-2} = 1, q_{-1} = 0, q_n = a_n q_{n-1} + q_{n-2}, \text{ for all } n \geqslant 0.$$

The theorem can be proved easily by induction, since

$$\begin{aligned}\left[a_0, a_1, a_2, \ldots, a_n, a_{n+1}\right] &= \left[a_0, a_1, a_2, \ldots, a_n + \frac{1}{a_{n+1}}\right] \\ &= \frac{\left(a_n + \frac{1}{a_{n+1}}\right)p_{n-1} + p_{n-2}}{\left(a_n + \frac{1}{a_{n+1}}\right)q_{m-1} + q_{n-2}} \\ &= \frac{a_{n+1}(a_n p_{n-1} + p_{n-2}) + p_{n-1}}{a_{n+1}(a_n q_{n-1} + q_{n-2}) + q_{n-1}} \\ &= \frac{a_{n+1}p_n + p_{n-1}}{a_{n+1}q_n + q_{n-1}} = \frac{p_{n+1}}{q_{n+1}}.\end{aligned}$$

https://doi.org/10.1017/mag.2017.125 Published online by Cambridge University Press

Sometimes we want to investigate the sub-sequences $\{p_{2n}\}$, $\{p_{2n-1}\}$, $\{q_{2n}\}$, $\{q_{2n-1}\}$ and the following theorem is useful in those scenarios.

Theorem 6: The convergents $\left[a_0, a_1, a_2, \ldots, a_n\right] = \dfrac{p_n}{q_n}$ satisfy the following:

$$p_n = \left(a_n a_{n-1} + \frac{a_n}{a_{n-2}} + 1\right)p_{n-2} - \frac{a_n}{a_{n-2}}p_{n-4}, \text{ for all } n \geqslant 2,$$

$$q_n = \left(a_n a_{n-1} + \frac{a_n}{a_{n-2}} + 1\right)q_{n-2} - \frac{a_n}{a_{n-2}}q_{n-4}, \text{ for all } n \geqslant 2.$$

Proof: By the Euler-Wallis formula,

$$p_{n-2} = a_{n-2}p_{n-3} + p_{n-4}, \text{ for all } n \geqslant 2,$$

$$p_{n-3} = \frac{1}{a_{n-2}}p_{n-2} - \frac{1}{a_{n-2}}p_{n-4},$$

$$\begin{aligned} p_{n-1} &= a_{n-1}p_{n-2} + p_{n-3} \\ &= a_{n-1}p_{n-2} + \frac{1}{a_{n-2}}p_{n-2} - \frac{1}{a_{n-2}}p_{n-4}, \end{aligned}$$

$$\begin{aligned} p_n &= a_n p_{n-1} + p_{n-2} \\ &= a_n a_{n-1}p_{n-2} + \frac{a_n}{a_{n-2}}p_{n-2} - \frac{a_n}{a_{n-2}}p_{n-4} + p_{n-2}, \end{aligned}$$

$$p_n = \left(a_n a_{n-1} + \frac{a_n}{a_{n-2}} + 1\right)p_{n-2} - \frac{a_n}{a_{n-2}}p_{n-4}.$$

The relation for q_n is proved similarly.

Finding HCF (1)

The following theorem establishes closed form formulas for the numerator p_n and the denominator q_n of the convergents of *HCF* (1). Our Theorem 3 follows from this theorem.

Theorem 7: The numerator p_n and the denominator q_n of the convergents of *HCF* (1) are

$$p_{2n} = \prod_{i=1}^{n} \frac{2i+1}{2i}, \text{ for all } n \geqslant 0,$$

$$p_{2n+1} = \prod_{i=1}^{n+1} \frac{2i+1}{2i}, \text{ for all } n \geqslant 0,$$

https://doi.org/10.1017/mag.2017.125 Published online by Cambridge University Press

$$q_{2n} = \left(\frac{1}{2n+2}\frac{2^2 4^2 \ldots (2n+2)^2}{1^2 3^2 \ldots (2n+1)^2} - 1\right)\prod_{i=1}^{n}\frac{2i+1}{2i}, \text{ for all } n \geqslant 0,$$

$$q_{2n+1} = \left(\frac{1}{2n+3}\frac{2^2 4^2 \ldots (2n+2)^2}{1^2 3^2 \ldots (2n+1)^2} - 1\right)\prod_{i=1}^{n+1}\frac{2i+1}{2i}, \text{ for all } n \geqslant 0.$$

Proof: By Theorem 6

$$p_n = \left(\frac{1}{n(n+1)} + \frac{n-1}{n+1} + 1\right)p_{n-2} - \frac{n-1}{n+1}p_{n-4}$$

$$= \frac{2n^2+1}{n(n+1)}p_{n-2} - \frac{n-1}{n+1}p_{n-4}, \text{ for all } n \geqslant 2.$$

So

$$p_{2n} = \frac{8n^2+1}{2n(2n+1)}p_{2n-2} - \frac{2n-1}{2n+1}p_{2n-4}, \text{ for all } n \geqslant 1,$$

$$p_{2n+1} = \frac{8n^2+8n+3}{(2n+1)(2n+2)}p_{2n-1} - \frac{2n}{2n+2}p_{2n-3}, \text{ for all } n \geqslant 1.$$

For each $n \geqslant -1$, let

$$p_{2n} = p'_{2n}\prod_{i=1}^{n}\frac{2i+1}{2i} \quad \text{and} \quad p_{2n+1} = p'_{2n+1}\prod_{i=1}^{n+1}\frac{2i+1}{2i},$$

then

$$\frac{2n+1}{2n}\frac{2n-1}{2n-2}p'_{2n} = \frac{8n^2+1}{2n(2n+1)}\frac{2n-1}{2n-2}p'_{2n-2} - \frac{2n-1}{2n+1}p'_{2n-4}, \text{for all } n \geqslant 2,$$

$$\frac{2n+3}{2n+2}\frac{2n+1}{2n}p'_{2n+1} = \frac{8n^2+8n+3}{(2n+1)(2n+2)}\frac{2n+1}{2n}p'_{2n-1} - \frac{2n}{2n+2}p'_{2n-3}, \text{for all } n \geqslant 1.$$

This simplifies to

$$(2n+1)^2 p'_{2n} = \left(8n^2+1\right)p'_{2n-2} - 2n(2n-2)p'_{2n-4}, \text{for all } n \geqslant 2,$$

$$(2n+3)(2n+1)p'_{2n+1} = \left(8n^2+8n+3\right)p'_{2n-1} - 4n^2 p'_{2n-3}, \text{for all } n \geqslant 1,$$

and so

$$(2n+1)^2\left(p'_{2n} - p'_{2n-2}\right) = 2n(2n-2)\left(p'_{2n-2} - p'_{2n-4}\right), \text{for all } n \geqslant 2,$$

$$(2n+3)(2n+1)\left(p'_{2n+1} - p'_{2n-1}\right) = 4n^2\left(p'_{2n-1} - p'_{2n-3}\right), \text{for all } n \geqslant 1.$$

Therefore

$$p'_{2n} - p'_{2n-2} = \frac{2n(2n-2)}{(2n+1)^2}\left(p'_{2n-2} - p'_{2n-4}\right), \text{ for all } n \geqslant 2, \qquad (1)$$

$$p'_{2n+1} - p'_{2n-1} = \frac{4n^2}{(2n+3)(2n+1)}\left(p'_{2n-1} - p'_{2n-3}\right), \text{for all } n \geqslant 1. \qquad (2)$$

https://doi.org/10.1017/mag.2017.125 Published online by Cambridge University Press

We have $p_{-1} = 1, p_0 = 1, p_1 = \frac{3}{2}, p_2 = \frac{3}{2}$, so

$$p'_{-1} = p'_0 = p'_1 = p'_2 = 1.$$

It follows that

$$p'_n = 1, \text{ for all } n \geqslant -1.$$

This gives us the desired closed form formula for p_n:

$$p_{2n} = \prod_{i=1}^{n} \frac{2i+1}{2i} \qquad \text{and} \qquad p_{2n+1} = \prod_{i=1}^{n+1} \frac{2i+1}{2i}, \text{ for all } n \geqslant 0.$$

Defining a sequence $\{q'_n\}$ for $\{q_n\}$ similar to $\{p'_n\}$ for $\{p_n\}$, we seek for $\{q'_n\}$ equations corresponding to (1) and (2). With $q_{-1} = 0$, $q_0 = 1$, $q_1 = \frac{1}{2}, q_2 = \frac{7}{6}$, we have

$$q'_{-1} = 0, q'_0 = 1, q'_1 = \frac{1}{3}, q'_2 = \frac{7}{9}.$$

This gives us

$$\begin{aligned} q'_{2n} - q'_{2n-2} &= \frac{2n(2n-2)}{(2n+1)^2} \cdots \frac{(4)(2)}{5^2}(q'_2 - q'_0) \\ &= -\frac{2n(2n-2)}{(2n+1)^2} \cdots \frac{(4)(2)}{5^2}\frac{2}{9} \\ &= -\frac{1}{2n}\frac{2^2 4^2 \ldots (2n)^2}{1^2 3^2 \ldots (2n+1)^2}, \text{ for all } n \geqslant 1 \\ q'_{2n+1} - q'_{2n-1} &= \frac{(2n)^2}{(2n+3)(2n+1)} \cdots \frac{2^2}{(5)(3)}(q'_1 - q'_{-1}) \\ &= \frac{(2n)^2}{(2n+3)(2n+1)} \cdots \frac{2^2}{(5)(3)}\frac{1}{3} \\ &= \frac{1}{2n+3}\frac{2^2 4^2 \ldots (2n)^2}{1^2 3^2 \ldots (2n+1)^2}, \text{ for all } n \geqslant 1. \end{aligned}$$

Simple algebraic manipulation gives us

$$q'_{2n} - q'_{2n-2} = \frac{1}{2n+2}\frac{2^2 4^2 \ldots (2n+2)^2}{1^2 3^2 \ldots (2n+1)^2} - \frac{1}{2n}\frac{2^2 4^2 \ldots (2n)^2}{1^2 3^2 \ldots (2n-1)^2}, \text{for all } n \geqslant 1,$$

$$q'_{2n+1} - q'_{2n-1} = \frac{1}{2n+3}\frac{2^2 4^2 \ldots (2n+2)^2}{1^2 3^2 \ldots (2n+1)^2} - \frac{1}{2n+1}\frac{2^2 4^2 \ldots (2n)^2}{1^2 3^2 \ldots (2n-1)^2}, \text{for all } n \geqslant 1.$$

https://doi.org/10.1017/mag.2017.125 Published online by Cambridge University Press

Replacing n by i and summing over i, from 1 to n, we find that

$$q'_{2n} - q'_0 = \frac{1}{2n+2}\frac{2^2 4^2 \ldots (2n+2)^2}{1^2 3^2 \ldots (2n+1)^2} - \frac{1}{2}\frac{2^2}{1^2}, \text{ for all } n \geqslant 0,$$

$$q'_{2n+1} - q'_1 = \frac{1}{2n+3}\frac{2^2 4^2 \ldots (2n+2)^2}{1^2 3^2 \ldots (2n+1)^2} - \frac{1}{3}\frac{2^2}{1^2}, \text{ for all } n \geqslant 0.$$

Therefore

$$q'_{2n} = \frac{1}{2n+2}\frac{2^2 4^2 \ldots (2n+2)^2}{1^2 3^2 \ldots (2n+1)^2} - 1, \text{ for all } n \geqslant 0,$$

$$q'_{2n+1} = \frac{1}{2n+3}\frac{2^2 4^2 \ldots (2n+2)^2}{1^2 3^2 \ldots (2n+1)^2} - 1, \text{ for all } n \geqslant 0.$$

From here we obtain the desired closed form formula for q_n.

Proof of Theorem 1: By Theorem 7,

$$\frac{q_{2n}}{p_{2n}} = \frac{1}{2n+2}\frac{2^2 4^2 \ldots (2n+2)^2}{1^2 3^2 \ldots (2n+1)^2} - 1,$$

$$\frac{q_{2n+1}}{p_{2n+1}} = \frac{1}{2n+3}\frac{2^2 4^2 \ldots (2n+2)^2}{1^2 3^2 \ldots (2n+1)^2} - 1.$$

Now

$$\begin{aligned}\Gamma\left(n+\frac{3}{2}\right) &= \left(n+\frac{1}{2}\right)\Gamma\left(n+\frac{1}{2}\right) = \left(n+\frac{1}{2}\right)\left(n-\frac{1}{2}\right)\Gamma\left(n-\frac{1}{2}\right) \\ &= \left(n+\frac{1}{2}\right)\left(n-\frac{1}{2}\right)\ldots\frac{1}{2}\Gamma\left(\frac{1}{2}\right) \\ &= \frac{2n+1}{2}\frac{2n-1}{2}\ldots\frac{1}{2}\sqrt{\pi}.\end{aligned}$$

It follows that

$$\frac{q_{2n}}{p_{2n}} = \frac{\pi}{2}\frac{(n+1)\Gamma(n+1)^2}{\Gamma\left(n+\frac{3}{2}\right)^2} - 1 \to \frac{\pi}{2} - 1.$$

The above limit is due to the fact [9, 10, 11] that, for any complex number a,

$$\lim_{n\to\infty}\frac{n^a\Gamma(n)}{\Gamma(n+a)} = 1;$$

we have used $a = \frac{1}{2}$.

Further

$$\lim_{n\to\infty}\frac{q_{2n+1}}{p_{2n+1}} = \lim_{n\to\infty}\frac{q_{2n}}{p_{2n}} = \frac{\pi}{2} - 1.$$

The reciprocal of this limit shows that $HCF(1) = \dfrac{2}{\pi - 2} \approx 1.75.$

https://doi.org/10.1017/mag.2017.125 Published online by Cambridge University Press

Finding HCF (2)

The following theorem establishes closed form formulas for the numerator p_n and the denominator q_n of the convergents of *HCF* (2). Our Theorem 4 follows from this theorem.

Theorem 8: For any $n \geqslant 0$, the numerator p_n and the denominator q_n of the convergents of *HCF* (2) are

$$p_n = n + 2,$$

$$q_n = (n + 2) \sum_{i=0}^{n} \frac{(-1)^i}{(i + 1)(i + 2)}.$$

Proof: By the Euler-Wallis formula, the convergents $\frac{p_n}{q_n}$ are determined by the following recurrence relation

$$p_{-2} = 0,\ p_{-1} = 1,\ p_n = \frac{2}{n + 1} p_{n-1} + p_{n-2}, \text{ for all } n \geqslant 0,$$

$$q_{-2} = 1,\ q_{-1} = 0,\ q_n = \frac{2}{n + 1} q_{n-1} + q_{n-2}, \text{ for all } n \geqslant 0.$$

We have

$$(n + 1)p_n = 2p_{n-1} + (n + 1)p_{n-2}, \text{ for all } n \geqslant 0,$$

$$(n + 1)p_n + np_{n-1} = (n + 2)p_{n-1} + (n + 1)p_{n-2}, \text{ for all } n \geqslant 0,$$

$$(-1)^n(n + 1)p_n - (-1)^{n-1}np_{n-1} = (-1)^n(n + 2)p_{n-1} - (-1)^{n-1}(n + 1)p_{n-2},$$

$$\text{for all } n \geqslant 0.$$

Taking summation, we have

$$(-1)^n(n + 1)p_n = (-1)^n(n + 2)p_{n-1} - (-1)^{-1}1p_{-2}, \text{for all } n \geqslant 0,$$

$$(n + 1)p_n = (n + 2)p_{n-1} + (-1)^n p_{-2}, \text{for all } n \geqslant 0.$$

Since $p_{-2} = 0$, it follows that

$$(n + 1)p_n = (n + 2)p_{n-1}, \text{ for all } n \geqslant 0,$$

$$\frac{p_n}{n + 2} = \frac{p_{n-1}}{n + 1} = \ldots = \frac{p_{-1}}{1} = 1, \text{ for all } n \geqslant 0,$$

$$p_n = n + 2, \text{ for all } n \geqslant 0.$$

Similarly,

$$(n + 1)q_n = (n + 2)q_{n-1} + (-1)^n q_{-2}, \text{ for all } n \geqslant 0.$$

Since $q_{-2} = 1$, it follows that

$$(n + 1)q_n = (n + 2)q_{n-1} + (-1)^n, \text{ for all } n \geqslant 0,$$

https://doi.org/10.1017/mag.2017.125 Published online by Cambridge University Press

$$\frac{q_n}{n+2} = \frac{q_{n-1}}{n+1} + \frac{(-1)^n}{(n+1)(n+2)}, \text{ for all } n \geqslant 0.$$

Replacing n by i and summing from $i = 0$ to $i = n$, we obtain

$$\frac{q_n}{n+2} = \frac{q_{-1}}{1} + \sum_{i=0}^{n} \frac{(-1)^i}{(i+1)(i+2)} = \sum_{i=0}^{n} \frac{(-1)^i}{(i+1)(i+2)}, \text{for all } n \geqslant 0,$$

and this gives the desired closed form formula for q_n.

Proof of Theorem 2: By Theorem 8,

$$\frac{q_n}{p_n} = \sum_{i=0}^{n} \frac{(-1)^i}{(i+1)(i+2)}.$$

So

$$\lim_{n\to\infty} \frac{q_n}{p_n} = \lim_{n\to\infty} \sum_{i=0}^{n} \frac{(-1)^i}{(i+1)(i+2)} = \lim_{n\to\infty} \sum_{i=0}^{n} (-1)^i \left(\frac{1}{i+1} - \frac{1}{i+2}\right)$$

$$= \lim_{n\to\infty} \sum_{i=0}^{n} \left(\frac{(-1)^i}{i+1} + \frac{(-1)^{i+1}}{i+2}\right)$$

$$= -1 + 2 \lim_{n\to\infty} \sum_{i=0}^{n} \frac{(-1)^i}{i+1} = 2\ln 2 - 1.$$

The reciprocal of this limit gives $HCF(2) = \dfrac{1}{2\ln 2 - 1} \approx 2.59$.

Conclusion

We have established explicit formulas for the convergents of the first two harmonic continued fractions $HCF(1)$ and $HCF(2)$ in Theorem 3 and Theorem 4, respectively. From these formulas, we derive the values of $HCF(1)$ and $HCF(2)$ as in Theorem 1 and Theorem 2.

We can see that the result for $HCF(2)$ is relatively easy to establish compared to that for $HCF(1)$, as in Theorem 4 we need only one formula for any length of convergent but in Theorem 3 we have one formula for the odd-length convergents and another one for the even-length convergents of $HCF(1)$.

We have not seen any result on $HCF(t)$ for a general t, but we conjecture that when t is an odd integer, we may need two formulas for odd-length and even-length convergents, and that the value of $HCF(t)$ may involve π when t is odd and involve $\ln 2$ when t is even. Obviously, $HCF(0) = 1$ and $HCF(t) \approx t + \dfrac{2}{t}$ as $t \to \infty$.

Acknowledgement

The authors wish to thank the referee for many helpful suggestions.

https://doi.org/10.1017/mag.2017.125 Published online by Cambridge University Press

References

1. C. D. Olds, *Continued fractions*, The Mathematical Association of America (1963).
2. M. Wiener, Cryptanalysis of short RSA secret exponents, *IEEE Transactions on Information Theory* **36** (1990) pp. 553-558.
3. M. Bunder and J. Tonien, A new improved attack on RSA, *Proceedings of the 5th International Cryptology and Information Security Conference* (2016) pp. 101-110.
4. L. Seidel, *Untersuchungen über die Konvergenz und Divergenz der Kettenbrüche*, Habilschrift München (1846).
5. M. A. Stern, Über die Kennzeichen der Konvergenz eines Kettenbruchs, *Journal für die Reine und Angewandte Mathematik* **37** (1848) pp. 255-272.
6. L. Lorentzen and H. Waadeland, *Continued fractions, Volume 1, Convergence theory*, Atlantis Press (2008).
7. A. F. Beardon and I. Short, The Seidel, Stern, Stolz and Van Vleck theorems on continued fractions, *Bulletin of the London Mathematical Society* **42**(3) (2010) pp. 457-466.
8. L. Euler, De fractionibus continuis observationes, *Commentarii Academiae Scientiarum Imperialis Petropolitanae* **11** (1750) pp. 32-81.
9. J. S. Frame, An approximation to the quotient of gamma function, *The American Mathematical Monthly* **56**(8) (1949) pp. 529-535.
10. F. G. Tricomi and A. Erdélyi, The asymptotic expansion of a ratio of gamma functions, *Pacific Journal of Mathematics* **1**(1) (1951) pp. 133-142.
11. M. Abramowitz and I. Stegun, *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, National Bureau of Standards (1964).

10.1017/mag.2017.125

MARTIN BUNDER

School of Mathematics and Applied Statistics, University of Wollongong, Australia

e-mail: *mbunder@uow.edu.au*

JOSEPH TONIEN

Institute of Cybersecurity and Cryptology, School of Computing and Information Technology, University of Wollongong, Australia

e-mail: *joseph_tonien@uow.edu.au*

https://doi.org/10.1017/mag.2017.125 Published online by Cambridge University Press