J. Plasma Physics (1997), *vol.* 57, *part* 1, *pp.* 83–87

Copyright © 1997 Cambridge University Press

Reduced MHD in nearly potential magnetic fields

H. R. STRAUSS

Courant Institute of Mathematical Sciences, New York University,
251 Mercer Street, New York, New York 10012, USA

(Received 25 March 1996)

Reduced, approximate MHD equations are derived for the case where the magnetic field is close to a potential field. The potential field can have an arbitrary three-dimensional structure, as long as it is non-vanishing. Finite current and pressure effects are included.

1. Introduction

Reduced MHD equations (Strauss 1976, 1977, 1980, 1982, 1983, 1984) have been useful for numerical solution (Dahlburg *et al.* 1985, 1986; Strauss and Hossain 1990), as well as being convenient for linear and nonlinear analysis. An important feature of reduced MHD systems is that they eliminate the fast compressional Alfvén wave, leaving the shear Alfvén branch and slow magnetosonic waves.

All reduced sets of equations have essentially the same structure. The main time-dependent variables are the magnetic field components transverse to the background, vacuum magnetic field, as well as velocity components transverse to the vacuum field. These transverse fields are approximately incompressible, and hence are conveniently represented by streamfunctions: the poloidal magnetic flux ψ and the electric potential ϕ. Additional equations may be used for the density ρ, the parallel velocity $v_\parallel$, and the pressure p or temperature.

Several versions of reduced MHD have been developed, appropriate to various magnetic field geometries. These versions are appropriate for tokamaks (Strauss 1977, 1983), stellarators (Strauss 1980), mirror machines (Strauss 1982) and reversed-field pinches (Strauss 1984). Several of these can be unified by the assumptions that (i) the magnetic field is close to a vacuum field and (ii) the perturbations of interest are nearly resonant with the magnetic field.

The first assumption implies weak currents and low β. This is the case in many magnetic fusion experiments, including tokamaks, stellerators and tandem mirrors. It is also the case in solar and stellar coronal magnetic fields, and in regions of the geomagnetic field.

A deficiency of some reduced models is that they do not include flow along the magnetic field. As pointed out by Montgomery (1992), this precludes equilibria with helical flows. This has been corrected in extensions of the reduced MHD model, such as the four-field model (Hazeltine *et al.* 1985). A better approximation can be based on low β, and small ratio of poloidal to toroidal magnetic field (ϵ/q) (Strauss 1983; Izzo *et al.* 1985). As well as including parallel flow, this approach yields the exact two dimensional Grad–Shafranov

https://doi.org/10.1017/S0022377896005296 Published online by Cambridge University Press

equilibrium equation, as well as the correct Mercier interchange stability criterion. A similar two-fluid model was developed by Drake and Antonsen (1984).

Another weakness of some reduced models is that they conserve a non-standard energy. The equations presented here conserve the usual energy.

2. MHD equations and ordering

The starting point is the usual MHD equations, which may be expressed in the form

$$\mathbf{B} = \nabla \times \mathbf{A}, \tag{1}$$

$$\mathbf{J} = \nabla \times \mathbf{B}, \tag{2}$$

$$\frac{\partial \mathbf{A}}{\partial t} = \mathbf{v} \times \mathbf{B} + \nabla \phi, \tag{3}$$

$$\rho \frac{d\mathbf{v}}{dt} = \mathbf{J} \times \mathbf{B} - \nabla p, \tag{4}$$

$$\frac{\partial \rho}{dt} = -\nabla \cdot (\rho \mathbf{v}). \tag{5}$$

The pressure is assumed to be adiabatic:

$$r = c\rho^{\gamma}, \tag{6}$$

where c is a constant. It is straightforward to use a more general energy equation. This system conserves energy $\mathscr{E}$, given by

$$\mathscr{E} = \frac{1}{2} \int d^3x \left(\rho |\mathbf{v}|^2 + |\mathbf{B}|^2 + 2 \frac{p}{\gamma - 1} \right), \tag{7}$$

assuming that the normal components of $\mathbf{B}$ and $\mathbf{v}$ vanish on the boundary of the domain of integration.

In the following reduced model the magnetic field is close to a potential or 'vacuum' field $\mathbf{B}_v$, which satisfies

$$\nabla \cdot \mathbf{B}_v = \nabla \times \mathbf{B}_v = 0. \tag{8}$$

Hence $\mathbf{B}_v$ can be expressed in the form

$$\mathbf{B}_v = \nabla \chi, \tag{9}$$

where χ is the magnetic potential. The total magnetic field deviates from a potential field by

$$\mathbf{B} = \mathbf{B}_v + \mathbf{B}_1. \tag{10}$$

The ratio

$$\lambda = \frac{B_1}{B_v} \tag{11}$$

will be used as a small expansion parameter, where $B_v = |\mathbf{B}_v|$ and $B_1 = |\mathbf{B}_1|$.

We now introduce an ordering in λ. The potential field strength B_v, the density, and the basic length scale L are of order λ^0. The Alfvén speed $v_A = B_v/\rho^{1/2}$ and Alfvén time L/v_A are also λ^0.

https://doi.org/10.1017/S0022377896005296 Published online by Cambridge University Press

The fundamental assumption is that the relevant scale length along $\mathbf{B}_v$ is $L_\parallel \sim L/\lambda$, so that

$$\mathbf{B}_v \cdot \boldsymbol{\nabla} \sim \lambda \frac{B_v}{L} \sim \frac{|\mathbf{B}_1|}{L}. \tag{12}$$

If we do not make this assumption, we shall only find linear shear Alfvén and slow magnetosonic waves. It is only when the gradient length along the vacuum field is long that the relatively weaker non-potential magnetic field effects can be important. This occurs near resonances of the magnetic field lines. Resonant perturbations satisfy the magnetic differential equation

$$\mathbf{B} \cdot \boldsymbol{\nabla} \phi = 0 \tag{13}$$

on a resonant field line. Since any field line is, in effect, close to a resonance, we adopt a maximal ordering valid for both resonant and non-resonant field lines.

Having introduced a longer length scale parallel to $\mathbf{B}_v$ we also have a longer time scale, $L/\lambda v_A$. Thus the time derivatives will be taken to be of order λ:

$$\frac{\partial}{\partial t} = \lambda \frac{\partial}{\partial t_1}. \tag{14}$$

In order that the convective terms be of the same order as the time derivatives, the velocity should be of no higher than order λ:

$$\mathbf{v} = \lambda \mathbf{v}_1. \tag{15}$$

Finally, the pressure p will be of second order:

$$p = \lambda^2 p_2, \tag{16}$$

so that $\beta = 2p/B_v^2 \sim \lambda^2$.

From the low-β and low-current assumptions, the force must vanish in order λ. This implies that the components of the current perpendicular to $\mathbf{B}_v$ should be of order λ^2. This requires that the magnetic vector potential be parallel to $\mathbf{B}_v$ in leading order:

$$\mathbf{A}_1 = \psi \mathbf{B}_v. \tag{17}$$

From (10) and (17), $\mathbf{B}_1$ may be expressed as

$$\mathbf{B}_1 = \boldsymbol{\nabla} \psi \times \mathbf{B}_v. \tag{18}$$

3. Reduced MHD equations

Taking the interior product of (3) with $\mathbf{B}$ yields, in order λ^2,

$$\frac{\partial \psi}{\partial t} = \frac{1}{B_v^2} \mathbf{B} \cdot \boldsymbol{\nabla} \phi = \frac{\mathbf{B}_v}{B_v^2} \cdot (\boldsymbol{\nabla} \phi + \boldsymbol{\nabla} \phi \times \boldsymbol{\nabla} \psi). \tag{19}$$

Next, taking the exterior product of (3) with $\mathbf{B}_v/B_v^2$ gives

$$v_\perp = \boldsymbol{\nabla} \phi \times \frac{\mathbf{B}_v}{B_v^2}. \tag{20}$$

There is also an order-λ^2 component of $\mathbf{v}$ parallel to $\mathbf{B}$, which does not contribute to $\mathbf{v} \times \mathbf{B}$ in (3). The total velocity field will be given by

$$\mathbf{v} = \mathbf{v}_\perp + \frac{v_\parallel}{B_v} \mathbf{B}. \tag{21}$$

https://doi.org/10.1017/S0022377896005296 Published online by Cambridge University Press

The reason that the last term in (21) is taken proportional to $\mathbf{B}$ rather than just $\mathbf{B}_v$ is so that the convective derivative is

$$\frac{d}{dt} = \frac{\partial}{\partial t} + \mathbf{v}_\perp \cdot \boldsymbol{\nabla} + \frac{v_\parallel}{B_v} \mathbf{B} \cdot \boldsymbol{\nabla}, \tag{22}$$

and (12) is assumed.

Similarly, (4) can be solved for the current $\mathbf{J}$:

$$\mathbf{J} = \left(\rho \frac{d\mathbf{v}}{dt} + \boldsymbol{\nabla} p \right) \times \frac{\mathbf{B}_v}{B_v^2} + \frac{\mathbf{J} \cdot \mathbf{B}_v}{B_v^2} \mathbf{B}. \tag{23}$$

Taking the curl of $\mathbf{B}_1$ and the interior product with $\mathbf{B}_v$ yields the parallel current

$$\frac{\mathbf{B}_v \cdot \boldsymbol{\nabla} \times \mathbf{B}_1}{B_v^2} = \Delta^* \psi = \frac{1}{B_v^2} \boldsymbol{\nabla}_\perp \cdot B_v^2 \boldsymbol{\nabla}_\perp \psi. \tag{24}$$

where $\boldsymbol{\nabla}_\perp$ is given by

$$\boldsymbol{\nabla}_\perp = \boldsymbol{\nabla} - \frac{\mathbf{B}_v \mathbf{B}_v}{B_v^2} \cdot \boldsymbol{\nabla} = -\frac{\mathbf{B}_v}{B_v^2} \times (\mathbf{B}_v \times \boldsymbol{\nabla}). \tag{25}$$

From (2), the divergence of (23) is zero. Taking the divergence of (23) and using (22) and (24) yields

$$\boldsymbol{\nabla}_\perp \cdot \frac{\rho}{B_v^2} \boldsymbol{\nabla}_\perp \frac{\partial \phi}{\partial t} = \mathbf{B} \cdot \boldsymbol{\nabla} \Delta^* \psi - \frac{1}{B_v^4} \boldsymbol{\nabla} B_v^2 \times \boldsymbol{\nabla} p \cdot \mathbf{B}_v - \boldsymbol{\nabla} \left(\frac{\rho}{B_v^2} \mathbf{v} \cdot \boldsymbol{\nabla} \mathbf{v}_\perp \right) \times \mathbf{B}_v. \tag{26}$$

Taking the $\mathbf{B}$ component of (4) yields

$$\rho \frac{dv_\parallel}{dt} = -\frac{1}{B_v} \mathbf{B} \cdot \boldsymbol{\nabla} p. \tag{27}$$

Now the continuity equation (5) has to be evolved:

$$\frac{\partial \rho}{\partial t} = -\boldsymbol{\nabla} \frac{\rho}{B_v^2} \times \boldsymbol{\nabla} \phi \cdot \mathbf{B}_v - \mathbf{B} \cdot \boldsymbol{\nabla} \left(\frac{\rho v_\parallel}{B} \right). \tag{28}$$

This completes the reduced system. These equations conserve the correct energy $\mathscr{E}$, given by

$$\mathscr{E} = \frac{1}{2} \int d^3x \left[\rho \left(\frac{\boldsymbol{\nabla} \phi \times \mathbf{B}_v}{B_v^2} \right)^2 + \rho v_\parallel^2 + (\boldsymbol{\nabla} \psi \times \mathbf{B}_v)^2 + 2 \frac{p}{\gamma - 1} \right]. \tag{29}$$

This can be verified by multiplying (26) by ϕ, integrating over volume, integrating by parts, and using the flux equation (19), the continuity equation (28) and pressure equation (6). This assumes the same boundary conditions as in (7). Comparing with (7), note that the vacuum magnetic field energy is separately conserved and does not appear. This is clear physically, since the only free energy is stored in the non-potential part of the field.

The boundary conditions are straightforward. At flux surfaces, which are tangent to the magnetic field, the variables ψ, ϕ, ρ, p and $v_\parallel$ are all constant (or zero). Where the magnetic field intersects a material surface, there are other possibilities. Both ψ and ϕ cannot vanish, because of (19). If perturbations of

https://doi.org/10.1017/S0022377896005296 Published online by Cambridge University Press

ϕ vanish, one has a line-tying condition, which is strongly stabilizing. If ψ perturbations vanish, one has an insulating boundary.

4. Conclusions

A general set of reduced MHD equations has been derived, consisting of (19), (21), (26)–(28) and (6). The equations conserve energy correctly. The reduced system is appropriate when the magnetic field is nearly a potential field, with small current and pressure effects. It should be useful for low-aspect-ratio tokamaks, stellarators and torsatrons, mirrors, solar and stellar coronae, and magnetospheric magnetohydrodynamics. The validity requirements are that the deviations from a potential field be sufficiently small, and that the nonlinear motions of interest be resonant or nearly resonant with the magnetic field lines.

References

Dahlburg, J. P., Montgomery, D. and Mattheus, W. H. 1985 Turbulent disruptions from the Strauss equations. *J. Plasma Phys.* **34**, 1.

Dahlburg, J. P., Montgomery, D., Doolen, G. D. and Mattheus, W. H. 1986 Large-scale disruptions in a current-carrying magnetic fluid. *J. Plasma Phys.* **35**, 1.

Drake, J. F. and Antonsen, T. M. 1984 Nonlinear reduced fluid equations for toroidal plasmas. *Phys. Fluids* **27**, 898.

Hazeltine, R. D., Kotschenreuther, M. and Morrison, P. J. 1985 *Phys. Fluids* **28**, 2466.

Izzo, R., Monticello, D. A., Park, W., Strauss, H., Manickam, J., Grimm, R. and McGuire, K. 1983 Effects of toroidicity on resistive tearing modes. *Phys. Fluids* **26**, 2240.

Montgomery, D. 1992 'Reduced' magnetohydrodynamics and minimum dissipation rates. *Phys. Fluids* **B4**, 292.

Strauss, H. 1976 Nonlinear three dimensional dynamics of noncircular tokamaks. *Phys. Fluids* **19**, 134.

Strauss, H. 1977 Dynamics of high beta tokamaks. *Phys. Fluids* **20**, 1354.

Strauss, H. 1980 Stellarator equations of motion. *Plasma Phys.* **22**, 733.

Strauss, H. 1982 Reduced MHD equations for mirror machines. *Nucl. Fusion* **22**, 893.

Strauss, H. 1983 Finite aspect ratio MHD equations for tokamaks. *Nucl. Fusion* **23**, 649.

Strauss, H. 1984 Dynamical equations for the reversed field pinch. *Phys. Fluids* **27**, 2580.

Strauss, H. and Hossain, M. 1990 Profile consistency from quasilinear and nonlinear magneto-hydrodynamic computations. *Plasma Phys.* **32**, 327.

https://doi.org/10.1017/S0022377896005296 Published online by Cambridge University Press