Ergod. Th. & Dynam. Sys. (2005), **25**, 1881–1907
doi:10.1017/S0143385705000350
© *2005 Cambridge University Press*
Printed in the United Kingdom

Multifractal analysis for countable Markov shifts

GODOFREDO IOMMI

Departamento de Matemática, Instituto Superior Técnico, Av. Rovisco Pais, 1949-001 Lisboa, Portugal
(e-mail: giommi@math.ist.utl.pt)

(*Received* 12 *November* 2004 *and accepted in final form* 23 *June* 2005)

Abstract. We study the multifractal analysis of the pointwise dimension for equilibrium measures on countable Markov shifts. The main difficulty is that the space is not compact. In order to overcome this, we use an approximation argument based on the theory of convergence of Fenchel pairs developed by Wijsman. The results of Pesin and Weiss on multifractal analysis for compact spaces are used as well. We also prove a Bowen formula for countable Markov shifts. It turns out that, in this setting, this formula provides the Hausdorff dimension of the set of recurrent points.

1. *Introduction*

This paper is devoted to the study of multifractal analysis for countable Markov shifts. In a general setting multifractal analysis can be described as follows. Let μ be a finite measure on a metric space (Σ, d) and $B(x, r) := \{y \in \Sigma : d(x, y) < r\}$. The *pointwise dimension* of μ at the point $x \in \Sigma$ is defined by

$$d_\mu(x) := \lim_{r \to 0} \frac{\log \mu(B(x, r))}{\log r},$$

provided the limit exists. This function describes the power-law behaviour of $\mu(B(x, r))$ as $r \to 0$. The pointwise dimension quantifies how concentrated a measure is around a point: the larger it is the less concentrated the measure is around that point. Note that if μ is an atomic measure supported at the point x_0 then $d_\mu(x_0) = 0$ and if $x_1 \neq x_0$ then $d_\mu(x_1) = \infty$. This function induces a decomposition of the space into level sets:

$$J_\alpha = \{x \in \Sigma : d_\mu(x) = \alpha\}, \quad J' = \{x \in \Sigma : \text{the limit } d_\mu(x) \text{ does not exist}\}.$$

The set J' is called the *irregular set*. The decomposition

$$\Sigma = \Big(\biguplus_\alpha J_\alpha \Big) \uplus J'$$

is called *multifractal decomposition*. The *multifractal spectrum* is defined by

$$f_\mu(\alpha) = \dim_{\mathrm{H}}(J_\alpha),$$

https://doi.org/10.1017/S0143385705000350 Published online by Cambridge University Press

where $\dim_H$ denotes the Hausdorff dimension. Several authors have successfully studied the case when Σ is a finite state Markov shift, see [**9**] for a general account. Pesin and Weiss [**10**] developed techniques to describe the multifractal spectrum in very general settings. They define an auxiliary function, usually denoted by $T(q)$. It is possible to describe this function by means of the thermodynamical formalism. It turns out that $T(q)$ and the multifractal spectrum form a Legendre pair, therefore it is possible to characterize the multifractal spectrum based on the properties of $T(q)$. In the finite-state case the multifractal spectrum is real analytic, concave and with bounded domain [**10, 20**].

The countable alphabet case has been studied by Nakaishi [**8**], Pollicott and Weiss [**11**] and Hanus *et al* [**5**], among others. They noticed that the multifractal spectrum does not necessarily have the same properties as in the compact setting. In particular, it can have points of non-analyticity or can have unbounded domain.

Let Σ denote the symbolic space and μ the *equilibrium measure* corresponding to the potential $\log \phi$ (see §2). Consider

$$T(q) := \begin{cases} \inf\{t : P_G(-t \log \psi + q \log \phi) \leq 0\} & \text{if } q \geq q^*, \\ \infty & \text{if } q < q^*, \end{cases}$$

where P_G denotes the Gurevich pressure (Definition 2.5) and $\log \psi$ a *metric potential* (see Definition 2.10), for precise statements see Definitions 4.2 and 4.1. We prove the following.

THEOREM 4.1. *The multifractal spectrum f_μ is the Fenchel transform of T.*

Our approach is based on an approximation argument. Subsets K of Σ that are compact and invariant will be considered. In those sets the Pesin and Weiss techniques can be applied. For each such set K we obtain a Legendre pair $(T_K, f_{\mu,K})$, where $f_{\mu,K}$ is the multifractal spectrum of the measure associated to the potential restricted to K. Using the infimal convergence introduced by Wijsman in [**22**] and [**23**], we prove the convergence of $(T_K, f_{\mu,K})$ to another Legendre pair. We prove that this limit corresponds to the function $T(q)$ and to the multifractal spectrum of the measure μ.

This approach allows us to explain the different properties of the multifractal spectrum for countable Markov shifts in terms of the Fenchel transform relation between $T(q)$ and f_μ. In particular, if $q^* = 0$ then f_μ has unbounded domain (see Proposition 4.5). The existence of points of non-analyticity of f_μ can also be understood in terms of the behaviour of $T(q)$ near q^* (see Proposition 4.6, Theorem 4.7). Sufficient conditions will be given in order for the multifractal spectrum to have unbounded domain and points of non-analyticity (see Proposition 4.7, Theorem 5.2). The case in which the function $T(q)$ is linear for $q > q' \geq 1$ is also studied (see Theorem 5.3).

This approach also provides a framework to develop the multifractal analysis on spaces that do not satisfy the big images and preimages (BIP) property (see § 5).

In the first part of this paper we prove a Bowen-type formula (see [**2**]) for countable Markov shifts. In this setting this formula provides the Hausdorff dimension of the *recurrent set* (Definition 3.1), which we denote by $\mathcal{R}$. Assume the potential $\log \psi$ to be a metric potential (see Definition 2.10). We prove the following theorem.

https://doi.org/10.1017/S0143385705000350 Published online by Cambridge University Press

THEOREM 3.1.

$$\dim_{\mathrm{H}}(\mathcal{R}) = \inf\{t : P_{\mathrm{G}}(-t \log \psi) \leq 0\}.$$

This formula was first proved for finite Markov shifts by Bowen [**2**] and Ruelle [**13**]. Mauldin and Urbanski [**7**] generalized it to the case of countable Markov shifts with the BIP property (Definition 2.9). They noticed, in contrast to the finite-state case, that the Bowen equation might not have a root and that conformal measures might not exist. We treat the case of arbitrarily topologically mixing countable Markov shifts (Definition 2.2). In this setting the Bowen formula can only account for the Hausdorff dimension of the Recurrent set. Note that it is possible for the Hausdorff dimension of this set to be strictly smaller than the Hausdorff dimension of the repeller (Example 3.3). The problem of the existence of conservative conformal measures is also treated (see Propositions 3.1 and 3.2).

2. *Preliminaries*

The purpose of this section is to define the objects and constructions needed later. Let S be a countable (infinite) set and $T = (t_{ij})_{S\times S}$ a matrix of zeros and ones. Let

$$\Sigma := \{x \in S^{\mathbb{N}\cup\{0\}} : t_{x_i x_{i+1}} = 1 \text{ for every } i \geq 0\}.$$

A cylinder of length n is defined by

$$C_{i_0 i_1 \ldots i_{n-1}} := \{x \in \Sigma : x_j = i_j \text{ for } 0 \leq j \leq n-1\}.$$

An admissible word is an element $\underline{a} \in S^n$ such that $C_{\underline{a}} \neq \emptyset$.

Definition 2.1. The function $\sigma : \Sigma \to \Sigma$ defined by $(\sigma x)_i = x_{i+1}$ is called the left shift. The dynamical system (Σ, σ) is called a topological Markov chain and the matrix T is called the transition matrix. Note that T can be represented by a directed graph G.

Definition 2.2. A topologically Markov shift (Σ, σ) is said to be topologically mixing if for every $a, b \in S$ there exists N such that for all $n \geq N$ there exists an admissible word $\underline{a}$ of length n such that $a_0 = a$ and $a_{n-1} = b$.

Unlike the finite-state case, this does not imply that some power of the transition matrix is positive. Throughout this paper we will always assume the system to be topologically mixing.

Definition 2.3. Let $\log \phi : \Sigma \to \mathbb{R}$. The variations of $\log \phi$ are defined by

$$V_n := \sup\{|\log \phi(x) - \log \phi(y)| : x, y \in \Sigma, x_i = y_i, 0 \leq i \leq n-1\}.$$

Definition 2.4. Let $\log \phi : \Sigma \to \mathbb{R}$. We say that $\log \phi$ is locally Hölder continuous (with parameter θ) if there exists $B > 0$ and $\theta \in (0, 1)$ such that, for all $n \geq 1$,

$$V_n(\log \phi) \leq B\theta^n.$$

This does not imply that $\log \phi$ is bounded. In this setting it is possible to define the pressure. The so-called Gurevich pressure, which was defined by Sarig [**15**] based on the work of Gurevich [**3**], is the most appropriate to our purposes. Let $\log \phi : \Sigma \to \mathbb{R}$ be a locally Hölder potential and fix a symbol $i_0 \in S$.

https://doi.org/10.1017/S0143385705000350 Published online by Cambridge University Press

Definition 2.5. The Gurevich pressure of $\log \phi$ is defined by

$$P_{\mathrm{G}}(\log \phi) = \lim_{n\to\infty} \frac{1}{n} \log \sum_{\sigma^n x = x} \exp\left(\sum_{i=0}^{n-1} \log \phi(\sigma^i x)\right) 1_{C_{i_0}}(x),$$

where 1_{C_i} is the indicator function of the cylinder C_i. The Gurevich pressure is convex and if $\mathcal{K} := \{K \subset \Sigma : K \text{ compact and } \sigma\text{-invariant}, K \neq \emptyset\}$ then

$$P_{\mathrm{G}}(\log \phi) = \sup\{P(\log \phi | K) : K \in \mathcal{K}\}, \tag{1}$$

where $P(\log \phi | K)$ is the topological pressure of $\log \phi$ restricted to the compact set K (for definition and properties see [**21**, Ch. 9]). If the system is topologically mixing then its value does not depend on the cylinder C_{i_0} considered in the indicator function.

Definition 2.6. The first return time map to C_i is defined by

$$\varphi_i(x) := 1_{C_i}(x) \inf\{n \geq 1 : \sigma^n x \in C_i\}.$$

Let

$$[\varphi_1 = n] := \{x \in \Sigma : \varphi_1(x) = n\}.$$

A locally Hölder continuous potential $\log \phi$ can be classified according to its recurrence properties (see [**17**]). In order to do so, consider the first return time map φ_1.

Definition 2.7. Assume that $\log \phi$ has finite Gurevich pressure $P_{\mathrm{G}}(\log \phi) = \log A$. We call $\log \phi$:

(1) *recurrent* if

$$\sum_{n\geq 1} A^{-n} \sum_{\sigma^n x = x} \exp\left(\sum_{i=0}^{n-1} \log \phi(\sigma^i x)\right) 1_{C_1}(x) = \infty$$

and *transient* otherwise;

(2) *positive recurrent* if it is recurrent and

$$\sum_{n\geq 1} nA^{-n} \sum_{\sigma^n x = x} \exp\left(\sum_{i=0}^{n-1} \log \phi(\sigma^i x)\right) 1_{[\varphi_1 = n]}(x) < \infty;$$

(3) *null recurrent* if it is recurrent and

$$\sum_{n\geq 1} nA^{-n} \sum_{\sigma^n x = x} \exp\left(\sum_{i=0}^{n-1} \log \phi(\sigma^i x)\right) 1_{[\varphi_1 = n]}(x) = \infty.$$

Remark 2.1. If we consider the potential $\log \phi = 0$ then we say that a graph G is recurrent (positive or null) or transient according to the previous definition. The number $P_{\mathrm{G}}(0)$ is called the Gurevich entropy. See [**3, 4**] for properties and related concepts.

In [**17**] Sarig generalizes the Ruelle–Perron–Frobenius theorem to countable Markov shifts. Let $\log \phi$ be a locally Hölder continuous potential of finite pressure $\log A$, and let

$$L_{\log \phi} g(x) := \sum_{\sigma y = x} \exp(\log \phi(y)) g(y)$$

denote the Ruelle operator (see [**17**]). Sarig proves that if $\log \phi$ is:

https://doi.org/10.1017/S0143385705000350 Published online by Cambridge University Press

(1) positive recurrent, then there exist a conservative conformal measure ν and a continuous function g such that $L^*_{\log\phi}\nu = A\nu$, $L_{\log\phi} g = Ag$ and $\int g\,d\nu < \infty$;
(2) null recurrent, then there exist a conservative conformal measure ν and a continuous function g such that $L^*_{\log\phi}\nu = A\nu$, $L_{\log\phi} g = Ag$ and $\int g\,d\nu = \infty$;
(3) transient, then there is no conservative conformal measure.

In the positive recurrent case the measure $m = g\,d\nu$ is an equilibrium measure provided $-\int \log\phi\,dm < \infty$.

The inducing procedure in the context of topological Markov chains will be summarized. Let (Σ, σ) be a topologically mixing Markov chain with transition matrix $T = (t_{ij})_{S\times S}$. The *induced system* on the symbol 1 will be denoted by $(\Sigma_1, \overline{\sigma})$. It is defined as the full-shift on the new alphabet

$$\{C_{\overline{a}} : a_i = 1 \text{ if and only if } i = 0, C_{\overline{a}1} \neq \emptyset\}.$$

Recall that φ_1 is the first return map to C_1. For every locally Hölder potential $\log\phi : \Sigma \to \mathbb{R}$ set

$$\overline{\log\phi} := \left(\sum_{k=0}^{\varphi_1 - 1} \log\phi \circ \sigma^k\right) \circ \pi,$$

where $\pi : \Sigma_1 \to C_1$ is defined by

$$\pi(C_{\underline{a}_0}, C_{\underline{a}_1}, \dots) = (\underline{a}_0, \underline{a}_1, \dots).$$

The pair $(\Sigma_1, \overline{\log\phi})$ is called the induced system and $\overline{\log\phi}$ is called the induced potential. Note that if the potential $\log\phi$ is locally Hölder then $\overline{\log\phi}$ is locally Hölder. This follows from the estimates $V_n(\overline{\log\phi}) \leq \sum_{j=n+1}^{\infty} V_j(\log\phi)$.

Next we define the notion of Gibbs measure.

Definition 2.8. A probability measure μ is called a *Gibbs measure* for the potential $\log\phi$ if there exists two constants M and P such that, for every cylinder $C_{i_0\dots i_{n-1}}$ and every $x \in C_{i_0,\dots i_{n-1}}$,

$$\frac{1}{M} \leq \frac{\mu(C_{i_0\dots i_{n-1}})}{\exp(-nP + \sum_{j=0}^{n-1} \log\phi(\sigma^j x))} \leq M.$$

Let us consider a topologically mixing graph G with transition matrix $T = (t_{ij})$. The following definition appears in [**19**].

Definition 2.9. We say that the graph G (or the system Σ) satisfies the big images and preimages (BIP) property if there exist $\{b_1, b_2 \dots b_n\}$ in the alphabet S such that

$$\text{for all } a \in S \text{ there exists } i,\ j \text{ such that } t_{b_i a} t_{a b_j} = 1. \tag{2}$$

Sarig [**19**] proved that a locally Hölder potential $\log\phi$ of finite Gurevich pressure has an invariant Gibbs measure if and only if the system has the BIP property. Moreover, if Σ satisfies the BIP property and $P_{\mathrm{G}}(\log\phi) < \infty$ then the function $t \to P_{\mathrm{G}}(t\log\phi)$ is real analytic for $t > 1$. Note that the induced system satisfies the BIP property.

Let d be a metric on Σ.

https://doi.org/10.1017/S0143385705000350 Published online by Cambridge University Press

Definition 2.10. A positive, locally Hölder potential $\log \psi : \Sigma \to \mathbb{R}$ with the property that there exists a constant $C > 0$ such that

$$\frac{1}{C}\prod_{j=0}^{n-1}(\psi(\sigma^j x))^{-1} \leq \operatorname{diam}(C_{i_0\dots i_{n-1}}) \leq C\prod_{j=0}^{n-1}(\psi(\sigma^j x))^{-1}, \tag{3}$$

for every $x = (i_0, i_1, \dots) \in C_{i_0,\dots i_{n-1}}$, will be called a *metric potential*.

We now describe part of the theory of sequences of Fenchel pairs developed by Wijsman in [**22, 23**].

Definition 2.11. Let h be a convex function; we say that (h, g) form a *Fenchel pair* if

$$g(p) = \sup_x\{px - h(x)\}.$$

Remark 2.2. If h is a convex C^2 function then the function g is also called a Legendre transform. In this case,

$$g(\alpha) = h(q) + q\alpha,$$

where $\alpha(q) = -h'(q)$.

Let h be a convex function, $\rho > 0$ and

$${}_\rho h(x) := \inf\{h(y) : |x - y| < \rho\}.$$

Definition 2.12. (Wijsman [**22, 23**]) Let $\{h_n\}$ be a sequence of convex functions. We say that $\{h_n\}$ converges *infimally* to the function h if

$$\lim_{\rho\to 0}\liminf_{n\to\infty}{}_\rho h_n(x) = \lim_{\rho\to 0}\limsup_{n\to\infty}{}_\rho h_n(x) = h(x).$$

We will write in this case $h_n \xrightarrow{\inf} h$.

Remark 2.3. (Wijsman [**23**]) There are examples where this notion of convergence does not coincide with the pointwise convergence. In general, we have that if $\{h_n\}$ converges infimally to h and $\{h_n\}$ converges pointwise to h_p then $h(x) \leq h_p(x)$.

THEOREM 2.1. (Wijsman [**23**]) *Let (h_n, g_n) and (h, g) be Fenchel pairs. Then*

$$h_n \xrightarrow{\inf} h \text{ if and only if } g_n \xrightarrow{\inf} g.$$

3. *A dimension formula for the Gurevich pressure*

In this section we prove a Bowen-type formula (see [**2**]) for the Gurevich pressure. In the setting of countable Markov shifts this formula relates the pressure function with the Hausdorff dimension of the recurrent set.

Definition 3.1. A point $x \in \Sigma$ is said to be *Recurrent* if there exists a sequence of non-equal natural numbers n_i, such that

$$\lim_{i\to\infty}\sigma^{n_i}x = x.$$

Denote by $\mathcal{R}$ be the set of recurrent points.

https://doi.org/10.1017/S0143385705000350 Published online by Cambridge University Press




FIGURE 1. Possible behaviour of $t \to P(-t \log \psi)$.

Let (Σ, σ) be a topologically mixing Markov chain and $\log \psi$ a locally Hölder metric potential (see Definition 2.10).

THEOREM 3.1.

$$\dim_{\mathrm{H}}(\mathcal{R}) = \inf\{t : P_{\mathrm{G}}(-t \log \psi) \leq 0\}.$$

In the finite-state case, Ruelle [**13**] proved that the Bowen equation

$$P(-t \log \psi) = 0$$

always has a root and that it is equal to the Hausdorff dimension of the repeller. In [**7**] Mauldin and Urbanski generalized this result to the case of infinite iterated function systems. They first noticed that the Bowen equation might not have a root (see Figure 1) and that conservative conformal measures might not exist. We consider a broader class of systems and prove that the pressure function can only account for the recurrent set. We give an example of a piecewise linear map of the interval such that the Hausdorff dimension of the recurrent set is strictly less than the Hausdorff dimension of the repeller (see Example 3.2).

Proof of Theorem 3.1. Let $d := \inf\{t : P_{\mathrm{G}}(-t \log \psi) \leq 0\}$. We begin by proving that $\dim_{\mathrm{H}}(\mathcal{R}) \geq d$. Let $K \subset \Sigma$ be a topologically mixing finite Markov shift. There exists a unique $t_K > 0$ such that $P_K(-t_K \log \psi) = 0$ and $\dim_{\mathrm{H}}(K \cap \mathcal{R}) = t_K$ (see [**9**]). By the approximation property of the Gurevich pressure (1) we have that $t_K \leq d$. Moreover,

$$\sup\{t_K : K \in \mathcal{K}\} = d. \tag{4}$$

In fact, assume by way of a contradiction that there exists $\epsilon > 0$ such that

$$\sup\{t_K : K \in \mathcal{K}\} = d - \epsilon,$$

then

$$\sup\{P_K((d - \epsilon/2) \log \psi) : K \in \mathcal{K}\} \leq 0 < P_{\mathrm{G}}((d - \epsilon/2) \log \psi),$$

which contradicts (1). From the Bowen equation for finite shifts and equation (4) we obtain that

$$\dim_{\mathrm{H}}(\mathcal{R}) \geq d.$$

https://doi.org/10.1017/S0143385705000350 Published online by Cambridge University Press

In order to prove the other inequality we will consider the induced system. Denote

$$\overline{\psi_t}(x) := \prod_{k=0}^{\varphi_1(x)-1} (\psi \circ \sigma^k)^t \circ \pi(x).$$

Note that the diameter of a cylinder of length n on the induced system is such that

$$\operatorname{diam}(C_{i_0 \dots i_{n-1}}) \leq C\overline{\psi_1}(x), \tag{5}$$

where $x \in C_{i_0 \dots i_{n-1}}$. Let $t > d$. Associated to the induced potential $\overline{-t \log \psi}$, there exists a Gibbs measure m (the induced system satisfies the BIP property). Thus, there exists $M > 0$ such that

$$\frac{1}{M} \leq \frac{m(C_{i_0 \dots i_{n-1}})}{\exp(-nP + \log \overline{\psi_t}(x))} \leq M,$$

where $P = P_{\mathrm{G}}(\overline{-t \log \psi})$. Since $P \leq 0$ we have

$$\frac{1}{M}\overline{\psi_t}(x) \leq m(C_{i_0 \dots i_{n-1}}). \tag{6}$$

Combining (5) and (6) we obtain

$$\operatorname{diam}(C_{i_0 \dots i_{n-1}})^t \leq M_1 m(C_{i_0 \dots i_{n-1}}).$$

Denote by $C(n)$ the set of cylinders belonging to the induced system of length n. We have

$$\sum_{C \in C(n)} (\operatorname{diam}(C))^t \leq \sum_{C \in C(n)} M_1 m(C) \leq M_1.$$

Note that $C(n)$ is a cover and the diameters of the cylinders tend to zero as n tends to infinity. Therefore, the t-Hausdorff measure of $\mathcal{R}$ is finite. Since $t > d$ was arbitrary, this completes the proof. □

Remark 3.1. Since the system is topologically mixing the result does not depend on the symbol where we induce.

PROPOSITION 3.1. *If* Σ *has finite Gurevich entropy and* $\dim_{\mathrm{H}}(\Sigma) < \infty$*, then there exists* $d > 0$ *such that* $P_{\mathrm{G}}(-d \log \psi) = 0$*. Moreover,*

(1) *if the potential* $-d \log \psi$ *is recurrent then there exists a conservative* d*-conformal measure;*
(2) *if the potential* $-d \log \psi$ *is transient then there is no conservative* d*-conformal measure.*

Proof. Since $P_{\mathrm{G}}(0) < \infty$ the existence of d is a consequence of the continuity of the pressure. The (non) existence of a conservative d-conformal measure follows from the generalization of the Ruelle–Perron–Frobenius theorem (see §2). □

PROPOSITION 3.2. *Assume* Σ *to have infinite Gurevich entropy and* $\dim_{\mathrm{H}}(\Sigma) < \infty$.

(1) *If there exists* $d > 0$ *such that* $P_{\mathrm{G}}(-d \log \psi) = 0$ *then:*
   (a) *if the potential* $-d \log \psi$ *is recurrent then there exists a conservative* d*-conformal measure;*

https://doi.org/10.1017/S0143385705000350 Published online by Cambridge University Press

(b) *if the potential* $-d\log\psi$ *is transient then there is no conservative* d*-conformal measure.*

(2) *If* $P_{\mathrm{G}}(-d\log\psi) < 0$ *then there is no conservative* d*-conformal measure.*

Proof. It follows from the Ruelle–Perron–Frobenius theorem for countable Markov shifts (see §2). □

Remark 3.2. Assume that Σ satisfies the BIP property. If there exists $d > 0$ such that $P_{\mathrm{G}}(-d\log\psi) = 0$, then $-d\log\psi$ is recurrent; therefore, there exists a d-conformal measure. Mauldin and Urbanski [**7**] called such a system *regular*. If there is no such d then there is no conservative d-conformal measure. In the terminology of Mauldin and Urbanski, these systems are called *irregular*.

In what follows we give examples that illustrate Theorem 3.1. We give them in the context of one-dimensional dynamics, but they can be translated to the countable Markov shift setting.

Example 3.1. (No root for the Bowen equation) Let $k' \in \mathbb{N}$ be such that

$$\sum_{k=k'}^{\infty} \frac{1}{k(\log(2k))^2} < 1.$$

Let $\{I_k : k \geq k'\}$ be a sequence of closed intervals contained in $[0, 1]$ with disjoint interiors. Assume that $|I_k| = 1/k(\log(2k))^2$, where $|\cdot|$ denotes the length of the interval. Define the map $F : \bigcup_{k=k'}^{\infty} I_k \to [0, 1]$ such that $F|I_k$ is linear, onto $[0, 1]$ and of positive slope for every $k \geq k'$. The pressure function is given by

$$P_{\mathrm{G}}(-t\log|F'|) = \log \sum_{k=k'}^{\infty} 1/(k^t(\log 2k)^{2t}).$$

If $t < 1$ then $P_{\mathrm{G}}(-t\log|F'|) = \infty$ and if $t \geq 1$ then $P_{\mathrm{G}}(-t\log|F'|) < 0$. Therefore, the Bowen equation has no root and

$$\dim_{\mathrm{H}}(\mathcal{R}) = 1 = \inf\{t : P_{\mathrm{G}}(-t\log|F'|) \leq 0\}.$$

This potential was first used by Sarig [**16**] for another purpose. Mauldin and Urbanski [**7**] produced other examples where the Bowen equation has no root.

Example 3.2. (Infinite number of roots for the Bowen equation) The Manneville–Pomeau transformation $F : [0, 1] \to [0, 1]$ is defined by

$$F(x) := x + x^{1+\alpha},$$

where $\alpha \in (0, 1)$. This map has two branches defined on the intervals, $[0, a], [a, 1]$. The induced map on $[a, 1]$ satisfies the assumptions of Theorem 3.1. The pressure function $P_{\mathrm{G}}(-t\log|F'|)$ is positive for $t < 1$ and equal to zero for $t \geq 1$ (see [**12**]). Therefore,

$$\dim_{\mathrm{H}}(\mathcal{R}) = 1 = \inf\{t : P_{\mathrm{G}}(-t\log|F'|) \leq 0\}.$$

https://doi.org/10.1017/S0143385705000350 Published online by Cambridge University Press

Example 3.3. (Small recurrent set) We construct a topologically mixing interval map such that the Hausdorff dimension of the recurrent set is strictly less than the Hausdorff dimension of the repeller. Recall that the repeller of a map F is defined by

$$J := \{x \in [0,1] : F^n(x) \text{ is well defined for all } n \in \mathbb{N}\}.$$

Let $r = \frac{1}{2}$ and $I_1 = [0, r]$. If $n > 1$ and $n \in \mathbb{N} \setminus \{2^{i+1} : i \in \mathbb{N}\}$, let $I_n := [\sum_{i=1}^{n-1} r^i, \sum_{i=1}^{n} r^i]$. For each positive integer $n > 1$ there exists a unique interval in $[0,1] \setminus \biguplus I_i$ of length r^{2^n}. Divide this interval into two intervals,

$$I_{a_n} = \left[\sum_{i=1}^{2^n-1} r^i, a_n + \sum_{i=1}^{2^n-1} r^i\right] \quad \text{and} \quad I_{2^n} = \left[a_n + \sum_{i=1}^{2^n-1} r^i, \sum_{i=1}^{2^n} r^i\right].$$

In this way we obtain a partition of the unit interval $[0, 1]$. Let F be a piecewise linear map defined by

$$F(I_n) = \begin{cases} I_1 & \text{if } n \in \{a_{i+1} : i \in \mathbb{N}\}, \\ I_n \cup I_{a_n} \cup I_{n+1} & \text{if } n \in \{2^{i+1} - 1 : i \in \mathbb{N}\}, \\ I_n \cup I_{n+1} & \text{in any other case.} \end{cases} \tag{7}$$

Let $0 < c < 1$. Consider a strictly increasing sequence (s_i) such that $s_i \in (0, c)$ and $c = \sup\{s_i\}$. Denote by F_1 the restriction of F to the set $I_{a_2} \bigcup \left(\bigcup_{j=1}^{4} I_j\right)$. Note that if $a_2 = 0$ then the Hausdorff dimension of the recurrent set for F_1 is zero. If $a_2 = r^4$ the Hausdorff dimension of the recurrent set for F_1 is one. Choose a_2 such that the Hausdorff dimension of the recurrent set is s_1, that is $P(-s_1 \log |F_1'|) = 0$.

We define a_n inductively. Suppose that we have already defined $\{a_2 \dots a_{n-1}\}$. Denote by F_n the restriction of F to $\left(\bigcup_{j=2}^{n} I_{a_j}\right) \bigcup \left(\bigcup_{j=1}^{2^n} I_j\right)$. Note that if $a_n = 0$ then the Hausdorff dimension of the recurrent set of F_n is s_{n-1}, and if $a_n = r^{2^n}$ then the Hausdorff dimension of the recurrent set of F_n is one. Choose a_n such that

$$P(-s_n \log |F_n'|) = 0.$$

Note that $c = \inf\{t : P_{\mathrm{G}}(-t \log |F'|) \leq 0\}$. Therefore, $\dim_{\mathrm{H}}(\mathcal{R}) = c < 1 = \dim_{\mathrm{H}}(J)$ as required.

4. *Multifractal analysis, the BIP case*

The purpose of this section is to develop techniques that enable us to carry out the multifractal analysis for countable Markov shifts. Recall that Pesin and Weiss [**10**] studied the case in which the space is compact (the finite-state case) and μ is an equilibrium measure for a Hölder potential. They proved that the multifractal spectrum is concave and real analytic. Schmeling [**20**] proved that it has bounded domain. Note that the domain is defined by $\{\alpha \in \mathbb{R} : J_\alpha \neq \emptyset\}$. As mentioned in the introduction, the multifractal analysis for countable Markov shifts has also been studied in [**5**], [**8**] and [**11**]. Our techniques are slightly different. The lower bounds for the Hausdorff dimension of the level sets are obtained by means of an approximation argument that relies on some tools of convex analysis (the infimal convergence). This approach allows us to develop the multifractal analysis in more general symbolic spaces, gives a simple explanation to the differences

https://doi.org/10.1017/S0143385705000350 Published online by Cambridge University Press

between the finite- and the countable-state cases and enables us to give sufficient conditions in order for the multifractal spectrum to have unbounded domain and points of non-analyticity. The main ingredients that will be used are:

- the infimal convergence of Wijsman;
- the multifractal analysis over compact spaces;
- the approximation property of the Gurevich pressure (1).

In this section we will consider a countable Markov shift that satisfies the BIP property and a metric potential $\log \psi$ (see Definition 2.10). Throughout the rest of the paper we will work on the recurrent set of Σ. Let $\log \phi : \Sigma \to \mathbb{R}$ be a locally Hölder potential of zero pressure, not cohomologous to $\log \psi$. Denote by μ the corresponding Gibbs measure.

Remark 4.1. The metric that we are considering is such that

$$d_\mu(x) := \lim_{r\to 0} \frac{\log \mu(B(x,r))}{\log r} = \lim_{n\to\infty} \frac{\log \mu(C_{i_0\dots i_{n-1}}(x))}{\log \prod_{i=0}^{n-1} \big(\psi(\sigma^i x)\big)^{-1}}.$$

Definition 4.1.

$$q^* := \inf\{q : \text{there exists } t \in \mathbb{R} \text{ such that } P_{\mathrm{G}}(-t \log \psi + q \log \phi) \le 0\}.$$

LEMMA 4.1. *If* $\dim_{\mathrm{H}}(\Sigma) < \infty$ *then either* $q^* = 0$ *or* $q^* = -\infty$.

Proof. Note that $P_{\mathrm{G}}(-\dim_{\mathrm{H}}(\Sigma) \log \psi) \le 0$, therefore $q^* \le 0$.

Assume by way of a contradiction that $q^* \in (-\infty, 0)$. Let $0 < \epsilon, \epsilon' < |q^*|/2$. Since $q^* + \epsilon > q^*$ and $q^* + \epsilon' > q^*$, there exist $T(\epsilon)$ and $T(\epsilon')$ such that

$$P_{\mathrm{G}}(-T(\epsilon) \log \psi + (q^* + \epsilon) \log \phi) \le 0 \quad \text{and} \quad P_{\mathrm{G}}(-T(\epsilon') \log \psi + (q^* + \epsilon') \log \phi) \le 0.$$

Since $P_{\mathrm{G}}(f + g) \le P_{\mathrm{G}}(f) + P_{\mathrm{G}}(g)$ then

$$P_{\mathrm{G}}(-(T(\epsilon) + T(\epsilon')) \log \psi + (2q^* + \epsilon + \epsilon') \log \phi) \le 0.$$

Therefore, $\epsilon + \epsilon' > |q^*|$. This contradiction proves the statement. □

Remark 4.2. If $\dim_{\mathrm{H}}(\Sigma) = \infty$ it is possible for q^* to be larger than zero. See Example 4.3.

Definition 4.2. The function $T(q)$ is implicitly defined by

$$T(q) := \begin{cases} \inf\{t : P_{\mathrm{G}}(-t \log \psi + q \log \phi) \le 0\} & \text{if } q \ge q^*, \\ \infty & \text{if } q < q^*. \end{cases} \tag{8}$$

We prove the following.

THEOREM 4.1. *The multifractal spectrum* f_μ *is the Fenchel transform of* T.

Equivalent definitions of $T(q)$ will be required to prove the theorem. Let $K \in \mathcal{K}$ and let $P_K(\cdot)$ denote the topological pressure restricted to the set K.

PROPOSITION 4.1.

$$T(q) = \sup\{t : P_K(-t \log \psi + q \log \phi) = 0,\ K \in \mathcal{K}\}.$$

https://doi.org/10.1017/S0143385705000350 Published online by Cambridge University Press

Proof. The result is clear for $q < q^*$. Let $q \geq q^*$. For every compact invariant set K and for every $q \in \mathbb{R}$ there exists $T_K(q)$ such that (see [**10**])

$$P_K(-T_K(q)\log\psi + q\log\phi) = 0.$$

Since $K \subset \Sigma$, by (1) we have $P_K(-t\log\psi + q\log\phi) \leq P_{\mathrm{G}}(-t\log\psi + q\log\phi)$ and therefore $T_K(q) \leq T(q)$. Note that the pressure is monotone, that is, if $K_1 \subset K_2$ then

$$P_{K_1}(-t\log\psi + q\log\phi) \leq P_{K_2}(-t\log\psi + q\log\phi).$$

In particular, $T_{K_1}(q) \leq T_{K_2}(q)$. Assume by way of a contradiction that

$$A := \sup\{T_K(q) : K \in \mathcal{K}\} < T(q). \tag{9}$$

Let $B \in (A, T(q))$. Since the pressure function is decreasing in t, we have that

$$P_{\mathrm{G}}(-B\log\psi + q\log\phi) > 0.$$

On the other hand, because of the approximation property (1) and the assumption (9),

$$P_{\mathrm{G}}(-B\log\psi + q\log\phi) \leq 0.$$

This contradiction proves the proposition. □

PROPOSITION 4.2.

$$T(q) = \sup\{t : P_K(-t\log\psi + q(\log\phi - P_K(\log\phi))) = 0, K \in \mathcal{K}\}.$$

Proof. Again the result is clear for $q < q^*$. Let $K \in \mathcal{K}$ and $q \geq q^*$. Define $T_K^*(q)$ implicitly by

$$P_K(-T_K^*(q)\log\psi + q(\log\phi - P_K(\log\phi))) = 0. \tag{10}$$

Note that

$$|P_K(-t\log\psi + q\log\phi - qP_K(\log\phi)) - P_K(-t\log\psi + q\log\phi)| \leq |qP_K(\log\phi)|. \tag{11}$$

Since $P_{\mathrm{G}}(\log\phi) = 0$ and because of (1) we have that

$$\sup\{qP_K(\log\phi) : K \in \mathcal{K}\} = 0. \tag{12}$$

The result follows from (11), (12), (1) and Proposition 4.1. □

PROPOSITION 4.3. *The function $T(q)$ is convex and decreasing.*

Proof. Let $K \in \mathcal{K}$. Pesin and Weiss [**10**], using the fact that the pressure is analytic and considering the formulae for its derivatives, showed that the function $T_K^*(q)$ is analytic, decreasing and convex. Since

$$T(q) = \sup\{T_K^*(q) : K \subset \Sigma \text{ compact invariant}\}$$

the desired properties follow. □

https://doi.org/10.1017/S0143385705000350 Published online by Cambridge University Press

The next proposition has been proved in different settings, in particular see [**10**]. It relates the pointwise dimension to the Birkhoff sums.

PROPOSITION 4.4. *If $x \in \Sigma$ then*

$$d_\mu(x) = \lim_{n\to\infty} \frac{\log \mu(C_{i_0\dots i_{n-1}}(x))}{\log \prod_{i=0}^{n-1} (\psi(\sigma^i x))^{-1}} = \lim_{n\to\infty} \frac{\sum_{i=0}^{n-1} \log \phi(\sigma^i(x))}{\sum_{i=0}^{n-1} \log(\psi(\sigma^i x))^{-1}},$$

whenever one of the limits exists.

Proof. The proof follows from the Gibbs property of the measure μ and from Remark 4.1. □

Let $q > q^*$. Associated to the potential $\phi_q := \phi_{T(q),q} = -T(q)\log\psi + q\log\phi$ there exists a (normalized) Gibbs measure μ_q (see [**19**]).

Definition 4.3. Let $q > q^*$ and consider

$$\alpha(q) := -\frac{\int \log\phi \, d\mu_q}{\int \log\psi \, d\mu_q}.$$

Note that, since the system satisfies the BIP property, for $q > q^*$ the potential ϕ_q is strongly positive recurrent (see [**16**]). In particular, the function $P_{\mathrm{G}}(-t\log\psi + q\log\phi)$ is real analytic in each variable (when finite). Moreover, $\log\psi \in L^1_{\mu_q}$ and $\log\phi \in L^1_{\mu_q}$. Using the formula for the derivative of the pressure (see [**9**]) we have $-T'(q) = \alpha(q)$.

Remark 4.3. Note that μ_q-almost every $x \in \Sigma$ is such that

$$d_\mu(x) = \lim_{n\to\infty} -\frac{\sum_{i=0}^{n-1} \log\phi(\sigma^i(x))}{\sum_{i=0}^{n-1} \log\psi(\sigma^i(x))} = -\frac{\int \log\phi \, d\mu_q}{\int \log\psi \, d\mu_q} =: \alpha(q).$$

Let (K_n) be a sequence in $\mathcal{K}$, such that $T^*_{K_n}(q) := T^*_n(q)$ converges pointwise to $T(q)$. We prove that $T^*_n(q)$ converges infimally to $T(q)$.

LEMMA 4.2. $T^*_n(q) \xrightarrow{\inf} T(q)$.

Proof. Fix a positive integer n. The function $T^*_n(q)$ is decreasing in q. Thus,

$${}_\rho T^*_n(x) := \inf\{T^*_n(y) : |x-y| < \rho\} = T^*_n(x+\rho).$$

Since $T^*_n(q)$ converges pointwise to $T(q)$, we have

$$\lim_{n\to\infty} {}_\rho T^*_n(x) = T(x+\rho).$$

If $q > q^*$ then $T(q)$ is continuous and convex, so $\lim_{\rho\to 0} T(q+\rho) = T(q)$. Assume $q^* = 0$. If $q < 0$ then for ρ small enough we have $T(q+\rho) = \infty$, so $\lim_{\rho\to 0} T(q+\rho) = T(q)$. If $q = 0$ then the result follows because $T(0) = \lim_{q\to 0^+} T(q)$. Therefore,

$$T^*_n(q) \xrightarrow{\inf} T(q). \qquad \square$$

Proof of Theorem 4.1. Denote by L the Fenchel transform of T. We first prove that if $q > q^*$ then $\dim_{\mathrm{H}}(J_{\alpha(q)}) = T(q) + q\alpha(q)$. Recall that $\mu_q(J_{\alpha(q)}) = 1$ and that μ_q is a

https://doi.org/10.1017/S0143385705000350 Published online by Cambridge University Press

Gibbs measure. The arguments developed by Pesin and Weiss [**10**, Lemma 2] apply and yield

$$d_{\mu_q}(x) \leq T(q) + q\alpha(q)$$

for every $x \in J_{\alpha(q)}$. Therefore, $\dim_{\mathrm{H}}(J_{\alpha(q)}) \leq T(q) + q\alpha(q)$. Note that in order for the argument to work we just need $P_{\mathrm{G}}(\phi_q) \leq 0$. In particular, we have proved that $f_\mu \leq L$.

Let $f_{\mu,n}(\alpha)$ be the Fenchel transform of $T_n^*(q)$. We have already proved that $T_n^*(q) \xrightarrow{\inf} T(q)$. By Wijsman's theorem (see Theorem 2.1) we have

$$f_{\mu,n} \xrightarrow{\inf} L.$$

Note that $f_{\mu,n}$ is an increasing sequence and $f_{\mu,n}(\alpha) \leq \alpha$. Thus, there exists a function f_p which corresponds to the pointwise limit of $f_{\mu,n}$. By Wijsman's remark (see Remark 2.3) we have $L \leq f_p$. But by definition we have that $f_p \leq f_\mu$, that is

$$L \leq f_p \leq f_\mu,$$

therefore $L = f_\mu$.

If $q^* = 0$ and $\lim_{q\to 0^+} -T'(q) = \infty$, the result follows from the previous argument. If $q^* = 0$ and $\lim_{q\to 0^+} -T'(q) < \infty$, then for every $\alpha > \lim_{q\to 0^+} -T'(q)$ we have $f_\mu(\alpha) = \dim_{\mathrm{H}} \mathcal{R}$. This concludes the proof. □

Remark 4.4. It is possible to prove that $f_p = L$ just using the definition of infimal convergence and the fact that the sequence of concave functions $(f_{\mu,n})$ is monotonic.

In the next two theorems we prove that the behaviour of the multifractal spectrum for countable Markov shifts can be different from the finite-state case (see Figures 2 and 3).

PROPOSITION 4.5. *If $q^* = 0$, then the multifractal spectrum f_μ has unbounded domain.*

Proof. If $q^* = 0$, then $T(q) = \infty$ for every $q < 0$. Recall that

$$\lim_{n\to\infty} T_n(q) = T(q)$$

and that the function $T_n(\cdot)$ is an analytic function for every $n \in \mathbb{N}$.

Recall that, in this setting, when finite the pressure is analytic. Therefore $T(q)$ is analytic on (q^*, ∞). Assume first that the Hausdorff dimension of the system is finite, that is $T(0) < \infty$. Assume as well that the function $T(q)$ has bounded derivative for $q > q^*$, that is

$$-T'(0) = \lim_{q\to 0^+} -T'(q) < \infty.$$

The mean value theorem implies that for every $\alpha > -T'(0)$ there exist $n \in \mathbb{N}$ and $q < 0$ such that $-T_n'(q) = \alpha$. That is $f_{\mu,n}(\alpha) > 0$ and therefore $f_\mu(\alpha) > 0$.

Assume now that

$$-T'(0) = \lim_{q\to 0^+} -T'(q) = \infty.$$

Then, for every $\alpha > -T'(1)$, there exists $q \in (0, 1)$ such that $-T'(q) = \alpha$. Therefore $f_\mu(\alpha) > 0$, which concludes the proof. □

https://doi.org/10.1017/S0143385705000350 Published online by Cambridge University Press




FIGURE 2. Possible behaviour of the function $T(q)$.

https://doi.org/10.1017/S0143385705000350 Published online by Cambridge University Press




FIGURE 3. Possible behaviour of the multifractal spectrum.

Remark 4.5. Note that when considering compact spaces the domain is always a bounded interval (see [**20**]).

Remark 4.6. Note that Proposition 4.5 follows from the Fenchel relation between $T(q)$ and f_μ.

If the multifractal spectrum has unbounded domain then two possible behaviours can occur, as the following proposition shows (see Figures 2 and 3).

https://doi.org/10.1017/S0143385705000350 Published online by Cambridge University Press

PROPOSITION 4.6. *Assume* $q^* = 0$.

(1) *If* $\lim_{q\to 0^+} -T'(q) < \infty$ *then there exists* $\alpha(0)$ *such that for every* $\alpha > \alpha(0)$ *we have* $f_\mu(\alpha) = \dim_{\mathrm{H}}(\mathcal{R})$.

(2) *If* $\lim_{q\to 0^+} -T'(q) = \infty$ *then the multifractal spectrum is strictly increasing.*

Proof. First assume $T(0) < \infty$ and that

$$\alpha(0) = -T'(0) = \lim_{q\to 0^+} -T'(q) < \infty.$$

Since $q^* = 0$, for every $q < 0$ we have $T(q) = \infty$. In particular, the multifractal spectrum has unbounded domain. Recall that

$$f_\mu(\alpha) = \inf_{-\infty\le q\le\infty}\{T(q) + \alpha q\} = \inf_{0\le q\le\infty}\{T(q) + \alpha q\}.$$

Let $\alpha > -T'(0)$. The function $T(q)$ is convex and $-\alpha q + T(0)$ is a support line of $T(q)$. Hence, for every $q > 0$ we have

$$T(q) > -\alpha q + T(0).$$

In particular,

$$T(q) + \alpha q > T(0).$$

Therefore,

$$f_\mu(\alpha) = \inf_{0\le q\le\infty}\{T(q) + \alpha q\} = T(0) = \dim_{\mathrm{H}}(\mathcal{R}).$$

Assume that

$$-T'(0) = \lim_{q\to 0^+} -T'(q) = \infty.$$

Then,

$$f_\mu(\alpha) = \inf_{-\infty\le q\le\infty}\{T(q) + \alpha q\} = \inf_{0\le q\le\infty}\{T(q) + \alpha q\} = T(q') - T'(q')q',$$

where $q' > 0$ is such that $-T'(q') = \alpha$. □

Remark 4.7. Note that in case (1) of the previous proposition the multifractal spectrum has a point of non-analyticity. Recall that in the finite-state case the multifractal spectrum is analytic (see [**10**]).

Remark 4.8. Note that in order for $\alpha(0)$ to be infinity we need $-\int \log\phi\, d\mu_0 = \infty$. Recall that μ_0 is a Gibbs measure for $-\dim_H(\mathcal{R})\log\psi$ and that it does not depend on $\log\phi$. Therefore, the integrability of $\log\phi$ depends on the relation between $\psi^{-\dim_H(\mathcal{R})}$ and $\log\phi$.

In the next proposition we provide sufficient conditions in order for the multifractal spectrum to have bounded or unbounded domain.

PROPOSITION 4.7. *Assume* $\dim_{\mathrm{H}}(\Sigma) < \infty$.

(1) *If, for every* $x \in \Sigma$,

$$(\phi(x))^{-1} \le \psi(x),$$

then the multifractal spectrum of μ *has bounded domain.*

https://doi.org/10.1017/S0143385705000350 Published online by Cambridge University Press

(2) *If, for every* $q < 0$, $t > 0$ *and* $x \in \Sigma$,

$$(\phi(x))^q > (\psi(x))^t,$$

then the multifractal spectrum of μ *has unbounded domain. Moreover:*

(a) *if* $\lim_{q\to 0^+} -T'(q) < \infty$ *then there exists a point* $\alpha(0)$ *in the domain of* f_μ *such that, for* $\alpha < \alpha(0)$, *the multifractal spectrum is strictly increasing and, for* $\alpha > \alpha(0)$, $f_\mu(\alpha) = \dim_{\mathrm{H}}(\mathcal{R})$*;*

(b) *if* $\lim_{q\to 0^+} -T'(q) = \infty$ *then the multifractal spectrum is strictly increasing.*

Proof. Note that if $q^* = -\infty$ then f_μ has bounded domain. If $q < 0$, then from the condition given in part (1) of the theorem we have

$$-t \log \psi + q \log \phi \le -(t+q) \log \psi.$$

Therefore,

$$P_{\mathrm{G}}(-t \log \psi + q \log \phi) \le P_{\mathrm{G}}(-(t+q) \log \psi).$$

If $t > (\dim_{\mathrm{H}}(\mathcal{R}) - q)$ then

$$P_{\mathrm{G}}(-t \log \psi + q \log \phi) \le 0.$$

Thus $q^* = -\infty$.

Note that if $q^* = 0$ then f_μ has unbounded domain. Recall that since the system satisfies the BIP property, then $P_{\mathrm{G}}(0) = \infty$. The condition given in part (2) of the theorem ensures that for every $q < 0$ and for every $t \in \mathbb{R}$ we have

$$-t \log \psi + q \log \phi > 0.$$

Therefore $q^* = 0$. If $\lim_{q\to 0^+} -T'(q) < \infty$ and $q^* = 0$, then f_μ is concave, has unbounded domain and $f_\mu(-T'(0)) = \dim_{\mathrm{H}}(\mathcal{R})$. Note that f_μ is bounded above by $\dim_{\mathrm{H}}(\mathcal{R})$. Therefore, if $\alpha > -T'(0)$, then $f_\mu(\alpha) = \dim_{\mathrm{H}}(\mathcal{R})$.

If $\lim_{q\to 0^+} -T'(q) = \infty$ and $q^* = 0$, then f_μ is concave, has unbounded domain and does not have a maximum. □

Example 4.1. Let Σ be the full shift over the positive integers. Let $\psi|_{C_n} = 2^n$ and let $\phi|_{C_n} = (n!)^{-s}$, where $s \in \mathbb{R}^+$ is such that $P_{\mathrm{G}}(\log \phi) = 0$. Let $q < 0$. Then for fixed t and n large enough,

$$\phi^q\big|_{C_n} = (n!)^{s|q|} > 2^{nt} = \psi|_{C_n}.$$

That is, the multifractal spectrum for the equilibrium measure μ corresponding to the potential $\log \phi$, has unbounded domain.

Example 4.2. Let Σ be the full shift over the positive integers. Let

$$\psi|_{C_n} = 3^n \quad \text{and} \quad \phi|_{C_n} = 2^{-n}.$$

Clearly $\phi^{-1} \le \psi$, therefore the multifractal spectrum for the equilibrium measure μ corresponding to the potential $\log \phi$ has bounded domain.

If $\dim_{\mathrm{H}}(\mathcal{R}) = \infty$ the situation is different as the following example shows.

https://doi.org/10.1017/S0143385705000350 Published online by Cambridge University Press

Example 4.3. Assume Σ satisfies the BIP property and let $\log\psi = \log\lambda$, where $\lambda > 1$. Since the system has infinite Gurevich entropy then $\dim_{\mathrm{H}}(\mathcal{R}) = \infty$. If $\log\phi$ is a potential of zero pressure and μ is the corresponding Gibbs measure, then the multifractal spectrum of μ:

(1) has unbounded domain;
(2) has unbounded image;
(3) is an increasing concave function.
(4) If $P_{\mathrm{G}}(q^*\log\phi) < \infty$ then there exists an associated equilibrium measure μ_{q^*}. We have:
   (a) if $\log\phi \in L^1_{\mu_{q^*}}$ then there exists a point $-T'(q^*)$ such that for $\alpha < -T'(q^*)$ the multifractal spectrum is strictly concave, and for $\alpha > -T'(q^*)$ it is linear with slope q^*;
   (b) if $\log\phi \notin L^1_{\mu_{q^*}}$ then the multifractal spectrum is strictly concave and one of the asymptotes has slope q^*.
(5) If $P_{\mathrm{G}}(q^*\log\phi) = \infty$ then the multifractal spectrum is strictly concave and one of the asymptotes has slope q^*.

Let (Σ, σ) be the full-shift on a countable infinite alphabet. Consider the following potentials that illustrate the possibilities in this example. For the locally constant potential defined by $\log\phi_1|_{C_n} = -\log(n(n+1))$ we have $q^* = 1/2$. For the potential $\log\phi_2(x) = -\log(x_0(\log 2x_0)^2)^r$, where $r > 1$ is such that $P_{\mathrm{G}}(\log\phi_2) = 0$, the multifractal spectrum of the corresponding measure is strictly concave for $\alpha < -T'(1)$ and for $\alpha > -T'(1)$ it is linear with slope one. The potential $\log\phi_3|_{C_n} = \log n^{-nr}$, where r is a positive number such that $P_{\mathrm{G}}(\log\phi_3) = 0$, is such that the multifractal spectrum of the corresponding measure is strictly concave.

Remark 4.9. Note that $\dim_{\mathrm{H}}(\mathcal{R}) = \sup\{\dim_{\mathrm{H}}(K) : K \in \mathcal{K}\}$. Let $J'_K := \{x \in K :$ the limit $d_\mu(x)$ does not exist$\}$ be the irregular set of K. Barreira and Schmeling proved in [**1**] that the set J'_K has full Hausdorff dimension. Therefore,

$$\dim_{\mathrm{H}}(\mathcal{R}) = \sup\{\dim_{\mathrm{H}}(J'_K) : K \in \mathcal{K}\}.$$

That is, the irregular set J' has the same Hausdorff dimension as the recurrent set.

5. *Multifractal analysis, the non-BIP case*

This section will be devoted to the case in which the graph G does not satisfy the BIP property. Therefore, locally Hölder potentials do not have corresponding Gibbs measures. As in the previous section, we will assume the potential $\log\phi$ to be locally Hölder continuous of zero pressure not cohomologous to the metric potential $\log\psi$. We also assume that it is positive recurrent. Our approach is based on the study of the induced system Σ_1 (see §2). Note that the induced system satisfies the BIP property. Therefore, the results of the previous sections (with some modifications) can be translated to this setting.

Definition 5.1. Let $T_i(q)$ be defined implicitly by

$$P_{\mathrm{G}}(-T_i(q)\,\overline{\log\psi} + q\,\overline{\log\phi}) = 0, \tag{13}$$

provided such a number exists.

https://doi.org/10.1017/S0143385705000350 Published online by Cambridge University Press

Denote $\overline{\phi_q} := -T_i(q)\,\overline{\log\psi} + q\,\overline{\log\phi}$ and denote by μ_q the associated Gibbs measure (recall that this potential is defined on the induced system).

Remark 5.1. Three conditions will be required on the potentials. Assume that there exists an interval I of values of q for which:

(1) the function $T_i(q)$ is well defined (equation (13) has a root);
(2) the potential $-T_i(q)\,\log\psi + q\,\log\phi$ on the original system is positive recurrent;
(3) the potentials $\overline{\log\psi}$, $\overline{\log\phi}$ are integrable with respect to the Gibbs measure associated to $-T_i(q)\,\overline{\log\psi} + q\,\overline{\log\phi}$.

Proposition 4.1 is still valid. So the function $T_i(q)$ is convex and decreasing. Denote by $f_{\mu,i}$ the Fenchel transform of $T_i(q)$. As proved in the previous section, this is the multifractal spectrum of the Gibbs measure μ_i associated to $\overline{\log\phi}$.

Remark 5.2. The potential $\overline{\phi_q}$ has zero pressure and is positive recurrent. This implies that the potential on the original system, $-T_i(q)\,\log\psi + q\,\log\phi$, has zero pressure and is recurrent (see [**16**]). Therefore, if $q \in I$ then the function $T_i(q)$ defined on the induced system coincides with the function $T(q)$ defined in the original system.

Denote the Fenchel transform of $T(q)$ by f_μ and the level sets of the induced system by $J_{\alpha,i}$.

THEOREM 5.1. *If $q \in I$ then* $\dim_{\mathrm{H}}(J_{\alpha(q)}) = T(q) + q(-T'(q))$.

Proof. The lower bound $\dim_{\mathrm{H}}(J_{\alpha(q)}) \geq T(q)+q(-T'(q))$ can be obtained using the same arguments (infimal convergence) as in the proof of Theorem 4.1.

In order to obtain the upper bound, note that $J_{\alpha(q)} \subset \pi(J_{\alpha(q),i})$. Since we have that $\dim_{\mathrm{H}}(J_{\alpha(q),i}) = T(q) + q(-T'(q))$ and the projection π is a Lipschitz function, we have

$$\dim_{\mathrm{H}}(J_{\alpha(q)}) \leq \dim_{\mathrm{H}}(\pi(J_{\alpha(q),i})) \leq T(q) + q(-T'(q)). \tag{14}$$

□

Remark 5.3. As in the previous section, the domain of the multifractal spectrum f_μ can be unbounded. Moreover, assume $\dim_{\mathrm{H}}(\mathcal{R}) = d < \infty$ and that f_μ has unbounded domain. Since (f_μ, T) form a Fenchel pair, we have two possible behaviours. Either f_μ is strictly increasing and concave, or there exists a point α^* such that for $\alpha < \alpha^*$ the multifractal spectrum is strictly increasing and strictly concave and for $\alpha > \alpha^*$ we have $f_\mu(\alpha) = d$.

In the next theorem we give sufficient conditions in order for the multifractal spectrum to have unbounded domain. Before stating it we need the following definitions. Let $\mathcal{A}_n := \{C_{1i_1\ldots i_n} : i_j \neq 1,\ j \in \{1,\ldots,n\}, t_{i_n,1} = 1\}$, where $1i_1\ldots i_n$ are admissible words. Denote by $\Sigma_{\mathcal{A}_n}$ and $\Sigma_{\bigcup_{i=1}^n \mathcal{A}_i}$ the full-shifts over the alphabet $\mathcal{A}_n$ and $\bigcup_{i=1}^n \mathcal{A}_i$, respectively.

THEOREM 5.2. *Let (b_n) be an increasing sequence of real numbers with $b_n > 1$. Assume that for every $x \in [\varphi_1 = n]$ we have $\overline{\log\psi(x)} < \log b_n$. If, for every $q < 0$,*

$$\lim_{n\to\infty} \frac{P_{\Sigma_{\bigcup_{i=1}^n \mathcal{A}_i}}(q\,\overline{\log\phi})}{\log b_n} = \infty, \tag{15}$$

then the multifractal spectrum of μ has unbounded domain.

https://doi.org/10.1017/S0143385705000350 Published online by Cambridge University Press

Proof. Note that

$$0 = P_{\Sigma_{\bigcup_{i=1}^{n}\mathcal{A}_i}}(-T_n(q)\,\overline{\log\psi} + q\,\overline{\log\phi}) \geq -T_n(q)\,\log b_n + P_{\Sigma_{\bigcup_{i=1}^{n}\mathcal{A}_i}}(q\,\overline{\log\phi}).$$

Therefore,

$$T_n(q) \geq \frac{P_{\Sigma_{\bigcup_{i=1}^{n}\mathcal{A}_i}}(q\,\overline{\log\phi})}{\log b_n}.$$

From equation (15) we deduce that for negative values of q

$$\lim_{n\to\infty} T_n(q) = T(q) = \infty.$$

Recall that for every positive integer n the function $T_n(q)$ is analytic. Applying the mean value theorem, we have that for every $\alpha > \alpha(0)$ there exists n and $q_n < 0$ such that

$$-T_n'(q_n) = \alpha,$$

therefore $\dim_{\mathrm{H}}(J_{(\alpha,i)}) > 0$. This implies that for every $\alpha > \alpha(0)$ the multifractal spectrum $f_{\mu_i}(\alpha) > 0$. Hence it has unbounded domain. □

Remark 5.4. Note that if the system has finite entropy then the sequence (b_n) is always well defined.

Remark 5.5. Let (a_n) be a sequence of positive numbers satisfying

$$\lim_{n\to\infty} \frac{\log a_n}{\log b_n} = -\infty. \tag{16}$$

If there exists $n^* \in \mathbb{N}$ such that for every $n > n^*$ there exists a periodic point $x_n \in [\varphi_1 = n]$ such that

$$\prod_{i=0}^{n-1} \phi(\sigma^i(x_n)) \leq a_n, \tag{17}$$

then, from the variational principle, for every $q < 0$

$$\frac{P_{\Sigma_{\mathcal{A}_n}}(q\,\overline{\log\phi})}{\log b_n} \geq \frac{q\,\log a_n}{\log b_n}.$$

From equation (16) we obtain that $T(q) = \infty$. Hence the multifractal spectrum of μ has unbounded domain.

Remark 5.6. Assume the system to have finite entropy $\log h$. Recall that the induced measure μ_i on C_1 is Gibbs and let $C_{1i_1\ldots i_n} \in \mathcal{A}_n$. If, for every $x \in [\varphi_1 = n]$, we have

$$\frac{P_{\Sigma_{\mathcal{A}_n}}(q\,\overline{\log\phi})}{\log b_n} \geq \frac{q\,\log a_n}{\log b_n},$$

where (a_n), (b_n) are as in Remark 5.5, then there exists a constant C that does not depend on n or on the cylinder, such that

$$\mu_i(C_{1i_1\ldots i_n}) \leq C a_n.$$

https://doi.org/10.1017/S0143385705000350 Published online by Cambridge University Press

Since the system has finite entropy $\log h$, the number of cylinders of length n is bounded above by h^n (see [**6**, p. 213]). Therefore,

$$\mu_i([\varphi_1(x) = n]) \leq Ch^n a_n.$$

This allows us to estimate the tail

$$\mu([\varphi_1 > n]) \leq \mu(C_1)C \sum_{i=n+1}^{\infty} h^i a_i.$$

Example 5.1. The renewal graph G is defined over the positive integers by the transition matrix $T = (t_{ij})$, whose entries t_{11}, t_{1i} and $t_{i,i-1}$ for $i = 2, 3, \ldots$ are one, and the rest of the entries are equal to zero. This is a topologically mixing graph with finite entropy (actually $\log 2$) and is positive recurrent.

The combinatorial structure of the induced system on the cylinder C_1 is simple. For every $n \geq 1$ we have that $\mathcal{A}_n = \{C_{1n(n-1)\ldots 32}\}$. A generalization of this graph arises naturally when studying tower extensions. Let (d_i) be an increasing sequence of positive integers. The transition matrix $T = (t_{ij})$ associated to this kind of graph has entries $t_{11}, t_{i,i-1}$ and t_{1,d_i} equal to one. The rest of the entries are equal to zero. Depending on d_n the graph is transient or recurrent. If it is recurrent, that is

$$\sum_{n\geq 1} h^{-n} \sum_{\sigma^n x = x} 1_{[1]}(x) = \infty,$$

where $P_{\mathrm{G}}(0) = \log h$ is the entropy of the system, then the graph is positive recurrent. In fact, note that

$$\sum_{n\geq 1} \frac{n}{h^n} \sum_{\sigma^n x = x} 1_{[\varphi_1 = n]} \leq \sum_{n\geq 1} \frac{n}{h^n}$$

which converges because $h > 1$ (the system has positive entropy). Hence, there exists a probability measure of maximal entropy.

Consider the case in which $d_n = 2n + 1$. Let $\log \psi = \log \lambda$, where $\lambda > 1$. The potential is defined by

$$\log \phi(x) = \begin{cases} \log n^{-r} & \text{if } x \in C_n \text{ and } n > 1, \\ 1 & \text{if } x \in C_1. \end{cases} \tag{18}$$

The constant r is such that $P_{\mathrm{G}}(\log \phi) = 0$. In this case the multifractal spectrum of the equilibrium measure μ corresponding to $\log \phi$ has unbounded domain. Moreover, since

$$\sum_{n=1}^{\infty} \frac{r \log(2n+1)!}{\lambda^{nT(0)}} < \infty$$

there exists a point $\alpha(0)$ such that for all $\alpha > \alpha(0)$ the multifractal spectrum is constant.

Example 5.2. (Nakaishi) In [**8**] Nakaishi studied the multifractal spectrum of the measure of maximal entropy of maps of the interval with a parabolic fixed point. He proved that it has unbounded domain. His example can be understood in our setting as follows. Let G be the renewal graph and $\log \phi(x) = \log 2^{-n}$ for $x \in C_n$. The metric considered is such that $\log \psi\big|_{C_n} = \log n$. Since

$$\lim_{n\to\infty} \frac{-n \log 2}{\log n} = -\infty, \tag{19}$$

the multifractal spectrum has unbounded domain.

https://doi.org/10.1017/S0143385705000350 Published online by Cambridge University Press




FIGURE 4. The behaviour of the multifractal spectrum of concentrated measures.

The following example can be understood as a model for a piecewise linear geometrical construction.

Example 5.3. Let (λ_i) be an increasing sequence with $\lambda_i > 1$. Assume $\log \psi|_{C_i} = \lambda_i$ and that $\operatorname{diam}(C_{i_0 \dots i_n}) = \prod_{j=0}^{n} \lambda_{i_j}^{-1}$. Let G be a positive recurrent graph of finite Gurevich entropy and let $\log \phi$ be a positive recurrent, locally Hölder continuous potential of zero pressure. Denote by μ the corresponding equilibrium measure. Let

$$M_n := \min\{\operatorname{diam}(C_{i_0 \dots i_n}) : C_{i_0 \dots i_n} \in \mathcal{A}_n\}.$$

Let (a_n) be a sequence of positive numbers such that

$$\lim_{n \to \infty} \frac{\log a_n}{\log M_n} = -\infty.$$

If there exists n^* such that for every $n > n^*$ and for every $x \in [\varphi_1 = n]$

$$\prod_{i=0}^{n-1} \phi(\sigma^i(x)) \leq a_n,$$

then the multifractal spectrum of μ is continuous, concave and has unbounded domain.

(1) If the potential $-T(0) \log \psi$ is positive recurrent we denote by μ_m the corresponding equilibrium measure. One of the following alternatives occurs:
   (a) if $\log \phi \in L^1_{\mu_m}$ then there exists a point $\alpha(0)$ in the domain of f_μ such that for $\alpha < \alpha(0)$ the multifractal spectrum is increasing and for $\alpha > \alpha(0)$ it is constant;
   (b) if $\log \phi \notin L^1_{\mu_m}$ then the multifractal spectrum is strictly increasing.

(2) If the potential $-T(0) \log \psi$ is null recurrent then the multifractal spectrum is strictly increasing.

The next theorem describes the multifractal spectrum for small values of α, when there exists $q' \geq 1$ such that the function $T(q)$ is linear for $q > q'$. Recall that those are the level sets where the measure is highly concentrated (see Figure 4).

Definition 5.2.

$$\alpha(q') := \lim_{q \to (q')^-} -T'(q).$$

https://doi.org/10.1017/S0143385705000350 Published online by Cambridge University Press

THEOREM 5.3. *Let $q' \geq 1$. Assume the function $T(q)$ to be strictly convex for $q \in (0, q')$ and linear with slope $-A < 0$ for $q > q'$. Then the multifractal spectrum of μ is linear on the interval $(A, \alpha(q'))$ with slope q'.*

Proof. Let $\alpha \in (A, \alpha(q'))$; we claim that the Fenchel transform of $T(q)$ is given by

$$\inf_{-\infty<q<\infty}\{T(q) + \alpha q\} = T(q') + q'\alpha.$$

Note that if $q \in (0, q')$ then

$$\frac{T(q') - T(q)}{q' - q} < -\alpha. \tag{20}$$

In fact, assume by way of a contradiction that

$$\frac{T(q') - T(q)}{q' - q} > -\alpha.$$

The mean value theorem implies that there exists $q_1 \in (q, q')$ such that $T'(q_1) > -\alpha$. This contradiction leads to the claim. In fact, from equation (20) we obtain

$$T(q') + \alpha q' < T(q) + \alpha q.$$

Assume now that $q > q'$. Let $C > 0$ be such that

$$T(q) = -Aq + C.$$

Note that $\alpha - A > 0$. Therefore, since $q' < q$, we have

$$q'(\alpha - A) + C \leq q(\alpha - A) + C,$$

that is

$$T(q') + \alpha q' \leq T(q) + \alpha q.$$

This implies

$$f_\mu(\alpha) = \inf_{-\infty<q<\infty}\{T(q) + \alpha q\} = T(q') + q'\alpha,$$

as required. □

PROPOSITION 5.1. *If the potential $\log\phi$ is such that $P_{\mathrm{G}}(q \log\phi) = 0$ for every $q \geq 1$ then the multifractal spectrum is given by*

$$f_\mu(\alpha) = \alpha,$$

for $\alpha \in (0, \alpha(1))$.

Proof. Note that if $q \geq 1$ then $P_{\mathrm{G}}(q \log\phi) = 0$, therefore $T(q) \leq 0$. We prove that it is equal to zero. Assume by way of a contradiction that $T(q) < 0$. Then the function

$$t \to P_{\mathrm{G}}(-t \log\psi + q \log\phi)$$

is equal to zero for $t \in (T(q), 0)$ and negative for $t > \dim_{\mathrm{H}}(\mathcal{R})$. This contradicts the convexity of the function. Therefore, if $q \geq 1$ then $T(q) = 0$. The result follows from Theorem 5.3. □

https://doi.org/10.1017/S0143385705000350 Published online by Cambridge University Press

Example 5.4. Let G be the renewal graph. Sarig proved in [**15**] that if $\log \phi$ is a locally Hölder potential then there exists a critical value q_c (it might be infinity) such that:

(1) the potential $q \log \phi$ is strongly positive recurrent for $q \in (0, q_c)$ and transient for $q > q_c$;
(2) the pressure function $P_G(q \log \phi)$ is real analytic in $(0, q_c)$ and linear in (q_c, ∞); it is continuous but not analytic at q_c;
(3) the potential $q_c \log \phi$ can be positive recurrent, null recurrent or transient.

This framework provides a family of examples that illustrate Proposition 5.1.

We finally give an example where the multifractal spectrum has bounded domain.

Example 5.5. (Gauss map and the renewal graph) The Gauss map $G : [0, 1] \to [0, 1]$ is defined by

$$G(x) := \frac{1}{x} - \left[\frac{1}{x}\right],$$

where $[x]$ is the integer part of x. Denote by F the restriction of the Gauss map G to the renewal graph, that is,

$$F(x) := \begin{cases} G(x) & \text{if } x \in I_{ij} \text{ and } t_{ij} = 1, \\ \text{not defined} & \text{any other case,} \end{cases} \tag{21}$$

where $I_{ij} := ((i+1)^{-1}, i^{-1}) \cap G^{-1}((j+1)^{-1}, j^{-1})$. Denote by f the induced system on the set $(\frac{1}{2}, 1)$. The combinatorics of the induced graph is simple, there is just one cylinder of length n, which is $C_{1n(n-1)\ldots 2}$. Note that if $x \in (1/(n+1), 1/n)$ then

$$\left(\frac{n-1}{n^2-n+1}\right)^2 \le |F'(x)|^{-1} \le \left(\frac{n}{n^2+1}\right)^2.$$

If $x \in (\frac{1}{2}, 1)$ and $F(x) \in (1/(n+1), 1/n)$ then

$$x \in \left(\frac{n^2+1}{n^2+n+1}, \frac{n^2-n+1}{n^2}\right).$$

Using the combinatorics of the induced system and the above estimates we can bound the derivatives of the induced system f. That enables us to bound the pressure function,

$$\log\left((1/4)^t + \sum_{n=2}^{\infty}\left(\left(\frac{n^2+1}{n^2+n+1}\right)^2 \prod_{i=2}^{n}\left(\frac{i-1}{i^2-i+1}\right)^2\right)^t\right)$$
$$\le P_G(-t \log |f'|) \le \log\left((4/9)^t + \sum_{n=2}^{\infty}\left(\left(\frac{n^2-n+1}{n^2}\right)^2 \prod_{i=2}^{n}\left(\frac{i}{i^2+1}\right)^2\right)^t\right).$$

Computer estimates of the root d of the equation $P_G(-t \log |f'|) = 0$ and Theorem 3.1 yield

$$\dim_H(\mathcal{R}) = d \in (0.4, 0.6).$$

Note that $\mathcal{R}$ is the set of recurrent points such that their continued fraction expansion is determined by the renewal graph.

https://doi.org/10.1017/S0143385705000350 Published online by Cambridge University Press

The potential $-d\log|F'|$ has zero pressure and is recurrent (see Remark 5.2). Using the bounds on the derivative of F we obtain

$$\sum_{n\geq 1} n \sum_{F^n x = x} \exp\left(\sum_{i=0}^{n-1} (-d\log|F'(F^i x)|) 1_{[\varphi_1 = n]}(x)\right) \leq C + \sum_{n\geq 1}\left(\frac{1}{n!}\right)^{2d},$$

where C is a constant. This series converges, therefore the potential $-d\log|F'|$ is positive recurrent and there exists a (finite) equilibrium measure m. Since the Gurevich entropy is $\log 2$ and $0.4 \leq d$, using the variational principle (see [**15, 21**]) we can estimate the Lyapunov exponent,

$$\lambda_m := \int \log|F'|\,dm \leq \frac{5}{2}\log 2 \sim 1.732.$$

Note that λ_m measures the exponential speed of approximation of a number by its approximants. Recall that, Lebesgue almost everywhere, the Lyapunov exponent is $\pi^2/6\log 2 \sim 2.373$.

Let $\log\phi = -\log 2$ be a constant potential. It has pressure zero, is positive recurrent and the corresponding equilibrium measure is the *measure of maximal entropy*. Note that $q^* = -\infty$. Therefore, the multifractal spectrum f_μ is concave, has a unique maximum at $-T'(0)$ and has bounded domain.

Acknowledgements. This is part of the author's PhD thesis, written under the supervision of Omri Sarig and Peter Walters. Part of this paper was written while the author was a Marie Curie fellow at the Banach Center, Warsaw. The author was partially supported by FCT/POCTI/FEDER.

References

[1] L. Barreira and J. Schmeling, Sets of 'Non-Typical' points have full topological entropy and full Hausdorff dimension. *Israel J. Math.* **116** (2000), 29–70.

[2] R. Bowen. Hausdorff dimension of quasi-circles. *Publ. Math. Inst. Hautes Études Sci.* **50** (1979), 11–26.

[3] B. M. Gurevič. Topological entropy for denumerable Markov chains. *Soviet. Math. Dokl.* **10** (1969), 911–915.

[4] B. M. Gurevič. Shift entropy and Markov measures in the path space of a denumerable graph. *Soviet. Math. Dokl.* **11** (1970), 744–747.

[5] P. Hanus, D. Mauldin and M. Urbanski. Thermodynamic formalism and multifractal analysis of conformal infinite iterated function systems. *Acta. Math. Hungarica* **96** (2002), 27–98.

[6] B. P. Kitchens. *Symbolic Dynamics*. Springer, 1998.

[7] D. Mauldin and M. Urbanski. Dimension and measures in infinite iterated function systems. *Proc. London Math. Soc.* **73** (1996), 105–154.

[8] K. Nakaishi. Multifractal formalism for some parabolic maps. *Ergod. Th. & Dynam. Sys.* **20** (2000), 843–857.

[9] Y. Pesin. *Dimension Theory in Dynamical Systems*. Cambridge University Press, Cambridge, 1997.

[10] Y. Pesin and H. Weiss. A multifractal analysis of equilibrium measures for conformal expanding maps and Moran-like geometric constructions. *J. Stat. Phys.* **86** (1997), 233–275.

[11] M. Pollicott and H. Weiss. Multifractal analysis of Lyapunov exponent for continued fraction and Manneville Pomeau transformations and applications to Diophantine approximations. *Comm. Math. Phys.* **207** (1999), 145–171.

https://doi.org/10.1017/S0143385705000350 Published online by Cambridge University Press

[12] T. Prellberg and J. Slawny. Maps of intervals with indifferent fixed points: thermodynamic formalism and phase transitions. *J. Stat. Phys.* **66** (1992), 503–514.
[13] R. Ruelle. Repellers for real analytic maps. *Ergod. Th. & Dynam. Sys.* **2** (1982), 99–107.
[14] S. Ruette. On the Vere-Jones classification and existence of maximal measures for topological Markov chains. *Pacific J. Math.* **209** (2003), 365–380.
[15] O. Sarig. Thermodynamic formalism for countable Markov shifts. *Ergod. Th. & Dynam. Sys.* **19** (1999), 1565–1593.
[16] O. Sarig. Thermodynamic formalism for countable Markov shifts. *PhD Thesis*, Tel-Aviv University, 2000.
[17] O. Sarig. Thermodynamic formalism for null recurrent potentials. *Israel J. Math.* **121** (2001), 285–311.
[18] O. Sarig. Phase transitions for countable Markov shifts. *Comm. Math. Phys.* **217** (2001), 555–577.
[19] O. Sarig. Characterization of existence of Gibbs measures for countable Markov shifts. *Proc. Amer. Math. Soc.* **131**(6) (2003), 1751–1758.
[20] J. Schmeling. On the completeness of the multifractal spectra. *Ergod. Th. & Dynam. Sys.* **19** (1999), 1–22.
[21] P. Walters. *An Introduction to Ergodic Theory*. Springer, 1982.
[22] R. A. Wijsman. Convergence of sequence of convex sets, cones and functions. *Bull. Amer. Math. Soc.* **70** (1964), 186–188.
[23] R. A. Wijsman. Convergence of sequence of convex sets,cones and functions. II. *Trans. Amer. Math. Soc.* **123** (1966), 32–45.

https://doi.org/10.1017/S0143385705000350 Published online by Cambridge University Press