*Animal* (2011), **5:6**, pp 927–937 © The Animal Consortium 2011
doi:10.1017/S1751731110002685

# Podolian beef production on pasture and in confinement

A. Braghieri[1†], C. Pacelli[1], G. De Rosa[2], A. Girolami[1], P. De Palo[3] and F. Napolitano[1]

[1]*Dipartimento di Scienze delle Produzioni animali, Università degli Studi della Basilicata, Via dell'Ateneo Lucano 10, 85100 Potenza, Italy;* [2]*Dipartimento di Scienze del Suolo, della Pianta, dell'Ambiente e delle Produzioni animali, Università di Napoli 'Federico II', Via Università 133, 80055 Portici (NA), Italy;* [3]*Dipartimento di Sanità Pubblica e Zootecnia, Università di Bari, S. P. per Casamassima Km 3, 70010 Valenzano (BA), Italy*

(Received 13 May 2010; Accepted 10 October 2010; First published online 14 January 2011)

*From February to August 2007 the effect of rearing system (confined (C) v. free ranging (FR)) and season (spring v. summer) was evaluated on behaviour, immune and blood parameters and beef production (experiment 1). From February to August 2008, the effect of rearing system was only evaluated on beef production (experiment 2). A total of 12 Podolian young bulls were used each year. They were slaughtered at 18 and 15 months of age in 2007 and 2008, respectively. Herbage mass and pasture composition were monitored during the 2 years. Pasture availability, in terms of herbage mass and composition, as well as its seasonal development, was similar in the 2 years. In the first experiment, FR animals spent more time walking ($P < 0.05$), feeding ($P < 0.001$) and standing ($P < 0.01$) and showed a lower number of agonistic ($P < 0.05$) and non-agonistic social interaction than C animals ($P < 0.01$). Significant lower concentrations of serum urea nitrogen ($P < 0.001$) and creatinine ($P < 0.10$) in FR animals indicated a lower protein nutritional status due to inadequate protein availability at pasture. As a consequence, average daily gains ($P < 0.05$), slaughter weight ($P < 0.05$) and body condition score ($P < 0.01$) were lower in grazing animals as compared with C bulls. Cellular immune responsiveness was higher in FR animals ($P < 0.05$). Similarly, antibody titre to keyhole limpet hemocyanin was higher in FR bulls at the 2nd and 3rd month after antigen injection ($P < 0.05$), whereas it tended to be higher at the 4th month ($P < 0.10$). In both experiments, grazing negatively affected meat colour in terms of lightness. Eighteen-month-old bulls also showed lower final weight, weight gain and body conditions when kept outdoor: a possible consequence of nutrient deficits, as suggested by the metabolic status of FR animals. The same animals, however, benefited from FR in terms of natural behaviour expression and immune responsiveness. When the experiment was replicated the subsequent year, on animals slaughtered at 15 months of age, no differences between the performances of FR and C animals were detected. The earlier slaughter age system was also proportionally less dependent on external inputs as grazing was not extended to the dry season when herbage mass availability was lower.*

**Keywords:** pasture, Podolian cattle, beef production, animal welfare, behaviour

## Implications

Behavioural and immune indicators showed that bulls can benefit from an extensive rearing system based on pasture, although animal performances were lower than in confinement when animals were finished up to 18 months. These differences were not detectable when the age at slaughter was reduced to 15 months. In addition, an early slaughter age reduced the degree of competition with human nutrition, as the system proportionally rely more on pasture. A possible negative impact of free ranging was observed on meat colour in terms of reduced lightness.

† E-mail: ada.braghieri@unibas.it

## Introduction

Many beef cattle are kept in less-favoured areas. These areas are often characterized by being upland and with soils of low fertility. Therefore, most of these animals are finished in tie stalls, free stalls or feedlot. Pasture-based finishing of beef can potentially lower the cost of production and the concentration of manure, reduce the competition with humans in terms of food, increase animal welfare and be perceived by consumers as ethically sound (Napolitano *et al.*, 2005 and 2007).

On one hand, natural pastures provide an environment where animals can express their own natural behaviour. Apart from wild ancestors, which for cattle are extinct, and few examples of feral populations (e.g. Hernandez *et al.*, 1999), domestic herds kept in natural environments represent the main source of information about natural behaviour.



https://doi.org/10.1017/S1751731110002685 Published online by Cambridge University Press

Podolian cattle are considered the most direct descendants of the ancient wild bovine with a genetic relationship with similar breeds from the Balkans, Anatolia and the Middle East. They have been subjected to a lower selective pressure (artificial insemination has been rarely used) and a higher natural selection (animals kept in natural environments, where food search, avoidance of predators, maternal care, etc. were essential) as compared with other breeds (Napolitano *et al.*, 2005). This breed is characterized by a high rusticity and is well adapted to live in arid environment with poor vegetation, as suggested by skin pigmentation, well-developed dewlap and sturdy hooves. The most common rearing system is based on pasture with either no- or nocturnal shelter. Most often, cows and calves are not supplemented, whereas young adults may receive concentrates in the finishing period.

On the other hand, herbage production from natural pastures located in less-favoured areas is characterized by being unevenly distributed throughout the year, which leads to greater reliance on conserved forages and concentrates. Therefore, optimizing the nutrient supply from forage is also a key component in the development of sustainable finishing systems. In addition, this beef production system is not well received by meat industry, which tends to discount the value of the product on the basis of increased incidence of dark cutting, reduced carcass yield and poor conformation.

Therefore, the aim of this study was to test the effect of rearing system (free ranging (FR) *v.* confined (C)) from two points of view: performance and behavioural adaptation.

## Material and methods

*Experimental design*

The study was conducted in a farm located in Basilicata (southern Italy) at 338 m above sea level (40°45′2.16″N; 16°14′11.76″E). The mean annual rainfall in this area is below 600 mm.

The first experiment was conducted from February to August 2007 on 12 Podolian young bulls. They were kept in a grazing area with their mothers until 10 months of age and then abruptly weaned and divided in two groups: C and FR. Animals were aged about 11 months at the start of the experiment (329.83 ± 9.95 and 339.83 ± 10.00 days, respectively). FR animals were allowed to graze on a natural fenced pasture similar to the previous grazing area (18 ha of grassland and 2 ha of shrub vegetation) until slaughter (18 months). They were supplemented with 2 and 4 kg/day per bull of flour that was dispensed in a covered area from February to mid-June and from mid-June to August, respectively. The ingredients and chemical composition of the flour are reported in Table 1. Group C was kept in a loose barn with a straw-bedded resting area and an uncovered exercise area with a total space allowance of 13.4 $m^2$/bull. They received the same flour (6.0 and 7.5 kg/day per bull in spring and summer, respectively) and 2 kg/day per bull of straw.

The environmental parameters (temperature and relative humidity) recorded with a digital portable thermo-hygrometer (Model 1750-1/QM, Filotecnica, Salmoiraghi, Milan, Italy) are reported in Table 2.

**Table 1** *Ingredients and chemical composition of flour offered to C and FR Podolian young bulls in both experiments (2007 and 2008)*

| | |
|---|---|
| Ingredients (%) | |
| Oat | 31 |
| Barley | 31 |
| Field bean (*Vicia faba minor*) | 31 |
| Linseed | 3 |
| Mineral mix[1] | 1 |
| Chemical composition (%) | |
| DM | 91.4 |
| Ash (% DM) | 4.4 |
| CP (% DM) | 17.4 |
| Ether extract (% DM) | 2.7 |
| Crude fibre (% DM) | 7.9 |
| NDF (% DM) | 34.1 |
| MFU (kg DM)[2] | 1.11 |

C = confined; FR = free range; DM = dry matter; MFU = meat forage unit.
[1]Vitasol Spa, Brescia, Italy.
[2]INRA (1988).

In order to verify whether a rearing system is less dependent on concentrates, and mainly based on pasture could be used, the experiment was replicated from February to June 2008 in the same field. The experiment conditions were similar to those used in the previous experiment, a part from the fact that the bulls were slaughtered at 15 months, before the dry season reduced herbage mass availability on pasture.

Average temperature and relative humidity in spring and summer 2008 were 12.27°C ± 0.98°C and 66.63% ± 2.45%, and 20.9°C ± 1.56°C and 57.12% ± 4.57%, respectively.

Similarly to the previous experiment, 12 male Podolian animals were divided in two groups, C and FR, aging approximately 10 months (303.40 ± 2.34 *v.* 300 ± 2.5 days, respectively). FR group received 3 kg/day per bull of flour supplementation from May to the end of June. The ingredients and the chemical composition of the flour were similar to those reported in the first experiment (Table 1). Group C received the same flour (5.0 and 7.0 kg/day per bull in spring and summer, respectively) and 2 kg/day per bull of straw. In both experiments water was always available to all the animals.

Data on pasture composition and herbage mass, as well as the performance of grazing and C animals, were recorded in both years. Behaviour measurements, immune responses and biochemical serum profile were monitored only during the first year.

*Pasture composition and herbage mass*

Grass availability (dry matter (DM)/ha) was measured from December 2006 to August 2007 and from December 2007 to June 2008 in experiments 1 and 2, respectively. Fifteen iron-fenced net boxes (1.5 × 1.5 m, 1 m height) were randomly placed on grazing area (Meijs *et al.*, 1982; Martiniello *et al.*, 2007). Herbage was harvested at ground level from three 1 × 1 m areas placed into boxes. The total herbage mass was weighted and sub-sampled to evaluate DM and chemical



https://doi.org/10.1017/S1751731110002685 Published online by Cambridge University Press

**Table 2** *Environmental parameters recorded at pasture and in the barn throughout the behavioural observations performed in experiment 1 (2007)*

| Behavioural session | I | II | III | IV | V | VI | VII | VIII | IX |
|---|---|---|---|---|---|---|---|---|---|
| Barn | | | | | | | | | |
| Temperature (°C) | 8.7 | 21.0 | 16.4 | 23.5 | 20.9 | 37.9 | 40.09 | 26.4 | 34.9 |
| Relative humidity (%) | 85.0 | 70.4 | 83.3 | 41.8 | 69.6 | 22.0 | 26.0 | 40.4 | 26.6 |
| Pasture | | | | | | | | | |
| Temperature (°C) | 15.0 | 19.7 | 16.3 | 28.5 | 19.5 | 38.6 | 33.0 | 22.3 | 34.0 |
| Relative humidity (%) | 73.0 | 74.9 | 85.0 | 37.8 | 67.0 | 13.6 | 28.0 | 41.0 | 27.8 |

composition (200 to 300 g) and to estimate the botanical composition (grass, legumes, composites and others).

The samples of herbage were dried in a forced-air oven at 60°C until reaching constant weight and ground with a 1-mm screen for subsequent chemical analysis. For herbage and flour, the DM content was determined by drying samples at 105°C until reaching constant weight. Ash content was determined in a muffle furnace at 550°C overnight. Crude protein (CP) was determined using the Kjeldahl method by multiplying the concentration of nitrogen (N) by a factor of 6.25 and ether extract with the Soxhlet method (Association of Official Analytical Chemists (AOAC), 1990). Neutral detergent fibre (NDF) was determined according to Van Soest *et al*. (1991).

*Behavioural measurements*

Two observers carried out both the familiarization procedure and the observations. The observers spent several days with the experimental animals in order to accustom the groups to their presence. The success of the habituation was measured in terms of the distance the animals kept from the observers. The familiarization procedure ended when the observer was able to move around at a distance of approximately 0.5 to 1.5 m. Behavioural observations were performed in nine sessions through continuous focal animal recording method (Martin and Bateson, 2007). The observers were balanced across groups and time of the day. Six-hour period of continuous observations were alternatively conducted from 0600 to 1200 h and from 1200 to 1800 h. In each session a different animal was chosen. Therefore, all the animals were included in the observations, whereas three animals for each group were sampled twice.

The duration (accuracy: 1 s) of the observed behaviours was recorded. The nine behavioural observations were divided into two seasons: spring (mid-February to mid-June; five recordings) and summer (mid-June to August; four recordings). The following parameters were recorded: posture (standing or lying) and activity such as grazing/feeding (biting or chewing the herbage, and, only for group FR, walking with muzzle close to the ground), walking, resting (opened or closed eyes, but no other overt activity), ruminating and alert, whereas all other behaviours (e.g. drinking, vocalization and self-grooming) were recorded as 'other'. The proportion of time spent on each behaviour was calculated for each of the observation session. In addition, behaviours such as self-grooming, licking objects, vocalization, agonistic (pushing, butting or threatening co-specifics) and non-agonistic interactions (sniffing, horning or nuzzling co-specifics) were recorded as number of events.

The distance covered during the observation session was recorded by means of pedometer attached to the waist of the observer.

Ingestive behaviour was studied only for group FR. During the 6-h period, every hour, the number of bites and bite rate (number of bites per grazing minute) were recorded (using a mechanical counter) along with herbage height by means of a herb metre. Bite mass and botanical preferences were determined in hand-plucked samples (15 simulated plucks/h). Observers mimicked the bite size of tame animals by visual observation of the bites and clipping or hand plucking simulated bites from the vegetation. This technique allowed estimating intake rate (bite mass $\times$ bite rate) of FR animals (Gordon, 1995). Samples were sealed in polythene bags and taken to the laboratory in an icebox. Subsequently, they were weighted and botanical composition (grass, legumes, composites and others) was evaluated. Then, herbage samples were dried and analysed for chemical composition as previously described.

*Immune responses and biochemical serum profile*

At 3 months after grouping phytohemagglutinin (PHA; 1 mg, Sigma, Milan, Italy) dissolved in 1 ml of sterile saline solution was injected subcutaneously into the middle of 2-cm wide circles marked on shaved skin on the upper side of each shoulder. The skin-fold thickness (mm) was determined with a caliper before PHA injection and 24 h after. For each animal, a mean increase in skin-fold thickness (24-h thickness–pre-injection thickness) was calculated using the two measurements gathered from shoulders.

The animals were injected subcutaneously with 10 mg (5 mg per shoulder) of Keyhole Limpet Hemocyanin (KLH, Sigma, Milan, Italy) dissolved in 2 ml of sterile saline solution and emulsified in an equal volume of incomplete Freund's adjuvant a month after grouping. Another injection without adjuvant was repeated after 1 month. Antibody titre was evaluated at monthly intervals on serum collected by coccygeal venipuncture using vacutainer tubes. The concentrations of IgG anti-KLH in serum were determined by an ELISA test performed in 96-well, U-bottomed microtitre plates. In short, wells were coated with the antigen (0.5 mg of KLH/ml of phosphate buffered saline (PBS)) and incubated overnight at 4°C. PBS (pH 7.2) and 0.05% Tween 20 was used as washing buffer and 10% reconstituted milk was incubated



https://doi.org/10.1017/S1751731110002685 Published online by Cambridge University Press

to block non-specific binding. Then the serum (1 : 1600 dilution in PBS; 100 ml per well) was added and incubated at 37°C for 1 h. The extent of antibody binding was detected using a rabbit-anti-bovine IgG-horseradish peroxidase conjugate (Sigma Aldrich, Milan, Italy) diluted at 1 : 30 000 (100 ml per well). Optical density was measured at a wavelength of 450 nm using Rosys Anthos 2020 ELISA reader (Diessechem, Milan, Italy). The intra- and inter-assay CV was 5.66% and 4.92%, respectively. The assay was optimized in our laboratory for concentrations of coating antigen, serum and detector antibody.

For the determination of biochemical metabolites including glucose, cholesterol, total proteins, aspartate aminotransferase (AST), alanine aminotransferase (ALT), lactate dehydrogenase (LDH), creatine kinase (CK), triglyceride, phosphorus, chloride, calcium, urea and creatinine, required quantity of blood serum was taken. Standard commercial kits were used for analysis and procedures were adopted as recommended by the manufacturer of kits (PKL, Pokler, Italy, Geneva). After processing samples and standards provided with the kits, absorbance of the standard and the samples was determined and the concentrations of respective metabolites in samples were computed, using the formula: concentration of a metabolite = absorbance of sample divided by absorbance of standard and multiplied by standard concentration. Glucose concentrations were determined by the enzymatic kinetic colorimetric test (glucose oxidase-peroxidase antiperoxidase (GOD-PAP)). The standard had glucose concentration of 100 mg/dl. For cholesterol, enzymatic colorimetric test using diagnostic kit was applied. The concentration of cholesterol in the standard was 200 mg/dl. Total proteins content in the serum were measured by the Biuret method. AST and ALT activities in blood serum samples were determined by the International Federation of Clinical Chemistry and Laboratory Medicine (IFCC) colorimetric method. For triglycerides, the standard contained 200 mg/dl of triglycerides. LDH activity in serum samples was determined by the French Society of Clinical Biology/optical standard method (SFBC/OPT). For CK activity, the creatinine kinase N-acetylcysteine method was performed. Phosphorus concentration was determined by enzymatic kinetic colorimetric test. The standard had phosphorus concentration of 4 mg/dl. For chloride, the enzymatic colorimetric test (mercury thyocianate method) was used. The concentration of chloride in the standard was 100 mEq/l. Calcium content in the serum samples was determined by the enzymatic kinetic colorimetric test (Arsenazo III). The concentration of calcium in the standard was 10 mg/dl. Urea concentrations were determined by the enzymatic UV test glutamate dehydrogenase (Urease-GLDH), using a commercial kit. The standard had urea concentration of 50 mg/dl. For creatinine, an enzymatic colorimetric test was performed. Both in chromatic methods and in kinetics methods a UV spectrophotometer (Beckman Coulter Inc., DU800, Fullerton, CA, USA) was used.

*Animal performance and post-slaughter measurements*

Animals were individually weighted at monthly intervals in order to estimate the average daily gain (ADG, g/day) and body condition score (BCS) on a 9-point scale (Richards *et al.*, 1986). Animals from experiments 1 and 2 were slaughtered at 18 and 15 months of age, respectively. They were weighed on the morning of slaughter, after 12 h of fasting, to record full-body weight and transported to a commercial slaughter facility. Slaughter occurred according to industrial routines used in Italy and in accordance to the EU Directive 93/119/CE. One hour after slaughter, each carcass was weighed and dressing percentage was calculated.

Colour was measured using a colour-meter Minolta CR 200 (D65: illuminant; Konica Minolta Sensing Inc., Sakai-ku, Sakai, Osaka, Japan) on 1 cm thick steaks from *Longissimus dorsi* muscle, removed from the right half-carcass 24 h *post mortem*. Prior to the measurement, meat samples were allowed to bloom for 1 h at 3°C ± 1°C, stored in a plastic tray and over wrapped with a polyethylene film.

*Statistical analyses*

Data were analysed with statistical analysis software (Statistical Analysis Software (SAS), 1990). Behavioural data were analysed using the observation session as the experimental unit. These data were subjected to analysis of variance (ANOVA) with season (spring and summer), group (C and FR) and their interactions as factors.

Data on animal performance, cellular immune response and meat quality were subjected to ANOVA with one factor (group), whereas the antibody titre and blood metabolites were analysed using the PROC MIXED with time as repeated factor and group as non-repeated factor. Both analyses used the animal as the experimental unit.

Only for group FR, data on ingestive behaviour were subjected to ANOVA with one factor (season), whereas chemical and botanical composition of pasture and hand-plucked samples were analysed using ANOVA with season, source (pasture and hand plucking) and their interactions as factors. In addition, for this statistical model the animal was used as the experimental unit.

In addition, Pearson correlation coefficients between herbage height *v.* bite rate, bite mass and intake rate were computed.

## Results and discussion

*Pasture composition and herbage mass*

The main features of herbage collected in 2007 from different sources (pasture and hand-plucked samples) are reported in Table 3 in relation to the season. No significant interactions between the source and the season were observed.

Biomass availability was much lower in summer than in spring, although differences were not significant due to the high variability of samples. Compared with botanical composition, season significantly affected percent Grasses, which were lower in summer ($P < 0.05$) and Composites ($P < 0.05$), which increased in summer. In agreement with Hessle *et al.* (2008), in both pasture and hand-plucked samples chemical composition was affected by season. In particular, DM ($P < 0.001$), crude fibre ($P < 0.01$) and NDF ($P < 0.05$) increased during summer



https://doi.org/10.1017/S1751731110002685 Published online by Cambridge University Press

**Table 3** *Botanical and chemical composition (means ± s.e.) of pasture and hand-plucked samples in experiment 1 (2007)*

| | Pasture | | Hand-plucked samples | | *P* | |
|---|---|---|---|---|---|---|
| | Spring | Summer | Spring | Summer | Season | Source |
| Herbage mass (g/m$^2$) | 849.72 ± 399.74 | 568.03 ± 399.74 | – | – | ns | – |
| Grass (%) | 81.83 ± 5.87 | 60.63 ± 5.87 | 78.62 ± 5.87 | 61.26 ± 5.87 | <0.05 | ns |
| Legumes (%) | 5.99 ± 2.12 | 6.36 ± 2.12 | 4.62 ± 2.12 | 3.27 ± 2.12 | ns | ns |
| Composites (%) | 7.24 ± 6.14 | 23.15 ± 6.14 | 2.98 ± 6.14 | 21.32 ± 6.14 | <0.05 | ns |
| Others (%) | 4.94 ± 3.03 | 9.86 ± 3.03 | 13.78 ± 3.03 | 14.15 ± 3.03 | ns | ns |
| DM (%) | 29.98 ± 6.12 | 66.72 ± 6.12 | 26.97 ± 6.12 | 61.82 ± 6.12 | <0.001 | ns |
| CP (% DM) | 9.06 ± 1.31 | 5.18 ± 1.31 | 13.40 ± 1.31 | 10.96 ± 1.31 | <0.05 | <0.01 |
| Ether extract (% DM) | 1.34 ± 0.17 | 1.20 ± 0.17 | 1.35 ± 0.17 | 1.58 ± 0.17 | ns | ns |
| Crude fibre (% DM) | 24.34 ± 2.24 | 33.95 ± 2.24 | 21.04 ± 2.24 | 30.58 ± 2.24 | <0.01 | ns |
| Ash (% DM) | 13.29 ± 2.53 | 9.98 ± 2.53 | 16.78 ± 2.53 | 8.06 ± 2.53 | <0.05 | ns |
| NDF (% DM) | 47.09 ± 4.07 | 59.47 ± 4.07 | 45.35 ± 4.07 | 56.10 ± 4.07 | <0.05 | ns |
| MFU (kg DM)[1] | 0.64 ± 0.03 | 0.51 ± 0.03 | 0.68 ± 0.03 | 0.57 ± 0.03 | <0.01 | <0.10 |

DM = dry matter; MFU = meat forage unit.
[1]INRA (1988).

**Table 4** *Botanical and chemical composition (means ± s.e.) of pasture in experiment 2 (2008)*

| | Spring | Summer | *P* |
|---|---|---|---|
| Herbage mass (g/m$^2$) | 643.65 ± 224.67 | 455.75 ± 275.17 | ns |
| Grass (%) | 58.63 ± 5.30 | 43.92 ± 6.50 | ns |
| Legumes (%) | 15.37 ± 1.37 | 6.13 ± 1.68 | <0.05 |
| Composites (%) | 13.44 ± 1.38 | 23.15 ± 1.69 | <0.05 |
| Others (%) | 13.23 ± 4.15 | 29.69 ± 5.08 | <0.10 |
| DM (%) | 21.58 ± 11.35 | 59.95 ± 13.91 | ns |
| CP (% DM) | 13.50 ± 0.68 | 6.34 ± 0.84 | <0.01 |
| Ether extract (% DM) | 1.32 ± 0.25 | 1.11 ± 0.30 | ns |
| Crude fibre (% DM) | 24.45 ± 3.45 | 37.60 ± 4.22 | ns |
| Ash (% DM) | 13.66 ± 1.82 | 9.95 ± 2.23 | ns |
| NDF (% DM) | 46.85 ± 4.45 | 65.50 ± 5.45 | <0.10 |
| MFU (kg DM)[1] | 0.62 ± 0.05 | 0.45 ± 0.06 | ns |

DM = dry matter; MFU = meat forage unit.
[1]INRA (1988).

as compared with spring. Conversely, in summer CP ($P < 0.05$), ash ($P < 0.05$) percentages and meat forage unit (MFU) values ($P < 0.01$) were lower.

Measurements recorded during the second year are reported in Table 4. Herbage mass was lower in summer than in spring, although, as for the previous year, differences were not significant.

A predominance of the grass family was observed but with no significant seasonal differences, whereas legumes were significantly ($P < 0.05$) higher in spring than in summer, and composites ($P < 0.05$) and others ($P < 0.10$) were higher in summer than in spring.

As to chemical composition (Table 4), seasonal differences were significant ($P < 0.01$) only for CP percentage that was higher in spring as compared with summer, whereas NDF fraction tended to increase ($P < 0.10$) in summer, as already observed in the previous experiment. On the whole, pasture availability, in terms of herbage mass, composition and seasonal development, was similar to that observed in the previous experiment (2007).

**Table 5** *Ingestive behaviour (means ± s.e.) expressed by free ranging Podolian young bulls during experiment 1 (2007)*

| | Spring | Summer | *P* |
|---|---|---|---|
| Bite rate (n/min) | 45.79 ± 2.99 | 26.75 ± 5.17 | <0.01 |
| Bite mass (g DM) | 1.12 ± 0.15 | 1.19 ± 0.11 | ns |
| Intake rate (g DM/min) | 54.88 ± 4.89 | 32.02 ± 8.48 | <0.05 |
| Sward height (cm) | 29.12 ± 4.29 | 42.87 ± 7.43 | <0.10 |

DM = dry matter.

*Ingestive behaviour*

No significant differences in botanical composition were detected between pasture and hand-plucked samples (Table 3). However, the diet selected by the animals, as assessed by hand plucking, was higher in CP percentages in comparison with pasture samples (12.17% ± 0.87% *v.* 7.11% ± 0.93%, respectively; $P < 0.01$) and tended to have higher MFU values (0.63 ± 0.02 *v.* 0.58 ± 0.02 MFU/kg DM; $P < 0.10$). This may be ascribed to the 'foraging strategy' of the animal (Gordon and Illius, 1992). According to this theory, grazing herbivores make a series of short-term decisions (e.g. about which plants or plant portions to select and how long to search) in order to ingest a diet adequate to satisfy their nutrient requirements for maintenance, growth and reproduction (Gordon, 1995; Dumont *et al.*, 2007). No significant effect of source was detected for the other chemical components.

The ingestive behaviour of FR animals is depicted in Table 5. While foraging, an animal takes a series of bites varying in size from the herbage on offer. Combination of bite size and short-term rate of biting is defined as the short-term intake rate. These variables are related, as larger bite size need more time for its chewing and swallowing. Usually



https://doi.org/10.1017/S1751731110002685 Published online by Cambridge University Press

ruminants prefer plants with faster ingestion (De Rosa *et al.*, 1997). In this experiment, bite rate resulted higher ($P < 0.01$) in spring than in summer. Similarly, Lamoot *et al.* (2005) reported that bite rate of cattle were higher in spring than in the other seasons, probably because in this period plant growth starts and provides the herbivores with more palatable and digestible forage. In this study crude fibre and NDF percentages were lower in spring. Therefore, seasonal changes in forage quality parameters were reflected in adaptations of ingestive behaviour. No significant differences between the seasons were observed for bite mass. Therefore, as a consequence of a higher bite rate during spring, intake rate was higher in spring than in summer ($P < 0.05$). Cattle usually exhibit very typical selectivity patterns preferentially selecting against reproductive grass patches (Wallis De Vries and Daleboudt, 1994; Dumont *et al.*, 2007), especially as these become more mature (Ginane *et al.*, 2003).

Herbage height tended ($P < 0.10$) to be higher in summer. In agreement with Gibb *et al.* (1997) and Burns and Sollenberger (2002), this variable was negatively correlated with bite rate ($r = -0.60$; $P < 0.001$) and positively related to bite mass ($r = 0.55$; $P < 0.01$). In particular, in a Boval *et al.*'s (2007) experiment, bite mass increased by 9 mg per additional centimetre of extended tiller length. A lower biting rate would be a consequence of greater harvested leaf length or a higher bite mass requiring more time for prehension, manipulation and mastication (Cosgrove, 1997).

It has been suggested that bovids are morphologically constrained to graze on very short swards because of the lack of the upper incisors (Illius and Gordon, 1987). Nevertheless, they do prefer shorter herbage if less mature (Wallis De Vries and Daleboudt, 1994; Ginane *et al.*, 2003). Accordingly, as already reported, in our experiment bulls showed higher intake rates in spring than in summer because of the lower DM content.

*Maintenance and social behaviour*

The activity budget observed in 2007 and expressed as percentage of time is reported in Table 6. FR animals walked longer distances in comparison with C animals ($P < 0.05$). This is closely related to the lower inactivity ($P < 0.01$) observed in FR bulls, which needed to walk in order to feed, as also suggested by the higher percentage time spent standing ($P < 0.01$) and feeding ($P < 0.001$). This latter activity included walking with muzzle close to the ground in the group FR that was precluded to the group C. In general, cattle allowed to graze are perceived as animals with higher welfare standards because they can express natural behaviours, such as grazing and exploration (Hemsworth *et al.*, 1995). The distance covered by FR animals in this study was similar to that reported by Krohn *et al.* (1992) in dairy cows kept in extensive environment (1.7 km), whereas group C showed a walking distance similar to that covered by dairy cows (0.6 km) kept in loose housing systems (Kempkens and Boxberger, 1987). Although no effects of season and group × season were observed on walked distance, the mean distance travelled was lower in summer as compared with spring. Accordingly, in response to high ambient temperature (Table 1) walking ($P < 0.001$) and standing ($P < 0.05$) were lower in summer, whereas inactivity was higher ($P < 0.05$). Dumont *et al.* (2007) observed that daily distance walked by cattle either increased or remained relatively constant over the grazing season for animals at a low stocking rate, which suggests that they kept on exploring the grazing area. However, these results are not necessarily in contrast with our results, as in this experiment the higher supplementation given to the grazing animals could have reduced the motivation to explore and ingest material of poor quality from pasture (Table 3). In fact, a significant interaction season × rearing system was observed for walking ($P < 0.05$). This result can be attributed to the fact that group FR walked more than group C in spring (8.42% ± 0.90% *v.* 3.39% ± 0.90%, respectively; $P < 0.01$), whereas no differences between groups were observed in summer (0.94% ± 1.02% *v.* 1.00% ± 1.02%, respectively). The same interaction only tended to be significant for feeding ($P < 0.10$), as differences between groups were much higher in spring (43.60% ± 3.90% *v.* 14.87% ± 3.90%, respectively; $P < 0.001$) than in summer (28.72% ± 4.36% *v.* 14.43% ± 4.36%, respectively; $P < 0.05$). Accordingly, it has been observed that both low availability of grass in

**Table 6** *Activity budget (means ± s.e.) of Podolian young bulls expressed over 6-h periods and recorded in nine sessions during experiment 1 (2007)*

| | RS | | S | | *P* | | |
|---|---|---|---|---|---|---|---|
| | C | FR | Spring | Summer | RS | S | RS × S |
| Distance travelled (km) | 0.60 ± 0.37 | 1.92 ± 0.37 | 1.48 ± 0.35 | 1.04 ± 0.39 | <0.05 | ns | ns |
| Standing[1] | 44.17 ± 3.87 | 55.51 ± 3.87 | 59.12 ± 3.65 | 40.56 ± 4.08 | <0.01 | <0.05 | ns |
| Feeding[1] | 14.65 ± 2.92 | 36.16 ± 2.92 | 29.24 ± 2.75 | 21.57 ± 3.08 | <0.001 | ns | <0.10 |
| Walking[1] | 2.20 ± 0.68 | 4.68 ± 0.68 | 5.90 ± 0.64 | 0.97 ± 0.71 | <0.05 | <0.001 | <0.05 |
| Inactivity[1] | 50.90 ± 4.05 | 31.23 ± 4.05 | 33.87 ± 3.82 | 48.25 ± 4.27 | <0.01 | <0.05 | ns |
| Alert[1] | 4.14 ± 2.07 | 4.12 ± 2.07 | 6.83 ± 1.95 | 1.43 ± 2.18 | ns | ns | ns |
| Ruminating[1] | 23.73 ± 2.89 | 21.95 ± 2.89 | 20.11 ± 2.72 | 25.57 ± 3.04 | ns | ns | ns |
| Other[1] | 4.38 ± 0.97 | 1.85 ± 0.97 | 4.03 ± 0.91 | 2.20 ± 1.02 | ns | ns | ns |

RS = rearing system; S = season; C = confined; FR = free range.
[1]Data expressed as percentage of time.



https://doi.org/10.1017/S1751731110002685 Published online by Cambridge University Press

**Table 7** *Other activities (means ± s.e.) of Podolian young bulls observed over 6-h periods and recorded in nine sessions during experiment 1 (2007)*

| | RS | | S | | *P* | |
|---|---|---|---|---|---|---|
| | C | FR | Spring | Summer | RS | S |
| Self-grooming[1] | 9.02 ± 1.78 | 8.60 ± 1.78 | 12.50 ± 1.68 | 5.12 ± 1.88 | ns | <0.05 |
| Allo-grooming[1] | 7.80 ± 1.71 | 3.75 ± 1.71 | 6.30 ± 1.62 | 5.25 ± 1.80 | ns | ns |
| Agonistic interactions[1] | 8.17 ± 1.61 | 3.00 ± 1.61 | 5.30 ± 1.52 | 5.87 ± 1.70 | <0.05 | ns |
| Non-agonistic interactions[1] | 18.62 ± 1.77 | 9.95 ± 1.77 | 13.70 ± 1.67 | 14.87 ± 1.87 | <0.01 | ns |
| Object licking[1] | 6.55 ± 1.10 | 2.47 ± 1.10 | 6.40 ± 1.03 | 2.62 ± 1.15 | <0.05 | <0.05 |
| Vocalizations[1] | 0.55 ± 0.92 | 3.40 ± 0.92 | 3.70 ± 0.87 | 0.25 ± 0.98 | <0.05 | <0.05 |

RS = rearing system; S = season; C = confined; FR = free range.
[1]Data expressed as number of events.

summer (Ruckebusch and Bueno, 1978) and feeding supplements (Sarker and Holmes, 1974) reduce grazing time.

As to behavioural categories expressed as number of events (Table 7), group FR showed lower agonistic ($P < 0.05$) and non-agonistic ($P < 0.01$) interactions in comparison with group C, as a possible consequence of higher inter-animal distance and different priorities (i.e. more time needed to eat and lower spare time). A lower number of non-agonistic social interactions, such as sniffing and rubbing, have been also observed by Krohn (1994) in extensively kept dairy cows as compared with animals in intensive systems. The mean number of agonistic events recorded in this study was much lower than that reported in previous experiments (Wierenga, 1984; Miller and Wood-Gush, 1991), possibly because the bulls were all part of one herd since their birth, consequently their social status was well established, whereas agonistic interactions prominently occur in occasion of establishing the social order in dynamic groups (Bouissou *et al.*, 2001; Bouissou and Boissy, 2005).

Self-grooming was higher in spring than in summer ($P < 0.01$). This comfort behaviour is closely related to hygiene and possibly associated to the higher rainfalls and body dirtiness usually observed in this season.

Object licking was higher in spring and in C animals as compared with summer and FR animals ($P < 0.05$). In agreement with Huber *et al.* (2008), the lower object licking observed in group FR may be explained by the higher walking and feeding activities performed at pasture, whereas group C expressed more object licking while spending less time in walking and feeding. Abnormal oral behaviours are linked with the lack of opportunities to express explorative behaviours and to spend time feeding (Bergeron *et al.*, 2006). Accordingly, Redbo and Nordblad (1997) showed that increasing dietary forage results in longer feeding duration along with reduced abnormal oral behaviour. Recently, Keyserlingk *et al.* (2009) reviewed papers showing that cattle do prefer pasture even over a well-designed free-stall barn.

Vocalization were higher both in FR animals ($P < 0.05$) and in spring ($P < 0.05$). Vocal signals are used to keep contact with and locate other herd members, therefore they were more frequently expressed in FR conditions and in spring when animals moved more often (Watts and Stookey, 2000).





**Figure 1** Antibody titres to keyhole limpet hemocyanin (KLH) of confined (C) and free range (FR) Podolian young bulls were evaluated before the first antigen administration and then at 30-day intervals using ELISA techniques. Group FR had a humoral immune response constantly higher than group C, with significant differences 2 and 3 months after antigen injection ($P < 0.05$), whereas differences only tended to be significant at 3 months ($P < 0.10$) and were not significant at 4 and 5 months.

*Immune responses*

*In vivo* cellular immune response as recorded in experiment 1 was higher in the group FR than in the group C (4.51 ± 0.97 *v.* 1.07 ± 0.97 mm, respectively; $P < 0.05$). Similar results were obtained in different ruminant species such as sheep (Braghieri *et al.*, 2007) and buffaloes (De Rosa *et al.*, 2007) in which grazing animals showed higher skin thickening as compared with C animals.

The antibody titres are presented in Figure 1. For this variable the interaction between time and group tended to be significant ($P < 0.10$). Group FR had a humoral immune response constantly higher than group C, with significant differences between groups observed 2 and 3 months after antigen injection ($P < 0.05$); differences only tended to be significant at 3 months ($P < 0.10$) and were not significant at 4 and 5 months. It is generally recognized that environmental or management stressors are associated to impair immunological functions in farm animals. Previous studies indicated that space allowance did not affect humoral immune response of beef cattle (Fisher *et al.*, 1997a and 1997b). However, in those studies animals were all kept



https://doi.org/10.1017/S1751731110002685 Published online by Cambridge University Press

indoor, whereas in our experiment indoor animals were compared with animals kept outdoor. Therefore, the housing system may have affected more markedly the immune status of the animals than a different space allocation within a single housing system (indoor). Accordingly, in an earlier study the housing system affected the immune function of calves with or without access to a fenced area (Cummins and Brunner, 1991). In this study, the effects of housing system on immune functions indicated a differential success in coping of animals of different groups, thus providing a measurement for the potential higher disease susceptibility and lower welfare of animals kept indoor. Accordingly, some aspects of cattle health, such as lameness and mastitis, can be improved by free access to grazing areas (reviewed by Keyserlingk *et al*., 2009).

*Metabolic profile and weight gains*

Haematological parameters as recorded in experiment 1 are presented in Table 8. Most blood metabolites levels were not affected by time and the interaction group × time. The rearing system did not change the energetic metabolism of Podolian young bulls in terms of glucose, triglycerides and cholesterol levels. Similar results were obtained by Marino *et al*. (2009) in animals of the same breed and age.

Blood total proteins were also unaffected by the rearing system, thus indicating an overall similar nutritional state of the animals from the two experimental groups (Doornenbal *et al*., 1988). However, serum urea-N levels were lower ($P < 0.001$) and creatinine tended to be lower ($P < 0.10$) in the group FR than in the group C. In addition, creatinine increased with time ($P < 0.001$). In young growing ruminants, these two metabolites are deemed to be important indicators of the protein nutritional state (Cabaraux *et al*., 2005). In particular, the amount of creatinine released in the blood is related to the growth rate of the muscular tissue, as the precursor creatinine is mainly contained in the skeletal muscles (Doornenbal *et al*., 1988). In addition, urea-N levels can provide information about the quality of the pasture in terms of protein availability. In a previous study, Tucker and Hentges (1983) observed lower urea-N levels in beef cattle grazing on a pasture characterized by low protein contents. In our study the herbage ingested by FR animals from pasture had low protein (CP, 12.17% DM) and energetic contents (0.63 MFU/kg DM). Although these animals also received a feeding supplementation (2 and 4 kg/day per bull of flour in spring and summer, respectively), they were likely to be unable to fully express their growth potential, thus showing lower weight gains, slaughter weight and body condition scores as compared with C animals (Table 9).

The higher levels of alkaline phosphatase observed in group C ($P < 0.05$) also indicated a higher growth potential in C animals, as this enzyme is related to bone growth (Freedland and Szepesi, 1971).

Phosphorous concentration was higher in group C than in group FR ($P < 0.05$). Serum phosphorous level is only partly dependent on its dietary intake (Karn, 2001), and the lower levels observed in FR animals may be attributed to their higher physical activity and thermoregulation needs. Although concentrations of serum calcium are more closely physiologically regulated than phosphorous, the higher levels observed in group FR as compared with group C ($P < 0.01$) can be due to the fact that concentrates are low in calcium, whereas forages are generally satisfactory sources of this element (Suttle, 2009).

Animal performances recorded in the second experiment are illustrated in Table 10. Although both groups achieved lower performances compared with the previous experiment possibly because of the lower initial weight, the rearing system did not affect final weight, ADGs and BCSs. These results show that FR system can produce performances similar to those obtained in C conditions when the animals are slaughtered at an earlier age. The ratio (kilogram of concentrate fed to the animals × days of feedings supplementation)/total amount of meat in kilogram was much

**Table 8** *Blood metabolites (means ± s.e.) of C and FR Podolian young bulls in experiment 1 (2007)*

| Group | C | FR | *P* |
|---|---|---|---|
| Glucose (mmol/l) | 3.56 ± 0.08 | 3.37 ± 0.08 | ns |
| Triglycerides (mmol/l) | 0.46 ± 0.01 | 0.45 ± 0.01 | ns |
| Cholesterol (mmol/l) | 2.59 ± 1.12 | 2.60 ± 0.13 | ns |
| LDH (IU/l) | 991.30 ± 35.13 | 1015.89 ± 35.81 | ns |
| Total protein (g/l) | 69.08 ± 0.51 | 68.86 ± 0.51 | ns |
| BUN (mmol/l) | 12.57 ± 0.56 | 8.39 ± 0.57 | <0.001 |
| Creatinine (mmol/l) | 120.19 ± 3.90 | 110.03 ± 4.00 | <0.10 |
| Creatine kinase (IU/l) | 173.89 ± 9.70 | 166.17 ± 9.67 | ns |
| Phosphorus (mmol/l) | 2.23 ± 0.05 | 2.02 ± 0.05 | <0.05 |
| Calcium (mmol/l) | 2.34 ± 0.02 | 2.50 ± 0.02 | <0.001 |
| AST (IU/l) | 79.47 ± 4.18 | 77.17 ± 4.18 | ns |
| ALT (IU/l) | 25.77 ± 0.96 | 23.80 ± 0.96 | ns |
| AP (IU/l) | 87.58 ± 4.09 | 74.61 ± 4.07 | <0.05 |

C = confined; FR = free range; LDH = lactate dehydrogenase; BUN = blood urea nitrogen; AST = aspartate aminotransferase; ALT = alanine aminotransferase; AP = alkaline phosphatase.

**Table 9** *Performance infra vitam and post-slaughter measurements (means ± s.e.) of Podolian young bulls in experiment 1 (2007)*

| | RS | | |
|---|---|---|---|
| | C | FR | *P* |
| Initial weight (kg) | 400.00 ± 7.67 | 383.33 ± 7.67 | ns |
| Final weight (kg) | 662.75 ± 20.30 | 599.47 ± 20.30 | <0.05 |
| Average daily gain (kg) | 1.35 ± 0.10 | 1.05 ± 0.10 | <0.05 |
| BCS | 5.45 ± 0.08 | 5.02 ± 0.08 | <0.01 |
| Dressing percentage | 54.93 ± 0.99 | 53.32 ± 0.99 | ns |
| pH 1 | 6.35 ± 0.10 | 6.61 ± 0.10 | ns |
| pH 24 | 5.30 ± 0.12 | 5.60 ± 0.12 | ns |
| *L** | 34.81 ± 0.25 | 33.94 ± 0.25 | <0.05 |
| *a** | 22.09 ± 0.26 | 22.50 ± 0.26 | ns |
| *b** | 2.04 ± 0.12 | 1.95 ± 0.12 | ns |

RS = rearing system; C = confined; FR = free range; BCS = body condition score; *L** = lightness; *a** = redness; *b** = yellowness.



https://doi.org/10.1017/S1751731110002685 Published online by Cambridge University Press

**Table 10** *Performance infra vitam and post-slaughter measurements (means ± s.e.) of Podolian young bulls in experiment 2 (2008)*

| | RS | | |
|---|---|---|---|
| | C | FR | *P* |
| Initial weight (kg) | 317.09 ± 17.91 | 304.00 ± 19.62 | ns |
| Final weight (kg) | 458.17 ± 19.83 | 433.33 ± 19.83 | ns |
| Average daily gain (kg) | 0.94 ± 0.05 | 0.86 ± 0.06 | ns |
| BCS | 5.00 ± 0.01 | 5.00 ± 0.01 | ns |
| Dressing percentage | 58.80 ± 2.58 | 54.35 ± 2.58 | ns |
| pH 1 | 6.29 ± 0.05 | 6.74 ± 0.05 | <0.001 |
| pH 24 | 5.48 ± 0.13 | 5.70 ± 0.13 | ns |
| *L** | 36.98 ± 1.20 | 32.74 ± 1.20 | <0.05 |
| *a** | 20.94 ± 0.82 | 21.41 ± 0.82 | ns |
| *b** | 8.60 ± 0.83 | 3.03 ± 0.83 | <0.001 |

RS = rearing system; C = confined; FR = free range; BCS = body condition score; *L** = lightness; *a** = redness; *b** = yellowness.

higher for 18-month-old bulls kept on pasture (0.96) than 15-month-old bulls kept in the same conditions (0.46). This result can be attributed both to the length of the finishing period (7 *v.* 5 months) and to the fact that younger bulls were slaughtered before the dry season pressed to increase the amount of supplement to be offered to the animals. Thus, this system was proportionally relying more on pasture as compared with 18-month-old bulls. Although we did not compare the two systems (15- *v.* 18-month-old bulls) from an economic point of view, the lower amount of concentrate fed to the animals suggests that FR 15-month-old bulls may be more profitable, in terms of feed and labour, than FR 18-month-old bulls.

*Post-slaughter measurements*

In both experiments no significant differences were detected between the two rearing systems for dressing percentage (Tables 9 and 10). pH values were not affected by rearing system in the first experiment (Table 9), thus indicating similar glycogenolysis post-slaughter in animals from both groups. In the second experiment, meat pH 1 h after slaughter was higher in group FR than group C group ($P < 0.05$), whereas no significant differences were observed between the two groups for ultimate pH (Table 10). The higher early *post-mortem* pH of FR bulls may be indicative of slower muscle metabolism immediately before and/or after slaughter. This may be indicative of lower stress (Terlouw *et al.*, 2009; Bourguet *et al.*, 2010), or of higher proportions of oxidative fibres in these animals (Gentry *et al.*, 2004). The increased redness (*a**) observed in muscles of outdoor raised cattle may be due to the increased proportions of oxidative muscle fibres induced by greater exercise (Vestergaard *et al.*, 2000). In addition, redness is negatively correlated with early *post-mortem* pH (Lindahl *et al.*, 2006).

In both experiments, FR significantly affected *L** index ($P < 0.05$; Tables 9 and 10) with higher values in meat from group C. In the second experiment, the meat produced by FR animals had a *b** value ($P < 0.001$; Table 10) lower than group C. These results are in agreement with Huuskonen *et al.* (2010) who found higher *L** and *b** values in meat from grass silage-finished bulls compared with meat from grazed-finished animals. Lindahl *et al.* (2006) found that low initial pH increased both the *L** and *b** values in pork meat. Yellowness (*b**) value generally varies in the same direction of *L** (Franck *et al.*, 2000). According to these authors, the higher *L** and *b** values could be explained by a less reductive environment that promotes oxygenation of myoglobin to $MbO_2$ (Lindahl *et al.*, 2001).

## Conclusion

The activity budget of FR bulls was characterized by high locomotor and feeding activities. These results along with the immune indicators suggest that bulls can benefit from an extensive rearing system based on pasture. Pasture availability, in terms of herbage mass and composition, as well as its seasonal development, was similar in the 2 years. Blood parameters indicated that adequate feeding supplementation should be provided in summer when plant physiological stage adversely affects diet quality and rate of ingestion, thus inhibiting growth and performances of 18-month-old bulls. In the second experiment, when the animals were slaughtered at an earlier age (15 months), the FR system produced results similar to those obtained in confinement. However, in both experiments minor negative effects of FR were detected on meat lightness (*L**). In addition the system based on FR 15-month-old bulls can be considered more sustainable in terms of resource sufficiency, as a lower amount of concentrate, per kilogram of meat was needed.

Although the number of animals per group was small and the nature of the production system was specific, the present results can serve as an indicative example for Mediterranean beef cattle production enterprises based on pasture utilization.

## Acknowledgements

The authors thank G. Migliori and A. M. Riviezzi for their expert technical assistance. They also thank the two referees for improving the quality of the manuscript. This research was funded by the Italian Ministry of Agriculture (TECOPA) and Regione Marche (EQUIZOOBIO).

## References

Association of Official Analytical Chemists (AOAC) 1990. Official methods of analysis, 15th edition. AOAC, Arlington, VA, USA.

Bergeron R, Badnell-Waters AJ, Lambton S and Mason GJ 2006. Stereotypic oral behaviour in captive ungulates: foraging, diet and gastrointestinal function. In Stereotypic behaviour: fundamentals and applications to welfare (ed. G Mason), pp. 19–57. CAB International, Wallingford, UK.

Bouissou MF and Boissy A 2005. The social behaviour of cattle and its consequences on breeding. Productions Animales 18, 87–99.

Bouissou MF, Boissy A, Le Neindre P and Veissier I 2001. The social behaviour of cattle. In Social behaviour in farm animals (ed. LJ Keeling and HW Gonyou), pp. 113–144. CABI Publishing, CAB International, Wallingford, UK.

Bourguet C, Deiss V, Gobert M, Durand D, Boissy A and Terlouw EMC 2010. Characterising the emotional reactivity of cows to understand and predict their



https://doi.org/10.1017/S1751731110002685 Published online by Cambridge University Press

stress reactions to the slaughter procedure. Applied Animal Behaviour Science 125, 9–21.

Boval M, Fanchone A, Archimede H and Gibb MJ 2007. Effect of structure of a tropical pasture on ingestive behaviour, digestibility of diet and daily intake by grazing cattle. Grass Forage Science 62, 44–54.

Braghieri A, Pacelli C, Verdone M, Girolami A and Napolitano F 2007. Effect of grazing and homeopathy on milk production and immunity of Merino derived ewes. Small Ruminant Research 69, 95–102.

Burns JC and Sollenberger LE 2002. Grazing behaviour of ruminants and daily performance from warm-season grasses. Crop Science 47, 873–881.

Cabaraux JF, Dufrasne I, Istasse L and Hornick JL 2005. Variation of plasma parameters and nitrogen metabolism in finishing Belgian Blue double-muscled cull females. Journal of Animal Physiology Animal Nutrition 89, 55–62.

Cosgrove GP 1997. Grazing behaviour and forage intake. In International Symposium on Animal Production under Grazing, Departmento de Zootechnia, Universite Federal de Vicosa, Brazil, pp. 59–80.

Cummins KA and Brunner CJ 1991. Effect of calf housing on plasma ascorbate and endocrine and immune function. Journal of Dairy Science 74, 1582–1588.

De Rosa G, Fedele V, Napolitano F, Gubitosi L, Bordi A and Rubino R 1997. Dietary preferences in adult and juvenile goats. Animal Science 65, 457–463.

De Rosa G, Napolitano F, Saltalamacchia F, Bilancione A, Sabia E, Grasso F and Bordi A 2007. The effect of rearing system on behavioural and immune responses of buffalo heifers. Italian Journal of Animal Science 6 (suppl. 2), 1260–1263.

Doornenbal H, Tong AKW and Murray NL 1988. Reference values of blood parameters in beef cattle of different ages and stages of lactation. Canadian Journal of Veterinary Research 52, 99–105.

Dumont B, Garel JP, Ginane C, Decuq F, Farruggia A, Pradel P, Rigolot C and Petit M 2007. Effect of cattle grazing a species-rich mountain pasture under different stocking rates on the dynamics of diet selection and sward structure. Animal 1, 1042–1052.

EU Directive 93/119/CE 1993. On the protection of animals at the time of slaughter or killing. Official Journal L340, 21–34.

Fisher AD, Crowe MA, O'Kiely P and Enright WJ 1997a. Growth, behaviour, adrenal and immune responses of finishing beef heifers housed on slatted floors at 1.5, 2.0, 2.5 or 3.0 $m^2$ space allowance. Livestock Production Science 51, 245–254.

Fisher AD, Crowe MA, Prendiville DJ and Enright WJ 1997b. Indoor space allowance: effects on growth, behaviour, adrenal and immune responses of finishing beef heifers. Animal Science 64, 53–62.

Franck M, Figwer P and Poirel MT 2000. Efft du type genetique et du temps d'attente sur la frequence du carctere destructure de la viande de porc. 8eme Journees des Sciences du Muscle et tecnologies de la viande, Paris, pp. 51–53.

Freedland RA and Szepesi B 1971. Control of enzyme activity: nutritional factors. In Enzyme synthesis and degradation in mammalian systems (ed. M Recheigkl Jr) University Park Press, Baltimore, USA.

Gentry JG, McGlone JJ, Miller MF and Blanton JR Jr 2004. Environmental effects on pig performance, meat quality, and muscle characteristics. Journal of Animal Science 82, 209–217.

Gibb MJ, Huckle CA, Nuthall R and Rook AJ 1997. Effect of sward surface height on intake and grazing behaviour by lactating Holstein Friesian cows. Grass and Forage Science 52 (suppl. 3), 309–321.

Ginane C, Petit M and D'hour P 2003. How do grazing heifers choose between maturing reproductive and tall or short vegetative swards. Applied Animal Behaviour Science 83, 15–27.

Gordon IJ 1995. Animal-based techniques for grazing ecology research. Small Ruminant Research 16, 203–214.

Gordon LJ and Illius AW 1992. Foraging strategy: from monoculture to mosaic. In Progress in sheep and goat research (ed. AW Speedy), pp. 153–170. CAB International, Wallingford, UK.

Hemsworth PHH, Barnett JL, Beveridge L and Matthews LR 1995. The welfare of extensively managed dairy cattle: a review. Applied Animal Behaviour Science 42, 161–182.

Hernandez L, Barral H, Halffter G and Colon SS 1999. A note on the behaviour of feral cattle in the Chihuahan Deser of Mexico. Applied Animal Behaviour Science 63, 259–267.

Hessle A, Rutter M and Wallin K 2008. Effect of breed, season and pasture moisture gradient on foraging behaviour in cattle on semi-natural grasslands. Applied Animal Behaviour Science 111, 108–119.

Huber R, Baumung R, Wurzinger M, Semambo D, Mwai O and Winckler C 2008. Grazing, social and comfort behaviour of Ankole and crossbred heifers on pasture in south western Uganda. Applied Animal Behaviour Science 112, 223–234.

Huuskonen A, Jansson S, Honkavaara M, Tuomisto L, Kauppinen R and Joki-Tokola E 2010. Meat colour, fatty acid profile and carcass characteristics of Hereford bulls finished on grazed pasture or grass silage-based diets with similar concentrate allowance. Livestock Science 131, 125–129.

Illius AW and Gordon IJ 1987. The allometry of food intake in grazing ruminants. Journal of Animal Ecology 56, 989–999.

INRA 1988. Alimentation des bovins ovins & caprins (ed. R Jarrige). Institut National de la Recherche Agronomique, Paris, France.

Karn JF 2001. Phosphorus nutrition of grazing cattle: a review. Animal Feed Science Technology 89, 133–153.

Kempkens K and Boxberger J 1987. Locomotion of cattle in loose housing system. In Cattle housing systems, lameness and behaviour (ed. HK Wierenga and JD Peterse), pp. 107–118. Martinus Nijhoff, Dordrecht, The Netherlands.

Keyserlingk MAG, Rushen J, de Passillé AM and Weary DM 2009. Invited review: the welfare of dairy cattle – key concepts and the role of science. Journal of Dairy Science 92, 4101–4111.

Krohn CC 1994. Behaviour of dairy cows kept in extensive (loose housing/pasture) or intensive (tie stall) environments. III. Grooming, exploration and abnormal behaviour. Applied Animal Behaviour Science 42, 73–86.

Krohn CC, Munksgaard L and Jonasen B 1992. Behaviour of dairy cows kept in extensive (loose housing/pasture) or intensive (tie stall) environments, I. Experimental procedure, facilities, time budgets – diurnal and seasonal conditions. Applied Animal Behaviour Science 34, 37–47.

Lamoot I, Meert C and Hoffmann M 2005. Habitat use of ponies and cattle foraging together in a coastal dune area. Biological Conservation 122, 523–536.

Lindahl G, Lundstrom K and Tornberg E 2001. Contribution of pigment content, myoglobin forms and internal reflectance to the colour of pork loin and ham from pure breed pigs. Meat Science 59, 141–151.

Lindahl G, Henckel P, Karlsson AH and Andersen HJ 2006. Significance of early postmortem temperature and pH decline on colour characteristics of pork loin from different crossbreeds. Meat Science 72, 613–623.

Marino R, Braghieri A, Albenzio M, Caroprese M, Girolami A, Santillo A and Sevi A 2009. Effect of rearing system and of dietary protein level on leptin, growth, and carcass composition in young Podolian bulls. Journal of Animal Science 87, 3097–3104.

Martin P and Bateson P 2007. Measuring behaviour, 3rd edition. Cambridge University Press, Cambridge, UK.

Martiniello P, Gesualdo G, Sabia E, Terzano GM, Pacelli C and Berardo N 2007. Intensive rainfed and irrigated forage crop production for Mediterranean Italian Buffalo feeding. Italian Journal of Animal Science 6 (suppl. 2), 1226–1229.

Meijs JAC, Walters RJK and Keen A 1982. Sward methods. In Herbage intake handbook (ed. JD Leaver), pp. 11–36. British Grassland Society, Berkshire, UK.

Miller K and Wood-Gush DGM 1991. Some effects of housing on the social behaviour of dairy cows. Animal Production 53, 271–278.

Napolitano F, Caporale G, Carlucci A and Monteleone E 2007. Effect of information about animal welfare and product nutritional properties on acceptability of meat from Podolian cattle. Food Quality Preference 18, 305–312.

Napolitano F, Pacelli C, De Rosa G, Braghieri A and Girolami A 2005. Sustainability and welfare of Podolian cattle. Livestock Production Science 92, 323–331.

Redbo I and Nordblad A 1997. Stereotypies in heifers are affected by feeding regime. Applied Animal Behaviour Science 53, 193–202.

Richards MW, Spitzer JC and Warner MB 1986. Effect of varying levels of postpartum nutrition and body condition at calving on subsequent reproductive performance in beef cattle. Journal of Animal Science 62, 300–306.

Ruckebusch Y and Bueno L 1978. An analysis of ingestive behaviour and activity of cattle under field conditions. Applied Animal Ethology 4, 301–313.

Sarker AB and Holmes W 1974. The influence of supplementary feeding on the herbage intake and grazing behaviour of dry cows. Journal of the British Grassland Society 29, 141–143.



https://doi.org/10.1017/S1751731110002685 Published online by Cambridge University Press

Statistical Analysis Software (SAS) 1990. SAS/STAT user's guide, version 6, 4th edition. SAS Institute, Cary, NC, USA.

Suttle NF 2009. Mineral nutrition of livestock. CABI Publishing, Wallingford, Oxon, UK.

Terlouw C, Berne A and Astruc T 2009. Effect of rearing and slaughter conditions on behaviour, physiology and meat quality of Large White and Duroc-sired pigs. Livestock Science 122, 199–213.

Tucker JF and Hentges JF Jr 1983. Effects of breed and pasture location on blood serum components of beef cattle. Retrieved November 11, 2009, from http://www.animal.ufl.edu/extension/beef/pubs_beefreports_1983.shtml.

Van Soest PJ, Robertson DA and Lewis BA 1991. Methods for dietary fiber, neutral detergent fiber and non-starch polysaccharides in relation to animal nutrition. Journal of Dairy Science 74, 3583–3597.

Vestergaard M, Oksbjerg N and Henckel P 2000. Influence of feeding intensity, grazing and finishing feeding on muscle fibre characteristics and meat colour of semitendinosus, longissimus dorsi and supraspinatus muscles of young bulls. Meat Science 54, 177–185.

Wallis De Vries MF and Daleboudt C 1994. Foraging strategy of cattle in patchy grassland. Oecologia 100, 98–106.

Watts JM and Stookey JM 2000. Vocal behaviour in cattle: the animal's commentary on its biological processes and welfare. Applied Animal Behaviour Science 67, 15–33.

Wierenga HK 1984. The social behaviour of dairy cows: some differences between pasture and cubicle system. In Proceedings of International Congress on Applied Ethology (ed. J Unshelm, G van Putten and K Zeeb), pp. 135–138. Kiel, Germany.



https://doi.org/10.1017/S1751731110002685 Published online by Cambridge University Press