Proceedings of the Edinburgh Mathematical Society (2021) **64**, 298–316
doi:10.1017/S0013091521000158

SCATTERING FOR CRITICAL WAVE EQUATIONS WITH VARIABLE COEFFICIENTS

SHI-ZHUO LOOI AND MIHAI TOHANEANU

Department of Mathematics, University of Kentucky, Lexington, KY 40506, USA
(shizhuo.looi@uky.edu; mihai.tohaneanu@uky.edu)

(Received 2 June 2020; first published online 30 April 2021)

Abstract We prove that solutions to the quintic semilinear wave equation with variable coefficients in $\mathbb{R}^{1+3}$ scatter to a solution to the corresponding linear wave equation. The coefficients are small and decay as $|x| \to \infty$, but are allowed to be time dependent. The proof uses local energy decay estimates to establish the decay of the L^6 norm of the solution as $t \to \infty$.

Keywords: scattering; defocusing; variable coefficients; nonlinear wave equation; hyperbolic equation; power-type nonlinearity; power-type non-linearity; semilinear wave equations

2020 *Mathematics subject classification:* Primary 35L05; 35L71

1. Introduction

In Minkowski space, solutions of the equation $\Box u = |u|^{p-1}u$ with $\Box = -\partial_t^2 + \Delta$ have a conserved and positive-definite energy

$$E(t) = \int_{\mathbb{R}^3} \frac{1}{2}|\nabla u(t,x)|^2 + \frac{1}{p+1}|u(t,x)|^{p+1}\,\mathrm{d}x$$

and the scaling symmetry

$$u(t,x) \mapsto \lambda^{2/(1-p)} u\left(\frac{t}{\lambda}, \frac{x}{\lambda}\right).$$

In three dimensions, the exponent $p = 5$ is called the energy-critical exponent, because solutions of the equation have an energy that is invariant under the scaling symmetry.

For the Cauchy problem with initial data in the energy space $\dot{H}^1 \times L^2$, local well-posedness is proven for $1 < p \leq 5$ by Strichartz estimates. Global existence for small initial data is a straightforward adaptation of the proof of local existence. In addition, there is global existence for large initial data due to the existence of a blow-up criterion, which informally says that the energy cannot concentrate at any point in spacetime. Moreover, given any finite-energy initial data, there is a unique global solution with finite

© The Author(s), 2021. Published by Cambridge University Press on Behalf of The Edinburgh Mathematical Society

https://doi.org/10.1017/S0013091521000158 Published online by Cambridge University Press

energy lying in $L^4L^{12}([0,\infty)\times\mathbb{R}^3)$; these solutions are known as strong (or Shatah–Struwe) solutions. See [5–8, 13, 28–33, 36] for details and more. The results in [1, 2] then combine to prove scattering of solutions with finite-energy initial data using a profile decomposition, which describes the failure of a sequence of uniformly bounded solutions to the free wave equation to be compact in the sense of Strichartz estimates. A similar result holds for the focusing equation with energy below that of the ground state: see [16].

This paper considers the equation

$$\begin{cases} Pu(t,x) = u(t,x)^5(t,x) \in (0,\infty)\times\mathbb{R}^3, \quad P = \partial_\alpha g^{\alpha\beta}\partial_\beta \\ u[0] \in \dot{H}^1 \times L^2 \end{cases} \tag{1.1}$$

Global existence and uniqueness of strong solutions (lying in $C(\mathbb{R}_t,\dot{H}^1)\cap L^5_{\mathrm{loc}}L^{10}$) was shown in [9] in the stationary setting. A similar result for classical solutions in the non-stationary setting was shown in [20]. These results require minimal assumptions on the coefficients, as eliminating the blow-up scenario only requires local-in-time arguments.

Our main theorem establishes scattering of strong solutions to (1.1) for certain small, asymptotically flat perturbations of the Minkowski metric. To the authors' knowledge, this is the first such result for small perturbations of the Minkowski metric m with variable coefficients.

Definition 1.1 (scattering in the energy space). We say that the solution u to (1.1) *scatters in the energy space* if there exists $(f,g)\in\dot{H}^1\times L^2$ such that

$$\lim_{t\to\infty}\|u[t]-S(t,0)(f,g)\|_{\dot{H}^1\times L^2}=0.$$

Definition 1.2. We define $\underline{L} := \sum_{i=1}^3 (x^i/|x|)\partial_{x_i} - \partial_t$.

Theorem 1.3. *Let $g^{\alpha\beta}(t,x)$ be a Lorentzian metric, let $P=\partial_\alpha g^{\alpha\beta}\partial_\beta$, and let $h := g-m$ denote the perturbative terms of the metric g. The unique global strong solution to the Cauchy problem (1.1) scatters in the energy space $\dot{H}^1\times L^2$ provided that*

$$|h| \lesssim \epsilon \frac{\langle t-|x|\rangle^{1/2}}{\langle x\rangle^\gamma \langle t+|x|\rangle^{1/2}}, \tag{1.2}$$

$$|h^{\underline{LL}}| \lesssim \epsilon \frac{\langle t-|x|\rangle}{\langle x\rangle^\gamma \langle t+|x|\rangle}, \tag{1.3}$$

$$|\partial^J h| \lesssim \epsilon \frac{1}{\langle x\rangle^{|J|+\gamma}} \quad \text{for } |J|=1 \text{ and } |J|=2 \tag{1.4}$$

where $\gamma>0$ is an arbitrarily small constant and $\epsilon>0$ is a sufficiently small constant. In these assumptions, $\partial^J h$ denotes $\partial^J h^{\alpha\beta}$ for all multi-indices α and β, and $h^{\underline{LL}} = h^{\alpha\beta}\underline{L}_\alpha\underline{L}_\beta$, where we lower indices with respect to the Minkowski metric.

This says that the unique global solution of the nonlinear problem on small, asymptotically flat perturbations of Minkowski space that have appropriate decay at infinity behaves, in the asymptotic sense, like the solution to the linear homogeneous problem $Pu=0$, at least in the energy space.

https://doi.org/10.1017/S0013091521000158 Published online by Cambridge University Press

Remark 1.4. The assumptions (1.2), (1.3), and (1.4) are satisfied by metrics that arise as solutions to Einstein's Vacuum Equations when expressed in harmonic coordinates, see [22].

Remark 1.5. One of the key ingredients in our proof is the fact that Strichartz estimates for the linear problem hold. Assuming

$$|\partial^J h| \lesssim \epsilon \frac{1}{\langle x\rangle^{|J|+\gamma}} \quad \text{for } 0 \leq |J| \leq 2$$

this was proved by Metcalfe–Tataru [25]. Our assumptions are the same, except that we require more decay of h (but not its derivatives) near the light cone. This is due to the fact that we need to control certain boundary terms that appear when multiplying the equation by $t\partial_t + r\partial_r$. In particular, the extra decay requirement (1.3) is needed to control the term $|\underline{L}u|^2$ on the boundary, and geometrically it implies that the light cones of the perturbed metric are comparable to those of the Minkowski metric.

Remark 1.6. Theorem 1.3 also holds if we replace P by the geometric wave operator

$$\Box_g = \frac{1}{\sqrt{|g|}}\partial_\alpha \sqrt{|g|} g^{\alpha\beta}\partial_\beta, \quad |g| := |\det g^{\alpha\beta}|.$$

Indeed, in the estimates below, one integrates with respect to the volume form $\sqrt{|g|}dtdx$, and uses the fact that $\sqrt{|g|}g^{\alpha\beta} \approx g^{\alpha\beta}$. There are extra error terms of the form $\partial\sqrt{|g|}u^6$ arising, which can be absorbed since by (1.4) we have

$$\partial\sqrt{|g|} \lesssim \frac{\epsilon}{\langle x\rangle^{1+\gamma}}$$

Remark 1.7. A key tool in proving scattering on variable-coefficient backgrounds is local energy decay. Such an estimate was proven in [15, 27, 35] for Minkowski space and in [23, 24] for perturbations of Minkowski space and became a valuable tool in the study of both linear and nonlinear problems. In particular, they imply Strichartz estimates on certain variable-coefficient backgrounds, see [25]. Our result is one of several showing that local energy decay is fruitful for understanding the long-time behaviour and asymptotics of solutions to nonlinear dispersive equations on variable-coefficient backgrounds.

Remark 1.8. For the energy-critical problem on Minkowski space, global a priori estimates were proven in [28, 29], from which scattering for the wave and Klein–Gordon equations were deduced. Analogous results in the exterior of obstacles were obtained in [3, 4, 15, 34]. Scattering on Riemannian manifolds for a class of non-trapping obstacles close to the two-convex framework has been shown as well for the energy-critical nonlinear problem [19]. Profile decompositions akin to [1] have been shown for waves on hyperbolic space in [21]; a similar result was shown for $(\Box + a|x|^{-2})u = u^5$ in [26]. Finally, for the equation $(\Box + V(x))u = u^5$, scattering to steady states was shown in [11, 12].

For the nonlinear Schrödinger equation, the energy-critical problem for the defocusing quintic problem with initial data in the energy space for small, compactly supported perturbations of the Euclidean metric also exhibits scattering to linear solutions for

https://doi.org/10.1017/S0013091521000158 Published online by Cambridge University Press

all finite-energy data, as shown in [10]. There are many other known results for the energy-critical energy Schrödinger with potential, and for the exterior of a strictly convex obstacle, see [17, 18, 37], etc.

2. Notation and preliminaries

We fix the spatial dimension to be $d = 3$ and define $\Box = -\partial_t^2 + \Delta$ and $P = \partial_\alpha g^{\alpha\beta}\partial_\beta$ where $g = g(t,x)$ is a Lorentzian metric. We write either $X \lesssim Y$ or $X = O(Y)$ to indicate that

$$|X| \leq CY$$

(rather than $X \leq CY$) for some absolute constant C which may vary by line. Similarly, $X \approx Y$ means that there are constants $0 < C_1 < C_2$ so that

$$C_1|X| \leq |Y| \leq C_2|X|.$$

We let $\langle r\rangle = \langle x\rangle = (1+|x|^2)^{1/2}$. We write $\nabla = (\partial_t, \nabla_x)$ for the spacetime gradient. Throughout the paper, we use the Einstein summation convention, and we let Greek (respectively Latin) indices denote spacetime (respectively space) indices. We write $u[T] = (u(T,x),\ \partial_t u(T,x))$.

The energy of the solution u is defined to be

$$E(t) = \int_{\mathbb{R}^3} \frac{1}{2}|\nabla u(t,x)|^2 + \frac{1}{6}u(t,x)^6\,\mathrm{d}x.$$

We will also use the notation

$$E_K(t) := \int_K \frac{1}{2}|\nabla u(t,x)|^2 + \frac{1}{6}u(t,x)^6\,\mathrm{d}x$$

for some subset K of $\mathbb{R}^3$.

For any $(f,g) \in \dot{H}^1 \times L^2$, we denote by $S(t,s)(f,g)$ the unique solution $u \in C(\mathbb{R}_t, \dot{H}^1)$ with $\partial_t u \in C(\mathbb{R}_t, L^2)$ to the equation

$$\begin{cases} Pu = 0 \quad (t,x) \in (s,\infty)\times\mathbb{R}^3 \\ u[s] = (f,g) \end{cases} \tag{2.1}$$

Let

$$X = (C(\mathbb{R}_t, \dot{H}^1) \cap L^5_{\mathrm{loc}}L^{10}) \times C(\mathbb{R}_t, L^2)$$

and for any closed, finite interval

$$X(I) = (C(I, \dot{H}^1) \cap L^5(I)L^{10}) \times C(I, L^2)$$

Consider the Cauchy problem

$$\begin{cases} Pu(t,x) = u(t,x)^5 \quad (t,x) \in (0,\infty)\times\mathbb{R}^3 \\ u[0] \in \dot{H}^1 \times L^2 \end{cases} \tag{2.2}$$

https://doi.org/10.1017/S0013091521000158 Published online by Cambridge University Press

By Duhamel's formula, classical solutions to (2.2) satisfy

$$u(t) = S(t,0)u[0] + \int_0^t \frac{1}{g^{00}} S(t,s)(0,u^5(s))\,\mathrm{d}s \tag{2.3}$$

We can thus define a strong solution to be a solution of (2.3) so that $(u, \partial_t u)$ also lies in X.

The results of [9, 20] show that, for smooth initial data, there is a unique global classical solution to (2.2) that is also a strong solution. Moreover, this result is extended to initial data in the energy space in [9] for time-independent coefficients, and the same argument can be used to prove it in the time-dependent case. We will be interested in studying the asymptotic properties of the unique strong solution in the energy space, in particular, the fact that it approaches a solution to the linear equation in the energy space.

3. Energy conservation and local energy decay in Minkowski space

In order to motivate the discussion that follows, we devote this section toward certain key results for the linear problem in the setting of Minkowski space. Consider the Cauchy problem

$$\begin{cases} \Box u = 0 \quad (t,x) \in (0,\infty) \times \mathbb{R}^3 \\ u[0] \in \dot{H}^1 \times L^2 \end{cases} \tag{3.1}$$

The energy of the solution u to (3.1) is defined to be

$$E^{\mathrm{lin}}(t) = \int_{\mathbb{R}^3} \frac{1}{2} |\nabla u(t,x)|^2 \,\mathrm{d}x$$

and it is conserved: for all $T \geq 0$, $E^{\mathrm{lin}}(T) = E^{\mathrm{lin}}(0) = E$.

The solution u to (3.1) also satisfies the local energy estimate

$$\iint_{[0,T]\times\mathbb{R}^3} \frac{|\nabla u|^2}{\langle r\rangle^{1+\gamma}} + \frac{u^2}{\langle r\rangle^{3+\gamma}} \mathrm{d}x\,\mathrm{d}t \lesssim E$$

where $\gamma > 0$ is an arbitrarily small fixed constant. This estimate is proven by multiplying both sides of the equation by $a(r)u + b(r)\partial_r u + C\partial_t u$ in the region $[0,T] \times \mathbb{R}^3$, with:

$$b(r) = \sum_{j=0}^{\infty} 2^{-j\gamma} \frac{r}{r+2^j},$$

where for each j, this is a function catered to the region $r \approx 2^j$ with the factor $2^{-j\gamma}$, where the small number $\gamma > 0$ is introduced in order to obtain convergence of the series; $a(r) = b(r)/r$; and $C > 0$ is a constant chosen to be sufficiently large.

https://doi.org/10.1017/S0013091521000158 Published online by Cambridge University Press

More precisely, one obtains

$$\begin{aligned}&\iint_{[0,T]\times\mathbb{R}^3} \Box u b(r)\partial_r u \,\mathrm{d}x\,\mathrm{d}t\\ &\quad= -\iint b'(\partial_r u)^2 + \frac{b}{r}|\partial_\omega u|^2 + \frac{1}{2}\left(b' + 2\frac{b}{r}\right)(u_t^2 - |\nabla_x u|^2)\,\mathrm{d}x\,\mathrm{d}t\\ &\qquad+ \int_{\mathbb{R}^3} -b\partial_r u\partial_t u|_0^T \,\mathrm{d}x,\end{aligned}$$

where $|\partial_\omega u|^2 := |\nabla_x u|^2 - |\partial_r u|^2$, and

$$\iint_{[0,T]\times\mathbb{R}^3} \Box u a(r) u \,\mathrm{d}x\,\mathrm{d}t = \iint a(u_t^2 - |\nabla_x u|^2) + \frac{1}{2}\Delta a u^2 \,\mathrm{d}x\,\mathrm{d}t + \frac{1}{2}\int_{\mathbb{R}^3} -au\partial_t u|_0^T \,\mathrm{d}x.$$

Since $a = O(\langle r\rangle^{-1})$ and $b = O(1)$, Hardy's inequality $\int_{\mathbb{R}^3} u^2/r^2 \,\mathrm{d}x \lesssim \int_{\mathbb{R}^3} |\nabla_x u|^2 \,\mathrm{d}x$ shows that there exists a sufficiently large constant $C > 0$ such that

$$\int_{\{t\}\times\mathbb{R}^3} b\partial_t u\partial_r u + a\partial_t u u + C|\nabla u|^2 \,\mathrm{d}x \approx E^{\mathrm{lin}}(t)$$

for all $t \geq 0$. We obtain

$$E^{\mathrm{lin}}(T) + \iint_{[0,T]\times\mathbb{R}^3} \frac{1}{2}b'(u_r^2 + u_t^2) - \left(\frac{1}{2}b' - \frac{b}{r}\right)|\partial_\omega u|^2 - \frac{\Delta a}{2}u^2 \,\mathrm{d}x\,\mathrm{d}t \lesssim E^{\mathrm{lin}}(0)$$

One can check directly that

$$b' \gtrsim \langle r\rangle^{-1-\gamma}, \quad b/r - \frac{1}{2}b' \gtrsim \langle r\rangle^{-1-\gamma}, \quad -\Delta a \gtrsim \langle r\rangle^{-3-\gamma}$$

and thus

$$\frac{1}{2}b'(u_r^2 + u_t^2) + \left(-\frac{1}{2}b' + \frac{b}{r}\right)|\partial_\omega u|^2 - \frac{\Delta a}{2}u^2 \gtrsim \frac{|\nabla u|^2}{\langle r\rangle^{1+\gamma}} + \frac{u^2}{\langle r\rangle^{3+\gamma}} \tag{3.2}$$

which finishes the proof.

4. Uniform energy bounds and local energy decay for the nonlinear problem on perturbations

We now come to certain key tools, analogous to the results presented in Section 3, that will be used in the proof of scattering for the nonlinear problem on certain perturbations of Minkowski space that have appropriate decay at infinity.

If $g(t,x)$ is a non-stationary metric that satisfies certain decay conditions, and u is a solution of the Cauchy problem (2.2), then we have uniform energy bounds: with the

https://doi.org/10.1017/S0013091521000158 Published online by Cambridge University Press

energy now defined to be

$$E(t) := \int_{\{t\}\times\mathbb{R}^3} \frac{1}{2}|\nabla u|^2 + \frac{1}{6}|u|^6 \,\mathrm{d}x,$$

if g and its derivatives satisfy certain decay conditions, then

$$E(T) \lesssim E := E(0)$$

for some implicit constant that is independent of T. In fact, we may prove local energy decay and uniform energy bounds in one fell swoop for (2.2), as the following proposition shows.

Theorem 4.1 (integrated local energy decay for the nonlinear Cauchy problem). *Let u be a solution of (2.2) and let $|J| \leq 1$ be a multi-index. If $\partial^J h^{\alpha\beta} \lesssim \epsilon\langle r\rangle^{-|J|-\gamma}$ where $\gamma > 0$ is an arbitrarily small constant and $\epsilon > 0$ is a sufficiently small constant then*

$$\|u\|^2_{LE^1[T_1,T_2]} + E(T_2) \lesssim E(T_1) \tag{4.1}$$

for some implicit constant that is independent of T_1 and T_2, where

$$\|u\|^2_{LE^1[T_1,T_2]} = \iint_{[T_1,T_2]\times\mathbb{R}^3} \frac{|\nabla u|^2}{\langle r\rangle^{1+\gamma}} + \frac{u^2}{\langle r\rangle^{3+\gamma}} + \frac{|u|^6}{\langle r\rangle}\mathrm{d}x\,\mathrm{d}t$$

Let us first assume that u is a classical solution to the equation

$$Pu = u^5 + F$$

Following [24] and the discussion in the previous section, we multiply the equation by $a(r)u + b(r)\partial_r u + C\partial_t u$, with

$$b(r) = \sum_{j=0}^{\infty} 2^{-j\gamma}\frac{r}{r+2^j}, \quad a(r) = b(r)/r.$$

Upon integrating by parts, we obtain

$$E(T_2) + \iint_{[T_1,T_2]\times\mathbb{R}^3} \frac{1}{2}b'(u_r^2+u_t^2) - \left(\frac{1}{2}b' - \frac{b}{r}\right)|\partial_\omega u|^2 - \frac{\Delta a}{2}u^2 + Err$$
$$+ \left(\frac{2}{3}a(r) - \frac{1}{6}b'(r)\right)u^6 \,\mathrm{d}x\,\mathrm{d}t \lesssim E(T_1) + \iint_{[T_1,T_2]\times\mathbb{R}^3} |F|\left(|\nabla u| + \frac{|u|}{\langle r\rangle}\right)\mathrm{d}x\,\mathrm{d}t.$$

where the error satisfies

$$\mathrm{Err} \lesssim \left(\frac{|h|}{\langle x\rangle} + |\nabla h|\right)\left(|\nabla u|^2 + |\nabla u|\frac{|u|}{\langle x\rangle}\right)$$

Since $\partial^J h^{\alpha\beta} \lesssim \epsilon\langle r\rangle^{-|J|-\gamma}$, we can estimate by Cauchy–Schwarz

$$\mathrm{Err} \lesssim \epsilon\left(\frac{|\nabla u|^2}{\langle r\rangle^{1+\gamma}} + \frac{u^2}{\langle r\rangle^{3+\gamma}}\right)$$

https://doi.org/10.1017/S0013091521000158 Published online by Cambridge University Press

Moreover,

$$\frac{2}{3}a(r) - \frac{1}{6}b'(r) = \sum_{j=0}^{\infty} 2^{-j\gamma}\left(\frac{1}{2}\frac{1}{r+2^j} + \frac{1}{6}\frac{r}{(r+2^j)^2}\right) \gtrsim \frac{1}{\langle r\rangle}$$

Taking (3.2) into account, and applying Hölder and Hardy to control the inhomogeneity, we get

$$\|u\|^2_{LE^1[T_1,T_2]} + E(T_2) \lesssim E(T_1) + \|F\|_{L^1[T_1,T_2]L^2}\|\nabla u\|_{L^\infty[T_1,T_2]L^2} \tag{4.2}$$

Consider now a strong solution u, and a sequence of classical solutions u_n so that $u_n(T_1) \to u(T_1)$ in the energy norm. After dividing the interval $I = [T_1, T_2]$ into finitely many intervals so that the L^5L^{10} norm of u is suitably small on each interval, a contraction argument shows that $u_n \to u$ in $X(I)$. In particular, this implies that u_n^5 is a Cauchy sequence in $L^1[T_1, T_2]L^2$, and thus by (4.2) we must have $u_n \to u$ in $LE^1[T_1, T_2]$. The desired conclusion (4.1) now follows.

5. L^6 norm decay of solutions in Minkowski space

In order to motivate the next section, which contains the main result and its proof, in this section, we shall present the highlights of the proof of L^6 norm decay in Minkowski space for the nonlinear problem, as done in [2].

5.1. More notation

First, we fix some notation which will be used for the rest of the paper. Let

$$\Gamma = \{(t, x) : |x| - c < t, t > 0\}$$

be a forward solid light cone with $c \geq 0$ to be determined and let $\Gamma(I) = \Gamma \cap (I \times \mathbb{R}^3)$ where $I \subset [0, \infty)$ is a time interval. Let $D(T) = \{(t, x) : t = T,\ |x| - c < t\}$ denote its $t = T$ slices and let $L(I) = \{(t, x) : t \in I,\ |x| - c = t\}$ denote the lateral boundary of $\Gamma(I)$ with

$$L_c(I) := L(I)$$

also used for emphasis, but usually we shall simply write $L(I)$.

Consider next the Cauchy problem

$$\begin{cases} \Box u = u^5 \quad (t, x) \in (0, \infty) \times \mathbb{R}^3 \\ u[0] \in \dot{H}^1 \times L^2 \end{cases} \tag{5.1}$$

We now sketch a proof of the L^6 norm decay in the energy space for solutions to (5.1) (see [1, 33] for more details). We will adapt this proof to the variable-coefficient case in the next section.

Given $\delta > 0$, pick c sufficiently large so that the energy in the exterior region $|x| > c$ is less than $\delta/2$. Now given any $c \geq 0$, the flux on the time interval I is defined to be

https://doi.org/10.1017/S0013091521000158 Published online by Cambridge University Press

the integral on the lateral boundary arising from multiplying the equation $\Box u = u^5$ by $\partial_t u$, namely

$$\text{flux}(I) = \int_{L(I)} \frac{1}{2}|\partial_\omega u|^2 + \frac{1}{2}|\partial_t u + \partial_r u|^2 + \frac{1}{6}u^6 \frac{\mathrm{d}\sigma}{\sqrt{2}}$$

It is clear that the flux is non-negative. As the upper and lower limits of I approach infinity, the flux decays, as an application of the divergence theorem in the interior of the $\Gamma(I)$ region shows. More precisely, if $I = [T_1, T_2]$, then in Minkowski space, one obtains for the arbitrary number $c \geq 0$ chosen

$$E_{|x|<c+T_2}(T_2) = E_{|x|<c+T_1}(T_1) + \text{flux}([T_1, T_2]).$$

Thus $E_{|x|<c+t}(t)$ is monotone non-decreasing; moreover, it is bounded; therefore it converges to a limit as $t \to \infty$, as claimed. In particular, for all T_2 such that $T_2 > T_1$, and any $c \geq 0$,

$$\lim_{T_1 \to \infty} \lim_{T_2 \to \infty} \text{flux}([T_1, T_2]) = 0.$$

We now multiply (5.1) by

$$Xu := (t+c)\partial_t u + x^i \partial_i u + u = Su + c\partial_t u + u$$

where $S := t\partial_t + \sum_{i=1}^3 x^i \partial_i u$ and apply the divergence theorem in $\Gamma(I)$. We obtain

$$P(T_2) + \iint_{\Gamma(I)} \frac{u^6}{3} \mathrm{d}x\,\mathrm{d}t = P(T_1) + \int_{L(I)} (t+c)\left(\frac{Xu}{t+c}\right)^2 \frac{\mathrm{d}\sigma}{\sqrt{2}} \tag{5.2}$$

where

$$P(T) := \int_{D(T)} \frac{t+c}{2}\left[\left(\frac{Xu}{t+c}\right)^2 + \left(|\nabla_x u|^2 - \left(\frac{x}{t+c}\cdot\nabla_x u\right)^2\right)\right] + \frac{u^2}{t+c} + \frac{t+c}{6}u^6\,\mathrm{d}x$$

Recall that on $L(I)$ we have $r = t + c$, enabling us to write $Xu/(t+c) = \partial_t u + \partial_r u + u/(t+c)$. By Cauchy–Schwarz and Hölder, we obtain

$$\begin{aligned}\int_{L(I)} (t+c)\left(\frac{Xu}{t+c}\right)^2 \frac{\mathrm{d}\sigma}{\sqrt{2}} &\lesssim (T_2+c)\int_{L(I)} (\partial_t u + \partial_r u)^2\,\mathrm{d}\sigma + \int \frac{u^2}{t+c}\mathrm{d}\sigma \\ &\lesssim (T_2+c)(\|(\partial_t+\partial_r)u\|^2_{L^2(L(I))} + \|u\|^2_{L^6(L(I))})\end{aligned}$$

In summary,

$$\begin{aligned}T_2 \int_{D(T_2)} u^6\,\mathrm{d}x &\lesssim P(T_2) + \iint_{\Gamma(I)} u^6\,\mathrm{d}x\,\mathrm{d}t \lesssim P(T_1) + (T_2+c)G(\text{flux}([T_1,T_2])) \\ &\lesssim (T_1+c)E_{|x|<T_1+c}(T_1) + (T_2+c)G(\text{flux}([T_1,T_2]))\end{aligned}$$

and

$$G(\theta) := \theta + \theta^{1/3}$$

https://doi.org/10.1017/S0013091521000158 Published online by Cambridge University Press

is a function which decays to zero as its argument decays to zero. Take $T_1 = \delta T_2$ to see that, since δ was arbitrary and the flux decays,

$$\limsup_{t\to\infty} \|u(t,\cdot)\|_{L^6(\mathbb{R}^3)} = 0.$$

6. L^6 norm decay and scattering of solutions on small asymptotically flat perturbations of Minkowski space

We now come to the main result and its proof.

Theorem 6.1 (main theorem). *Let u be the unique global strong solution of (2.2).*

(1) We make the following assumptions on the perturbation h:

$$|\partial h| \lesssim \epsilon \langle x\rangle^{-1-\gamma} \tag{6.1}$$

$$|h| \lesssim \epsilon \frac{\langle t-r\rangle^{1/2}}{\langle x\rangle^{\gamma}\langle t+r\rangle^{1/2}} \tag{6.2}$$

$$|h^{\underline{L}\underline{L}}| \lesssim \epsilon \frac{\langle t-r\rangle}{\langle x\rangle^{\gamma}\langle t+r\rangle} \tag{6.3}$$

where $\gamma > 0$ is an arbitrarily small constant and $\epsilon > 0$ is a sufficiently small constant.
Then

$$\limsup_{t\to\infty} \|u(t,\cdot)\|_{L^6(\mathbb{R}^3)} = 0. \tag{6.4}$$

(2) If in addition

$$|\partial^J h| \lesssim \epsilon \langle x\rangle^{-2-\gamma}, \quad |J| = 2, \tag{6.5}$$

then u scatters in the energy space.

Recall that we define the normal derivative to the cone

$$\underline{L} = \frac{x^i}{|x|}\partial_i - \partial_0 = \frac{x^i}{|x|}\partial_i - \partial_t$$

and

$$h^{\underline{L}\underline{L}} = h^{\alpha\beta}\underline{L}_\alpha \underline{L}_\beta = h^{00} - 2\sum_i h^{0i}\frac{x^i}{|x|} + \sum_{i,j} h^{ij}\frac{x^i x^j}{|x|^2}.$$

We also remark that the decay rates on h and ∂h are consistent with the ones required for local energy decay Theorem 4.1 except near the cone $t \approx |x|$, where we need better decay rates to close the argument.

We now sketch the proof of the main theorem. Let us first assume that u is a classical solution to the equation

$$Pu = u^5 + F \tag{6.6}$$

https://doi.org/10.1017/S0013091521000158 Published online by Cambridge University Press

The main estimate of the paper is the following:

Proposition 6.2. *If u solves (6.6), and for any R, T_1 and T_2 so that $R \geq 0$, $1 < T_1$*[*], *and $T_1 + 2R < T_2$, we have*

$$\int_{\mathbb{R}^3} u^6(T_2, x)\,\mathrm{d}x \lesssim \frac{T_1 + 2R}{T_2} E_{\{|x|T_1+R\}}(T_1) + \|u\|^2_{LE^1[T_1,T_2]} + \|F\|_{L^1[T_1,T_2]L^2}\|\nabla u\|_{L^\infty[T_1,T_2]L^2}) \tag{6.7}$$

where $E := E(0)$.

We now make the key observation that, unlike in the case of Minkowski, flux(I) is not necessarily non-negative on arbitrary light cones. Instead, by averaging, we obtain that there is $c \in [R, 2R]$ so that

$$\int_{L_c([T_1,T_2])} \frac{|\nabla u|^2}{\langle x\rangle^{1+\gamma}}\mathrm{d}\sigma \lesssim R^{-1}\|u\|^2_{LE^1[T_1,T_2]}. \tag{6.8}$$

For the rest of this proof, fix c as above. Note that the hypothesis $T_1 + 2R < T_2$ implies that $T_2 \approx T_2 + c$. Moreover, in this case, c depends on T_1, so we make sure that we carefully track the dependence on c in our estimates. In fact, the implicit constants do not depend on c, T_1, T_2, or R.

Proposition 6.2 follows from the results of §§ 6.1 and 6.2. We then finish the proof of Theorem 6.1 in § 6.3.

6.1. L^6 norm decay of solutions on spacelike slices exterior to the cone

The next lemma shows that we can control both the outside energy and the flux through L_c. Note that, unlike in the Minkowski case, it is not clear that this can be done for all c.

Lemma 6.3. *Let u solve (6.6). Then*

$$\begin{aligned} E_{\{|x|>T_2+c\}}(T_2) + \mathit{flux}([T_1,T_2]) &\lesssim E_{\{|x|>T_1+c\}}(T_1) + \|u\|^2_{LE^1[T_1,T_2]} \\ &+ \|F\|_{L^1[T_1,T_2]L^2}\|\nabla u\|_{L^\infty[T_1,T_2]L^2}. \end{aligned} \tag{6.9}$$

Here,

$$\mathit{flux}([T_1,T_2]) := \int_{L_c([T_1,T_2])} \frac{1}{2}|\bar\partial u|^2 + \frac{1}{6}u^6\frac{\mathrm{d}\sigma}{\sqrt{2}}$$

and

$$\bar\partial u := \{Lu, (r\sin\phi)^{-1}\partial_\theta u, r^{-1}\partial_\phi u\}, \quad L = \frac{x^i}{|x|}\partial_i + \partial_t$$

denote the tangential derivatives of u to the light cone.

[*] We shall only be interested in certain sufficiently large values of T_1 and R.

https://doi.org/10.1017/S0013091521000158 Published online by Cambridge University Press

Proof. Let $I = [T_1, T_2]$. Multiplying both sides of the equation in (6.6) by $\partial_t u$, we obtain the identity

$$\partial_\alpha(g^{\alpha\beta}\partial_\beta u\partial_t u) - \tfrac{1}{2}\partial_t(g^{\alpha\beta}\partial_\beta u\partial_\alpha u) + \tfrac{1}{2}\partial_t g^{\alpha\beta}\partial_\beta u\partial_\alpha u = \tfrac{1}{6}\partial_t(u^6) + F\partial_t u.$$

Define

$$\Gamma^{\mathrm{ext}}(I) := \{T_1 \le t \le T_2, |x| > t + c\}, \quad D(T)^c := \Gamma^{\mathrm{ext}}(I) \cap \{t = T\}$$

Applying the divergence theorem within the region $\Gamma^{\mathrm{ext}}(I)$ leads to

$$\begin{aligned}\iint_{\Gamma^{\mathrm{ext}}(I)} F\partial_t u \,\mathrm{d}x\,\mathrm{d}t = \iint_{\Gamma^{\mathrm{ext}}(I)} \frac{1}{2}\partial_t g^{\alpha\beta}\partial_\beta u\partial_\alpha u \,\mathrm{d}x\,\mathrm{d}t \\ + \int_{\partial\Gamma^{\mathrm{ext}}(I)} \nu_\alpha g^{\alpha\beta}\partial_\beta u\partial_t u - \frac{1}{2}\nu_0 g^{\alpha\beta}\partial_\beta u\partial_\alpha u - \frac{1}{6}\nu_0 u^6 \,\mathrm{d}\sigma. \quad (6.10)\end{aligned}$$

Next, let BD_h denote the part of the energy density on the boundary of $\Gamma^{\mathrm{ext}}(I)$ arising from h

$$\mathrm{BD}_h := \nu_\alpha h^{\alpha\beta}\partial_\beta u\partial_t u - \tfrac{1}{2}\nu_0 h^{\alpha\beta}\partial_\beta u\partial_\alpha u.$$

Note that BD_h depends on the domain of integration. Expanding (6.10), we have

$$\begin{aligned}E_{\{|x|>T_2+c\}}(T_2) + \int_{D(T_2)^c} \mathrm{BD}_h \,\mathrm{d}x + \mathrm{flux}([T_1, T_2]) + \iint_{\Gamma^{\mathrm{ext}}(I)} \frac{1}{2}\partial_t h^{\alpha\beta}\partial_\alpha u\partial_\beta u \,\mathrm{d}x\,\mathrm{d}t \\ = \iint_{\Gamma^{\mathrm{ext}}(I)} F\partial_t u \,\mathrm{d}x\,\mathrm{d}t + E_{\{|x|>T_1+c\}}(T_1) + \int_{D(T_1)^c} \mathrm{BD}_h \,\mathrm{d}x + \int_{L_c(I)} \mathrm{BD}_h \,\mathrm{d}\sigma \quad (6.11)\end{aligned}$$

The space-time term is easy to estimate by (6.1)

$$\iint_{\Gamma^{\mathrm{ext}}(I)} \frac{1}{2}\partial_t h^{\alpha\beta}\partial_\alpha u\partial_\beta u \,\mathrm{d}x\,\mathrm{d}t \lesssim \iint_{\Gamma^{\mathrm{ext}}(I)} \frac{|\nabla u|^2}{\langle x\rangle^{1+\gamma}}\mathrm{d}x\,\mathrm{d}t \le \|u\|^2_{LE^1[T_1,T_2]}. \quad (6.12)$$

Similarly, using that $|h| \lesssim \epsilon$, we obtain

$$\int_{D(T_j)^c} \mathrm{BD}_h \,\mathrm{d}x \lesssim \epsilon E_{\{|x|>T_j+c\}}(T_j), \quad j = 1, 2. \quad (6.13)$$

which can be absorbed in $E_{\{|x|>T_j+c\}}(T_j)$ for small enough ϵ.

https://doi.org/10.1017/S0013091521000158 Published online by Cambridge University Press

Finally, we need to estimate the perturbative error term on the lateral boundary; this is where we will use (6.2) and (6.3). We write

$$\nu_\alpha h^{\alpha\beta}\partial_\beta = -\tfrac{1}{2}h^{\underline{L}\underline{L}}\underline{L} + O(h)\bar{\partial}$$
$$\partial_t = \tfrac{1}{2}(L - \underline{L})$$
$$h^{\alpha\beta}\partial_\alpha u\partial_\beta u = \tfrac{1}{4}h^{\underline{L}\underline{L}}(\underline{L}u)^2 + O(h)\bar{\partial}u\partial u$$

Note that, due to (6.3) and (6.2), we have that on $L(I)$

$$h^{\underline{L}\underline{L}} \lesssim \frac{R}{\langle x\rangle^{1+\gamma}}, \quad h \lesssim \epsilon\frac{R^{1/2}}{\langle x\rangle^{1/2+\gamma}} \tag{6.14}$$

and thus by Cauchy–Schwarz

$$\begin{aligned}\int_{L_c(I)} \mathrm{BD}_h \,\mathrm{d}\sigma &\lesssim \int_{L_c(I)} |h^{\underline{L}\underline{L}}|(\underline{L}u)^2 + |h||\bar{\partial}u||\partial u|\,\mathrm{d}\sigma \\ &\lesssim \epsilon\mathrm{flux}([T_1,T_2]) + R\int_{L_c(I)}\frac{|\nabla u|^2}{\langle x\rangle^{1+\gamma}}\mathrm{d}\sigma \end{aligned}\tag{6.15}$$

The conclusion of the lemma now follows from (6.8), (6.11), (6.12), (6.13), and (6.15). □

6.2. L^6 norm decay of solutions on interior spacelike slices of the cone

The objective within this section is to show that solutions to (6.6) satisfy the following estimate.

Lemma 6.4. *Let u solve (6.6). Then*

$$\begin{aligned}\int_{D(T_2)} u^6(T_2,x)\,\mathrm{d}x \lesssim{}& \frac{T_1+2R}{T_2}E_{\{|x|<T_1+c\}}(T_1) + \frac{E}{T_2^\gamma} + G(\mathit{flux}([T_1,T_2])) \\ &+ \|u\|^2_{LE^1[T_1,T_2]} + \|F\|_{L^1[T_1,T_2]L^2}\|\nabla u\|_{L^\infty[T_1,T_2]L^2}\end{aligned}\tag{6.16}$$

We remark that Proposition 6.2 easily follows from Lemmas 6.3 and 6.4.

Proof. To prove (6.16), we multiply both sides of (6.6) by Xu and obtain

$$\begin{aligned}&\partial_\alpha(g^{\alpha\beta}\partial_\beta u Xu) - \frac{1}{2}\partial_t((t+c)g^{\alpha\beta}\partial_\beta u\partial_\alpha u) - \frac{1}{2}\partial_i(x^i g^{\alpha\beta}\partial_\beta u\partial_\alpha u) \\ &\quad + \frac{1}{2}((X-1)g^{\alpha\beta})\partial_\alpha u\partial_\beta u = \partial_t\left((t+c)\frac{u^6}{6}\right) + \partial_i\left(x^i\frac{u^6}{6}\right) - \frac{u^6}{3} + FXu.\end{aligned}\tag{6.17}$$

Indeed, (6.17) follows by the following computations and the symmetry of $g^{\alpha\beta}$:

$$\partial_\alpha(g^{\alpha\beta}\partial_\beta u\partial_t u) - \tfrac{1}{2}\partial_t(g^{\alpha\beta}\partial_\beta u\partial_\alpha u) + \tfrac{1}{2}\partial_t g^{\alpha\beta}\partial_\alpha u\partial_\beta u = (Pu)\partial_t u;$$

https://doi.org/10.1017/S0013091521000158 Published online by Cambridge University Press

similarly, we have

$$\partial_\alpha(g^{\alpha\beta}\partial_\beta ut\partial_t u) - \tfrac{1}{2}\partial_t(tg^{\alpha\beta}\partial_\beta u\partial_\alpha u) - g^{0\beta}\partial_\beta u\partial_t u + \tfrac{1}{2}g^{\alpha\beta}\partial_\alpha u\partial_\beta u + \tfrac{1}{2}t\partial_t g^{\alpha\beta}\partial_\beta u\partial_\alpha u = (Pu)t\partial_t u$$

and

$$\partial_\alpha(g^{\alpha\beta}\partial_\beta ux^j\partial_j u) - g^{j\beta}\partial_\beta u\partial_j u - \tfrac{1}{2}\partial_j(g^{\alpha\beta}\partial_\beta ux^j\partial_\alpha u) + \tfrac{1}{2}x^j\partial_j g^{\alpha\beta}\partial_\beta u\partial_\alpha u + \tfrac{3}{2}g^{\alpha\beta}\partial_\beta u\partial_\alpha u = (Pu)(x^j\partial_j u)$$

as well as

$$\partial_\alpha(g^{\alpha\beta}u\partial_\beta u) - g^{\alpha\beta}\partial_\beta u\partial_\alpha u = (Pu)u.$$

The nonlinear term follows in a similar manner. Upon summing these terms, we obtain (6.17).

We now integrate (6.17) on $\Gamma(I)$ and apply the divergence theorem. We obtain

$$\iint_{\Gamma(I)} \frac{u^6}{3} + \frac{1}{2}((X-1)g^{\alpha\beta})\partial_\alpha u\partial_\beta u - FXu\,\mathrm{d}x\,\mathrm{d}t = -\int_{\partial\Gamma(I)} \nu_\alpha g^{\alpha\beta}\partial_\beta uXu - \frac{1}{2}\nu\cdot(t+c,x)g^{\alpha\beta}\partial_\beta u\partial_\alpha u - \frac{1}{2}\nu\cdot(t+c,x)\frac{u^6}{6}\,\mathrm{d}\sigma$$

Recall that on $L(I)$ the outward unit normal vector ν to $L(I)$ is $(-1, x/|x|)/\sqrt{2}$, and thus $\nu\cdot(t+c,x) = 0$ on $L(I)$. The boundary term can now be written more explicitly as

$$-P(T_2) + P(T_1) + \mathrm{flux}([T_1,T_2]) - BDR_h$$

where the first three terms come from the Minkowski case, and BDR_h denotes the part of the energy density on the boundary arising from h:

$$BDR_h := \int_{D(T_2)} h^{0\beta}\partial_\beta uXu - \frac{1}{2}(t+c)h^{\alpha\beta}\partial_\beta u\partial_\alpha u\,\mathrm{d}x - \int_{D(T_1)} h^{0\beta}\partial_\beta uXu - \frac{1}{2}(t+c)h^{\alpha\beta}\partial_\beta u\partial_\alpha u\,\mathrm{d}x + \int_{L(I)} \nu_\alpha h^{\alpha\beta}\partial_\beta uXu\,\mathrm{d}\sigma$$

As explained in § 5, we know that

$$P(T_2) \gtrsim T_2\int_{D(T_2)} u^6(T_2,x)\,\mathrm{d}x$$

$$P(T_1) \lesssim (T_1+c)E_{\{|x|<T_1+c\}}(T_1)$$

We can also make the trivial estimate

$$\iint_{\Gamma(I)} FXu\,\mathrm{d}x\,\mathrm{d}t \lesssim T_2\|F\|_{L^1[T_1,T_2]L^2}\|\nabla u\|_{L^\infty[T_1,T_2]L^2}$$

Moreover, our assumptions on $h^{\alpha\beta}$ immediately imply that

$$(X-1)h^{\alpha\beta} \lesssim t\langle x\rangle^{-1-\gamma}$$

https://doi.org/10.1017/S0013091521000158 Published online by Cambridge University Press

and thus

$$\iint_{\Gamma(I)} |(X-1)g^{\alpha\beta}\partial_\alpha u \partial_\beta u| \,\mathrm{d}x\,\mathrm{d}t \lesssim T_2 \iint_{\Gamma(I)} \frac{|\nabla u|^2}{\langle x\rangle^{1+\gamma}} \mathrm{d}x\,\mathrm{d}t \leq T_2 \|u\|^2_{LE^1[T_1,T_2]} \tag{6.18}$$

The conclusion (6.16) will follow if we show that

$$BDR_h \lesssim \epsilon(P(T_2)+P(T_1)) + T_2^{1-\gamma}E + T_2(G(\mathrm{flux}([T_1,T_2])) + \|u\|^2_{LE^1[T_1,T_2]})$$

Let us write

$$D(T_2) = D_{\mathrm{int}}(T_2) \cup D_{\mathrm{ext}}(T_2)$$

where

$$D_{\mathrm{int}}(T_2) = D(T_2) \cap \left\{|x| \leq \frac{T_2+c}{2}\right\}, \quad D_{\mathrm{ext}}(T_2) = D(T_2) \cap \left\{|x| \geq \frac{T_2+c}{2}\right\},$$

Since $|h| \lesssim \epsilon$ in D_{int}, we have that

$$\int_{D_{\mathrm{int}}(T_2)} h^{0\beta}\partial_\beta u X u - \frac{1}{2}(t+c)h^{\alpha\beta}\partial_\beta u \partial_\alpha u \,\mathrm{d}x \lesssim \epsilon P(T_2)$$

On the other hand, $|h| \lesssim 1/T_2^\gamma$ in D_{ext}, and thus by the boundedness of energy

$$\int_{D_{\mathrm{ext}}(T_2)} h^{0\beta}\partial_\beta u X u - \frac{1}{2}(t+c)h^{\alpha\beta}\partial_\beta u \partial_\alpha u \,\mathrm{d}x \lesssim T_2^{1-\gamma}E(T_2) \lesssim T_2^{1-\gamma}E$$

Adding the last two inequalities, we obtain

$$\int_{D(T_2)} h^{0\beta}\partial_\beta u X u - \frac{1}{2}(t+c)h^{\alpha\beta}\partial_\beta u \partial_\alpha u \,\mathrm{d}x \lesssim \epsilon P(T_2) + T_2^{1-\gamma}E$$

Similarly, we can show that

$$\int_{D(T_1)} h^{0\beta}\partial_\beta u X u - \frac{1}{2}(t+c)h^{\alpha\beta}\partial_\beta u \partial_\alpha u \,\mathrm{d}x \lesssim \epsilon P(T_1) + T_1^{1-\gamma}E$$

We are left with dealing with the lateral terms. We will show that

$$\int_{L(I)} \nu_\alpha h^{\alpha\beta}\partial_\beta u X u \,\mathrm{d}\sigma \lesssim T_2(G(\mathrm{flux}([T_1,T_2])) + \|u\|^2_{LE^1[T_1,T_2]}) \tag{6.19}$$

We first remark that $Xu = (rL+1)u$ on $L(I)$, and we again write

$$\nu_\alpha h^{\alpha\beta}\partial_\beta u = -\tfrac{1}{2}h^{\underline{L}\underline{L}}\underline{L}u + O(h)\bar{\partial}u \tag{6.20}$$

Note that (6.14), in particular, implies the weaker estimates

$$h^{\underline{L}\underline{L}} \lesssim \frac{R^{1/2}}{\langle x\rangle^{1/2+\gamma}}, \quad h \lesssim 1$$

https://doi.org/10.1017/S0013091521000158 Published online by Cambridge University Press

We can now estimate by Cauchy–Schwarz, (6.14) and the fact that $r \leq T_2 + c \lesssim T_2$:

$$\begin{aligned}\int_{L(I)} |h^{\underline{L}\underline{L}} \underline{L}u(rLu)| \, \mathrm{d}\sigma &\lesssim T_2 \int_{L(I)} \left| \frac{R^{1/2}}{\langle x \rangle^{1/2+\gamma}} \underline{L}u Lu \right| \mathrm{d}\sigma \\ &\leq T_2 \left(R \int_{L(I)} \frac{|\nabla u|^2}{\langle x \rangle^{1+\gamma}} \mathrm{d}\sigma + \int_{L(I)} |Lu|^2 \, \mathrm{d}\sigma \right) \\ &\lesssim T_2 (\|u\|^2_{LE^1[T_1,T_2]} + \mathrm{flux}([T_1, T_2]))\end{aligned}$$

where in the last inequality we used (6.8).

Similarly,

$$\begin{aligned}\int_{L(I)} |h\bar{\partial}u(rLu)| \, \mathrm{d}\sigma &\lesssim T_2 \int_{L(I)} |\bar{\partial}uLu| \, \mathrm{d}\sigma \\ &\leq T_2 \int_{L(I)} |\bar{\partial}u|^2 \, \mathrm{d}\sigma \lesssim T_2 \mathrm{flux}([T_1, T_2])\end{aligned}$$

Thirdly, by an application of (6.14), Cauchy–Schwarz and then Hölder's inequality,

$$\begin{aligned}\int_{L(I)} |h^{\underline{L}\underline{L}} \underline{L}uu| \, \mathrm{d}\sigma &\lesssim \int_{L(I)} \frac{R^{1/2}}{\langle x \rangle^{1/2+\gamma}} |\underline{L}u||u| \, \mathrm{d}\sigma \\ &\lesssim \left(\int_{L(I)} R \frac{|\nabla u|^2}{\langle x \rangle^{1+\gamma}} \mathrm{d}\sigma \right)^{1/2} \left(\int_{L(I)} u^2 \, \mathrm{d}\sigma \right)^{1/2} \\ &\lesssim \left(\int_{L(I)} R \frac{|\nabla u|^2}{\langle x \rangle^{1+\gamma}} \mathrm{d}\sigma \right)^{1/2} \|u\|_{L^6(L(I))} T_2 \\ &\lesssim T_2 (\|u\|^2_{LE^1[T_1,T_2]} + \mathrm{flux}([T_1, T_2])^{1/3})\end{aligned}$$

where again we used (6.8).

Finally,

$$\begin{aligned}\int_{L(I)} |h\bar{\partial}uu| \, \mathrm{d}\sigma &\lesssim \left(\int_{L(I)} |\bar{\partial}u|^2 \, \mathrm{d}\sigma \right)^{1/2} \left(\int_{L(I)} u^2 \, \mathrm{d}\sigma \right)^{1/2} \\ &\lesssim \left(\int_{L(I)} |\bar{\partial}u|^2 \, \mathrm{d}\sigma \right)^{1/2} \left(\int_{L(I)} |u|^6 \, \mathrm{d}\sigma \right)^{1/6} T_2 \\ &\lesssim T_2 G(\mathrm{flux}[T_1, T_2])\end{aligned}$$

The last four estimates now imply (6.19), which finishes the proof of (6.16). □

6.3. Proof of Theorem 6.1

Proof. Assume now that Proposition 6.2 holds. A similar argument as the one in § 4 allows us to pass to the limit and deduce the following:

https://doi.org/10.1017/S0013091521000158 Published online by Cambridge University Press

Proposition 6.5. *Let u be the strong solution to (2.2). For any $R \geq 1$, $1 < T_1$, and $T_1 + 2R < T_2$, we have*

$$\int_{\mathbb{R}^3} u^6(T_2, x)\, \mathrm{d}x \lesssim \frac{T_1 + 2R}{T_2} E_{\{|x|T_1+R\}}(T_1) + \|u\|^2_{LE^1[T_1,T_2]}) \tag{6.21}$$

Let us now prove (6.4). Pick any $\tilde{\epsilon} > 0$, and let T_1 be large enough such that

$$\|u\|^2_{LE^1[T_1,\infty)} < \tilde{\epsilon};$$

such a number may be found because of the local energy estimate (4.1). Next pick R large enough so that

$$E_{\{|x|>T_1+R\}}(T_1) < \tilde{\epsilon}$$

Now let $T_2 \to \infty$ in (6.21). We obtain

$$\limsup_{T_2\to\infty} \int_{\mathbb{R}^3} u^6(T_2, x)\, \mathrm{d}x \lesssim G(\tilde{\epsilon})$$

and (6.4) follows by letting $\tilde{\epsilon} \to 0$.

To obtain part (2) of Theorem 6.1, note that if $\partial^J h \lesssim \epsilon \langle x\rangle^{-|J|-\gamma}$ where $|J| \leq 2$ (which are implied by our assumptions in the main theorem), then global Strichartz estimates are implied by a refinement of the local energy decay estimates (see Theorem 2 in [25]†). Then, for any $\eta > 0$, by choosing a sufficiently large number $T > 0$, we obtain $\|w\|_{L^5L^{10}([T,\infty)\times\mathbb{R}^3)} \leq \eta$ where w solves (1.1). For any $\tilde{w}$ with $\|\tilde{w}\|_{L^5L^{10}([T,\infty)\times\mathbb{R}^3)} \leq \eta$, let W be the solution to

$$PW = (w + \tilde{w})^5$$

with

$$\lim_{t\to\infty} \|\nabla W(t,\cdot)\|_{L^2(\mathbb{R}^3)} = 0.$$

As $\eta > 0$ was arbitrary, we may select η sufficiently small so that the map $\tilde{w} \mapsto W$ is a contraction mapping, so that for any finite-energy solution w of (1.1), there exists a unique solution to (2.1) such that their difference vanishes in the $\dot{H}^1 \times L^2$ norm as $t \to \infty$, and we conclude that the solution scatters in the energy space. □

References

1. H. Bahouri and P. Gérard, High frequency approximation of solutions to critical nonlinear wave equations, *Amer. J. Math.* **121** (1999), 131–175.
2. H. Bahouri and J. Shatah, Decay estimates for the critical semilinear wave equation, *Ann. Inst. H. Poincare Anal. Non Lineaire* **15** (1998), 783–789.
3. N. Burq, G. Lebeau and F. Planchon, Global existence for energy critical waves in 3-D domains, *J. Amer. Math. Soc.* **21** (2008), 831–845.
4. T. Duyckaerts and D. Lafontaine, *Scattering for critical radial Neumann waves outside a ball*, arXiv:2004.08576.
5. J. Ginibre, A. Soffer and G. Velo, The global Cauchy problem for the critical nonlinear wave equation, *Jour. Func. Anal.* **110** (1992), 96–130.

† In the current arXiv version of this paper, see Theorem 6.1 instead.

https://doi.org/10.1017/S0013091521000158 Published online by Cambridge University Press

6. M. Grillakis, A priori estimates and regularity of nonlinear waves, in *Proceedings of the International Congress of Mathematicians*, pp. 1187–1194 (Birkhauser, 1994).
7. M. Grillakis, Regularity and asymptotic behavior of the wave equation with a critical nonlinearity, *Ann. Math.* **132** (1990), 485–509.
8. M. Grillakis, Regularity for the wave equation with a critical nonlinearity, *Comm. Pure Appl. Math.* **45** (1992), 749–774.
9. S. Ibrahim and M. Majdoub, Solutions globales de l'equation des ondes semilineaire critique 'a coefficients variables, *Bull. Soc. Math. France. (1)* **131** (2003), 1–22.
10. C. Jao, The quintic NLS on perturbations of $\mathbb{R}^3$, *Amer. J. Math.* **141** (2019), 981–1035.
11. H. Jia, B. Liu and G. Xu, Long time dynamics of defocusing energy critical $3+1$ dimensional wave equation with potential in the radial case, *Comm. Math. Phys.* **339**(2) (2015), 353–384.
12. H. Jia, B. Liu, W. Schlag and G. Xu, Global center stable manifold for the defocusing energy critical wave equation with potential, *Amer. J. Math.* to appear.
13. L. Kapitanski, Global and unique weak solutions of nonlinear wave equations, *Math. Res. Lett.* **1** (1994), 211–223.
14. M. Keel and T. Tao, Endpoint Strichartz estimates, *Amer. J. Math.* **120** (1998), 955–980.
15. M. Keel, H. Smith and C. Sogge, Almost global existence for some semilinear wave equations, *J. d'Anal. Math.* **87** (2002), 265–279.
16. C. Kenig and F. Merle, Global well-posedness, scattering and blow-up for the energy-critical focusing non-linear wave equation, *Acta Math.* **201**(2) (2008), 147–212.
17. R. Killip, M. Visan and X. Zhang, Quintic NLS in the exterior of a strictly convex obstacle, *Amer. J. Math.* **138** (2016), 1193–1346.
18. R. Killip, C. Miao, M. Visan, J. Zhang and J. Zheng, The energy-critical NLS with inverse- square potential, *Discrete Contin. Dyn. Syst.* **37**(7) (2017), 3831–3866.
19. D. Lafontaine, *About the wave equation outside two strictly convex obstacles*, arXiv:1711.09734.
20. N.-A. Lai. and Y. Zhou, Global existence of critical nonlinear wave equation with time dependent variable coefficients, *Comm. Part. Diff. Equ.* **11** (2012), 2171–2183.
21. A. Lawrie, S.-J. Oh and S. Shahshahani, Profile decompositions for wave equations on hyperbolic space with applications, *Math. Ann.* **365**(1–2) (2016), 707–803.
22. H. Lindblad and I. Rodnianski, The global stability of Minkowski space- time in harmonic gauge, *Ann. Math.* **171** (2010), 1401–1477.
23. J. Metcalfe and C. Sogge, Long-time existence of quasilinear wave equations exterior to star-shaped obstacles via energy methods, *SIAM J. Math. Anal.* **38** (2006), 188–209.
24. J. Metcalfe and D. Tataru, Decay estimates for variable coefficient wave equations in exterior domains, in *Advances in phase space analysis of partial differential equations*, Progr. Nonlinear Differential Equations Appl., 78, pp. 201–216 (2009).
25. J. Metcalfe and D. Tataru, Global parametrices and dispersive estimates for variable coefficient wave equations, *Math. Ann.* **353**(4) (2012), 1183–1237.
26. C. Miao, J. Murphy and J. Zheng, The energy-critical nonlinear wave equation with an inverse-square potential, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **37**(2) (2020), 417–456.
27. C. Morawetz, Time decay for the nonlinear Klein-Gordon equation, *Proc. Roy. Soc. Lon.* **306** (1968), 291–296.
28. K. Nakanishi, Scattering theory for nonlinear Klein-Gordon equation with Sobolev critical power, *Int. Math. Res. Not.* **1** (1999), 31–60.

https://doi.org/10.1017/S0013091521000158 Published online by Cambridge University Press

29. K. Nakanishi, Unique global existence and asymptotic behaviour of solutions for wave equations with non-coercive critical nonlinearity, *Comm. Partial Diff. Equ.* **24** (1999), 185–221.
30. H. Pecher, Nonlinear small data scattering for the wave and Klein-Gordon equations, *Math. Z.* **185** (1984), 261–270.
31. J. Rauch, The u^5 Klein-Gordon equation. II. Anomalous singularities for semilinear wave equations, in *Nonlinear partial differential equations and their applications*. College de France Seminar, pp. 335–364, Research Notes in Mathematics, Volume 51 (Pitman Publ., 1981).
32. J. Shatah and M. Struwe, Regularity results for nonlinear wave equations, *Ann. Math.* **138** (1993), 503–518.
33. J. Shatah and M. Struwe, Well-posedness in the energy space for semilinear wave equation with critical growth, *Int. Math. Res. Not.* **1994** (1994), 303–309.
34. H. F. Smith and C. D. Sogge, On the critical semilinear wave equation outside convex obstacles, *J. Amer. Math. Soc.* **8**(4) (1995), 879–916.
35. J. Sterbenz and I. Rodnianski, Angular regularity and strichartz estimates for the wave equation, *Int. Math. Res. Not.* **4** (2005), 187–231.
36. M. Struwe, Globally regular solutions to the u^5 Klein-Gordon equation, *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* **15** (1988), 495–513.
37. J. Zhang and J. Zheng, Scattering theory for nonlinear Schrödinger with inverse-square potential, *J. Funct. Anal.* **267** (2014), 2907–2932.

https://doi.org/10.1017/S0013091521000158 Published online by Cambridge University Press