Ergod. Th. & Dynam. Sys. (2005), **25**, 1739–1765
doi:10.1017/S0143385705000179
© *2005 Cambridge University Press*
Printed in the United Kingdom

Livšic regularity for Markov systems

HENK BRUIN†, MARK HOLLAND† and MATTHEW NICOL‡
† *Department of Mathematics and Statistics, University of Surrey, Guildford, Surrey GU2 7XH, UK*
(e-mail: h.bruin@surrey.ac.uk, mark.holland@surrey.ac.uk)
‡ *Department of Mathematics, University of Houston, Houston, TX 77204-3008, USA*
(e-mail: nicol@math.uh.edu)

(*Received* 7 *January* 2005 *and accepted in revised form* 19 *April* 2005)

Abstract. We prove measurable Livšic theorems for dynamical systems modelled by Markov towers. Our regularity results apply to solutions of cohomological equations posed on Hénon-like mappings and a wide variety of non-uniformly hyperbolic systems. We consider both Hölder cocycles and cocycles with singularities of prescribed order.

1. *Introduction*

In this paper we study the regularity of solutions ψ of the cohomological equation

$$\varphi = (\psi \circ T)\psi^{-1} \quad \mu\text{-a.e.} \tag{1}$$

where (T, X, μ) is a dynamical system and $\varphi : X \to G$ is a cocycle taking values in a Lie group G. Measurable rigidity in this context means that a measurable solution ψ must have a higher degree of regularity, in many contexts inheriting the regularity of φ and/or T. Such cohomological equations arise in different applications: they are used to determine whether certain observables have positive variance in the context of the Central Limit Theorem and related distributional theorems. In the context of group extensions, they decide on (stable) ergodicity and weak-mixing of the system. In other contexts, cohomological equations play a role in the question of whether two dynamical systems are (Hölder or smoothly) conjugate to each other.

Fundamental work on the regularity of measurable solutions to cohomological equations was done by Livšic [**24, 25**] who established rigidity theorems for Abelian group extensions of Anosov systems with an absolutely continuous invariant measure. Dynamical rigidity theorems are often called Livšic theorems in the literature because of this. Parry and Pollicott [**33**], using a transfer operator approach, extended Livšic's results to prove Hölder regularity of coboundary and transfer functions for compact Lie group extensions of subshifts of finite type and hence, via Markov partitions, Axiom A systems.

https://doi.org/10.1017/S0143385705000179 Published online by Cambridge University Press

Further generalizations for uniformly hyperbolic smooth systems are given in [**28, 29, 34, 38, 39**]. In the Anosov setting de la Llave *et al* [**13**] proved a C^∞ version of Livšic's theorem and also C^∞ dependence of solutions upon parameters.

There are only a few results on Livšic theorems for non-uniformly hyperbolic or discontinuous systems. Pollicott and Yuri [**35**] have established Livšic theorems for Hölder $\mathbb{R}$-extensions of β-transformations ($f : [0,1) \longrightarrow [0,1)$, $f(x) = \beta x \pmod 1$ where $\beta > 1$) via transfer operator techniques but the regularity they obtain is bounded variation rather than Hölder. Jenkinson [**18**] has proved that essentially bounded measurable coboundaries ψ (i.e. solutions to $\varphi = \psi \circ f - \psi$) for $\mathbb{R}$-valued smooth cocycles φ over smooth expanding maps f have smooth versions.

Nicol and Scott [**30**] have obtained Livšic theorems for certain discontinuous hyperbolic systems, showing that coboundary solutions taking values in Lie groups satisfying a pinching condition (to ensure that the system is partially hyperbolic) are Lipschitz if the cocycle is Lipschitz. The same techniques show that, for such systems, measurable transfer functions taking values in compact matrix groups have Lipschitz versions. These results were applied to prove stable ergodicity for semisimple and Abelian compact group extensions of certain uniformly hyperbolic systems with singularities, including the β-transformation, Markov maps and mixing Lasota–Yorke maps.

Aaronson and Denker [**1**, Corollary 2.3] have shown that if $(f, X, \mu, \mathcal{P})$ is a mixing Gibbs–Markov map preserving a probability measure μ with countable Markov partition $\mathcal{P}$ and $\varphi : X \to \mathbb{R}^d$ is Lipschitz (i.e. $d(\varphi(x), \varphi(y)) < C\rho(x, y)$ for all $x, y \in f(a), a \in \mathcal{P}$ for metrics d on $\mathbb{R}^d$ and ρ on X derived from the symbolic dynamics) then any measurable solution $\psi : X \to \mathbb{R}^d$ to $\varphi = \psi \circ f - \psi$ has a version $\tilde{\varphi}$ which is Lipschitz continuous.

In [**2**] there is a study of the statistical properties of fibred systems and rigidity results are given which provide checkable conditions for the aperiodicity of cocycles (i.e. non-existence of solutions ψ) which allow one to establish, for example, de Moivre's approximation for various systems, including the β-transformation. A related result is given in [**14**, Lemma 6.1.2].

In two influential papers [**41, 42**], Young describes properties of a class of Markov extensions (which we call Young towers) which are useful to establish rates of decay of correlations and the CLT in non-uniformly hyperbolic systems. Scott [**36, 37**] has recently proved measurable Livšic theorems for certain Lie group valued Hölder cocycles over a class of unimodal maps modelled by a Young tower [**41**]. More precisely suppose that (f, X, μ) (X a finite collection of intervals) is a unimodal map (belonging to a certain class) and $g : X \to G$ is a Lie group valued Hölder cocycle (satisfying a pinching condition if G is non-compact). If ψ is a measurable solution to $g = (\psi \circ f)\psi^{-1}$ μ-a.e. then ψ is Hölder on an arbitrarily large open set (i.e. given $\varepsilon > 0$ there exists an open set U such that ψ is Hölder on U and $\mu(U) > 1 - \varepsilon$). Similar measurable Livšic theorems for other types of cohomological equations are given.

This paper extends the results of Scott in several directions. We prove measurable Livšic theorems for more general Markov extensions and for cocycles with singularities of prescribed order. We also obtain regularity results for measurable conjugacies between certain non-uniformly hyperbolic systems and measurable Livšic theorems for certain non-uniformly hyperbolic systems (in particular Hénon-like mappings).

https://doi.org/10.1017/S0143385705000179 Published online by Cambridge University Press

After writing this paper we learnt that Gouëzel [**16**] has obtained similar results for cocycles into Abelian groups over one-dimensional Gibbs–Markov systems and Young towers. From [**16**] we learnt of a Martingale Density Theorem (see Appendix A) which allows a more elegant approach in part of our proof than the argument using Lebesgue density points adapted from [**30**].

2. *Cohomological equations and group extensions*

In this section we collect some facts about group extensions and cohomological equations that set the framework of this paper. All Lie groups are assumed connected and finite-dimensional.

Suppose that $f : X \to X$ is an ergodic dynamical system with respect to an invariant measure μ. Let G be a compact Lie group and let dh denote Haar measure on G. Suppose $\varphi : X \to G$ is measurable.

PROPOSITION 1. [**23, 31**]

(1) *The compact group extension $T(x, g) = (f(x), \varphi(x)g)$ is ergodic with respect to $\mu \times dh$ if and only if the equation*

$$\psi(fx) = R(\varphi(x))\psi(x) \quad \mu\text{-a.e.} \tag{2}$$

where R is an irreducible (unitary) representation of degree d and $\psi : X \longrightarrow \mathbb{C}^d$ is measurable, is only satisfied when ψ is constant or R is the trivial representation.

(2) *Suppose $f : X \to X$ is weak-mixing and $T : X \times G \to X \times G$ is ergodic. Then $T(x, g) = (f(x), \varphi(x)g)$ is weak-mixing with respect to $\mu \times dh$ if and only if for any $e^{i\alpha} \neq 1$ and any non-trivial one-dimensional representation χ of G the equation*

$$\psi(fx) = e^{i\alpha}\chi(\varphi(x))\psi(x) \quad \mu\text{-a.e.} \tag{3}$$

has no non-trivial measurable solution $\psi : X \to \mathbb{C}$.

Note that the aperiodicity condition of [**2**]

$$\gamma \circ \varphi = \lambda\psi/\psi \circ f$$

where γ is a character of G (a locally compact Abelian polish group), $\lambda \in S^1$ is a special case of (3).

Suppose that $\varphi_i : X \to G$, $i = 1, 2$ are two compact Lie-group valued cocycles over a system (f, X, μ). A measurable function $\psi : X \to G$ conjugates the G extensions $T(x, g) = (fx, \varphi_i(x)g)$ $(i = 1, 2)$ if

$$\psi(fx)\varphi_1(x) = \varphi_2(x)\psi(x) \quad \mu\text{-a.e.} \tag{4}$$

We call such a conjugating function ψ a *transfer function*.

If G is compact we may identify G with a subgroup of $U(d)$, the group of $d \times d$ unitary matrices. In this representation we may identify a G-valued $\varphi : X \to G$ with $\varphi : X \to \mathbb{C}^{d^2}$. Define $\theta(x) : M(d) \to M(d)$, a mapping from the space of $d \times d$ complex matrices to itself by

$$\theta(x) : A \to \varphi_2(x)A\varphi_1(x)^*.$$

https://doi.org/10.1017/S0143385705000179 Published online by Cambridge University Press

It is possible to show that $\theta(x) : M(d) \to M(d)$ is unitary. There is a standard way, see [**33**, Theorem 1] and [**18**, Theorem A], to rewrite $\psi(fx)\varphi_1(x) = \varphi_2(x)\psi(x)$ in the form $\psi(fx) = \theta(x)\psi(x)$ where $\psi : X \to \mathbb{C}^{d^2}$, $\theta(x) \in U(d)$. Hence, the question of the regularity of conjugacies between compact group extensions may be reduced to those of the regularity of solutions to (2).

The proof of our coboundary Livšic regularity results, such as Theorem 1, may be slightly modified (as in [**30**, §2.1]), to establish the same degree of regularity for solutions ψ to (2), (3) or (4) posed over the same dynamical system. We omit the straightforward proof of this and refer the reader to [**30**, §2.1].

2.1. *Lie groups.* Let G be a connected Lie group with Lie algebra denoted by $L(G)$ which we identify with the tangent space at the identity, T_eG. We let r_g denote right multiplication by $g \in G$. Given a norm $\|\cdot\|$ on T_eG we define a norm on T_gG by $\|\mathbf{v}\|_g = \|r_{g^{-1}}\mathbf{v}\|_e$. This norm induces a right invariant metric d_G on G so that $d_G(gk, hk) = d_G(g, h)$, see [**34**, §4]. Throughout this paper we will write $d(\cdot,\cdot)$ instead of $d_G(\cdot,\cdot)$ when it is clear from context that we mean the metric on G. For a general reference on Lie groups, see [**7**].

We define the adjoint map $\mathrm{Ad}\colon G \to \mathrm{Aut}(L(G))$, for $g \in G$ and $X \in T_eG$ by

$$\mathrm{Ad}(g)\mathbf{v} = \frac{d}{dt}(g\exp(t\mathbf{v})g^{-1}) \text{ at } t = 0.$$

Note that when G is a matrix group this action is conjugation, i.e. $v \to gvg^{-1}$. A calculation [**34**, §4] shows that

$$d(gh, gk) \le \|\mathrm{Ad}\,(g)\|d(h, k). \tag{5}$$

Suppose that $\varphi\colon M \to G$ is Hölder of exponent $\alpha > 0$. Define

$$\mu_u := \lim_{n\to\infty}\Big(\sup_{x\in X}\|\mathrm{Ad}\,(\varphi_n(x))\|\Big)^{1/n},$$

where $\varphi_n(x) = \varphi(f^{n-1}x)\ldots\varphi(fx)\varphi(x)$.

If G is Abelian or compact, then $\sup_x \|\mathrm{Ad}\,(\varphi_n(x))\|$ is bounded in n, whereas if G is nilpotent, $\sup_x \|\mathrm{Ad}\,(\varphi_n(x))\|$ can grow at most at a polynomial rate. In these three cases, $\mu_u = 1$.

3. *Axiomatic approach for non-uniformly expanding maps*

Let (M, ρ, μ) be a metric space endowed with a non-atomic Borel probability measure μ. We assume that M can be decomposed as $M = \bigcup_k M_k \bmod \mu$, where each M_k is connected and $\sup_k(\mathrm{diam}(M_k)) \le 1$. Let $f : \bigcup_k M_k \to M$ be a map such that $f|M_k$ is continuous for each k and such that μ is f-invariant and ergodic.

Let $\mathcal{P}_0$ be the partition of M into the sets M_k, and $\mathcal{P}_n = \bigvee_{i=0}^{n-1} f^{-i}(\mathcal{P}_0)$. For $x \in M$, let $\mathcal{P}_n[x]$ be the partition element (cylinder set) in $\mathcal{P}_n$ containing x.

Consider the natural extension $(\hat{M}, \hat{f}, \hat{\mu})$ of (M, f, μ): each point $\hat{x} \in \hat{M}$ is a sequence

$$\hat{x} = (x_0, x_1, x_2, \ldots) \quad \text{with } M \ni x_i = f(x_{i+1}) \text{ for all } i \ge 0.$$

https://doi.org/10.1017/S0143385705000179 Published online by Cambridge University Press

The measure $\hat{\mu}$ is defined in the standard way [**20**] and, in particular, for each n

$$\hat{\mu}(\{\hat{x} \in \hat{M} : x_n \in A\}) = \mu(A)$$

for each μ-measurable set A.

We assume the following.

(1) For all k, $f : M_k \to f(M_k)$ is one-to-one and $f(M_k)$ is equal to a union of components M_l mod μ (Markov property).

(2) There exists $\lambda > 1$ and for $\hat{\mu}$-a.e. $\hat{x}$ a number $K(\hat{x})$ such that

$$\rho(y_n, z_n) \le K(\hat{x})\lambda^{-n}\rho(y_0, z_0) \tag{6}$$

for all $n \ge 0$ and $y_n, z_n \in \mathcal{P}_n[x_n]$. (In dimension one this assumption can be weakened, see §4).

(3) Let $J_\mu(x)$ denote the Jacobian of μ at x. For $\hat{x} \in \hat{M}$, define $J_\mu^n(x_n) = \prod_{i=1}^n J_\mu(x_i)$. For $\hat{\mu}$-a.e. $\hat{x}$, there exists a constant $C(\hat{x})$ such that if $\hat{y}, \hat{z} \in \hat{M}$ are such that $y_i, z_i \in \mathcal{P}_i[x_i]$ for all i, then

$$\left|\frac{J_\mu^n(y_n)}{J_\mu^n(z_n)}\right| \le C(\hat{x}). \tag{7}$$

The Jacobian J_μ of μ is γ-Hölder if there exists C and $\gamma \in (0, 1]$ such that

$$\left|\frac{J_\mu(x)}{J_\mu(y)} - 1\right| \le C \cdot \rho(f(x), f(y))^\gamma. \tag{8}$$

Hölderness of the Jacobian implies a result stronger than (7).

LEMMA 1. *Assume J_μ is γ-Hölder with coefficient C. For $\hat{\mu}$-a.e. $\hat{x}$, there exists a constant $B = B(\hat{x}, C, \lambda^\gamma)$ such that*

$$\left|\frac{J_\mu^n(y_n)}{J_\mu^n(z_n)}\right| \le 1 + B\rho(y_0, z_0)^\gamma.$$

Proof. Using (8) and (6), we obtain

$$\begin{aligned}
\left|\frac{J_\mu^n(y)}{J_\mu^n(z)}\right| &= \prod_{i=0}^{n-1} \frac{|J_\mu(y_i)|}{|J_\mu(z_i)|} \\
&\le \prod_{i=0}^{n-1} (1 + C\rho(f(y_i), f(z_i))^\gamma) \\
&\le \exp\left(C \sum_{i=1}^{n} \rho(y_i, z_i)^\gamma\right) \\
&\le \exp\left(C \cdot K(\hat{x}) \sum_{i=1}^{n} \lambda^{-i\gamma} \rho(y_0, z_0)^\gamma\right) \\
&\le \exp\left(\frac{C \cdot K(\hat{x})}{\lambda^\gamma - 1} \rho(y_0, z_0)^\gamma\right),
\end{aligned}$$

which is smaller than $1 + B\rho(y_0, z_0)^\gamma$ for some B depending only on $\hat{x}$, C, λ^γ and the diameter of the component M_k containing x_0.

In fact, the above computation only requires that $\sum_{i=0}^{n-1} \operatorname{diam}(\mathcal{P}_i[x_i])^\gamma \le K_0(\hat{x}) < \infty$ for μ-a.e. $\hat{x}$ independently of n, which is an estimate valid under a less strict assumption than (6). □

https://doi.org/10.1017/S0143385705000179 Published online by Cambridge University Press

Cocycle assumptions. Let $\varphi\colon M \to G$ be Hölder of exponent $\alpha > 0$. Recall that

$$\mu_u := \lim_{n\to\infty} \Big(\sup_{x\in M} \|\mathrm{Ad}\,(\varphi_n(x))\| \Big)^{1/n},$$

where $\varphi_n(x) = \varphi(f^{n-1}x)\dots\varphi(fx)\varphi(x)$ with $\varphi_0(x) = e$. If G is Abelian, $\|\mathrm{Ad}\,(\varphi_n(x))\| = 1$ and $\|\mathrm{Ad}\,(\varphi_n(x))\|$ is bounded if G is compact. For nilpotent groups G, $\|\mathrm{Ad}\,(\varphi_n(x))\|$ grows at most at a polynomial rate in n, so $\mu_u = 1$. For the general case, we impose a partial hyperbolicity condition on the group extension:

$$1 \le \mu_u < \lambda^\alpha \tag{PH}$$

where λ is from (6).

THEOREM 1. *Assume that (M, f, μ) is a measure-preserving Markov system as above and let $M_k \in \mathcal{P}_0$. Let $\varphi : M \to G$ be a Lie group valued α-Hölder observable (i.e. $d(\varphi(x), \varphi(y)) \le C\rho(x, y)^\alpha$) satisfying the partial hyperbolicity condition (PH). Let $\psi : M \to G$ be a μ-measurable solution of the cohomological equation*

$$\psi \circ f(x) = \varphi(x) \cdot \psi(x) \quad \mu\text{-a.e.}$$

Then there is a version $\tilde\psi$ of ψ (i.e. $\psi = \tilde\psi$ μ-a.e.) such that $\tilde\psi$ is α-Hölder on M_k.

COROLLARY 1. *If $f^j\big(\bigcup_{k\in S} M_k\big) = M$ for some $j > 0$ and finite collection of indices S then there is a version which is α-Hölder on M.*

Proof of Corollary 1. By considering the cohomological equation $\psi \circ f^n(x) = \varphi(f^{n-1}x)\cdot \varphi(f^{n-1}x)\dots\varphi(x)\cdot\psi(x)$ we may extend the version of ψ as a Hölder function to any image $f^j(M_k)$. □

Remark 1. It is easy to show that given $\varepsilon > 0$ there is a version of ψ which is α-Hölder on a finite union of sets $\bigcup_{k\in S} M_k$ such that $\mu\big(\bigcup_{k\in S} M_k\big) > 1 - \varepsilon$. The Hölder coefficient depends, in general, upon S but the exponent is uniform.

Remark 2. A slight modification of the proof shows that the same regularity results hold for solutions ψ to (2), (3) or (4).

Proof of Theorem 1. Choose any $\Lambda := M_k \in \mathcal{P}_0$ such that $\mu(\Lambda) > 0$. Let $0 < \delta < 1$. As a consequence of the Martingale Density Theorem (see Appendix A) for $\hat\mu$-a.e. $\hat x \in \hat M$ and for infinitely many n

$$\frac{\mu\{y_n \in \mathcal{P}_n[x_n] : d(\psi(y_n), \psi(x_n)) < \delta\}}{\mu(\mathcal{P}_n[x_n])} > 1 - \delta.$$

Let $\hat x$ be such a point with $x_0 \in \Lambda$. For simplicity of notation in the rest of the proof we will not indicate the dependence of constants upon $\hat x$. We consider points $\hat y = (y_0, y_1, \dots y_n \dots)$ and $\hat z = (z_0, z_1, \dots z_n \dots) \in \hat M$ such that $y_n, z_n \in \mathcal{P}_k[x_n]$ for all $n = 0, 1, \dots$. Hence, $\hat x$, $\hat y$ and $\hat z$ are all paths in the 'same inverse branch' of f.

On Λ we define a function $\Phi : \Lambda \to G$ by

$$\Phi(y_0) = \lim_{n\to\infty} \varphi_n(y_n)\varphi_n(x_n)^{-1},$$

https://doi.org/10.1017/S0143385705000179 Published online by Cambridge University Press

where $\varphi_n(x_n) = \varphi(x_1)\dots\varphi(x_n)$. This function is well defined since

$$\begin{aligned}
&d(\varphi_{n+1}(y_{n+1})\varphi_{n+1}(x_{n+1})^{-1}, \varphi_n(y_n)\varphi_n(x_n)^{-1}) \\
&\quad = d(\varphi_n(y_n)\varphi(y_{n+1})\varphi(x_{n+1})^{-1}\varphi_n(x_n)^{-1}, \varphi_n(y_n)\varphi_n(x_n)^{-1}) \\
&\quad = d(\varphi_n(y_n)\varphi(y_{n+1})\varphi(x_{n+1})^{-1}\varphi_n(x_n)^{-1}, \varphi_n(y_n)\varphi(x_{n+1})\varphi(x_{n+1})^{-1}\varphi_n(x_n)^{-1}) \\
&\quad = d(\varphi_n(y_n)\varphi(y_{n+1}), \varphi_n(y_n)\varphi(x_{n+1})) \quad \text{(right invariance)} \\
&\quad \le \|\mathrm{Ad}\,(\varphi_n(y_n))\|\, d(\varphi(y_{n+1}), \varphi(x_{n+1})) \quad \text{(by (5))} \\
&\quad \le C \cdot K(\hat{x})((\mu_u)\lambda^{-\alpha})^n \quad \text{(by (6))} \\
&\quad \le C \cdot K(\hat{x}) \cdot \kappa^n,
\end{aligned}$$

where $\kappa \in (0, 1)$ by (PH). Thus, the sequence $\varphi_n(y_n)\varphi_n(x_n)^{-1}$ is Cauchy and so converges.

Next we show that Φ is Hölder. Let $y_0, z_0 \in \Lambda$, then

$$\begin{aligned}
&d(\varphi_n(y_n)\varphi_n(x_n)^{-1}, \varphi_n(z_n)\varphi_n(x_n)^{-1}) \\
&\quad = d(\varphi_n(y_n), \varphi_n(z_n)) \\
&\quad \le \sum_{i=0}^{n-1} d(\varphi_i(y_i)\varphi(y_{i+1})\varphi_{n-i-1}(z_{i+1}), \varphi_i(y_i)\varphi(z_{i+1})\varphi_{n-i-1}(z_{i+1})) \\
&\quad \le \sum_{i=0}^{n-1} \|\mathrm{Ad}\,(\varphi_i(y_i))\| d(\varphi(y_{i+1}), \varphi(z_{i+1})) \\
&\quad \le \sum_{i=0}^{n-1} C \cdot K(\hat{x})(\mu_u)^{i+1}\lambda^{-(i+1)\alpha}\rho(y_0, z_0)^\alpha. \qquad (9)
\end{aligned}$$

Letting $n \to \infty$ gives $d(\Phi(y_0), \Phi(z_0)) \le C \cdot K(\hat{x}) \cdot \rho(y_0, z_0)^\alpha$. It is clear that if φ is Lipschitz (i.e. $\alpha = 1$) then Φ is also Lipschitz.

Define

$$\Psi_n(y_0) = \varphi_n(y_n)\varphi_n(x_n)^{-1}.$$

Then

$$\begin{aligned}
\psi(y_0) &= \varphi_n(y_n)\psi(y_n) \\
&= \Psi_n(y_0)\varphi_n(x_n)\psi(x_n)\psi(x_n)^{-1}\psi(y_n) \\
&= \Psi_n(y_0)\psi(x_0)\psi(x_n)^{-1}\psi(y_n).
\end{aligned}$$

Thus

$$\begin{aligned}
d(\psi(y_0), \psi(z_0)) \le\ & d(\Psi_n(y_0)\psi(x_0)\psi(x_n)^{-1}\psi(y_n), \Psi_n(y_0)\psi(x_0)) \\
&+ d(\Psi_n(y_0)\psi(x_0)\psi(x_n)^{-1}\psi(z_n), \Psi_n(y_0)\psi(x_0)) \\
&+ d(\Psi_n(y_0)\psi(x_0)\psi(x_n)^{-1}\psi(z_n), \Psi_n(z_0)\psi(x_0)\psi(x_n)^{-1}\psi(z_n)).
\end{aligned}$$

By right-invariance of the metric the last term may be written as

$$d(\Psi_n(y_0), \Psi_n(z_0)).$$

As a function of y_0, $\Psi_n(y_0)$ converges to the α-Hölder function $\Psi(y_0)$. Thus, letting $n \to \infty$, we obtain $d(\Psi(y_0), \Psi(z_0)) \le C\rho(y_0, z_0)^\alpha$.

https://doi.org/10.1017/S0143385705000179 Published online by Cambridge University Press

Given $\eta > 0$ there exists $\delta_\eta > 0$ such that $d(\psi(z_n), \psi(y_n)) \le \delta_\eta$ implies

$$d(\Psi_n(y_0)\psi(x_0)\psi(x_n)^{-1}\psi(y_n), \Psi_n(y_0)\psi(x_0)) \le \frac{\eta}{2},$$

$$d(\Psi_n(y_0)\psi(x_0)\psi(x_n)^{-1}\psi(z_n), \Psi_n(y_0)\psi(x_0)) \le \frac{\eta}{2}.$$

Choose n sufficiently large so that

$$\frac{\mu\{y_n \in \mathcal{P}_n[x_n] : d(\psi(y_n), \psi(x_n)) < \delta_\eta\}}{\mu(\mathcal{P}_n[x_n])} > 1 - \delta_\eta.$$

Now we estimate $\mu(f^n(\{y_n \in \mathcal{P}_n[x_n] : d(\psi(y_n), \psi(x_n)) < \delta_\eta\}))$ relative to $\mu(\Lambda)$. By boundedness of distortion of the Jacobian of f^n we have that

$$\frac{\mu(f^n(\{y_n \in \mathcal{P}_n[x_n] : d(\psi(y_n), \psi(x_n)) < \delta_\eta\}))}{\mu(f^n\mathcal{P}_n[x_n])} \le \mathcal{O}(1)\frac{\mu(\{y_n \in \mathcal{P}_n[x_n] : d(\psi(y_n), \psi(x_n)) < \delta_\eta\})}{\mu(\mathcal{P}_n[x_n])}. \tag{10}$$

Hence, for the above $\eta > 0$, choosing δ_η smaller if necessary, we have

$$\mu(f^n(\{y_n \in \mathcal{P}_n[x_n] : d(\psi(y_n), \psi(x_n)) < \delta_\eta\})) > (1 - \eta)\mu(\Lambda).$$

Since $d(\psi(y_n), \psi(z_n)) < \delta_\eta$ implies $d(\psi(z_0), \psi(y_0)) < \eta + K\rho(z_0, y_0)^\alpha$ we have shown that

$$\mu\times\mu\{(y_0, z_0) \in \Lambda\times\Lambda : d(\psi(z_0), \psi(y_0)) < 2\eta+2K\rho(z_0, y_0)^\alpha\} > (1-2\eta)\mu\times\mu(\Lambda\times\Lambda).$$

Since η was arbitrary, $\psi \mid \Lambda$ has a Hölder version. □

3.1. *Cocycles with singularities.* Let $\varphi : M \to G$ be a cocycle which is Hölder except for discontinuities and singularities concentrated on a finite set $\mathcal{C}$. Let λ be as in (6), which is defined μ-a.e., and for a fixed $\delta > 0$ let $B(c, \delta)$ denote a δ-neighbourhood of c for $c \in \mathcal{C}$. We consider the following three scenarios.

(1) *Bounded discontinuity.* The cocycle $\varphi(x)$ is bounded and γ-Hölder in the complement of $\mathcal{C}$ but for each $c \in \mathcal{C}$ we have

$$\lim_{x\to c^+} \varphi(x) \ne \lim_{x\to c^-} \varphi(x).$$

(2) *Logarithmic singularity.* For each $c \in \mathcal{C}$ inside $B(c; \delta)$ we have $\|\varphi(x)\| \approx |\log \rho(x, c)|$, where $\|\cdot\|$ denotes the norm of the group element. Moreover there is a sequence $\{\varepsilon_n\}$, such that $\sum_n \mu(B(c; \varepsilon_n)) < \infty$, and

$$\limsup_{n\to\infty} \frac{\log \varepsilon_n^{-1}}{n \log \lambda} < 1.$$

(3) *Pole.* For each $c \in \mathcal{C}$ there is $p > 0$ so that on $B(c; \delta)$, $\|\varphi(x)\| \approx \rho(x, c)^{-p}$, and for some sequence $\{\varepsilon_n\}$, such that $\sum_n \mu(B(c; \varepsilon_n)) < \infty$, we have

$$\limsup_{n\to\infty} \frac{(p + 1) \log \varepsilon_n^{-1}}{n \log \lambda} < 1.$$

https://doi.org/10.1017/S0143385705000179 Published online by Cambridge University Press

In each case, we will assume that the Holder exponent of φ restricted to the complement of $\bigcup_{c\in\mathcal{C}} B(c;\delta)$ is γ. We state the following result.

THEOREM 2. *Assume that (M, f, μ) is a measure-preserving smooth Markov system as defined in §3, and $M_k \in \mathcal{P}_0$. Let $\varphi : M \to G$ be a Lie group valued observable which has a singularity set $\mathcal{C} \subset M \setminus \bigcup_k \overline{M_k}$, characterized by cases (1), (2) or (3) above. Let ψ be a μ-measurable solution of the cohomological equation*

$$\psi \circ f(x) = \varphi(x)\psi(x) \quad \mu\text{-a.e.}$$

Then there is a version $\tilde{\psi}$ of ψ, with $\tilde{\psi} = \psi$ μ-a.e. such that $\tilde{\psi}$ is α-Hölder on M_k, for some $\alpha \in (0, 1)$.

Remark 3. The condition that $\mathcal{C} \subset M \setminus \bigcup_k \overline{M_k}$ is unnecessarily strong, but is shared by many examples. For multimodal maps and Lorenz maps on the interval, induced maps F to a neighborhood Y of the critical point c are common constructions [**9**]. The interval Y has the decomposition $Y = \bigcup_i Y_i \bmod \mu$ where $F|Y_i = f^{r_i} : Y_i \to Y$ is monotone onto for an appropriate $r_i > 0$. For each i, $\overline{f^j(Y_i)} \not\ni c$ for $0 \le j < r_i$. Hence, if φ has only singularities at c, then using the above theorem and by the argument of Corollary 1, we can conclude that ψ has a Hölder version on Y.

Remark 4. Also in cases where $\mathcal{C} \subset M \setminus \bigcup_k \overline{M_k}$ fails, the proof below can still be used to get partial results. Given an element M_k the proof constructs an N and a component J (where $f^N(J) = M_k$) of the preimage $f^{-N}(M_k)$ such that ψ has a Hölder version on J. If J can be chosen such that $\overline{f^i(J)} \cap \mathcal{C} = \emptyset$ for $0 \le i < N$, then there is a version of ψ which is Hölder on M_k. If this condition is not satisfied then using the proof of Corollary 1 it is possible to show that for any $\varepsilon > 0$, M_k contains an open set $U \subset M_k$, $\mu(M_k \setminus U) < \varepsilon$, and ψ has a version which is Hölder on U.

Remark 5. The dependence of α on the exponent γ, the asymptotics of the sequence ε_n, and the type of singularity are apparent from the proof.

Remark 6. The same regularity is forced upon solutions ψ to (2), (3) or (4).

Proof. As in the proof of Theorem 1, choose any $\Lambda = M_k \in \mathcal{P}_0$ such that $\mu(\Lambda) > 0$ and let $(\hat{M}, \hat{f}, \hat{\mu})$ denote the natural extension of (M, f, μ). We have to check that there are sufficiently many backward paths that avoid passing too close to the singularity. We do this by using a Borel–Cantelli argument. Let $B_n = B(c; \varepsilon_n)$, then $\sum_{n\ge1} \mu(B_n) < \infty$, and hence we deduce that for $\hat{\mu}$-a.e. $\hat{x} \in \hat{M}$, there exists $N = N(\hat{x})$ such that $x_n \notin B_n$ for all $n \ge N$. Combining all these facts we obtain that for $\hat{\mu}$-a.e. backward orbit there exists $N(\hat{x})$ such that for all $n \ge N(\hat{x})$ the following hold simultaneously:

$$x_0 \in \Lambda \text{ and } f^n(\mathcal{P}_n[x_n]) = \Lambda; \tag{11}$$

$$\operatorname{diam}(f^{-n}(\Lambda) \cap \mathcal{P}_n[x_n]) \le \frac{1}{C(\hat{x})}\lambda^{-n}; \tag{12}$$

$$\rho(x_n, c) \ge 2\varepsilon_n. \tag{13}$$

The last two observations show that $\varphi \mid f^{-n}(\Lambda) \cap \mathcal{P}_n[x_n]$ is a Hölder function for $n \ge N$. We now consider the Hölder properties of φ in the cases that we are interested in.

https://doi.org/10.1017/S0143385705000179 Published online by Cambridge University Press

Suppose that for $n \geq N$ we have $y, z \in f^{-n}(\Lambda) \cap \mathcal{P}_n[x_n]$. In the case of a logarithmic singularity we have (inside $B(c;\delta)$)

$$d(\varphi(y),\varphi(z)) \approx |\log \rho(y,c) - \log \rho(z,c)| \leq \log\left(1 + \frac{\rho(y,z)}{\rho(z,c)}\right)$$
$$\leq \rho(y,z)^{1-\alpha} \frac{(\lambda^{-n}/C(\hat{x}))^{\alpha}}{\rho(z,c)} \leq \frac{C(\hat{x})^{-\alpha}}{\varepsilon_n \lambda^{n\alpha}} \rho(y,z)^{1-\alpha}$$

for some constant $\alpha > 0$. For $\iota > 0$ arbitrary, we then obtain the estimate

$$d(\varphi(y),\varphi(z)) \leq \tilde{C}(\hat{x})\rho(y,z)^{1-\tilde{\alpha}-\iota} \quad \text{with } \tilde{\alpha} = \limsup_{n\to\infty} \frac{\log \varepsilon_n^{-1}}{n \log \lambda} < 1.$$

Outside $B(c;\delta)$, the function φ will be γ-Hölder.

Now consider the case where φ has a finite-order pole. Arguing as in the case of a logarithmic singularity we obtain (inside $B(c,\delta)$)

$$d(\varphi(y),\varphi(z)) \approx |\rho(y,c)^{-p} - \rho(z,c)^{-p}|$$
$$\leq \max\{p\rho(z,c)^{-p-1}, p\rho(y,c)^{-p-1}\}\rho(y,z)$$
$$\leq \frac{C(\hat{x})^{-\alpha}}{(\varepsilon_n)^{p+1}\lambda^{n\alpha}} \rho(y,z)^{1-\alpha}$$

and hence for arbitrary $\iota > 0$ we obtain

$$d(\varphi(y),\varphi(z)) \leq \tilde{C}(\hat{x})\rho(y,z)^{1-\tilde{\alpha}-\iota} \quad \text{with } \tilde{\alpha} = \limsup_{n\to\infty} \frac{(p+1)\log \varepsilon_n^{-1}}{n \log \lambda} < 1.$$

Outside $B(c;\delta)$, φ will be γ-Hölder.

In the case of a bounded discontinuity, the Borel–Cantelli argument is simpler, since we only have to worry about $f^{-n}(\Lambda) \cap P_n[x_n]$ intersecting $\mathcal{C}$, which is impossible by the assumption that $\mathcal{C} \cap \overline{M}_k = \emptyset$.

So we have now proved that ψ has a Hölder version on $f^{-N}(\Lambda) \cap P_N[x_N]$. To show that $\psi \mid \Lambda$ has a Hölder version, we argue as Corollary 1. The fact that $\mathcal{C}$ is disjoint from each $\overline{M}_k$ implies that the version $\tilde{\psi}$ will be Hölder. □

4. *One-dimensional systems*

In this section, we consider C^2 one-dimensional systems for which ρ is Euclidean distance and μ is an invariant measure which is absolutely continuous with respect to Lebesgue. The assumption that there exists a function $K(\hat{x})$ and $\lambda > 0$ such that

$$\rho(y_n, z_n) \leq K(\hat{x})\lambda^{-n}\rho(y_0, z_0)$$

for all $n \geq 0$ and $y_n, z_n \in \mathcal{P}_n[x_n]$ can be replaced by the following two conditions which are commonly assumed in the literature.

(1) f is C^2 and has bounded distortion uniformly over all iterates: there exists a function $K(\hat{x})$ such that

$$\frac{|Df^n(y)|}{|Df^n(z)|} \leq K(\hat{x}) \tag{14}$$

for all $n \geq 0$ and $y, z \in \mathcal{P}_n[x_n]$.

https://doi.org/10.1017/S0143385705000179 Published online by Cambridge University Press

(2) Positive Lyapunov exponent, i.e. $\lambda(\mu) = \exp \int \log |Df| \, d\mu > 1$.

Instead of assuming (6) we may use (14) and $\lambda(\mu) > 1$ to prove the following lemma.

LEMMA 2. *For $\hat{\mu}$-a.e. $\hat{x} \in \hat{M}$, there exists a constant $C(\hat{x})$ such that*

$$\operatorname{diam}(\mathcal{P}_n[x_n]) \leq C(\hat{x})\lambda^{-n} \quad \textit{for all } n \geq 0. \tag{15}$$

Proof. The measure $\hat{\mu}$ is invariant in forward and backward time, in particular $\hat{\mu}(A) = \hat{\mu}(\hat{f}(A))$. Let $D\hat{f}^{-n}$ denote the derivative of f^{-n} restricted to an inverse branch. By the Birkhoff Ergodic Theorem,

$$\lim_{n\to\infty} \frac{1}{n} \log |D\hat{f}^{-n}(\hat{x})| = \int \log |D\hat{f}^{-1}| \, d\hat{\mu} = \int -\log |Df| \, d\mu = -\lambda(\mu)$$

for $\hat{\mu}$-a.e. $\hat{x}$, so $|Df^n(x_n)| \geq (1/C_0)\lambda^n$ for some $C_0 = C_0(\hat{x}) < \infty$. Using (14), we find that

$$\operatorname{diam}(\mathcal{P}_n[x_n]) \leq K \frac{|\mathcal{P}_0[x_0]|}{|Df^n(x_n)|} \leq C(\hat{x})\lambda^{-n}$$

as required. □

As a consequence of (14) for some $K > 0$

$$\frac{1}{K^2} \leq \frac{\rho(y_{i+1}, z_{i+1})\rho(\tilde{y}_0, \tilde{z}_0)}{\rho(\tilde{y}_{i+1}, \tilde{z}_{i+1})\rho(y_0, z_0)} \leq K^2 \tag{16}$$

for all $y_n, \tilde{y}_n, z_n, \tilde{z}_n \in \mathcal{P}_n[x_n]$. In dimension one it is sufficient to bound

$$\sum_{i=0}^{n-1} \|\operatorname{Ad}(\varphi_i(y_i))\| d(\varphi(y_{i+1}), \varphi(z_{i+1}))$$

by

$$\sum_{i=0}^{n-1} C \cdot K(\hat{x})(\mu_u)^{i+1}\lambda^{-(i+1)\alpha}\rho(y_0, z_0)^{\alpha}$$

in (9) of the proof of the main theorem. Using the two observations (15) and (16) the proof goes through as in Theorem 1.

4.1. *Smooth measures.* The assumption that the Jacobian J_μ is Hölder is used by, e.g., Young [**41**], and enables us to apply the technique to Gibbs measures and equilibrium states of suitable potentials. However, in the case that μ is absolutely continuous with respect to Lebesgue measure, it suffices to assume that the density $h \in L^1(\text{Leb})$. Indeed, (10) can be derived as follows. Let $\varepsilon > 0$ and $\eta > 0$. Choose n sufficiently large that $\text{Leb}(\mathcal{P}_n[x_n] \cap \{y \in \mathcal{P}_n[x_n] : d(\psi(x_n), \psi(y)) < \eta\}^c) \leq \eta \, \text{Leb}(\mathcal{P}_n[x_n])$. Boundedness of distortion gives that

$$\frac{\text{Leb}(f^n(\mathcal{P}_n[x_n] \cap \{y \in \mathcal{P}_n[x_n] : d(\psi(x_n), \psi(y)) < \eta\}^c))}{\text{Leb}(f^n(\mathcal{P}_n[x_n]))} \leq \mathcal{O}(1) \, \frac{\text{Leb}(\mathcal{P}_n[x_n] \cap \{y \in \mathcal{P}_n[x_n] : d(\psi(x_n), \psi(y)) < \eta\}^c)}{\text{Leb}(\mathcal{P}_n[x_n])}.$$

https://doi.org/10.1017/S0143385705000179 Published online by Cambridge University Press

Now write

$$\begin{aligned}\mu(\Lambda) &= \mu(f^n(\mathcal{P}_n[x_n] \cap \{y \in \mathcal{P}_n[x_n] : d(\psi(x_n), \psi(y)) < \eta\})) \\ &\quad + \mu(f^n(\mathcal{P}_n[x_n] \cap \{y \in \mathcal{P}_n[x_n] : d(\psi(x_n), \psi(y)) < \eta\}^c)) \\ &= \int_{f^n(\mathcal{P}_n[x_n] \cap \{y \in \mathcal{P}_n[x_n] : d(\psi(x_n), \psi(y)) < \eta\})} h(x)\,dx \\ &\quad + \int_{f^n(\mathcal{P}_n[x_n] \cap \{y \in \mathcal{P}_n[x_n] : d(\psi(x_n), \psi(y)) < \eta\}^c)} h(x)\,dx\end{aligned}$$

and note that since the density $h(x) \in L^1(\text{Leb})$, and $\text{Leb}(f^n(\mathcal{P}_n[x_n] \cap \{y \in \mathcal{P}_n[x_n] : d(\psi(x_n), \psi(y)) < \eta\}^c)) \le \mathcal{O}(\eta)\,\text{Leb}(\Lambda)$ we can assume that $\eta > 0$ is such that $\mu(f^n(\mathcal{P}_n[x_n] \cap \{y \in \mathcal{P}_n[x_n] : d(\psi(x_n), \psi(y)) < \varepsilon\}^c)) \le \varepsilon\mu(\Lambda)$. Here we have used the fact that for a $L^1(\text{Leb})$ function g: given $\delta_1 > 0$, there exists a $\delta_2 > 0$ so that $\text{Leb}(A) < \delta_2$ implies $\int_A g < \delta_1$.

5. *Refinements and applications*

5.1. *Young towers.* We will show that Theorem 1 implies Hölder regularity for measurable solutions to a broad class of cohomological equations on Young towers that arise in applications.

Suppose that $T : X \to X$ is a $C^{1+\gamma}$ mapping of a Riemannian manifold X and Leb denotes Lebesgue measure. Let ρ_X denote the corresponding metric. A Young tower for T has the following properties.

- There exists a set $\Lambda \subset X$, decomposed as $\Lambda = \bigcup_j \Lambda_j \mod \text{Leb}$.
- For each j, there exists $R_j \ge 1$ such that $T^{R_j} : \Lambda_j \to \Lambda$ is bijective. Denote the *induced map* $T^{R_j}|\Lambda_j$ by F.
- The distortion is bounded, i.e. there exists $K < \infty$ such that for all $n \ge 0$

$$\left|\frac{\text{Jac}\, DF_x^n}{\text{Jac}\, DF_y^n}\right| \le K$$

  for all $x, y \in U$ and sets U on which F^n is a diffeomorphism.
- There exists $\lambda_0 > 1$ such that $\min_{|v|=1} |DF_x v|/|v| \ge \lambda_0$ for $x \in \Lambda$.

The Folklore theorem [**26**] states that F has an invariant probability μ, which is equivalent to Lebesgue and the Radon–Nikodym derivative h is bounded and bounded away from 0.

The measure μ can be pulled back to a T-invariant measure ν:

$$\nu(A) = \sum_j \sum_{i=0}^{R_j-1} \mu(T^{-i}A \cap \Lambda_j). \tag{17}$$

The measure ν is finite if and only if

$$\mathcal{R} := \int_\Lambda R\,d\mu = \sum_j R_j \mu(\Lambda_j) < \infty. \tag{18}$$

This set-up can be viewed as a Markov system as follows.

https://doi.org/10.1017/S0143385705000179 Published online by Cambridge University Press

- M is the disjoint union $\bigsqcup_j \bigsqcup_{i=0}^{R_j-1}(\Lambda_j, i)$ where each (Λ_j, i) is a copy of Λ_j. The set $\Lambda = \bigcup_j(\Lambda_j, 0)$ is called the *base* of the tower. Each set (Λ_j, i) is a component M_k of M.
- For the metric ρ on M, there are at least two choices. Take $(x, i) \in (\Lambda_j, i)$ and $(\tilde{x}, \tilde{i}) \in (\Lambda_{\tilde{j}}, \tilde{i})$
  - $\rho_1((x, i), (\tilde{x}, \tilde{i})) = \begin{cases} \rho_X(T^i(x), T^{\tilde{i}}(\tilde{x})) & \text{if } j = \tilde{j} \text{ and } i = \tilde{i}; \\ 1 & \text{otherwise.} \end{cases}$
    This metric is induced from the metric ρ_X on X. The metric ρ_1 is used in Corollary 2.
  - $\rho_2((x, i), (\tilde{x}, \tilde{i})) = \begin{cases} \rho_X(x, \tilde{x}) & \text{if } j = \tilde{j} \text{ and } i = \tilde{i}; \\ 1 & \text{otherwise.} \end{cases}$
    This metric is the *tower metric* and is induced from the metric on the base Λ.
- Define $f : M \to M$ as

$$f(x, i) = \begin{cases} (x, i+1) & \text{if } x \in \Lambda_j \text{ and } i < R_j - 1, \\ (T^{R_j}x, 0) & \text{if } x \in \Lambda_j \text{ and } i = R_j - 1. \end{cases}$$

  There is a projection $\pi : M \to X$ given by $\pi(x, i) = T^i(x)$ and $\pi \circ f = T \circ \pi$.
- Extend the definition of μ from Λ to M as $\mu((A, i)) = \mu(A)$ for each measurable set $A \subset \Lambda_j$ and $0 \le i \le R_j - 1$. By (18) μ is finite and hence can be normalized.

Let $\mathcal{P}_0$ denote the partition of M into sets (Λ_j, i) and set $\mathcal{P}_n = \bigvee_{i=0}^{n-1} f^{-i}(\mathcal{P}_0)$. For each $x \in M$ let $\mathcal{P}_n[x]$ be the partition element (cylinder set) in $\mathcal{P}_n$ containing x. Let $(\hat{M}, \hat{f}, \hat{\mu})$ denote the natural extension of (M, f, μ). For $\hat{x} = (x_0, x_1, \dots, x_n, \dots)$ let $n_0 < n_1 < \cdots$ denote the indices such that x_{n_i} belongs to the base.

LEMMA 3. *Let (M, f, μ) be a Young tower satisfying (18). Assume one of the following three conditions:*

- *$\rho = \rho_1$ and there is $\delta_0 > 0$ such that*

$$\|DT^{R_j-k}(T^k(x))\| \ge \delta_0 \quad \text{for all } j, x \in \Lambda_j \text{ and } 0 \le k < R_j; \tag{19}$$

- *$\rho = \rho_2$; or*
- *$\log^+ \|Df^{-1}\| \in L^1(\mu)$, where the derivative is taken with respect to the metric used.*

Then for $\hat{\mu}$-a.e. $\hat{x}$ there exists a number $K(\hat{x})$ and $\lambda > 1$ such that

$$\rho(y_n, z_n) < K(\hat{x})\lambda^{-n}\rho(y_0, z_0)$$

for all $n \ge 0$ and $y_n, z_n \in \mathcal{P}_n[x_n]$.

Proof. Write $R(x) = R_j$ for $x \in \Lambda_j$, so by Birkhoff's and Kac's theorems $\mathcal{R}^{-1} = \lim_{k\to\infty}(1/k)\sum_{i=1}^{k} R(x_{n_i})$ for $\hat{\mu}$-a.e. $\hat{x}$. Let $\tilde{\Lambda}_i = \cup\{\Lambda_j : R_j = i\}$, so (18) gives

https://doi.org/10.1017/S0143385705000179 Published online by Cambridge University Press

$\sum_i i\mu(\tilde{\Lambda}_i) < \infty$. Given $\varepsilon > 0$, we have

$$\begin{aligned}\sum_k \hat{\mu}\{\hat{x} : R(x_{n_{k+1}}) > \varepsilon n_k\} &\le \sum_k \hat{\mu}\Big\{\hat{x} : x_{n_{k+1}} \in \bigcup_{i\ge\varepsilon k} \tilde{\Lambda}_i\Big\}\\ &\le \sum_k \sum_{i\ge\varepsilon k} \mu(\tilde{\Lambda}_i)\\ &\le \left\lceil \frac{1}{\varepsilon} \right\rceil \sum_k k\mu(\tilde{\Lambda}_k) < \infty.\end{aligned}$$

The Borel–Cantelli lemma therefore implies that for $\hat{\mu}$-a.e. $\hat{x}$, there is $k_0 = k_0(\hat{x})$ such that $n_{k+1} - n_0 \le (1+\varepsilon)(n_k - n_0)$ for all $k \ge k_0$. Take $n_k \le n < n_{k+1}$, then

$$\|Df^{-(n-n_k)}(x_{n_k})\| \le \begin{cases}\delta_0^{-1} & \text{if } \rho = \rho_1 \text{ by (19);}\\ \lambda_0^{-1} < 1 & \text{if } \rho = \rho_2.\end{cases}$$

Write $B(\hat{x}) = \|Df^{-n_0}(x_0)\|$. Direct calculation gives

$$\begin{aligned}\lim_{n\to\infty}\left(\left\|\prod_{i=0}^{n-1} Df^{-1}(x_i)\right\|\right)^{1/n} &\le \lim_{k\to\infty}\left(\frac{B(\hat{x})}{\delta_0}\left\|\prod_{i=0}^{n-1} Df^{-1}(x_i)\right\|\right)^{1/(1+\varepsilon)n_k}\\ &= \lim_{k\to\infty}\left(\frac{B(\hat{x})}{\delta_0}\left\|\prod_{i=1}^{k} DF^{-1}(x_{n_i})\right\|\right)^{1/(n_0+(1+\varepsilon)\sum_{i=1}^k R(x_{n_i}))}\\ &= \lim_{k\to\infty}\left(\left(\frac{B(\hat{x})}{\delta_0}\left\|\prod_{i=1}^{k} DF^{-1}(x_{n_i})\right\|\right)^{1/k}\right)^{k/(n_0+(1+\varepsilon)\sum_{i=1}^k R(x_{n_i}))}\\ &\le \lambda_0^{-1/(1+\varepsilon)\mathcal{R}}. \qquad (20)\end{aligned}$$

As $\varepsilon > 0$ was arbitrary, we get $\lim_{n\to\infty}\left(\|\prod_{i=0}^{n-1} Df^{-1}(x_i)\|\right)^{1/n} \le \lambda_0^{-1/\mathcal{R}}$ as well.

Alternatively, assume that $\log^+\|Df^{-1}\| \in L^1(\mu)$. The estimate of (20) holds for the sequence (n_k). Oseledec's Multiplicative Theorem then implies that $\lim_{n\to\infty}\left(\|\prod_{i=0}^{n-1} Df^{-1}(x_i)\|\right)^{1/n}$ exists $\hat{\mu}$-a.e. $\hat{x}$. This limit equals the limit along the subsequence, so

$$\lim_{n\to\infty}\left(\left\|\prod_{i=0}^{n-1} Df^{-1}(x_i)\right\|\right)^{1/n} \le \lambda_0^{1/\mathcal{R}}.$$

Since these estimates hold uniformly over $y_i, z_i \in \mathcal{P}_i[x_i]$, we obtain $\rho(y_n, z_n) < K(\hat{x})\lambda^{-n}\rho(y_0, z_0)$ for $\lambda = \lambda_0^{1/\mathcal{R}}$. □

Remark 7. The condition $\log^+\|Df^{-1}\| \in L^1(\mu)$ relates to the reference measure m on X as follows: if μ has density h, and $h \in L^{1+\varepsilon}(m)$ for some $\varepsilon > 0$, then by the Hölder inequality, $\log^+\|Df^{-1}\| \in L^q(m)$ for $q = (1+\varepsilon)/\varepsilon$ implies that $\log^+\|Df^{-1}\| \in L^1(\mu)$.

THEOREM 3. *Let (M, f, μ) be a Young tower with base Λ, satisfying (18). Assume that $\log^+\|Df^{-1}\| \in L^1(\mu)$ or that (19) holds. Let $\varphi : M \to G$ be a Lie group valued α-Hölder observable and define*

$$\mu_u := \lim_{n\to\infty}\Big(\sup_{(x,0)\in M} \|\mathrm{Ad}\,(\varphi(f^{n-1}(x,0)\dots\varphi(f(x,0))\varphi((x,0))))\|\Big)^{1/n}.$$

https://doi.org/10.1017/S0143385705000179 Published online by Cambridge University Press

If $1 \leq \mu_u < \lambda_0^{\alpha/\mathcal{R}}$, *then any* μ*-measurable solution to the cohomological equation*

$$\varphi = (\psi \circ f)\psi^{-1} \quad \mu\text{-a.e.}$$

has a version which is α*-Hölder on* Λ.

Remark 8. If G is Abelian, compact or nilpotent then $\mu_u = 1$ and the spectral condition $1 \leq \mu_u < \lambda_0^{\alpha/\mathcal{R}}$ is automatically satisfied.

Remark 9. For any n there exists an α-Hölder version on $\bigcup_{j=0}^{n} f^j(\Lambda)$ but the Hölder constant may increase with n.

Remark 10. The same result holds for solutions ψ to (2), (3) or (4).

Proof. First observe that μ is an ergodic f-invariant measure. If $\rho = \rho_2$, and $x \in \Lambda_i$, then $Df(f^i(x)) = \mathrm{Id}$ if $i < R_i - 1$ and $Df(f^{R_i(x)}) = DF(x)$. If $\rho = \rho_1$, then $Df(f^i(x)) = DT(T^i(x))$. In either case, $Df(f^{R_i(x)}) = DF(x)$.

A computation similar to (20) yields that for μ-a.e. $x \in \Lambda$:

$$\lambda(\mu) := \lim_{n\to\infty} \left(\left\| \prod_{i=0}^{n-1} Df(f^i(x,0)) \right\| \right)^{1/n} = \lambda_0^{1/\mathcal{R}}. \tag{21}$$

Therefore, the assumption $\mu_u < \lambda_0^{\alpha/\mathcal{R}}$ implies partial hyperbolicity (PH). Lemma 3 shows that condition (6) holds. Hence, the proof of Theorem 1 applies and we obtain a Hölder version on each (Λ_j, i). Using the argument from Corollary 1 we obtain Livšic regularity on the base Λ as well. □

COROLLARY 2. *Suppose that* (T, X) *is modelled by a Young tower* (M, f, μ) *over base* $\Lambda \subset X$, *satisfying (18). Assume that* $\log^+ \|Df^{-1}\| \in L^1(\mu)$ *or that (19) holds. Let* ν *be the* T *invariant ergodic pulled back measure, as in (17). Let* $\tilde{\varphi} : X \to G$ *be* α*-Hölder and satisfy* $\mu_u < \lambda_0^{\alpha/\mathcal{R}}$. *Then for any* $0 \leq k \leq \mathcal{R}_j$, *any* ν*-measurable solution to the cohomological equation* $\tilde{\varphi} = (\tilde{\psi} \circ T)\tilde{\psi}^{-1}$ *has a version which is Hölder on* $T^k(\Lambda_j)$.

Proof. Suppose that $\tilde{\varphi} : X \to G$ is Hölder and $\tilde{\varphi} = (\tilde{\psi} \circ T)\tilde{\psi}^{-1}$, ν-a.e. Let $\pi : M \to X$ be defined as $\pi((x,i)) = T^i(x)$, so that $T \circ \pi = \pi \circ f$. Then $\tilde{\varphi}, \tilde{\psi} : X \to G$ lift to the tower as $\varphi = \tilde{\varphi} \circ \pi$, $\psi = \tilde{\psi} \circ \pi$ and satisfy $\varphi = (\psi \circ f)\psi^{-1}$, μ-a.e. Moreover, φ is α-Hölder with respect to the metric ρ_1. Since derivatives $Df(f^i(x))$ on the tower agree with derivatives $DT(T^i(x))$ on (X, T),

$$\left(\left\| \prod_{i=0}^{n-1} Df(f^i(x)) \right\| \right)^{1/n} \geq \lambda_0^{\alpha/\mathcal{R}} > \mu_u$$

for μ-a.e. x, verifying (PH) on (M, f, μ).

Now Theorem 3 gives α-Hölder (with respect to ρ_1) solution ψ' to the cohomological equation $\varphi = (\psi' \circ T)\psi'^{-1}$, such that $\psi' = \psi$ μ-a.e. Since ψ' takes the same value on every point of $\pi^{-1}(x)$ for ν-a.e. x the projection $\tilde{\psi}' = \psi \circ \pi$ is well defined, α-Hölder and satisfies $\tilde{\varphi} = (\tilde{\psi}' \circ T)\tilde{\psi}'^{-1}$, ν-a.e. □

https://doi.org/10.1017/S0143385705000179 Published online by Cambridge University Press

Remark 11. A priori, a larger class of observables $\varphi : M \to G$ is Hölder with respect to the tower metric ρ_2 than with respect to ρ_1. For metric ρ_2, however, the projection $\pi : M \to X$, $(x, i) \mapsto T^i(x)$ need not preserve the Hölderness of solutions of the cohomologous equation on (M, f). If the projection π preserves continuity or is only used on the base Λ, this may still suffice for applications.

5.2. *The Manneville–Pomeau family.* In the previous results, the assumptions that μ is finite and/or has positive Lyapunov exponents can be weakened for some group extensions.

For the Manneville–Pomeau family the Jacobian J_μ of μ is Hölder, i.e. there exists C and $\gamma \in (0, 1]$ such that

$$\left| \frac{J_\mu(x)}{J_\mu(y)} - 1 \right| \le C \cdot \rho(f(x), f(y))^\gamma. \tag{22}$$

To prove Lemma 1, assumption (6) can be weakened to

$$\sum_{n\ge 0} \rho(y_n, z_n)^\gamma < K(\hat{x})\rho(y_0, z_0)^\gamma, \tag{23}$$

where $\gamma > 0$ is the Hölder exponent in (8). In this case $\lambda(\mu)$ can be 0, and (PH) fails, but if G is Abelian or compact, or if G is nilpotent with a dominated growth rate so that for inverse branches

$$\sum_{n\ge 0} \|\mathrm{Ad}\,(\varphi_n(y_0))\| d(\varphi(y_{n+1}), \varphi(x_{n+1}))^\alpha < \infty, \tag{24}$$

then the estimates (9) hold. Furthermore, the pulled back measure ν can be at most σ-finite if (18) fails. According to (21), this implies that $\lambda(\mu) \le 1$.

This scenario is found in the well-known Manneville–Pomeau maps. This is a family of maps on $[0, 1]$ which have a neutral fixed point (where we take ρ to be Euclidean distance), parameterized by parameter $p \in (0, \infty)$. For $p \ge 1$ these maps admit a σ-finite absolutely continuous invariant measure. However, the measure is not a probability measure, and the map has zero Lyapunov exponents for Lebesgue almost all initial conditions. We have the following regularity result in this setting.

THEOREM 4. *Consider the Manneville–Pomeau map (for $p \ge 0$):*

$$T(x) = \begin{cases} x + 2^p x^{1+p} & \text{if } x \in [0, \frac{1}{2}), \\ 2x - 1 & \text{if } x \in [\frac{1}{2}, 1]. \end{cases}$$

If G is an Abelian or compact group and $\tilde{\varphi} : [0, 1] \to G$ is a G-valued α-Hölder observable for $\alpha > p/(1 + p)$, then any Lebesgue measurable solution to the cohomological equation $\varphi = (\psi \circ T)\psi^{-1}$ has a version which is α-Hölder on $[0, 1]$.

Remark 12. The same regularity is forced upon solutions ψ to (2), (3) or (4).

Proof. For $p \in [0, 1)$, T has an invariant probability $\nu \ll$ Leb, whereas ν is only σ-finite if $p \ge 1$. However, the first return map F to $[\frac{1}{2}, 1]$ has always an invariant probability $\mu \ll$ Leb, and F^n has bounded distortion, independently of n. Therefore, we can use Theorem 3

https://doi.org/10.1017/S0143385705000179 Published online by Cambridge University Press

provided we can show that $\varphi|(\Lambda_j, i) = \tilde{\varphi} \circ \pi^{-1}|\Lambda_j = \varphi \circ T^i|\Lambda_j$ is Hölder. This is done as follows. It is not hard to check that $T^{-n}(\frac{1}{2}) = \frac{1}{2}(pn)^{-1/p} + o(n^{-1/p})$, where T^{-n} indicates the nth inverse of the left branch of T. Consequently, $\operatorname{diam}(T^{-n}(J)) = \mathcal{O}(n^{-(1+p)/p})$.

Even if ν is σ-finite, $\nu([\frac{1}{2}, 1]) < \infty$. Therefore, the subset $\hat{M}_0 := \{\hat{x} \in \hat{M} : x_0 \in [\frac{1}{2}, 1]\}$ has finite $\hat{\mu}$-measure. Suppose that $\hat{x} \in \hat{M}_0$ is a backward orbit (chosen as in Theorem 1), and $\hat{y}, \hat{z} \in M$ are such that $y_n, z_n \in \mathcal{P}_n[x_n]$ for each n. Let $n_0 = 0$ and $n_k = \min\{n > n_{k-1} : x_n \in \Lambda = [\frac{1}{2}, 1]\}$. Since $|DF| \geq 2$, $\rho(y_{n_k}, z_{n_k}) \leq 2^{-k}$ and, hence, using the fact that $\alpha > p/(1+p)$,

$$\begin{aligned}
\sum_{n\geq 0} d(\varphi(y_n), \varphi(z_n)) &\leq \sum_{k\geq 0} \sum_{n=n_k}^{n_{k+1}-1} \mathcal{O}(1)\rho(y_n, z_n)^\alpha \\
&\leq \sum_{k\geq 0} \rho(y_0, z_0)^\alpha 2^{-\alpha k} \sum_{n=0}^{n_{k+1}-n_k-1} \mathcal{O}(1)\operatorname{diam}(T^{-n}([\tfrac{1}{2}, 1]))^\alpha \\
&\leq \rho(y_0, z_0)^\alpha \sum_{k\geq 0} 2^{-\alpha k} \sum_{n=0}^{n_{k+1}-n_k-1} \mathcal{O}(1) n^{-\alpha(1+p)/p} \\
&\leq \mathcal{O}(1)\rho(y_0, z_0)^\alpha.
\end{aligned}$$

This calculation replaces (9) in the proof of Theorem 1. Continuing the proof as in Theorem 1, we get that ψ has an α-Hölder version on $[\frac{1}{2}, 1]$. As $T : [\frac{1}{2}, 1] \to [0, 1]$ is smooth by using the cohomological equation $\psi \circ T(x) = \varphi(x) \cdot \psi(x)$, it follows that ψ has an α-Hölder version on $[0, 1]$, cf. Corollary 1. □

Remark 13. The arguments above can also be used to establish Hölder regularity for nilpotent groups G, provided that (24) holds. Having $\mu_u = 1$ alone is not enough to establish Livšic regularity, but if $\|\operatorname{Ad}(\varphi_n(y_0))\|$ grows at a rate 'dominated' by the polynomial contraction of $d(\varphi(y_{n+1}), \varphi(x_{n+1}))^\alpha$, which in our case is $\mathcal{O}(n^{-\alpha(1+p)/p})$, then the conclusion of Theorem 4 will remain valid.

5.3. *Interval maps with critical points.* So far, the Markov systems used in the examples were Young towers, even though we allowed a σ-finite measure for the Manneville–Pomeau map. Theorem 1 also applies to different kinds of Markov systems. In this section we discuss the consequences of the theory to systems modelled by so-called Hofbauer towers. The metric ρ is Euclidean distance in this section.

An interval map $T : [0, 1] \to [0, 1]$ is called piecewise continuous (piecewise C^r) if there exists a finite set of points $0 = a_0 < a_1 < \cdots < a_{k-1} < a_k = 1$ such that $T|(a_{i-1}, a_i)$ has a continuous (C^r) extension to $[a_{i-1}, a_i]$. We call $a_0, \ldots, a_k$ the *critical* points of T and denote this set by $\mathcal{C}$. Let $\mathcal{P}_1$ denote the partition $\{[a_0, a_1], \ldots, [a_{k-1}, a_k]\}$ and let $\mathcal{P}_n = \bigvee_{i=0}^{n-1} T^{-i}(\mathcal{P}_1)$ be the partition into n-cylinders. For $x \in I \setminus \bigcup_{i=0}^{n-1} T^{-i}(\mathcal{C})$, let $\mathcal{P}_n[x]$ denote the n-cylinder containing x.

Due to the presence of critical points, T^{-1} need not be differentiable at boundary points of $T(P)$, $P \in \mathcal{P}_1$, and hence boundedness of distortion (i.e. condition (14)) cannot be realized globally. For this reason, we call J a *core* interval if it is compactly contained

https://doi.org/10.1017/S0143385705000179 Published online by Cambridge University Press

in $T^n(P)$ for some $n \geq 1$ and $P \in \mathcal{P}_n$. For example, if $T(x) = 1 - ax^2$ is a non-renormalizable unimodal map, then any interval compactly contained in $[T^2(0), T(0)]$ is a core interval.

THEOREM 5. *Let T be a piecewise C^3 interval map onto the unit interval with negative Schwarzian derivative. Let ν be a T-invariant probability measure such that $\int \log|T'|d\nu > 0$. Assume that the Jacobian J_ν is Hölder or $\nu \ll$ Leb. Let φ be a Lie group valued piecewise α-Hölder observable, with discontinuities (if any) only at the points a_i, and satisfying the partial hyperbolicity condition (PH). If $\varphi = (\psi \circ T)\psi^{-1}$ for some ν-measurable function, then for every* core *interval J with $\nu(J) > 0$, $\psi|J$ has an α-Hölder version.*

Remark 14. Assume that the core interval J is compactly contained in $T^n(P)$ for some $P \in \mathcal{P}_n$, and $n \geq 1$. Although the Hölder exponent is independent of J, the Hölder coefficient in the Hölder property of the Hölder version of $\psi|J$ depends on $\min\{\operatorname{diam}(J)/\operatorname{diam}(L), \operatorname{diam}(J)/\operatorname{diam}(R)\}$, where L and R are the components of $T^n(P) \setminus J$. As this minimum tends to 0, the Hölder coefficient tends to infinity.

Remark 15. The assumption that T is onto is not a severe restriction. Since ν is assumed to have a positive Lyapunov exponent, ν cannot be supported on a non-repelling periodic orbit, so $\bigcap_j T^j([0,1])$ is a non-trivial interval. For the same reason, ν cannot be supported on a wandering interval, i.e. an interval W such that $T^n(W) \bigcap T^m(W) = \emptyset$ for all $n \neq m \geq 0$. By restricting and rescaling, we can assume that $T : [0,1] \to [0,1]$ is onto.

Remark 16. The same result holds for solutions ψ to (2), (3) or (4).

Proof. We construct a Markov system, introduced by Hofbauer [**17**] as the *canonical Markov extension* and sometimes called a *Hofbauer tower*. This is the system (M, f), where M is a disjoint union of closed intervals. We call $B = [0,1]$ the *base of the tower*. Then

$$M = B \bigsqcup \Big(\bigsqcup_{n \geq 1} \bigsqcup_{P \in \mathcal{P}_n} \overline{T^n(P)} \Big) \Big/ \sim$$

where the sets $T^n(P) \sim T^{n'}(P')$ if they are the same interval. Let $\pi : M \to [0,1]$ be the natural projection. The action f is defined on M as follows. If $x \in M$ belongs to the component D, then

$$f(x) = \pi^{-1}(T(\pi(x))) \cap \tilde{D},$$

where component $\tilde{D} := \overline{T(D \cap \mathcal{P}_1[\pi(x)])}$ is again a component of M. Obviously (M, f) is Markov and $T \circ \pi = \pi \circ f$. Due to the Markov property, the following holds for any component D of M:

$$T^n(\mathcal{P}_n[x]) = \pi(D) \quad \text{if and only if } f^n(\pi^{-1}(x) \cap B) \in D.$$

If ν is T-invariant, then we can construct a measure μ as follows. Let μ_0 be the measure ν lifted to the level B and set $\mu_n = (1/(n+1)) \sum_{i=0}^{n} \mu_0 \circ f^{-i}$. Clearly $\nu = \mu_n \circ \pi^{-1}$ for each n. As shown in [**22**], μ_n converges vaguely. We call the limit measure μ. If ν is ergodic, then μ is either a probability measure on M, in which case we call μ *liftable*,

https://doi.org/10.1017/S0143385705000179 Published online by Cambridge University Press

or it is identically 0 on M. In this case the mass 'has escaped to infinity'. Keller's result [**22**, Theorem 3] states if $\int \log |T'| \, d\nu > 0$, then ν is liftable to an invariant measure μ on the Markov extension. Moreover, μ is ergodic if ν is.

If J is a core interval and $\nu(J) > 0$, then there is some level $D \in M$ compactly containing a lifted copy $\tilde{J} := D \cap \pi^{-1}(J)$, and $\mu(\tilde{J}) > 0$. Let $\delta > 0$ be such that D contains a $\delta|J|$-neighbourhood of $\tilde{J}$. Using negative Schwarzian derivative and the Koebe principle, see [**27**], we find that for every $x \in M$ with $f^n(x) \in \tilde{J}$, that f^n has bounded distortion on the component of $f^{-n}(\tilde{J})$ containing x. In fact, the distortion depends only on δ.

Finally, we will show that $\psi|J$ has a Hölder version. $\tilde{\varphi} = \varphi \circ \pi$ is an observable on the Markov extension. The coboundary ψ lifts to a coboundary $\tilde{\psi} = \psi \circ \pi$. Therefore, we can apply Theorem 1 to it to find a version of $\tilde{\psi}$ that is α-Hölder. Projecting it back to the interval, we find the desired α-Hölder version of ψ. (Note that since $\tilde{\psi}$ takes the same value on every point in $\pi^{-1}(x)$, we find that the Hölder version of ψ does not depend on the level $D \in M$ that we lift J to.) □

Let $T : I \to I$ be a C^3 S-multimodal map having critical set $\mathcal{C}$. Assume each critical point $c \in \mathcal{C}$ has order ℓ_c, $1 < \ell_c < \infty$. We assume for simplicity that T is locally eventually onto, i.e. for every non-degenerate subinterval $U \subset I$, $T^n(U) = I$ for some $n \geq 0$. This excludes that g is renormalizable, or has a non-expanding periodic orbit. (Also wandering intervals are excluded, but this is already a corollary of the smoothness, see [**27**].) In this case the T-invariant measure ν that we will be considering is supported on I.

THEOREM 6. *Assume that $T : I \to I$ is C^3 multimodal with negative Schwarzian derivative and non-flat critical points. Write $c_n = T^n(c)$ and $\ell_{\max} = \max\{\ell_c : c \in \mathcal{C}\}$. Assume that T satisfies the summability condition*

$$\sum_{c \in \mathcal{C}} \sum_{k \geq 1} |DT^{k-1}(c_1)|^{-1/\ell_{\max}} < \infty, \tag{25}$$

and hence possesses an acip ν (cf. [**10, 32**]*). Let φ be a piecewise α-Hölder $L^1(\nu)$ observable, with discontinuities of types (1)–(3) of §3.1, at critical points only. If $\varphi = \psi \circ T - \psi$ for some ν-measurable function, then for every core interval J, $\psi|J$ is $\tilde{\alpha}$-Hölder for any $\tilde{\alpha} \in (0, \alpha)$.*

Proof. Assume as above that T is locally eventually onto. We build a Markov extension (M, f) as in Theorem 5. Since T admits an acip ν (with necessarily positive Lyapunov exponent, cf. [**21**] and also [**8**]), it can be lifted to an acip μ on (M, f). Let D be any level in M that compactly contains a lifted copy $\tilde{J} = D \cap \pi^{-1}(J)$ of J and such that $\nu(J) > 0$.

Let $(\hat{M}, \hat{f}, \hat{\mu})$ be the natural extension of (M, f, μ). By the Koebe principle, we have a uniform distortion bound for $f^n|\mathcal{P}_n[x_n] \cap f^{-n}(J)$ for each $x \in \tilde{J}$ (i.e. J is a core interval as introduced before).

The next thing to check is that there are sufficiently many backward paths that avoid passing close to the singularities of φ at $\mathcal{C}$. This argument is similar to that in the proof of Theorem 2. The proof that (I, f) has an acip was given in the multimodal case in [**10**],

https://doi.org/10.1017/S0143385705000179 Published online by Cambridge University Press

based on the well-known result of Nowicki and van Strien [**32**]. An important estimate in [**10**] is that for some $C = C(T) < \infty$,

$$\nu(A) \le C \cdot \mathrm{Leb}(A)^{1/\ell} \quad \text{for all measurable } A \subset I.$$

It follows that if $B_n = (c - n^{-2\ell}, c + n^{-2\ell})$, then

$$\nu(B_n) \le C \cdot \mathrm{Leb}(B_n)^{1/\ell} \le \frac{2C}{n^2},$$

and hence, lifted to the tower

$$\mu(T^{-n}(\tilde{J}) \cap B_n) \le \nu(B_n) \le \frac{2C}{n^2}.$$

It follows from the Borel–Cantelli lemma that for $\hat{\mu}$-a.e. $\hat{x} \in \hat{M}$, there exists $N = N(\bar{x})$ such that $x_n \notin B_n$ for all $n \ge N$. From now on, the argument is the same as in Theorem 2. □

In view of questions raised in, e.g., [**11**], we are particularly interested in the potential $\varphi := \int_I \log|T'|\,d\mu - \log|T'|$. This potential is smooth, except for logarithmic singularities at the critical points. Theorem 6 shows that any ν-measurable solution of the cohomological equation $\varphi = \psi - \psi \circ T$ has a version which is α-Hölder for any $\alpha \in (0, 1)$ on each interval that is compactly contained in $[c_2, c_1]$. We can apply it to the quadratic family $f_a(x) = 1 - ax^2$. It is known that for Leb-a.e. $a \in [0, 2]$, f_a has either an attracting periodic orbit, or has a positive Lyapunov exponent at the critical value, and both parameter sets have positive measure. Below we make a weaker assumption on the growth rate of derivatives along the critical orbit.

COROLLARY 3. *Let $f : I \to I$ be a C^3 S-unimodal map with critical order $\ell < \infty$, satisfying the summability condition $\sum_n |Df^n(c_1)|^{-1/\ell} < \infty$ (and hence possessing an acip ν). Then $\varphi = \log|f'| - \int \log|f'|\,d\nu$ can only be a measurable coboundary if there exists a periodic interval $J \subseteq I$ of period k such that $c \in J$ and $J = [f^{2k}(c), f^k(c)]$.*

Proof. First assume that f is non-renormalizable. In this case, ψ is Hölder continuous and hence bounded on any interval compactly contained in $[c_2, c_1]$. Since φ is bounded except at c, we easily derive that ψ has to be unbounded at every forward image of the critical point. This is only possible if $\bigcup_{n\ge1} f^n(c) \subset \{c_1, c_2\}$.

Next assume that f is finitely renormalizable, say $J \ni c$ is a periodic interval, $f^k(J) \subset J$, $f^k(\partial J) \subset \partial J$ where $k \ge 2$ is the period of renormalization, and $f^k|J$ is unimodal and non-renormalizable. Then the above argument shows that $\bigcup_{n\ge1} f^n(c) \cap J^\circ = \emptyset$. Therefore, $J = [f^{2k}(c), f^k(c)]$, and $f^k|J$ is conjugate to $x \mapsto 1 - 2x^2$. □

In the context of parametrized S-unimodal families f_a, the conclusion of this corollary is that under the summability condition $\sum_n |Df^n(c_1)|^{-1/\ell} < \infty$, φ is not a measurable coboundary except, possibly, for countably many parameter values. If f_a is infinitely renormalizable, then the summability condition fails. If f_a is finitely renormalizable, then equivalently, there is a smallest periodic interval $J \ni c$ with $f^k(J) \subset J$. Only when the renormalized map $f^k : J \to J$ is conjugate to the full unimodal map $x \mapsto 1 - 2x^2$ (and for each $k \ge 1$ and configuration of $J, f(J), \ldots, f^{k-1}(J)$, this usually holds for only

https://doi.org/10.1017/S0143385705000179 Published online by Cambridge University Press

one parameter), then it is possible that φ is a measurable coboundary. However, even in this situation, it seems extremely unlikely that φ is a measurable coboundary as it requires that the multipliers of all periodic orbits are the same.

Example. If $f(x) = 1 - ax^2$ is the quadratic map, then $\varphi = \log|f'| - \int \log|f'|\,dv$ is a measurable coboundary for $a = 2$, but not for the parameter $a \approx 1.543\,689\,01$ at which f is renormalizable of period 2, and $f^3(c)$ is the orientation reversing fixed point.

Proof. For $a = 2$, then f is a Chebychev polynomial, and hence $h(x) = -\cos 2x$ conjugates T with the tent map $T(x) = \min\{2x, 2(1-x)\}$: $h \circ T = f \circ h$. It follows that $\varphi = \log|f'| - \log 2 = \psi - \psi \circ f$ for $\psi = -\log|h' \circ h^{-1}|$.

Next assume that f has a k-periodic interval J as in the proof of Corollary 3, such that $F := f^k|J$ is conjugate to $x \mapsto 1 - 2x^2$. Applying the cohomological equation to F, we get $\log|F'| - k\int \log|f'|\,dv = \psi - \psi \circ F$. If $h : [0,1] \to J$ is defined by $\psi = -\log|h' \circ h^{-1}|$, then $F \circ h = h \circ T$ for the tent map T as above. In particular, $k\int \log|f'|\,d\mu = \log 2$. By the cohomological equation, each periodic point $y \in J^\circ$ must have multiplier $\log 2$.

For $k = 2$, $F|J$ has an orientation reversing fixed point q_1, and $\{q_1, q_2 := f(q_1)\}$ is the corresponding period 2 orbit under f. We find $\log|F'(q_1)| = \log|f'(q_1) \cdot f'(q_2)| = \log 4|1 - a|$. So $\log|F'(q_1)| = \log 2$ only if $a = \frac{1}{2}$ or $\frac{3}{2}$, but neither parameter value corresponds to the required renormalizable map. □

6. *Livšic theorems for non-uniformly hyperbolic systems*

In this section we use Young towers [**41, 42**] to prove measurable Livšic theorems for Lie group valued cocycles over non-uniformly hyperbolic systems. In particular we are able to prove Livšic theorems for Hénon maps [**5**]. These maps take the form $f(x, y) = (1 - ax^2 + y,\ bx)$, with $a \simeq 2$ and $b \simeq 0$, where for a positive Lebesgue measure subset of parameter space (a, b), it is proved that these maps admit a non-trivial attracting set with an ergodic Sinai–Ruelle–Bowen measure supported on it, see [**12**]. A Markov extension can be associated with such maps, as we describe below. Our results are also applicable to other non-uniformly hyperbolic systems which can be shown to admit a Young tower; see, for example, [**3, 15, 40**].

We discuss diffeomorphisms for which there exists a stable foliation. For the non-uniformly expanding case drop all references to the stable foliation. We refer to Young [**41, 42**] and Baladi [**4**] for more details. In a non-uniformly hyperbolic system the unstable leaves are not invariant under the return map to a reference set and this introduces some complications to the analysis. The proof in this section is based on that of Theorem 3 and Corollary 2.

In this section we use the notation that is commonly used for Young towers. Let $T : X \to X$ be a $C^{1+\varepsilon}$ diffeomorphism, where X is a compact manifold with metric ρ_X. Suppose there exists $\Lambda \subset X$ with a hyperbolic product structure [**41**, Definition 1] $\Lambda = \{(\bigcup \gamma^u) \cap (\bigcup \gamma^s) : \gamma^u \in \Gamma^u, \gamma^s \in \Gamma^s\}$, where Γ^u, Γ^s are two families of C^1 disks in X with the following properties: (i) disks in Γ^u are pairwise disjoint and the disks in Γ^s are pairwise disjoint; (ii) every $\gamma^u \in \Gamma^u$ meets every $\gamma^s \in \Gamma^s$ in exactly one point; (iii) there exists a lower bound on the angle between γ^u and γ^s at the point of intersection;

https://doi.org/10.1017/S0143385705000179 Published online by Cambridge University Press

and (iv) each $\gamma^u \in \Gamma^u$ satisfies $m_\gamma(\gamma^u \cap \Lambda) > 0$, where m_γ is the measure on γ^u induced by the Riemannian structure of X.

Under assumptions P1–P5 [**41**, §1], Young constructs a Markov extension (Young tower) (F, Δ) over $T : X \to X$ with base Λ. The set Λ is decomposed as $\Lambda = \bigcup_j \Lambda_j$ and there is a return function $R : \Lambda \to \mathbb{N}$, with constant value R_j on each Λ_j. Define $T^R(x) = T^{R(x)}(x)$.

We have $\Delta := \{(x, l) : x \in \Lambda;\ l = 0, 1, \ldots, R(x) - 1\}$ where $\Delta_0 = \{(x, 0) : x \in \Lambda\} = \Lambda$ in a natural identification. Define $F : \Delta \to \Delta$ as

$$F(x, i) = \begin{cases} (x, i + 1) & \text{if } x \in \Lambda_j \text{ and } i < R_j - 1, \\ (T^{R_j} x, 0) & \text{if } x \in \Lambda_j \text{ and } i = R_j - 1. \end{cases}$$

There is a projection $\pi : \Delta \to X$ given by $\pi(x, i) = T^i(x)$ and $\pi \circ F = T \circ \pi$. The tower (F, Δ) is then reduced to an expanding map $\bar{F}^R : \bar{\Delta}_0 \longrightarrow \bar{\Delta}_0$, where $\bar{\Delta}_0$ is the quotient of Δ_0 under the equivalence relation that two points are equivalent if and only if they belong to the same local stable leaf γ^s.

We need the following properties.

(A1) There exists an F-invariant probability measure ν with conditional measures $\{\nu^\gamma\}$ on $\gamma^u \cap \Lambda$ leaves with densities $\{\rho^\gamma\}$ with respect to m_γ, which satisfy $1/C \le \rho^\gamma \le C$ for some $C > 0$ (see [**42**, §2]). The measure ν induces a T-invariant measure ν_X on X by $\nu_X = \pi^* \nu$.

(A2) There is a countable partition $\mathcal{P}_0$ of $\Lambda = \Delta_0$ into elements Λ_j together with a return function $R : \Lambda \to \mathbb{N}$, with $R\,|_{\Lambda_i} = R_i$. Moreover, F^{R_i} maps Λ_i bijectively onto Λ (see [**42**, §1.1]). There is a countable partition $\mathcal{P}_\Delta$ with elements $(\Lambda_{i,j})$ with $0 \le j < R_i$ of Δ.

(A3) There exists $K > 0$ such that if $y \in \gamma^u(x)$,

$$\frac{1}{K} \le \frac{\|D^u(F^j x)\|}{\|D^u(F^j y)\|} \le K,$$

for all $j = 0, \ldots, s(x, y)$ [**42**, P4(b)]. Here D^u denotes the derivative along unstable leaves γ^u, and $s(x, y)$ is the first time n for which $F^n x$ and $F^n y$ lie in different elements of $\mathcal{P}_\Delta$.

(A4) For $\gamma, \gamma' \in \Gamma^u$, if $\Theta : \gamma \cap \Lambda \longrightarrow \gamma' \cap \Lambda$ is defined by $\Theta(x) = \gamma^s(x) \cap \gamma'$, then Θ is absolutely continuous, and there exists a $C_1 > 0$ such that

$$\frac{d(\Theta_*^{-1} \mu_{\gamma'})}{d\mu_\gamma}(x) \le C_1,$$

for all $x \in \gamma^u$ (see [**42**, P5(b)]).

(A5) There exists $\lambda_u > 1$ such that for each $x \in \Lambda$, $|D^u F^R(x)| \ge \lambda_u$ (see [**42**, §3.1]).

(A6) There exists $\lambda_s < 1$ such that for all $\gamma^s \in \Gamma^s$ and every x, y in the same γ^s, $\rho_X(F^j x, F^j y) \le C \lambda_s^j$ (cf. [**42**, P3]). Here $C > 0$ is a uniform constant.

(A7) Let $\mathcal{P}_n[x]$ denote the element of the partition $(T^R)^{-n}\mathcal{P}_0$ that contains $x \in \Lambda$. For $y \in \mathcal{P}_n[x]$ define $\tau(y) = R(y) + R(T^R y) + \cdots + R((T^R)^{n-1} y)$. Note that $\tau(y) = \tau(z)$ if $y, z \in \mathcal{P}_n[x]$. Let A_n be an element of the partition $(T^R)^{-n}\mathcal{P}_0$. Given $x_0 \in \Lambda$ and $x_{\tau_n} \in A_n$ with $T^{\tau_n} x_{\tau_n} = x_0$ define $x_i = T^{\tau_n - i} x_0 \to x_{\tau_n}$.

https://doi.org/10.1017/S0143385705000179 Published online by Cambridge University Press

Let $\hat{x} = (x_0, x_{\tau_1}, x_{\tau_2}, \dots, x_{\tau_n}, \dots)$ be a point in the natural extension of $T^R : \Lambda \to \Lambda$ with corresponding invariant measure $\hat{\nu}_0 \to \hat{\nu}$. We assume either:

(1) a one-dimensional unstable manifold; or

(2) for $\hat{\nu}$-a.e. $\hat{x}$ there exists $C(\hat{x})$ such that for all $z_{\tau_n}, y_{\tau_n} \in A_n$, $0 \le i \le \tau_n$,

$$\rho_X(y_i, z_i) \le C(\hat{x})\lambda_u^i \rho_X(y_0, z_0). \tag{26}$$

Cocycle assumptions. Let $\varphi\colon X \to G$ be Hölder of exponent $\alpha > 0$, and define constants μ_u (as in §2) and μ_s by

$$\mu_u := \lim_{n\to\infty} \left(\sup_{x\in X} \|\mathrm{Ad}\,(\varphi_n(x))\| \right)^{1/n},$$
$$\mu_s := \lim_{n\to\infty} \left(\sup_{x\in X} \|\mathrm{Ad}\,(\varphi_n(x))^{-1}\| \right)^{-1/n}$$

where $\varphi_n(x) = \varphi(T^{n-1}x)\dots\varphi(x)$. Assume a partial hyperbolicity condition (PH*) on the group extension

$$\lambda_s^\alpha < \mu_s \le 1 \le \mu_u < \tilde{\lambda}_u^\alpha \tag{PH*}$$

where $\tilde{\lambda}_u = \lambda_u^{1/\mathcal{R}}$.

We define

$$\Theta_{PH} := \max\left\{ \frac{\log \mu_u}{\log \tilde{\lambda}_u}, \frac{\log \mu_s}{\log \lambda_s} \right\}.$$

The assumption (PH*) implies that $\Theta_{PH} < \alpha$.

THEOREM 7. *Assume that (T, X, ν) is modelled by a tower over a base set $\Lambda \subset X$. Suppose, in addition, that T has a one-dimensional unstable direction or condition (26). Let $\varphi : X \to G$ be Hölder of exponent α and suppose condition (PH*) holds. If $\psi(Tx) = \varphi(x)\psi(x)$ ν_x-a.e. for some measurable function $\psi : X \to G$, then $\psi \mid \Lambda$ is γ-Hölder for some $\gamma \in (0, 1)$.*

Remark 17. The same regularity is forced upon solutions ψ to (2), (3) or (4).

Remark 18. Condition (PH*) is automatic if G is Abelian, compact or nilpotent.

Proof. We start with a lemma tackling the stable direction.

LEMMA 4. *There exists $\psi' = \psi$ ν_x-a.e. and ψ' is Hölder when restricted to each $\gamma^s \in \Lambda$ (with uniform constant and exponent).*

Proof. Choose a version of ψ and $\gamma^u \in \Lambda$ so that for ν_{γ^u}-a.e. $z \in \gamma^u$, $\psi(Tz) = \varphi(z)\psi(z)$. For each $z \in \gamma^u$, each $x \in \gamma^s(z) \subset \Lambda$ define

$$\psi'(x) = \lim_{n\to\infty} \varphi(x)^{-1} \dots \varphi(T^n x)^{-1}\varphi(T^n z)\dots\varphi(z)\psi(z).$$

By conditions (PH*) and (A6), an argument similar to that of §3 can be used to show that ψ' restricted to each $\gamma^s(z)$ is uniformly Hölder. Furthermore, $\psi'(Tx) = \varphi(x)\psi'(x)$ for ν-a.e. $x \in \Lambda$ and, hence, $\psi' = \alpha\psi$ ν-a.e. for some constant group element α. As $\nu_x = \pi^*\nu$, the result follows. □

https://doi.org/10.1017/S0143385705000179 Published online by Cambridge University Press

From now on we assume that ψ has the properties specified in the lemma above, namely ψ restricted to each $\gamma^s \in \Lambda$ is uniformly Hölder. Now we need only show that ψ restricted to each γ^u is Hölder since the local product structure implies in this case that ψ is Hölder on Λ. In fact, to show that ψ restricted to each γ^u is Hölder we only need to show that there is a $\gamma^u \in \Lambda$ such that ψ restricted to γ^u is Hölder, since the fact that the holonomy is Hölder and ψ restricted to each γ^s is Hölder implies the result for all $\gamma^u \in \Lambda$.

Recall that $\mathcal{P}_0$ is the partition of Λ into $\{\Lambda_j\}$, and each Λ_j contains whole stable leaves γ^s. We will, to simplify notation, denote $(x, 0) \in \Lambda_0$ as x. For $i \geq 1$ let $\mathcal{P}_i = \bigvee_{j=0}^{i-1} (F^R)^{-j} \mathcal{P}_0$. Refine the partition $\{\mathcal{P}_i\}$ in the stable direction by partitioning the stable manifolds into leaves of length at most 2^{-i} to form a partition $\mathcal{Q}_i$ of Λ. Partition Λ in such a way that if $A \subset B$, $A \in \mathcal{Q}_i$, $B \in \mathcal{P}_i$ then $\gamma^u \cap B = \gamma^u \cap A$ for each γ^u for each γ^u such that $\gamma^u \cap A \neq \emptyset$.

The σ-algebra generated by $\bigvee_i \mathcal{Q}_i$ generates the Borel σ-algebra on Λ. By the Martingale Density Theorem, given $\eta > 0$ there exists n and an element of the partition component $A_n \in \mathcal{Q}_n$ so that for some $x \in A_n$

$$\frac{\nu\{y \in A_n : d(\psi(x), \psi(y)) < \eta\}}{\nu(A_n)} > 1 - \eta.$$

For $x \in A_n$ define $\tau_n = R(x) + R(T^R x) + \cdots + R((T^R)^{n-1} x)$ and note that $T^{\tau_n}(x) = (T^R)^n x$. Given $x_0 \in \Lambda$, $x_{\tau_n} \in A_n$ with $T^{\tau_n} x_{\tau_n} = x_0$ define x_i by $T^{\tau_n - i} x_{\tau_n} = x_i$. By (A1) and (A4) we may choose a portion of leaf $\gamma_n = \gamma^u \cap A_n$ and x_{τ_n} such that

$$\frac{m_{\gamma_n}\{y \in \gamma_n \cap A_n : d(\psi(x_{\tau_n}), \psi(y)) < \eta\}}{m_{\gamma_n}(\gamma_n)} > 1 - \mathcal{O}(\eta).$$

Then $\tilde{\gamma}_n := (T^R)^n \gamma_n$ is an unstable leaf which crosses Λ completely in the unstable direction. As a consequence of (A3)

$$\frac{m_{\tilde{\gamma}_n}\{y_0 \in \tilde{\gamma}_n : d(\psi(x_{\tau_n}), \psi(y_{\tau_n})) < \eta\}}{m_{\tilde{\gamma}_n}(\tilde{\gamma}_n)} > 1 - \mathcal{O}(\eta). \tag{27}$$

On $\tilde{\gamma}_n$ define a function $\Psi_n : \tilde{\gamma}_n \longrightarrow G$ by

$$\Psi_n(y_0) = \varphi_{\tau_n}(y_{\tau_n})[\varphi_{\tau_n}(x_{\tau_n})]^{-1},$$

where $\varphi_i(x_j) = \varphi(x_{j-i+1}) \ldots \varphi(x_{i+j})$ and $\varphi_0(x) = e$.

Take points $z_0, w_0 \in \tilde{\gamma}_n$. Then by the cohomological equation

$$\begin{aligned}\psi(z_0) &= \varphi_{\tau_n}(z_{\tau_n})\psi(z_{\tau_n}) \\ &= \Psi_n(z_0)[\varphi_{\tau_n}(x_{\tau_n})]\psi(x_{\tau_n})\psi(x_{\tau_n})^{-1}\psi(z_{\tau_n}) \\ &= \Psi_n(z_0)\psi(x_0)\psi(x_{\tau_n})^{-1}\psi(z_{\tau_n}).\end{aligned}$$

By the right-invariance of the metric and the triangle inequality we have

$$\begin{aligned}d(\psi(z_0), \psi(w_0)) \leq\ & d(\Psi_n(z_0)\psi(x_0)\psi(x_{\tau_n})^{-1}\psi(z_{\tau_n}), \Psi_n(z_0)\psi(x_0)) \\ &+ d(\Psi_n(z_0)\psi(x_0)\psi(x_{\tau_n})^{-1}\psi(w_{\tau_n}), \Psi_n(z_0)\psi(x_0)) \\ &+ d(\Psi_n(z_0)\psi(x_0)\psi(x_{\tau_n})^{-1}\psi(w_{\tau_n}), \Psi_n(w_0)\psi(x_0)\psi(x_{\tau_n})^{-1}\psi(w_{\tau_n})).\end{aligned}$$

https://doi.org/10.1017/S0143385705000179 Published online by Cambridge University Press

We claim that Ψ_n is Hölder on $\tilde{\gamma}_n$ with uniform Hölder constant and exponent (the uniformity is over n in the construction).

We calculate

$$\begin{aligned} d(\varphi_{\tau_n}(z_{\tau_n})\varphi_{\tau_n}(x_{\tau_n})^{-1}, \varphi_{\tau_n}(w_{\tau_n})\varphi_{\tau_n}(x_{\tau_n})^{-1}) &= d(\varphi_{\tau_n}(z_{\tau_n}), \varphi_{\tau_n}(w_{\tau_n})) \\ &\le \sum_{i=0}^{\tau_n-1} \|\mathrm{Ad}\,(\varphi_i(z_i))\| d(\varphi(z_{i+1}), \varphi(w_{i+1})) \\ &\le \sum_{i=0}^{\tau_n-1} C(\mu_u)^{i+1}\lambda_u^{-(i+1)\alpha}\rho_X(z_0, w_0)^{\alpha}, \end{aligned}$$

where in passing from the third to fourth line, we use condition (26). Equivalently for the unstable direction we could have used the existence of the positive Lyapunov exponent $\tilde{\lambda}_u$, together with bounded distortion as in the proof of Theorem 1. The series converges uniformly because of condition (PH*). Equation (27) gives

$$\frac{m_{\tilde{\gamma}_n}\{y_0 \in \tilde{\gamma}_n : d(\psi(x_0), \psi(y_0)) < C\rho_X(x_0, y_0)^{\alpha}\}}{m_{\tilde{\gamma}_n}(\tilde{\gamma}_n)} > 1 - \mathcal{O}(\eta). \tag{28}$$

Recall that ψ restricted to each stable leaf in Λ is uniformly Hölder. The holonomy map along stable leaves is absolutely continuous (A4) and the density of ν with respect to Lebesgue is bounded away from zero and above by (A1). Hence [**19**, Proposition 19.1.1] implies that

$$\frac{\nu\{(x, y) \in \Lambda \times \Lambda : d(\psi(x), \psi(y)) < \tilde{C}\rho_X(x, y)^{\alpha}\}}{\nu \times \nu(\Lambda \times \Lambda)} > 1 - \mathcal{O}(\eta). \tag{29}$$

Since $\nu_x = \pi^*\nu$ and η is arbitrary, it follows that $\psi|_{\Lambda}$ has a Hölder version, thus proving Theorem 7. □

Acknowledgements. We would like to thank Ian Melbourne and Gerhard Keller for fruitful suggestions. M. Holland also acknowledges the support of an EPSRC grant No. GR/S11862/01. M. Nicol acknowledges the support of NSF grant DMS-0244529.

A. *Appendix*

Suppose that $\psi : X \to G$ is a measurable function from a metric measure space into a connected finite-dimensional matrix Lie group G endowed with a right invariant metric d_G. Let $\pi_{i,j} : G \to \mathbb{R}$, $1 \le i, j \le 2d$ be the local coordinate chart functions. The proof of the proposition below clearly generalizes to any finite number of real-valued measurable functions and hence establishes the Martingale Convergence Theorem, since continuity is a local property.

PROPOSITION 2. *Suppose that (X, μ) is a probability space and $\{\mathcal{P}_n\}$ is an increasing sequence of partitions of X and let $\mathcal{P}_n[x]$ denote the partition element of $\mathcal{P}_n$ which*

https://doi.org/10.1017/S0143385705000179 Published online by Cambridge University Press

contains $x \in X$. Suppose the Borel σ-algebra is generated by $\bigvee_n \mathcal{P}_n$. Let $\varphi : X \to \mathbb{R}$ be μ-measurable and $\eta > 0$. For μ-a.e. $x \in X$,

$$\lim_{n\to\infty} \frac{\mu\{y \in \mathcal{P}_n[x] : d(\varphi(x), \varphi(y)) < \eta\}}{\mu(\mathcal{P}_n)} > 1 - \eta. \tag{30}$$

Proof. First suppose that $\varphi \in L^1(\mu)$. Let $\mathcal{F}_n$ denote the σ-algebra generated by the partition $\mathcal{P}_n$. Then

$$\lim_{n\to\infty} \mathbb{E}[\varphi|\mathcal{F}_n](x) = \varphi(x), \quad \mu\text{-a.e.}$$

by [**6**, Corollary 5.22]. Note that $\mathbb{E}[\varphi|\mathcal{F}_n](x)$ is constant on $\mathcal{P}_n$. Choose a sequence $\{\delta_i\}$ such that $\sum_i \delta_i < \infty$. Given $\delta_i > 0$ take N_i sufficiently large that $d(\mathbb{E}[\varphi|\mathcal{F}_n](x), \varphi(x)) < \eta$ except for a set of measure at most $\delta_i \eta^2$ for all $n \geq N_i$. For all N_i, the union U_i of the set of atoms $A \in \mathcal{P}_{N_i}$ for which

$$\frac{\mu\{y \in A : d(\mathbb{E}[\varphi|\mathcal{F}_{N_i}](y), \varphi(y)) > \eta\}}{\mu(A)} > \eta,$$

satisfies $\mu(U_i) < \eta^2 \delta_i$. By the Borel–Cantelli lemma, μ-a.e. $x \in X$ lies in only finitely many U_i. Finally to remove the assumption that φ is integrable note that given $\varepsilon > 0$ there exists an integrable function ψ such that $\psi(x) = \varphi(x)$ except for a set of measure at most ε. An argument using approximating functions and the Borel–Cantelli lemma gives the same result for measurable φ. □

References

[1] J. Aaronson and M. Denker. Local limit theorems for partial sums of stationary sequences generated by Gibbs–Markov maps. *Stochast. Dynam.* **1** (2001), 193–237.

[2] J. Aaronson, M. Denker, O. Sarig and R. Zweimüller. Aperiodicity of cocycles and stochastic properties of non-Markov maps. *Stochast. Dynam.* **4** (2004), 31–62.

[3] J. Alves, S. Luzzatto and V. Pinheiro. Markov structures and decay of correlations for non-uniformly expanding dynamical systems. *Ann. Inst. Henri Poincaré* to appear.

[4] V. Baladi. *Positive Transfer Operators and Decay of Correlations (Adv. Series Nonlin. Dynam., 16)*. World Scientific, Singapore, 2000.

[5] M. Benedicks and L. Carleson. The dynamics of the Hénon map. *Ann. Math.* **133** (1991), 73–69.

[6] L. Breiman. *Probability (Classics Appl. Math., 7)*. SIAM, Philadelphia, PA, 1992.

[7] T. Bröcker and T. Dieck. *Representations of Compact Lie Groups*. Springer, New York, 1985.

[8] H. Bruin and S. Luzzatto. Topological invariance of the sign of the Lyapunov exponents in one-dimensional maps. *Proc. Amer. Math. Soc.* to appear.

[9] H. Bruin, S. Luzzatto, and S. van Strien. Decay of correlations in one-dimensional dynamics. *Ann. Sci. École Norm. Sup.* **36** (2003), 621–646.

[10] H. Bruin and S. van Strien. Existence of acips for multimodal maps. *Global Analysis of Dynamical Systems, Festschrift* dedicated to Floris Takens for his 60th birthday (2001).

[11] H. Bruin and S. Vaienti. Return time statistics for unimodal maps. *Fund. Math.* **176** (2003), 77–94.

[12] M. Benedicks and L. S. Young. Markov extensions and decay of correlations of certain Hénon maps. *Astérisque* **261** (2000), 13–56.

[13] R. de la Llave, J. M. Marco and E. Moriyon. Canonical perturbation theory of Anosov systems and regularity results for the Livsic cohomological equation. *Ann. Math.* **123** (1986), 537–611.

[14] S. Gouëzel. Sharp polynomial estimates for the decay of correlations. *Israel J. Math.* **139** (2004), 29–65.

[15] S. Gouëzel. Decay of correlations for non-uniformly expanding dynamical systems. *Bull. Soc. Math. France* to appear.

https://doi.org/10.1017/S0143385705000179 Published online by Cambridge University Press

[16] S. Gouëzel. Regularity of coboundaries for non uniformly expanding Markov maps. *Proc. Amer. Math. Soc.* to appear.
[17] F. Hofbauer. On intrinsic ergodicity of piecewise monotonic transformations with positive entropy. *Israel J. Math.* **34** (1979), 213–237.
[18] O. Jenkinson. Smooth cocycle rigidity for expanding maps and an application to Mostow rigidity. *Math. Proc. Camb. Phil. Soc.* **132** (2002), 439–452.
[19] A. Katok and B. Hasselblatt. *Introduction to the modern theory of dynamical systems (Encyclopedia Math. Appl., 54)*. Cambridge University Press, Cambridge, 1995.
[20] G. Keller. *Equilibrium States in Ergodic Theory*. Cambridge University Press, Cambridge, 1988.
[21] G. Keller. Exponents, attractors and Hopf decompositions for interval maps. *Ergod. Th. & Dynam. Sys.* **10** (1990), 717–744.
[22] G. Keller. Lifting measures to Markov extensions. *Monatsh. Math.* **108** (1989), 183–200.
[23] H. Keynes and D. Newton. Ergodic measures for non-abelian compact group extensions. *Compositio Math.* **32** (1976), 53–70.
[24] A. N. Livšic. Cohomology of dynamical systems. *Math. USSR Izv.* **6** (1972), 1278–1301.
[25] A. N. Livšic. Homology properties of Y-systems. *Math. Notes* **10** (1971), 758–763.
[26] R. Mañé, *Ergodic Theory and Differentiable Dynamics*. Springer, Berlin, 1987.
[27] W. de Melo and S. van Strien. *One-dimensional Dynamics*. Springer, Berlin, 1993.
[28] M. Nicol and M. Pollicott. Measurable cocycle rigidity for some noncompact groups. *Bull. Lond. Math. Soc.* **31** (1999), 592–600.
[29] M. Nicol and M. Pollicott. Livšic theorems for semisimple Lie groups. *Ergod. Th. & Dynam. Sys.* **21** (2001), 1501–1509.
[30] M. Nicol and A. Scott. Livšic theorem and stable ergodicity for group extensions of hyperbolic systems with discontinuities. *Ergod. Th. & Dynam. Sys.* **23** (2003), 1867–1889.
[31] M. Noorani. Ergodicity and weak-mixing of homogeneous extensions of measure-preserving transformations with applications to Markov shifts. *Monatsh. Math.* **123** (1997), 149–170.
[32] T. Nowicki and S. van Strien. Invariant measures exist under a summability condition. *Invent. Math.* **105** (1991), 123–136.
[33] W. Parry and M. Pollicott. The Livsic cocycle equation for compact Lie group extensions of hyperbolic systems. *J. London Math. Soc. (2)* **56** (1997), 405–416.
[34] M. Pollicott and C. Walkden. Livšic theorems for connected Lie groups. *Trans. Amer. Math. Soc.* **353** (2001), 2879–2895.
[35] M. Pollicott and M. Yuri. Regularity of solutions to the measurable Livsic equation. *Trans. Amer. Math. Soc.* **351** (1999), 559–568.
[36] A. Scott. Livšic theorems for unimodal maps. *Preprint*, 2004.
[37] A. Scott. Livšic theorems and the stable ergodicity of compact group extensions of systems with some hyperbolicity. *PhD Thesis*, University of Surrey, 2003.
[38] C. Walkden. Livsic theorems for hyperbolic flows. *Trans. Amer. Math. Soc.* **352** (2000), 1299–1313.
[39] C. Walkden. Livsic regularity theorems for twisted cocycle equations over hyperbolic systems. *J. London Math. Soc.* **61** (2000), 286–300.
[40] Q. Wang and L.-S. Young. Strange attractors with one direction of instability. *Comm. Math. Phys.* **218** (2001), 1–97.
[41] L.-S. Young. Recurrence times and rates of mixing. *Israel J. Math.* **110** (1999), 153–188.
[42] L.-S. Young. Statistical properties of dynamical systems with some hyperbolicity. *Ann. Math.* **147** (1998), 585–650.

https://doi.org/10.1017/S0143385705000179 Published online by Cambridge University Press