© NIAB 2011
ISSN 1479-2621
Plant Genetic Resources: Characterization and Utilization (2011) **9**(3); 464–477
doi:10.1017/S1479262111000670

Diversity, characterization and utilization of ginger: a review

Jaleel Kizhakkayil[1] and B. Sasikumar[2]*

[1]*Department of Presidential Affairs, Abu Dhabi, Al-Ain, PO 47087, United Arab Emirates and* [2]*Indian Institute of Spices Research, PO Marikunnu, Calicut 673 012, Kerala, India*

Received 24 January 2011; Accepted 14 April 2011 – First published online 1 June 2011

Abstract

Ginger (*Zingiber officinale* Rosc.), originated in the Indo-Malayan region, is now widely distributed across the tropics of Asia, Africa, America and Australia. It was domesticated in India and China, which represent the centre of origin of the species. Cultivated ginger though sterile, exhibits variations in rhizome and vegetative characters. The crop is gaining importance as a curative agent for a variety of ailments. Yield and quality traits (such as essential oil, fibre and oleoresin contents) along with volatile and non-volatile constituents are important determinants of the commodity's end product. Cultivar diversity for yield and morphological features is well known in ginger with few primitive types having excellent quality. However, the common name(s) of ginger cultivars are confusing and have resulted in geographical bias in its *ex situ* conservation. Most of the molecular/biochemical marker studies reported in ginger show low levels of polymorphism in contrast to plentiful phenotypic variability recorded in the species. The large influence of environmental factors on the content of key compounds, lack of seed set and confusion of the common names are the leading constraints to further varietal improvement of ginger. As tailor-made ginger varieties assume future significance these aspects will have relevance.

Keywords: antiquity; biochemical; medicinal use; molecular; morphological; taxonomy

Introduction

Ginger (*Zingiber officinale* Rosc.), originated in the Indo-Malayan region, is now widely distributed across many countries as a spice and medicinal plant (Purseglove *et al.*, 1981; Burkill, 1996; Park and Pizzuto, 2002). Traders took ginger from India to Mediterranean region during the 1st century CE (Current Era). The Arabs introduced ginger to East Africa in the 13th century CE and the Portugese spread it to West Africa and the Pacific islands for commercial cultivation (Ravindran *et al.,* 2006).

The major ginger growing countries include Australia, Brazil, Bangladesh, Cameroon, China, Costa Rica, Fiji, Ghana, Guatemala, Hawaii, India, Indonesia, Jamaica, Mauritius, Malaysia, Nepal, New Zealand, Nigeria, Philippines, Sierra Leone, Sri Lanka, Taiwan, Thailand, Trinidad and Uganda covering a total area of 387,300 ha with a production of 1,476,900 MT. India is the world's largest producer of ginger at present (Table 1). Export of ginger from India was 7500 MT during 2006–2007. 'Cochin ginger', 'Wayanadan ginger' (India), 'Chinese ginger' (China), 'Buderim Gold' (Australia) 'Jamaican' (Jamaica) are globally traded products. Ginger in various forms is used as food flavorant, antioxidant and antimicrobial besides as deodorizing agent in food. Chinese, Indian, Tibetan and Arabic systems of medicine recognized the medicinal value of ginger since ancient times (Atman and Marcussen, 2001).

This review is intended to provide an overview of the diversity of cultivated ginger, its characterization and utilization.

* Corresponding author. E-mail: bhaskaransasikumar@yahoo.com

https://doi.org/10.1017/S1479262111000670 Published online by Cambridge University Press

Table 1. Area, production and productivity of ginger in the world (2006)

Country	Area (ha)	Production (tons)	Productivity (tons/ha)
India	105.50	517.8	4.9
China	24.50	279.0	11.3
Indonesia	18.20	159.0	8.7
Nepal	12.90	154.1	12.0
Nigeria	191.00	134.0	2.4
Bangladesh	7.70	49.4	6.4
Thailand	14.00	34.0	2.4
Philippines	3.90	27.2	6.9
Cameroon	1.30	7.5	5.7
USA	0.04	1.9	47.5
Rest of the World	8.20	71.5	8.7
World	387.30	1476.9	3.8

Source: http://faostat.fao.org/site/567/DesktopDefault.aspx?PageID=567.

Antiquity

Ayurveda, the Indian 'science of life' of approximately 5000 years of antiquity, documented the medicinal value of ginger in treatises such as *Charaka Samhita*, *Shushrutha Samhita* and *Ashtangahridaya*. In *Ayurveda*, ginger is known as *Mahaoushadhi* meaning great medicine. It is *Sringavera* in Sanskrit, the ancient Indian language, which has given way to the Greek Zingiberi and to the Latin *Zingiber* (Rosengarten, 1969; Purseglove *et al*., 1981). Ginger is also mentioned in Koran (76: 15–17) though it does not figure in Bible. Ginger was first documented by van Rheede (1692) as *inschi* (*inchi*) in 'Hortus Indicus Malabaricus' – the first printed account of the plants of Western Coast (Malabar Coast) of India.

Taxonomy

The genus *Zingiber* belongs to the family *Zingiberaceae* under the order *Zingiberales* and the tribe *Zingibereae* (Holtum, 1950). *Zingiberaceae* includes three other tribes: *Hedychieae*, *Alpinieae* and *Globbeae* (Petersen, 1889; Burtt and Smith, 1972). The tribe *Zingibereae* has seven other genera: *Boesenbergia*, *Camptandra*, *Roscoea*, *Kaempferia*, *Amomum*, *Hedychium* and *Curcuma* (Kress *et al*., 2002). Jatoi *et al*. (2006) using rice microsatellite markers assessed the genetic variability among three genera of the family *Zingiberaceae*: *Zingiber*, *Alpinia* and *Curcuma* and found the origin of the genera diverse covering eight Asian countries. *Zingiber* contains 150 species and four sections distributed throughout tropical Asia, China, Japan and tropical Australia besides the subspecies (varieties): *Z. officinale* var. *rubra* and *Z. officinale* var. *rubrum* (Muda *et al*., 2004). *Z. officinale* is included in the section II, *Lampuzium* (Baker, 1882; Sabu, 2003; Larsen and Larsen, 2006).

Diversity

Species level

The genus *Zingiber* includes many economically important species besides *Z. officinale* as given in Table 2. The Indian species reported include *Z. chrysanthum* Rosc., *Z. rubens* Roxb., *Z. roseum* Rosc., *Z. nimmonii* Dalz., *Z. wightianum* Thw., *Z. barbatum* Wall., *Z. squarrosum* Roxb., *Z. ligulatum* Roxb., *Z. cernuum* Dalz., *Z. pandur-atum* Roxb., *Z. pardocheilum* Wall., *Z. intermedium* Baker, *Z. officinale* Rosc., *Z. griffithii* Baker, *Z. gracile* Jack., *Z. zerumbet* Smith, *Z. cylindricum* Moon,

Table 2. Some of the economically important *Zingiber* species

Species/subspecies	Occurrence	Use
Z. officinale Roscoe	Tropical countries, China, USA	Spice, condiment, medicinal
Z. officinale var. *rubrum*	Malaysia	Medicinal, spice
Z. officinale var. *rubra*	Malaysia	Medicinal, spice
Z. mioga Roscoe	Japan	Vegetable. Shoot and flower are edible
Z. zerumbet (L.) Smith	Tropical Asia	Medicinal, ornamental
*Z. montanum (*Koenig*)* Link ex Dietr	India, Malaysia, SriLanka, Java	Used in traditional medicine
Z. *clarkii* King	Sikkim Himalayas	Ornamental
Z. *aromaticum* Val	Tropical Asia	Ornamental, medicinal, flavouring
Z. rubens Roxb	Indo-Malaya	Medicinal, ornamental
Z. griffithii Baker	Malaysia	Ornamental
Z. ottensii Valet	South East Asia	Medicinal, ornamental
Z. corallinum Hance	South East Asia	Chinese medicine, ornamental
Z. americanus Bl.	South East Asia	Medicinal, vegetable
Z. argenteum (J. Mood and I. Theilade)	Sarawak, Malaysia	Ornamental

https://doi.org/10.1017/S1479262111000670 Published online by Cambridge University Press

Table 3. Cultivar diversity of ginger in India (revised from Sasikumar *et al.*, 1999)

S. No.	Cultivar/variety
1	Ambalavayalan (Kerala)
2	Anamika
3	Arippa
4	Assam (Assam)
5	Athira[a] (Kerala)
6	Bahrica (Orissa)
7	Bajpai
8	Bazar local (Arunachal Pradesh)
9	Bhaise (Sikkim)
10	Bola (Assam)
11	Burdwan (West Bengal)
12	China[b] (Kerala)
13	Edappalayam (Kerala)
14	Ellakallan (Kerala)
15	Ernad-Chernad (Kerala)
16	Ernad-Manjeri (Kerala)
17	Gorubathani (Sikkim)
18	Himachal (Kerala, Himachal Pradesh)
19	Himgiri[a] (Himachal Pradesh)
20	Jamaica[b] (Kerala)
21	Jatia (Assam)
22	Jorathangey (Sikkim)
23	Jorhat (Assam)
24	Jugijan (North East India)
25	Kakkakalan (Kerala)
26	Karakkal (Kerala)
27	Karthika[a] (Kerala)
28	Keki (Arunachal Pradesh)
29	Khasi local (Meghalaya)
30	Kozhikkalan (Kerala)
31	Kunduli local (Orissa)
32	Kunnamangalam (Kerala)
33	Kuruppampady (Kerala)
34	Mahim (Maharashtra)
35	Mahima[a] (all over India)
36	Majhauley (Sikkim)
37	Mananthody (Kerala)
38	Maran/Moran (Assam, Kerala)
39	Mowshom (NorthEast India)
40	Nadia (West Bengal, Kerala, Meghalaya)
41	Naga Shing (Manipur)
42	Nangrey (Sikkim)
43	Narasapattam (Andhra Pradesh)
44	Poona
45	Rajagarh local
46	Rejatha[a] (Kerala)
47	Rio-de-Janeiro[b] (Kerala, Karnataka)
48	Sargi guda (Orissa)
49	Saw thing laidum (North East India)
50	Sawthing Pui (North East India)
51	Saying Makhim/pink ginger (Meghalaya)
52	Shing Bhoi (Meghalaya)
53	Shing Bhukir (Meghalaya)
54	Sierra Leone[b]
55	Singhihara
56	Suprabha[a] (all over India)
57	Suravi[a] (Orissa)
58	Suruchi[a] (Orissa)
59	Taffingiva
60	Taiwan[b]
61	Thang-Chang (Mizoram)
62	Thing laidum (Mizoram)
63	Thingaria (Mizoram)
64	Thingpui (Manipur, Mizoram)
65	Thodupuzha (Kerala)
66	Tura (Meghalaya)
67	Uttarpradesh (Uttarpradesh)
68	Valluvanad (Kerala)
69	Varada[a] (all over India)
70	Vellinchi (Kerala)
71	Vengara (Kerala)
72	Wynad Kunnmangalam (Kerala)
73	Wynad local (Kerala)
74	Zahirabad

Words in parenthesis denote important states of cultivation.
[a] Improved variety, [b] Exotic.

Z. macrostachyum Dalz., *Z. spectabile* Griff., *Z. casumunar* Roxb., *Z. parishii* Hook., *Z. clarkii* King, *Z. capitatum* Roxb., *Z. marginatum* Roxb. (Baker, 1882).

Most of the *Zingiber* species are diploid ($2n = 22$) except *Z. mioga* ($2n = 55$) and set seed. In cultivated ginger (*Z. officinale)* $2n = 24$ and $22 + 2B$ are also reported besides the normal diploid number of $2n = 22$ (Ravindran *et al.*, 2005). The species is sterile.

Cultivar level

Ginger is a rhizome propagated crop. Cultivar diversity abounds in India and China, which represents the centre of origin of this species, unlike Simmonds (1979) observed in many other crops. Geographical spread of ginger clones coupled with mutation and selection are considered to be responsible for the cultivar diversity (Ravindran *et al.*, 1994). About 74 cultivars, most of them with vernacular names, possessing varying levels of quality attributes and yield occur in India (Table 3; Sasikumar *et al.*, 1999). Major Chinese cultivars include 'Gandzhou', 'Shandong', 'Zaoyang', 'Zungi big ginger', 'Chenggu yellow', 'Kintoki', 'Sanshu' and 'Oshoga' are three major ginger cultivars from Japan (Ravindran *et al.*, 2005). Ridley (1912) has reported three types of ginger from Malaysia viz. 'halyia betle', 'halyia udang', 'halyia bara', 'Native' and 'Hawaiin' are cultivars from the Philippines. Nepal has about 50 cultivars (Sasikumar *et al.*, 1999). Ginger varieties of Guyana, mainly from the hinterlands of that country, are bold or very bold and lack any varietal names. An obsolete cultivar characterized by very dwarf stature and extremely slender, fibrous rhizome is also grown in Guyana (Sasikumar,

https://doi.org/10.1017/S1479262111000670 Published online by Cambridge University Press

2008). 'Sidda' and 'China' are the cultivars of Sri Lanka (Macloed and Pieris, 1984) and Jamaica has only one popular cultivar (Lawrence, 1984).

Varietal improvement in ginger has been limited to germplasm selection for high yield and better quality attributes besides some mutation breeding and ploidy breeding. So far nine improved ginger varieties ('Varada,' 'Mahima', 'Rejatha', 'Suruchi', 'Suravi', 'Suprabha', 'Himgiri', 'Athira' and 'Karthika') were released in India (Sasikumar *et al*., 2003, 2007) and one ('Buderim Gold') from Australia (Smith *et al*., 2004).

Characterization of ginger

Morphological and anatomical characterization

Variability for yield, quality traits and rhizome features has been reported in ginger (Khan, 1959; Thomas, 1966; Krishnamurthy *et al*., 1972; Muralidharan and Kamalam, 1973; Mohanty and Sharma, 1979; Nybe and Nair, 1979; Nybe *et al*., 1980; Sreekumar *et al*., 1980; Kumar *et al*., 1980; Mohanty *et al*., 1981; Rattan *et al*., 1988; Arya and Rana, 1990; Sasikumar *et al*., 1992; Zachariah *et al*., 1993; Ravindran *et al*., 1994; Sasikumar *et al*., 1994, 1999, 2003; Aiyadurai, 1996; John and Ferreira, 1997; Chandra and Govind, 1999; Yadav, 1999; Zachariah *et al*., 1999; Singh *et al*., 1999; Singh *et al*., 2000; Gowda and Melanta, 2000; Mohandas *et al*., 2000; Naidu *et al*., 2000; Singh, 2001; Tiwari, 2003; Abraham and Latha, 2003; Rana and Korla, 2007; Lincy *et al*., 2008) (Fig. 1). Primitive types (obsolete cultivars) in general are characterized by small rhizomes; poor yield and better quality although the cultivars/improved varieties possess attractive bold rhizomes with good yield and mixed quality traits. Ginger cultivars/varieties suited for various end uses are given as Table S1, available online only at http://journals.cambridge.org. However, yield and quality levels in ginger also vary with genotypes, soil types, locations, seasons, cultural practices and climatic conditions.

Ginger (*Z. officinale*) has distinct anatomical features such as absence of periderm, short lived functional cambium, occurrence of xylem vessels with scalariform thickening in the rhizome compared with *Z. zerumbet*, *Z. roseum* and *Z. macrostachyum* (Ravindran *et al*., 1998).



Fig. 1. Variability for rhizome features in Indian ginger germplasm (a) 'Varada', (b) 'Mahima', (c) 'Rejatha', (d) 'Suprabha', (e) 'Sabarimala', (f) 'Kozhikkalan', (g) 'Kakakalan', (h) 'Ellakallan', (i) 'Nadia', (j) 'Rio-de-Janeiro', (k) 'Silent valley' and (l) 'Himachal' (A colour version of this figure can be found online at journals.cambridge.org/pgr).

https://doi.org/10.1017/S1479262111000670 Published online by Cambridge University Press

Molecular characterization

RAPD (random amplified polymorphic DNA), AFLP (amplified fragment length polymorphism), ISSR (inter simple sequence repeats), simple sequence repeats markers and isozymes have been used to characterize ginger germplasm. Sasikumar *et al.* (2000) evaluated 14 accessions of ginger from different regions of India for variations in acid phosphatase, polyphenol oxidase, super oxide dismutase (SOD) and peroxidase (PRX). Among them, acid phosphatase showed maximum number of bands followed by SOD, while PRX had the least bands. Though variability in the population was very low, the accessions from Kerala region (southern India) were distinct from those collected from the north-east Indian states suggesting a geographical bias of the germplasm studied. A total of 28 ginger cultivars from China were compared for PRX isoenzyme patterns by fuzzy cluster analysis (Chengkun *et al.*, 1995). The cultivars differed in zonal number, activity intensity and isoenzyme pattern, which were related to rhizome size, growth intensity and blast (blight?) (*Ralstonia solanacearum*) resistance. The cultivars from the Fujian Province were more diverse.

Muda *et al.* (2004) studied the genetic variation among three *Z. officinale* cultivars/subspecies i.e. *Z. officinale* var. *officinale*, *Z. officinale* var. *rubra* ('halia bara') and *Z. officinale* var. *rubrum* ('halia padi') using RAPD analysis and cultivar-specific markers were obtained. Similarly, Nayak *et al.* (2005) based on their study of 16 ginger cultivars from India showed that RAPD primers OPC02, OPD20 and OPN06 had strong resolving powers and were able to distinguish all cultivars. However, Palai and Rout (2007) found that of the eight Indian landraces/varieties of ginger ('Surabhi', 'Tura Local', 'Jugijan', 'Nadia', 'S-558', 'Gorubathani', 'ZO-17' and 'Nangrey') all but one formed a single cluster in the dendrogram. Molecular characterization of seven ginger accessions including primitive, exotic and elite ones using 14 ISSR and 16 RAPD markers by Prem *et al.* (2007) also showed that the four primitive gingers ('Kozhikkalan', 'Kakkakalan', 'Ellakkallan' and 'Sabarimala') formed one cluster while all others were separately clustered in the dendogram. In another molecular characterization study using 25 RAPD primers Zhen-wei *et al.* (2006) reported a narrow genetic base among 20 Chinese cultivars.

AFLP markers were used to study the genetic relationship among three Indonesian type gingers (big ginger, small ginger and red ginger) (Wahyuni *et al.*, 2003). In this analysis, 28 accessions including those from Africa and Japan were used. Red ginger was genetically distinct from the big ginger, but close to some accessions of small ginger. There was no clear genetic differentiation between the small and big types ginger. Lee *et al.* (2007) isolated and characterized eight polymorphic microsatellite markers. A total of 34 alleles were detected across 20 ginger accessions, with an average of 4.3 alleles per locus. This study revealed the existence of moderate level of genetic diversity among the ginger accessions genotyped.

Recently, Kizhakkayil and Sasikumar (2010) characterized 46 accessions using 30 RAPD and 17 ISSR markers. Unweighted pair group method with arithmetic mean dendrograms based on three similarity coefficients (Jaccard's, Sorensen–Dice and Simple Matching) placed the accessions in four similar clusters and revealed less genetic distance among the accessions. Improved varieties/cultivars were grouped together with primitive types. In the clustering pattern of the accessions, a geographical bias was also evident implying that germplasm collected from nearby locations with local identity may not be genetically distinct. The clustering of the accessions was largely independent of its agronomic features.

The different molecular markers in general revealed low level of polymorphism (Figs. 2 and 3) and only moderate variability with some exceptions, among the ginger accessions studied probably due to the confounded effect of the duplicate accessions in the genebank.

Biochemical characterization

Characterization of ginger for the major biochemical constituents viz. oleoresin, essential oil and crude fibre levels has lead to identification of cultivars rich in one or other of these constituents (Table S1, available online only at http://journals.cambridge.org). Oleoresin content of ginger varied from 3 to 11% depending on the genotype, solvent extraction condition, state of rhizome, place of origin and harvest season (Ratnambal *et al.*, 1987; Vernin and Parkanyl, 2005).

Crude fibre content of dried ginger ranged from 4.8 to 9% (Natarajan *et al.*, 1972) although essential oil content of ginger varied from 0.2 to 3%, depending on the origin and state of rhizome (van Beek *et al.*, 1987; Ekundayo *et al.*, 1988). Characterization of 46 ginger accessions revealed that the primitive type gingers such as 'Sabarimala', 'Kozhikkalan' and 'Kakakalan' as well as few landraces have higher levels of oleoresin and essential oil compared with the improved varieties. (Kizhakkayil and Sasikumar, 2009). Menon (2007) too has observed that primitive ginger types such as 'Kozhikkalan' and 'Vellinchi' are rich in essential oil content. *Shing Bhukir*, a primitive type ginger from Meghalaya, India, is sold at a premium price for its medicinal value (Rahman *et al.*, 2009). 'Kintoki', a landrace from Japan, has very low fibre content (Kizhakkayil, 2008). Indian ginger cultivars are known to vary for the pungent and non-pungent constituents: gingerol and shogaol (Zachariah *et al.*,

https://doi.org/10.1017/S1479262111000670 Published online by Cambridge University Press




Fig. 2. RAPD profile of the DNA isolated from different ginger accessions amplified with primer OPA 07. M-marker; lane 1–46 'Varada', 'Mahima', 'Rejatha', 'Suruchi', 'Suprabha', 'Himachal', 'Maran', 'Nadia', 'Karakkal', 'Mananthodi', 'Sabarimala', 'Kozhikkalan', 'Ellakkallan', 'Kakakalan', 'Pakistan', 'Oman', 'Brazil', 'Jamaica', 'Rio-de-Janeiro', 'Pink ginger', 'Bakthapur', 'Kintoki', 'Nepal', 'China', 'Jugijan', 'Acc. no. 50', 'Pulpally', 'Acc. no. 95', 'Ambalawayalan', 'Kozhikode', 'Thodupuzha-1', 'Konni local', 'Angamali', 'Thodupuzha-2', 'Kottayam', 'Palai', 'Silent valley', 'Wayanad local', 'Vizagapatnam-1', 'Vizagapatnam-2', 'Fiji', 'Gorubathani', 'Bhaise', 'Naval parasi', 'Neyyar', 'Jolpaiguri', respectively.

1993) though lack of variability for gingerol and shogaol has been observed in Australian ginger (Wohlmuth *et al.*, 2005). High levels of *ar*-curcumene and β-bisabolene with reasonable levels of citral isomers and very low levels of zingiberene were observed in dried Sri Lankan ginger (Macloed and Pieris, 1984).

In short, the volatile and/or non-volatile compounds of ginger are affected by the environment and the state of rhizome (Connel and Jordan, 1971; Macloed and Pieris, 1984; Ekundayo *et al.*, 1988; Menut *et al.*, 1994; Vernin and Parkanyl, 1994; Fakim *et al.*, 2002; Wohlmuth *et al.* 2006). However, a wide variability in the essential oil profile of ginger is unseen though there are few odd cultivars/primitive types excelling in one or the other compounds including the pungent constituents.

Characterization against pests

Bacterial wilt (*R. solanacearum*), soft rot (*Pythium* spp.) and shoot borer (*Conogethes punctiferalis*) are three major pests of ginger. No stable bacterial wilt resistant ginger cultivar is known to exist. Consistent with this, 600 accessions screened for bacterial wilt tolerance using soil inoculation method were found susceptible (Kumar and Hayward, 2005). However, Shylaja *et al.* (2010) reported that the ginger clone 'Athira' is relatively more tolerant to bacterial wilt and soft rot than its mother clone 'Maran'. Clones 'Maran' 'Suprabha' and 'Himachal' have been reported to show field tolerance to ginger rot (*Pythium aphanidermatum*) (Indrasenan and Paily, 1974; Setty *et al.*, 1995).

Nybe and Nair (1979a) while recording shoot borer incidence in 25 ginger cultivars reported cultivar 'Rio de Janeiro' with minimum incidence. Devasahayam *et al.* (2010) could not find any shoot borer resistant lines in 592 ginger accessions screened though 49 accessions were rated as moderately resistant.

Uses

Products and end uses

Ginger imparts flavour and pungency to food and beverages (Arctangder, 1960; Pruthy, 1993; Bakhru, 1999).

https://doi.org/10.1017/S1479262111000670 Published online by Cambridge University Press




Fig. 3. ISSR profile of the DNA isolated from different ginger accessions amplified with primer ISSR3. M-marker; lane 1–46 'Varada', 'Mahima', 'Rejatha', 'Suruchi', 'Suprabha', 'Himachal', 'Maran', 'Nadia', 'Karakkal', 'Mananthodi', 'Sabarimala', 'Kozhikkalan', 'Ellakkallan', 'Kakakalan', 'Pakistan', 'Oman', 'Brazil', 'Jamaica', 'Rio-de-Janeiro', 'Pink ginger', 'Bakthapur', 'Kintoki', 'Nepal', 'China', Jugijan', 'Acc. No. 50', 'Pulpally', 'Acc. No. 95', 'Ambalawayalan', 'Kozhikode', 'Thodupuzha-1', 'Konni local', 'Angamali', 'Thodupuzha-2', 'Kottayam', 'Palai', 'Silent valley', Wayanad local', 'Vizagapatnam-1', 'Vizagapatnam-2', 'Fiji', 'Gorubathani', 'Bhaise', 'Naval parasi', 'Neyyar', 'Jolpaiguri', respectively.

Flavour properties of ginger depend on both volatile and non-volatile fractions. Fresh succulent baby pink ginger, salted ginger, crushed fresh ginger, dry ginger, ginger powder, ginger oil, ginger oleoresin, dry soluble ginger, ginger paste and ginger emulsion are used in different preparations. Fresh young ginger, succulent in nature with low fibre content is preferred for products such as ginger candy, preserves, and salted ginger while mature fresh ginger is used for preparing ginger shreds, the ethnic ginger chutney, ginger tea and ginger curry (*Inchi puli*). Ginger is an indispensable component of curry powder, sauces, ginger bread and ginger flavoured carbonated drinks (Hirara and Takesma, 1998) in addition to its use in biscuits, pickles, confectionaries and other dietary preparations.

Ginger in medicine

Ayurveda recommends ginger as a carminative, diaphoretic, antispasmodic, expectorant, peripheral circulatory stimulant, astringent, appetite stimulant, anti-inflammatory agent, diuretic and digestive aid (Warrier, 1989). Recent pharmacological studies attest this age-old medicinal value of ginger (Aimbire *et al*., 2007; Ansari and Bhandari, 2008; Morakinyo *et al*., 2008; Heimes *et al*., 2009; Nammi *et al*., 2009, 2010; Lakshmi and Sudhakar, 2010). Ginger is recommended to expecting women as an antenatal medicine as well as to alleviate the 'morning sickness' during pregnancy (Ozgoli *et al*., 2009; Broussard *et al*., 2010) and motion sickness. *Trikatu* is a favourite carminative remedy prepared with equal portions of *Z. officinale*, *Piper nigrum* and *Piper longum* (Anonymous, 1982). Ginger is the major constituent in formulations including *Chatarbhada Kvatha* used against fever and respiratory disorders; in antitussive drug *Lavangadhichurn* and in *Hingvastaka churna* recommended for indigestion (Singh, 1983; Thakur *et al*., 1989).

Ginger is also a major ingredient in *Unani* preparations such as 'Hub-gul-pista', an expectorant, 'Sufuf-shirin' an anti-dysenteric drug and 'Majun Izuroqui' a geriatric tonic as well as in the ginger based carminatives 'Qurs Pudina' and 'Murraba Adrak' (Singh, 1983; Thakur *et al*., 1989).

Chinese medicine too value ginger since 4th century BC (Wagner and Hikino, 1965). The Chinese administer ginger for a wide variety of medical problems such as stomachache, headache, diarrhoea, nausea, cholera, asthma,

https://doi.org/10.1017/S1479262111000670 Published online by Cambridge University Press

heart conditions, respiratory disorders, toothache and rheumatic complaints (Wagner and Hikino, 1965; Grant and Lutz, 2000). Africans and West Indians also use ginger for many ailments (Tyler *et al.*, 1981). Ginger has been used in the Mediterranea for treatment of arthritis, rheumatological problems and muscular discomfort (Langner *et al.,* 1998; Sharma and Clark, 1998). It has also been recommended for the treatment of various other conditions including atherosclerosis, migraine, rheumatoid arthritis, high cholesterol, ulcers, depression and impotence (Liang, 1992).

In folk/ethnic medicine too ginger is indispensable. Fresh ginger juice is administered to pregnant women just before childbirth for easy delivery (Rao and Jamir, 1982). Ginger is an ethnic remedy for pain, rheumatism, mad convulsion, collapse, scabies, constipation, indigestion, prolepsis, fistula, cholera, throat pain, tuberculosis, cold, fever and cough (Jain and Tarafder, 1970; Anonymous, 1978). Ginger along with *P. longum* is used as an abortifacient in some tribes (Tarafder, 1983).

In veterinary medicine too ginger is important against many livestock maladies. Iqbal *et al.* (2006) reported the anthelmintic activity of crude powder and crude aqueous extract of dried ginger in sheep naturally infected with gastrointestinal nematodes.

Pharmacology

Both the volatile and non-volatile compounds of ginger are credited with medicinal properties besides imparting pungency and aroma to ginger as a spice (Fig. S1, available online only at http://journals.cambridge.org). The main volatile compounds are mono- and sesqui-terpenes, camphene, phellandrene, curcumene, cineole, geranyl acetate, terphineol, terpenes, borneol, geraniol, limonene, linalool, zingiberene, sesqui-phellandrene, bisabolene and farnesene. Many of these compounds are credited with curative properties. (Feng and Lipton, 1987; Zebovitz, 1989; Keeler and Tu, 1991; Yamahara *et al.*, 1992; Hansel *et al.*, 1992; Reddey and Lokesh, 1992; Martin *et al.*, 1993; Denyer *et al.*, 1994; Leung and Foster, 1995; Newall *et al.*, 1996; Obeng-Ofori and Reichmuth, 1997; Blascheck *et al.*, 1998; Atsumi *et al.*, 2000; Coppen, 2002; Murakami *et al.*, 2003; Zhou *et al.*, 2004; Juergens *et al.*, 2004; Masuda *et al.*, 2004; Ali *et al.*, 2005; Hsu *et al.*, 2005; Kuo *et al.*, 2005; Shin *et al.*, 2005; Rao and Rao, 2005; Riyazi *et al.*, 2007; Chen *et al.*, 2007; Shukla and Singh, 2007; Lee *et al.*, 2009; El-Baroty *et al.*, 2010; Hsu *et al.*, 2010). β-Phellandrene and zingiberene are the major compounds in Indian commercial gingers. Ginger oil *per se* is also having antiulcer, antidepressant and anti-inflammatory properties (Sharma *et al.*, 1997; Khushtar *et al.*, 2009; Qiang *et al.*, 2009).

The major non-volatile pungent compounds of ginger, the gingerols and shogaols, too possess analgesic, antipyretic, cardio tonic, antioxidant, and anti-inflammatory properties besides suppressing cytokine formation and promoting angiogenesis. (Mascolo *et al.*, 1989; Yamahara *et al.*, 1989; Yamahara and Huang, 1990; Huang *et al.*, 1991; Mustafa *et al.*, 1993; Lee and Surh, 1998; Surh *et al.*, 1998; Surh, 1999; Bhattarai *et al.*, 2001; Chung *et al.*, 2001; Ficker *et al.*, 2003; Jolad *et al.*, 2004; Isa *et al.*, 2008; Pan *et al.*, 2008; Park *et al.*, 2009; Koh *et al.*, 2009; Imm *et al.*, 2010). Gingerol, particularly 6-gingerol, has been found to be the most active compound biologically (Yamahara *et al.*, 1989; Mascolo *et al.*, 1989; Huang *et al.*, 1991; Mustafa *et al.*, 1993; Aeschbach *et al.*, 1994; Agarwal *et al.*, 2001; Ficker *et al.*, 2003; Kim *et al.*, 2005; Tripathi *et al.*, 2006; Lam *et al.*, 2007; Sjmonatj, 2009; Jeong *et al.*, 2009; Wu *et al.*, 2010; Yang *et al.*, 2010).

Biologically active compounds of ginger are given in the Table S2, available online only at http://journals.cambridge.org.

Ginger has strong antibacterial and to some extent antifungal properties too (Kapoor, 1997; Habsah *et al.*, 2000; Srinivasan *et al.*, 2001). *In vitro* studies have shown that active constituents of ginger inhibit multiplication of colon bacteria (Bakhru, 1999). Ginger also inhibits the growth of *Escherichia coli*, *Proteus* spp., *Staphylococci*, *Streptococci* and *Salmonella* (Gugnani and Ezenwanze, 1985; James *et al.*, 1999). O'Mahony *et al.* (2005) reported curative effects against *Helicobacter pylori*. Fresh ginger juice showed inhibitory action against *Aspergillus niger*, *Saccharomyces cerevisiae*, *Mycoderma* spp. and *Lactobacillus acidophilus* at 4, 10, 12 and 14%, respectively, at ambient temperatures (Nanir and Kadu, 1987; Meena, 1992; Kapoor, 1997). Martins *et al.* (2001) demonstrated antimicrobial activity of essential oil against gram-positive and gram-negative bacteria using agar diffusion method.

Conclusion

Ginger is one of the most important and ancient spices. Though ginger is sterile, plentiful cultivar diversity, recognized mostly by local names exists. However, most of the molecular/biochemical characterization studies attempted so far failed to reveal significant polymorphism commensurating with the observed morphological variability or yield. A geographical bias in the ginger germplasm conserved in the *ex situ* genebanks, probably due to local (vernacular) identity problems during germplasm collection and the geographical effect on key constituents are the hallmark of most of the germplasm characterization studies. Variability for the essential oil profile as well as the pungent constituents is also not that remarkable though few primitive

https://doi.org/10.1017/S1479262111000670 Published online by Cambridge University Press

cultivars with excellent quality traits have been reported from different countries. The primitive types (obsolete cultivars), invariably poor yielders on the verge of extinction, require special attention in long-term conservation strategy especially because convergent improvement or hybridization is rather difficult in a sterile ginger. Pharmacokinetic studies on the long-term effect of the curative compounds are still in infancy. Taxonomic and pharmacological studies on the subspecies of *Z. officinale* will also be rewarding. In future, there may be premium for tailor-made ginger varieties to meet a specific end use. Thus, research on the effect of environmental factors including the changing climate on quality profile and aroma of ginger, pharmacokinetics of the medicinally important constituents, problem of sterility and above all a comprehensive characterization study on the global germplasm should assume priority.

References

Abraham Z and Latha M (2003) Correlation and path analysis in ginger (*Zingiber officinale* Rosc.). *Journal of Spices and Aromatic Crops* 12: 187–189.

Aeschbach R, Löliger J, Scott BC, Murcia A, Butler J, Halliwell B and Aruoma OI (1994) Antioxidant actions of thymol, carvacrol, 6-gingerol, zingerone and hydroxytyrosol. *Food Chemistry and Toxicology* 32: 31–36.

Agarwal M, Walia S, Dhingra S and Khambay BP (2001) Insect growth inhibition, antifeedant and antifungal activity of compounds isolated/derived from *Zingiber officinale* Roscoe (ginger) rhizomes. *Pest Management Science* 57: 289–300.

Aimbire F, Penna SC, Rodrigues M, Rodrigues KC, Lopes-Martins RAB and Sertié JAA (2007) Effect of hydroalcoholic extract of *Zingiber officinalis* rhizomes on LPS-induced rat airway hyper reactivity and lung inflammation. *Prostaglandins Leukotrienes and Essential Fatty Acids* 77: 129–138.

Aiyadurai SGA (1996) *Review of Research on Spice and Cashewnut in India. Regional Office (Spices and Cashewnut)*. Ernakulam: Indian Council of Agricultural Research, p. 228.

Ali SM, Khan AA, Ahmed I, Musaddiq M, Ahmed KS, Polasa H, Rao V, Habibullah CM and Sechi LA (2005) Antimicrobial activities of eugenol and cinnamaldehyde against the human gastric pathogen *Helicobacter pylori*. *Annals of Clinical Microbiology and Antimicrobes* 4: 20–27.

Anonymous (1978) *Hand Book of Domestic Medicine and Common Ayurvedic Remedies*. New Delhi: Central Council for Research in Ayurveda and Siddha (CCRAS).

Anonymous (1982) Pharmacognosy of indigenous drugs (Vol-E). Raghunathan K and Roma Mitra (eds), vol. I. New Delhi: Central Council for Research in Ayurveda and Siddha, pp. 52–76.

Ansari MN and Bhandari U (2008) Cardioprotective activity of *Zingiber officinale* Roscoe in rats. *Hamdard Medicus* 51: 38–41.

Arctangder S (1960) *Perfume and Flavour Materials of Natural Origin*. New Jersey: Elizabeth, p. 275.

Arya PS and Rana KS (1990) Performance of ginger varieties in Himachal Pradesh. *Indian Cocoa, Arecanut and Spices Journal* 14: 16–20.

Atman RD and Marcussen KC (2001) Effects of a ginger extract on knee pain in patients with osteoarthritis. *Arthritis and Rheumatism* 44: 2531–2538.

Atsumi T, Fujisawa S, Satoh K, Sakagami H, Iwakura I, Ueha T, Sugita Y and Yokoe I (2000) Cytotoxicity and radical intensity of eugenol, isoeugenol or related dimers. *Anti Cancer Research* 20: 2519–2524.

Baker JG (1882) Scitaminae. In: Hooker JD (ed.) *The Flora of British India*. vol. VI. Dehra Dun: Bishen Singh, Mahendrapal Singh, pp. 198–264.

Bakhru HK (1999) *Herbs that Heal: Natural Remedies for Good Health*. New Delhi: Oriental Paper Backs, A Division of Vision Books Pvt. Ltd, p. 97.

Bhattarai S, Tran VH and Duke CC (2001) The stability of gingerol and shogoal in aqueous solutions. *Journal of Pharmaceutical Science* 90: 1658–1664.

Blascheck W, Hansel R, Keller K, Reichling J, Rimpler H and Schneider G (eds) (1998) Hager's Handbuch der Pharmazeutischen Praxis, Auflage Band 2 (A–K), 909 pp., (L–Z), 858 pp. Berlin: Springer-Verlag.

Broussard CS, Louik C, Honein MA and Mitchell AA (2010) Herbal use before and during pregnancy. *American Journal of Obstetrics and Gynecology* 202: 443e1–443e6.

Burkill IH (1996) *A Dictionary of the Economic Products of the Malay Peninsula*. Kuala Lumpur: Ministry of Agriculture and Co-operatives, p. 2444.

Burtt BL and Smith RM (1972) Tentative keys to the subfamilies, tribes and genera of the Zingiberales. *Notes from the Royal Botanical Garden Edinburgh* 31: 171–176.

Chandra R and Govind S (1999) Genetic variability and performance of ginger genotypes under mid hills of Meghalaya. *Indian Journal of Horticulture* 56: 274–278.

Chen JC, Huang LJ, Wu SL, Kuo SC, Ho TY and Hsiang CY (2007) Ginger and its bioactive component inhibit enterotoxigenic *Escherichia coli* heat-labile Enterotoxin-induced diarrhea in mice. *Journal of Agriculture and Food Chemistry* 55: 8390–8397.

Chengkun H, Jiashen L, Suzhi G, Min Z and Wensheng C (1995) The relationship between geographic distribution and the genetic difference of peroxidase isozyme of ginger germplasm in Fujian. *Acta Horticulturae* 402: 125–132.

Chung WY, Jung YJ, Surh YJ, Lee SS and Park KK (2001) Antioxidative and antitumor promoting effects of [6]-paradol and its homologs. *Mutation Research, Genetic Toxicology and Environmental Mutagenesis* 496: 199–206.

Connel DW and Jordan RA (1971) Composition and distinctive volatile flavour characteristics of the essential oil from Australian-grown ginger (*Zingiber officinale*). *Journal of Science Food and Agriculture* 22: 93–95.

Coppen JJW (2002) *Eucalyptus: The Genus Eucalyptus (Mediacinal and Aromatic Plants – Industrial Profiles)*. London and NewYork: Taylor and Francis, p. 440.

Denyer CV, Jackson P, Loakes DM, Ellis MR and Yound DAB (1994) Isolation of antirhinoviral sesqui-terpenes from ginger (*Zingiber officinale*). *Journal of Natural Products* 57: 658–662.

Devasahayam S, Jacob TK, Koya KMA and Sasikumar B (2010) Screening of ginger (*Z. officinale* R) germplasm for resistnace to shoot borer (*Conogethes punctiferalis)*. *Journal of Medicinal and Aromatic Plant Science* 32: 137–138.

https://doi.org/10.1017/S1479262111000670 Published online by Cambridge University Press

Ekundayo O, Laakso I and Hiltunen R (1988) Composition of ginger (*Zingiber officinale* Roscoe) volatile oils from Nigeria. *Flavour and Fragrance Journal* 3: 85–90.

El-Baroty GS, El-Baky HHA, Farag RS and Saleh MA (2010) Characterization of antioxidant and antimicrobial compounds of cinnamon and ginger essential oils. *African Journal of Biochemistry Research* 4: 167–174.

Fakim AG, Maudarbaccus N, Leach D, Doimo L and Wohlmuth H (2002) Essential oil composition *Zingiberaceae* species from Mauritius. *Journal of Essential Oil Research* 14: 271–273.

Feng J and Lipton JM (1987) Eugenol: antipyretic activity in rabbits. *Neuropharmacology* 26: 1775–1778.

Ficker C, Smith ML, Akpagana K, Gbeassor M, Zhang J, Durst T, Assabgui R and Arnason JT (2003) Bioassay-guided isolation and identification of antifungal compounds from ginger. *Phototherapy Research* 17: 897–902.

Gowda KK and Melanta KR (2000) Varietal performance of ginger in Karnataka. In: Muraleedharan N and Rajkumar R (eds) *Recent Advances in Plantation Crops Research*. New Delhi: Allied Publishers, pp. 92–93.

Grant KL and Lutz RB (2000) Alternative therapies: ginger. *American Journal of Health-System Pharmacy* 57: 945–947.

Gugnani HC and Ezenwanze EC (1985) Antibacterial activity of extracts of ginger (*Zingiber officinale*) and African oil bean seed (*Pentaclethora macrophylla*). *Journal of Communicable Diseases* 17: 233–236.

Habsah M, Amran M, Mackeen MM, Lajis NH, Kikuzaki H, Nakatani N, Rahman AA, Ghafar and Ali AM (2000) Screening of *Zingiberaceae* extracts for antimicrobial and antioxidant activities. *Journal of Ethnopharmacology* 72: 403–410.

Hansel R, Keller K, Rimpler H and Schneider G (eds) (1992) Hager's Handbuch der Pharmazeutischen Praxis, Drogen (A–D). Berlin: Springer-Verlag, p. 1209.

Heimes K, Feistel B and Verspohl EJ (2009) Impact of the 5-HT_3 receptor channel system for insulin secretion and interaction of ginger extracts. *European Journal of Pharmacology* 624: 58–65.

Hirara K and Takesma M (1998) *Spices – Science and Technology*. New York, NY: Marcel Deckker.

Holtum RE (1950) The *Zingiberaceae* of the Malay Peninsula. *Gardens Bulletin (Singapore)* 13: 1–50.

Hsu MH, Kuo SC, Chen CJ, Chung JG, Lai YY and Huang LJ (2005) 1-(3,4-Dimethoxyphenyl)-3,5-dodecenedeone (16) induces GI arrest and apoptosis in human promyclocytic leukemia HL-60 cells. *Leukemia Research* 29: 1399–1406.

Hsu YL, Chen CY, Hou MF, Tsai EM, Jong YJ, Hung CH and Kuo PL (2010) 6-Dehydrogingerdione, an active constituent of dietary ginger, induces cell cycle arrest and apoptosis through reactive oxygen species/c-Jun N-terminal kinase pathways in human breast cancer cells. *Molecular Nutrition and Food Research* 54: 1307–1317.

Huang QR, Iwamoto M, Aoki S, Tanaka N, Tajima K, Yamahara J, Takaishi Y, Yoshida M, Tomimatsu T and Tamai Y (1991) Anti-5-hydroxitryptamine 3 effect of galanolactone, diterpenoid isolated from ginger. *Chemical Pharma Bulletin* 39: 397–399.

Imm JY, Zhang GD, Chan LY, Nitteranon V and Parkin KL (2010) [6]-Dehydroshogaol, a minor component in ginger rhizome, exhibits quinone reductase inducing and anti-inflammatory activities that rival those of curcumin. *Food Research International* 43: 2208–2213.

Indrasenan G and Paily PV (1974) Studies on the soft rot of ginger (*Z. officinale*) caused by *Pythium aphanidermatum* (Edson) Fitz. *Agricultural Research Journal of Kerala* 11: 53–56.

Iqbal Z, Latheef M, Akhtar MS, Ghayur MN and Gilani AH (2006) *In vivo* anthelmintic activity of ginger against gastrointestinal nematodes of sheep. *Journal of Ethanopharmacology* 106: 285–287.

Isa Y, Miyakawa Y, Yanagisawa M, Goto T, Kang MS, Kawada T, Morimitsu Y, Kubota K and Tsuda T (2008) 6-Shogaol and 6-gingerol, the pungent of ginger, inhibit TNF-α mediated down regulation of adiponectin expression via different mechanisms in 3T3-L1 adipocytes. *Biochemical and Biophysical Research Communications* 373: 429–434.

Jain SK and Tarafder CR (1970) Medicinal plant lore of the Santals. A revival of P.O. Bodding's Work. *Economic Botany* 24: 241–278.

James ME, Nannapaneni R and Johnson M (1999) Identification and characterization of two bacteriocin producing bacteria isolated from garlic and ginger root. *Journal of Food Protection* 62: 899–904.

Jatoi SA, Kikuchi A, Yi SS, Naing KW, Yamanaka S, Watanabe JA and Watanabe KN (2006) Use of rice SSR markers as RAPD markers for genetic diversity analysis in Zingiberaceae. *Breeding Science* 56: 107–111.

Jeong CH, Bode AM, Pugliese A, Cho YY, Kim HG, Shim JH, Jeon YJ, Li HL, Jiang HL and Dong ZG (2009) Gingerol suppresses colon cancer growth by targetting leukotriene A_4 hydrolase. *Cancer Research* 69: 5584–5591.

John RP and Ferreira DI (1997) Yield and quality evaluation of five ginger (*Zingiber officinale* Roscoe) selections in the Northern Province of South Africa. *Journal of the Southern African Society for Horticultural Sciences* 7: 26–29.

Jolad SD, Lantz RC, Solyom AM, Chen GJ, Bates RB and Timmermann BN (2004) Fresh organically grown ginger (*Zingiber officinale*): composition and effects on LPS-induced PGE2 production. *Phytochemistry* 65: 1937–1954.

Juergens U, Engelen T, Racké K, Stöber M, Gillissen A and Vetter H (2004) Inhibitory activity of 1,8-cineol (eucalyptol) on cytokine production in cultured human lymphocytes and monocytes. *Pulmonary Pharmacology and Therapeutics* 17: 281–287.

Kapoor A (1997) Antifungal activities of fresh juice and aqueous extracts of turmeric and ginger (*Zingiber officinale*). *Journal of Phytological Research* 10: 59–62.

Keeler RF and Tu AT (1991) *Handbook of Natural Toxins*. vol. 6. New York, NY: Marcel Dekker, Inc., p. 665.

Khan KI (1959) Ensure two fold ginger yields. *Indian Farming* 8: 10–14.

Khushtar M, Kumar V, Javed K and Uma Bhandari (2009) Protective effect of ginger oil on aspirin and pylorus ligation-induced gastric ulcer model in rats. *Indian Journal of Pharmaceutical Sciences* 71: 554.

Kim SO, Kundu JK, Shin YK, Park JH, Cho MH, Kim TY and Surh YJ (2005) [6]-Gingerol inhibits COX-2 expression by blocking the activation of p38 MAP kinase and NF-kappaB in phorbol ester-stimulated mouse skin. *Oncogene* 24: 2558–2567.

Kizhakkayil J (2008) Molecular and biochemical characterization of ginger (*Z. officinale* R) germplasm. PhD Thesis. Guntur: Acharya Nagarjuna University, p. 173.

Kizhakkayil J and Sasikumar B (2009) Variability for quality traits in a global germplasm collection of ginger (*Z. officinale* R.). *Current Trends in Biotechnology and Pharmacy* 3: 254–259.

https://doi.org/10.1017/S1479262111000670 Published online by Cambridge University Press

Kizhakkayil J and Sasikumar B (2010) Molecular characterisation of ginger (*Zingiber officinale* Rosc.) germplasm – a medicinal spice. *Scientia Horticulture* 125: 73–76.

Koh EM, Kim HJ, So HK, Choi WH, Choi YH, Ryu SY, Kim YS, Koh WS and Park SY (2009) Modulation of macrophage functions by compounds isolated from *Zingiber officinale*. *Planta Medica* 75: 148–151.

Kress WJ, Prince LM and Williams KJ (2002) The phylogeny and a new classification of the gingers (*Zingiberaceae*): evidence from molecular data. *American Journal of Botany* 89: 1682–1696.

Krishnamurthy N, Mathew AG, Nambudiri ES and Lewis YS (1972) Essential oils and oleoresin from major spices in India. *Journal of Plantation Crop, (Suppl)*: 181–183.

Kumar A and Hayward AC (2005) Bacterial diseases of ginger and their control. In: Ravindran PN and Nirmal Babu K (eds) *Ginger – The Genus Zingiber*. Boca Raton, FL: CRC Press, pp. 341–346.

Kumar VS, Balasubramanian PP, Kumar KV and Mammen MK (1980) Comparative performance of six varieties of ginger for second crop in Wynad. *Indian Spices* 17: 111–114.

Kuo PC, Dhamu AG, Cheng CY, Teng CM, Lee EJ and Wu TS (2005) Isolation of a natural antioxidant, dehydrozingerone from *Zingiber officinale* and synthesis of its analogues for recognition of effective antioxidant and antityrosinase agent. *Archives of Pharmacology Research* 28: 518–528.

Lakshmi BVS and Sudhakar MA (2010) Protective effect of *Z. officinale* on gentamicin induced nephrotoxicity in rats. *International Journal of Pharmacology* 6: 58–62.

Lam RYY, Woo AYH, Leung PS and Cheng CHK (2007) Antioxidant actions of phenolic compounds found in dietary plants on low-density lipoprotein and erythrocytes in vitro. *Journal of the American College of Nutrition* 26: 233–242.

Langner E, Greifenberg S and Gruenwald J (1998) Ginger: history and use. *Advances in Therapy* 15: 25–44.

Larsen K and Larsen SS (2006) *Zingiber Miller. Ginger of Thailand*. Chiang Mai: Queen Sirikit Botanic Garden, p. 184.

Lawrence BM (1984) Major tropical spices – ginger (*Zingiber officinale* Rosc.). *Perfumer and Flavourist* 9: 1–40.

Lee E and Surh YJ (1998) Induction of apoptosis in HL-60 cells by pungent vanilloids, [6]-gingerol and [6]-paradol. *Cancer Letters* 134: 163–168.

Lee SY, Fai WK, Zakaria M, Ibrahim H, Othman RY, Gwang JG, Rao VR and Park YJ (2007) Characterization of polymorphic microsatellite markers, isolated from ginger (*Zingiber officinale* Rosc.). *Molecular Ecology Note*. Doi: 10.1111/j.1471-8286.2007.01757.x.

Lee JY, Kang HS, Park BE, Moon HJ, Sim SS and Kim CJ (2009) Inhibitory effects of Geijigajakyak-Tang on trinitrobenzene sulfonic acid-induced colitis. *Journal of Ethnopharmacology* 126: 244–251.

Leung AY and Foster S (1995) *Encyclopedia of Common Natural Ingredients*. 2nd edn. New York, NY: John Wiley and Sons, p. 649.

Liang MH (1992) From America: cookbook medicine or food for thought: practice guidelines development in USA. *Annals of the Rheumatic Diseases* 51: 1257–1258.

Lincy AK, Jayarajan K and Sasikumar B (2008) Relationship between vegetative and rhizome characters and final rhizome yield in micro propagated ginger plants (*Zingiber officinale* Rosc.) over two generations. *Scientia Horticulture* 118: 70–73.

Macloed AJ and Pieris NM (1984) Volatile aroma constituents of Sri Lankan ginger. *Phytochemistry* 23: 353–360.

Martin S, Padilla E, Ocete MA, Galvez J, Jimenez J and Zarzuelo A (1993) Anti-inflammatory activity of the essential oil of *Bupleurum fruticescens*. *Planta Medica* 59: 533–536.

Martins AP, Salgueiro L, Goncalves MJ, da Cunha AP, Vila R and Canigueral S (2001) Essential oil composition and antimicrobial activity of three *Zingiberaceae* from S.Tome e Principe. *Planta Medica* 67: 580–584.

Mascolo N, Jain R, Jain SC and Capasso F (1989) Ethnopharmacological investigation of ginger (*Zingiber officinale*). *Journal of Ethnopharmacology* 27: 129–140.

Masuda Y, Kikuzaki H, Hisamoto M and Nakalani N (2004) Antioxidant properties of gingerol related compounds from ginger. *Biofactors* 21: 129–140.

Meena MR (1992) *Studies on antimicrobial activity of various spices and their oils*. MSc Thesis. New Delhi: Indian Agricultural Research Institute.

Menon AN (2007) Chemical composition of the essential oil from leaves and roots of ginger (*Zingiber officinale*) from Kerala. *Journal of Medicinal and Aromatic Plant Sciences* 29: 78–80.

Menut C, Lamaty G, Bessiere JM and Koudou J (1994) Aromatic plants of tropical Central Africa. XIII. Rhizomes volatile components of two *Zingiberales* from the Central African Republic. *Journal of Essential Oil Research* 6: 161–164.

Mohandas TP, Pradeep Kumar T, Mayadevi P, Aipe KC and Kumaran K (2000) Stability analysis in ginger (*Zingiber officinale Rosc.*) genotypes. *Journal of Spices and Aromatic Crops* 9: 165–167.

Mohanty DC and Sharma YN (1979) Genetic variability and correlation for yield and other variables in ginger germplasm. *Indian Journal of Agricultural Sciences* 49: 250–253.

Mohanty DC, Das RC and Sharma YN (1981) Variability of agronamic of ginger. *Orissa Journal of Horticuture* 9: 15–17.

Morakinyo AO, Adeniyi OS and Arikawe AP (2008) Effects of *Zingiber officinale* on reproductive functions in the male rat. *African Journal of Biomedical Research* 11: 329–333.

Muda MM, Halijah I and Norzulaani K (2004) Differentiation of three varieties of *Zingiber officinale* Rosc. by RAPD fingerprinting. *Malaysian Journal of Science* 23: 135–139.

Murakami Y, Shoji M, Hanazawa S, Tanaka S and Fujisawa S (2003) Preventive effect of bis-eugenol, a eugenol ortho dimer, on lipopolysaccharide-stimulated nuclear factor kappa B activation and inflammatory cytokine expression in macrophages. *Biochemical Pharmacology* 66: 1061–1066.

Muralidharan A and Kamalam N (1973) Improved ginger means foreign exchange. *Indian Farming* 22: 37–39.

Mustafa T, Srivastava KC and Jensen KB (1993) Drug development report. 9. Pharmacology of ginger, *Zingiber officinale*. *Journal of Drug Development* 6: 25–39.

Naidu MM, Padma M, Yuvaraj KM and Murty PSS (2000) Evaluation of ginger varieties for high altitude and tribal area of Andra Pradesh. Spices and Aromatic Plants: Challenges and opportunities in the new century. Centennial conference on spices and aromatic plants, 20–23 Sept. 2000, Calicut, Kerala, India, pp. 50–51.

Nammi S, Satyanarayana S and Roufogalis BD (2009) Protective effects of ethanolic extract of *Zingiber officinale* rhizome on the development of metabolic syndrome in high-fat diet-fed rats. *Basic and Clinical Pharmacology and Toxicology* 104: 366–373.

Nammi S, Kim MS, Gavande NS, Li GQ and Roufogalis BD (2010) Regulation of low-density lipoprotein receptor and 3-hydroxy-3-methylglutaryl coenzyme A reductase expression by *Zingiber*

https://doi.org/10.1017/S1479262111000670 Published online by Cambridge University Press

officinale in the liver of high-fat diet-fed rats. *Basic and Clinical Pharmacology and Toxicology* 106: 389–395.

Nanir SP and Kadu BB (1987) Effect of medicinal plant extracts on some fungi. *Acta Botanica Indica* 15: 170–175.

Natarajan CP, Kuppuswamy S, Shankaracharya NB, Padmabai R, Raghavan B, Krishnamurthy MN, Khan F, Lewis YS and Govindarajan VS (1972) Chemical composition of ginger varieties and dehydration studies of ginger. *Journal of Food Science and Technology* 9: 120–124.

Nayak S, Naik PK, Acharya L, Mukherjee AK, Panda PC and Das P (2005) Assessment of genetic diversity among 16 promising cultivars of ginger using cytological and molecular markers. *Zeitschrift fur Naturforschung* 60C: 485–492.

Newall CA, Anderson LA and Phillipson JD (1996) *Herbal Medicine – A Guide for Health-care Professionals*. London: The Pharmaceutical Press, p. 296.

Nybe EV and Nair PCS (1979) Studies on the morphology of the ginger types. *Indian Cocoa, Arecanut and Spices Journal* 3: 7–13.

Nybe EV and Nair PCS (1979a) Field tolerance of ginger types to important pests and diseases. *Indian Cocoa, Arecanut and Spices Journal* 2: 109–111.

Nybe EV, Nair PCS and Mohanakumaran N (1980) Assessment of quality components in ginger. In: Nair MK, Premkumar T, Ravindran PN and Sarma YR (eds) *Proceedings, National Seminar on Ginger and Turmeric, 8–9 April 1980, Calicut*. Central Plantation Crops Research Institute, Kasargod, India, pp. 24–29.

Obeng-Ofori D and Reichmuth C (1997) Bioactivityof eugenol, a major component of essential oil of *Ocimum suave* (Wild.) against four species of stored-product Coleoptera. *International Journal of Pest Management* 43: 89–94.

O'Mahony R, Al-Khtheeri H, Weerasekera D, Fernando N, Vaira D, Holton J and Christelle BC (2005) Bactericidal and anti-adhesive properties of culinary and medicinal plants against *Helicobacter pylori*. *World Journal Gastroenterology* 11: 7499–7507.

Ozgoli G, Goli M and Simbar M (2009) Effects of ginger capsules on pregnancy, nausea, and vomiting. *Journal of Alternative and Complementary Medicine* 15: 243–246.

Palai SK and Rout GR (2007) Identification and genetic variation among eight varieties of ginger by using random amplified polymorphic DNA markers. *Plant Biotechnology* 24: 417–420.

Pan MH, Hsieh MC, Kuo JM, Lai CS, Wu H, Sang SM and Ho CT (2008) 6-Shog gaol induces apoptosis in human colorectal carcinoma cells *via* ROS production, caspase activation, and GADD 153 expression. *Molecular Nutrition and Food Researh* 52: 527–537.

Park EJ and Pizzuto JM (2002) Botanicals in cancer chemoprevention. *Cancer Metastasis Review* 21: 231–255.

Park SJ, Lee MY, Son BS and Youn HS (2009) K1-targeted suppression of TRIF-dependent signaling pathway of Toll-like receptors by 6-Shogaol, an active component of ginger. *Bioscience, Biotechnology and Biochemistry* 73: 1474–1478.

Petersen OG (1889) Musaceae, Zingiberaceae, Cannaceae, Marantaceae. In: Engler A and Prant K (eds) *Die Naturlichen Pflanzenfamilien*. vol. 2. Leipzig: Verlag von Wilhelm Engelmann, pp. 1–43.

Prem J, Kizhakkayil J, Thomas E, Dhanya K, Syamkumar S and Sasikumar B (2007) Molecular characterization of primitive, elite and exotic ginger genotypes to protect the biowealth of elite ginger accessions. *SYMSAC IV. Threats and Solution to Spices and Aromatic Crops Industry, National symposium on spices and aromatic crops, 25–26, November 2007*. Bhuvaneswar, Orissa, India, p. 337.

Pruthy JS (1993) *Major spices of India: crop management post harvest technology*. New Delhi: Indian Council of Agricultural Research, pp. 244–288.

Purseglove JW, Brown EG, Green CL and Robbins SRJ (1981) *Spices*. vol. 2. New York, NY: Longman Inc.

Qiang LQ, Wang CP, Wang FM, Pan Y, Yi LT, Zhang X and Kong LD (2009) Combined administration of the mixture of honokiol and magnolol and ginger oil evokes antidepressant-like synergism in rats. *Archives of Pharmacal Research* 32: 1281–1292.

Rahman H, Karuppaiyan R, Kishore K and Denzongpa R (2009) Traditional practices of ginger cultivation in north east India. *Indian Journal of Traditional Knowledge* 8: 23–28.

Rana N and Korla BN (2007) Evaluation of ginger germplasm for yield and quality under mid hill conditions of Himachal Pradesh. *Haryana Journal of Horticultural Sciences* 36: 393–394.

Rao RR and Jamir NS (1982) Ethnobotanical studies in Nagaland-1. Medicinal Plants. *Economic Botany* 36: 176–181.

Rao BN and Rao BSS (2005) Antagonistic effects of Zingerone, a phenolic alkanone against radiation-induced cytotoxicity, genotoxicity, apoptosis and oxidative stress in Chinese hamster lung fibroblast cells growing *in vitro*. *Mutagenesis* 25: 577–587.

Ratnambal MJ, Gopalam A and Nair MK (1987) Quality evaluation in ginger (*Zingiber officinale* Rosc.) in relation to maturity. *Journal of Plantation Crops* 15: 108–117.

Rattan RS, Korla BN and Dohroo NP (1988) Performance of ginger varieties in Solan area of Himachal pradesh. In: Satyanarayana G, Reddy MS, Rao MR, Azam KM and Naidu R (eds) *Proceedings of National seminar on Chillies, Ginger and Turmeric*. Cochin: Spices Board, pp. 71–73.

Ravindran PN, Sasikumar B, George JK, Ratnambal MJ, Nirmal Babu K, Zachariah TJ and Nair RR (1994) Genetic recourses of ginger (*Zingiber officinale* Rosc.) and its conservation in India. *Plant Genetic Resources News Letter* 98: 1–4.

Ravindran PN, Remasree AB and Sherlija KK (1998) *Developmental Morphology of Rhizomes of Ginger and Turmeric. Final Report. ICAR Ad hoc Scheme*. Calicut: Indian Institute of Spices Research.

Ravindran PN, Nirmal Babu K and Shiva KN (2005) Botany and crop improvement of ginger. In: Ravindran PN and Nirmal Babu K (eds) *Ginger – The Genus Zingiber*. Boca Raton, FL: CRC Press, pp. 15–85.

Ravindran PN, Shiva KN, Nirmal Babu K and Korla BN (2006) Ginger. In: Ravindran PN, Nirmal Babu K, Shiva KN and Johny AK (eds) *Advances in Spices Research*. Jodhpur: Agrobios, pp. 365–432.

Reddey AC and Lokesh BR (1992) Studies on spice principles as antioxidants in the inhibition of lipid peroxidation of rat liver microsomes. *Molecular and Cellular Biochemistry* 111: 117–124.

Ridley HN (1912) *Spices*. London: Mcmillan.

Riyazi A, Hensel A, Bauer K, Geissler N, Schaaf S and Verspohl EJ (2007) The effect of the volatile oil from ginger rhizomes (*Zingiber officinale*), its fractions and isolated compounds on the 5-HT_3 receptor complex and the serotoninergic system of the rat ileum. *Planta Medica* 73: 355–362.

Rosengarten FJ (1969) *The Book of Spices*. London: Macmillan and Co.

https://doi.org/10.1017/S1479262111000670 Published online by Cambridge University Press

Sabu M (2003) Revision of the genus *Zingiber* in South India. *Folia Malaysiana* 4: 25–52.

Sasikumar B, Nirmal Babu K, Abraham J and Ravindran PN (1992) Variability, correlation and path analysis in ginger germplasm. *Indian Journal of Genetics and Plant breeding* 52: 428–431.

Sasikumar B, Ravindran PN and George JK (1994) Breeding ginger and turmeric. *Indian Cocoa, Arecanut and Spices Journal* 18: 10–12.

Sasikumar B, Saji KV, Ravindran PN and Peter KV (1999) Genetic resources of ginger (*Zingiber officinalle* Rosc.) and its conservation in India. In: Sasikumar B, Krishnamurthy B, Rema J, Ravindren PN and Peter KV (eds) *Proceedings of the National Seminar on Biodiversity, Conservation and Utilisation of Spices, Medicinal and Aromatic Plants*. IISR, Calicut, pp. 96–100.

Sasikumar B, John T Zachariah and Shamina A (2000) Characterisation of ginger germplasm based on isozyme polymorphism. In: Muraleedharan N and Raj Kumar R (eds) *Recent Advances in Plants. Crops. Res.* (PLACROSYM XIII). NewDelhi: Allied Publishers, pp. 251–255.

Sasikumar B, Saji KV, Antony A, George JK, John Zachariah T and Eapen SJ (2003) IISR Mahima and IISR Rejatha – two new high yielding quality ginger (*Zingiber officinale* Rosc.). *Journal of Spices and Aromatic Crops* 12: 34–37.

Sasikumar B, Krishnamurthy KS, Kandiannan K and Srinivasan V (2007) Turmeric. In: Dinesh R, Shiva KN, Prasath D, Babu KN and Parthasarathy VA (eds) *Threats and Solutions to Spices and Aromatic Crops Industry. National Symposium on Spices and Aromatic Crops, Bhubaneswar, Orissa, India*, pp. 229–259.

Sasikumar B (2008) Ginger in Guyana. *Spice India* 21: 7.

Setty TAS, Guruprasad TR, Mohan E and Reddy MNN (1995) Susceptibility of ginger cultivars to rhizome rot at west coast of conditions of Karnataka. *Environmental Ecology* 13: 443–444.

Sharma JN, Ishak FI, Yusof APM and Srivastava KC (1997) Effects of eugenol and ginger oil on adjuvant arthritis and the kallikreins in rats. *Asia Pacific Journal of Pharmacology* 12: 9–14.

Sharma H and Clark C (1998) *Contemporary Ayurveda*. London: Churchill Living Stone, pp. 100–101.

Shin SG, Kim JY, Chung HY and Jeong JC (2005) Zingerone as an antioxidant against peroxynitrite. *Journal of Agricultural and Food Chemistry* 53: 7617–7622.

Shukla Y and Sing M (2007) Cancer preventive properties of ginger: a brief review. *Food and Chemical Toxicology* 45: 683–690.

Shylaja MR, Paul R, Nybe EV, Abraham K, Nazeem PA, Nazeem PA, Valsala PA and Krishnan S (2010) Two new ginger varieties from Kerala Agricultural University. *Indian Journal of Arecanut, Spices and Medicinal Plants* 12: 3–4.

Simmonds NW (1979) *Principles of Crop Improvement*. New York, NY: Longman Group Ltd.

Singh RS (1983) *Vanausadhi Nidarshika (Ayurvedic Pharmacopia)*. II edn. Lucknow: U.P. Hindi Sansthan Rajarshi Puruhottam Das Tandan Hindi Bhavan, Mahatma Gandhi Marg, p. 389.

Singh AK (2001) Correlation and path analysis for certain metric traits in ginger. *Annals of Agricultural Research* 22: 285–286.

Singh PP, Singh VB, Singh A and Singh HB (1999) Evaluation of different ginger cultivars for growth, yield and quality character under Nagaland condition. *Journal of Medicinal and Aromatic Plant Sciences* 21: 716–718.

Singh PP, Singh VB, Singh HP and Rajan S (2000) Genetic diversity in ginger (*Zingiber officinale* Rosc.) with reference to essential oil content. *Journal of Spices and Aromatic Crops* 9: 161–164.

Sjmonatj CN (2009) 6-Gingerol content and atibioactive properties of ginger extracts from supercritical CO_2 extraction. Documentar study. *Ingineria Mediului* 14: 609–615.

Smith MK, Hamill SD, Gogel BJ and Severn-Ellis AA (2004) Ginger (*Zingiber officinale*) autotetraploids with improved processing quality produced by an *in vitro* colchicine treatment. *Australian Journal of Experimental Agriculture* 44: 1065–1072.

Sreekumar V, Indrasenan G and Mamman MK (1980) Studies on the quantitative and qualitative attributes of ginger cultivars. In: Nair MK, Premkumar T, Ravindran PN and Sarma YR (eds) *Proceedings, National Seminar on Ginger and Turmeric, 8–9. April 1980, Calicut*. Central Plantation Crops Research Institute, Kasargod, India, pp. 47–49.

Srinivasan D, Nathan S, Suresh T and Lakshmana Perumalsamy P (2001) Antimicrobial activity of certain Indian medicinal plants used in folkloric medicine. *Journal of Ethnopharmacology* 74: 217–220.

Surh Y (1999) Molecular mechanisms of chemo preventive effects of selected dietary and medicinal phenolic substances. *Mutation Research* 428: 305–327.

Surh YJ, Lee ME and Lee JM (1998) Chemoprotective properties of some pungent ingredients present in red pepper and ginger. *Mutation Research* 402: 259–267.

Tarafder CR (1983) Ethnogynaecology in relation to plants-II. Plants used in abortion. *Journal of Economic and Taxonomic Botany* 4: 507–516.

Thakur RS, Puri HS and Akhtar Husain (1989) Major Medicinal Plants of India. Lucknow: CIMAP, pp. 540–542.

Thomas KM (1966) Rio-de-Janeiro will double your ginger yield. *Indian Farming* 15: 15–18.

Tiwari SK (2003) Evaluation of ginger genotypes for yield and quality attributes under rain fed and irrigated conditions. *Annals of Agricultural Research* 24: 512–515.

Tripathi S, Tripathi S, Maier K, Bruch D and Kittur D (2006) Ginger and its active ingredient 6-gingerol down regulate pro-inflammatory cytokine release by macrophages. *Journal of Surgical Research* 130: 318–324.

Tyler VE, Brady LR and Robbers JE (1981) *Pharmacognosy*. 8th edn. Philadelphia, PA: Lea and Febiger, p. 156.

van Beek TA, Posthumus MA, Lelyveld GP, Hoang VP and Yen BT (1987) Investigation of the essential oil of Vietnamese ginger. *Phytochemistry* 26: 3005–3010.

van Rheede (1692) *Hortus Indicus Malabaricus*. vol II. Amsterdam: Joannis van Someren, et Joannis van Dyck.

Vernin G and Parkanyl C (1994) Ginger oil (*Zingiber officinale* Roscoe). In: Charalambous G (ed.) *Spices, Herbs and Edible Fungi*. Amsterdam: Elsevier Science, pp. 579–594.

Vernin G and Parkanyl C (2005) Chemistry of ginger. In: Ravindran PN and Nirmal Babu K (eds) *Ginger – The Genus Zingiber*. Boca Raton, FL: CRC Press, pp. 87–180.

Wagner H and Hikino H (1965) *Economic and Medicinal Plants Research*. vol. 1. New York, NY: Academic Press, p. 62.

Wahyuni S, Xu DH, Bermawie N, Tsunematsu H and Ban T (2003) Genetic relationships among ginger accessions based on AFLP marker. *Journal Bioteknologi Pertanian* 8: 60–68.

Warrier PK (1989) Spices in ayurveda. In: George CK, Sivadasan CR, Devakaran D and Sreekumari KP (eds) *Strategies*

https://doi.org/10.1017/S1479262111000670 Published online by Cambridge University Press

for Export Development of Spices. Cochin/Geneva: Spices Board/International Trade Centre, p. 28.

Wohlmuth H, Leach DN, Smith MK and Myers SP (2005) Gingerol content of diploid and tetraploid clones of ginger (*Zingiber officinale* Roscoe). *Journal of Agricultural and Food Chemistry* 53: 5772–5778.

Wohlmuth H, Smith MK, Brooks LO, Myers SP and Leach DN (2006) Essential oil composition of diploid and tetraploid clones of ginger (*Zingiber officinale* Roscoe) grown in Australia. *Journal of Agricultural and Food Chemistry* 53: 5772–5778.

Wu H, Hsieh MC, Lo CY, Liu CB, Sang SM, Ho CT and Pan MH (2010) 6-Shogaol is more effective than 6-gingerol and curcumin in inhibiting 12-*O*-tetradecanoylphorbol 13-acetate-induced tumor promotion in mice. *Molecular Nutrition and Food Research* 54: 1296–1306.

Yadav RK (1999) Genetic variability in ginger (*Zingiber officinale* Rosc.). *Journal of Spices and Aromatic Crops* 8: 81–83.

Yamahara J and Huang Q (1990) Gastrointestinal motility enhancing effect of ginger and its active constituents. *Chemical and Pharmaceutical Bulletin* 38: 430–431.

Yamahara J, Rong HQ, Iwamoto M, Kobayashi G, Matsuda H and Fujimura H (1989) Active components of ginger exhibiting antiserotonergic action. *Phototherapy Research* 3: 70–71.

Yamahara J, Hatakeyama S, Taniguichi K, Kawamura M and Yoshikawa M (1992) Stomachic principles in ginger. II. Pungent and anti-ulcer effects of low polar constituents isolated from ginger, the dried rhizoma of *Zingiber officinale* Roscoe, cultivated in Taiwan. The absolute stereostructure of a new diarylheptanoid. *Yakugaku Zasshi* 112: 645–655.

Yang G, Zhong LF, Jiang LP, Geng CY, Cao J, Sun XC and Ma YF (2010) Genotoxic effect of 6-gingerol on human hepatoma G2 cells. *Chemico-Biological Interactions* 185: 12–17.

Zachariah TJ, Sasikumar B and Ravindran PN (1993) Variability in gingerol and shogaol content of ginger accession. *Indian Perfumer* 37: 87–90.

Zachariah TJ, Sasikumar B and Nirmal Babu K (1999) Variation for quality components in ginger and turmeric and their interaction with environments. In: Sasikumar B, Krishnamurthy B, Rema J, Ravindren PN and Peter KV (eds) *Proceedings of National Seminar on Biodiversity, Conservation and Utilisation of spices, Medicinal and Aromatic Plants*. Calicut: IISR, pp. 116–120.

Zebovitz TC (1989) Part VII. Flavor and fragrance substances. In: Keith LH and Walters DB (eds) *Compendium of Safety Data Sheets for Research and Industrial Chemicals*. New York, NY: VCH Publishers, pp. 3560–4253.

Zhen-wei, Shou-Jin Fan, De-min GAO, Zhen-wei LIU and Shou-jin FAN (2006) RAPD analysis of genetic diversity among *Zingiber officinale* cultivars. *Journal of Agricultural Biotechnology* 2: 134–138.

Zhou J, Tang F, Mao G and Bian R (2004) Effect of alpha-pinene on nuclear translocation of NF-B in THP-1 cells. *Acta Pharmacologica Sinica* 25: 480–484.

https://doi.org/10.1017/S1479262111000670 Published online by Cambridge University Press