THE CANADIAN ENTOMOLOGIST

Index to Volume 135 (2003)



THE ENTOMOLOGICAL SOCIETY OF CANADA, OTTAWA

https://doi.org/10.4039/ent135index Published online by Cambridge University Press

Alyokhin A. See Boiteau *et al*., 1
Aotsuka T. See Chen and Aotsuka, 143
Archibald SB. See Engel and Archibald, 63
See Makarkin *et al*., 637
Argauer RA. See Shapiro and Argauer, 85
Ashfaq M. See Erlandson *et al*., 71
Baguette M. See Schtickzelle *et al*., 313
Baldwin D. See Erlandson *et al*., 71
Barbagallo S, Cocuzza GE. Morphological discrimination of six species of the genus *Anuraphis* (Hemiptera: Aphididae), including description of a new species, 839
Baute T. See Hunt *et al*., 879
Behringer AM. See Vernon *et al*., 379
Bentz BJ. See Hansen and Bentz, 697
Bergh JC, Leskey TC. Biology, ecology, and management of dogwood borer in eastern apple orchards, 615
Bjørnson S, Raworth DA. Effects of plant nutrition on the expression of abdominal discoloration in *Phytoseiulus persimilis* (Acari: Phytoseiidae), 129
Boiteau G, Alyokhin A, Ferro DN. The Colorado potato beetle in movement, 1
Boivin G, Nénon J-P. Variability and inheritance of the chorionic tubercles on eggs of *Listronotus oregonensis* (Coleoptera: Curculionidae), 765
Borden JH. See Wardle and Borden, 733
Boudreault C. See Mason *et al*., 405
Bourchier RS. Receptor characterization of nontarget butterflies for risk assessment of biological control with the egg parasitoid *Trichogramma minutum* (Hymenoptera: Trichogrammatidae), 449
See Lyons *et al*., 685
Branson DH. Reproduction and survival in *Melanoplus sanguinipes* (Orthoptera: Acrididae) in response to resource availability and population density: the role of exploitative competition, 415
Braun L. See Erlandson *et al*., 71
Braun MP. See Olfert *et al*., 303
Broadbent AB, Rhainds M, Shipp L, Murphy G, Wainman L. Pupation behaviour of western flower thrips (Thysanoptera: Thripidae) on potted chrysanthemum, 741
Buddle CM, Hammond HEJ. Comparison of ground bettles (Coleoptera: Carabidae) and spiders (Araneae) collected in pan and pitall traps, 609
Chandler BW. See Prokopy *et al*., 909
Charlet LD. See Foster *et al*., 713
Chen H-w, Aotsuka T. Survey of the genus *Leucophenga* (Diptera, Drosophilidae) from Iriomote-jima, Japan, with descriptions of three new species, 143
Chen J-x. See Wang *et al*., 823
Chen X-S. Key to genera of the tribe Tropidocephalini (Hemiptera: Fulgoroidea: Delphacidae) from the People's Republic of China, with description of a new genus, 811
Christiansen K. See Wang *et al*., 823
Cocuzza GE. See Barbagallo and Cocuzza, 839
Cook J. See Peck and Cook, 775
Currie RW. See Gatien and Currie, 749
de Groot P, Poland TM. Attraction of *Hylastes opacus* (Coleoptera: Scolytidae) to nonanal, 309
Dixon PL. See Hillier *et al*., 139
Doane JF. See Olfert *et al*., 303
Douglas H. Revision of *Cardiophorus* (Coleoptera: Elateridae) species of eastern Canada and United States of America, 493
Duchateau MJ. See Kwon *et al*., 893
Eberhard WG. Sexual behaviour and morphology of *Themira minor* (Diptera: Sepsidae) males and the evolution of male sternal lobes and genitalic surstyli, 569
Eigenbrode SD. See Rutledge and Eigenbrode, 93
Elliott B. See LeSage and Elliott, 587
Engel MS, Archibald SB. An Early Eocene bee (Hymenoptera: Halictidae) from Quilchena, British Columbia, 63
Erlandson M, Braun L, Baldwin D, Soroka J, Ashfaq M, Hegedus D. Molecular markers for *Peristenus* spp. (Hymenoptera: Braconidae) parasitoids associated with *Lygus* spp. (Hemiptera: Miridae), 71
Eveleigh ES, Pollock SJ, Goulet H, Huber J. Identification of cocoons of *Apanteles* and *Dolichogenidea* (Hymenoptera: Braconidae) species attacking *Choristoneura fumiferana* (Lepidoptera: Tortricidae) and associated Microlepidoptera, 677
Ferro DN. See Boiteau *et al*., 1
Filotas MJ, Hajek AE, Humber RA. Prevalence and biology of *Furia gastropachae* (Zygomycetes: Entomophthorales) in populations of forest tent caterpillar (Lepidoptera: Lasiocampidae), 359

https://doi.org/10.4039/ent135index Published online by Cambridge University Press

Finn EM. See Liburd *et al*., 427
Floate KD. Field trials of *Trichomalopsis sarcophagae* (Hymenoptera: Pteromalidae) in cattle feedlots: a potential biocontrol agent of filth flies (Diptera: Muscidae), 599
Foottit R. See Hunt *et al.*, 879
Foster SP, Morris BD, Charlet LD, Gross T, Grugel S. Identification of the sex pheromone of *Cochylis arthuri* (Lepidoptera: Cochylidae), 713
Gagnier D. See Hunt *et al*., 879
Galloway TD. See Mironov and Galloway, 655
Gatien P, Currie RW. Timing of acaracide treatments for control of low-level populations of *Varroa destructor* (Acari: Varroidae) and implications for colony performance of honey bees, 749
Gillette NE. See Sun *et al*., 721
Goulet H. See Eveleigh *et al*., 677
Gross T. See Foster *et al*., 713
Grossrieder M. See Mason *et al*., 405
Grugel S. See Foster *et al*., 713
Hajek AE. See Filotas *et al*., 359
Hammond HEJ. See Buddle and Hammond, 609
Hansen EM, Bentz BJ. Comparison of reproductive capacity among univoltine, semivoltine, and re-emerged parent spruce beetles (Coleoptera: Scolytidae), 697
Hegedus D. See Erlandson *et al*., 71
Helson BV. See Lyons *et al*., 685
Hillier NK, Dixon PL, Larson DJ. Efficacy of five pheromone trap designs in trapping male *Grapholita libertina* (Lepidoptera: Tortricidae), 139
Huber JT, Moreau G. New synonymy and new sawfly host records for *Cirrospilus vittatus* (Hymenoptera: Eulophidae) in North America, 669
See Eveleigh *et al*., 677
Humber RA. See Filotas *et al*., 359
Humble LM. See Zilahi-Balogh *et al*., 103
Hunt D, Foottit R, Gagnier D, Baute T. First Canadian records of *Aphis glycines* (Hemiptera: Aphididae), 879
Jones GC. See Lyons *et al*., 685
Kabaluk JT. See Vernon *et al*., 379
Kang L. See Sun *et al*., 721
Klimaszewski J. Aleocharinae rove beetles (Coleoptera: Staphylinidae) of the ancient Sitka spruce forest on Vancouver Island, British Columbia, Canada: new synonymy and generic considerations, 867
Klimaszewski J, Pohl G, Pelletier, G. Revision of the Nearctic *Silusa* (Coleoptera, Staphylinidae, Aleocharinae), 159
Kok LT. See Zilahi-Balogh *et al*., 103
Kuhlmann U. See Mason *et al*., 405
Kwon YJ, Saeed S, Duchateau MJ. Control of *Plodia interpunctella* (Lepidoptera: Pyralidae), a pest of *Bombus terrestris* (Hymenoptera: Apidae) colonies, 893
Lafontaine JD. See Troubridge and Lafontaine, 53
Lafontaine JD, Troubridge JT. Review of the genus *Cosmia* (Lepidoptera: Noctuidae) in North America, with description of a new species, 325
Lafontaine JD, Mikkola K. New species of *Xanthia* (Lepidoptera: Noctuidae) from North America, 549
Lamb AB. See Zilahi-Balogh *et al*., 103
Larson DJ. See Hillier *et al*., 139
Le Boulengé É. See Schtickzelle *et al*., 313
LeSage L, Elliott B. Major range extension of the lily leaf beetle (Coleoptera: Chrysomelidae), a pest of wild and cultivated Liliaceae, 587
Leskey TC. See Bergh and Leskey, 615
Li S. Notes on larval instars and adult antennae of *Neodiprion abietis* (Hymenoptera: Diprionidae), 745
Liburd OE, Finn EM, Pettit KL, Wise JC. Response of blueberry maggot fly (Diptera: Tephritidae) to imidacloprid-treated spheres and selected insecticides, 427
Lowery DT. See Smirle *et al*., 117
Lyons DB, Helson BV, Bourchier RS, Jones GC, McFarlane JW. Effects of azadirachtin-based insecticides on the egg parasitoid *Trichogramma minutum* (Hymenoptera: Trichogrammatidae), 685
MacGregor JD. See Thurston and MacGregor, 583
Makarkin VN, Archibald SB, Oswald JD. New Early Eocene brown lacewings (Neuroptera: Hemerobiidae) from western North America, 637
Mason PG, Olfert O, Sluchinski L, Weiss RM, Boudreault C, Grossrieder M, Kuhlmann U. Actual and potential distribution of an invasive canola pest, *Meligethes viridescens* (Coleoptera: Nitidulidae), in Canada, 405

https://doi.org/10.4039/ent135index Published online by Cambridge University Press

Matheson FO. See Turnock *et al*., 391
McBrien HL, Millar JG. Substrate-borne vibrational signals of the Consperse stink bug (Hemiptera: Pentatomidae), 555
McFarlane JW. See Lyons *et al*., 685
McKee J. See Pearce *et al*., 337
McKenney D. See Pearce *et al*., 337
Miao Z. See Sun *et al*., 721
Mier Durante MP, Nieto Nafría JM, Ortego J. Aphidini (Hemiptera: Aphididae) living on *Senecio* (Asteraceae), with descriptions of a new genus and three new species, 187
———Aphidini (Hemiptera: Aphididae) living on *Senecio* (Asteraceae), with descriptions of a new genus and three new species — ERRATUM, 613
Mikkola K. See Lafontaine and Mikkola, 549
Millar JG. See McBrien and Millar, 555
Mironov SV, Galloway TD. Two new feather mites (Acari: Astigmata) from the Turkey Vulture (Ciconiiformes: Cathartidae) in Canada, 655
Moreau G. See Huber and Moreau, 669
Morón M-A. Revision of the *Phyllophaga* s.s. *schizorhina* species group (Coleoptera: Melolonthidae: Melolonthinae), 213
Morón MA, Paucar-Cabrera A. Larvae and pupae of species of the genus *Macraspis* (Coleoptera: Scarabaeidae: Ruelinae: Ruelini), 467
Morris BD. See Foster *et al*., 713
Murphy G. See Broadbent *et al*., 741
Muza AJ. See Nagarkatti *et al*., 589
Nagarkatti S, Tobin PC, Saunders MC, Muza AJ. Release of native *Trichogramma minutum* to control grape berry moth, 589
Nealis VG, Turnquist R. Predicting defoliation by *Choristoneura biennis* (Lepidoptera: Tortricidae), 903
Nénon J-P. See Boivin and Nénon, 765
Nieto Nafría JM. See Meir Durante *et al*., 187, 613
Ocampo FC. A new genus and two new species of Hybosoridae (Coleoptera: Scarabaeoidea) from Costa Rica, Panama, and Colombia, 41
Olfert O, Doane JF, Braun MP. Establishment of *Platygaster tuberosula*, an introduced parasitoid of the wheat midge, *Sitodiplosis mosellana*, 303
See Mason *et al*., 405
Ortego J. See Meir Durante *et al*., 187, 613
Oswald JD. See Makarkin *et al*., 637
Ovruski SM, Schliserman P. Opiine and Eucoiline (Hymenoptera: Braconidae and Figitidae, respectively) parasitoids of *Anastrepha fraterculus* (Diptera: Tephritidae) in citrus orchard areas in Corrientes, northeastern Argentina, 863
Owen DR. See Sun *et al*., 721
Paucar-Cabrera A. See Moron and Paucar-Cabrera, 467
Pearce JL, Venier LA, McKee J, Pedlar J, McKenney D. Influence of habitat and microhabitat on carabid (Coleoptera: Carabidae) assemblages in four stand types, 337
Peck SB, Cook J. Neotropical species of *Dietta* (Coleoptera: Leiodidae: Leiodinae: Estadiini), 775
Pedlar J. See Pearce *et al*., 337
Pelletier G, Piché C. Species of *Telenomus* (Hymenoptera: Scelionidae) associated with the hemlock looper (Lepidoptera: Geometridae) in Canada, 23
See Klimaszewski *et al*., 159
Pettit KL. See Liburd *et al*., 427
Piché C. See Pelletier and Piché, 23
Pohl G. See Klimaszewski *et al*., 159
Poland TM. See de Groot and Poland, 309
Pollock SJ. See Eveleigh *et al*., 677
Prokopy RJ, Chandler BW, Wright SE. Improved sugar delivery onto pesticide-treated spheres for controlling *Rhagoletis pomonella* (Diptera: Tephritidae), 909
Raworth DA. See Bjørnson and Raworth, 129
Rhainds M. See Broadbent *et al*., 741
Roden DB. Influence of burlap-band colour on larval, pupal, and egg-mass counts of *Lymantria dispar* (Lepidoptera: Lymantriidae), 869
Roland J. See Schmidt and Roland, 439
Rutledge CE, Eigenbrode SD. Epicuticular wax on pea plants decreases instantaneous search rate of *Hippodamia convergens* larvae and reduces attachment to leaf surfaces, 93
Saeed S. See Kwon *et al*., 893
Salom SM. See Zilahi-Balogh *et al*., 103

https://doi.org/10.4039/ent135index Published online by Cambridge University Press

Saunders MC. See Nagarkatti *et al*., 589
Schliserman P. See Ovruski and Schliserman, 863
Schmidt BC, Roland J. Developing techniques for monitoring forest tent caterpillar populations using synthetic pheromones, 439
Schtickzelle N, Baguette M, Le Boulengé É. Modelling insect demography from capture–recapture data: comparison between the constrained linear models and the Jolly–Seber analytical method, 313
Shapiro M, Argauer RA. Stilbenes as enhancers for gypsy moth (Lepidoptera: Lymantriidae) nucleopolyhedrovirus, 85
Shipp L. See Broadbent *et al*., 741
Sluchinski L. See Mason *et al*., 405
Smirle MJ, Lowery DT, Zurowski CL. Variation in response to insecticides in two species of univoltine leafrollers (Lepidoptera: Tortricidae), 117
Soroka J. See Erlandson *et al*., 71
Stein JD. See Sun *et al*., 721
Stephen FM. See VanLaerhoven and Stephen, 737
Sun J, Gillette NE, Miao Z, Kang L, Zhang Z, Owen DR, Stein JD. Verbenone interrupts attraction to host volatiles and reduces attack on *Pinus tabuliformis* (Pinaceae) by *Dendroctonus valens* (Coleoptera: Scolytidae) in the People's Republic of China, 721
Thurston GS, MacGregor JD. Body size – realized fecundity relationship of white-marked tussock moth, 583
Tobin PC. See Nagarkatti *et al*., 589
Troubridge JT, Lafontaine JD. A review of the pine-feeding *Lithophane lepida* species group (Lepidoptera: Noctuidae), with descriptions of two new species, 53
See Lafontaine and Troubridge, 325
Turnock WJ, Wise IL, Matheson FO. Abundance of some native coccinellines (Coleoptera: Coccinellidae) before and after the appearance of *Coccinella septempunctata*, 391
Turnquist R. See Nealis and Turnquist, 903
VanLaerhoven SL, Stephen FM. Host species influences body size and egg load of the bark beetle parasitoid *Roptrocerus xylophagorum* (Hymenoptera: Pteromalidae), 737
Venier LA. See Pearce *et al*., 337
Vernon RS, Kabaluk JT, Behringer AM. Aggregation of *Agriotes obscurus* (Coleoptera: Elateridae) at cereal bait stations in the field, 379
Wainman L. See Broadbent *et al*., 741
Wang F, Chen J-x, Christiansen K. Taxonomy of the genus *Lepidocyrtus* s.l. (Collembola: Entomobryidae) in East and Southeast Asia and Malaysia, with description of a new species from the People's Republic of China, 823
Wardle AR, Borden JH. Sexual attraction among *Lygus* (Hemiptera: Miridae) species, 733
Weiss RM. See Mason *et al*., 405
Whittington R, Winston ML. Are bumble bee colonies in tomato greenhouses obtaining adequate nutrition?, 883
Winston ML. See Whittington and Winston, 883
Wise JC. Liburd *et al*., 427
Wise IL. See Turnock *et al*., 391
Wright SE. See Prokopy *et al.*, 909
Zhang Z. See Sun *et al*., 721
Zilahi-Balogh GMG, Humble LM, Lamb AB, Salom SM, Kok LT. Seasonal abundance and synchrony between *Laricobius nigrinus* (Coleoptera: Derodontidae) and its prey, the hemlock woolly adelgid (Hemiptera: Adelgidae), 103
Zurowski CL. See Smirle *et al*., 117

https://doi.org/10.4039/ent135index Published online by Cambridge University Press