Econometric Theory, **24**, 2008, 1628–1662. Printed in the United States of America.
doi:10.1017/S026646660808064X

ADAPTIVE DENSITY ESTIMATION FOR GENERAL ARCH MODELS

F. COMTE
Université Paris Descartes

J. DEDECKER
Université Paris 6

M.L. TAUPIN
IUT de Paris 5 and Université Paris Sud

We consider a model $Y_t = \sigma_t \eta_t$ in which (σ_t) is not independent of the noise process (η_t) but σ_t is independent of η_t for each t. We assume that (σ_t) is stationary, and we propose an adaptive estimator of the density of $\ln(\sigma_t^2)$ based on the observations Y_t. Under a new dependence structure, the τ-dependency defined by Dedecker and Prieur (2005, *Probability Theory and Related Fields* 132, 203–236), we prove that the rates of this nonparametric estimator coincide with the rates obtained in the independent and identically distributed (i.i.d.) case when (σ_t) and (η_t) are independent. The results apply to various linear and nonlinear general autoregressive conditionally heteroskedastic (ARCH) processes. They are illustrated by simulations applying the deconvolution algorithm of Comte, Rozenholc, and Taupin (2006, *Canadian Journal of Statistics* 34, 431–452) to a new noise density.

1. INTRODUCTION

In this paper, we consider the following general autoregressive conditionally heteroskedastic–type (ARCH-type) model: $((Y_t, \sigma_t))_{t\in\mathbb{N}}$ is a strictly stationary sequence of $\mathbb{R} \times \mathbb{R}^+$-valued random variables, satisfying the equation

$$Y_t = \sigma_t \eta_t, \tag{1.1}$$

where $(\eta_t)_{t\in\mathbb{Z}}$ is a sequence of independent and identically distributed (i.i.d.) random variables with mean zero and finite variance and for each $t \geq 0$, the random vector $(\sigma_i, \eta_{i-1})_{0\leq i\leq t}$ is independent of the sequence $(\eta_i)_{i\geq t}$.

Such models are classically encountered in finance, when $(\sigma_t^2)_{t\in\mathbb{Z}}$, the volatility process of interest, is unobserved. The only available data are the

We thank Yves Rozenholc for his precious help in adding a new type of noise in the deconvolution Matlab programs. We are also grateful to the associate editor and the referees who helped us to improve the paper. Their questions and remarks were most helpful. Address correspondence to F. Comte, Université Paris Descartes, MAP5, UMR CNRS 8145, 45, rue des Saints-Pères, 75006 Paris, France; e-mail: fabienne.comte@univ-paris5.fr.

 © 2008 Cambridge University Press 0266-4666/08 $15.00

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

demeaned or detrended log-return process (Y_t) of an asset. A large variety of parametric models have been proposed since the first ARCH(1) model of Engle (1982), such as the GARCH(p,q) models of Bollerslev (1986), and other extensions to be found in Duan (1997). In general, it is not possible to compute, even in those parametric cases, the stationary density of σ_t. Here, we want to use flexible and powerful nonparametric tools to obtain information on the properties of the hidden process. More precisely, we shall build an estimator of the density of $\ln(\sigma_t^2)$ based on the data $(Y_t)_{1\le t\le n}$. This allows us to look for its possible bimodality properties or to determine precisely the localization of the peaks. Such a representation is of importance for modelization purposes.

Model (1.1) is classically rewritten via a logarithmic transformation:

$$Z_t = X_t + \varepsilon_t, \tag{1.2}$$

where $Z_t = \ln(Y_t^2)$, $X_t = \ln(\sigma_t^2)$, and $\varepsilon_t = \ln(\eta_t^2)$. In the context derived from the model (1.1), X_t and ε_t are independent for a given t, whereas the processes $(X_t)_{t\ge 0}$ and $(\varepsilon_t)_{t\in\mathbb{Z}}$ are not independent.

Our aim is the adaptive estimation of g, the common distribution of the unobserved variables $X_t = \ln(\sigma_t^2)$, when the density f_ε of $\varepsilon_t = \ln(\eta_t^2)$ is known. More precisely we shall build an estimator of g without any prior knowledge on its smoothness, using the observations $Z_t = \ln(Y_t^2)$ and the knowledge of the convolution kernel f_ε. Because X_t and ε_t are independent for each t, the common density f_Z of the Z_t's is given by the convolution equation $f_Z = g * f_\varepsilon$, which justifies the term *deconvolution* for the estimation of g.

It is often assumed that the noise process (η_t) is Gaussian (e.g., van Es, Spreij, and van Zanten, 2005), but general distributions can be considered. This may be of interest if heavier or thinner tails are suspected to be relevant. Nevertheless, for identifiability of the statistical problem, the density of ε_t is required to be known. This assumption cannot be easily removed: even if the density of ε_t is completely known up to a scale parameter, model (1.2) may be nonidentifiable as soon as the unknown density g of X_t is smoother than the density of ε_t (see Butucea and Matias, 2005, Sect. 1). For instance, the model in which g is an unknown normal distribution and ε_t is a Gaussian random variable with unknown variance is nonidentifiable.

In density deconvolution of i.i.d. variables the X_t's and the ε_t's are i.i.d., and the sequences $(X_t)_{t\ge 0}$ and $(\varepsilon_t)_{t\in\mathbb{Z}}$ are independent (for brevity we shall refer to this case as the i.i.d. case). In the i.i.d. case, the slowest rates of convergence for estimating g are obtained for most regular error densities. For instance, when ε_t is Gaussian or the log of a squared Gaussian and g belongs to some Sobolev class, the minimax rates are negative powers of $\ln(n)$ (see Fan, 1991). Nevertheless, it has been noticed by several authors (see Pensky and Vidakovic, 1999; Butucea, 2004; Butucea and Tsybakov, 2008a, 2008b; Comte, Rozenholc, and Taupin, 2006) that the rates are improved if g has stronger smoothness properties.

In the setting of model (1.2), the classical assumptions of independence between the processes $(X_t)_{t\ge 0}$ and $(\varepsilon_t)_{t\in\mathbb{Z}}$ are no longer satisfied, and the tools

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

for deconvolution have to be revisited. Our estimator of g is constructed by minimizing an appropriate penalized contrast function depending only on the observations and on f_ε. It is chosen in a purely data-driven way among a collection of nonadaptive estimators. We start by studying those nonadaptive estimators and show that their mean integrated squared error (MISE) has the same order as in the i.i.d. case. Next we prove that the MISE of our adaptive estimator is of the same order as the MISE of the best nonadaptive estimator in the collection, up to some possible negligible logarithmic loss in one case.

In their 2005 paper, van Es et al. (2005) have considered the case where η_t is Gaussian, the density g of X_t is twice differentiable, and the process (Z_t, X_t) is α-mixing. Here we show that, if g happens to be more regular (e.g., if g is a Gaussian density, or the log of a squared Gaussian exactly like the noise process ε_t), then their procedure is suboptimal, and the bandwidth they propose is not the best one. This is the reason why we do not make any assumption on the smoothness of g: this is the advantage of the adaptive procedure.

We also consider a new type of dependence property, which is satisfied by many ARCH processes. We can prove all results under the classical β-mixing assumption, satisfied by general ARCH models, as recalled in Doukhan (1994) and described in more detail in Carrasco and Chen (2002). But we choose to illustrate that new recent coefficients can be used in our context, which allow an easy characterization of the dependence properties in function of the parameters of the models. Those new dependence coefficients, recently defined and studied in Dedecker and Prieur (2005), are interesting and powerful because they require much lighter conditions on the models. Such ideas have been popularized by Ango Nzé and Doukhan (2004) and Doukhan, Teyssière, and Winant (2006). For instance, these coefficients allow us to deal with the general ARCH(∞) processes defined by Giraitis, Kokoszka, and Leipus (2000).

To be complete, we also give a small simulation study when ε_t is the log of the square of a standard Gaussian. We check the effect of misspecification of the error distribution, and we show that the Comte, Rozenholc, and Taupin (2007) algorithm applies to ARCH processes.

The paper is organized as follows. The estimator is defined in Section 2. The MISE bounds are given in Section 3 under some dependence properties. Examples and simulation results are described in Section 4. All the proofs are given in Section 5.

2. THE ESTIMATORS

For two complex-valued functions u and v in $\mathbb{L}_2(\mathbb{R}) \cap \mathbb{L}_1(\mathbb{R})$, let $u^*(x) = \int e^{itx} u(t)\,dt$, $u * v(x) = \int u(y)v(x-y)\,dy$, and $\langle u,v\rangle = \int u(x)\bar{v}(x)\,dx$ with $\bar{z}$ the conjugate of a complex number z. We also denote by $\|u\|_1 = \int |u(x)|\,dx$, $\|u\|^2 = \int |u(x)|^2\,dx$, and $\|u\|_\infty = \sup_{x\in\mathbb{R}} |u(x)|$.

In what follows, we consider model (1.1) and its equivalent representation (1.2), and we describe the estimation strategy for g, the density of the X_i's based

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

on observations Z_i, $i = 1, \ldots, n$ and on the knowledge of f_ε, the density of the ε_i's.

2.1. Construction of the Minimum Contrast Estimators

First, the heuristics for the construction of the estimators are the following. Mean-square-type contrasts are determined by following the idea that we look for a function t such that its $\mathbb{L}_2$-distance to the function g is minimum. Because $\|t - g\|^2 = \|t\|^2 - 2\langle t, g\rangle + \|g\|^2$, this amounts to minimizing $\|t\|^2 - 2\langle t, g\rangle$ in a well-chosen class of functions t. With Parseval identity, $\langle t, g\rangle = \langle t^*, g^*\rangle/2\pi$. Now, if f_Z is the common density of the Z_i's, then $f_Z = f_\varepsilon * g$, so that $f_Z^* = f_\varepsilon^* g^*$ and $g^* = f_Z^*/f_\varepsilon^*$, where f_ε^* is assumed to be known. Moreover $f_Z^*(x) = \mathbb{E}(e^{ixZ})$ can be replaced by its empirical counterpart $(1/n)\sum_{k=1}^n e^{ixZ_k}$. We can now define the contrast function $\|t\|^2 - (\pi n)^{-1}\sum_{k=1}^n \int e^{ixZ_k} t^*(-x)/f_\varepsilon^*(x)\, dx$, which is denoted by

$$\gamma_n(t) = \frac{1}{n}\sum_{k=1}^{n}[\|t\|^2 - 2u_t^*(Z_k)], \quad \text{with } u_t(x) = \frac{1}{2\pi}\left(\frac{t^*(-x)}{f_\varepsilon^*(x)}\right),$$

under the assumption that

$$f_\varepsilon \quad \text{belongs to } \mathbb{L}_2(\mathbb{R}) \text{ and is such that } f_\varepsilon^*(x) \neq 0 \quad \text{for any } x \text{ in } \mathbb{R}. \tag{2.1}$$

We have $\mathbb{E}[u_t^*(Z_i)] = (2\pi)^{-1}\langle f_Z^*/f_\varepsilon^*, t^*\rangle = \langle t, g\rangle$, where the last equality follows from the Parseval identity. It follows that $\mathbb{E}(\gamma_n(t)) = \|t - g\|^2 - \|g\|^2$ is minimal when $t = g$.

Clearly the functions t must be chosen such that $u_t^*(x)$ is well defined. Because $1/f_\varepsilon^*$ can be nonintegrable (think of a Gaussian density), a solution is to choose the functions t such that t^* exists and has compact support.

The simplest spaces that suit that aim are often studied in preliminary wavelet courses (see, e.g., Meyer, 1990, p. 22) and are the following. Let $\varphi(x) = \sin(\pi x)/(\pi x)$. The Fourier transform of φ is obtained by noticing that the Fourier transform of $\mathbb{1}_{[-\pi,\pi]}(x)/2\pi$ is equal to φ and by using the inverse Fourier formula, and thus $\varphi^*(x) = \mathbb{1}_{[-\pi,\pi]}(x)$. For $m \in \mathbb{N}$ and $j \in \mathbb{Z}$, set $\varphi_{m,j}(x) = \sqrt{m}\varphi(mx - j)$. The functions $\{\varphi_{m,j}\}_{j\in\mathbb{Z}}$ constitute an orthonormal system in $\mathbb{L}^2(\mathbb{R})$: indeed $\varphi_{m,j}^*(x) = e^{ijx/m}\varphi^*(x/m)/\sqrt{m}$ and $\langle \varphi_{m,j}, \varphi_{m,k}\rangle = (2\pi)^{-1}\langle \varphi_{m,j}^*, \varphi_{m,k}^*\rangle = \delta_{j,k}$. Therefore, if we define

$$S_m = \overline{\text{span}}\{\varphi_{m,j}, j \in \mathbb{Z}\}, \qquad m \in \mathbb{N},$$

the space S_m is exactly the subspace of $\mathbb{L}_2(\mathbb{R})$ of functions having a Fourier transform with compact support contained in $[-\pi m, \pi m]$. The orthogonal projection of g on S_m is $g_m = \sum_{j\in\mathbb{Z}} a_{m,j}(g)\varphi_{m,j}$ where $a_{m,j}(g) = \langle \varphi_{m,j}, g\rangle$. Now, we cannot describe practical algorithms involving infinite representations. Thus, to obtain representations having a finite number of "coordinates," we introduce

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

$$S_m^{(n)} = \overline{\text{span}}\{\varphi_{m,j}, |j| \leq n\}.$$

The family $\{\varphi_{m,j}\}_{|j|\leq n}$ is an orthonormal basis of $S_m^{(n)}$, and the orthogonal projection of g on $S_m^{(n)}$ is given by $g_m^{(n)} = \sum_{|j|\leq n} a_{m,j}(g)\varphi_{m,j}$. Subsequently a space $S_m^{(n)}$ will be referred to as a "model" and also as a "projection space."

Then, for an arbitrary fixed integer m, an estimator of g belonging to $S_m^{(n)}$ is defined by

$$\hat{g}_m^{(n)} = \arg \min_{t \in S_m^{(n)}} \gamma_n(t). \tag{2.2}$$

Moreover, it is easy to see that

$$\hat{g}_m^{(n)} = \sum_{|j|\leq n} \hat{a}_{m,j}\varphi_{m,j} \quad \text{with } \hat{a}_{m,j} = \frac{1}{n}\sum_{i=1}^{n} u^*_{\varphi_{m,j}}(Z_i), \quad \text{and}$$

$$\mathbb{E}(\hat{a}_{m,j}) = \langle g, \varphi_{m,j}\rangle = a_{m,j}(g).$$

2.2. Minimum Penalized Contrast Estimator

For the sake of completeness in the description of our estimation strategy, let us also give the last step. We show in what follows that a "good" choice of m exists but that it results from a squared bias-variance compromise in which the bias is unknown. The end of the procedure is the analogue of a bandwidth selection strategy in kernel estimation and gives a criterion for selecting m.

The minimum penalized estimator of g is defined as $\tilde{g} = \hat{g}_{\hat{m}_g}^{(n)}$ where $\hat{m}_g$ is chosen in a purely data-driven way. The main point of the estimation procedure lies in the choice of $m = \hat{m}$ (or equivalently in the choice of a model $S_{\hat{m}}^{(n)}$) involved in the estimators $\hat{g}_m^{(n)}$ given by (2.2), to mimic the oracle parameter

$$\breve{m}_g = \arg \min_m \mathbb{E}\|\hat{g}_m^{(n)} - g\|^2. \tag{2.3}$$

The model selection is performed in an automatic way, using the following penalized criteria:

$$\tilde{g} = \hat{g}_{\hat{m}}^{(n)} \quad \text{with } \hat{m} = \arg \min_{m\in\{1,\ldots,m_n\}} [\gamma_n(\hat{g}_m^{(n)}) + \text{pen}(m)], \tag{2.4}$$

where $\text{pen}(m)$ is a penalty function that depends on $f_\varepsilon^*(\cdot)$ through $\Delta(m)$ defined by

$$\Delta(m) = \frac{1}{2\pi}\int_{-\pi m}^{\pi m} \frac{1}{|f_\varepsilon^*(x)|^2}\, dx. \tag{2.5}$$

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

The key point in the dependent context is to find a penalty function not depending on the dependence coefficients such that

$$\mathbb{E}\|\tilde{g} - g\|^2 \le C \inf_{m\in\{1,\dots,m_n\}} \mathbb{E}\|\hat{g}_m^{(n)} - g\|^2.$$

In that way, the estimator $\tilde{g}$ is adaptive because it achieves the best rate among the estimators $\hat{g}_m^{(n)}$, without any prior knowledge on the smoothness on g.

3. DENSITY ESTIMATION BOUNDS

3.1. Mixing Assumptions

Clearly, the process of interest after the logarithmic transformation ($Z_t = \ln(Y_t^2)$, $X_t = \ln(\sigma_t^2)$) is

$$(W_t)_{t\in\mathbb{Z}} = ((Z_t, X_t))_{t\in\mathbb{Z}} \tag{3.1}$$

and involves dependent variables. We omit the presentation of the classical β-mixing properties and describe instead a new τ-dependence notion that reveals its usefulness for a large class of models.

Let $(\Omega, \mathcal{A}, \mathbb{P})$ be a probability space. Let W be a random vector with values in a Banach space $(\mathbb{B}, \|\cdot\|_{\mathbb{B}})$ and let $\mathcal{M}$ be a σ-algebra of $\mathcal{A}$. Let $\mathbb{P}_{W|\mathcal{M}}$ be a conditional distribution of W given $\mathcal{M}$ and let $\mathbb{P}_W$ be the distribution of W. Let $\Lambda_1(\mathbb{B})$ be the set of 1-Lipschitz functions from $(\mathbb{B}, \|\cdot\|_{\mathbb{B}})$ to $\mathbb{R}$. If $\mathbb{E}(\|W\|_{\mathbb{B}}) < \infty$, define

$$\tau(\mathcal{M}, W) = \mathbb{E}\left(\sup_{f\in\Lambda_1(\mathbb{B})} |\mathbb{P}_{W|\mathcal{M}}(f) - \mathbb{P}_W(f)| \right),$$

as in Dedecker and Prieur (2005).

Let $(W_t)_{t\ge0}$ be a strictly stationary sequence of $\mathbb{R}^2$-valued random variables. On $\mathbb{R}^2$, we put the norm $\|x - y\|_{\mathbb{R}^2} = |x_1 - y_1| + |x_2 - y_2|$. For any $k \ge 0$, define the coefficients

$$\text{if} \quad \mathbb{E}(\|W_0\|_{\mathbb{R}^2}) < \infty, \qquad \tau_1(k) = \tau(\sigma(W_0), W_k). \tag{3.2}$$

On $(\mathbb{R}^2)^l$, we put the norm $\|x - y\|_{(\mathbb{R}^2)^l} = l^{-1}(\|x_1 - y_1\|_{\mathbb{R}^2} + \cdots + \|x_l - y_l\|_{\mathbb{R}^2})$. Let $\mathcal{M}_i = \sigma(W_k, 0 \le k \le i)$. If $\mathbb{E}(\|W_1\|_{\mathbb{R}^2}) < \infty$, the coefficients $\tau_\infty(k)$ are defined by

$$\tau_\infty(k) = \sup_{i\ge0} \sup_{l\ge1} \{\tau(\mathcal{M}_i, (W_{i_1}, \dots, W_{i_l})), i + k \le i_1 < \cdots < i_l\}. \tag{3.3}$$

We say that the process $(W_t)_{t\ge0}$ is τ-dependent if the coefficients $\tau_\infty(k)$ tend to zero as k tends to infinity. We say that it is geometrically τ-dependent if there exist $a < 1$ and $C > 0$ such that $\tau_\infty(k) \le Ca^k$ for all $k \ge 1$.

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

From now on, the dependence coefficients are defined as in (3.2) and (3.3) with $(W_t)_{t\in\mathbb{Z}} = ((Z_t, X_t))_{t\in\mathbb{Z}}$. Moreover, we summarize the dependency assumptions for model (1.2) as follows:

The ε_i's are i.i.d.

The random vector $(X_i, \varepsilon_{i-1})_{0\le i\le t}$ is independent of the sequence $(\varepsilon_i)_{i\ge t}$.

The process $(W_t)_{t\in\mathbb{Z}}$ is strictly stationary and τ-dependent.

(3.4)

3.2. Risk Bound of the Minimum Contrast Estimators $\hat{g}_m^{(n)}$

Subsequently, the density g is assumed to satisfy the following assumption:

$$g \in \mathbb{L}_2(\mathbb{R}), \quad \text{and there exists } M_2 > 0, \qquad \int x^2 g^2(x)\,dx \le M_2 < \infty. \tag{3.5}$$

For instance, (3.5) is fulfilled if g is bounded by M_0 and $\mathbb{E}(X_1^2) \le M_1 < +\infty$, with $M_2 = M_0 M_1$. Assumption (3.5) is due to the construction of the estimator on $S_m^{(n)}$ instead of S_m and is not very restrictive.

The order of the MISE of $\hat{g}_m^{(n)}$ is given in the following proposition.

PROPOSITION 3.1. *If (2.1) and (3.5) hold, then $\hat{g}_m^{(n)}$ defined by (2.2) satisfies*

$$\mathbb{E}\|g - \hat{g}_m^{(n)}\|^2 \le \|g - g_m\|^2 + \frac{m^2(M_2+1)}{n} + \frac{2\Delta(m)}{n} + \frac{2R_m}{n},$$

where $\Delta(m)$ is defined by (2.5) and

$$R_m = \frac{1}{\pi}\sum_{k=2}^{n}\int_{-\pi m}^{\pi m}\left|\frac{\mathrm{Cov}(e^{ixZ_1}, e^{ixX_k})}{f_\varepsilon^*(-x)}\right| dx. \tag{3.6}$$

Moreover, $R_m \le R_{m,\tau}$, where $R_{m,\tau} = \pi m \Delta_{1/2}(m)\sum_{k=1}^{n-1}\tau_1(k)$, with τ_1 defined by (3.2), and where

$$\Delta_{1/2}(m) = \frac{1}{2\pi}\int_{-\pi m}^{\pi m}\frac{1}{|f_\varepsilon^*(x)|}\,dx. \tag{3.7}$$

This proposition requires several comments.

The order of the risk is given by a bias term $\|g_m - g\|^2 + m^2(M_2+1)/n$ and a variance term $2\Delta(m)/n + 2R_m/n$.

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

The main part of the bias term is $\|g - g_m\|^2$. It is noteworthy that

$$\|g - g_m\|^2 = \int_{|x| \geq \pi m} |g^*(x)|^2 \, dx, \tag{3.8}$$

so that the order of the bias depends on the rate of decay of $g^*(x)$. The additional term $m^2(M_2 + 1)/n$ is negligible with respect to the variance term.

The variance term $2\Delta(m)/n + 2R_m/n$ depends on the rate of decay of the Fourier transform of f_ε. It is the sum of the variance term appearing in density deconvolution for i.i.d. variables, $2\Delta(m)/n$, and of an additional term, $2R_m/n$. This last term R_m involves the dependency coefficients and the quantity $\Delta_{1/2}(m)$, which is specific to the ARCH problem. The point is that the main order term in the variance part is $\Delta(m)/n$, which does not involve the dependency coefficients. In other words, the dependency coefficients only appear in front of the additional and negligible term, specific to ARCH models, $\Delta_{1/2}(m)/n$.

3.3. Discussion about the Rates

Now, we need to parameterize the rate of decay of f_ε^* and g^* to evaluate the order of the variance term $\Delta(m)/n$ and of the main part of the bias term given by (3.8).

More precisely, we assume that f_ε is such that there exist nonnegative numbers κ_0, γ, μ, and δ such that the Fourier transform f_ε^* of f_ε satisfies

$$\kappa_0(x^2 + 1)^{-\gamma/2} \exp\{-\mu|x|^\delta\} \leq |f_\varepsilon^*(x)| \leq \kappa_0'(x^2 + 1)^{-\gamma/2} \exp\{-\mu|x|^\delta\}. \tag{3.9}$$

Because f_ε is known, the constants μ, δ, κ_0, and γ defined in (3.9) are known. The class described by (3.9) is very general. By convention, we set $\delta = 0$ if $\mu = 0$. When $\delta = 0$ in (3.9), the errors are called "ordinary smooth" errors. When $\mu > 0$ and $\delta > 0$, they are called "super smooth." An example of ordinary smooth density is the Laplace distribution, for which $f_\varepsilon^*(x) = 1/(1 + x^2)$ so that $\delta = \mu = 0$ and $\gamma = 2$. The standard examples for super smooth densities are Gaussian ($f_\varepsilon^*(x) = e^{-x^2/2}$ so that $\gamma = 0$, $\delta = 2$, $\mu = \frac{1}{2}$) or Cauchy ($f_\varepsilon^*(x) = e^{-|x|}$, so that $\gamma = 0$, $\delta = 1$, $\mu = 1$) distributions. When $\varepsilon_t = \ln(\eta_t^2)$ with $\eta_t \sim \mathcal{N}(0,1)$ as in van Es et al. (2005), then $f_\varepsilon^*(x) = 2^{ix}\Gamma(\frac{1}{2} + ix)/\sqrt{\pi}$ and with Stirling formula $|f_\varepsilon^*(x)| \sim_{+\infty} \sqrt{2/e}e^{-\pi|x|/2}$. Then ε_t is super smooth with $\delta = 1$, $\gamma = 0$, and $\mu = \pi/2$. Note that, in that case, $\mathbb{E}(\varepsilon_1) = -\ln(2) - \gamma$ where γ is the Euler constant and $\mathrm{Var}(\varepsilon_1) = \pi^2/2$.

Moreover, the square integrability of f_ε and (3.9) require that $\gamma > \frac{1}{2}$ when $\delta = 0$.

Now, concerning the main variance term, if f_ε^* satisfies (3.9), then $\Delta(m)$ given by (2.5) has the same order as

$$\Gamma(m) = (1 + (\pi m)^2)^\gamma (\pi m)^{1-\delta} \exp\{2\mu(\pi m)^\delta\},$$

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

up to some constant bounded by

$$\lambda_1(f_\varepsilon, \kappa_0) = \frac{1}{\kappa_0^2 \pi R(\mu, \delta)}, \quad \text{where } R(\mu, \delta) = \mathbb{1}_{\{\delta=0\}} + 2\mu\delta \mathbb{1}_{\{\delta>0\}}. \tag{3.10}$$

Also, we describe the smoothness properties of g by the set

$$\mathcal{S}_{s,r,b}(C_1) = \left\{ \psi \quad \text{such that } \int_{-\infty}^{+\infty} |\psi^*(x)|^2 (x^2+1)^s \exp\{2b|x|^r\}\, dx \le C_1 \right\} \tag{3.11}$$

for s, r, b unknown nonnegative numbers. By convention, if $b = 0$ or $r = 0$, we set both to 0. When $r = 0$, the class $\mathcal{S}_{s,r,b}(C_1)$ corresponds to a Sobolev ball. Heuristically, a function ψ admitting k continuous derivatives is such that $|\psi^*(x)| = O(|x|^{-k})$ when x tends to infinity and thus belongs to $\mathcal{S}_{s,0,0}(C_1)$ for $s < k - \frac{1}{2}$. When $r > 0$, $b > 0$ functions belonging to $\mathcal{S}_{s,r,b}(C_1)$ are infinitely many times differentiable. For instance $\psi(x) = e^{-x^2/2}$ belongs to $\mathcal{S}_{s,b,2}(C_1)$ for any $b < \frac{1}{2}$ and any $s \ge 0$.

Now, concerning the main bias term, it follows from (3.8) that if g belongs to a space $\mathcal{S}_{s,r,b}(C_1)$, then

$$\|g - g_m\|^2 \le \frac{C}{2\pi} (m^2\pi^2 + 1)^{-s} \exp\{-2b\pi^r m^r\},$$

where C is a constant depending on C_1.

As a consequence, the rates resulting from Proposition 3.1 under (3.9) and (3.11) are deduced from the following corollary.

COROLLARY 3.1. *Assume that (3.9), (2.1), and (3.5) hold and that g belongs to $\mathcal{S}_{s,r,b}(C_1)$ defined by (3.11). Assume that $\delta = 0$, $\gamma > 1$, or $\delta > 0$ in (3.9) and that $\sum_{k\ge1} \tau_1(k) < +\infty$. Then $\hat{g}_m^{(n)}$ defined by (2.2) satisfies*

$$\mathbb{E}\|g - \hat{g}_m^{(n)}\|^2 \le \frac{C_1}{2\pi} (m^2\pi^2 + 1)^{-s} \exp\{-2b\pi^r m^r\} + \frac{2\lambda_1(f_\varepsilon, \kappa_0)\Gamma(m)}{n} + \frac{C_2}{n} \Gamma(m) o_m(1), \tag{3.12}$$

where C_1 and C_2 are finite constants. The constant C_2 depends on $\sum_{k\ge1} \tau_1(k)$.

The rate of convergence of $\hat{g}_{\check{m}}^{(n)}$ is the same as the rate for density deconvolution for i.i.d. sequences. Moreover, our context encompasses the particular case considered by van Es et al. (2005). Table 1 gives a summary of these rates obtained when minimizing the right-hand side of (3.12). The $\check{m}_g$ denotes the corresponding minimizer (see eqn. (2.3)).

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

TABLE 1. Choice of $\breve{m}_g$ and corresponding rates under assumptions (3.9) and (3.11)

	f_ε	
	$\delta = 0$ Ordinary smooth	$\delta > 0$ Super smooth
g		
$r = 0$	$\pi\breve{m}_g = O(n^{1/(2s+2\gamma+1)})$	$\pi\breve{m}_g = [\ln(n)/(2\mu+1)]^{1/\delta}$
Sobolev(s)	rate $= O(n^{-2s/(2s+2\gamma+1)})$	rate $= \mathrm{O}((\ln(n))^{-2s/\delta})$
$r > 0$	$\pi\breve{m}_g = [\ln(n)/2b]^{1/r}$	$\breve{m}_g$ solution of
$\mathcal{C}^\infty$	rate $= O\left(\dfrac{\ln(n)^{(2\gamma+1)/r}}{n}\right)$	$\breve{m}_g^{2s+2\gamma+1-r}\exp\{2\mu(\pi\breve{m}_g)^\delta + 2b\pi^r\breve{m}_g^r\}$ $= O(n)$

If g belongs to a classical Sobolev class ($r = 0$), the rates range from negative powers of n when the errors are ordinary smooth to negative powers of $\ln(n)$ if the errors are super smooth. But if the function to estimate also becomes super smooth (case $r > 0$), the rates become much better in both cases. When $r > 0$, $\delta > 0$ the value of $\breve{m}_g$ is not explicitly given. It is obtained as the solution of the equation

$$\breve{m}_g^{2s+2\gamma+1-r}\exp\{2\mu(\pi\breve{m}_g)^\delta + 2b\pi^r\breve{m}_g^r\} = O(n).$$

Consequently, the rate of $\hat{g}^{(n)}_{\breve{m}_g}$ is not easy to give explicitly and depends on the ratio r/δ. We refer to Comte et al. (2006) for further discussion about those rates. We refer to Lacour (2006) for explicit formulas for the rates in the special case $r > 0$ and $\delta > 0$.

Example

Consider the case of the van Es et al. (2005) paper, where f_ε^* satisfies (3.9) with $\delta = 1$, $\mu = \pi/2$, and $\gamma = 0$. The authors consider that the regularity of g is such that g belongs to $\mathcal{S}_{s,r,b}(C_1)$ with s at most $\frac{3}{2}$ and $r = 0$. This is the case in the last column of the first row in Table 1 and leads to a logarithmic rate. Now, if, for instance, $g^*(x)$ is of order $e^{-\pi|x|/2}$ like the noise, then the squared bias-variance compromise $C_1 e^{-\pi m} + C_2 e^{\pi m}/n$ leads us to choose $\breve{m} = \ln(n)/(2\pi)$, and the rate is of order $1/\sqrt{n}$, which is better than logarithmic. If g is Gaussian, then the squared bias-variance compromise $C_1 m^{-1} e^{-\pi^2 m^2/2} + C_2 e^{\pi m}/n$ leads us to choose $\breve{m} = 2\sqrt{\ln(n)}/\pi$, and the rate is of order $e^{2\sqrt{\ln(n)}}/n = o(1/n^{1-\epsilon})$ for any $\epsilon > 0$, which is even better than the previous one. This example is another way to see the value of the adaptive procedure, which enables us to face any regularity for g without being required to know either g or its regularity.

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

3.4. Adaptive Bound

Theorem 3.1, which follows, gives a general bound that holds under weak dependence conditions for ε being either ordinary or super smooth.

For $a > 1$, let $\text{pen}(m)$ be defined by

$$\text{pen}(m) = \begin{cases} 48a\,\dfrac{\Delta(m)}{n} & \text{if } 0 \le \delta < \dfrac{1}{3}, \\ 16a\lambda_3\,\dfrac{\Delta(m)(\pi m)^{\min((3\delta/2-1/2)_+,\,\delta)}}{n} & \text{if } \delta \ge \dfrac{1}{3}, \end{cases} \tag{3.13}$$

where $\Delta(m)$ is defined by (2.5). The constant $\lambda_1(f_\varepsilon, \kappa_0)$ is defined in (3.10), and

$$\lambda_3 = 1 + \frac{98\mu}{\lambda_1(f_\varepsilon, \kappa_0')}\left((\sqrt{2}+8)\|f_\varepsilon\|_\infty \kappa_0^{-1}\sqrt{\lambda_1(f_\varepsilon,\kappa_0)}\,\mathbb{I}_{0\le\delta\le 1} + 2\lambda_1(f_\varepsilon,\kappa_0)\mathbb{I}_{\delta>1}\right). \tag{3.14}$$

The important point here is that λ_3 is known. Hence the penalty is explicit up to a numerical multiplicative constant. This procedure has already been studied by simulation experimetns for independent sequences $(X_t)_{t\ge1}$ and $(\varepsilon_t)_{t\ge1}$ in Comte et al. (2006, 2007). In particular, the practical implementation of the penalty functions and the calibration of the constants have been studied in the two previously mentioned papers. Moreover, it is shown therein that the estimation procedure is robust to various types of dependence, whether the errors ε_i's are ordinary or super smooth (see Tables 4 and 5 in Comte et al., 2007).

To bound up $\text{pen}(m)$, we impose the condition that

$$\pi m_n \le \begin{cases} n^{1/(2\gamma+1)} & \text{if } \delta = 0 \\ \left[\dfrac{\ln(n)}{2\mu} + \dfrac{2\gamma+1-\delta}{2\delta\mu}\ln\left(\dfrac{\ln(n)}{2\mu}\right)\right]^{1/\delta} & \text{if } \delta > 0. \end{cases} \tag{3.15}$$

Subsequently we set

$$C_a = \max(\kappa_a^2, 2\kappa_a) \quad \text{where } \kappa_a = (a+1)/(a-1). \tag{3.16}$$

THEOREM 3.1. *Consider model (1.2) under (3.1)–(3.4). Assume that f_ε satisfies (3.9) and (2.1), that g satisfies (3.5), and that m_n satisfies (3.15). Consider the collection of estimators $\hat{g}_m^{(n)}$ defined by (2.2) with $1 \le m \le m_n$ and* $\text{pen}(m)$ *defined by (3.13). Assume either that*

(i) $\delta = 0$, $\gamma \ge \frac{3}{2}$ in (3.9) and $\tau_\infty(k) = O(k^{-(1+\theta)})$ for some $\theta > 3 + 2/(1+2\gamma)$
(ii) or $\delta > 0$ in (3.9) and $\tau_\infty(k) = O(k^{-(1+\theta)})$ for some $\theta > 3$,

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

where τ_∞ is defined as in (3.3). Then the estimator $\tilde{g} = \hat{g}_{\hat{m}}^{(n)}$ defined by (2.4) satisfies

$$\mathbb{E}(\|g - \tilde{g}\|^2) \le C_a \inf_{m \in \{1,\dots,m_n\}} \left[\|g - g_m\|^2 + \mathrm{pen}(m) + \frac{m^2(M_2+1)}{n} \right] + \frac{\bar{C}}{n}, \tag{3.17}$$

where C_a is defined in (3.16) and $\bar{C}$ is a constant depending on f_ε, a, and $\sum_{k\ge 1} \tau_\infty(k)$.

Remark. The result of Theorem 3.1 holds under a similar (but slightly weaker) β-mixing condition.

The estimator $\tilde{g}$ is adaptive in the sense that it is purely data-driven. This is due to the fact that $\mathrm{pen}(\cdot)$ is explicitly known. In particular, its construction does not require any prior smoothness knowledge on the unknown density g and does not use the dependency coefficients. This point is important because all quantities involving dependency coefficients are usually not tractable in practice.

Moreover, one can compare the order of the penalty to the variance order $\Delta(m)/n$ in light of Table 1. They are equal for $0 \le \delta \le \frac{1}{3}$: this means that, asymptotically, the right-hand side of (3.17) realizes an automatic squared bias-variance compromise. For $\delta > \frac{1}{3}$, the order penalty has the order of the variance increased by a power of m, but this does not change the choice of $\pi \check{m}_g$ if $r = 0$, and thus the rate is unchanged then; for $r > 0$, the optimal choice of $\check{m}$ in the case $\delta > 0$ is of order $\ln(n)$, and the rate is always faster than logarithmic. Therefore, if a loss occurs, it is negligible when compared to the rate.

To summarize, the main result in Theorem 3.1 shows that the MISE of $\tilde{g}$ automatically achieves the best squared bias-variance compromise (possibly up to some logarithmic factor) among the estimators $\hat{g}_m^{(n)}$. Consequently, it achieves the best rate among the rates of the $\hat{g}_m^{(n)}$, even from a nonasymptotical point of view. This nonasymptotic feature is important because the m's selected in practice are small and far away from asymptotic. For practical illustration of this point in the case of density deconvolution of i.i.d. variables, we refer to Comte et al. (2006, 2007).

To conclude, the estimator $\tilde{g}$ has the same rate as in the i.i.d. case, with an explicit penalty function not depending on the dependence coefficients.

3.5. Further Comments

We first show how the density f of σ_t^2 can be estimated. Because for $u > 0$, $f(u) = g(\ln(u))/u$, we choose the estimator

$$\hat{f}(u) = \frac{\tilde{g}(\ln(u))}{u}, \quad \text{for } u > 0.$$

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

A change of variables gives the equality

$$\mathbb{E}\left(\int_0^\infty |\hat{f}(t)-f(t)|^2 t\,dt\right) = \mathbb{E}(\|\tilde{g}-g\|^2).$$

Note that the term on the left-hand side is a MISE for $\hat{f}$ with respect to the measure tdt. In particular, it follows that for any $a > 0$,

$$\mathbb{E}\left(\int_a^\infty |\hat{f}(t)-f(t)|^2\,dt\right) \le \frac{1}{a}\,\mathbb{E}(\|\tilde{g}-g\|^2),$$

which shows that the usual MISE for $\hat{f}$ on any interval $[a,\infty[$ tends to zero at least with the same rate as $\mathbb{E}(\|\tilde{g}-g\|^2)$.

Second, we mention that the procedure can be generalized to estimate the joint distribution of (σ_t,σ_{t+1}) or more precisely (X_t,X_{t+1}). If G denotes the bivariate density of (X_t,X_{t+1}), then the extension is conducted as follows. Let

$$V_T(x,y) = \frac{1}{4\pi^2}\iint e^{ixu+iyv}\,\frac{T^*(-u,-v)}{f_\varepsilon^*(u)f_\varepsilon^*(v)}\,dudv,$$

$$T^*(u,v) = \iint e^{iux+ivy}T(x,y)\,dxdy,$$

and

$$\Gamma_n(T) = \frac{1}{n}\sum_{k=1}^n [\|T\|^2 - 2V_T(Z_k,Z_{k+1})],$$

for a function T square integrable on $\mathbb{R}^2$. Then we would define $\hat{G}_m = \arg\min \Gamma_n(T)$ for T belonging to product spaces spanned by $(\varphi_{m,j}\otimes\varphi_{m,k})(x,y) := \varphi_{m,j}(x)\varphi_{m,k}(y)$. In other words,

$$\hat{G}_m(x,y) = \sum_{j,k}\hat{A}_{j,k}\varphi_{m,j}(x)\varphi_{m,k}(y), \qquad \hat{A}_{j,k} = \frac{1}{n}\sum_{i=1}^n V_{\varphi_{m,j}\otimes\varphi_{m,k}}(Z_i,Z_{i+1}).$$

Model selection is then performed via penalization. In the context where the process X is mixing, but globally independent of the noise process (ε_t), the procedure is studied by Lacour (2008). It is likely that the multivariate procedure would work in our context also, but the technical price sets this question beyond the scope of the present paper.

Last, it is possible to deal with dependent errors, provided that the processes $(X_t)_t$ and $(\varepsilon_t)_t$ are independent. This context is not developed because then the presence of the unknown mixing coefficients in the variance bound and then in the penalty function is unavoidable. The procedure is then not theoretically satisfactory.

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

4. EXAMPLE MIXING PROCESSES AND ILLUSTRATION OF THE METHOD

4.1. Examples of Mixing Models

In this section, we give examples of ARCH models for which the result of Theorem 3.1 holds.

A particular case of model (1.1) is

$$Y_t = \sigma_t \eta_t, \quad \text{with } \sigma_t = f(\eta_{t-1}, \eta_{t-2}, \ldots) \tag{4.1}$$

for some measurable function f. Another important case is

$$Y_t = \sigma_t \eta_t, \quad \text{with } \sigma_t = f(\sigma_{t-1}, \eta_{t-1}) \quad \text{and } \sigma_0 \text{ independent of } (\eta_t)_{t \geq 0}, \tag{4.2}$$

that is, σ_t is a stationary Markov chain. For the sake of simplicity, we shall always assume here that $\mathbb{E}(\eta_0^2) = 1$.

We begin with models satisfying a recursive equation whose stationary solution satisfies (4.1). The original ARCH model was introduced by Engle (1982) and generalized by Bollerslev (1986) with the class of GARCH(p,q) models defined by $Y_t = \sigma_t \eta_t$ and

$$\sigma_t^2 = a + \sum_{i=1}^{p} a_i Y_{t-i}^2 + \sum_{j=1}^{q} b_j \sigma_{t-j}^2, \tag{4.3}$$

where the coefficients $a, a_i, i = 1, \ldots, p$ and $b_j, j = 1, \ldots, q$ are all positive real numbers. Those processes were studied from the point of view of existence and stationarity of solutions by Bougerol and Picard (1992a, 1992b) and Ango Nzé (1992). Under the condition $\sum_{i=1}^{p} a_i + \sum_{j=1}^{q} b_j < 1$, this model has a unique stationary solution of the form (4.1).

Many extensions have been proposed since then. A general linear example of the model is given by the ARCH(∞) model described by Giraitis et al. (2000):

$$\sigma_t^2 = a + \sum_{j=1}^{\infty} a_j Y_{t-j}^2, \tag{4.4}$$

where $a \geq 0$ and $a_j \geq 0$. Again if $\sum_{j \geq 1} a_j < 1$, then there exists a unique strictly stationary solution to (4.4) of the form (4.1).

For the models satisfying (4.2), let us cite first the so-called augmented GARCH$(1,1)$ models introduced by Duan (1997):

$$\Lambda(\sigma_t^2) = c(\eta_{t-1}) \Lambda(\sigma_{t-1}^2) + h(\eta_{t-1}), \tag{4.5}$$

where Λ is an increasing and continuous function on $\mathbb{R}^+$. We refer to Duan (1997) for numerous examples of more standard models belonging to this class. There exists a stationary solution to (4.5), provided c satisfies the condition A_2^* given in Carrasco and Chen (2002) (this condition is satisfied as soon as

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

$\mathbb{E}(|c(\eta_0)|^s) < 1$ and $\mathbb{E}(|h(\eta_0)|^s) < \infty$ for integer $s \geq 1$; see the condition A_2 of the same paper).

An example of the model (4.5) is the threshold ARCH model (see Zakoïan, 1993):

$$\sigma_t = a + b\sigma_{t-1}\eta_{t-1}\mathbb{1}_{\{\eta_{t-1}>0\}} - c\sigma_{t-1}\eta_{t-1}\mathbb{1}_{\{\eta_{t-1}<0\}}, \quad a, b, c > 0, \tag{4.6}$$

for which $c(\eta_{t-1}) = b\eta_{t-1}\mathbb{1}_{\{\eta_{t-1}>0\}} - c\eta_{t-1}\mathbb{1}_{\{\eta_{t-1}<0\}}$ and $h = a$. In particular, the condition for the stationarity is satisfied as soon as $b \vee c < 1$.

Other models satisfying (4.2) are the nonlinear ARCH models (see Doukhan, 1994, pp. 106–107), for which

$$\sigma_t = f(\sigma_{t-1}\eta_{t-1}). \tag{4.7}$$

There exists a stationary solution to (4.7) provided that the density of η_0 is positive on a neighborhood of zero and $\limsup_{|x|\to\infty}|f(x)/x| < 1$.

Now, for models (4.3)–(4.7), the following results can be deduced from the literature. If we assume that in all cases the η_t's are centered with unit variance and admit a density with respect to the Lebesgue measure, then all of the following processes have dependence properties under which the risk bound (3.17) of Theorem 3.1 holds, provided that the other assumptions are fulfilled:

- the process $((Y_t, \sigma_t))_{t\in\mathbb{Z}}$ defined by model (4.3) under $\sum_{i=1}^p a_i + \sum_{j=1}^q b_j < 1$ (see Carrasco and Chen, 2000, 2002)
- the process $((Y_t, \sigma_t))_{t\in\mathbb{Z}}$ defined by model (4.5) under: the density of η_0 is positive on an open set containing 0; c and h are polynomial functions; there exists an integer $s \geq 1$ such that $|c(0)| < 1$, $\mathbb{E}(|c(\eta_0)|^s) < 1$, and $\mathbb{E}(|h(\eta_0)|^s) < \infty$ (see Prop. 5 in Carrasco and Chen, 2002)
- the process $((Y_t, \sigma_t))_{t\in\mathbb{Z}}$ defined by model (4.6) if $0 < b \vee c < 1$
- the process $((Y_t, \sigma_t))_{t\in\mathbb{Z}}$ defined by model (4.7) if the density of η_0 is positive on a neighborhood of 0 and $\limsup_{|x|\to+\infty}|f(x)/x| < 1$ (see Doukhan, 1994, Prop. 6, p. 107).

More precisely, the processes are then geometrically β-mixing.

Note that some other extensions to nonlinear models having stationarity and dependency properties can be found in Lee and Shin (2005).

Concerning more specifically the τ-dependence, we can prove the following result that relies on a more general property given in Section 5.

PROPOSITION 4.1. *Let Y_t and σ_t satisfy either (4.1) or (4.2). Assume that σ_0^2 and η_0^2 admit bounded densities and that $\mathbb{E}(\eta_0^2) = 1$.*

(i) For the ARCH(∞) model (4.4), the following rates for $((X_t, Z_t))_{t\geq 0}$ hold.

If $a_j = 0$, for $j \geq J$, then $((X_t, Z_t))_{t\geq 0}$ is geometrically τ-dependent.
If $a_j = O(b^j)$ for some $b < 1$ then $\tau_\infty(n) = O(\kappa^{\sqrt{n}})$ for some $\kappa < 1$.
If $a_j = O(j^{-b})$ for some $b > 1$ then $\tau_\infty(n) = O(n^{-b/2}(\ln(n))^{b+2})$.

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

(ii) For model (4.2), if there exists $\kappa < 1$ *such that*

$$\mathbb{E}(|(f(x,\eta_0))^2 - (f(y,\eta_0))^2|) \le \kappa|x^2 - y^2|, \tag{4.8}$$

then $((X_t, Z_t))_{t>0}$ *is geometrically* τ*-dependent with* $\tau_\infty(n) = O(n(\sqrt{\kappa})^n)$.

For more general models than (4.4), we refer to Doukhan et al. (2006). An example of a Markov chain satisfying (4.8) is the autoregressive model $\sigma_t^2 = h(\sigma_{t-1}^2) + r(\eta_{t-1})$ for some κ-Lipschitz function h.

4.2. Illustration of the Method

In this section, we use Matlab programs that can be found on Yves Rozenholc's page: http://www.math-info.univ-paris5.fr/~rozen/.

In their 2005 paper, Comte et al. (2007) provide a simulation study for deconvolution in the i.i.d. case. In this paper, the authors have considered the estimation of various signal densities g and two types of errors: Gaussian and Laplace. We add here the case $\varepsilon_t \rightsquigarrow \ln[(\mathcal{N}(0,1))^2]$. We refer to Comte et al. (2007) for the values of the constants in the penalty, and similarly, we take for the last case

$$\text{pen}(m) = \left(1 + \frac{\ln(\pi m)^{2.5}}{(1+\sigma_\varepsilon^2)\pi m} + \pi m\right) \int_{-\pi m}^{\pi m} \frac{dx}{|f_\varepsilon^*(x)|^2},$$

where σ_ε^2 is the (known) variance of ε_1. Many replications have been done in this context, and the reader is referred to Comte et al. (2006, 2007) for details.

Here, we just give a few illustrations for the case $\varepsilon_t \rightsquigarrow \ln[(\mathcal{N}(0,1))^2]$ (which implies that $|f_\varepsilon^*(x)| = |\Gamma(1+ix)/\sqrt{\pi}|$). More precisely, we consider three independent and three dependent models. In the three independent models, $Z_i = X_i + \ln(\eta_i^2)$ with i.i.d. $\mathcal{N}(0,1)$ η_i's and i.i.d. X_i's.

M1. X_i has density $0.6g_1 + 0.4g_2$ with $g_1 \rightsquigarrow \mathcal{N}(-2,4)$ and $g_2 \rightsquigarrow \mathcal{N}(6,4)$.

M2. X_i has density $0.6g_1 + 0.4g_2$ with $g_1 \rightsquigarrow \mathcal{N}(-2,1)$ and $g_2 \rightsquigarrow \mathcal{N}(6,1)$.

M3. $X_i \rightsquigarrow \mathcal{N}(0,9)$.

The first two models are chosen to compare the impact of the empirical signal to noise ratio defined by $s2n = \widehat{\text{var}}(Z)/\sigma_\varepsilon^2 - 1$, where $\widehat{\text{var}}(Z)$ is the empirical variance of the observations (note that $s2n$ is an estimation of the true signal to noise ratio $\text{var}(X)/\sigma_\varepsilon^2$). Model M1 has $s2n = 5.6$ and $s2n = 1.94$ for M2, in a difficult bimodal case. For M3, $s2n = 1.66$, but the curve is easier to estimate. The dependent models are as follows.

M4. GARCH(1,1) process (i.e., $p = q = 1$ in (4.3)) with parameter values $a = 1, a_1 = 0.7, b_1 = 0.2$ (as in van Es et al., 2005).

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press




FIGURE 1. Left: $x \mapsto |f_1^*(x)| = 1/(1+x^2)$, solid line; $x \mapsto |f_2^*(x)| = e^{-x^2/2}$, dotted line; $x \mapsto |f_\varepsilon^*(x)| = |\Gamma(\frac{1}{2} + ix)/\sqrt{\pi}|$, dashed-dotted line. Right: Deconvolution of mixed normals, $n = 1{,}000$ data. True density, solid line. Estimator computed with the right characteristic function f_ε^*, small dotted line. Estimator computed with f_2^* instead of f_ε^*, large dotted line. Estimator computed with f_1^* instead of f_ε^*, dashed-dotted line.

M5. GARCH(1,1) process ($p = q = 1$ in (4.3)) with parameter values $a = 5$, $a_1 = 0.79$, and $b_1 = 0.2$.

M6. $Y_t = \sigma_t \eta_t$ and $\sigma_t^2 = \tau^2 \sigma_{t-1}^2 + 1/\eta_{t-1}^2$, with i.i.d. $\mathcal{N}(0,1)$ variables η_t and $\tau^2 = 0.5$ or $\tau^2 = 0.8$ (the de Vries, 1991, example, eqn. (10)).

First, we consider the problem of misspecification of the noise density. The left panel of Figure 1 plots the modulus of the characteristic function of the Laplace distribution f_1 of the $\mathcal{N}(0,1)$ distribution f_2 and of the noise density f_ε. The right panel of Figure 1 shows the true density g and the estimators computed with f_ε^* (true characteristic function of f_ε) and f_1^* and f_2^* (wrong characteristic functions of f_ε) for M1. The left panel of Figure 2 and the right panel of




FIGURE 2. Deconvolution of a mixed normal (right) and of a normal (left) density, $n = 1{,}000$ data. True density, solid line. Estimator computed with the right characteristic function f_ε^*, small dotted line. Estimator computed with f_2^* instead of f_ε^*, large dotted line. Estimator computed with f_1^* instead of f_ε^*, dashed-dotted line.

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

Figure 2 show the same picture for M2 and M3, respectively. The left panel of Figure 2 shows clearly that if the estimator is computed with a wrong characteristic function of f_ε, then it can be very far from the true distribution g. This is not surprising because we have recalled in the Introduction that if f_ε is not completely known, then the model may be nonidentifiable.

For the dependent processes, as the true stationary density is generally unknown, we compare the estimated curve with the optimal histogram selected with the Birgé and Rozenholc (2006) adaptive procedure using the direct data $\ln(\sigma_t^2)$. This histogram is a reliable indication of the shape of the true density, all the more because it is based on the direct data (which are available only in the simulation context). For M4, we obtain a good result. The top left panel of Figure 3 shows the estimated curve based on the $\ln(Y_t^2)$ data. We can see that the peak is slightly cut. The result is better in the top right panel of Figure 3 (model M5), when the signal to noise ratio increases (it reaches 0.35 instead of the ratio of 0.13 obtained in the previous case). For M6, we choose




FIGURE 3. Deconvolution of two GARCH processes (top) and of two de Vries models (bottom). Estimated curve and optimal histogram on the direct data $\ln(\sigma_t^2)$ for $n = 1{,}000$ simulated data.

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

$\tau^2 = 0.5$ (Figure 3, bottom left panel) and $\tau^2 = 0.8$ (Figure 3, bottom right panel) and obtain a signal to noise ratio that is equal to 0.6 (left) and 1.34 (right) in the bottom two panels of Figure 3. Here, the procedure works well, which shows that the penalty is well calibrated.

5. PROOFS

5.1. Two Useful Tools in the Dependent Context

We first recall the coupling property associated with the dependence coefficients. Assume that Ω is rich enough, which means that there exists U uniformly distributed over $[0,1]$ and independent of $\mathcal{M} \vee \sigma(W)$. There exists an $\mathcal{M} \vee \sigma(U) \vee \sigma(W)$-measurable random variable $W^\star$ distributed as W and independent of $\mathcal{M}$ such that

$$\tau(\mathcal{M}, W) = \mathbb{E}(\|W - W^\star\|_{\mathbb{B}}). \tag{5.1}$$

Equality in (5.1) has been established in Dedecker and Prieur (2005, Sect. 7.1).

As consequence of the coupling property (5.1), we have the following covariance inequality. Let $\|\cdot\|_{\infty,\mathbb{P}}$ be the $\mathbb{L}^\infty(\Omega,\mathbb{P})$-norm. For two measurable functions f, h from $\mathbb{R}$ to $\mathbb{C}$, and if $\mathrm{Lip}(h)$ is the Lipschitz coefficient of h,

$$|\mathrm{Cov}\,(f(Y), h(X))| \le \|f(Y)\|_{\infty,\mathbb{P}} \mathrm{Lip}(h) \tau(\sigma(Y), X). \tag{5.2}$$

Thus, using that $t \to e^{ixt}$ is $|x|$-Lipschitz, we obtain the bound

$$|\mathrm{Cov}(e^{ixZ_1}, e^{ixX_k})| \le |x| \tau_1(k-1). \tag{5.3}$$

Proof of Proposition 3.1 The proof of Proposition 3.1 follows the same lines as in the independent framework (see Comte et al., 2006). The main difference lies in the control of the variance term. We keep the same notation as in Section 2.1. According to (2.2), for any given m belonging to $\{1, \ldots, m_n\}$, $\hat{g}_m^{(n)}$ satisfies, $\gamma_n(\hat{g}_m^{(n)}) - \gamma_n(g_m^{(n)}) \le 0$. For a random variable T with density f_T, and any function ψ such that $\psi(T)$ is integrable, set $\nu_{n,T}(\psi) = n^{-1} \sum_{i=1}^n [\psi(T_i) - \langle \psi, f_T \rangle]$. In particular,

$$\nu_{n,Z}(u_t^*) = \frac{1}{n} \sum_{i=1}^n [u_t^*(Z_i) - \langle t, g \rangle]. \tag{5.4}$$

Because

$$\gamma_n(t) - \gamma_n(s) = \|t - g\|^2 - \|s - g\|^2 - 2\nu_{n,Z}(u_{t-s}^*), \tag{5.5}$$

we infer that

$$\|g - \hat{g}_m^{(n)}\|^2 \le \|g - g_m^{(n)}\|^2 + 2\nu_{n,Z}(u^*_{\hat{g}_m^{(n)} - g_m^{(n)}}). \tag{5.6}$$

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

Writing that $\hat{a}_{m,j} - a_{m,j} = \nu_{n,Z}(u^*_{\varphi_{m,j}})$, we obtain that

$$\nu_{n,Z}(u^*_{\hat{g}_m^{(n)} - g_m^{(n)}}) = \sum_{|j| \le k_n} (\hat{a}_{m,j} - a_{m,j}) \nu_{n,Z}(u^*_{\varphi_{m,j}}) = \sum_{|j| \le k_n} [\nu_{n,Z}(u^*_{\varphi_{m,j}})]^2.$$

Consequently, $\mathbb{E}\|g - \hat{g}_m^{(n)}\|^2 \le \|g - g_m^{(n)}\|^2 + 2\sum_{j \in \mathbb{Z}} \mathbb{E}[(\nu_{n,Z}(u^*_{\varphi_{m,j}}))^2]$. According to Comte et al. (2006),

$$\|g - g_m^{(n)}\|^2 = \|g - g_m\|^2 + \|g_m - g_m^{(n)}\|^2 \le \|g - g_m\|^2 + \frac{(\pi m)^2 (M_2 + 1)}{k_n}. \quad \textbf{(5.7)}$$

The variance term is studied by using first that for $f \in \mathbb{L}_1(\mathbb{R})$,

$$\nu_{n,Z}(f^*) = \int \nu_{n,Z}(e^{ix\cdot}) f(x)\, dx. \quad \textbf{(5.8)}$$

Now, we use (5.8) and apply Parseval's formula to obtain

$$\begin{aligned} \mathbb{E}\left(\sum_{j \in \mathbb{Z}} (\nu_{n,Z}(u^*_{\varphi_{m,j}}))^2 \right) &= \frac{1}{4\pi^2} \sum_{j \in \mathbb{Z}} \mathbb{E}\left(\int \frac{\varphi^*_{m,j}(-x)}{f^*_\varepsilon(x)} \nu_{n,Z}(e^{ix\cdot})\, dx \right)^2 \\ &= \frac{1}{2\pi} \int_{-\pi m}^{\pi m} \frac{\mathbb{E}|\nu_{n,Z}(e^{ix\cdot})|^2}{|f^*_\varepsilon(x)|^2}\, dx. \end{aligned} \quad \textbf{(5.9)}$$

Because $\nu_{n,Z}$ involves centered and stationary variables, we have

$$\mathbb{E}|\nu_{n,Z}(e^{ix\cdot})|^2 = \operatorname{Var}|\nu_{n,Z}(e^{ix\cdot})| = \frac{1}{n} \operatorname{Var}(e^{ixZ_1}) + \frac{1}{n^2} \sum_{1 \le k \ne l \le n} \operatorname{Cov}(e^{ixZ_k}, e^{ixZ_l}). \quad \textbf{(5.10)}$$

It follows from the structure of the model that, for $k < l$, ε_l is independent of (X_l, Z_k), so that $\mathbb{E}(e^{ixZ_k}) = f^*_\varepsilon(x) g^*(x)$ and $\mathbb{E}(e^{ix(Z_l - Z_k)}) = f^*_\varepsilon(x) \mathbb{E}(e^{ix(X_l - Z_k)})$. Thus, for $k < l$,

$$\operatorname{Cov}(e^{ixZ_k}, e^{ixZ_l}) = f^*_\varepsilon(x) \operatorname{Cov}(e^{ixZ_k}, e^{ixX_l}). \quad \textbf{(5.11)}$$

From (5.10) and the stationarity of $(X_i)_{i \ge 1}$, we obtain that

$$\mathbb{E}|\nu_{n,Z}(e^{ix\cdot})|^2 \le \frac{1}{n} + \frac{2}{n} \sum_{k=2}^{n} |\operatorname{Cov}(e^{ixZ_1}, e^{ixX_k})|\, |f^*_\varepsilon(x)|. \quad \textbf{(5.12)}$$

The first part of Proposition 3.1 follows from the stationarity of the X_i's and from (5.6), (5.7), (5.9), and (5.12). The proof of $R_m \le R_{m,\tau}$, where $R_{m,\tau}$ is defined in Proposition 3.1, comes from inequality (5.3). Hence we get the result. ■

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

Proof of Corollary 3.1 According to Butucea and Tsybakov (2008a, 2008b), under (3.9), we have

$$\lambda_1(f_\varepsilon,\kappa_0')\Gamma(m)(1+o_m(1)) \le \Delta(m) \le \lambda_1(f_\varepsilon,\kappa_0)\Gamma(m)(1+o_m(1)) \quad \text{as } m\to\infty,$$
$$\text{where} \quad \Gamma(m) = (1+(\pi m)^2)^{\gamma}(\pi m)^{1-\delta}\exp\{2\mu(\pi m)^{\delta}\}, \qquad \textbf{(5.13)}$$

where λ_1 is defined in (3.10). In the same way

$$\overline{\lambda_1}(f_\varepsilon,\kappa_0')\bar{\Gamma}(m)(1+o_m(1))$$
$$\le \Delta_{1/2}(m) \le \overline{\lambda_1}(f_\varepsilon,\kappa_0)\bar{\Gamma}(m)(1+o_m(1)) \quad \text{as } m\to\infty,$$

where

$$\bar{\Gamma}(m) = (1+(\pi m)^2)^{\gamma/2}(\pi m)^{1-\delta}\exp(\mu(\pi m)^{\delta}),$$
$$\overline{\lambda_1}(f_\varepsilon,\kappa_0) = [\kappa_0^2\pi(\mathbb{1}_{\{\delta=0\}}+\mu\delta\mathbb{1}_{\{\delta>0\}})]^{-1}.$$

It is easy to see that $\Delta_{1/2}(m) \le \sqrt{m\Delta(m)}$ and hence $\Delta_{1/2}(m) = \Gamma(m)o_m(1)$. Now, as soon as $\gamma > 1$ when $\delta = 0$, $m\Delta_{1/2}(m) = \Gamma(m)o_m(1)$. Set m_1 such that for $m \ge m_1$ we have

$$0.5\lambda_1(f_\varepsilon,\kappa_0')\Gamma(m) \le \Delta(m) \le 2\lambda_1(f_\varepsilon,\kappa_0)\Gamma(m) \qquad \textbf{(5.14)}$$

and

$$0.5\overline{\lambda_1}(f_\varepsilon,\kappa_0')\bar{\Gamma}(m) \le \Delta_{1/2}(m) \le 2\overline{\lambda_1}(f_\varepsilon,\kappa_0)\bar{\Gamma}(m). \qquad \textbf{(5.15)}$$

If $\sum_{k\ge1}\tau_1(k) < +\infty$, if $\gamma > 1$ when $\delta = 0$, if (3.9) and (3.5) hold, and if $k_n \ge n$, then we have the upper bound: for $m \ge m_1$,

$$\mathbb{E}\|g-\hat{g}_m^{(n)}\|^2 \le \|g-g_m\|^2 + \frac{m^2(M_2+1)}{n} + \frac{2\lambda_1\Gamma(m)}{n}$$
$$+ 2\pi\overline{\lambda_1}\sum_{k\ge1}\tau_1(k)\frac{m\bar{\Gamma}(m)}{n}$$
$$\le \|g-g_m\|^2 + \frac{m^2(M_2+1)}{n} + \frac{2\lambda_1\Gamma(m)}{n}$$
$$+ \frac{C\left(\sum_{k\ge1}\tau_1(k)\right)\Gamma(m)}{n}o_m(1).$$

Because $\gamma > 1$ when $\delta = 0$, the residual term $n^{-1}m^2(M_2+1)$ is negligible with respect to the variance term.

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

Finally, g_m being the orthogonal projection of g on S_m, we get $g_m^* = g^* \mathbb{1}_{[-m\pi, m\pi]}$, and therefore

$$\|g - g_m\|^2 = \frac{1}{2\pi} \|g^* - g_m^*\|^2 = \frac{1}{2\pi} \int_{|x| \geq \pi m} |g^*|^2(x)\, dx.$$

If g belongs to the class $\mathcal{S}_{s,r,b}(C_1)$ defined in (3.11), then

$$\|g - g_m\|^2 \leq \frac{C_1}{2\pi} (m^2\pi^2 + 1)^{-s} \exp\{-2b\pi^r m^r\}.$$

The corollary is proved. ■

Proof of Theorem 3.1. By definition, $\tilde{g}$ satisfies that for all $m \in \{1, \ldots, m_n\}$,

$$\gamma_n(\tilde{g}) + \text{pen}(\hat{m}) \leq \gamma_n(g_m) + \text{pen}(m).$$

Therefore, by using (5.5) we get

$$\|\tilde{g} - g\|^2 \leq \|g_m^{(n)} - g\|^2 + 2\nu_{n,Z}(u^*_{\tilde{g} - g_m^{(n)}}) + \text{pen}(m) - \text{pen}(\hat{m}),$$

where $\nu_{n,Z}$ is defined in (5.4). If $t = t_1 + t_2$ with t_1 in $S_m^{(n)}$ and t_2 in $S_{m'}^{(n)}$, t^* has its support in $[-\pi \max(m, m'), \pi \max(m, m')]$, and t belongs to $S^{(n)}_{\max(m,m')}$. Set $B_{m,m'}(0,1) = \{t \in S^{(n)}_{\max(m,m')} \,/\, \|t\| = 1\}$ and write

$$|\nu_{n,Z}(u^*_{\tilde{g} - g_m^{(n)}})| \leq \|\tilde{g} - g_m^{(n)}\| \sup_{t \in B_{m,\hat{m}}(0,1)} |\nu_{n,Z}(u_t^*)|.$$

Using that $2uv \leq a^{-1}u^2 + av^2$ for any $a > 1$ leads to

$$\|\tilde{g} - g\|^2 \leq \|g_m^{(n)} - g\|^2 + a^{-1}\|\tilde{g} - g_m^{(n)}\|^2 + a \sup_{t \in B_{m,\hat{m}}(0,1)} (\nu_{n,Z}(u_t^*))^2 + \text{pen}(m) - \text{pen}(\hat{m}).$$

Therefore, we have

$$\|\tilde{g} - g\|^2 \leq \kappa_a^2 \|g_m^{(n)} - g\|^2 + a\kappa_a \sup_{t \in B_{m,\hat{m}}(0,1)} |\nu_{n,Z}(u_t^*)|^2 + \kappa_a(\text{pen}(m) - \text{pen}(\hat{m})), \tag{5.16}$$

where κ_a is defined in (3.16).

The main point is to control $\sup_{t \in B_{m,\hat{m}}(0,1)} |\nu_{n,Z}(u_t^*)|^2$, the supremum of a centered empirical process $\nu_{n,Z}(u_t^*)$. To handle this supremum, we use coupling methods to replace the dependent random variables involved in this process by block-independent random variables to apply Talagrand's inequality. Heuristically, we replace the expectation of the supremum by the (negligible)

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

price of coupling plus the expectation of a term $p(m,\hat{m})$ imposed by Talagrand's Inequality, which in turn will fix the choice of the penalty function.

We use the coupling method recalled in Section 5.1 to build approximating variables for the $W_i = (Z_i, X_i)$'s. More precisely, we build variables $W_i^\star$ such that if $n = 2p_nq_n + r_n$, $0 \le r_n < q_n$ and $\ell = 0, \dots, p_n - 1$,

$$E_\ell = (W_{2\ell q_n+1}, \dots, W_{(2\ell+1)q_n}), \qquad F_\ell = (W_{(2\ell+1)q_n+1}, \dots, W_{(2\ell+2)q_n}),$$

$$E_\ell^\star = (W^\star_{2\ell q_n+1}, \dots, W^\star_{(2\ell+1)q_n}), \qquad F_\ell^\star = (W^\star_{(2\ell+1)q_n+1}, \dots, W^\star_{(2\ell+2)q_n}).$$

The variables $E_\ell^\star$ and $F_\ell^\star$ are such that

- $E_\ell^\star$ and E_ℓ are identically distributed, and $F_\ell^\star$ and F_ℓ are identically distributed;
- $\sum_{i=1}^{q_n} \mathbb{E}(\|W_{2\ell q_n+i} - W^\star_{2\ell q_n+i}\|_{\mathbb{R}^2}) \le q_n\tau_\infty(q_n)$, $\sum_{i=1}^{q_n} \mathbb{E}(\|W_{(2\ell+1)q_n+i} - W^\star_{(2\ell+1)q_n+i}\|_{\mathbb{R}^2}) \le q_n\tau_\infty(q_n)$;
- and therefore $E_\ell^\star$ is independent of $\mathcal{M}_{(2\ell-1)q_n}$, and the same holds for the blocks $F_\ell^\star$.

For the sake of simplicity we assume that $r_n = 0$. We denote by $(Z_i^\star, X_i^\star) = W_i^\star$ the new couple of variables. Using the notation of (5.4), we denote by $\nu^\star_{n,Z}(u_t^*)$ the empirical contrast computed on the $Z_i^\star$. Then we write

$$\begin{aligned}\|\tilde{g} - g\|^2 \le{}& \kappa_a^2\|g - g_m^{(n)}\|^2 + 2a\kappa_a \sup_{t\in B_{m,\hat{m}}(0,1)} |\nu^\star_{n,Z}(u_t^*)|^2 \\ &+ \kappa_a(\mathrm{pen}(m) - \mathrm{pen}(\hat{m})) + 2a\kappa_a \sup_{t\in B_{m,\hat{m}}(0,1)} |\nu^\star_{n,Z}(u_t^*) - \nu_{n,Z}(u_t^*)|^2.\end{aligned}$$

Set

$$T_n^\star(m,m') := \left[\sup_{t\in B_{m,m'}(0,1)} |\nu^\star_{n,Z}(t)|^2 - p(m,m')\right]_+, \tag{5.17}$$

where $p(m,m')$ will defined further. Hence

$$\begin{aligned}\|\tilde{g} - g\|^2 &\le \kappa_a^2\|g - g_m^{(n)}\|^2 + 2a\kappa_a T_n^\star(m,\hat{m}) \\ &\quad + \kappa_a(2ap(m,\hat{m}) + \mathrm{pen}(m) - \mathrm{pen}(\hat{m})) \\ &\quad + 2a\kappa_a \sup_{t\in B_{m,\hat{m}}(0,1)} |\nu_{n,Z}(u_t^*) - \nu^\star_{n,Z}(u_t^*)|^2 \\ &\le \kappa_a^2\|g - g_m^{(n)}\|^2 + 2\kappa_a\mathrm{pen}(m) \\ &\quad + 2a\kappa_a \sup_{t\in B_{m,\hat{m}}(0,1)} |\nu_{n,Z}(u_t^*) - \nu^\star_{n,Z}(u_t^*)|^2 \\ &\quad + 2a\kappa_a T_n^\star(m,\hat{m}),\end{aligned} \tag{5.18}$$

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

where $\mathrm{pen}(m)$ is chosen such that

$$2ap(m,m') \le \mathrm{pen}(m) + \mathrm{pen}(m'). \tag{5.19}$$

Now write

$$\begin{aligned}\nu_{n,Z}(u_t^*) - \nu_{n,Z}^\star(u_t^*) &= \frac{1}{2\pi}\frac{1}{n}\sum_{k=1}^{n}\int [e^{ixZ_k} - e^{ixZ_k^\star}]\,\frac{t^*(-x)}{f_\varepsilon^*(x)}\,dx \\ &= \frac{1}{2\pi}\int [\nu_{n,Z}(e^{ix\cdot}) - \nu_{n,Z}^\star(e^{ix\cdot})]\,\frac{t^*(-x)}{f_\varepsilon^*(x)}\,dx.\end{aligned}$$

Consequently,

$$\begin{aligned}&\mathbb{E}\left[\sup_{t\in B_{m,\dot m}(0,1)} |\nu_{n,Z}(u_t^*) - \nu_{n,Z}^\star(u_t^*)|^2\right] \\ &\quad\le \frac{1}{2\pi}\int_{-\pi m_n}^{\pi m_n} \mathbb{E}[|\nu_{n,Z}(e^{ix\cdot}) - \nu_{n,Z}^\star(e^{ix\cdot})|^2]\,\frac{1}{|f_\varepsilon^*(x)|^2}\,dx.\end{aligned} \tag{5.20}$$

Next, the bound $|e^{-ixt} - e^{-ixs}| \le |x||t-s|$ implies that

$$\sum_{i=1}^{q_n} \mathbb{E}(|e^{-ixZ_{2\ell q_n+i}} - e^{-ixZ_{2\ell q_n+i}^\star}|^2) \le 2q_n|x|\tau_\infty(q_n).$$

It follows that

$$\begin{aligned}\mathbb{E}\left[\sup_{t\in B_{m,\dot m}(0,1)} |\nu_{n,Z}(u_t^*) - \nu_{n,Z}^\star(u_t^*)|^2\right] &\le \frac{\tau_\infty(q_n)}{\pi}\int_{-\pi m_n}^{\pi m_n}\frac{|x|}{|f_\varepsilon^*(x)|^2}\,dx \\ &\le 2\tau_\infty(q_n)m_n\Delta(m_n).\end{aligned} \tag{5.21}$$

By gathering (5.18) and (5.21) we get

$$\begin{aligned}\mathbb{E}\|\tilde g - g\|^2 \le \kappa_a^2\|g - g_m^{(n)}\|^2 &+ 2a\kappa_a\sum_{m'=1}^{m_n}\mathbb{E}[T_n^\star(m,m')] \\ &+ 2\kappa_a\mathrm{pen}(m) + 2a\kappa_a\tau_\infty(q_n)m_n\Delta(m_n).\end{aligned}$$

Therefore we infer that, for all $m \in \{1,\dots,m_n\}$,

$$\mathbb{E}\|g - \tilde g\|^2 \le C_a[\|g - g_m^{(n)}\|^2 + \mathrm{pen}(m)] + 2a\kappa_a(C_1 + C_2)/n, \tag{5.22}$$

provided that

$$\Delta(m_n)m_n\tau_\infty(q_n) \le C_1/n \quad\text{and}\quad \sum_{m'=1}^{m_n}\mathbb{E}(T_n^\star(m,m')) \le C_2/n. \tag{5.23}$$

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

Using (5.14), we conclude that the first part of (5.23) is fulfilled as soon as

$$m_n^{2\gamma+2-\delta}\exp\{2\mu\pi^\delta m_n^\delta\}\tau_\infty(q_n) \le C_1'/n. \tag{5.24}$$

To ensure that our estimators converge, we only consider models with bounded penalty, and therefore (5.24) requires that $m_n\tau_\infty(q_n) \le C_1'/n^2$. For $q_n = [n^c]$ and $\tau_\infty(k) = O(n^{-1-\theta})$, we obtain the condition

$$m_n n^{-c(1+\theta)} = O(n^{-2}). \tag{5.25}$$

If f_ε satisfies (3.9) with $\delta > 0$, and if $\theta > 3$, one can find $c \in]0, \frac{1}{2}[$ such that (5.25) is satisfied. Now, if $\delta = 0$ and $\gamma \ge \frac{3}{2}$ in (3.9) and if $\theta > 3 + 2/(1+2\gamma)$, then one can find $c \in]0, \frac{1}{2}[$, such that (5.25) is satisfied. These conditions ensure that (5.24) holds.

To prove the second part of (5.23), we split $T_n^\star(m,m')$ into two terms:

$$T_n^\star(m,m') \le (T_{n,1}^\star(m,m') + T_{n,2}^\star(m,m'))/2,$$

where $p_1(m,m') = p_2(m,m') = p(m,m')/2$, and for $k = 1, 2$

$$T_{n,k}^\star(m,m') = \left[\sup_{t\in B_{m,m'}(0,1)}\left|\frac{1}{p_n q_n}\sum_{\ell=1}^{p_n}\sum_{i=1}^{q_n}(u_t^*(Z_{(2\ell+k-1)q_n+i}^\star) - \langle t,g\rangle)\right|^2 - p_k(m,m')\right]_+. \tag{5.26}$$

We only study $T_{n,1}^\star(m,m')$ and conclude for $T_{n,2}^\star(m,m')$ analogously. The study of $T_{n,1}^\star(m,m')$ consists in applying a concentration inequality to $\nu_{n,1}^\star(t)$ defined by

$$\nu_{n,1}^\star(t) = \frac{1}{p_n q_n}\sum_{\ell=1}^{p_n}\sum_{i=1}^{q_n}(u_t^*(Z_{2\ell q_n+i}^\star) - \langle t,g\rangle) = \frac{1}{p_n}\sum_{\ell=1}^{p_n}\nu_{q_n,\ell}^\star(u_t^*). \tag{5.27}$$

The random variable $\nu_{n,1}^\star(u_t^*)$ is considered the sum of the p_n independent random variables $\nu_{q_n,\ell}^\star(t)$ defined as

$$\nu_{q_n,\ell}^\star(u_t^*) = (1/q_n)\sum_{j=1}^{q_n}u_t^*(Z_{2\ell q_n+j}^\star) - \langle t,g\rangle. \tag{5.28}$$

Let $m^* = \max(m,m')$. Let $M_1^\star(m^*)$, $v^\star(m^*)$, and $H^\star(m^*)$ be some terms such that $\sup_{t\in B_{m,m'}(0,1)}\|\nu_{q_n,\ell}^\star(u_t^*)\|_\infty \le M_1^\star(m^*)$, $\sup_{t\in B_{m,m'}(0,1)}\mathrm{Var}(\nu_{q_n,\ell}^\star(u_t^*)) \le v^\star(m)$, and last $\mathbb{E}(\sup_{t\in B_{m,m'}(0,1)}|\nu_{n,1}^\star(u_t^*)|) \le H^\star(m^*)$.

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

According to Lemma 5.2 in Section 5.2, we take

$$(H^\star(m^*))^2 = \frac{2\Delta(m^*)}{n}, \qquad M_1^\star(m^*) = \sqrt{\Delta(m^*)}, \quad \text{and}$$

$$v^\star(m^*) = \frac{C_{v^*}\sqrt{\Delta_2(m^*, f_Z)}}{2\pi q_n},$$

where $\Delta_2(m, f_Z)$ is defined by

$$\Delta_2(m, f_Z) = \int_{-\pi m}^{\pi m}\int_{-\pi m}^{\pi m} \frac{|f_Z^*(x-y)|^2}{|f_\varepsilon^*(x) f_\varepsilon^*(y)|^2}\, dxdy \tag{5.29}$$

and where

$$C_{v^*} = 2\left[\mathbb{1}_{\delta>0} + \frac{\sqrt{2}\pi^{3/2}(2\pi)^{3/2}}{\sqrt{3}} \sum_{k\geq 1} \tau_1(k)\mathbb{1}_{\delta=0}\right]. \tag{5.30}$$

From the definition of $T_{n,1}^\star(m, m')$, by taking $p_1(m, m') = 2(1 + 2\xi^2)(H^\star)^2(m^*)$, we get

$$\mathbb{E}(T_{n,1}^\star(m, m')) \leq \mathbb{E}\left[\sup_{t\in B_{m,m'}(0,1)} \left|\nu_{n,1}^\star(u_t^*) - 2(1+2\xi^2)(H^\star)^2(m^*)\right|\right]_+. \tag{5.31}$$

According to the condition (5.19), we thus take

$$\begin{aligned}\text{pen}(m) &= 2ap(m,m) = 2a(p_1(m,m) + p_2(m,m)) = 4ap_1(m,m)\\ &= 8a(1+2\xi^2)(2n^{-1}\Delta(m)) = 16a(1+2\xi^2)n^{-1}\Delta(m),\end{aligned} \tag{5.32}$$

where ξ^2 is suitably chosen. Set m_2 and m_3 as defined in Lemma 5.2 and set m_1 such that for $m^* \geq m_1$, $\Delta(m^*)$ satisfies (5.14). Take $m_0 = m_1 \vee m_2 \vee m_3$. We split the sum over m' in two parts and write

$$\sum_{m'=1}^{m_n} \mathbb{E}(T_{n,1}^\star(m, m')) = \sum_{m'|m^*\leq m_0} \mathbb{E}(T_{n,1}^\star(m, m')) + \sum_{m'|m^*\geq m_0} \mathbb{E}(T_{n,1}^\star(m, m')). \tag{5.33}$$

By applying Lemma 5.3 in Section 5.2, we get $\mathbb{E}(T_{n,1}^\star(m, m')) \leq K[I(m^*) + II(m^*)]$, where

$$I(m^*) = \frac{\sqrt{\Delta_2(m^*, f_Z)}}{p_n} \exp\left\{-K_1\xi^2 \frac{\Delta(m^*)}{q_n v^\star(m^*)}\right\},$$

$$II(m^*) = \frac{\Delta(m^*)}{p_n^2} \exp\left\{-\frac{2K_1\xi C(\xi)^2}{7}\frac{\sqrt{n}}{q_n}\right\}.$$

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

When $m^* \le m_0$, with m_0 finite, we get that, for all $m \in \{1, \dots, m_n\}$,

$$\sum_{m' | m^* \le m_0} \mathbb{E}(R^{\star}_{n,1}(m, m')) \le \frac{C(m_0)}{n}.$$

We now come to the sum over m' such that $m^* \ge m_0$. It follows from Comte et al. (2006) that

$$v^{\star}(m^*) = \frac{C_{v^*}\sqrt{\Delta_2(m^*, f_Z)}}{2\pi q_n} \le \lambda_2^{\star}(f_\varepsilon, \kappa_0)\, \frac{\Gamma_2(m^*)}{q_n}, \tag{5.34}$$

with

$$\lambda_2^{\star}(f_\varepsilon, \kappa_0) = \kappa_0^{-1} C_{v^*}\sqrt{2\pi\lambda_1}\, \|f_{\varepsilon^*}\|\, \mathbb{1}_{\delta \le 1} + \mathbb{1}_{\delta > 1}, \tag{5.35}$$

where $\lambda_1 = \lambda_1(f_\varepsilon, \kappa_0)$ is defined in (3.10) and

$$\begin{aligned}\Gamma_2(m) &= (1 + (\pi m)^2)^{\gamma} (\pi m)^{\min((1/2-\delta/2),(1-\delta))} \exp(2\mu(\pi m)^{\delta}) \\ &= (\pi m)^{-(1/2-\delta/2)_+} \Gamma(m). \end{aligned} \tag{5.36}$$

By combining the left-hand sides of (5.14) and (5.34), we get that, for $m^* \ge m_0$,

$$I(m^*) \le \frac{\lambda_2^{\star}(f_\varepsilon, \kappa_0)\Gamma_2(m^*)}{2n} \exp\left\{ - \frac{K_1 \xi^2 \lambda_1(f_\varepsilon, \kappa_0')}{\lambda_2^{\star}(f_\varepsilon, \kappa_0)} (\pi m^*)^{(1/2-\delta/2)_+} \right\}$$

$$\text{and} \quad II(m^*) \le \frac{\Delta(m^*) q_n^2}{n^2} \exp\left\{ - \frac{2K_1 \xi C(\xi^2)}{7} \frac{\sqrt{n}}{q_n} \right\}.$$

Study of $\sum_{m' | m^* \ge m_0} II(m^*)$. According to the choices for $v^{\star}(m^*)$, $(H^{\star}(m^*))^2$, and $M_1^{\star}(m^*)$, we have

$$\begin{aligned}\sum_{m' | m^* \ge m_0} II(m^*) &\le \sum_{m' \in \{1, \dots, m_n\}} \frac{\Delta(m^*) q_n^2}{n^2} \exp\left\{ - \frac{2K_1 \xi C(\xi^2)}{7} \frac{\sqrt{n}}{q_n} \right\} \\ &= O\left[m_n \exp\left\{ - \frac{2K_1 \xi C(\xi^2)}{7} \frac{\sqrt{n}}{q_n} \right\} \frac{\Delta(m_n) q_n^2}{n^2} \right].\end{aligned}$$

Because $\Delta(m_n)/n$ is bounded, then $q_n = [n^c]$ with c in $]0, \frac{1}{2}[$ ensures that

$$\sum_{m'=1}^{m_n} m_n \exp\left\{ - \frac{2K_1 \xi C(\xi^2)}{7} \frac{\sqrt{n}}{q_n} \right\} \frac{\Delta(m_n) q_n^2}{n^2} \le \frac{C}{n}. \tag{5.37}$$

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

Consequently

$$\sum_{m' | m^* \geq m_0} II^\star(m^*) \leq \frac{C}{n}. \tag{5.38}$$

Study of $\sum_{m' | m^* \geq m_0} I(m^*)$. Denote by $\psi = 2\gamma + \min(\frac{1}{2} - \delta/2, 1 - \delta)$, $\omega = (\frac{1}{2} - \delta/2)_+$, and $K' = K_1 \lambda_1(f_\varepsilon, \kappa_0')/(2\lambda_2^\star(f_\varepsilon, \kappa_0))$. For $a, b \geq 1$, we use that

$$\begin{aligned}
&\max(a,b)^\psi e^{2\mu\pi^\delta \max(a,b)^\delta} e^{-K'\xi^2 \max(a,b)^\omega} \\
&\quad \leq (a^\psi e^{2\mu\pi^\delta a^\delta} + b^\psi e^{2\mu\pi^\delta b^\delta}) e^{-(K'\xi^2/2)(a^\omega + b^\omega)} \\
&\quad \leq a^\psi e^{2\mu\pi^\delta a^\delta} e^{-(K'\xi^2/2)a^\omega} e^{-(K'\xi^2/2)b^\omega} + b^\psi e^{2\mu\pi^\delta b^\delta} e^{-(K'\xi^2/2)b^\omega}.
\end{aligned} \tag{5.39}$$

Consequently,

$$\begin{aligned}
\sum_{m' | m^* \geq m_0} I(m^*) &\leq \sum_{m'=1}^{m_n} \frac{\lambda_2^\star(f_\varepsilon, \kappa_0)\Gamma_2(m^*)}{2n} \\
&\quad \times \exp\left\{ - \frac{K_1 \xi^2 \lambda_1(f_\varepsilon, \kappa_0')}{\lambda_2^\star(f_\varepsilon, \kappa_0)} (\pi m^*)^{(1/2-\delta/2)_+} \right\} \\
&\leq \frac{\lambda_2^\star(f_\varepsilon, \kappa_0)\Gamma_2(m)}{n} \exp\left\{ - \frac{K'\xi^2}{2} (\pi m)^{(1/2-\delta/2)_+} \right\} \\
&\quad \times \sum_{m'=1}^{m_n} \exp\left\{ - \frac{K'\xi^2}{2} (\pi m')^{(1/2-\delta/2)_+} \right\} \\
&\quad + \sum_{m'=1}^{m_n} \frac{\lambda_2^\star(f_\varepsilon, \kappa_0)\Gamma_2(m')}{n} \exp\left\{ - \frac{K'\xi^2}{2} (\pi m')^{(1/2-\delta/2)_+} \right\}.
\end{aligned} \tag{5.40}$$

Case $0 \leq \delta < \frac{1}{3}$. In that case, because $\delta < (\frac{1}{2} - \delta/2)_+$, the choice $\xi^2 = 1$ ensures that $\Gamma_2(m)\exp\{-(K'\xi^2/2)(\pi m)^{(1/2-\delta/2)}\}$ is bounded, and thus the first term in (5.40) is bounded by C/n. Clearly the term $\sum_{m'=1}^{m_n} \Gamma_2(m') \exp\{-(K'/2)(m')^{(1/2-\delta/2)}\}/n$ is bounded by C'/n, and hence

$$\sum_{m' | m^* \geq m_0} I(m^*) \leq \frac{C}{n}.$$

According to (5.19), the result follows by choosing $\text{pen}(m) = 2ap(m,m) = 48a\Delta(m)/n$.

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

Case $\delta = \frac{1}{3}$. According to the inequality (5.40), we choose ξ^2 such that $(K'\xi^2/2)(\pi m)^\delta = (2+\epsilon)\mu(\pi m)^\delta$ for some $\epsilon > 0$, for instance,

$$\xi^2 = \frac{49\mu\lambda_2^\star(f_\varepsilon, \kappa_0)}{\lambda_1(f_\varepsilon, \kappa_0')}.$$

Arguing as for the case $0 \le \delta < \frac{1}{3}$, this choice ensures that $\sum_{m'|m^* \ge m_0} I(m^*) \le C/n$. The result follows by taking $p(m,m') = 8(1+2\xi^2)\Delta(m^*)/n$ and

$$\text{pen}(m) = 16a(1+2\xi^2)\frac{\Delta(m)}{n} = 16a\left(1 + \frac{98\mu\lambda_2^\star(f_\varepsilon, \kappa_0)}{\lambda_1(f_\varepsilon, \kappa_0')}\right)\frac{\Delta(m)}{n}.$$

Case $\delta > \frac{1}{3}$. In that case $\delta > (\frac{1}{2} - \delta/2)_+$. According to the inequality (5.40), we choose ξ^2 such that $(K'\xi^2/2)(\pi m)^\omega = (2+\epsilon)\mu(\pi m)^\delta$ for some $\epsilon > 0$, for instance,

$$\xi^2 = \xi^2(m) = \frac{49\mu\lambda_2^\star(f_\varepsilon, \kappa_0)}{\lambda_1(f_\varepsilon, \kappa_0')}(\pi m)^{\min((3\delta/2-1/2)_+, \delta)}.$$

Consequently, we have $\sum_{m'|m^* \ge m_0} I(m^*) \le C/n$. The result follows by choosing $p(m,m') = 8(1+2\xi^2(m,m'))\Delta(m)/n$, associated to

$$\begin{aligned}\text{pen}(m) &= 16a(1+2\xi^2(m))\frac{\Delta(m)}{n} \\ &= 16a\left(1 + \frac{98\mu\lambda_2^\star(f_\varepsilon, \kappa_0)}{\lambda_1(f_\varepsilon, \kappa_0')}(\pi m^*)^{\min((3\delta/2-1/2)_+, \delta)}\right)\frac{\Delta(m)}{n}.\end{aligned}$$

■

5.2. Technical Lemmas

LEMMA 5.1.

$$\left\| \sum_{j \in \mathbb{Z}} |u^*_{\varphi_{m,j}}|^2 \right\|_\infty \le \Delta(m). \tag{5.41}$$

Proof of Lemma 5.1. The proof of Lemma 5.1 can be found in Comte et al. (2006).

LEMMA 5.2. *Assume that $\sum_{k \ge 1} \tau_1(k) < +\infty$. Assume either that $\delta = 0$, $\gamma \ge \frac{3}{2}$ in (3.9) or $\delta > 0$ in (3.9). Then we have*

$$\sup_{t \in B_{m,m'}(0,1)} \|\nu^\star_{q_n,\ell}(u_t^*)\|_\infty \le \sqrt{\Delta(m^*)}. \tag{5.42}$$

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

Moreover, there exist m_2 and m_3 such that

$$\mathbb{E}\left[\sup_{t\in B_{m,m'}(0,1)}|\nu^{\star}_{n,1}(u^*_t)|\right]\le\sqrt{2\Delta(m^*)/n}\quad \textit{for } m^*\ge m_2,$$

and

$$\sup_{t\in B_{m,m'}(0,1)}\mathrm{Var}(\nu^{\star}_{q_n,\ell}(u^*_t))\le C_{\upsilon^*}\sqrt{\Delta_2(m^*,f_Z)}/(2\pi q_n)\quad \textit{for } m^*\ge m_3,$$

where $\Delta(m)$ and $\Delta_2(m,f_Z)$ are defined by (2.5) and (5.29) and where C_{υ^*} is defined in (5.30).

Proof of Lemma 5.2. Arguing as in Lemma 5.1 and by using the Cauchy–Schwarz inequality and the Parseval formula, we obtain that the first term $\sup_{t\in B_{m,m'}(0,1)}\|\nu^{\star}_{q_n,\ell}(u^*_t)\|_\infty$ is bounded by

$$\sup_{t\in B_{m,m'}(0,1)}\|\nu^{\star}_{q_n,\ell}(u^*_t)\|_\infty\le\sqrt{\sum_{j\in\mathbb{Z}}\int\left|\frac{\varphi^*_{m^*,j}(x)}{f^*_\varepsilon(x)}\right|^2dx}=\sqrt{\Delta(m^*)}.$$

Next

$$\mathbb{E}\left[\sup_{t\in B_{m,m'}(0,1)}|\nu^{\star}_{n,1}(u^*_t)|\right]=\mathbb{E}\left[\sup_{t\in B_{m,m'}(0,1)}\left|\frac{1}{p_nq_n}\sum_{\ell=1}^{p_n}\sum_{i=1}^{q_n}u^*_t(Z^{\star}_{2\ell q_n+i})-\langle t,g\rangle\right|\right]$$
$$\le\sqrt{\sum_{j\in\mathbb{Z}}\mathrm{Var}(\nu^{\star}_{n,1}(u^*_{\varphi_{m^*,j}}))}.$$

By using (5.9) we obtain

$$\sqrt{\sum_{j\in\mathbb{Z}}\mathrm{Var}(\nu^{\star}_{n,1}(u^*_{\varphi_{m^*,j}}))}$$
$$=\sqrt{\sum_{j\in\mathbb{Z}}\frac{1}{p_n^2}\sum_{\ell=1}^{p_n}\mathrm{Var}(\nu^{\star}_{q_n,\ell}(u^*_{\varphi_{m^*,j}}))}=\sqrt{\sum_{j\in\mathbb{Z}}\frac{1}{p_n^2}\sum_{\ell=1}^{p_n}\mathrm{Var}(\nu_{q_n,\ell}(u^*_{\varphi_{m^*,j}}))}$$
$$=\sqrt{\sum_{j\in\mathbb{Z}}\frac{1}{p_n}\mathrm{Var}(\nu_{q_n,1}(u^*_{\varphi_{m^*,j}}))}=\sqrt{\frac{1}{2\pi p_n}\int_{-\pi m^*}^{\pi m^*}\frac{\mathbb{E}|\nu_{q_n,1}(e^{ix\cdot})|^2}{|f^*_\varepsilon(x)|^2}dx}.$$

Now, according to (5.12) and (5.2)

$$\mathbb{E}|\nu_{q_n,1}(e^{ix\cdot})|^2\le\frac{1}{q_n}+\frac{1}{q_n}\sum_{k=1}^{n-1}\tau_1(k)|x||f^*_\varepsilon(x)|.$$

This implies that

$$\mathbb{E}^2\left[\sup_{t\in B_{m,m'}(0,1)}|\nu^{\star}_{n,1}(u^*_t)|\right]\le\frac{1}{p_n}\left(\frac{1}{q_n}\Delta(m^*)+\frac{2\pi}{q_n}\sum_{k=1}^{n-1}\tau_1(k)m\Delta_{1/2}(m^*)\right).$$

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

Because $2\pi \sum_{k\geq 1} \tau_1(k) m \Delta_{1/2}(m) \leq \Delta(m)$ for m large enough, we get that for m^* large enough

$$\mathbb{E}^2\left[\sup_{t\in B_{m,m'}(0,1)} |\nu^{\star}_{n,1}(u_t^*)|\right] \leq 2\Delta(m^*)/n.$$

Now, for $t \in B_{m,m'}(0,1)$ we write

$$\begin{aligned}
&\mathrm{Var}\left(\frac{1}{q_n}\sum_{i=1}^{q_n} u_t^*(Z^{\star}_{2\ell q_n+i})\right)\\
&\quad = \mathrm{Var}\left(\frac{1}{q_n}\sum_{i=1}^{q_n} u_t^*(Z_i)\right)\\
&\quad = \frac{1}{q_n^2}\left[\sum_{k=1}^{q_n} \mathrm{Var}(u_t^*(Z_k)) + 2\sum_{1\leq k<l\leq q_n} \mathrm{Cov}(u_t^*(Z_k), u_t^*(Z_l))\right].
\end{aligned}$$

According to (5.8), (5.11), and (5.2), we have

$$\begin{aligned}
|\mathrm{Cov}(u_t^*(Z_k), u_t^*(Z_l))| &= \left|\int_{-\pi m^*}^{\pi m^*}\int_{-\pi m^*}^{\pi m^*} \frac{f_\varepsilon^*(-y)\mathrm{Cov}(e^{ixZ_k}, e^{iyX_l}) t^*(x)t^*(y)}{f_\varepsilon^*(x) f_\varepsilon^*(-y)}\, dxdy\right|\\
&\leq \int_{-\pi m^*}^{\pi m^*}\int_{-\pi m^*}^{\pi m^*} \frac{|y|\tau_1(k)|t^*(x)t^*(y)|}{|f_\varepsilon^*(x)|}\, dxdy.
\end{aligned}$$

Hence,

$$\begin{aligned}
\mathrm{Var}\left(\frac{1}{q_n}\sum_{i=1}^{q_n} u_t^*(Z^{\star}_{2\ell q_n+i})\right) \leq \frac{1}{q_n}\Bigg(&\int_{-\pi m^*}^{\pi m^*}\int_{-\pi m^*}^{\pi m^*} \frac{f_Z^*(u-v)t^*(u)t^*(-v)}{f_\varepsilon(u)f_\varepsilon(-v)}\, dudv\\
&+ 2\sum_{k=1}^{q_n}\tau_1(k)\int_{-\pi m^*}^{\pi m^*}\int_{-\pi m^*}^{\pi m^*}\left|\frac{ut^*(u)t^*(v)}{f_\varepsilon^*(v)}\right| dudv\Bigg).
\end{aligned}$$

Following Comte et al. (2006) and applying Parseval's formula, the first integral is less than $\sqrt{\Delta_2(m^*, f_Z)}/2\pi$. For the second one, write

$$\begin{aligned}
&\int_{-\pi m^*}^{\pi m^*}\int_{-\pi m^*}^{\pi m^*}\left|\frac{ut^*(u)t^*(v)}{f_\varepsilon^*(v)}\right| dudv\\
&\quad \leq \frac{\sqrt{2}\pi^{3/2}}{\sqrt{3}}(m^*)^{3/2}\|t^*\|\sqrt{\int |t^*(v)|^2\, dv \int_{-\pi m^*}^{\pi m^*}\frac{dv}{|f_\varepsilon^*(v)|^2}},
\end{aligned}$$

that is,

$$\int_{-\pi m^*}^{\pi m^*}\int_{-\pi m^*}^{\pi m^*}\left|\frac{t^*(u)t^*(v)}{f_\varepsilon^*(v)}\right| dudv \leq \frac{\sqrt{2}\pi^{3/2}}{\sqrt{3}}(2\pi)^{3/2}\sqrt{(m^*)^3\Delta(m^*)}.$$

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

If $\delta > 0$, then $\sqrt{(m^*)^3\Delta(m^*)} = o_m\sqrt{\Delta_2(m^*, f_Z)}$. If $\gamma > \frac{3}{2}$ and $\delta = 0$, we get that $\sqrt{(m^*)^3\Delta(m^*)} = o_m\sqrt{\Delta_2(m^*, f_Z)}$. Last, if $\gamma = \frac{3}{2}$ and $\delta = 0$, we get that $\sqrt{(m^*)^3\Delta(m^*)} \le \sqrt{\Delta_2(m^*, f_Z)}$, and the result follows for m large enough. ■

LEMMA 5.3. *Let $Y_1, \dots, Y_n$ be independent random variables and let $\mathcal{F}$ be a countable class of uniformly bounded measurable functions. Then for $\xi^2 > 0$*

$$\mathbb{E}\left[\sup_{f\in\mathcal{F}}|\nu_{n,Y}(f)|^2 - 2(1+2\xi^2)H^2\right]_+ \le \frac{2}{K_1}\left(\frac{v}{n}e^{-K_1\xi^2(nH^2/v)} + \frac{49M_1^2}{4K_1n^2C^2(\xi^2)}e^{-(2\sqrt{2}K_1C(\xi^2)\xi/7)(nH/M_1)}\right),$$

with $C(\xi^2) = (\sqrt{1+\xi^2} - 1) \wedge 1$, $K_1 = \frac{1}{6}$, and

$$\sup_{f\in\mathcal{F}}\|f\|_\infty \le M_1, \qquad \mathbb{E}\left[\sup_{f\in\mathcal{F}}|\nu_{n,Y}(f)|\right] \le H, \qquad \sup_{f\in\mathcal{F}}\frac{1}{n}\sum_{k=1}^n \operatorname{Var}(f(Y_k)) \le v.$$

This inequality comes from a concentration inequality in Klein and Rio (2005) and arguments that can be found in Birgé and Massart (1998; see the proof of Cor. 2, p. 354). Usual density arguments show that this result can be applied to the class of functions $\mathcal{F} = B_{m,m'}(0,1)$.

Proof of Proposition 4.1. We describe here a general method to handle the models (4.1) and (4.2) and prove the following result that implies Proposition 4.1 (for related results, see Ango Nzé and Doukhan, 2004; Doukhan et al., 2006).

PROPOSITION 5.1. *Let Y_t and σ_t satisfy either (4.1) or (4.2). For model (4.1), let $(\eta'_t)_{t\in\mathbb{Z}}$ be an independent copy of $(\eta_t)_{t\in\mathbb{Z}}$ and for $t > 0$, let $\sigma_t^* = f(\eta_{t-1}, \dots, \eta_1, \eta'_0, \eta'_{-1}, \dots)$. For model (4.2), let σ_0^* be a copy of σ_0 independent of $(\sigma_0, \eta_t)_{t\in\mathbb{Z}}$ and for $t > 0$ let $\sigma_t^* = f(\sigma_{t-1}^*, \eta_{t-1})$. Let δ_n be a nonincreasing sequence such that*

$$2\mathbb{E}(|\sigma_n^2 - (\sigma_n^*)^2|) \le \delta_n. \tag{5.43}$$

Then

(i) The process $((Y_t^2, \sigma_t^2))_{t\ge0}$ is τ-dependent with $\tau_\infty(n) \le \delta_n$.

(ii) Assume that Y_0^2, σ_0^2 have densities satisfying $\max(f_{\sigma^2}(x), f_{Y^2}(x)) \le C|\ln(x)|^\alpha x^{-\rho}$ in a neighborhood of zero, for some $\alpha \ge 0$ and $0 \le \rho < 1$. The process $((X_t, Z_t))_{t\ge0}$ is τ-dependent with $\tau_\infty(n) = O((\delta_n)^{(1-\rho)/(2-\rho)}|\ln(\delta_n)|^{(1+\alpha)/(2-\rho)})$.

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

Consider model (4.4) with $\mathbb{E}(\eta_0^2) = 1$ *and assume that* $c = \sum_{j\geq 1} a_j < 1$. *Then let* $((Y_t, \sigma_t))_{t\in\mathbb{Z}}$ *be the unique strictly stationary solution of the form (4.1). Then (5.43) holds with*

$$\delta_n = O\left(\inf_{1\leq k\leq n}\left\{c^{n/k} + \sum_{i=k+1}^{\infty} a_i\right\}\right).$$

Let us first explain how Proposition 5.1 implies Proposition 4.1. First, if σ_0^2 and η_0^2 have bounded densities, then $f_{Y^2}(x) \leq C|\ln(x)|$ in a neighborhood of zero, so that Proposition 4.1(ii) holds with $\rho = 0$ and $\alpha = 1$.

Under the assumptions of Proposition 5.1(ii), we obtain straightforwardly the rates given in the first two cases for model (4.4) and in the third case, the general rate $\tau_\infty(n) = O(n^{-b(1-\rho)/(2-\rho)}(\ln(n))^{(b+2)(1+\alpha)/2})$. Taking here $\rho = 0$ and $\alpha = 1$ gives the result.

For model (4.2), if there exists $\kappa < 1$ such that (4.8) is satisfied, then one can take $\delta_n = 4\mathbb{E}(\sigma_0^2)\kappa^n$. Hence, under the assumptions of Proposition 4.1(ii), $((X_t, Z_t))_{t>0}$ is geometrically τ dependent, and substituting δ_n gives the order of $\tau_\infty(n)$.

Proof of Proposition 5.1. To prove (i), let for $t > 0$, $Y_t^* = \eta_t\sigma_t^*$. Note that the sequence $((Y_t^*, \sigma_t^*))_{t\geq 1}$ is distributed as $((Y_t, \sigma_t))_{t\geq 1}$ and independent of $\mathcal{M}_i = \sigma(\sigma_j, Y_j, 0 \leq j \leq i)$. Hence, by the coupling properties of τ (see eqn. (5.1)), we have that, for $n + i \leq i_1 < \cdots < i_l$,

$$\tau(\mathcal{M}_i, (Y_{i_1}^2, \sigma_{i_1}^2), \ldots, (Y_{i_l}^2, \sigma_{i_l}^2)) \leq \frac{1}{l}\sum_{j=1}^{l} \|(Y_{i_j}^2, \sigma_{i_j}^2) - ((Y_{i_j}^*)^2, (\sigma_{i_j}^*))^2\|_{\mathbb{R}^2} \leq \delta_n,$$

and (i) follows.

To prove (ii), define the function $f_\epsilon(x) = \ln(x)\mathbb{1}_{x>\epsilon} + 2\ln(\epsilon)\mathbb{1}_{x\leq\epsilon}$ and the function $g_\epsilon(x) = \ln(x) - f_\epsilon(x)$. Clearly, for any $\epsilon > 0$ and any $n + i \leq i_1 < \cdots < i_l$, we have

$$\begin{aligned}\tau(\mathcal{M}_i, (Z_{i_1}, X_{i_1}), \ldots, (Z_{i_l}, X_{i_l}))&\\ \leq 2\mathbb{E}(|g_\epsilon(Y_0^2)| + |g_\epsilon(\sigma_0^2)|)&\\ + \tau(\mathcal{M}_i, (f_\epsilon(Y_{i_1}^2), f_\epsilon(\sigma_{i_1}^2)), \ldots, (f_\epsilon(Y_{i_l}^2), f_\epsilon(\sigma_{i_l}^2))).& \quad \textbf{(5.44)}\end{aligned}$$

For $0 < \epsilon < 1$, the function f_ϵ is $1/\epsilon$-Lipschitz. Hence, applying (i),

$$\tau(\mathcal{M}_i, (f_\epsilon(Y_{i_1}^2), f_\epsilon(\sigma_{i_1}^2)), \ldots, (f_\epsilon(Y_{i_l}^2), f_\epsilon(\sigma_{i_l}^2))) \leq \frac{\delta_n}{\epsilon}.$$

Because $\max(f_{\sigma^2}(x), f_{Y^2}(x)) \leq C|\ln(x)|^\alpha x^{-\rho}$ in a neighborhood of 0, we infer that for small enough ϵ,

$$\mathbb{E}(|g_\epsilon(Y_0^2)| + |g_\epsilon(\sigma_0^2)|) \leq K_1\epsilon^{1-\rho}|\ln(\epsilon)|^{1+\alpha},$$

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

for K_1 a positive constant. From (5.44), we infer that there exists a positive constant K_2 such that, for small enough ϵ,

$$\tau(\mathcal{M}_i,(Z_{i_1},X_{i_1}),\ldots,(Z_{i_l},X_{i_l})) \le K_2\left(\frac{\delta_n}{\epsilon}+\epsilon^{1-\rho}|\ln(\epsilon)|^{1+\alpha}\right).$$

The result follows by taking $\epsilon = (\delta_n)^{1/(2-\rho)}|\ln(\delta_n)|^{-(1+\alpha)/(2-\rho)}$.

Now, we go back to the model (4.4). If $\sum_{j=1}^{\infty} a_j < 1$, the unique stationary solution to (4.4) is given by Giraitis et al. (2000):

$$\sigma_t^2 = a + a\sum_{\ell=1}^{\infty}\sum_{j_1,\ldots,j_l=1}^{\infty} a_{j_1}\ldots a_{j_l}\eta_{t-j_1}^2\ldots\eta_{t-(j_1+\cdots+j_l)}^2.$$

For any $1 \le k \le n$, let

$$\sigma_t^2(k,n) = a + a\sum_{\ell=1}^{[n/k]}\sum_{j_1,\ldots,j_l=1}^{k} a_{j_1}\ldots a_{j_l}\eta_{t-j_1}^2\ldots\eta_{t-(j_1+\cdots+j_l)}^2.$$

Clearly $\mathbb{E}(|\sigma_n^2-(\sigma_n^*)^2|) \le 2\mathbb{E}(|\sigma_0^2-\sigma_0^2(k,n)|)$. Now

$$\mathbb{E}(|\sigma_0^2-\sigma_0^2(k,n)|) \le \left(\sum_{l=[n/k]+1}^{\infty} c^l + \sum_{l=1}^{\infty} c^{l-1}\sum_{j>k} a_j\right).$$

This being true for any $1 \le k \le n$, the proof of Proposition 5.1 is complete. ■

REFERENCES

Ango Nzé, P. (1992) Critères d'ergodicité de quelques modèles à représentation markovienne (Criteria of ergodicity for some models with a Markovian representation). *Comptes Rendus de l'Academie des Sciences de Paris Série I Mathematiques* 315, 1301–1304.

Ango Nzé, P. & P. Doukhan (2004) Weak dependence: Models and applications to econometrics. *Econometric Theory* 20, 995–1045.

Birgé, L. & P. Massart (1998) Minimum contrast estimators on sieves: Exponential bounds and rates of convergence. *Bernoulli* 4, 329–375.

Birgé, L. & Y. Rozenholc (2006) How many bins should be put in a regular histogram. *ESAIM Probability and Statistics* 10, 24–45 (electronic).

Bollerslev, T. (1986) Generalized autoregressive conditional heteroskedasticity. *Journal of Econometrics* 31, 307–327.

Bougerol, P. & N. Picard (1992a) Stationarity of GARCH processes and of some nonnegative time series. *Journal of Econometrics* 52, 115–127.

Bougerol, P. & N. Picard (1992b) Strict stationarity of generalized autoregressive processes. *Annals of Probability* 20, 1714–1730.

Butucea, C. (2004) Deconvolution of supersmooth densities with smooth noise. *Canadian Journal of Statistics* 32, 181–192.

Butucea, C. & C. Matias (2005) Minimax estimation of the noise level and of the deconvolution density in a semiparametric convolution model. *Bernoulli* 11, 309–340.

Butucea, C. & A.B. Tsybakov (2008a) Sharp optimality in density deconvolution with dominating bias. I. *Theory of Probability and Its Applications* 52, 24–39.

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press

Butucea, C. & A.B. Tsybakov (2008b) Sharp optimality in density deconvolution with dominating bias. II. *Theory of Probability and Its Applications* 52, 237–249.

Carrasco, M. & X. Chen (2000) β-mixing and moment properties of RCA models with application to GARCH(p,q). *Comptes Rendus de l'Academie des Sciences de Paris Série I Mathématiques* 331, 85–90.

Carrasco, M. & X. Chen (2002) Mixing and moment properties of various GARCH and stochastic volatility models. *Econometric Theory* 18, 17–39.

Comte, F., Y. Rozenholc, & M.-L. Taupin (2006) Penalized contrast estimator for adaptive density deconvolution. *Canadian Journal of Statistics* 34, 431–452.

Comte, F., Y. Rozenholc, & M.-L. Taupin (2007) Finite sample penalization in adaptive density deconvolution. *Journal of Statistical Computation and Simulation* 77, 977–1000.

de Vries, C.G. (1991) On the relation between GARCH and stable processes. *Journal of Econometrics* 48, 313–324.

Dedecker, J. & C. Prieur (2005) New dependence coefficients. Examples and applications to statistics. *Probability Theory and Related Fields* 132, 203–236.

Doukhan, P. (1994) *Mixing: Properties and Examples*. Lecture Notes in Statistics. Springer-Verlag.

Doukhan, P., G. Teyssière, & P. Winant (2006) A LARCH(∞) vector valued process. *Lecture Notes in Statistics* 187, 245–258.

Duan, J.-C. (1997) Augmented GARCH(p,q) process and its diffusion limit. *Journal of Econometrics* 79, 97–127.

Engle, R.F. (1982) Autoregressive conditional heteroscedasticity with estimates of the variance of United Kingdom inflation. *Econometrica* 50, 987–1007.

Fan, J. (1991) On the optimal rates of convergence for nonparametric deconvolution problems. *Annals of Statistics* 19, 1257–1272.

Giraitis, L., P. Kokoszka, & R. Leipus (2000) Stationary ARCH models: Dependence structure and central limit theorem. *Econometric Theory* 16, 3–22.

Klein, T. & E. Rio (2005) Concentration around the mean for maxima of empirical processes. *Annals of Probability* 33, 1060–1077.

Lacour, C. (2006) Rates of convergence for nonparametric deconvolution. *Comptes Rendus de l'Academie des Sciences de Paris Série I Mathematik* 342, 877–882.

Lacour, C. (2008) Adaptive estimation of the transition density of a particular hidden Markov chain. *Journal of Multivariate Analysis* 99, 784–814.

Lee, O. & D.W. Shin (2005) On stationarity and β-mixing property of certain nonlinear GARCH(p,q) models. *Statistics and Probability Letters* 73, 25–35.

Meyer, Y. (1990) *Ondelettes et opérateurs I: Ondelettes*. Actualités Mathématiques. Hermann, éditeurs des Sciences et des Arts.

Pensky, M. & B. Vidakovic (1999) Adaptive wavelet estimator for nonparametric density deconvolution. *Annals of Statistics* 27, 2033–2053.

van Es, B., P. Spreij, & H. van Zanten (2005) Nonparametric volatility density estimation for discrete time models. *Journal of Nonparametric Statistics* 17, 237–251.

Zakoïan, J.-M. (1993) Modèle autoregressif à un seuil. *Publications de l'Institut de Statistique de l'Université de Paris* 37, 85–113.

https://doi.org/10.1017/S026646660808064X Published online by Cambridge University Press