Animal (2012), **6:2**, pp 179–192 © The Animal Consortium 2011
doi:10.1017/S1751731111001637

Gene expression in Sinclair swine with malignant melanoma

M. Okomo-Adhiambo[1], A. Rink[1,2], W. M. Rauw[1,3] and L. Gomez-Raya[1,3†]

[1]*Department of Animal Biotechnology, University of Nevada, 1664 North Virginia Street, 89557 Reno, NV, USA;* [2]*Nevada Department of Agriculture, Animal Disease and Food Safety Laboratory, 350 Capitol Hill Avenue, 89502 Reno, NV, USA;* [3]*Departamento de Mejora Genética Animal, Instituto Nacional de Investigación y Tecnología Agraria y Alimentaria (INIA), Carretera de La Coruña km 7.5, 28040 Madrid, Spain*

(Received 18 February 2011; Accepted 24 June 2011; First published online 22 September 2011)

Sinclair swine develop an aggressive form of melanoma, which, in many cases, spontaneously regresses after a complete metastatic phase. We used Affymetrix GeneChip® Porcine Genome Arrays consisting of 24 123 probe sets to compare gene expression in white blood cells (WBCs) and various tissues including the liver, lungs, inguinal lymph nodes and spleen harvested from a Sinclair piglet afflicted by melanoma at birth and exhibiting metastatic lesions at weaning (6 weeks) with those from a full-sibling piglet that showed no incidence of melanoma at birth and weaning. The highest number (3489; ~14%) of significantly upregulated transcripts (fold change in gene expression ⩾2.0 and t*-test* P*-value ⩽0.05) was observed in the liver, while the spleen exhibited the lowest number of upregulated transcripts (528; ~2%). Among significantly downregulated genes, the highest numbers were observed in the inguinal lymph nodes (3651; ~15%) and the least in WBCs (730; ~3%). Differentially expressed transcripts included genes involved in melanoma pathogenesis including* SILV, TYR *and* RAB28. SILV *was over-expressed 784-, 430- and 164-fold, while* TYR *was over-expressed 138-, 81- and 28-fold in the liver, lungs and inguinal lymph nodes, respectively. Quantitative real-time RT-PCR (qRT-PCR) confirmed the microarray data of 12 selected differentially expressed sequences. These results suggest that significant changes in gene expression occur during metastasis of malignant melanoma in the Sinclair swine model. In addition, qRT-PCR analysis of the above 12 differentially expressed sequences was carried out on liver samples collected from 22 pigs (12 of which had melanoma during the first 6 weeks of life), and an ANOVA test contrasting absolute RNA expression between pigs with regressing, progressing and without tumors was significant for TYR, TACSTD1, MATP, GPNMB and CYP4A22, with* P*-values of 0.034, 0.015, 0.007, 0.050 and 0.022, respectively.*

Keywords: melanoma, metastasis, gene expression, Sinclair swine

Implications

This study provides new information on changes in the patterns of gene expression during the progression of malignant melanoma and its regression in Sinclair swine.

Introduction

The Sinclair Miniature Pig Melanoma model was established in the 1970s by breeding melanoma-bearing pigs to establish a line with up to 70% incidence of melanoma (Hook *et al*., 1979; Tissot *et al*., 1987). Melanoma in Sinclair swine usually develops *in utero* or within the first 6 weeks of life (Gomez-Raya *et al*., 2007), with lesions varying from benign flat nevi to large, deeply invasive, exophytic tumors possessing characteristics of malignant tumors (Das Gupta *et al*., 1989) as well as some *in vitro* characteristics of benign neoplastic cells found in childhood tumors such as retinoblastoma or neuroblastoma that are of neural or neural crest tissue origin as are melanocytes. Cutaneous malignant melanoma in Sinclair swine may have multiple primary sites and usually disseminates in the lymph nodes and visceral organs (Das Gupta *et al*., 1989; Greene *et al*., 1994; Greene *et al*., 1997). Some pigs may die because of widespread disease but a majority develop a cell-mediated immune response that causes complete regression of the tumors (Misfeldt and Grimm, 1994; Greene *et al*., 1997), although the immune response also attacks melanocytes in the skin, resulting in varying degrees of vitiligo that may progress to complete depigmentation (Misfeldt and Grimm, 1994).

Gene expression analysis using DNA microarrays allows the simultaneous monitoring of differential gene expression for thousands of genes in biological samples (Eisen and Brown, 1999). Microarray gene expression analysis provides a fresh insight into the genetic basis of cancer initiation and progression (Luo *et al*., 2001; Hoek *et al*., 2004; Talantov *et al*., 2005) and

† E-mail: gomez.luis@inia.es

https://doi.org/10.1017/S1751731111001637 Published online by Cambridge University Press

identification of genes that may be of prognostic value (Wang *et al*., 2004; Barrier *et al*., 2005) as well as genes that may be associated with aggressive tumor behavior (Bittner *et al*., 2000). In addition, microarray gene expression profiling has shown promise in evaluating the sensitivity of tumors to chemotherapy or other targeted approaches to therapy (Huang and Sadee, 2003; Mariadason *et al*., 2003; Alaoui-Jamali *et al*., 2004; Clarke *et al*., 2004). There are no data on gene expression changes in melanoma regression. However, Sinclair swine is a unique model for investigating molecular changes during regression, given their high frequency of cancer regression.

This study hypothesizes that genes are differentially expressed in Sinclair swine with melanoma compared with those not afflicted by the disease. The study also hypothesizes that genes are differentially expressed in melanoma *v.* non-melanoma pigs during tumor progression and regression. The main objective of this study was to investigate the expression profiles of genes that are modulated during development and metastasis of malignant melanoma in Sinclair swine at 6 weeks using an Affymetrix GeneChip® Porcine Genome Array (Affymetrix Inc., Santa Clara, CA, USA) and quantitative real-time RT-PCR (qRT-PCR), with the goal of improving the understanding of melanoma progression at molecular and gene-expression levels. This study also investigates gene expression at later ages, in an effort to gain further knowledge of the molecular changes that occur during the process of tumor progression and regression in Sinclair swine.

Material and methods

Animals

The Sinclair melanoma swine were derived from a herd maintained at the University of Nevada, Reno Agricultural Experiment Station. They were maintained under animal care and use protocols of the University of Nevada, Reno. The experimental piglets, one female afflicted with melanoma at birth, and its full female sibling that showed no incidence of melanoma at birth and at weaning, were sacrificed at 6 weeks (weaning age). Before euthanasia, the live piglets were bled by jugular venipuncture using a 20-gauge needle. A volume of 5 ml peripheral whole blood was collected in vacutainer tubes containing EDTA anticoagulant (Becton Dickinson Biosciences; Franklin Lakes, NJ, USA). Blood samples were spun in a centrifuge at 3000 × **g** for 15 min to separate buffy coats containing white blood cells (WBCs), which were stored in liquid nitrogen.

Euthanasia was conducted using sodium pentobarbital at a dosage of 1 ml/10 lb body weight. The *post-mortem* examination showed that the piglet afflicted by melanoma at birth had numerous metastatic lesions infiltrated into the internal visceral organs and lymph nodes (Figure 1 and Table 1) while the other piglet had no evident metastatic tumors. For the melanoma-afflicted pig, samples were collected from the lymph nodes and internal visceral organs that were infiltrated by tumors including the liver, lungs, spleen, kidneys, pancreas, stomach, small intestines and colon (Table 1). Only sections of the melanoma-afflicted tissues that exhibited almost 100% melanoma lesions were collected to minimize the presence of non-melanoma tissues in the sample. For the pig without melanoma, samples were obtained from equivalent lymph nodes and internal visceral organs, respectively. The harvested tissues were stored in liquid nitrogen in readiness for RNA extraction.



Figure 1 Sinclair piglet at 6 weeks of age exhibiting metastatic melanoma lesions in internal visceral organs.

Of note, skin samples with or without tumors were also collected from both piglets, respectively, but the quality of RNA extracted from samples with tumors was very low, because of the presence of high amounts of dead tissue. Consequently, skin samples were not included in the microarray and qRT-PCR analysis in this study.

RNA isolation

Total RNA was isolated from lungs, liver, inguinal lymph nodes, spleen and WBCs harvested from melanoma-afflicted and normal control pigs, respectively, using the Qiagen RNeasy mini kit (Qiagen Inc., Valencia, CA, USA) according to the manufacturer's instructions. RNA quantity and quality were determined by capillary electrophoresis using the Agilent 2100 Bioanalyzer (Agilent Technologies; Santa Clara, CA, USA) according to the manufacturer's instructions.

Microarray hybridization and data analysis

Labeled cDNA was prepared from total RNA and hybridized with the high-density oligonucleotide Affymetrix GeneChip® Porcine Genome Array (Affymetrix Inc.) consisting of 24 123 probe sets, according to standard manufacturer protocols. Arrays were scanned using Affymetrix protocols and scanners. For subsequent analysis, each of the ten hybridized arrays (five with melanoma and five with normal cDNA, respectively) was considered as an experimental sample, and each probe set on the array was considered as a separate gene. Microarray data acquisition and image analysis was performed using Affymetrix GeneChip® Operating Software (GCOS). GeneSpring 7.0 software (Agilent Technologies; Santa Clara, CA, USA) was used to normalize and analyze the GCOS output data to provide expression values for each gene. Gene names and accession numbers were updated according to the

https://doi.org/10.1017/S1751731111001637 Published online by Cambridge University Press

Table 1 *Characteristics of tumor samples collected from Sinclair swine*

				External skin tumors[a]			Internal tumors[b]	
Pig ID	Age	Sex	Weight (kg)	Location	Diameter (cm)	No. of tumors	Description	Location	Description
305039	6 weeks	Female	3.77	Above left eye	2.5	1	Elevated above skin	Inguinal lymph nodes	Complete infiltration of entire tissue with tumor
				Root of tail	1.0	1	Elevated above skin	Liver, lungs, spleen, kidney, pancreas	Numerous areas infiltrated by tumors; few areas unaffected
				Right inguinal skin	1.0	1	Elevated above skin	Stomach, small intestine, spiral colon	Numerous flat black patches of tumors, non-elevated; few areas unaffected
				Left knee	0.5	1	Elevated above skin		
				Right shoulder	1.0	1			
				Various locations on back, sides and abdomen	–	20	Non-elevated		
305040	6 weeks	Female	5.77	No tumors visible on the skin	–	–	–	No tumors visible in any internal organ	

Note: Peripheral blood was also collected from both pigs. Buffy coats containing white blood cells were separated and stored in liquid nitrogen along with other sampled tissues.
[a]For the melanoma-afflicted pig, skin samples were collected from areas with visible non-elevated tumors and consisted of biopsies measuring about 4 cm^2 each. For the normal pig, samples were taken from equivalent areas of the skin with no tumors. The skin samples were not analyzed in this study.
[b]Samples collected from internal organs of the melanoma-afflicted pig were taken from areas containing almost 100% visible tumors. For the normal pig, samples were taken from equivalent organs showing no tumors.

annotated version of the Affymetrix Porcine GeneChip® (Tsai *et al.*, 2006). Genes with expression values ⩾2.0 (upregulated ⩾2.0-fold) or expression values ⩽0.5 (downregulated ⩾2.0-fold) in at least one out of the 10 experimental samples were selected for further analysis, and a total of 20 177 genes (~84%) fulfilled this criterion. Genes selected for further analysis were classified into functional subgroups using Simplified Ontology, a feature of GeneSpring.

qRT-PCR

qRT-PCR was performed to validate the microarray data of 12 selected genes that were differentially expressed (up- or downregulated) in the melanoma compared with normal samples (Table 3). Primers and probes were designed using the Primer Express 2.0 software (Applied Biosystems [ABI]; Foster City, CA, USA). Reverse transcription (RT) was performed with 1 μg total RNA isolated from lungs, liver, inguinal lymph nodes, spleen and WBCs harvested from melanoma-afflicted and normal control pigs, using 300 ng/μl random primers and Superscript II Reverse Transcriptase™ (Invitrogen; Carlsbad, CA, USA) according to the manufacturer's instructions. qRT-PCR was performed using the 5′ Nuclease Taqman Assay (ABI) according to the manufacturer's instructions. In brief, 1 μl of cDNA obtained from the RT reaction were amplified in a 25 μl containing 2 × TaqMan® Universal PCR Mastermix (ABI; Foster City, CA, USA), 900 nmol/l of forward and reverse primers and 250 nmol/l of TaqMan hybridization probe. For each sampled time point, the target gene was amplified concurrently with the glyceraldehyde-3-phosphate dehydrogenase (GAPDH) control gene, in separate wells. Non-template controls were included in every run. All reactions were performed in triplicate. Thermal cycling was performed in an ABI PRISM 7000 Sequence Detection System unit (ABI; Foster City, CA, USA).

qRT-PCR data, comprising threshold value (Ct) values, were analyzed using the Relative Expression Software Tool (REST©; Pfaffl *et al.*, 2002), which uses a mathematical model based on the PCR efficiencies and the mean Ct deviation between the sample and the control group, where the ratio $R = E_{\text{target}}^{\Delta \text{Ct target (control}-\text{sample)}} / E_{\text{ref}}^{\Delta \text{Ct ref (control}-\text{sample)}}$. The quantification was carried out relative to a non-regulated reference housekeeping control gene (GAPDH). Differences in expression between the control and the treated samples for the investigated transcripts were assessed in group means for statistical significance using the Pair-Wise Fixed Reallocation Randomization Test© running within REST© (Pfaffl *et al.*, 2002).

Gene expression during tumor progression and regression

To analyze gene expression during tumor regression, additional frozen liver samples were obtained from the historical Sinclair herd sample collection, from 22 pigs (seven, five and ten with regressing, progressing and without melanoma, respectively). The Sinclair swine herd was established and maintained in the 1980s at Texas A & M University as a resource population for biomedical research. Records including the number of tumors, location and size were documented for

https://doi.org/10.1017/S1751731111001637 Published online by Cambridge University Press

each pig. A set of measurements were used to monitor tumor status of each piglet over time including length, width and height that were measured using a caliper. The recording of tumors was carried out weekly. Tissue samples (liver, tail, etc.) from culled animals were systematically collected and stored at −80°C for most pigs and information on those pigs was documented in a database. More information on this population and historical records can be found in Gomez-Raya *et al*. (2009). In this study, length was used as a criterion for estimating tumor size based on James *et al*. (1999), who concluded that unidimensional measurements for tumors are appropriate for estimating regression in solid tumors. Pigs whose tumors had reached the maximum length at collection time were considered as progressing animals. Pigs with tumors that were reduced in length (compared with the maximum registered length) at collection time were categorized as regressing animals. The average change for all tumors within each pig was used to assign piglets into progressing or regressing animals (Table 2). More information on the 22 liver samples used in the analyses of gene expression is given in Table 2. The frozen samples were used in qRT-PCR analysis of 12 genes that were upregulated during metastasis. A one-way ANOVA test was performed to test whether gene expression in piglets with regressing, progressing or without tumors was different.

For this tumor regression study, the same protocols were followed for RNA extraction from liver frozen samples obtained from the additional 22 pigs described above. qRT-PCR was performed as described earlier for the 12 selected genes (Table 3) that were differentially expressed in the melanoma compared with normal samples. qRT-PCR data comprising cDNA (ng) quantified for each gene were analyzed by one-way ANOVA (Minitab 15 software) contrasting absolute RNA expression for the 12 genes in the animals with melanoma (regression and progression) and without the disease.

Results

Differential gene expression in melanoma compared with normal samples

Microarray analysis compared the gene expression profiles of samples (lungs, liver, inguinal lymph nodes, spleen and WBCs) obtained from melanoma-afflicted and normal control piglets born from the same litter of Sinclair swine, at weaning. For the

Table 2 *Information of frozen samples obtained from pigs with regressing, progressing and without tumors*

Pig ID	Sex	Age at sample collection†	Age at first tumor†	No. of tumors recorded	
Regressing tumors					Average tumor reduction in length (cm)#
193047	M	134	0	15	−0.28
195113	F	145	21	1	−1.35
195136	F	95	0	4	−0.13
195166	F	85	0	2	−0.20
195208	F	164	0	9	−0.16
196009	M	156	7	3	−0.37
196026	F	153	7	5	−0.06
Progressing tumors					Maximum average tumor length (cm)
195209	F	164	7	5	2.23
195211	F	164	0	2	2.58
196007	M	152	28	1	2.10
196028	F	139	21	2	1.75
200093	M	61	42	2	1.56
Without tumors					
193122	F	1047		–	–
195123	F	135		–	–
195124	F	135		–	–
195134	M	143		–	–
195145	F	90		–	–
196001	M	−10		–	–
196002	F	−10		–	–
196006	M	210		–	–
203124	M	70		–	–
200133	F	129		–	–
196002	F	−10		–	–

†Age in days.
#Reduction in tumor length from the maximum recorded till the last recorded measurement.

https://doi.org/10.1017/S1751731111001637 Published online by Cambridge University Press

https://doi.org/10.1017/S1751731111001637 Published online by Cambridge University Press

Table 3 *Primer and probe sequences for quantitative real-time reverse transcription PCR*

Accession[a]	Gene name	Description	Forward primer (5′ to 3′)	Reverse primer (5′ to 3′)	Taqman probe (5′ to 3′)
CF180835	*CYP4A22*	Cytochrome P450 4B1	ACTGCTGCCCGATGCA	GCGCTTTTCCTTGGAGAATATGG	6FAM-CCACCCCTTTGCCTTC-MGBNFQ
CO949684	*GPNMB*	Transmembrane glycoprotein NMB	TGAGGAAAGCAGTTGGAAAACCA	TCAGAGGGTTGACATTCTTGTGTAAAT	6FAM-CTGGCACACATATTTG-MGBNFQ
CF359325	*KAI1*	CD82 antigen /suppressor of tumorigenicity-6	CCTGCTTCCCATGGTTTTGG	GGCACCCACTGATCTTTCCT	6FAM-TCGGCTCACAATCTG-MGBNFQ
AY604429.1	*LGALS1*	Galectin-1	CACGGAGACATTAACACCATCGT	CACCTCCACGACACTTCCA	6FAM-CCCCGCCGTCCTTG-MGBNFQ
BI182740	*LTBP4*	Latent transforming growth factor beta binding protein 4	AGGGCCCTGCCTAGGT	CCTGCCAGCCACAGTCT	6FAM-CACCAGCCCCGCCCCG-MGBNFQ
BX917105	*MATP*	Membrane-associated transporter protein	GTGATGTCCAGCACCCTGTA	GCGGTGGTACATGGCAATG	6FAM-CACCGTGCCCTTTAAC-MGBNFQ
BX915975	*MCAM*	Melanoma cell adhesion molecule MCAM	CATCCCAAGAAGGACCTCACATG	TGGTGGACAGGTCCCTGA	6FAM-CCTGGAGGCCCTCTCT-MGBNFQ
CN157469	*MRPL28*	Melanoma antigen p15	CCATTCCTCTGTTCAAGATCTTCGT	GGCCTGCTGCTGAAGCT	6FAM-CCGAGCAGCTCCTCC-MGBNFQ
CO955380	*SILV*	Melanocyte protein MEL17	GTCCACCTGGGTAAAGCATGA	GTGGGCAGCAGGTCTGA	6FAM-CAGCACGTGAGCCTAC-MGBNFQ
NM_214023.1	*SPP1*	Secreted phosphoprotein-I	CCCAAGGCCATCCTCGTT	GTCTCCTGACTGTCCTTCTCTTG	6FAM-CCCAGCGCCTGCACG-MGBNFQ
NM_214419.1	*TACSTD1*	Tumor-associated calcium signal transducer 1	CCAAAAGGATGGACCTGAGAGTAAA	CCTGCATTGAAAATTCAGGTGGTTT	6FAM-ATCCTGGTCAAACTTC-MGBNFQ
CN154304	*TYR*	Tyrosinase	TTTCCATAAGGAGTGGCTGCTTT	GCGGTTATGTTGTCGTCGAAAG	6FAM-CAGGAAGCCGCTTTCT-MGBNFQ
P00355	*GAPDH*	Glyceraldehyde-3-phosphate dehydrogenase	GTGACACTCACTCTTCTACCTTTGATG	GGTCCACCACCCTGTTGCT	6FAM-CAAGCTCATTTCCTGGTAC-MGBNFQ

6FAM = 6-carboxyfluorescein; MGBNFQ = minor groove binding non fluorescent quencher.
[a]GenBank database.

melanoma-afflicted piglet, metastatic lesions were visually observed in lungs, liver, inguinal lymph nodes and spleen from which samples were obtained (Figure 1). For the purpose of discussion, melanoma samples refer to those obtained from the melanoma-afflicted pig, while normal samples refer to those derived from the pig without melanoma.




Figure 2 Hierarchical clustering of 7012 genes that were modulated (up- or downregulated) ⩾2-fold in at least one out of the ten experimental samples after microarray analysis. Each column represents a sample and each row is a gene. The condition gene tree shows induced genes (red) and repressed genes (green), while genes whose expression did not change are black. The color code for melanoma samples is red and normal samples are yellow.

After microarray gene expression analysis, genes with expression values ⩾2.0 (upregulated ⩾2.0-fold) or expression values ⩽0.5 (downregulated ⩾2.0-fold) in at least one out of the 10 experimental samples (hybridized arrays) were selected for further analysis, with a total of 20 177 genes out of the original 24 123 (~ 84%) fulfilling the criterion.

Out of the originally selected 20 177 genes, a total of 7012 genes (displaying ⩾2.0-fold induction in a particular tissue and also ⩾2.0-fold repression in another tissue) were hierarchically clustered by Pearson's correlation using GeneSpring 7.0 software (Figure 2). Hierarchical clustering of gene expression values for the 7012 genes revealed two discrete gene modulation patterns: induced genes (red) and downregulated genes (green), with correlated expression of genes involved in diverse functions and cellular pathways. Hierarchical clustering also revealed a distinct separation between gene expression values for respective samples obtained from the melanoma-afflicted and non-melanoma pigs, with the exception of WBCs and spleen, where the least separation and variation of gene expression values were observed (Figure 2). The liver and inguinal lymph nodes exhibited the greatest separation and variation of gene expression values between melanoma and the non-melanoma pigs, followed by the lungs.

To determine differential gene expression ratios, the gene expression values for samples obtained from the melanoma-afflicted pig were divided by gene expression values for the corresponding samples derived from the animal without melanoma, for each of the selected 20 177 genes. Genes with differential gene expression ratios ⩾2.0 (upregulated by ⩾2.0-fold) or expression values ⩽0.5 (downregulated by ⩾2.0-fold) were considered as significantly induced or repressed in melanoma compared with normal samples. The liver exhibited the highest number of differentially expressed genes (3489) that were significantly upregulated ⩾2.0-fold in melanoma compared with normal samples (Table 4), while the spleen exhibited the least (528). Of the significantly downregulated genes, the inguinal lymph node exhibited the

Table 4 *Distribution of differentially genes modulated by >2-fold in melanoma compared with normal samples*

Tissue/organ[a]	Number of genes modulated by >2-fold in melanoma compared with normal samples		
	Induced genes[b]	Repressed genes[c]	Total modulated genes[d]
WBCs	1025	730	1755
Liver	3489	2428	5917
Lung	1608	1137	2745
Inguinal lymph node	2615	3651	6266
Spleen	528	1996	2524

WBCs = white blood cells.
[a]Source of total RNA for microarray hybridization.
[b]Significantly upregulated ⩾2-fold (gene expression ratio ⩾2.0; *t*-test *P*-value ⩽0.05).
[c]Significantly downregulated ⩾2-fold (gene expression ratio ⩽0.5; *t*-test *P*-value ⩽0.05).
[d]Sum of up- and downregulated genes.

https://doi.org/10.1017/S1751731111001637 Published online by Cambridge University Press

highest number of differentially expressed genes (3651), while WBCs exhibited the least number (730). The sample-specific variation in differential gene expression between the melanoma-afflicted and normal piglets was consistent with that observed after hierarchical clustering.

Diverse functional categories of genes are modulated during melanoma progression

Transcripts representing several functional categories were significantly modulated in melanoma compared with normal samples (Table 5). The modulated transcripts included genes directly involved in melanoma development and progression, as well as genes generally associated with cancer pathogenesis including transcription factors, signal transducers, cell cycle regulators and apoptosis-related genes, as well as genes involved in the immune response and energy metabolism (Table 5). These findings suggest that numerous cellular processes were transcriptionally altered during melanoma progression, with the majority of gene modulations (up- and downregulation) occurring in the liver and inguinal lymph nodes. Table 5 shows a partial list of genes significantly modulated in melanoma compared with normal samples; a complete list of these functionally classified genes can be found in the supplemental information.

A number of genes directly associated with cancer pathogenesis were differentially expressed in melanoma compared with normal samples including *SILV*, *TYR* and *RAB28* (Table 5). The *SILV* gene, which encodes melanocyte protein PMEL17, was the most highly differentially expressed gene in melanoma compared with normal samples, and was over-expressed 784-fold in the liver, 430-fold in the lungs and 164-fold in the inguinal lymph nodes, but remained unchanged in the WBCs and spleen samples (Table 5). qRT-PCR analysis (Table 6) confirmed the induced expression of *SILV* in the liver, lungs and inguinal lymph nodes, as well as its unaltered expression in the WBCs and spleen. The *TYR* gene which encodes tyrosinase protein, a copper-containing oxidase enzyme that catalyzes multiple steps in melanin biosynthesis in melanocytes and melanoma cells (Tripathi *et al.*, 1992; Battyani *et al.*, 1993; Kwon, 1993; Perez *et al.*, 2000), was over-expressed by 138-fold in the liver, by 81-fold in the lungs and 28-fold in the inguinal lymph nodes after microarray analysis (Table 5), and similar to *SILV*, remained unchanged in WBCs and spleen, findings that were confirmed by qRT-PCR (Table 6). Ras-related protein Rab-38 (*RAB38*) was induced by ~83-, 50- and 20-fold in the liver, lungs and inguinal lymph nodes with melanoma, respectively.

Genes encoding proteins associated directly with cellular transcription (Table 5) were among the majority of genes induced in melanoma compared with normal samples, including the proto-oncogene protein c-Fos (*FOS*) and the proto-oncogene c-jun (*JUN*). *FOS*, an immediate early gene expressed transiently in the G_1 phase of the cell cycle after antigen stimulation of T lymphocytes, which plays a role in cell proliferation (Ullman *et al.*, 1990; Angel and Karin, 1991), was significantly induced by ~3-fold in inguinal lymph nodes with melanoma, but remained unchanged in the rest of the melanoma samples. *JUN*, which encodes a protein that interacts directly with specific target DNA sequences to regulate gene expression (Bohmann *et al.*, 1987) and interacts with *FOS* to form a dimer, was upregulated 4-fold and 3-fold in the liver and inguinal lymph nodes with melanoma, respectively.

Several genes encoding signal transduction proteins were also upregulated in the samples with melanoma including tumor-associated calcium signal transducer 1 (*TACSTD1*) and mitogen-activated protein kinase 3 (*MAP3K2*). *TACSTD1,* which was induced 20-fold in the liver and 22-fold in the inguinal lymph nodes with melanoma, encodes a carcinoma-associated antigen that functions as a homotypic calcium-independent cell adhesion molecule (Simon *et al.*, 1990). The induced expression of *TACSTD1* was confirmed by qRT-PCR (Table 6). *MAP3K2*, upregulated >7-fold and >2-fold in the liver and lungs with melanoma, respectively, is a component of a protein kinase signal transduction cascade and regulates the JNK and ERK5 pathways by phosphorylating and activating MAP2K5 and MAP2K7 (Cheng *et al.*, 2000).

Malignant melanomas show impaired ability to undergo programmed cell death (apoptosis) in response to a wide range of external stimuli, conferring them with a selective advantage for progression and metastasis as well as resistance to therapy (Ivanov *et al.*, 2003; Gong *et al.*, 2005; Li *et al.*, 2005). In this study, we observed increased transcription of anti-apoptotic genes in the melanoma compared with normal samples (Table 5), including *BCL-2*, an integral outer mitochondrial membrane protein that blocks cell death by controlling mitochondrial membrane permeability and prevents translocation of pro-apoptotic caspases (Chandra *et al.*, 2002) that was upregulated by 2.5-fold in melanoma compared with normal liver, but was unchanged in the rest of the samples. Apoptosis inhibitor survivin (*BIRC5*) was induced 2.5-fold and 2.7-fold in liver and lungs with melanoma, respectively. Bcl-2 and other inhibitors of apoptosis such as survivin are elevated in a range of human cancers including melanomas, prostate carcinoma and gliomas (LaCasse *et al.*, 1998; Altieri, 2003; Chawla-Sarkar *et al.*, 2004).

In contrast, genes encoding pro-apoptotic proteins were repressed (Table 5), including DAP kinase-related apoptosis-inducing protein kinase 1 (*STK17A*), cell cycle and apoptosis regulatory protein 1 (*CCAR1*), as well as p53-regulated protein PA26 (Sestrin 1; *SESN1*). *STK17A*, which encodes a serine/threonine-protein kinase 17α that acts as a positive regulator of apoptosis (Sanjo *et al.*, 1998), was down-regulated ~6-fold in inguinal lymph nodes with malignant melanoma. *CCAR1*, downregulated 3-fold in inguinal lymph nodes with melanoma, mediates apoptosis induction, which involves sequestration of 14–3–3 protein(s) and altered expression of multiple cell cycle regulatory genes including *MYC*, *CCNB1* and *CDKN1A* (Rishi *et al.*, 2003). *SESN1*, a potential regulator of cellular growth (Velasco-Miguel *et al.*, 1999), was downregulated by ~3-fold and 2-fold in melanoma liver and lungs, respectively.

https://doi.org/10.1017/S1751731111001637 Published online by Cambridge University Press

https://doi.org/10.1017/S1751731111001637 Published online by Cambridge University Press

Table 5 *Genes differentially expressed between melanoma and normal porcine samples*

Accession[b]	Gene name	Gene description	Fold change difference in gene expression[a]				
			WBCs	Liver	Lungs	Inguinal lymph nodes	Spleen
Melanoma-associated proteins							
CO955380	*SILV**	Melanocyte protein Pmel 17 precursor	–	783.83	430.26	163.52	–
CN154304	*TYR**	Tyrosinase precursor	–	137.75	80.85	27.63	–
BF703346	*RAB38*	Ras-related protein Rab-38 (antigen NY-MEL-1)	−3.85	82.50	50.53	20.55	–
CN153410	*GPNMB**	Putative transmembrane protein NMB precursor	−3.57	66.84	15.18	47.49	–
BX917105	*MATP**	Membrane-associated transporter protein (AIM-1 protein)	–	60.09	53.38	21.30	–
AY604429.1	*LGALS1**	Galectin-1	–	58.62	4.27	5.40	–
NM_214023.1	*SPP1**	Osteopontin precursor	–	40.39	20.57	21.82	–
BI183561	*CITED1*	Cbp/p300-interacting transactivator 1 (melanocyte-specific protein 1)	–	73.83	12.55	13.03	–
CN157469	*MRPL28**	Melanoma antigen p15	–	5.45	4.32	3.62	–
BX915975	*MCAM**	Melanoma adhesion molecule	–	4.81	−2.17	–	–
CK451293	*KAI1**	CD82 antigen /suppressor of tumorigenicity-6	–	4.11	–	6.91	–
BI182740	*LTBP4**	Latent transforming growth factor beta binding protein 4	–	4.03	–	2.11	–
CF180835	*CYP4A22**	Cytochrome P450 4B1	–	17.29	–	26.78	–
Trancription							
NM_213880.1	*JUN*	Transcription factor AP-1 (proto-oncogene c-jun, p39)	–	4.30	–	3.09	–
CK459882	*MAFG*	Transcription factor MafG	–	4.16	–	–	–
CF365377	*FOS*	Proto-oncogene protein c-fos	–	–	–	2.65	–
CK457106	*DDX4*	DEAD-box protein 4	–	26.65	–	6.27	–
CO949206	*RUNX1*	Runt-related transcription factor 1 (oncogene AML-1)	–	2.98	–	–	–
BG833819	*TCF7L1*	Transcription factor 7-like 1	–	18.86	2.16	2.44	–
CK466336	*TRIM28*	Transcription intermediary factor 1-beta	–	5.01	2.30	–	–
BF080331	*CREBL1*	cAMP-responsive element binding protein-like 1	–	2.56	–	–	–
Signal transduction							
NM_214419.1	*TACSTD1**	Tumor-associated calcium signal transducer 1 precursor	–	20.32	–	21.73	–
BF193623	*MAP3K2*	Mitogen-activated protein kinase kinase kinase 3	–	7.30	2.45	1.34	–
M73237.1	*SYK*	Tyrosine-protein kinase SYK	–	20.91	2.35	0.35	–
CK453252	*GPR143*	G protein-coupled receptor 143	–	184.20	22.36	14.42	–
BX674263	*RAP2B*	Ras-related protein Rap-2b	–	82.36	4.52	–	–
BQ603688	*AGTPBP1*	ATP/GTP binding protein 1	23.88	–	–	–	–
BI402402	*CAGB*	Calgranulin B	–	60.24	2.50	37.84	–
CB475695	*CAGC*	Calgranulin C	–	11.26	4.47	17.33	–
Apoptosis							
NM_214141.1	*BIRC5*	Apoptosis inhibitor survivin	–	2.53	2.72	–	–
BF702022	*BCL2*	Apoptosis regulator Bcl-2.	–	2.52	–	–	–
CN153547	*CCAR1*	Cell-cycle and apoptosis regulatory protein 1	–	–	−2.41	−2.97	–
BF711044	*SESN1*	Sestrin 1 (p53-regulated protein PA26)	–	−2.66	−2.18	–	–
CN161837	*STK17A*	DAP kinase-related apoptosis-inducing protein kinase 1	–	–	–	−5.81	–
AJ669283	*APAF1*	Apoptotic protease activating factor 1	–	7.89	3.42	−4.00	−2.86
CO953639	*CDK11*	Cyclin-dependent kinase 11 (death-preventing kinase)	–	2.35	–	−2.70	–
Cell cycle							
CB286552	*CDK2*	Cell division protein kinase 2	–	20.15	2.49	–	–
CN166027	*CDC42SE1*	CDC42 small effector 1	–	9.39	–	–	–
CO946563	*Q9ULL6*	Cdc42 GTPase-activating protein	–	5.08	–	−3.85	−2.17

https://doi.org/10.1017/S1751731111001637 Published online by Cambridge University Press

Table 5 *Continued*

Accession[b]	Gene name	Gene description	Fold change difference in gene expression[a]				
			WBCs	Liver	Lungs	Inguinal lymph nodes	Spleen
AW785061	*MCM5*	DNA replication licensing factor MCM5	–	4.55	–	–	–
BI183543	*DOCK9*	Dedicator of cytokinesis protein 9	–	4.30	–	−3.85	–
BX675768	*CDC25B*	M-phase inducer phosphatase 2	–	3.17	–	–	–
BI403744	*ARHGEF2*	Rho guanine nucleotide exchange factor 2	–	4.86	–	–	–
CK458570	*DNAJC14*	Nuclear protein Hcc-1	–	3.58	–	–	–
CO940523	*PA2G4*	Proliferation-associated protein 2G4	–	3.46	–	–	–
AW359493	*BOP1*	Ribosome biogenesis protein BOP1	–	4.34	–	–	–
Metabolism							
CN166623	*PKM2*	Pyruvate kinase, isozymes M1/M2	–	25.17	2.71	–	−2.50
CF180857	*ALDOA*	Fructose-bisphosphate aldolase A	–	6.14	3.46	–	–
CO946907	*DPYD*	Dihydropyrimidine dehydrogenase [NADP+] precursor	–	19.41	3.60	−2.63	–
CN154388	*GSTP1*	Glutathione S-transferase P	–	19.34	2.77	2.77	–
BF712908	*LPL*	Lipoprotein lipase precursor	–	18.15	3.52	−6.25	−8.33
BF703588	*TDO2*	Tryptophan 2,3-dioxygenase	–	16.56	8.82	2.40	–
NM_213938.1	*OXCT1*	Succinyl-CoA:3-ketoacid-coenzyme A transferase 1, mitochondrial precursor	–	8.48	–	−2.44	−2.56
BX915001	*HK1*	Hexokinase, type I	–	8.37	–	–	–
CF180857	*ALDOA*	Fructose-bisphosphate aldolase A	–	6.14	3.46	–	–
NM_213827.1	*DCXR*	Dicarbonyl/L-xylulose reductase	–	60.21	–	51.38	–
D83766.1	*RENBP*	N-acylglucosamine 2-epimerase	–	14.22	9.52	8.18	−2.33
Immune response							
AY463542.1	*HLA-A*	MHC class I antigen A*3	–	20.70	2.20	–	–
AY102480.1	*HLA-DRB4*	MHC class I antigen DRB1*4	–	14.25	2.14	–	−2.86
AB032169.1	*HLA-DMA*	MHC class II antigen DMA	–	13.47	3.16	–	−2.94
NM_213774.1	*CD74*	CD74 antigen (MHC class II, gamma chain)	–	27.52	2.62	–	−2.08
NM_213773.1	*CXCR4*	C-X-C chemokine receptor type 4	2.51	17.76	2.50	–	–
BX918583	*LY96*	Lymphocyte antigen 96 precursor (MD-2 protein)	–	13.32	8.55	–	−2.94
BI337485	*TAP1*	Antigen peptide transporter 1 (APT1)	–	9.74	2.28	−2.08	–
AF156712.1	*ICAM1*	Intercellular adhesion molecule-1 precursor (ICAM-1)	–	9.55	–	–	–
AY312067.1	*CCL21*	Small inducible cytokine A21 precursor (CCL21)	–	8.78	2.56	–	–
NM_214083.1	*IL2RG*	Interleukin- 2 receptor gamma chain	–	7.94	2.45	−3.03	−2.70
CN163354	*TNFRSF5*	Tumor necrosis factor receptor superfamily member 5	–	7.30	2.33	−3.70	−2.50
BX671266	*IL2RB*	Interleukin-2 receptor beta chain precursor	–	5.49	3.08	–	−2.44
BI184849	*ANXA11*	Annexin A11	–	3.89	–	–	–
NM_213771.1	*IL10RB*	Interleukin-10 receptor beta chain precursor	–	3.12	–	–	–
CB287867	*IGJ*	Immunoglobulin J chain	–	26.46	4.26	–	–
U29948.1	*DF*	Complement factor D precursor	–	26.46	–	–	–

[a]Gene expression values of samples obtained from the melanoma-afflicted pig were divided by those of the corresponding samples from the non-melanoma pig, to provide differential gene expression ratios. Expression ratios ⩾2.0 (⩾2.0-fold upregulation) and ⩽0.05 (⩾2.0-fold downregulation) are considered significant.
[b]GenBank accession of cDNA elements spotted on Affymetrix oligonucleotide microarrays.
*Microarray data validated by quantitative real-time RT-PCR.
(–), no significant change in gene expression.

https://doi.org/10.1017/S1751731111001637 Published online by Cambridge University Press

Table 6 *Validation of microarray data by qRT-PCR*

			Fold changes in gene expression by:									
			Microarray analysis					qRT-PCR analysis				
Accession[a]	Gene	Gene description	WBCs	Liver	Lungs	Inguinal lymph nodes	Spleen	WBCs	Liver	Lungs	Inguinal lymph nodes	Spleen
CN154304	*TYR*	Tyrosinase	–	137.75	80.85	27.63	–	–	689.80	15.18	85.077	−52.05
NM_214419.1	*TACSTD1*	Tumor-associated calcium signal transducer 1	–	20.32	–	21.73	–	–	3.79	−7.39	45.49	−13.01
CO955380	*SILV*	Melanocyte protein MEL17	–	783.83	430.26	163.52	–	–	2654.81	170.88	17.03	−3.17
NM_214023.1	*SPP1*	Secreted phosphoprotein-I	–	40.39	20.57	21.82	–	–	169.93	5.23	14.12	−5.68
CN157469	*MRPL28*	Melanoma antigen p15	–	5.45	4.32	3.62	–	2.33	2.21	–	1.95	–
BX915975	*MCAM*	Melanoma cell adhesion molecule MCAM	–	4.81	−2.17	–	–	4.19	4.52	−5.15	–	−22.53
BX917105	*MATP*	Membrane-associated transporter protein (melanoma antigen AIM-1)	–	60.09	53.38	21.30	–	–	35.73	65.34	39.37	−5.87
BI182740	*LTBP4*	Latent transforming growth factor beta binding protein 4	–	4.03	–	2.11	–	3.38	8.23	–	3.02	−19.10
AY604429.1	*LGALS1*	Galectin-1	–	58.62	4.27	5.40	–	–	184.12	3.04	3.23	−5.16
CF359325	*KAI1*	CD82 antigen /suppressor of tumorigenicity-6	–	4.97	2.41	5.02	–	–	–	−2.01	–	−3.09
CO949684	*GPNMB*	Transmembrane glycoprotein NMB	–	44.71	29.20	47.78	–	–	97.48	5.04	29.67	−5.86
CF180835	*CYP4A22*	Cytochrome P450 4B1	–	17.29	–	26.78	–	9.13	111.62	−2.51	65.45	−8.13

qRT-PCR = quantitative real-time RT-PCR; WBCs = white blood cells.
[a]From GenBank database.
(–), no significant changes in gene expression.

https://doi.org/10.1017/S1751731111001637 Published online by Cambridge University Press

Table 7 *Sequence similarities between pig ESTs spotted on microarrays and human protein homologs of selected genes analyzed by microarrays and quantitative real-time RT-PCR*

Gene name	Gene description	Accession[a] of porcine EST	Accession[b] of human protein homolog	BLAST × sequence similarity (%)[c]	BLAST × bit score	BLAST × *E*-value	Human gene chromosomal location	Ref
TYR	Tyrosinase	CN154304	AAB60319	91	237	7.00E−61	11q14-q21	[d]
TACSTD1	Tumor-associated calcium signal transducer 1	NM_214419.1	CAG47078	90	461	5.00E−128	2p21	[e]
SPP1	Secreted phosphoprotein-I	NM_214023.1	NM_001040060	76	194	2.00E−47	4q21-q25	[f]
SILV	Melanocyte protein MEL17	CO955380	AAB31176	94	37.7	2.20E−01	12q12-q13	[g]
MRPL28	Melanoma antigen p15	CN157469	EAW85830	70	120	2.00E−45	16p13.3	[h]
MCAM	Melanoma cell adhesion molecule MCAM	BX915975	BAD93162	88	78.6	4.00E−14	11q23.3	[i]
MATP	Membrane-associated transporter protein/melanoma antigen AIM1	BX917105	NM_016180	91	210	3.00E−53	5p13.3	[j]
LTBP4	Latent transforming growth factor beta binding protein 4	BI182740	EAW56985	77	357	6.00E−97	19q13.1-q13.2	[k]
LGALS1	Galectin-1	AY604429.1	1713410A	92	253	2.00E−66	22q12-13.1	[l]
KAI1	CD82 antigen/suppressor of tumorigenicity-6	AK236958	CAG28578	94	230	4.00E−58	11p11.2	[m]
GPNMB	Transmembrane glycoprotein NMB	NM_001098584	AAH32783	84	817	0.00E+00	7p15	[n]
CYP4A22	Cytochrome P450 4B1	CF180835	AAQ21368	84	280	8.00E−74	1p33	[o]

EST = expressed sequence tags; BLAST = Basic Local Alignment Search Tool; NCBI = National Center for Biotechnology Information.
[a]From GenBank database.
[b]From NCBI protein database.
[c]Sequence similarity between pig ESTs spotted on microarrays and human proteins; performed by BLAST × analysis (Altschul *et al.*, 1997) against a non-redundant peptide sequence database collected from the NCBI.
[d]Perez *et al.* (2000), Kwon (1993), Battyani *et al.* (1993), Tripathi *et al.* (1992), Overwijk *et al.* (1999), Parkhurst *et al.* (1998) and Kawakami *et al.* (1998).
[e]Wallace *et al.* (2005) and Mitas *et al.* (2003).
[f]Talantov *et al.* (2005), Haqq *et al.* (2005) and Zhou *et al.* (2005).
[g]Nordlund *et al.* (1998) and Solano *et al.* (2000).
[h]Robbins *et al.* (1995).
[i]Leslie *et al.* (2007) and Watson-Hurst *et al.* (2006).
[j]Lewis *et al.* (2005).
[k]Giltay *et al.* (1997).
[l]van den Brûle (1995).
[m]Takaoka *et al.* (1998).
[n]Weterman *et al.* (1995).
[o]Savas *et al.* (2003).

Table 8 *Least square values of average absolute expression (ng cDNA) with standard error (in brackets), and corresponding* P*-value for contrasting expression of 12 genes for piglets with regressing, progressing and without tumors*

Gene name	Average absolute expression (ng cDNA)[a]			
	Piglets with regressing tumors ($n = 7$)	Piglets with progressing tumors ($n = 5$)	Piglets without tumors ($n = 10$)	*P*-value
TYR	818 (587)	948 (743)	1725 (752)	0.034
TACSTD1	1194 (1193)	1229 (956)	2580 (823)	0.015
SILV	293 (289)	972 (987)	2089 (2246)	0.097
SPP1	1641 (815)	2080 (797)	1866 (780)	0.642
MRPL28	458 (584)	911 (458)	734 (699)	0.450
MCAM	235 (226)	663 (512)	1127 (1157)	0.123
MATP	409 (311)	716 (379)	148 (188)	0.007
LTBP4	396 (269)	816 (568)	1568 (1719)	0.245
LGALS1	311 (341)	1016 (555)	1762 (2018)	0.143
KAI1	814 (929)	1482 (1297)	1214 (1620)	0.724
GPNMB	272 (266)	1090 (974)	1816 (1572)	0.050
CYP4A22	3122 (2232)	1657 (1753)	5616 (2925)	0.022

[a]To analyze gene expression during tumor regression, frozen liver samples from 22 pigs obtained from the historical Sinclair herd (Texas A & M University) were used in quantitative real-time RT-PCR analysis of 12 genes that were upregulated during tumor metastasis.

qRT-PCR analysis validated the microarray data of 12 genes that were differentially expressed (induced/repressed) or unchanged in melanoma compared with normal samples, and there was a qualitative agreement in the results of the microarray and qRT-PCR analyses (Table 6). The genes *TYR*, *SILV*, *SPP1*, *MRPL28*, *MATP*, *LGALS1* and *GPNMB* were over-expressed (⩾2-fold), based on microarray and qRT-PCR analyses, in the liver, lung and inguinal lymph node samples. *TACSTD1* was over-expressed based on both methods in the liver and inguinal lymph nodes, but in the lungs, it was unchanged by microarray and downregulated by qRT-PCR analysis. *MCAM* was upregulated in the liver, downregulated in the lungs and unchanged in the inguinal lymph nodes, while *LTBP4* was upregulated in the liver and inguinal lymph nodes, but remained unchanged in the lungs, based on both methods. However, in the spleen, all 12 genes were unchanged after microarray analysis but were downregulated based on qRT-PCR, with the exception of *MRPL28*, while in the WBCs, all 12 transcripts showed no significant changes in expression after microarray analysis, but after qRT-PCR, at least four transcripts were upregulated by ⩾2-fold (*MRPL28*, *MCAM*, *LTBP4* and *CYP4A22*). Similar to the microarray findings, after qRT-PCR analysis, the liver exhibited the highest fold change differences in gene expression in melanoma compared with normal samples, followed by the inguinal lymph nodes, lungs, spleen and then WBCs. Higher differential gene expression ratios were observed between melanoma and normal samples after qRT-PCR, compared with microarray analysis, a finding consistent with earlier observations that qRT-PCR has greater detection sensitivity than microarray technology (Yuen *et al.*, 2002).

Gene expression during tumor progression and regression

The results of the ANOVA test contrasting absolute RNA expression for 12 genes for piglets with regressing, progressing or without tumors are shown in Table 8. The results of the ANOVA test were significant for TYR, TACSTD1, MATP, GPNMB and CYP4A22, with *P*-values of 0.034, 0.015, 0.007, 0.050 and 0.022, respectively. With the exception of MATP, the other genes (TYR, TACSTD1, GPNMB and CYP4A22) were downregulated, suggesting that gene expression is under-regulated when comparing piglets with regressing *v.* progressing tumors. This finding is contrary to our earlier observation using animals with metastatic melanoma *v.* those that were disease free during the first 6 weeks of life, where TYR, TACSTD1, MATP, GPNMB and CYP4A22 were all upregulated in liver samples of animals with melanoma compared with those without the disease (Table 6). These results illustrate that changes in gene expression may occur in varying patterns during progression and regression of malignant melanoma.

Discussion

Our study identified genes from diverse functional classes that were differentially expressed in melanoma compared with normal tissue samples, suggesting that numerous cellular processes were transcriptionally modulated during the progression of melanoma in Sinclair swine (Table 5). Among tissues that were tested, the inguinal lymph node and liver had the highest number of genes (6266 and 5917, respectively) that were differentially expressed between melanoma and normal cells (Table 4), suggesting that these tissues are favored as predilection sites by metastatic melanocytes. The genes that were differentially expressed in this study showed ⩾80% sequence similar to genes modulated in human melanomas (Table 7). These molecular similarities are significant as some of the genes that were either induced or repressed in this study are used as biomarkers in human malignant melanoma. The *TYR* gene associated with

https://doi.org/10.1017/S1751731111001637 Published online by Cambridge University Press

melanoma pathogenesis and significantly upregulated in this study is expressed at various levels in human melanoma cells and is considered an important biochemical marker of melanocytes (Kawakami *et al.*, 1998; Parkhurst *et al.*, 1998; Overwijk *et al.*, 1999), while the *RAB28* gene, also over-expressed in this study, is expressed in melanocytes and is involved in melanosomal transport and docking (Jager *et al.*, 2000) as well as in the proper sorting of tyrosinase-related protein 1 (*TYRP1*). The increased transcription of anti-apoptotic genes in this study, including *BCL-2* that is elevated in a range of human cancers including melanomas, prostate carcinoma and gliomas (LaCasse *et al.*, 1998; Altieri, 2003; Chawla-Sarkar *et al.*, 2004), demonstrates further the similarity in the modulation of gene expression in Sinclair swine and human cancers.

The contrasting results observed in the expression of genes tested by microarrays and qRT-PCR in liver samples obtained from 6-week-old piglets, compared with gene expression tested by qRT-PCR in the frozen liver samples collected at an older age, suggests a better developed and more mature immune system in the older piglets. The downregulation of TYR, TACSTD1, GPNMB and CYP4A22 during regression and progression of the disease at a later age may be the result of a cell-mediated response that attacks metastatic melanocytes as the animals become older (Greene *et al.*, 1997), and is consistent with the observation that animals with melanoma develop vitiligo with age (Misfeldt and Grimm, 1994).

The findings in this study may facilitate further discovery of candidate genes and molecular markers that can provide a better understanding of the pathogenesis of metastatic melanoma in Sinclair swine.

Acknowledgments

The authors thank the Nevada Genomics Center for microarray and qRT-PCR assistance. The authors also acknowledge the Nevada Animal Disease and Food Safety Laboratory for their assistance with tissue sample collection and processing. The authors are very grateful to Prof. Craig W. Beattie for providing samples of Sinclair swine.

References

Alaoui-Jamali MA, Dupre I and Qiang H 2004. Prediction of drug sensitivity and drug resistance in cancer by transcriptional and proteomic profiling. Drug Resistance Updates 7, 245–255.

Altieri DC 2003. Validating survivin as a cancer therapeutic target. Nature Reviews Cancer 3, 46–54.

Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W and Lipman DJ 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. Nucleic Acids Research 25, 3389–3402.

Angel P and Karin M 1991. The role of Jun, Fos and the AP-1 complex in cell-proliferation and transformation. Biochimica et Biophysica Acta 1072, 129–157.

Barrier A, Lemoine A, Boelle PY, Tse C, Brault D, Chiappini F, Breittschneider J, Lacaine F, Houry S, Huguier M, Van der Laan MJ, Speed T, Debuire B, Flahault A and Dudoit S 2005. Colon cancer prognosis prediction by gene expression profiling. Oncogene 24, 6155–6164.

Battyani Z, Xerri L, Hassoun J, Bonerandi JJ and Grob JJ 1993. Tyrosinase gene expression in human tissues. Pigment Cell Research 6, 400–405.

Bittner M, Meltzer P, Chen Y, Jiang Y, Seftor E, Hendrix M, Radmacher M, Simon R, Yakhini Z, Ben-Dor A, Sampas N, Dougherty E, Wang E, Marincola F, Gooden C, Lueders J, Glatfelter A, Pollock P, Carpten J, Gillanders E, Leja D, Dietrich K, Beaudry C, Berens M, Alberts D and Sondak V 2000. Molecular classification of cutaneous malignant melanoma by gene expression profiling. Nature 406, 536–540.

Bohmann D, Bos TJ, Admon A, Nishimura T, Vogt PK and Tjian R 1987. Human proto-oncogene c-jun encodes a DNA binding protein with structural and functional properties of transcription factor AP-1. Science 238, 1386–1392.

Chandra D, Liu JW and Tang DG 2002. Early mitochondrial activation and cytochrome c upregulation during apoptosis. Journal of Biological Chemistry 277, 50842–50854.

Chawla-Sarkar M, Bae SI, Reu FJ, Jacobs BS, Lindner DJ and Borden EC 2004. Downregulation of Bcl-2, FLIP or IAPs (XIAP and survivin) by siRNAs sensitizes resistant melanoma cells to Apo2L/TRAIL-induced apoptosis. Cell Death and Differentiation 11, 915–923.

Cheng J, Yang J, Xia Y, Karin M and Su B 2000. Synergistic interaction of MEK kinase 2, c-Jun N-terminal kinase (JNK) kinase 2, and JNK1 results in efficient and specific JNK1 activation. Molecular and Cellular Biology 20, 2334–2342.

Clarke PA, te Poele R and Workman P 2004. Gene expression microarray technologies in the development of new therapeutic agents. European Journal of Cancer 40, 2560–2591.

Das Gupta TK, Ronan SG, Beattie CW, Shilkaitis A and Amoss MS Jr 1989. Comparative histopathology of porcine and human cutaneous melanoma. Pediatric Dermatology 6, 289–299.

Eisen MB and Brown PO 1999. DNA arrays for analysis of gene expression. Methods in Enzymology 303, 179–205.

Giltay R, Kostka G and Timpl R 1997. Sequence and expression of a novel member (LTBP-4) of the family of latent transforming growth factor-beta binding proteins. FEBS Letters 411, 164–168.

Gomez-Raya L, Okomo-Adhiambo M, Beattie C, Osborne K, Rink A and Rauw WM 2007. Modeling inheritance of malignant melanoma with DNA markers in Sinclair swine. Genetics 176, 585–597.

Gomez-Raya L, Amoss MS, Da Y, Beattie CW, Smith D, Ash O and Rauw WM 2009. Role of inbreeding, selection on the incidence of cutaneous malignant melanoma in Sinclair swine. Journal of Animal Breeding and Genetics 126, 242–249.

Gong J, Chen N, Zhou Q, Yang B, Wang Y and Wang X 2005. Melanoma inhibitor of apoptosis protein is expressed differentially in melanoma and melanocytic naevus, but similarly in primary and metastatic melanomas. Journal of Clinical Pathology 58, 1081–1085.

Greene JF, Townsend JS and Amoss MS 1994. Histopathology of regression in Sinclair swine model of melanoma. Laboratory Investigation 71, 17–24.

Greene JF Jr, Morgan CD, Rao A, Amoss MS Jr and Arguello F 1997. Regression by differentiation in the Sinclair swine model of cutaneous melanoma. Melanoma Research 7, 471–477.

Haqq C, Nosrati M, Sudilovsky D, Crothers J, Khodabakhsh D, Pulliam BL, Federman S, Miller JR III, Allen RE Singer MI *et al.* 2005. The gene expression signatures of melanoma progression. Proceedings of the National Academy of Sciences of the United States of America 102, 6092–6097.

Hoek K, Rimm DL, Williams KR, Zhao H, Ariyan S, Lin A, Kluger HM, Berger AJ, Cheng E Trombetta ES *et al.* 2004. Expression profiling reveals novel pathways in the transformation of melanocytes to melanomas. Cancer Research 64, 5270–5282.

Hook RR Jr, Aultman MD, Adelstein EH, Oxenhandler RW, Millikan LE and Middleton C 1979. Influence of selective breeding on the incidence of melanomas in Sinclair miniature swine. International Journal of Cancer 24, 668–672.

Huang Y and Sadee W 2003. Drug sensitivity and resistance genes in cancer chemotherapy: a chemogenomics approach. Drug Discovery Today 8, 356–363.

Ivanov VN, Bhoumik A and Ronai Z 2003. Death receptors and melanoma resistance to apoptosis. Oncogene 22, 3152–3161.

Jager D, Stockert E, Jager E, Gure AO, Scanlan MJ, Knuth A, Old LJ and Chen YT 2000. Serological cloning of a melanocyte rab guanosine 5′-triphosphate-binding protein and a chromosome condensation protein from a melanoma complementary DNA library. Cancer Research 60, 3584–3591.

James K, Eisenhauer E, Christian M, Terenziani M, Vena D, Muldal A and Therasse P 1999. Measuring response in solid tumors: unidimensional versus bidimensional measurement. Journal of the National Cancer Institute 91, 523–528.

https://doi.org/10.1017/S1751731111001637 Published online by Cambridge University Press

Kawakami Y, Robbins PF, Wang RF, Parkhurst M, Kang X and Rosenberg SA 1998. The use of melanosomal proteins in the immunotherapy of melanoma. Journal of Immunotherapy 21, 237–246.

Kwon BS 1993. Pigmentation genes: the tyrosinase gene family and the pmel 17 gene family. Journal of Investigative Dermatology 100, 134S–140S.

LaCasse EC, Baird S, Korneluk RG and MacKenzie AE 1998. The inhibitors of apoptosis IAPs and their emerging role in cancer. Oncogene 17, 3247–3259.

Leslie MC, Zhao YJ, Lachman LB, Hwu P and Bar-Eli M 2007. Immunization against MUC18/MCAM, a novel antigen that drives melanoma invasion and metastasis. Gene Therapy 14, 316–323.

Lewis TB, Robison JE, Bastien R, Milash B, Boucher K, Samlowski WE, Leachman SA, Dirk Noyes R, Wittwer CT, Perreard L and Bernard PS 2005. Molecular classification of melanoma using real-time quantitative reverse transcriptase-polymerase chain reaction. Cancer 104, 1678–1686.

Li H, Niederkorn JY, Neelam S and Alizadeh H 2005. Resistance and susceptibility of human uveal melanoma cells to TRAIL-induced apoptosis. Archives of Ophthalmology 123, 654–661.

Luo J, Duggan DJ, Chen Y, Sauvageot J, Ewing CM, Bittner ML, Trent JM and Isaacs WB 2001. Human prostate cancer and benign prostatic hyperplasia: molecular dissection by gene expression profiling. Cancer Research 61, 4683–4688.

Mariadason JM, Arango D, Shi Q, Wilson AJ, Corner GA, Nicholas C, Aranes MJ, Lesser M, Schwartz EL and Augenlicht LH 2003. Gene expression profiling-based prediction of response of colon carcinoma cells to 5-fluorouracil and camptothecin. Cancer Research 63, 8791–8812.

Minitab 15. Minitab. Inc., State College, PA.

Misfeldt ML and Grimm DR 1994. Sinclair miniature swine: an animal model of human melanoma. Veterinary Immunology and Immunopathology 43, 167–175.

Mitas M, Cole DJ, Hoover L, Fraig MM, Mikhitarian K, Block MI, Hoffman BJ, Hawes RH, Gillanders WE and Wallace MB 2003. Real-time reverse transcription-PCR detects KS1/4 mRNA in mediastinal lymph nodes from patients with non-small cell lung cancer. Clinical Chemistry 49, 312–325.

Nordlund JJ, Boissy RE, Hearing VJ, King L and Ortonne JP ed. 1998. The pigmentary system: physiology and pathophysiology. Oxford University Press, New York, NY.

Overwijk WW, Lee DS, Surman DR, Irvine KR, Touloukian CE, Chan CC, Carroll MW, Moss B, Rosenberg SA and Restifo NP 1999. Vaccination with a recombinant vaccinia virus encoding a 'self' antigen induces autoimmune vitiligo and tumor cell destruction in mice: requirement for CD4+ T lymphocytes. Proceedings of the National Academy of Sciences of the United States of America 96, 2982–2987.

Parkhurst MR, Fitzgerald EB, Southwood S, Sette A, Rosenberg SA and Kawakami Y 1998. Identification of a shared HLA-A*0201-restricted T-cell epitope from the melanoma antigen tyrosinase-related protein 2 TRP2. Cancer Research 58, 4895–4901.

Perez RP, Zhang P, Bosserhoff AK, Buettner R and Abu-Hadid M 2000. Expression of melanoma inhibitory activity in melanoma and nonmelanoma tissue specimens. Human Pathology 31, 1381–1388.

Pfaffl MW, Horgan G and Dempfle L 2002. Relative Expression Software Tool REST© for group wise comparison and statistical analysis of relative expression results in real-time PCR. Nucleic Acids Research 30, E36.

Rishi AK, Zhang L, Boyanapalli M, Wali A, Mohammad RM, Yu Y, Fontana JA, Hatfield JS, Dawson MI, Majumdar APN and Reichert U 2003. Identification and characterization of a cell cycle and apoptosis regulatory protein-1 as a novel mediator of apoptosis signaling by retinoid CD437. The Journal of Biological Chemistry 278, 33422–33435.

Robbins PF, El-Gamil M, Li YF, Topalian SL, Rivoltini L, Sakaguchi K, Appella E, Kawakami Y and Rosenberg SA 1995. Cloning of a new gene encoding an antigen recognized by melanoma-specific HLA-A24-restricted tumor-infiltrating lymphocytes. The Journal of Immunology 154, 5944–5950.

Sanjo H, Kawai T and Akira S 1998. DRAKs, novel serine/threonine kinases related to death-associated protein kinase that trigger apoptosis. Journal of Biological Chemistry 273, 29066–29071.

Savas U, Hsu MH and Johnson EF 2003. Differential regulation of human CYP4A genes by peroxisome proliferators and dexamethasone. Archives of Biochemistry and Biophysics 409, 212–220.

Simon B, Podolsky DK, Moldenhauer G, Isselbacher KJ, Gattoni-Celli S and Brand SJ 1990. Epithelial glycoprotein is a member of a family of epithelial cell surface antigens homologous to nidogen, a matrix adhesion protein. Proceedings of the National Academy of Sciences of the United States of America 87, 2755–2759.

Solano F, Martínez-Esparza M, Jiménez-Cervantes C, Hill SP, Lozano JA and García-Borrón JC 2000. New insights on the structure of the mouse silver locus and on the function of silver protein. Pigment Cell Research 13, 118–124.

Takaoka A, Hinoda Y, Sato S, Itoh F, Adachi M, Hareyama M and Imai K 1998. Reduced invasive and metastatic potentials of KAI1-transfected melanoma cells. Cancer Science 89, 397–404.

Talantov D, Mazumder A, Yu JX, Briggs T, Jiang Y, Backus J, Atkins D and Wang Y 2005. Novel genes associated with malignant melanoma but not benign melanocytic lesions. Clinical Cancer Research 11, 7234–7242.

Tissot RG, Beattie CW and Amoss MS Jr 1987. Inheritance of Sinclair swine cutaneous malignant melanoma. Cancer Research 47, 5542–5545.

Tripathi RK, Hearing VJ, Urabe K, Aroca P and Spritz RA 1992. Mutational mapping of the catalytic activities of human tyrosinase. Journal of Biological Chemistry 267, 23707–23712.

Tsai S, Cassady JP, Freking BA, Nonneman DJ, Rohrer GA and Piedrahita JA 2006. Annotation of the Affymetrix porcine genome microarray. Animal Genetics 37, 423–424.

Ullman KS, Northrop JP, Verweij CL and Crabtree GR 1990. Transmission of signals from the T lymphocyte antigen receptor to the genes responsible for cell proliferation and immune function: the missing link. Annual Review of Immunology 8, 421–452.

van den Brûle FA, Buicu C, Baldet M, Sobel ME, Cooper DN, Marschal P and Castronovo V 1995. Galectin-1 modulates human melanoma cell adhesion to laminin. Biochemical and Biophysical Research Communications 209, 760–767.

Velasco-Miguel S, Buckbinder L, Jean P, Gelbert L, Talbott R, Laidlaw J, Seizinger B and Kley N 1999. PA26, a novel target of the p53 tumor suppressor and member of the GADD family of DNA damage and growth arrest inducible genes. Oncogene 18, 127–137.

Wallace MB Block MI, Gillanders W, Ravenel J, Hoffman BJ, Reed CE, Fraig M, Cole D and Mitas M 2005. Accurate molecular detection of non-small cell lung cancer metastases in mediastinal lymph nodes sampled by endoscopic ultrasound-guided needle aspiration. Chest 127, 430–437.

Wang Y, Jatkoe T, Zhang Y, Mutch MG, Talantov D, Jiang J, McLeod HL and Atkins D 2004. Gene expression profiles and molecular markers to predict recurrence of Dukes' B colon cancer. Journal of Clinical Oncology 22, 1564–1571.

Watson-Hurst K and Becker D 2006. The role of N-cadherin, MCAM and beta3 integrin in melanoma progression, proliferation, migration and invasion. Cancer Biology and Therapy 5, 1375–1382.

Weterman MAJ, Ajubi N, van Dinter IMR, Degen WGJ, van Muijen GNP, Ruiter DJ and Bloemers HPJ 1995. Nmb, a novel gene, is expressed in low-metastatic human melanoma cell lines and xenografts. International Journal of Cancer 60, 73–81.

Yuen T, Wurmbach E, Pfeffer RL, Ebersole BJ and Sealfon SC 2002. Accuracy and calibration of commercial oligonucleotide and custom cDNA microarrays. Nucleic Acids Research 30, e48.

Zhou Y, Dai DL, Martinka M, Su M, Zhang Y, Campos EI, Dorocicz I, Tang L, Huntsman D, Nelson C, Ho V and Li G 2005. Osteopontin expression correlates with melanoma invasion. Journal of Investigative Dermatology 124, 1044–1052.

https://doi.org/10.1017/S1751731111001637 Published online by Cambridge University Press