Econometric Theory, **24**, 2008, 1443–1455. Printed in the United States of America.
doi:10.1017/S0266466608080675

NOTES AND PROBLEMS

A GENERAL BOUND FOR THE LIMITING DISTRIBUTION OF BREITUNG'S STATISTIC

JAMES DAVIDSON
University of Exeter

JAN R. MAGNUS
Tilburg University

JAN WIEGERINCK
University of Amsterdam

We consider the Breitung (2002, *Journal of Econometrics* 108, 343–363) statistic ξ_n, which provides a nonparametric test of the I(1) hypothesis. If ξ denotes the limit in distribution of ξ_n as $n \to \infty$, we prove (Theorem 1) that $0 \leq \xi \leq 1/\pi^2$, a result that holds under any assumption on the underlying random variables. The result is a special case of a more general result (Theorem 3), which we prove using the so-called cotangent method associated with Cauchy's residue theorem.

1. INTRODUCTION

Let $x_1, x_2, \ldots, x_n$ represent an arbitrary sequence of random variables and let $\bar{x}_n$ denote the sample mean. We consider

$$\xi_n := \frac{\sum_{t=1}^{n}\left(\sum_{j=1}^{t}(x_j - \bar{x}_n)\right)^2}{n^2\sum_{t=1}^{n}(x_t - \bar{x}_n)^2},$$

which is the statistic for the Breitung (2002) nonparametric test of the I(1) hypothesis. Let ξ denote the limit in distribution of ξ_n as $n \to \infty$. We wish to prove the following theorem.

We are grateful to Dieter Balkenborg, Nigar Hashimzade, Michael Jansson, Chris Müris, Morten Nielsen, Paolo Paruolo, and Katsuto Tanaka for helpful advice and discussions and to two referees for constructive comments. Figure 1 is taken from Davidson (2008). Address correspondence to Jan R. Magnus, Department of Econometrics and Operations Research, Tilburg University, P.O. Box 90153, 5000 LE Tilburg, The Netherlands; e-mail: magnus@uvt.nl.

© 2008 Cambridge University Press 0266-4666/08 $15.00

https://doi.org/10.1017/S0266466608080675 Published online by Cambridge University Press

THEOREM 1. *ξ is supported on the interval* $[0, 1/\pi^2]$.

This result holds under any assumption on $x_1, \ldots, x_n$ whatsoever. It is trivially true when the process is I(0) because then, as Breitung shows, the distribution is degenerate at 0. In fact, it can be shown that the same holds for a covariance stationary I(d) process when $|d| < \frac{1}{2}$. Under Breitung's null hypothesis, where x_t is an I(1) process subject to the usual regularity conditions, ξ would correspond to the functional

$$\frac{\int_0^1 \left(\int_0^t W\,dr - t \int_0^1 W\,dr \right)^2 dt}{\int_0^1 W^2\,dt - \left(\int_0^1 W\,dt \right)^2},$$

where W denotes standard Brownian motion. However, the theorem also holds when x_t is I(d), for *any* finite d. Some simulations are shown in Figure 1 for cases with $1 \leq d \leq 2$.

It will be convenient to write ξ_n in a different form. Let $x := (x_1, x_2, \ldots, x_n)'$ and define the $n \times n$ lower triangular "cumulation" matrix




FIGURE 1. Kernel density plots of Breitung's statistic computed from one million replications of I(d) series ($n = 1{,}000$).

https://doi.org/10.1017/S0266466608080675 Published online by Cambridge University Press

$$C := \begin{pmatrix} 1 & 0 & \dots & 0 \\ 1 & 1 & \dots & 0 \\ \vdots & \vdots & & \vdots \\ 1 & 1 & \dots & 1 \end{pmatrix},$$

as in Tanaka (1996, eqn. (1.3)). Then,

$$\xi_n = \frac{x'MC'CMx}{n^2 x'Mx},$$

where $M := I_n - (1/n)\iota\iota'$ and $\iota := (1,1,\dots,1)'$. Let $A := (1/n)C$ and let Q be an $n \times (n-1)$ matrix containing the eigenvectors associated with the $n-1$ unit eigenvalues of M, so that $M = QQ'$. Letting $y := Q'x$ we have

$$\xi_n = \frac{x'MA'AMx}{x'Mx} = \frac{y'Q'A'AQy}{y'y},$$

so that

$$\xi_n \leq \lambda_{\max}(Q'A'AQ) = \lambda_{\max}(AQQ'A') = \lambda_{\max}(AMA').$$

Hence, the theorem is true if and only if

$$\lambda_{\max}(AMA') \to \frac{1}{\pi^2} \quad \text{as } n \to \infty.$$

The plan of this note is as follows. In Section 2 we study the eigenvalues of the matrix AMA' and show that these can be found as the solutions of a particular equation (Theorem 2). In Section 3 we prove a generalization of Theorem 1, which states that, for any fixed j, the jth largest eigenvalue μ_j of AMA' converges to $1/(j^2\pi^2)$ (Theorem 3). The proof uses the so-called cotangent method associated with Cauchy's residue theorem. In Section 4 we discuss the speed of convergence and the behavior of the whole set of eigenvalues when n is large. Two Appendixes accompany this note. In Appendix A we discuss the determinant of the matrix $V - \omega\omega'$ where V is positive semidefinite and ω is a vector. In Appendix B we explain the cotangent method.

2. THE EIGENVALUES OF *AMA'*

Before proving the theorem we investigate what can be said about the eigenvalues of AMA'. We know that the matrix $A'A$ is positive definite and that its eigenvalues are given by $\lambda_1 > \lambda_2 > \cdots > \lambda_n > 0$, where

https://doi.org/10.1017/S0266466608080675 Published online by Cambridge University Press

$$\lambda_j = \frac{1}{4n^2 \sin^2(\alpha_j)}, \qquad \alpha_j := \frac{2j-1}{2n+1} \cdot \frac{\pi}{2}, \qquad j = 1, \ldots, n. \tag{1}$$

These eigenvalues were first obtained by Rutherford (1946); see also Tanaka (1996, eqn. (1.4)). The eigenvalues used in Dickey and Fuller (1979) are the same but presented in a different form. Because the sine function is monotonic on the interval $(0, \pi/2)$, the matrix $A'A$ has no multiple eigenvalues.

The matrix AMA' is positive semidefinite and has rank $n - 1$. Its ijth element is given by

$$(AMA')_{ij} = \frac{1}{n^2}\left(\min(i,j) - \frac{ij}{n}\right).$$

By Theorem 11.11 of Magnus and Neudecker (1988, p. 210) we know that its eigenvalues $\mu_1 \geq \mu_2 \geq \cdots \geq \mu_{n-1} > \mu_n = 0$ satisfy

$$\lambda_1 \geq \mu_1 \geq \lambda_2 \geq \mu_2 \geq \cdots \geq \lambda_{n-1} \geq \mu_{n-1} \geq \lambda_n.$$

For fixed j and large n we may use the approximation $\sin(\alpha_j) \approx \alpha_j$, giving the bounds

$$\frac{1}{j^2\pi^2} \cdot \frac{(2j)^2}{(2j+1)^2} \leq \mu_j \leq \frac{1}{j^2\pi^2} \cdot \frac{(2j)^2}{(2j-1)^2}. \tag{2}$$

These bounds are, however, not sharp. For example, when $j = 1$, we find $4/(9\pi^2) \leq \mu_1 \leq 4/\pi^2$ as $n \to \infty$, which is not very precise. On the other hand, the inequality

$$\frac{1}{4n^2} + \frac{j^2\pi^2}{4n^2(2n+1)^2} \leq \mu_{n-j} \leq \frac{1}{4n^2} + \frac{(j+1)^2\pi^2}{4n^2(2n+1)^2} \tag{3}$$

is precise and useful; see Section 4.

Because none of the eigenvalues of AMA' can be eigenvalues of $A'A$, the eigenvalue μ_j of AMA' is found as the unique solution of

$$\iota'(I_n - \mu_j(A'A)^{-1})^{-1}\iota = n, \qquad \lambda_{j+1} \leq \mu_j \leq \lambda_j; \tag{4}$$

see Lemma A3 in Appendix A.

The eigenvectors of $A'A$ are also known (Dickey and Fuller, 1979). Let s_j be the normalized eigenvector associated with λ_j. The ith element of s_j is given by

$$s_{ij} = \frac{2}{\sqrt{2n+1}} \cos\frac{\pi(2i-1)(2j-1)}{4n+2}, \qquad i = 1, \ldots, n. \tag{5}$$

https://doi.org/10.1017/S0266466608080675 Published online by Cambridge University Press

If we define $S := (s_1, s_2, \ldots, s_n)$, then $S'A'AS = \Lambda := \operatorname{diag}(\lambda_1, \lambda_2, \ldots, \lambda_n)$. Letting $q := S'\iota$, with components $q_1, \ldots, q_n$, we obtain, after some trigonometric simplifications,

$$q_j = \sum_{i=1}^{n} s_{ij} = \frac{2}{\sqrt{2n+1}} \sum_{i=1}^{n} \cos((2i-1)\alpha_j)$$

$$= \frac{1}{\sqrt{2n+1}} (-1)^{j-1} \cot(\alpha_j),$$

which implies that

$$\frac{q_j^2}{n} = \frac{4n^2\lambda_j - 1}{n(2n+1)}. \tag{6}$$

Note that $\iota'(I_n - \mu_j(A'A)^{-1})^{-1}\iota = n$ if and only if $q'(I_n - \mu_j\Lambda^{-1})^{-1}q = n$. The eigenvalues $\mu_1, \ldots, \mu_n$ of AMA' are thus found as the n solutions of

$$\sum_{j=1}^{n} \left(\frac{q_j^2}{n} \cdot \frac{\lambda_j}{\lambda_j - \mu} \right) = 1. \tag{7}$$

But because

$$\frac{q_j^2}{n} \cdot \frac{\lambda_j}{\lambda_j - \mu} = \frac{4n^2\lambda_j - 1}{n(2n+1)} \cdot \frac{\lambda_j}{\lambda_j - \mu}$$

$$= \frac{4n}{2n+1} \left(\lambda_j + \left(\mu - \frac{1}{4n^2} \right) \frac{\lambda_j}{\lambda_j - \mu} \right),$$

and $\sum_j \lambda_j = \operatorname{tr}(A'A) = (n+1)/(2n)$, we have proved the theorem that follows.

THEOREM 2. *The nonzero (positive) eigenvalues $\mu_1 > \cdots > \mu_{n-1}$ of AMA' are the $n-1$ solutions of the equation*

$$n \sum_{j=1}^{n} \frac{1}{g_n(j) - 1/\mu} = \frac{\mu n^2}{4\mu n^2 - 1},$$

where

$$g_n(j) := 4n^2 \sin^2(\alpha_j).$$

https://doi.org/10.1017/S0266466608080675 Published online by Cambridge University Press

3. GENERALIZATION AND PROOF OF THEOREM 1

To prove Theorem 1 we need to demonstrate that Theorem 2 holds asymptotically for $\mu = 1/\pi^2$, that is, that

$$n \sum_{j=1}^{n} \frac{1}{g_n(j) - \pi^2} \to \frac{1}{4}$$

as $n \to \infty$. We shall prove a more general result, namely, that Theorem 2 holds asymptotically for $\mu = 1/(k^2\pi^2)$. This implies that the kth largest eigenvalue μ_k of AMA' converges to $1/(k^2\pi^2)$ as $n \to \infty$. Theorem 1 is the special case when $k = 1$.

THEOREM 3. *Let k be a fixed positive integer and let*

$$S_k(n) := \sum_{j=1}^{n} \frac{1}{g_n(j) - k^2\pi^2}.$$

Then we have

$$nS_k(n) \to \frac{1}{4}$$

as $n \to \infty$.

Proof. The proof is based on Cauchy's residue theorem (see Appendix B) and consists of applying the so-called cotangent method. This method allows us to compute $S_k(n)$ in almost closed form. Let

$$g_n(z) := 4n^2 \sin^2\left(\frac{2z-1}{2n+1} \cdot \frac{\pi}{2}\right), \qquad f(z) := \frac{1}{g_n(z) - k^2\pi^2},$$

where we have suppressed the dependency of f on n. We will apply the theory of Appendix B to the function $f(z)$ on a rectangle Γ_n, with sides parallel to the real and imaginary axes, its vertical sides being intervals on the two lines $L_0 := \{x = \frac{1}{2}\}$ and $L_n := \{x = n + \frac{1}{2}\}$, respectively. It turns out that the integral over Γ_n is of the order $o(1/n)$ as $n \to \infty$. Thus the $S_k(n)$ will appear as a (finite) sum of residues plus a term of order $o(1/n)$.

Recall that $2i \sin(z) = e^{iz} - e^{-iz}$, so that $|\sin(x + iy)| \approx 2e^{|y|}$ for $|y|$ large. Taking orientation into account, it then follows from Remark B1 in Appendix B that we can shift the horizontal sides of our rectangle to ∞. Defining $F(z) := \pi \cot(\pi z) f(z)$, this gives

$$\int_{\Gamma_n} F(z)\, dz = \int_{L_n} F(z)\, dz - \int_{L_0} F(z)\, dz.$$

https://doi.org/10.1017/S0266466608080675 Published online by Cambridge University Press

On L_n we have $z = n + \frac{1}{2} + iy$, so that $\cot(\pi z) = -\tan(i\pi y)$, which is bounded by one in absolute value. Also,

$$g_n\left(n + \frac{1}{2} + iy\right) = 4n^2 \sin^2\left(\frac{2n + 2iy}{2n + 1} \cdot \frac{\pi}{2}\right)$$

$$= 4n^2 \cos^2\left(\frac{-1 + 2iy}{2n + 1} \cdot \frac{\pi}{2}\right). \tag{8}$$

We now show that $n\int_{L_n} F(z) \to 0$ as $n \to \infty$. Using the fact that $\cot(\pi z) = -\tan(i\pi y)$ and (8), we substitute $y = ns/\pi$ and obtain

$$n\int_{L_n} F(z)\,dz = n\int_{-\infty}^{\infty} \frac{-\pi \tan(i\pi y)}{4n^2 \cos^2\left(\left(\frac{-1}{2n+1} + \frac{2iy}{2n+1}\right)\frac{\pi}{2}\right) - k^2\pi^2}\, i\,dy$$

$$= \int_{-\infty}^{\infty} \frac{-\tan(ins)}{4\cos^2\left(\frac{-\pi}{2(2n+1)} + \frac{ins}{2n+1}\right) - \left(\frac{k\pi}{n}\right)^2}\, i\,ds.$$

Lebesgue's dominated convergence theorem applies so that we may interchange limit and integral. Thus,

$$\lim_{n\to\infty} n\int_{L_n} F(z)\,dz = \int_{-\infty}^{\infty} \frac{\operatorname{sgn}(s)}{4\cos^2(is/2)}\,ds = 0,$$

using the fact that $\lim_{n\to\infty} -i\tan(ins) = \operatorname{sgn}(s)$, the sign of s. Similarly, the integral over L_0 is $o(1/n)$.

It remains to compute the residues of F at the singular points that come from f, that is, at the zeros of $g_n(z) - k^2\pi^2$ that are located inside Γ_n. We write

$$g_n(z) - k^2\pi^2 = \left(2n \sin\left(\frac{2z-1}{2n+1} \cdot \frac{\pi}{2}\right) + k\pi\right)\left(2n \sin\left(\frac{2z-1}{2n+1} \cdot \frac{\pi}{2}\right) - k\pi\right).$$

The zeros of this equation are solutions of

$$\frac{2z-1}{2n+1} \cdot \frac{\pi}{2} = \pm \arcsin(k\pi/2n),$$

which we rewrite as

$$z = \frac{1}{2} \pm \frac{2n+1}{\pi} \arcsin(k\pi/2n) \approx \frac{1}{2} \pm \frac{k(2n+1)}{2n} \mod(2n+1).$$

https://doi.org/10.1017/S0266466608080675 Published online by Cambridge University Press

Of these solutions only the one close to $\frac{1}{2} + k$ is inside Γ_n. We compute the residue at the solution $z_1 \approx \frac{1}{2} + k$, using Remark B2, l'Hôpital's rule, and the fact that $2n \sin((2z_1 - 1)/(2n + 1) \cdot (\pi/2)) = k\pi$. This gives

$$\begin{aligned}
\mathrm{Res}(F, z_1) &= \lim_{z \to z_1} \frac{z - z_1}{2n \sin\left(\dfrac{2z-1}{2n+1} \cdot \dfrac{\pi}{2}\right) - k\pi} \cdot \frac{\pi \cot(\pi z)}{2n \sin\left(\dfrac{2z-1}{2n+1} \cdot \dfrac{\pi}{2}\right) + k\pi} \\
&= \frac{1}{2n \dfrac{\pi}{2n+1} \cos\left(\dfrac{2z_1-1}{2n+1} \cdot \dfrac{\pi}{2}\right)} \cdot \frac{\pi \cot(\pi z_1)}{2n \sin\left(\dfrac{2z_1-1}{2n+1} \cdot \dfrac{\pi}{2}\right) + k\pi} \\
&= \frac{1}{\dfrac{2n\pi}{2n+1} \sqrt{1 - \left(\dfrac{k\pi}{2n}\right)^2}} \cdot \frac{\pi \cot(\pi z_1)}{2k\pi}.
\end{aligned}$$

Hence the limit becomes

$$\begin{aligned}
\lim_{n \to \infty} n \, \mathrm{Res}(F, z_1) &= \lim_{n \to \infty} n \frac{\cot(\pi z_1)}{2k\pi} \\
&= \lim_{n \to \infty} \frac{-n \tan((2n+1)\arcsin(k\pi/(2n)))}{2k\pi} = -\frac{1}{4}.
\end{aligned}$$

Then, using (B.4) in Appendix B, we conclude that $\lim_{n \to \infty} n S_k(n) = \frac{1}{4}$, and this completes the proof of Theorem 3. ■

4. FURTHER DISCUSSION

In fact we can say a little more. Theorem 3 tells us that the jth largest eigenvalue μ_j of AMA' converges to $1/(j^2\pi^2)$ as $n \to \infty$. But it does not tell us how fast this convergence takes place, nor does it tell us what happens to the whole set of eigenvalues. Further analysis, not provided here, shows that, for $n \geq 4$,

$$\mu_j = \frac{1}{j^2\pi^2} + \frac{c_j}{n^2\pi^2} \qquad (j = 1, \ldots, n-1), \tag{9}$$

where, for each given n, c_j is a slowly increasing function of j bounded by $\underline{c} \leq c_j \leq \bar{c}$, with

$$\underline{c} := \lim_{n \to \infty} n^2(\pi^2\mu_1 - 1) = \frac{\pi^2}{12} \approx 0.8225$$

https://doi.org/10.1017/S0266466608080675 Published online by Cambridge University Press

and

$$\bar{c} := \lim_{n\to\infty} n^2\pi^2\mu_{n-1} - 1 = \frac{\pi^2}{4} - 1 \approx 1.4674$$

and satisfying $(1/n)\sum_j c_j \to 1$. In particular,

$$\max_{1\le j\le n-1}\left|\mu_j - \frac{1}{j^2\pi^2}\right| \le \frac{1}{n^2}\cdot\frac{\pi^2-4}{4\pi^2} \approx \frac{0.1487}{n^2}.$$

Hence we may say that "the set of eigenvalues $\{\mu_j\}$ of AMA' converges to the set $\{\mu_j^*\}$," where $\mu_j^* = 1/(j^2\pi^2)$ for $j = 1,\ldots,n-1$ and $\mu_n^* = 0$. Some caution is however required in interpreting this statement. For fixed j it means that μ_j approaches $1/(j^2\pi^2)$. But for j dependent on n, it only means that the difference approaches zero. Thus, $\mu_{n-j} \approx 1/(4n^2)$ (see eqn. (3)), whereas $\mu_{n-j}^* = 1/((n-j)^2\pi^2) \approx 1/(n^2\pi^2)$. Their difference is of order $O(1/n^2)$, but μ_{n-j} is better approximated by $1/(4n^2)$ than by μ_{n-j}^*.

We note that our findings agree with the facts that

$$\sum_{j=1}^{n}\mu_j = \mathrm{tr}(AMA') = \frac{n^2-1}{6n^2} = \frac{1}{6} + O\left(\frac{1}{n^2}\right)$$

and

$$\sum_{j=1}^{n}\mu_j^2 = \mathrm{tr}(AMA')^2 = \frac{(n^2-1)(2n^2+7)}{180n^4} = \frac{1}{90} + \frac{1}{36n^2} + O\left(\frac{1}{n^4}\right),$$

while we also know that

$$\sum_{j=1}^{n}\frac{1}{j^2} = \frac{\pi^2}{6} - \frac{1}{n} + O\left(\frac{1}{n^2}\right), \qquad \sum_{j=1}^{n}\frac{1}{j^4} = \frac{\pi^4}{90} + O\left(\frac{1}{n^3}\right).$$

Hence we obtain

$$n\left(\sum_{j=1}^{n}\mu_j - \sum_{j=1}^{n}\frac{1}{j^2\pi^2}\right) = \frac{1}{\pi^2} + O\left(\frac{1}{n}\right)$$

and

$$n^2\left(\sum_{j=1}^{n}\mu_j^2 - \sum_{j=1}^{n}\frac{1}{j^4\pi^4}\right) = \frac{1}{36} + O\left(\frac{1}{n}\right).$$

We conclude by noting that Theorem 1 also provides an asymptotic bound for various statistics related to ξ_n. Suppose first that M is replaced by M_Z, the projection matrix orthogonal to some collection of nonstochastic regressors Z, including the intercept. The obvious example is the inclusion of the linear trend.

https://doi.org/10.1017/S0266466608080675 Published online by Cambridge University Press

Because $M_Z = MM_ZM$, Theorem 11.11 of Magnus and Neudecker (1988) may be redeployed to show that the ordered eigenvalues of AM_ZA' are bounded by the corresponding eigenvalues of AMA'. On the other hand, if M is replaced by I so that we consider the version of ξ_n obtained from noncentered data, the exact bound for the ratio $x'A'Ax/x'x$ is provided, directly from (1), by the case $j = 1$ of the limits

$$\lambda_j \to \frac{1}{\pi^2\left(j - \frac{1}{2}\right)^2}.$$

As Breitung (2002) points out, ξ_n corresponds to n^{-1} times the so-called KPSS statistic of Kwiatkowski, Phillips, Schmidt, and Shin (1992), except that the kernel estimator of the long-run variance in the denominator of the latter statistic is replaced by the simple variance of the sample. A recently proposed variant of the KPSS test is the V/S test of Giraitis, Kokoszka, Leipus, and Teyssière (2003), in which the the numerator of the ratio is expressed in mean deviation form. The corresponding modification of Breitung's statistic takes the form $\xi_n^* = x'MA'MAMx/x'Mx$. If $\xi_n^* \to_d \xi^*$, observe that when, in particular, x is an I(1) vector,

$$\xi^* = \frac{\int_0^1\left(\int_0^t W\,dr - t\int_0^1 W\,dr\right)^2 dt - \left[\int_0^1\left(\int_0^t W\,dr - t\int_0^1 W\,dr\right)dt\right]^2}{\int_0^1 W^2\,dt - \left(\int_0^1 W\,dt\right)^2}.$$

In view of the general inequality $0 \le \xi_n^* \le \xi_n$ we are able to say that Theorem 1 also applies to ξ^*.

REFERENCES

Abadir, K.M. & J.R. Magnus (2005) *Matrix Algebra*. Cambridge University Press.

Breitung, J. (2002) Nonparametric tests for unit roots and cointegration. *Journal of Econometrics* 108, 343–363.

Conway, J.B. (1978) *Functions of One Complex Variable I*, 2nd ed., Graduate Texts in Mathematics. Springer-Verlag.

Davidson, J. (2008) When is a time series I(0)? In N. Shephard & J. Castle (eds.), *The Methodology and Practice of Econometrics*, Oxford University Press. Forthcoming.

Dickey, D.A. & W.A. Fuller (1979) Distribution of the estimates for autoregressive time series with a unit root. *Journal of the American Statistical Association* 74, 427–431.

Giraitis, L., P. Kokoszka, R. Leipus, & G. Teyssière (2003) Rescaled variance and related tests for long memory in volatility and levels. *Journal of Econometrics* 112, 265–294.

Kwiatkowski, D., P.C.B. Phillips, P. Schmidt, & Y. Shin (1992) Testing the null hypothesis of stationarity against the alternative of a unit root: How sure are we that economic time series have a unit root? *Journal of Econometrics* 54, 159–178.

Magnus, J.R. & H. Neudecker (1988) *Matrix Differential Calculus with Applications in Statistics and Econometrics*. Wiley. Rev. ed., 1999.

https://doi.org/10.1017/S0266466608080675 Published online by Cambridge University Press

Rutherford, D.E. (1946) Some continuant determinants arising in physics and chemistry. *Proceedings of the Royal Society of Edinburgh, Section A* 62, 229–236.
Tanaka, K. (1996) *Time Series Analysis: Nonstationary and Noninvertible Distribution Theory*. Wiley.

APPENDIX A: The Matrix $V - \omega\omega'$

We consider a positive semidefinite $n \times n$ matrix V and an $n \times 1$ vector ω.

LEMMA A1. *We have*

$$|V - \omega\omega'| = \begin{cases} |V|(1 - \omega' V^{-1}\omega) & \text{if } \mathrm{rk}(V) = n; \\ -(\omega' x)^2 \cdot p(V) & \text{if } rk(V) = n-1, Vx = 0, x'x = 1; \\ 0 & \text{if } rk(V) \le n-2, \end{cases}$$

where $p(V)$ denotes the product of the nonzero eigenvalues of V.

Proof. We prove the lemma first for a diagonal $n \times n$ matrix Λ with nonnegative diagonal elements and an $n \times 1$ vector a.

(i) If all diagonal elements of Λ are nonzero (hence positive), then

$$|\Lambda - aa'| = |\Lambda^{1/2}(I_n - \Lambda^{-1/2} aa' \Lambda^{-1/2})\Lambda^{1/2}| = |\Lambda|(1 - a'\Lambda^{-1}a).$$

(ii) If one of the diagonal elements of Λ is zero (say, the nth), then we partition

$$\Lambda = \begin{pmatrix} \Lambda_1 & 0 \\ 0 & 0 \end{pmatrix}, \qquad a = \begin{pmatrix} a_1 \\ a_2 \end{pmatrix}.$$

If $a_2 \neq 0$, then

$$|\Lambda - aa'| = \begin{vmatrix} \Lambda_1 - a_1 a_1' & -a_2 a_1 \\ -a_2 a_1' & -a_2^2 \end{vmatrix} = -a_2^2 |\Lambda_1|;$$

see Exercise 5.30(b) in Abadir and Magnus (2005). The result remains true when $a_2 = 0$, because both sides of the equality are then zero.

(iii) If two or more diagonal elements of Λ are zero, then

$$\mathrm{rk}(\Lambda - aa') \le \mathrm{rk}(\Lambda) + \mathrm{rk}(aa') \le (n-2) + 1 = n - 1,$$

and hence $|\Lambda - aa'| = 0$.

In the general case, we diagonalize V as $S'VS = \Lambda$ and define $a := S'\omega$. The results (i) and (iii) follow immediately. For (ii) we partition $S = (S_1, x)$, so that $\omega' x = a' S' x = a_1' S_1' x + a_2 x' x = a_2$. ■

https://doi.org/10.1017/S0266466608080675 Published online by Cambridge University Press

LEMMA A2. $|V - \omega\omega'| = 0$ *if and only if*

$$\begin{cases} \text{rk}(V) = n, & \omega' V^{-1}\omega = 1; \\ \text{rk}(V) = n-1, & \omega' x = 0, \quad Vx = 0; \\ \text{rk}(V) \leq n-2. \end{cases}$$

Proof. This follows directly from Lemma A1. ■

LEMMA A3. *Let C be a nonsingular $n \times n$ matrix and let $M := I_n - (1/n)\iota\iota'$. Then, μ is an eigenvalue of CMC' if and only if*

$$\begin{cases} \mu \text{ is not an eigenvalue of } CC', \ \iota'(I_n - \mu(C'C)^{-1})^{-1}\iota = n; \\ \mu \text{ is a simple eigenvalue of } CC' \text{ with associated eigenvector } x, \ \iota' C' x = 0; \\ \mu \text{ is a multiple eigenvalue of } CC'. \end{cases}$$

Proof. The eigenvalues of CMC' are given by

$$\left| CC' - \mu I_n - \frac{1}{n} C\iota\iota' C' \right| = 0.$$

Let $V =: CC' - \mu I_n$ and $\omega := C\iota/\sqrt{n}$ and apply Lemma A2. ■

APPENDIX B: Cauchy's Residue Theorem and the Cotangent Method

In this Appendix we state some results from complex function theory that are required in the proof of Theorem 3; see Conway (1978). Our proof depends heavily on the "cotangent method." Although this method is "well known," it is not easy to find a good reference for it. (Conway, 1978, has an exercise on p. 122.) Hence it will prove useful to state this method explicitly. Our starting point is Cauchy's famous theorem.

THEOREM B1 (Cauchy's residue theorem). *Let f be a function defined on a domain D and its boundary Γ. Assume that f is holomorphic (i.e., complex differentiable) except for a finite set of singular points $a_1, \ldots, a_k$ in D. Then,*

$$\frac{1}{2\pi i} \int_\Gamma f(z)\, dz = \sum_{h=1}^{k} \text{Res}(f, a_h), \tag{B.1}$$

where $\text{Res}(f, a)$ *denotes the residue of f at an isolated singularity a and is defined as*

$$\text{Res}(f, a) := \lim_{\epsilon \to 0} \frac{1}{2\pi i} \int_{|z-a|=\epsilon} f(z)\, dz.$$

https://doi.org/10.1017/S0266466608080675 Published online by Cambridge University Press

We note that, for ϵ sufficiently small, the preceding integral becomes independent of ϵ. Moreover, one can show that f may be developed into a convergent series near a:

$$f(z) = \sum_{l=-\infty}^{\infty} c_l (z-a)^l, \tag{B.2}$$

so that $\mathrm{Res}(f,a) = c_{-1}$.

Remark B1. If there are no singularities inside the boundary Γ, then the integral in (B.1) equals zero. This allows for changing the path of integration without altering the value of the integral as long as we do not "cross" any singularities.

Remark B2. It follows from (B.2) that if f has a singular point at a and if $L := \lim_{z\to a} f(z)(z-a)$ exists, then $\mathrm{Res}(f,a) = L$.

The cotangent method is based on the following expansion:

$$\pi \cot(\pi z) = \sum_{j=-n}^{n} \frac{1}{z-j} + g_n(z) = \sum_{j=-\infty}^{\infty} \frac{1}{z-j}.$$

Here g_n is a holomorphic function on $\mathbb{C}$ except for singularities at $\pm(n+1), \pm(n+2), \ldots$. (The last equality should be treated with care, because the sum is only conditionally convergent.)

Now, if f is a holomorphic function with singularities at $a_1, \ldots, a_k$, say, all less than n and none of which are integers, and if we let $F(z) := \pi \cot(\pi z) f(z)$, then by Theorem B1,

$$\frac{1}{2\pi i} \int_{\Gamma_n} F(z)\, dz = \sum_{a\in A} \mathrm{Res}(F,a), \tag{B.3}$$

where Γ_n is a square in $\mathbb{C}$ with vertices $(\pm(n+\frac{1}{2}), \pm(n+\frac{1}{2})i)$ and A is the set of singularities of F inside the square Γ_n. Because the residue $\mathrm{Res}(F,j)$ at an integer j equals $f(j)$, we can rewrite (B.3) as

$$\sum_{j=-n}^{n} f(j) = \frac{1}{2\pi i} \int_{\Gamma_n} F(z)\, dz - \sum_{h=1}^{k} \mathrm{Res}(F, a_h). \tag{B.4}$$

This is the cotangent method. Observing that the cotangent is uniformly bounded on the squares Γ_n, one can often show that the integral in (B.4) tends to zero as $n \to \infty$. For suitable f with finitely many singularities one can then compute the limit of the left-hand sum in (B.4).

https://doi.org/10.1017/S0266466608080675 Published online by Cambridge University Press