Canad. Math. Bull. Vol. **69** (1), 2026, pp. 137–141
http://dx.doi.org/10.4153/S0008439525100775
© The Author(s), 2025. Published by Cambridge University Press on behalf of Canadian Mathematical Society

B_h-sets of real and complex numbers

Melvyn B. Nathanson

Abstract. Let $K = \mathbf{R}$ or $\mathbf{C}$. An n-element subset A of K is a B_h-set if every element of K has at most one representation as the sum of h not necessarily distinct elements of A. Associated with the B_h-set $A = \{a_1, \ldots, a_n\}$ are the B_h-vectors $\mathbf{a} = (a_1, \ldots, a_n)$ in K^n. This article proves that "almost all" n-element subsets of K are B_h-sets in the sense that the set of all B_h-vectors is a dense open subset of K^n.

1 Sumsets and B_h-sets

Let A be a nonempty subset of an additive abelian group or semigroup G. For every positive integer h, the h-fold *sumset* of A is the set of all sums of h not necessarily distinct elements of A:

$$hA = \underbrace{A + \cdots + A}_{h \text{ summands}} = \{a'_1 + \cdots + a'_h : a'_i \in A \text{ for all } i \in [1, h]\}.$$

For integers $h \geq 2$ and $n \geq 2$, let

$$\mathcal{X}_{h,n} = \left\{ \mathbf{x} = (x_1, \ldots, x_n) \in \mathbf{N}_0^n : \sum_{i=1}^{n} x_i = h \right\}.$$

For all $\mathbf{x} = (x_1, \ldots, x_n) \in \mathcal{X}_{h,n}$ and $\mathbf{y} = (y_1, \ldots, y_n) \in \mathcal{X}_{h,n}$, we have

$$\|\mathbf{x} - \mathbf{y}\|_\infty = \max\{|x_i - y_i| : i = 1, \ldots, n\} \leq h. \tag{1.1}$$

Let $\mathbf{a} = (a_1, \ldots, a_n) \in G^n$ be a vector with distinct coordinates. Associated with the vector $\mathbf{a}$ is the subset $A = \{a_1, \ldots, a_n\}$ of G of cardinality n. For all $\mathbf{x} \in \mathcal{X}_{h,n}$ and $\mathbf{a} \in G^n$, we define the dot product

$$\mathbf{x} \cdot \mathbf{a} = \sum_{i=1}^{n} x_i a_i.$$

Then $\mathbf{x} \cdot \mathbf{a} \in G$. The h-fold sumset of A can be written in the form

$$hA = \{\mathbf{x} \cdot \mathbf{a} : \mathbf{x} \in \mathcal{X}_{h,n}\}.$$

Received by the editors March 5, 2025; revised March 5, 2025; accepted May 29, 2025.
Published online on Cambridge Core June 11, 2025.
Supported in part by PSC-CUNY Research Award Program grant 66197-00 54.
AMS subject classification: **11B05**, **11B13**, **11B30**, **11B34**, **11B75**, **54E52**.
Keywords: Sidon set, B_h-set, open dense subsets, Baire's theorem, additive number theory.

https://doi.org/10.4153/S0008439525100775 Published online by Cambridge University Press

The subset A of G is called a B_h-*set* if every element of G has at most one representation (up to permutation of the summands) as the sum of h not necessarily distinct elements of A. B_2-sets are also called *Sidon sets*. The study of B_h-sets of integers is a classical topic in combinatorial additive number theory (cf. [1–19]).

This article studies B_h-sets of real and complex numbers. Let $K = \mathbf{R}$ or $\mathbf{C}$. Associated with every vector $\mathbf{a} = (a_1, \ldots, a_n) \in K^n$ with distinct coordinates is the subset $A = \{a_1, \ldots, a_n\}$ of K of cardinality n. The vector $\mathbf{a}$ is called a B_h-*vector* if its associated set A is a B_h-set. Let $\mathcal{B}_h$ be the set of B_h-vectors. If $\mathbf{a} = (a_1, a_2, \ldots, a_n) \in \mathcal{B}_h$, then $\sigma\mathbf{a} = \left(a_{\sigma(1)}, a_{\sigma(2)}, \ldots, a_{\sigma(n)}\right) \in \mathcal{B}_h$ for every permutation σ of $\{1, \ldots, n\}$.

The object of this article is to prove that "almost all" n-element subsets of K are B_h-sets in the sense that the set $\mathcal{B}_h$ is an open and dense subset of K^n.

2 Open and dense subsets

Theorem 1 *Let $K = \mathbf{R}$ or $\mathbf{C}$. The set $\mathcal{B}_h$ is an open subset of K^n.*

Proof Let $\mathbf{a} = (a_1, \ldots, a_n) \in K^n$ be a B_h-vector. If $\mathbf{x}, \mathbf{y} \in \mathcal{X}_{h,n}$ with $\mathbf{x} \neq \mathbf{y}$, then $\mathbf{x} \cdot \mathbf{a} \neq \mathbf{y} \cdot \mathbf{a}$ and so

$$0 < \Delta = \min\left\{\|(\mathbf{x} - \mathbf{y}) \cdot \mathbf{a}\|_\infty : (\mathbf{x}, \mathbf{y}) \in \mathcal{X}_{h,n}^2 \text{ and } \mathbf{x} \neq \mathbf{y}\right\}.$$

We shall prove that, for all vectors $\mathbf{b} \in K^n$ with

$$0 < \|\mathbf{b}\|_\infty < \frac{\Delta}{h},$$

the vector

$$\mathbf{a} + \mathbf{b} = \mathbf{c}$$

is a B_h-vector and so the set of B_h-vectors contains the open ball with center at $\mathbf{a}$ and radius Δ/h.

Let $\mathbf{x}, \mathbf{y} \in \mathcal{X}_{h,n}$ with $\mathbf{x} \neq \mathbf{y}$. If $\mathbf{x} \cdot \mathbf{c} = \mathbf{y} \cdot \mathbf{c}$, then

$$\mathbf{x} \cdot (\mathbf{a} + \mathbf{b}) = \mathbf{y} \cdot (\mathbf{a} + \mathbf{b})$$

and

$$(\mathbf{x} - \mathbf{y}) \cdot \mathbf{a} = (\mathbf{y} - \mathbf{x}) \cdot \mathbf{b}.$$

Applying inequality (1.1), we obtain

$$\begin{aligned} \Delta \leq \|(\mathbf{x} - \mathbf{y}) \cdot \mathbf{a}\|_\infty &= \|(\mathbf{y} - \mathbf{x}) \cdot \mathbf{b}\|_\infty \\ &\leq \|\mathbf{y} - \mathbf{x}\|_\infty \|\mathbf{b}\|_\infty \\ &< h\left(\frac{\Delta}{h}\right) = \Delta \end{aligned}$$

which is absurd. Therefore, $\mathbf{x} \neq \mathbf{y}$ implies $\mathbf{x} \cdot \mathbf{c} \neq \mathbf{y} \cdot \mathbf{c}$ and so $\mathbf{c}$ is a B_h-vector. This completes the proof. ∎

Lemma 1 *For all $\delta > 0$ there is a B_h-vector $\mathbf{b}$ such that $\|\mathbf{b}\|_\infty < \delta$.*

https://doi.org/10.4153/S0008439525100775 Published online by Cambridge University Press

Proof If $\mathbf{w} = (w_1, \ldots, w_n)$ is any B_h-vector in $\mathbf{R}^n$ or $\mathbf{C}^n$ with associated B_h-set $W = \{w_1, \ldots, w_n\}$, then, for every $\lambda \neq 0$, the "contraction" $\lambda * W = \{\lambda w_i : i = 1, \ldots, n\}$ is a B_h-set and the corresponding vector $\lambda\mathbf{w} = (\lambda w_1, \ldots, \lambda w_n)$ is also a B_h-vector. Choosing $0 < \lambda < \delta/\|\mathbf{w}\|_\infty$ and $\mathbf{b} = \lambda\mathbf{w}$ gives

$$\|\mathbf{b}\|_\infty = \|\lambda\mathbf{w}\|_\infty = |\lambda|\|\mathbf{w}\|_\infty < \left(\frac{\delta}{\|\mathbf{w}\|_\infty}\right)\|\mathbf{w}\|_\infty = \delta.$$

This completes the proof. ∎

Theorem 2 *Let $K = \mathbf{R}$ or $\mathbf{C}$. The set $\mathcal{B}_h$ is a dense subset of K^n.*

Proof Let $\mathbf{a} \in K^n$ be a vector that is not a B_h-vector. We shall prove that, for every $\varepsilon > 0$, there is a B_h-vector $\mathbf{c} \in K^n$ such that $\|\mathbf{c} - \mathbf{a}\|_\infty < \varepsilon$.

We partition the set of pairs of distinct vectors in $\mathcal{X}_{h,n}$ into two disjoint finite sets as follows:

$$\mathcal{U} = \{(\mathbf{x}, \mathbf{y}) \in \mathcal{X}_{h,n}^2 : \mathbf{x} \neq \mathbf{y} \text{ and } \mathbf{x} \cdot \mathbf{a} \neq \mathbf{y} \cdot \mathbf{a}\}$$

and

$$\mathcal{V} = \{(\mathbf{x}, \mathbf{y}) \in \mathcal{X}_{h,n}^2 : \mathbf{x} \neq \mathbf{y} \text{ and } \mathbf{x} \cdot \mathbf{a} = \mathbf{y} \cdot \mathbf{a}\}.$$

The set $\mathcal{V}$ is nonempty because $\mathbf{a}$ is not a B_h-vector. The set $\mathcal{U}$ is nonempty because $n \geq 2$ and $(\mathbf{x}, \mathbf{y}) \in \mathcal{U}$ with $\mathbf{x} = (h, 0, 0, \ldots, 0)$ and $\mathbf{y} = (0, h, 0, \ldots, 0)$. Then

$$0 < \Delta = \min\{\|(\mathbf{x} - \mathbf{y}) \cdot \mathbf{a}\|_\infty : (\mathbf{x}, \mathbf{y}) \in \mathcal{U}\}.$$

By Lemma 1, for all $\varepsilon > 0$ there is a B_h-vector $\mathbf{b}$ in K^n such that

$$\|\mathbf{b}\|_\infty < \min\left(\varepsilon, \frac{\Delta}{h}\right).$$

Applying inequality (1.1), for all pairs $(\mathbf{x}, \mathbf{y}) \in \mathcal{X}_{h,n}^2$, we have

$$\|(\mathbf{x} - \mathbf{y}) \cdot \mathbf{b}\|_\infty \leq \|\mathbf{x} - \mathbf{y}\|_\infty \|\mathbf{b}\|_\infty < h\left(\frac{\Delta}{h}\right) = \Delta.$$

The vector

$$\mathbf{a} + \mathbf{b} = \mathbf{c}$$

satisfies

$$\|\mathbf{c} - \mathbf{a}\|_\infty = \|\mathbf{b}\|_\infty < \varepsilon.$$

We shall prove that $\mathbf{c}$ is a B_h-vector. Equivalently, we shall prove that $\mathbf{x} \cdot \mathbf{c} \neq \mathbf{y} \cdot \mathbf{c}$ for all pairs $(\mathbf{x}, \mathbf{y}) \in \mathcal{X}_{h,n}^2$ with $\mathbf{x} \neq \mathbf{y}$.

If $(\mathbf{x}, \mathbf{y}) \in \mathcal{U}$, then $\mathbf{x} \cdot \mathbf{a} \neq \mathbf{y} \cdot \mathbf{a}$ and

$$\begin{aligned}\|\mathbf{x} \cdot \mathbf{c} - \mathbf{y} \cdot \mathbf{c}\|_\infty &= \|(\mathbf{x} \cdot \mathbf{a} - \mathbf{y} \cdot \mathbf{a}) + (\mathbf{x} \cdot \mathbf{b} - \mathbf{y} \cdot \mathbf{b})\|_\infty \\ &\geq \|(\mathbf{x} - \mathbf{y}) \cdot \mathbf{a}\|_\infty - \|(\mathbf{x} - \mathbf{y}) \cdot \mathbf{b}\|_\infty \\ &> \Delta - \Delta = 0.\end{aligned}$$

https://doi.org/10.4153/S0008439525100775 Published online by Cambridge University Press

If $(\mathbf{x}, \mathbf{y}) \in \mathcal{V}$, then $\mathbf{x} \cdot \mathbf{a} = \mathbf{y} \cdot \mathbf{a}$. Because $\mathbf{x} \neq \mathbf{y}$ and $\mathbf{b}$ is a B_h-vector, we have $\mathbf{x} \cdot \mathbf{b} \neq \mathbf{y} \cdot \mathbf{b}$. It follows that

$$\begin{aligned} \|\mathbf{x} \cdot \mathbf{c} - \mathbf{y} \cdot \mathbf{c}\|_\infty &= \|(\mathbf{x} \cdot \mathbf{a} - \mathbf{y} \cdot \mathbf{a}) + (\mathbf{x} \cdot \mathbf{b} - \mathbf{y} \cdot \mathbf{b})\|_\infty \\ &= \|\mathbf{x} \cdot \mathbf{b} - \mathbf{y} \cdot \mathbf{b}\|_\infty \\ &> 0. \end{aligned}$$

This completes the proof. ∎

Combining Theorems 1 and 2 gives the following result.

Theorem 3 *Let $K = \mathbf{R}$ or $\mathbf{C}$. The set $\mathcal{B}_h$ is a dense open subset of K^n.*

Theorem 4 *Let $\mathcal{B}_h$ be the set of B_h-vectors in K^n. The set*

$$\mathcal{B}_\infty = \bigcap_{h=1}^{\infty} \mathcal{B}_h$$

is a dense subset of K^n.

Proof This is simply an application of Baire's theorem in functional analysis. ∎

3 $B_h[g]$-sets

Let g and h be positive integers. The subset A of an additive abelian semigroup G is called a $B_h[g]$-*set* if every element of G has at most g representations as the sum of h not necessarily distinct elements of A. The $B_2[1]$-sets are the B_2-sets. If A is a $B_h[g]$-set, then A is also a $B_h[g']$-set for all $g' \geq g$. In particular, every B_h-set is a $B_h[g]$-set.

Let $K = \mathbf{R}$ or $\mathbf{C}$. The vector $\mathbf{a} = (a_1, \ldots, a_n) \in K^n$ is a $B_h[g]$-*vector* if the set $A = \{a_1, \ldots, a_n\}$ is an n-element $B_h[g]$-set in K. Every B_h-vector is a $B_h[g]$-vector. Because the set of B_h-vectors is dense in K^n, it follows that the set of $B_h[g]$-vectors is also dense in K^n.

Problem Is the set of $B_h[g]$-vectors also open in K^n?

References

[1] R. C. Bose and S. Chowla, *Theorems in the additive theory of numbers*. Comment. Math. Helv. **37**(1962/63), 141–147.

[2] Y. C. Cheng, *Greedy Sidon sets for linear forms*. J. Number Theory. **266**(2025), 225–248.

[3] J. Cilleruelo, I. Ruzsa, and C. Vinuesa, *Generalized Sidon sets*. Adv. Math. **225**(2010), no. 5, 2786–2807.

[4] J. Cilleruelo, I. Z. Ruzsa, and C. Trujillo, *Upper and lower bounds for finite $B_h[g]$ sequences*. J. Number Theory **97**(2002), no. 1, 26–34.

[5] J. A. Dias da Silva and M. B. Nathanson, *Maximal Sidon sets and matroids*. Discrete Math. **309**(2009), no. 13, 4489–4494.

[6] P. Erdős and P. Turán, *On a problem of Sidon in additive number theory, and on some related problems*. J. London Math. Soc. **16**(1941), 212–215.

[7] M. N. Kolountzakis, *The density of $B_h[g]$ sequences and the minimum of dense cosine sums*. J. Number Theory **56**(1996), no. 1, 4–11.

[8] G. Martin and K. O'Bryant, *Constructions of generalized Sidon sets*. J. Combin. Theory Ser. A. **113**(2006), no. 4, 591–607.

https://doi.org/10.4153/S0008439525100775 Published online by Cambridge University Press

[9] M. B. Nathanson, *On the ubiquity of Sidon sets*. In: D. V. Chudnovsky, G. V. Chudnovsky, and M. B. Nathanson (eds.), Number theory (New York, 2003), Springer, New York, 2004, pp. 263–272.
[10] M. B. Nathanson, *Sidon sets and perturbations*. In: M. B. Nathanson (ed.) Combinatorial and Additive Number Theory IV, Springer Proc. Math. Stat., 347, Springer, Cham, 2021, pp. 401–408.
[11] M. B. Nathanson, *The Bose-Chowla argument for Sidon sets*. J. Number Theory. **238**(2022), 133–146.
[12] M. B. Nathanson, *An inverse problem for finite Sidon sets*, M. B. Nathanson (ed.) Combinatorial and additive number theory V, Springer Proc. Math. Stat., vol. 395, Springer, Cham, 2022, pp. 277–285. MR 4539832
[13] M. B. Nathanson, *Sidon sets for linear forms*. J. Number Theory. **239**(2022), 207–227.
[14] M. B. Nathanson, *The third positive element in a greedy B_h-set*. Palestine J. Math. **14**(2025), 213–216.
[15] M. B. Nathanson, **Q**-*independence and the construction of B_h-sets of integers and lattice points.* Discrete Math. (2025), to appear.
[16] M. B. Nathanson and K. O'Bryant, *The fourth positive element in the greedy B_h-set*. J. Integer Seq. **27**(2024), Article 24.7.3, pages 1–10.
[17] K. O'Bryant, *A complete annotated bibliography of work related to Sidon sequences*. Electron. J. Combin. **DS11**(2004), 39.
[18] K. O'Bryant, *Constructing thick B_h-sets*. J. Integer Seq. **27**(2024), no. 1, Paper No. 24.1.2, 17.
[19] I. Z. Ruzsa, *Solving a linear equation in a set of integers*. I. Acta Arith. **65**(1993), no. 3, 259–282.

Department of Mathematics, Lehman College (CUNY), Bronx, NY 10468, United States
e-mail: melvyn.nathanson@lehman.cuny.edu

https://doi.org/10.4153/S0008439525100775 Published online by Cambridge University Press