J. London Math. Soc. (2) 72 (2005) 91–109 © 2005 London Mathematical Society
doi:10.1112/S0024610705006423

DISCRETIZATION METHODS FOR HOMOGENEOUS FRAGMENTATIONS

JEAN BERTOIN AND ALAIN ROUAULT

Abstract

Homogeneous fragmentations describe the evolution of a unit mass that breaks down randomly into pieces as time passes. They can be thought of as continuous time analogues of a certain type of branching random walk, which suggests the use of time-discretization to shift known results from the theory of branching random walks to the fragmentation setting. In particular, this yields interesting information about the asymptotic behaviour of fragmentations.

On the other hand, homogeneous fragmentations can also be investigated using a powerful technique of discretization of space due to Kingman, namely, the theory of exchangeable partitions of $\mathbb{N}$. Spatial discretization is especially well suited to the direct development for continuous times of the conceptual method of probability tilting of Lyons, Pemantle and Peres.

1. *Introduction*

Homogeneous fragmentations form a family of random processes in continuous time which were introduced in [**4**]. Roughly, these are particle systems that model a mass that breaks down randomly into pieces as time passes. More precisely, each particle is identified with its mass (that is, it is specified by a positive real number), and the fragmentation property requires that different particles have independent evolutions. The homogeneity property means that the process started from a single particle with mass $x>0$ has the same distribution as x times the process started from a single particle with unit mass.

This verbal description has obvious similarities with that of branching random walks. More precisely, let us write $Z^{(t)}$ for the random point measure which assigns a Dirac point mass at $\log x$ for every x varying over the set of particles at time t. Taking logarithms transforms the fragmentation and homogeneity properties into the branching property for random point measures. More precisely, for every $t, t' \geqslant 0$, $Z^{(t+t')}$ is obtained from $Z^{(t)}$ by replacing each atom $z = \log x$ of $Z^{(t)}$ by a family $\{z + y, y \in \mathcal{Y}\}$, where $\mathcal{Y}$ is distributed as the family of the atoms of $Z^{(t')}$ for $Z^{(0)} = \delta_0$, and distinct atoms z of $Z^{(t)}$ correspond to independent copies of $\mathcal{Y}$.

Homogeneous fragmentations may be seen as extensions of so-called branching random walks in continuous time. The latter have been considered by Uchiyama [**27**], Biggins [**9**], Kyprianou [**18**], and others. Their main feature is that each particle has an exponentially distributed lifetime and at the instant of its death, scatters a random number of children-particles in space relative to its death point according to the point process. However the theory of branching processes in continuous time does not encompass homogeneous fragmentations, because usually each fragment starts to split instantaneously, which would correspond to particles with zero lifetime in the branching setting.

Received 18 March 2003; revised 5 October 2004.

2000 *Mathematics Subject Classification* 60J25, 60G09.

https://doi.org/10.1112/S0024610705006423 Published online by Cambridge University Press

The close connection between homogeneous fragmentations and branching random walks suggests that one should try to reduce the study of fragmentations to that of branching random walks by time-discretization. This is the path that we will follow in the first part of this work. Aside from some difficult technical problems (for instance, a most useful notion such as the first branching time has no analogue for fragmentations since, in general, dislocations occur instantaneously), this enables us to shift several deep results on branching random walks to the fragmentation setting. In particular, this yields interesting information about the asymptotic behaviour of fragmentations, which refine earlier results in [**6**].

There is another discretization method that will play an important role in this paper. The fundamental idea is due to Kingman [**15**], who pointed out that partitions of an object, say with a unit mass, can be fruitfully encoded by partitions of $\mathbb{N}$. In order to explain the coding, we introduce a sequence of independent and identically distributed random points $U_1, \ldots$ which are picked according to the mass distribution of the object. One then considers at each time $t \geqslant 0$ the random partition of the set of indices $\mathbb{N}$, such that two indices, say i and j, belong to the same block of the partition if and only if the points U_i and U_j belong to the same fragment of the object. By the law of large numbers, we see that the masses of the fragments can be recovered as the asymptotic frequencies of the blocks of the partition. Roughly, the fundamental point in Kingman's coding is that it translates the process to a discrete state-space. We refer to Pitman [**24**] for an important application of these ideas to a coalescent setting.

In the second part of this work, we shall present further applications of Kingman's idea to homogeneous fragmentations. In particular, we will show that spatial discretization is especially well suited to adapt the conceptual method of probability tilting introduced by Lyons, Pemantle and Peres (see for example [**21**]) to homogeneous fragmentations, which yields also some interesting limit theorems.

2. *Time-discretization for ranked fragmentations*

This section is devoted to the presentation of some applications of time-discretization to the asymptotic behaviour of ranked fragmentations as time tends to ∞. We shall first introduce some notation and definitions; then we shall merely translate results of Biggins on branching random walks in continuous time in the special case when the so-called dislocation measure of the fragmentation is finite. Finally, we shall extend the preceding results to the case when the dislocation measure is infinite, and also derive from the extension of a result of Rouault sharp estimates for the probability of presence of abnormally large fragments.

2.1. *Some notation and definitions*

Throughout this paper, we will work with the space of numerical sequences

$$\mathcal{S} := \left\{ \mathbf{s} = (s_1, \ldots) : s_1 \geqslant s_2 \geqslant \ldots \geqslant 0 \text{ and } \sum_1^\infty s_i \leqslant 1 \right\}$$

endowed with the uniform distance, which is a compact set. A configuration $s \in \mathcal{S}$ should be thought of as the ranked masses of the fragments resulting from the split of some object with unit total mass.

https://doi.org/10.1112/S0024610705006423 Published online by Cambridge University Press

We consider a family of Feller processes $X = (X_t, t \geqslant 0)$ with values in $\mathcal{S}$ and càdlàg paths. For every $a \in [0,1]$, we let $\mathbb{P}_a$ denote the law of X with initial distribution $(a, 0, \ldots)$ (that is, the process starts from a single fragment with mass a). We say that X is a (ranked) homogeneous fragmentation if the following two properties hold.

(i) *Homogeneity property*: For every $a \in [0,1]$, the law of aX under $\mathbb{P}_1$ is $\mathbb{P}_a$.

(ii) *Fragmentation property*: For every $\mathbf{s} = (s_1, \ldots) \in \mathcal{S}$, the process started from $X(0) = \mathbf{s}$ can be obtained as follows. Consider $X^{(1)}, \ldots$ a sequence of independent processes with respective laws $\mathbb{P}_{s_1}, \ldots$, and for every $t \geqslant 0$, let $\hat{X}(t)$ be the random sequence obtained by ranking in decreasing order the terms of the random sequences $X^{(1)}(t), \ldots$. Then $\hat{X}$ has the law of X started from $\mathbf{s}$.

It has been shown in [**3**, **4**] that homogeneous fragmentations result from the combination of two different phenomena: a continuous erosion and sudden dislocations. The erosion is a continuous deterministic mechanism; dealing with erosion is straightforward, and therefore we will only consider homogeneous fragmentations with no erosion in this work.

The dislocations occur randomly and can be viewed as the jump-component of the process. Roughly speaking, their distribution can be characterized by a measure ν on $\mathcal{S}$, called the dislocation measure. Informally ν specifies the rates at which a unit mass splits; see Subsection 2.2. It has to fulfil the conditions $\nu(\{(1,0,\ldots)\}) = 0$ and

$$\int_{\mathcal{S}} (1 - s_1)\nu(d\mathbf{s}) < \infty. \tag{1}$$

More precisely, (1) is the necessary and sufficient condition for a measure ν on $\mathcal{S}$ to be the dislocation measure of some homogeneous fragmentation (see [**3**, **4**]). We shall assume throughout this work that

$$\nu\left(\left\{\mathbf{s} \in \mathcal{S} : \sum_{i=1}^{\infty} s_i < 1\right\}\right) = 0, \tag{2}$$

which means that no mass is lost when a sudden dislocation occurs, and more precisely, entails that the total mass is a conserved quantity for the fragmentation process (that is, $\sum_{i=1}^{\infty} X_i(t) = 1$ for all $t \geqslant 0$, $\mathbb{P}_1$-almost surely). In the sequel, we shall also implicitly exclude the trivial case when $\nu \equiv 0$.

Given a real number $r > 0$, we say that a dislocation measure ν is r-geometric if ν is finite and is carried by the subspace of configurations $\mathbf{s} = (s_1, \ldots) \in \mathcal{S}$ such that $s_i \in \{r^{-n}, n \in \mathbb{N}\}$. This holds if and only if $\mathbb{P}_1(X_i(t) \in \{r^{-n}, n \in \mathbb{N}\}$ for every $i \in \mathbb{N}) = 1$ for all $t \geqslant 0$. We say that a dislocation measure is non-geometric if it is not r-geometric for any $r > 0$.

We now introduce analytic quantities defined in terms of ν which will have an important role in this work. First, we set

$$\underline{p} := \inf\left\{p \in \mathbb{R} : \int_{\mathcal{S}} \sum_{i=2}^{\infty} s_i^{p+1} \nu(d\mathbf{s}) < \infty\right\},$$

and then for every $q > \underline{p}$

$$\Phi(q) = \int_{\mathcal{S}} \left(1 - \sum_{i=1}^{\infty} s_i^{q+1}\right) \nu(d\mathbf{s}). \tag{3}$$

https://doi.org/10.1112/S0024610705006423 Published online by Cambridge University Press

The function Φ is a concave analytic increasing function; it is easy to see (cf. [**6**, Lemma 1]) that the equation

$$\Phi(q) = (q+1)\Phi'(q), \qquad q > \underline{p},$$

has a unique solution, which we denote by $\bar{p}$. More precisely, we then have

$$\Phi(q) - (q+1)\Phi'(q) < 0 \Longleftrightarrow q \in]\underline{p}, \bar{p}[\,, \tag{4}$$

and

$$\text{the map } q \longrightarrow \Phi(q)/(q+1) \text{ increases on }]\underline{p}, \bar{p}[\text{ and decreases on }]\bar{p}, \infty[. \tag{5}$$

To start with, we consider the simple sub-family of fragmentation processes with a finite dislocation measure. These can be reduced to continuous time branching random walks, and we specify some important results of Biggins [**9**] in this setting. Then we shall investigate the case when the dislocation measure is infinite by time-discretization.

2.2. *The case when the dislocation measure is finite*

Throughout this section, we assume that the dislocation measure ν is finite. It is easy to construct a fragmentation process $X = (X(t), t \geqslant 0)$ with dislocation measure ν. Let the process start, say from the state $\mathbf{1} := (1, 0, \ldots)$, and stay there for an exponential time with parameter $\nu(\mathcal{S})$. Then the process jumps independently of the waiting time to some random state in $\mathcal{S}$ distributed according to the probability measure $\nu(\cdot)/\nu(\mathcal{S})$. After this first split, each fragment has a similar evolution, independently of the other fragments. In words, a fragment with mass $x \in]0, 1[$ breaks after some exponential time with parameter $\nu(\mathcal{S})$, and produces a random sequence of smaller fragments, say xS, where S is a random variable in $\mathcal{S}$ with law $\nu(\cdot)/\nu(\mathcal{S})$.

Plainly, the empirical measure of the logarithms of the fragments

$$Z^{(t)} := \sum_{i=1}^{\infty} \delta_{\log X_i(t)}, \qquad t \geqslant 0, \tag{6}$$

can be viewed as a branching random walk in continuous time; see Uchiyama [**27**], Biggins [**9**], Kyprianou [**18**]. In this direction, let us identify two key quantities related to branching random walks in the fragmentation setting.

First, it is easy to see by an application of the Markov property at the first splitting (or branching) time, that the Laplace transform of the intensity of the point process $Z^{(t)}$ is given for $\theta > \underline{p} + 1$ by

$$m(\theta)^t := \mathbb{E}\left(\int_{\mathbb{R}} \mathrm{e}^{\theta x} Z^{(t)}(dx)\right) = \mathbb{E}\left(\sum_{i=1}^{\infty} X_i(t)^{\theta}\right) = \exp(-t\Phi(\theta - 1)). \tag{7}$$

Second, there is also the identification for the so-called additive martingale

$$W^{(t)}(\theta) := m(\theta)^{-t} \int_{\mathbb{R}} \mathrm{e}^{\theta x} Z^{(t)}(dx) = \exp(t\Phi(\theta - 1)) \sum_{i=1}^{\infty} X_i(t)^{\theta}.$$

These observations allow us to apply Biggins [**9**, Theorem 6] (see also [**8**]).

https://doi.org/10.1112/S0024610705006423 Published online by Cambridge University Press

PROPOSITION 1. *Assume that the dislocation measure ν is finite. Then, for every $p > \underline{p}$, the process*

$$M(p,t) := W^{(t)}(p+1) = \exp(t\Phi(p)) \sum_{i=1}^{\infty} X_i^{p+1}(t), \qquad t \geqslant 0,$$

is a martingale with càdlàg paths. This martingale converges uniformly on any compact subset of $]\underline{p}, \bar{p}[$, almost surely and in mean, as $t \to \infty$.

We also point out that for $p \geqslant \bar{p}$, the result of Biggins [**8**] entails that $\lim_{n\to\infty} M(p,n) = 0$ almost surely, and thus by the convergence theorem of càdlàg non-negative martingales, it holds that $\lim_{t\to\infty} M(p,t) = 0$ almost surely.

Proof of Proposition 1. Fix some compact interval $[a,b] \subset]\underline{p}, \bar{p}[$, and recall that the event that

$$\sum_{i=1}^{\infty} X_i^{a+1}(t) < \infty \qquad \text{for all } t \geqslant 0 \tag{8}$$

has probability 1. In the sequel, we shall always work on this event. We consider the random continuous function on $[a,b]$

$$M(t) : p \longrightarrow \exp(t\Phi(p)) \sum_{i=1}^{\infty} X_i^{p+1}(t),$$

which defines a martingale with values in the Banach space $\mathcal{C}([a,b],\mathbb{R})$.

Next, since $\sum_{i=1}^{\infty} s_i = 1$ for ν-almost every $\mathbf{s} \in \mathcal{S}$, observe that, for every $\theta > \underline{p}+1$, we have $\gamma(\theta-1) > \underline{p}$ for some $\gamma > 1$, and then Jensen's inequality implies that

$$\int_{\mathcal{S}} \left(\sum_{i=1}^{\infty} s_i^{\theta} \right)^{\gamma} \nu(d\mathbf{s}) \leqslant \int_{\mathcal{S}} \left(\sum_{i=1}^{\infty} s_i^{\theta\gamma+1-\gamma} \right) \nu(d\mathbf{s}) < \infty.$$

On the other hand, thanks to (5), for every $\theta \in]\underline{p}+1, \bar{p}+1[$, there exists $\alpha \in]1,\gamma[$ such that

$$m(\alpha\theta) < m(\theta)^{\alpha}.$$

By the argument used in Biggins [**9**, proof of Theorem 6] (see also Remark 2(1)), all that needs to be checked is that the martingale $(M(t), t \geqslant 0)$ has right-continuous paths. We stress that the argument we give does not rely on the assumption of finiteness for the dislocation measure.

In this direction, it is convenient to use an interval representation of the fragmentation (see [**5**]). Specifically, we consider the space of open subsets of the unit interval, endowed with the metric induced by the Hausdorff distance for the complementary closed set. One can construct a right-continuous family $(\Theta(t), t \geqslant 0)$ of random open subsets of the unit interval such that for each $t \geqslant 0$, $X(t)$ is the ranked sequence of the lengths of the interval components of $\Theta(t)$, and $\Theta(t') \subseteq \Theta(t)$ whenever $t \leqslant t'$. Recall also that assumption (2) ensures that each $\Theta(t)$ has full Lebesgue measure almost surely.

For every $x \in]0,1[$ and $q \in [a,b]$, write $f_{t,q}(x) = |I_x(t)|^q$, where $I_x(t)$ denotes the interval component of $\Theta(t)$ that contains x and $|I_x(t)|$ its length. In this setting,

https://doi.org/10.1112/S0024610705006423 Published online by Cambridge University Press

we thus have

$$\sum_{i=1}^{\infty} X_i^{q+1}(t) = \int_0^1 f_{t,q}(x)\,dx.$$

For every $t_0 \geqslant 0$ and $x \in \Theta(t_0)$, we get from the right-continuity of $(\Theta(t), t \geqslant 0)$ that $|I_x(t)|$ increases to $|I_x(t_0)|$ as t decreases to t_0. The obvious upper-bounds

$$f_{t,q}(x) \leqslant f_{t',a}(x), \qquad t \leqslant t' \text{ and } x \in]0,1[,\ q \leqslant 0$$
$$f_{t,q}(x) \leqslant f_{t_0,a}(x), \qquad t \geqslant t_0 \text{ and } x \in]0,1[,\ q \geqslant 0$$

combined with (8) enable us to apply the theorem of dominated convergence, and hence

$$\lim_{t \to t_0+} \int_0^1 f_{t,q}(x)\,dx = \int_0^1 f_{t_0,q}(x)\,dx.$$

This shows that with probability 1, the real-valued martingales $M(q,\cdot)$ have right-continuous paths for all $q \in [a,b]$.

To conclude, we observe that the random function $q \longrightarrow \sum_{i=1}^{\infty} X_i^{q+1}(t)$ is continuous and decreases as q increases. An appeal to Dini's theorem now shows that $(M(t), t \geqslant 0)$ is right-continuous at all $t \geqslant 0$, with probability 1. □

Remark 2. (1) It seems that the discretization argument in the proof of the almost sure convergence in [**9**, Theorem 6] might require a further explanation. Indeed, it is observed there that $W^{(n\delta)}$ converges almost surely when $n \to \infty$ through integers for any $\delta > 0$, and then it is claimed that this implies the almost sure convergence of $W^{(t)}$ when $t \to \infty$ through the rationals. The latter assertion does not look obvious, so we propose a slightly different argument. One works with the martingale $W^{(t)}$ with values in the space of continuous functions on some compact space, endowed with the supremum norm, $\|\cdot\|$. We know that this martingale converges in mean as $t \to \infty$. The norm is a convex map; therefore for every integer n, the process $\|W^{(t+n)} - W^{(n)}\|$ is a non-negative submartingale with regular paths. Doob's maximal inequality now entails that for every $\varepsilon > 0$,

$$\mathbb{P}\big(\exists\, t > 0 : \big\|W^{(t+n)} - W^{(n)}\big\| > \varepsilon\big) \leqslant \varepsilon^{-1} \sup_{t \geqslant 0} \mathbb{E}\big(\big\|W^{(t+n)} - W^{(n)}\big\|\big),$$

and the right-hand side converges to 0 as $n \to \infty$. An application of the Borel–Cantelli lemma now completes the proof of the almost sure convergence of $W^{(t)}$ as $t \to \infty$.

(2) We also mention that Proposition 1 can be extended to complex numbers p with $\underline{p} < \Re p < \bar{p}$. The key point is to check that the martingale $p \longrightarrow M(p,t)$ $(t \geqslant 0)$, viewed as a process with values in the space $\mathcal{C}(K, \mathbb{C})$ for some compact set $K \subset \{z : \underline{p} < \Re p < \bar{p}\}$, has right-continuous paths. We know from the proof of Proposition 1 that the latter holds when K is a real segment, and the general case follows easily.

Next, we derive an important consequence of the preceding analysis concerning almost sure large deviations for the empirical measure; see Biggins [**9**, Corollary 4 and the discussion on p. 150].

Corollary 3. *Assume that the dislocation measure ν is finite and non-geometric. For $p \in]\underline{p}, \bar{p}[$, let $M(p,\infty)$ be the terminal value of the uniformly*

https://doi.org/10.1112/S0024610705006423 Published online by Cambridge University Press

integrable martingale $M(p,\cdot)$. *If* $f:\mathbb{R}\longrightarrow\mathbb{R}$ *is a function with compact support which is directly Riemann integrable, then*

$$\lim_{t\to\infty}\sqrt{t}\,\mathrm{e}^{-t((p+1)\Phi'(p)-\Phi(p))}\int_{\mathbb{R}}f(t\Phi'(p)+y)Z^{(t)}(dy)$$
$$=\frac{M(p,\infty)}{\sqrt{2\pi|\Phi''(p)|}}\int_{-\infty}^{\infty}f(y)\mathrm{e}^{-(p+1)y}\,dy$$

uniformly for p *in compact subsets of* $]\underline{p},\bar{p}[$, *almost surely.*

We point out that this result applied for indicator functions of bounded intervals gives a sharp large deviation statement that extends [**6**, Corollary 2].

2.3. *The case when the dislocation measure is infinite*

We now drop the assumption of finiteness of the dislocation measure ν, and merely assume that (1) holds. As above we associate to the fragmentation X the empirical measures $Z^{(t)}$ defined in (6); note that when $\nu(\mathcal{S})=\infty$, the process $(Z^{(t)},t\geqslant 0)$ is no longer of the kind considered by Uchiyama [**27**].

THEOREM 4. *Proposition* 1 *and Corollary* 3 *hold when one relaxes the requirement of finiteness of the dislocation measure* ν *and merely assumes* (1).

Proof. The restriction of the process of empirical distribution to integers times $(Z^{(n)},n=0,1,\ldots)$ is a branching random walk; we aim at applying results of Biggins [**9**] in this setting.

The main difficulty is that the distribution of $Z^{(1)}$ is not explicitly known in terms of the dislocation measure ν. However, it is known that (7) still holds; see for example the identity [**6**, (6)]. Moreover, [**6**, proof of Theorem 2] shows that for every $p>\underline{p}$, there exists some $\gamma>1$ such that for every $t\geqslant 0$,

$$\mathbb{E}\left(\left(\sum_{i=1}^{\infty}X_i^{p+1}(t)\right)^{\gamma}\right)<\infty. \tag{9}$$

Biggins [**9**, Theorem 2] now shows that the conclusions of Proposition 1 hold provided that $t\to\infty$ through integers. We can complete the argument as in Biggins [**9**, Theorem 6] (using also the proof of Proposition 1 and Remark 2 in the present paper). The extension of Corollary 3 is proved by adapting the arguments of Biggins [**9**, p. 150]. □

In a different direction, one can also use the skeleton method to estimate the probability of presence of abnormally large fragments as time goes to infinity. Indeed, a similar problem has been solved for branching random walks, see [**25**] and the references therein. Informally in the so-called sub-critical region, the probability of presence of particles is of the same order as the mean number of particles in that region, and the asymptotic behaviour of the latter can be derived from the local central limit theorem. This incites us to fix two real numbers $\alpha<\beta$ and to introduce for every $t\geqslant 0$ and $x\in\mathbb{R}$ the notation

$$U(t,x):=\mathbb{P}(Z^{(t)}([x+\alpha,x+\beta])>0) \tag{10}$$
$$V(t,x):=\mathbb{E}(Z^{(t)}([x+\alpha,x+\beta])). \tag{11}$$

https://doi.org/10.1112/S0024610705006423 Published online by Cambridge University Press

THEOREM 5. *Assume that the dislocation measure ν is non-geometric.*

(i) *If $p > \underline{p}$, we have*

$$\lim_{t\to\infty} \sqrt{t}\, \mathrm{e}^{-t((p+1)\Phi'(p)-\Phi(p))} V(t, -t\Phi'(p)) = \frac{1}{\sqrt{2\pi|\Phi''(p)|}} (p+1)^{-1} \big(\mathrm{e}^{-(p+1)\alpha} - \mathrm{e}^{-(p+1)\beta}\big).$$

(ii) *If $p > \bar{p}$, there exists a positive finite constant K_p such that*

$$\lim_{t\to\infty} \frac{U(t, -t\Phi'(p))}{V(t, -t\Phi'(p))} = K_p.$$

We point out that in the range $p \in]\underline{p}, \bar{p}[$, (i) is the counterpart in mean of the result of Corollary 3, when $f = \mathbf{1}_{[\alpha,\beta]}$. It implies that the convergence there holds also in $L^1(\mathbb{P})$, thanks to Scheffé's theorem.

Proof of Theorem 5. The proof relies on a result of Rouault [**25**] for branching random walks, which has been extended under more general conditions in [**7**], and that we take here for granted. Recall also (9), which ensures that the assumptions of [**7**] are satisfied.

We apply the skeleton method. Let $h > 0$ be a time mesh; the fragmentation process observed at times nh ($n \in \mathbb{N}$) yields a branching random walk. Write

$$\widehat{Z^{(h)}}(\theta) := \int_{\mathbb{R}} \mathrm{e}^{\theta x} Z^{(h)}(dx) = \sum_i X_i(h)^\theta$$

and

$$\Lambda_h(\theta) := \log \mathbb{E}\big(\widehat{Z^{(h)}}(\theta)\big) = -h\Phi(\theta - 1).$$

In the case $h = 1$, we write for simplicity $\Lambda = \Lambda_1$ and set

$$a = -\Phi'(p), \quad \sigma_p^2 = -\Phi''(p), \quad \Lambda^*(a) = \Phi(p) - (p+1)\Phi'(p).$$

For an arbitrary mesh $h > 0$, we define by scaling

$$\Lambda_h^*(x) = h\Lambda^*(x/h).$$

It is immediately checked that if θ solves $a = \Lambda_h'(\theta)$, then $\Lambda_h^*(a) = \theta\Lambda_h'(\theta) - \Lambda_h(\theta)$. Applying [**7**, Theorem 2] we get first

$$\lim_n \sigma_p \sqrt{2\pi nh}\ \mathrm{e}^{nh\Lambda^*(a)} V(nh, anh) = (p+1)^{-1}\big(\mathrm{e}^{-(p+1)\alpha} - \mathrm{e}^{-(p+1)\beta}\big), \tag{12}$$

and then, if $\Lambda_h^*(a) > 0$, the limit

$$\lim_n \frac{U(nh, anh)}{V(nh, anh)} =: K_p^{(h)}$$

exists for each $h > 0$ and is positive. Now, it is easy to see that the functions

$$t \longmapsto \sigma_p \sqrt{2\pi t}\ \mathrm{e}^{t\Lambda^*(a)} V(t, at) \quad \text{and} \quad t \longmapsto \frac{U(t, at)}{V(t, at)} \tag{13}$$

are continuous. We apply the Croft–Kingman lemma ([**1**, A 9.1, p. 438], see also [**14**]). Both limits exist when $t \to \infty$. In the first case, it is of course the right-hand side of (12). In the second case, it is any $K_p^{(h)}$ since they are all equal. □

https://doi.org/10.1112/S0024610705006423 Published online by Cambridge University Press

3. *Spatial discretization and fragmentation of partitions*

This section is devoted to another useful discretization technique which has been sketched in the introduction and that we now recall for convenience. We may suppose that we are given a sequence of independent and identically distributed. random points $U_1, \ldots$ which are picked according to the mass distribution of the object. These random points are assumed to be independent of the fragmentation process. One then looks at the fragmentation process as the evolution as time t passes of the random partition $\Pi(t)$ of $\mathbb{N}$ which is given as follows. Two indices i and j are in the same block of $\Pi(t)$ if and only if the points U_i and U_j belong to the same fragment of the object at time t. Plainly, the random partition gets finer and finer as time passes.

We first present in Subsection 3.1 the necessary background on partitions of $\mathbb{N}$, and then in Subsection 3.2 the Poissonian construction of homogeneous fragmentations and the connection with subordinators, following closely [**4**]. Subsection 3.3 is devoted to the study of probability tilting based on additive martingales, adapting to the random partition setting the so-called conceptual method of Lyons and Pemantle and Peres [**21**, **22**]. Finally, as an example of application, we investigate in Subsection 3.4 the convergence of the so-called derivative martingale.

3.1. *Preliminaries*

A partition of $\mathbb{N} = \{1, \ldots\}$ is a sequence $\pi = (\pi_1, \pi_2, \ldots)$ of disjoint subsets, called blocks, such that $\bigcup \pi_i = \mathbb{N}$. The blocks π_i of a partition are enumerated in the increasing order of their least element, that is $\min \pi_i \leqslant \min \pi_j$ for $i \leqslant j$, with the convention that $\min \varnothing = \infty$. If π and π' are two partitions of $\mathbb{N}$, we say that π is finer than π' if every block of π is contained in some block of π'.

For every block $B \subseteq \mathbb{N}$, we denote by $\pi_{|B}$ the partition of B induced by π and an obvious restriction. For every integer k, the block $\{1, \ldots, k\}$ is denoted by $[k]$. A partition π is entirely determined by the sequence of its restrictions $(\pi_{|[k]}, k \in \mathbb{N})$, and conversely, if for every $k \in \mathbb{N}$, γ_k is a partition of $[k]$ such that the restriction of γ_{k+1} to $[k]$ coincides with γ_k (this will be referred to as the compatibility property in the sequel), then there exists a unique partition $\pi \in \mathcal{P}$ such that $\pi_{|[k]} = \gamma_k$ for every $k \in \mathbb{N}$.

The space of partitions of $\mathbb{N}$ is denoted by $\mathcal{P}$ and endowed with the hyper-distance

$$\operatorname{dist}(\pi, \pi') := 1/\max\left\{k \in \mathbb{N} : \pi_{|[k]} = \pi'_{|[k]}\right\},$$

with the convention $1/\max \mathbb{N} := 0$. This makes $\mathcal{P}$ compact.

One says that a block $B \subseteq \mathbb{N}$ has an asymptotic frequency if the limit

$$|B| := \lim_{n \to \infty} n^{-1} \operatorname{Card}(B \cap [n])$$

exists. When every block of some partition $\pi \in \mathcal{P}$ has an asymptotic frequency, we write $|\pi| = (|\pi_1|, \ldots)$, and then $|\pi|^{\downarrow} = (|\pi|_1^{\downarrow}, \ldots) \in \mathcal{S}$ for the decreasing rearrangement of the sequence $|\pi|$. (Ranking the asymptotic frequencies of the blocks of π in decreasing order is just a simple procedure to forget the labels of these blocks. In other words, we want to consider the family of the asymptotic frequencies without keeping the additional information provided by the way blocks are labelled.) In the case when some block of the partition π does not have an asymptotic frequency, we

https://doi.org/10.1112/S0024610705006423 Published online by Cambridge University Press

decide that $|\pi| = |\pi|^{\downarrow} = \partial$, where ∂ stands for some extra point added to $\mathcal{S}$. This defines a natural map $\pi \longrightarrow |\pi|^{\downarrow}$ from $\mathcal{P}$ to $\mathcal{S} \cup \{\partial\}$ which is not continuous.

We call nested partitions a collection $\Pi = (\Pi(t), t \geqslant 0)$ of partitions of $\mathbb{N}$ such that $\Pi(t)$ is finer than $\Pi(t')$ when $t' \leqslant t$. There is a simple procedure for the construction of a large family of nested partitions which we now describe and will use throughout the rest of this section.

We call discrete point measure on $\mathbb{R}_+ \times \mathcal{P} \times \mathbb{N}$ any measure ω which can be expressed in the form

$$\omega = \sum_{(t,\pi,k)\in\mathcal{D}}^{\infty} \delta_{(t,\pi,k)},$$

where $\mathcal{D}$ is a subset of $\mathbb{R}_+ \times \mathcal{P} \times \mathbb{N}$ such that the following two requirements hold.

(i) For every $t \in \mathbb{R}$, $\omega(\{t\} \times \mathcal{P} \times \mathbb{N}) = 0$ or 1.

(ii) For every real number $t' \geqslant 0$ and integer $n \geqslant 1$,

$$\mathrm{Card}\{(t,\pi,k) \in \mathcal{D} : t \leqslant t', \pi_{|[n]} \neq \mathrm{trivial}(n), k \leqslant n\} < \infty,$$

where $\mathrm{trivial}(n) = ([n], \varnothing, \varnothing, \ldots)$ stands for the partition of $[n]$ which has a single non-empty block. (Roughly, $\mathrm{trivial}(n)$ plays the role of a neutral element in the space of partitions of $[n]$.)

Starting from an arbitrary discrete point measure ω on $\mathbb{R}_+ \times \mathcal{P} \times \mathbb{N}$, we may construct nested partitions $\Pi = (\Pi(t), t \geqslant 0)$ as follows. Fix $n \in \mathbb{N}$; the assumption that the point measure ω is discrete enables us to construct a step-path $(\Pi(t,n), t \geqslant 0)$ with values in the space of partitions of $[n]$, which only jumps at times t at which the fibre $\{t\} \times \mathcal{P} \times \mathbb{N}$ carries an atom of ω, say (t,π,k), such that $\pi_{|[n]} \neq \mathrm{trivial}(n)$ and $k \leqslant n$. In that case, $\Pi(t,n)$ is the partition obtained by replacing the kth block of $\Pi(t-,n)$, namely $\Pi_k(t-,n)$, by the restriction $\pi_{|\Pi_k(t-,n)}$ of π to this block, and leaving the other blocks unchanged. Now it is immediate from this construction that for each time $t \geqslant 0$, the sequence $(\Pi(t,n), n \in \mathbb{N})$ is compatible, and hence there exists a unique partition $\Pi(t)$ such that $\Pi(t)_{|[n]} = \Pi(t,n)$ for each $n \in \mathbb{N}$.

3.2. *Poisson measures, homogeneous fragmentations, and subordinators*

We denote the space of discrete point measures on $\mathbb{R}_+ \times \mathcal{P} \times \mathbb{N}$ by Ω, and the sigma-field generated by the restriction to $[0,t] \times \mathcal{P} \times \mathbb{N}$ by $\mathcal{G}(t)$. Thus $(\mathcal{G}(t))_{t\geqslant 0}$ is a filtration, and the nested partitions $(\Pi(t),\ t \geqslant 0)$ are $(\mathcal{G}(t))_{t\geqslant 0}$-adapted. We shall also need to consider the sigma-field $\mathcal{F}(t)$ generated by the decreasing rearrangement $|\Pi(r)|^{\downarrow}$ of the sequence of the asymptotic frequencies of the blocks of $\Pi(r)$ for $r \leqslant t$, and $(\mathcal{F}(t))_{t\geqslant 0}$ is a sub-filtration of $(\mathcal{G}(t))_{t\geqslant 0}$.

Now consider a dislocation measure ν, that is, a measure on $\mathcal{S}$ which fulfils the requirements of Subsection 2.1. According to [**4**, Theorem 2] there exists a unique measure μ on $\mathcal{P}$ which is exchangeable (that is, invariant by the action of finite permutations on $\mathcal{P}$), and such that ν is the image of μ by the map $\pi \longrightarrow |\pi|^{\downarrow}$. An important fact which stems from exchangeability, is that the distribution of the asymptotic frequency of the first block $|\pi_1|$ under the measure μ is that of a size-biased picked term from the ranked sequence s under ν. In other words, there is the identity

$$\int_{\mathcal{P}} f(|\pi_1|)\mu(d\pi) = \int_{\mathcal{S}} \sum_{i=1}^{\infty} s_i f(s_i)\nu(d\mathbf{s}), \tag{14}$$

where $f : [0,1] \longrightarrow \mathbb{R}_+$ denotes a generic measurable function with $f(0) = 0$.

https://doi.org/10.1112/S0024610705006423 Published online by Cambridge University Press

Let $\mathbb{P}$ be the probability measure on Ω corresponding to the law of a Poisson point measure with intensity $dt \otimes \mu \otimes \#$, where $\#$ denotes the counting measure on $\mathbb{N}$. Assumption (1) on the dislocation measure ν ensures that ω is a discrete point measure $\mathbb{P}$-almost surely. The nested partitions $(\Pi(t),\ t \geqslant 0)$ constructed above from ω now form a Markov process; see [**4**, Section 3]. More precisely the Markov property is essentially a variation of the branching property; it can be stated as follows. For every $t, t' \geqslant 0$, the conditional distribution of $\Pi(t + t')$ given $\mathcal{G}(t)$ is the same as that of the random partition of $\mathbb{N}$ induced by the restrictions $\Pi^{(1)}(t')_{|B_1}, \Pi^{(2)}(t')_{|B_2}, \ldots$, where $\Pi^{(1)}, \ldots$ are independent copies of Π and $(B_1, \ldots) = \Pi(t)$ is the sequence of blocks of $\Pi(t)$. In the terminology of [**4**], we say that $\Pi = (\Pi(t),\ t \geqslant 0)$ is a (partition-valued) homogeneous fragmentation under $\mathbb{P}$.

Another crucial fact is that the partitions $(\Pi(t), t \geqslant 0)$ are exchangeable under $\mathbb{P}$, that is, their distribution is invariant under the action of finite permutations on $\mathbb{N}$; see [**4**, Section 3]. It follows from a celebrated theorem of Kingman [**15**] that $\mathbb{P}$-almost surely, $\Pi(t)$ has asymptotic frequencies for all $t \geqslant 0$; cf. [**4**, Theorem 3(i)]. The process of ranked asymptotic frequencies $|\Pi|^{\downarrow} := X$ is a Markov process with values in $\mathcal{S}$; it provides a version of the ranked fragmentation which we considered in Section 2; cf. [**3**].

The tagged fragment is the fragment of the object that contains the first tagged point U_1, that is, which corresponds to the first block $\Pi_1(\cdot)$. The process $|\Pi_1(\cdot)|$ of the asymptotic frequencies of the first block and its logarithm,

$$\xi(t) := -\log|\Pi_1(t)|, \qquad t \geqslant 0,$$

will have a special role in this study. A crucial point is that under $\mathbb{P}$, $\xi = (\xi(t),\ t \geqslant 0)$ is a subordinator with Laplace exponent Φ, which is given by (3); cf. [**6**, Theorem 3(ii)]. This means that $(\xi(t),\ t \geqslant 0)$ is a càdlàg process with independent and stationary increments, and the Laplace transform of its one-dimensional distribution is given by the identity

$$\mathbb{E}\left(\exp(-q\xi(t)\right) = \exp(-t\Phi(q)), \qquad q > \underline{p}.$$

More precisely, let us denote by $\mathcal{G}_1(t)$ the sigma-field generated by the restriction of the discrete point measure ω to the fibre $[0,t] \times \mathcal{P} \times \{1\}$. Thus $(\mathcal{G}_1(t))_{t \geqslant 0}$ is a sub-filtration of $(\mathcal{G}(t))_{t \geqslant 0}$, and the first block of $\Pi(t)$, $\Pi_1(t)$, and *a fortiori* its asymptotic frequency $\mathrm{e}^{-\xi(t)}$, are $\mathcal{G}_1(t)$-measurable. Let $\mathcal{D}_1 \subseteq [0, \infty[$ be the random set of times $r \geqslant 0$ for which the discrete point measure has an atom on the fibre $\{r\} \times \mathcal{P} \times \{1\}$, and for every $r \in \mathcal{D}_1$, denote the second component of this atom by $\pi(r)$. The construction of the nested partitions from the discrete point measure yields the identity

$$\exp(-\xi(t)) = |\Pi_1(t)| = \prod_{r \in \mathcal{D}_1 \cap [0,t]} |\pi_1(r)|, \tag{15}$$

for all $t \geqslant 0$, almost surely under $\mathbb{P}$; see for example [**4**, first remark at the end of Section 5]. Observe that taking logarithms turns identity (15) into the Lévy–Itô decomposition for subordinators.

Finally, the conditional distribution of the size of the tagged fragment, $|\Pi_1(t)| = \mathrm{e}^{-\xi(t)}$, given $\mathcal{F}(t)$ (the sigma-field generated by the ranked asymptotic frequencies) is that of a size-biased sample from the ranked sequence $|\Pi(t)|^{\downarrow}$. In other words,

https://doi.org/10.1112/S0024610705006423 Published online by Cambridge University Press

we have

$$\mathbb{E}(f(\exp(-\xi(t))) = \mathbb{E}\left(\sum_{i=1}^{\infty} |\Pi_i(t)| f(|\Pi_i(t)|)\right) = \mathbb{E}\left(\sum_{j=1}^{\infty} X_j(t) f(X_j(t))\right)$$

where $f : [0,1] \longrightarrow \mathbb{R}_+$ denotes a generic measurable function with $f(0) = 0$. More generally, exchangeability ensures that for every $t \geqslant 0$, the sequence $|\Pi(t)|$ of the asymptotic frequencies is a size-biased reordering of the ranked sequence $X(t) = |\Pi(t)|^{\downarrow}$.

3.3. *Additive martingales and tilted probability measures*

There are two simple martingales connected to fragmentations for every parameter $p > \underline{p}$. First, a well known fact for subordinators is that

$$\mathcal{E}(p,t) := \exp(-p\xi(t) + t\Phi(p)) = \mathrm{e}^{t\Phi(p)} |\Pi_1(t)|^p$$

is a positive $(\mathbb{P}, \mathcal{G}(t))$-martingale. Second, when we project $\mathcal{E}(p,t)$ on the sub-filtration $\mathcal{F}(t)$, we recover the additive martingale

$$M(p,t) = \exp(t\Phi(p)) \sum_{i=1}^{\infty} |\Pi_i(t)|^{p+1} = \exp(t\Phi(p)) \sum_{j=1}^{\infty} X_j^{p+1}.$$

We point out that, more precisely, $M(p,\cdot)$ is a $(\mathbb{P}, \mathcal{G}(t))$-martingale which is adapted to the sub-filtration $\mathcal{F}(t)$.

Following the genuine method of Lyons, Pemantle and Peres (see for example [**21**]), we introduce the tilted probability measure $\mathbb{P}^{(p)}$ on the space of discrete point measures Ω endowed with the filtration $(\mathcal{G}(t))_{t \geqslant 0}$ by

$$d\mathbb{P}^{(p)}_{|\mathcal{G}(t)} = \mathcal{E}(p,t)\, d\mathbb{P}_{|\mathcal{G}(t)}. \tag{16}$$

Observe that projections on the sub-filtration $\mathcal{F}(t)$ give the identity

$$d\mathbb{P}^{(p)}_{|\mathcal{F}(t)} = M(p,t)\, d\mathbb{P}_{|\mathcal{F}(t)}. \tag{17}$$

The effect of the change of probability is easy to describe, both at the level of the tagged fragment and that of the discrete point measure.

PROPOSITION 6. (i) *Under* $\mathbb{P}^{(p)}$, *the process* $\xi(t) = -\log|\Pi_1(t)|$ *is a subordinator with Laplace exponent*

$$\Phi^{(p)}(q) := \Phi(p+q) - \Phi(p), \qquad q > \underline{p} - p.$$

(ii) *Under* $\mathbb{P}^{(p)}$, *the discrete point measure* ω *is Poissonian. More precisely, the following hold*

(a) *The restriction of* ω *to* $\mathbb{R}_+ \times \mathcal{P} \times \{2,3,\ldots\}$ *has the same distribution as under* $\mathbb{P}$ *and is independent of the restriction to the fibre* $\mathbb{R}_+ \times \mathcal{P} \times \{1\}$.

(b) *In the notation of Subsection* 3.1, *the family* $\{(r, \pi(r)), r \in \mathcal{D}_1\}$ *is that of the atoms of a Poisson random measure on* $\mathbb{R}_+ \times \mathcal{P}$ *with intensity* $dr \otimes \mu^{(p)}$, *where*

$$\mu^{(p)}(d\pi) = |\pi_1|^p \mu(d\pi).$$

Proof. The first assertion stems from the classical Esscher transform; see for instance, Sato [**26**, Example 33.15]. The description of the law of the discrete point

https://doi.org/10.1112/S0024610705006423 Published online by Cambridge University Press

measure under $\mathbb{P}^{(p)}$ is easily seen from the formula (15), and classical properties of the exponential tilting for Poisson random measures. □

We stress that the tilting only affects the distribution of the discrete point measure on the fibre $\mathbb{R}_+ \times \mathcal{P} \times \{1\}$. In this direction, it is interesting to use Proposition 6(ii) and compare the evolution of the ranked-fragmentation $X = |\Pi(\cdot)|^{\downarrow}$ under $\mathbb{P}^{(p)}$ with the evolution under $\mathbb{P}$. For the sake of simplicity, we again suppose here that the dislocation measure ν is finite, so the evolution of the ranked fragmentation under $\mathbb{P}$ is described in Subsection 2.2.

First, we observe that by absolute continuity, the random partition $\Pi(t)$ obtained by evaluating the nested partitions at time t, possesses asymptotic frequencies almost surely under the tilted probability $\mathbb{P}^{(p)}$. The first block $\Pi_1(\cdot)$ has a special role in the definition of the tilted probability $\mathbb{P}^{(p)}$, and cannot be recovered from the ranked sequence $|\Pi(\cdot)|^{\downarrow}$ alone. Let us call marked the unique particle (or asymptotic frequency) at time t corresponding to $\Pi_1(t)$ and unmarked the other ones. Observe that under $\mathbb{P}$, the tagged fragment coincides with the marked particle, and that under $\mathbb{P}^{(p)}$, the unmarked particles follow the same evolution as under $\mathbb{P}$, that is, they split according to ν, independently of the others, and only produce unmarked particles. Under $\mathbb{P}^{(p)}$, the marked particle splits independently of the other particles, but with a different rate, namely

$$\nu^{(p)}(ds) := \left(\sum_{i=1}^{\infty} s_i^{p+1}\right)\nu(d\mathbf{s}).$$

Indeed, we deduce from Proposition 6(ii) that $\nu^{(p)}$ is the image of the intensity measure $\mu^{(p)}$ by the map $\pi \longrightarrow |\pi|^{\downarrow}$, and since under μ, $|\pi_1|$ can be viewed as a size-biased pick from the ranked sequence $|\pi|^{\downarrow}$ (recall identity (14)), this yields the formula above. The 'new' marked particle is picked at random amongst the particles produced by the splitting of the 'old' marked particle as follows. Let x denote the mass of the old marked particle and xs the ranked sequence of the masses of the particles produced after the splitting, where $s = (s_1, \ldots) \in \mathcal{S}$. Then the probability that the new marked particle has mass xs_j equals $s_j^{p+1}/\sum_{i=1}^{\infty} s_i^{p+1}$. In short, the marked particle $\Pi_1(\cdot)$ can be viewed as a canonic analogue in the fragmentation setting of the so-called spine in the branching random walk framework.

It may be interesting to point out the following connection with the so-called thinning of discrete point measures. Recall that, given some metric space A, a discrete point measure μ on A with atoms $x_1, \ldots$ (that is, $\mu = \sum_{i=1}^{\infty} \delta_{x_i}$), and a measurable function $f : A \longrightarrow [0,1]$, an f-thinning of μ is the random discrete point measure $\mu^{(r)}$ obtained by keeping each atom x of μ with probability $f(x)$, independently of the others. In other words,

$$\mu^{(f)} = \sum_{j=1}^{\infty} \mathbf{1}_{\{\chi_j = 1\}} \delta_{x_j},$$

where $\chi_1, \ldots$ is a sequence of independent Bernoulli variables with $\mathbb{P}(\chi_j = 1) = f(x_j)$. Informally, the following corollary shows that dislocations are less (respectively more) frequent under $\mathbb{P}^{(p)}$ than under $\mathbb{P}$ for $p > 0$ (respectively for $p < 0$).

https://doi.org/10.1112/S0024610705006423 Published online by Cambridge University Press

COROLLARY 7. *For every $q > 0$, let f_q be the map on $\mathbb{R}_+ \times \mathcal{P} \times \mathbb{N}$ such that $f_q(t, \pi, k) = |\pi_1|^q$ if the partition π possesses asymptotic frequencies and $k = 1$, and $f_q(t, \pi, k) = 0$ otherwise.*

(i) *For every $p > 0$, the image of $\mathbb{P}$ by an f_p-thinning of the discrete point measure ω is $\mathbb{P}^{(p)}$.*

(ii) *For every $p \in]\underline{p}, 0[$, the image of $\mathbb{P}^{(p)}$ by an f_{-p}-thinning of the discrete point measure ω is $\mathbb{P}$.*

Proof. The first statement is immediate from Proposition 6(ii) and properties of thinning (see for example Kingman [**16**, Chapter 5]). Suppose now that $\underline{p} < p < 0$, and work under $\mathbb{P}$. Let ω' be a random Poisson point measure on the fibre $\mathbb{R}_+ \times \mathcal{P} \times \{1\}$ with intensity $dt \otimes (|\pi_1|^p - 1)\mu(d\pi)$, which is independent of ω. By superposition of independent Poisson measures, $\mathbb{P}^{(p)}$ can be identified as the law of $\omega + \omega'$ under $\mathbb{P}$. It follows readily that the original probability measure $\mathbb{P}$ can be recovered from $\mathbb{P}^{(p)}$ by f_{-p}-thinning. □

3.4. *The derivative martingale*

We end this work by considering the so-called derivative martingale that we now introduce. Recall that $\bar{p} > 0$ is the critical value for the convergence in $L^1(\mathbb{P})$ of the additive martingales. The process

$$\mathcal{E}'(t) := (t\Phi'(\bar{p}) - \xi(t)) \exp(-\bar{p}\xi(t) + t\Phi(\bar{p})), \qquad t \geqslant 0,$$

is clearly a $(\mathbb{P}, \mathcal{G}(t))$-martingale; its projection on the sub-filtration $(\mathcal{F}(t))_{t \geqslant 0}$ is a $(\mathbb{P}, \mathcal{F}(t))$ martingale called the derivative martingale and given by

$$M'(t) = \sum_{i=1}^{\infty} (t\Phi'(\bar{p}) + \log(|\Pi_i(t)|)) \exp(t\Phi(\bar{p}))|\Pi_i(t)|^{\bar{p}+1}.$$

We stress that the derivative martingale is not always positive, which contrasts with the case of additive martingales. The idea of considering the derivative martingale at the critical value goes back to Neveu [**23**] for the branching Brownian motion. For the branching random walk, it has been considered by Kyprianou [**17**], Liu [**20**] with the help of a functional equation and by Biggins and Kyprianou [**10**] with the measure change method.

PROPOSITION 8. (i) *The martingale M' converges $\mathbb{P}$-almost surely to a finite non-positive limit $M'(\infty)$.*

(ii) $\mathbb{E}(M'(\infty)) = -\infty$.

(iii) $\mathbb{P}(M'(\infty) < 0) = 1$.

The proposition could be derived by time-discretization from its analogue for branching random walks. However, as checking the technical details may be rather involved in this instance, we shall present a direct proof based on probability tilting. This technique is due to Lyons, Pemantle and Peres [**22**]; it can be also applied to establish the uniform integrability of additive martingales (cf. Theorem 4). We also refer to Harris [**13**] and Kyprianou [**19**] for related treatments.

https://doi.org/10.1112/S0024610705006423 Published online by Cambridge University Press

Proof of Proposition 8. (i) Define for every $i \in \mathbb{N}$ and $s \leqslant t$, $\beta_{s,t}(i)$ as the unique block of $\Pi(s)$ containing $\Pi_i(t)$. For $a > 0$, let

$$\begin{cases} \Pi_i^{(a)}(t) = \Pi_i(t) & |\beta_{s,t}(i)| \leqslant \exp\{a - s\Phi'(\bar{p})\} \text{ for every } s \leqslant t \\ \Pi_i^{(a)}(t) = \varnothing & \text{otherwise.} \end{cases}$$

The family $\{\Pi_i^{(a)}(t) : i \in \mathbb{N}\}$ obviously possesses asymptotic frequencies. Moreover, it should be plain that as t varies in $[0, \infty[$, this family of partitions is nested. We denote by $(\mathcal{H}(t))_{t\geqslant 0}$ the filtration generated by the process of their ranked asymptotic frequencies, so $(\mathcal{H}(t))_{t\geqslant 0}$ is another sub-filtration of $(\mathcal{G}(t))_{t\geqslant 0}$.

Because $\Phi'(\bar{p}) = \Phi(\bar{p})/(\bar{p}+1)$ and the martingale $M(\bar{p}, t)$ converges to 0, $\mathbb{P}$-almost surely, we have $\sup_{t\geqslant 0}\{\exp(t\Phi'(\bar{p}))|\Pi(t)|_1^{\downarrow}\} < \infty$, $\mathbb{P}$-almost surely. It follows that

$$\lim_{a\to\infty} \mathbb{P}\big(\Pi_i^{(a)}(t) = \Pi_i(t) \text{ for all } i \in \mathbb{N} \text{ and for all } t \geqslant 0\big) = 1.$$

Thus, in order to prove the existence of a finite limit for M', it suffices to establish that if

$$M_a(t) := \sum_{i=1}^{\infty} (\log(1/|\Pi_i(t)|) - t\Phi'(\bar{p}) + a) \exp(t\Phi(\bar{p}))\big|\Pi_i^{(a)}(t)\big|^{\bar{p}+1}, \tag{18}$$

then $\lim_{t\to\infty} M_a(t) =: M_a(\infty)$ exists $\mathbb{P}$-almost surely for every $a > 0$.

From now on, we fix $a > 0$. Since the process $\xi(t) - \Phi'(\bar{p})t$ has no negative jumps,

$$\mathcal{M}_a(t) := (\xi(t) + a - t\Phi'(\bar{p})) \exp(-\bar{p}\xi(t) + t\Phi(\bar{p}))\mathbf{1}_{\{t<\zeta_a\}},$$

where $\zeta_a = \inf\{t \geqslant 0 : \xi(t) < t\Phi'(\bar{p}) - a\}$, can be viewed as a stopped (non-negative) $\mathbb{P}$-martingale. Its projection on the sub-filtration $(\mathcal{H}(t))_{t\geqslant 0}$ is $M_a(t)$, which therefore is a non-negative $(\mathbb{P}, \mathcal{H}(t))$ martingale, and thus possesses a finite limit as $t \to \infty$, $\mathbb{P}$-almost surely.

(ii) It is sufficient to show that for all $a > 0$, in the notation above,

$$\liminf_{t\to\infty} M_a(t) < \infty, \qquad \mathbb{Q}\text{-almost surely.} \tag{19}$$

Indeed (19) entails that the $\mathbb{P}$-martingale M_a is uniformly integrable (see Lyons [**21**]). Then $\mathbb{E}(M_a(t)) = a$, which yields $\mathbb{E}(-M'(\infty)) \geqslant a$ for every $a > 0$. In this direction, we introduce the tilted probability measure $\mathbb{Q}$ on Ω given by

$$d\mathbb{Q}_{|\mathcal{G}(t)} = a^{-1}\mathcal{M}_a(t)\, d\mathbb{P}_{|\mathcal{G}(t)}, \tag{20}$$

so we also have

$$d\mathbb{Q}_{|\mathcal{H}(t)} = a^{-1}M_a(t)\, d\mathbb{P}_{|\mathcal{H}(t)}.$$

The following lemma, which is a simple variation of Proposition 6, lies at the heart of the proof of (19).

LEMMA 9. (i) *Under* $\mathbb{Q}$, *the process*

$$\lambda(t) := (\xi(t) + a - t\Phi'(\bar{p}))\mathbf{1}_{\{t<\zeta_1\}}, \qquad t \geqslant 0,$$

is a centred Lévy process with no negative jumps started from a *and conditioned to stay positive for ever* (*see for instance* [**11**]).

(ii) *Under* $\mathbb{Q}$, *the restriction of* ω *to* $\mathbb{R}_+ \times \mathcal{P} \times \{2, 3, \ldots\}$ *has the same distribution as under* $\mathbb{P}$ *and is independent of the restriction to the fibre* $\mathbb{R}_+ \times \mathcal{P} \times \{1\}$.

https://doi.org/10.1112/S0024610705006423 Published online by Cambridge University Press

It is easily seen from Lemma 9(i) that

$$\inf\{\lambda(t), t \geqslant 0\} > 0 \quad \text{and} \quad \lim_{t\to\infty} \frac{\log \lambda(t)}{\log t} = 1/2, \qquad \mathbb{Q}\text{-almost surely.} \tag{21}$$

As a consequence,

$$\lim_{t\to\infty} (\log(1/|\Pi_1(t)|) - t\Phi'(\bar{p}) + a) \exp(t\Phi(\bar{p}))|\Pi_1(t)|^{\bar{p}+1} = 0, \qquad \mathbb{Q}\text{-almost surely,}$$

and since our goal is to check (19), this enables us to focus henceforth on

$$\begin{aligned}\tilde{M}(t) &= M_a(t) - (\log(1/|\Pi_1(t)|) - t\Phi'(\bar{p}) + a) \exp(t\Phi(\bar{p}))|\Pi_1(t)|^{\bar{p}+1} \\ &= \sum_{i=2}^{\infty} (\log(1/|\Pi_i(t)|) - t\Phi'(\bar{p}) + a) \exp(t\Phi(\bar{p}))\, |\Pi_i(t)|^{\bar{p}+1}, \quad \mathbb{Q}\text{-almost surely.}\end{aligned}$$

Next, we compute the conditional expectation of this quantity given $\mathcal{G}_1(\infty)$, the sigma-field generated by the restriction of the discrete point measure to the fibre $\mathbb{R}_+ \times \mathcal{P} \times \{1\}$, as follows. By construction of the fragmentation Π, each block $\Pi_i(t)$ for $i \geqslant 2$ is separated from 1 at some instant $r \in \mathcal{D}_1 \cap [0, t]$. More precisely, recall that at such an instant r, the block $\Pi_1(r-)$ splits into $\pi(r)_{|\Pi_1(r-)}$, and that the block after the split which contains 1 is $\Pi_1(r) = \pi_1(r) \cap \Pi_1(r-)$. Thus there is then some index $j \geqslant 2$ such that $\Pi_i(t) \subseteq \pi_j(r) \cap \Pi_1(r-)$, where $\pi_j(r)$ stands for the jth block of the partition $\pi(r)$. In other words, we may consider the partition of $\{2, \ldots\}$ whose blocks are of the type

$$B(r, j) = \{i \geqslant 2 : \Pi_i(t) \subseteq \pi_j(r) \cap \Pi_1(r-)\},$$

and then $(\Pi_i(t) : i \in B(r, j))$ forms a partition of $\pi_j(r) \cap \Pi_1(r-)$ which we now analyse.

Lemma 9(ii), standard properties of Poisson random measures, and the very construction of Π entail that for every $r \in [0, t]$ and $j \geqslant 2$, conditionally on $r \in \mathcal{D}_1$, $\Pi_1(r-)$ and $\pi_j(r)$, the partition $(\Pi_i(t) : i \in B(r, j))$ can be given in the form $\tilde{\Pi}(t - r)_{|\pi_j(r)\cap\Pi_1(r-)}$, where $\tilde{\Pi}$ is a homogeneous fragmentation distributed as Π under $\mathbb{P}$ and is independent of the sigma-field $\mathcal{G}_1(\infty)$. Recall from (7) that

$$\mathbb{E}\left(\sum_{i=1}^{\infty} |\Pi_i(t - r)|^{\bar{p}+1}\right) = \exp(-(t - r)\Phi(\bar{p})),$$

and (taking the derivative)

$$\mathbb{E}\left(\sum_{i=1}^{\infty} \log(1/|\Pi_i(t - r)|)|\Pi_i(t - r)|^{\bar{p}+1}\right) = (t - r)\Phi'(\bar{p}) \exp(-(t - r)\Phi(\bar{p})).$$

The analysis above now entails that

$$\begin{aligned}&\mathbb{Q}\left(\sum_{i=2}^{\infty} (\log(1/|\Pi_i(t)|) - t\Phi'(\bar{p}) + a) \exp(t\Phi(\bar{p}))\, |\Pi_i(t)|^{\bar{p}+1} \mid \mathcal{G}_1(\infty)\right) \\ &\quad = \sum_{r\in\mathcal{D}_1\cap[0,t]} \sum_{j=2}^{\infty} (\log(1/|\pi_j(r) \cap \Pi_1(r-)|) \\ &\qquad - r\Phi'(\bar{p}) + a) \exp(r\Phi(\bar{p}))|\pi_j(r) \cap \Pi_1(r-)|^{\bar{p}+1}.\end{aligned}$$

Now observe that there is the identity

$$|\pi_j(r) \cap \Pi_1(r-)| = |\pi_j(r)||\Pi_1(r-)| = |\pi_j(r)| \exp(-\xi(r-))$$

https://doi.org/10.1112/S0024610705006423 Published online by Cambridge University Press

for all $r \in \mathcal{D}_1$, $\mathbb{P}$-almost surely and hence also $\mathbb{Q}$-almost surely. Putting the pieces together, we get

$$\mathbb{Q}(\tilde{M}(t) \,|\, \mathcal{G}_1(\infty)) = \sum_{r \in \mathcal{D}_1 \cap [0,t]} \exp\{-(\bar{p}+1)(\lambda(r-) - a)\}\Sigma(r),$$

where

$$\Sigma(r) = \sum_{j=2}^{\infty} (\lambda(r-) - \log|\pi_j(r)|)\,|\pi_j(r)|^{\bar{p}+1}.$$

As pointed out by Lyons [**21**], by the conditional Fatou theorem, all that we need is to check that $\lim_{t\to\infty} \mathbb{Q}(\tilde{M}(t) \,|\, \mathcal{G}_1(\infty)) < \infty$, $\mathbb{Q}$-almost surely. In this direction, we compute the $(\mathbb{Q}, \mathcal{G}_1(t))$-predictable compensator corresponding to the point process $\{\Sigma(r), r \in \mathcal{D}_1\}$, and we find that

$$\begin{aligned}
&\lambda(r-)^{-1} \int_{\mathcal{P}} \mu(d\pi)(\lambda(r-) - \log|\pi_1|)|\pi_1|^{\bar{p}} \left(\sum_{j=2}^{\infty} (\lambda(r-) - \log|\pi_j|)|\pi_j|^{\bar{p}+1} \right) \\
&\qquad = \lambda(r-)^{-1} \int_{\mathcal{S}} \nu(d\mathbf{s}) \left\{ \left(\sum_{j=1}^{\infty} (\lambda(r-) - \log|s_j|)|s_j|^{\bar{p}+1} \right)^2 \right. \\
&\qquad\qquad \left. - \sum_{j=1}^{\infty} (\lambda(r-) - \log|s_j|)^2 |s_j|^{2\bar{p}} \right\}.
\end{aligned}$$

Using the fact that $\bar{p} > 0$, it is easily seen that this quantity can be bounded from above by $C\,(\lambda(r-) + 1 + 1/\lambda(r-))$ for some constant C that depends only on ν. All that we need now is to verify that the integral

$$\int_0^{\infty} (\lambda(r) + 1 + 1/\lambda(r)) \exp\{-(\bar{p}+1)(\lambda(r) - a)\}\, dr$$

converges $\mathbb{Q}$-almost surely, which is immediate from (21).

(iii) To ease the reading, let us denote

$$Y_i(t) = \exp(t\Phi(\bar{p}))(|\Pi(t)|_i^{\downarrow})^{\bar{p}+1}.$$

We first remark that for all $i \in \mathbb{N}$, $\lim_{t\to\infty} Y_i(t) = 0$, $\mathbb{P}$-almost surely, and we deduce from (18) that $M_a(t) \leqslant -M'(t) + aM(\bar{p}, t)$, for t large enough. Taking the limit as $t \to \infty$, we get

$$M_a(\infty) \leqslant -M'(\infty), \qquad \mathbb{P}\text{-almost surely}, \tag{22}$$

which proves that $-M'(\infty) \geqslant 0$, $\mathbb{P}$-almost surely.

From the fragmentation property at time 1, we may express $M'(1+t)$ in the form

$$-M'(1+t) = \sum_{i,j} Y_i(1) Y_{i,j}(t) \log \frac{1}{Y_i(1) Y_{i,j}(t)},$$

where $\{Y_{i,j}(\cdot), j \in \mathbb{N}\}$ for $i = 1, \ldots$ are independent copies of $\{Y_j(\cdot), j \in \mathbb{N}\}$, which are also independent of $\mathcal{G}(1)$. This yields

$$-M'(1+t) = \sum_i Y_i(1)(-M_i'(t)) + \sum_i \left(Y_i(1) \log \frac{1}{Y_i(1)} \right) M_i(t), \tag{23}$$

https://doi.org/10.1112/S0024610705006423 Published online by Cambridge University Press

where $\{M_i(\cdot), i \in \mathbb{N}\}$ (respectively, $\{M_i'(\cdot), i \in \mathbb{N}\}$) are independent copies of $M(\bar{p}, \cdot)$ (respectively $M'(\cdot)$) and independent of $\mathcal{G}(1)$. To get rid of the last infinite random combination of martingales converging to zero, we first establish the following technical result.

$$\lim_{t\to\infty} \sum_i \left(Y_i(1) \log \frac{1}{Y_i(1)} \right) M_i(t) = 0 \qquad \text{in probability under } \mathbb{P}. \tag{24}$$

Indeed, because the $|\Pi(1)|_i^{\downarrow}$, $i \in \mathbb{N}$ are ranked in decreasing order, and their sum is at most 1, we have $|\Pi(1)|_i^{\downarrow} \leqslant 1/i$ for every i, and thus $Y_i(1) < 1$ for $i > \mathrm{e}^{\Phi'(\bar{p})}$. The series $-M'(1) = \sum_i Y_i(1)|\log Y_i(1)|$ is absolutely convergent (and in L^1). Therefore, for every $\epsilon > 0$ there exists $k > \mathrm{e}^{\Phi'(\bar{p})}$ such that

$$\mathbb{E}\left(\sum_{k+1}^{\infty} Y_i(1)|\log Y_i(1)| \right) \leqslant \epsilon^2.$$

Since $\mathbb{E}(M_i(t)) = 1$ for all i, the Markov inequality enables us to write

$$\mathbb{P}\left(\sum_{k+1}^{\infty} (Y_i(1)|\log Y_i(1)|)\, M_i(t) > \epsilon \right) \leqslant \epsilon.$$

Since the sum of the k remaining terms converges $\mathbb{P}$-almost surely to 0, claim (24) is proved.

Now we are able to complete the proof of (iii). Assume that $\mathbb{P}(M'(\infty) = 0) > 0$. From (23) and (24) we may write

$$M'(\infty) = Y_1(1)M_1'(\infty) + Y_2(1)M_2'(\infty) + B,$$

where $B = \lim_t \sum_{i=3}^{\infty}$ is independent of $(M_1'(\infty), M_2'(\infty))$ conditionally on $\mathcal{G}_1$. Since $\mathbb{P}(M'(\infty) \leqslant 0) = 1$, this entails $\mathbb{P}(B \leqslant 0) = 1$ and

$$M'(\infty) \leqslant Y_1(1)M_1'(\infty) + Y_2(1)M_2'(\infty).$$

This implies that $\mathbb{P}(M'(\infty) = 0) \leqslant \mathbb{P}(M'(\infty) = 0)^2$, so we would have $\mathbb{P}(M'(\infty) = 0) = 1$, which contradicts (ii). □

Acknowledgements. We should like to thank an anonymous referee of this paper for insightful comments on a first draft of the work.

References

1. S. Asmussen and H. Hering, *Branching processes* (Birkhäuser, Boston, 1983).
2. G. Ben Arous and A. Rouault, 'Laplace asymptotics for reaction-diffusion equations', *Probab. Theory Related Fields* 97 (1993) 259–285.
3. J. Berestycki, 'Ranked fragmentations', *ESAIM Probab. Statist.* 6 (2002) 157–176.
4. J. Bertoin, 'Homogeneous fragmentation processes', *Probab. Theory Related Fields* 121 (2001) 301–318.
5. J. Bertoin, 'Self-similar fragmentations', *Ann. Inst. H. Poincaré* 38 (2002) 319–340.
6. J. Bertoin, 'The asymptotic behavior of fragmentation processes', *J. Europ. Math. Soc.* 5 (2003) 395–416.
7. J. Bertoin and A. Rouault, 'Asymptotical behaviour of the presence probability in branching random walks and fragmentations', 2004, arXiv:math.PR/0409547.
8. J. D. Biggins, 'Martingale convergence in the branching random walk', *J. Appl. Probab.* 14 (1977) 25–37.
9. J. D. Biggins, 'Uniform convergence of martingales in the branching random walk', *Ann. Probab.* 20 (1992) 137–151.
10. J. D. Biggins and A. E. Kyprianou, 'Measure change in multitype branching', *Adv. in Appl. Probab.* 36 (2004) 544–581.

https://doi.org/10.1112/S0024610705006423 Published online by Cambridge University Press

11. L. Chaumont, 'Sur certains processus de Lévy conditionnés à rester positifs', *Stochastics Stochastic Rep.* 47 (1994) 1–20.
12. B. Chauvin and A. Rouault, 'KPP equation and supercritical branching Brownian motion in the subcritical speed area. Application to spatial trees', *Probab. Theory Related Fields* 80 (1988) 299–314.
13. S. C. Harris, 'Travelling waves for the FKPP equation via probabilistic arguments', *Proc. Roy. Soc. Edinburgh Sect.* A 129 (1999) 503–517.
14. J. F. C. Kingman, 'Ergodic properties of continuous-time Markov processes and their discrete skeletons', *Proc. London Math. Soc.* 3 (1963) 593–604.
15. J. F. C. Kingman, 'The coalescent', *Stochastic Process. Appl.* 13 (1982) 235–248.
16. J. F. C. Kingman, *Poisson processes* (Clarendon Press, New York, 1993).
17. A. E. Kyprianou, 'Slow variation and uniqueness of solutions to the functional equation in the branching random walk', *J. Appl. Probab.* 35 (1998) 795–801.
18. A. E. Kyprianou, 'A note on branching Lévy processes', *Stochastic Process. Appl.* 82 (1999) 1–14.
19. A. E. Kyprianou, 'Travelling waves solutions to the K-P-P equation: alternative to Simon Harris' probabilistic argument', *Ann. Inst. H. Poincaré* (B) 40 (2004) 53–72.
20. Q. Liu, 'On generalized multiplicative cascades', *Stochastic Process. Appl.* 86 (2000) 263–286.
21. R. Lyons, 'A simple path to Biggins' martingale convergence for branching random walk', *Classical and modern branching processes* (ed. K. B. Athreya and P. Jagers, Springer, New York, 1997) 217–221.
22. R. Lyons, R. Pemantle and Y. Peres, 'Conceptual proofs of $L \log L$ criteria for mean behaviour of branching processes', *Ann. Probab.* 23 (1995) 1125–1138.
23. J. Neveu, 'Multiplicative martingales for spatial branching processes', *Seminar on Stochastic Processes* 1987, Progress in Probability Statistics 15 (ed. E. Cinlar, K. L. Chung and R. K. Getoor, Birkhäuser, 1988) 223–242.
24. J. Pitman, 'Coalescent with multiple collisions', *Ann. Probab.* 27 (1999) 1870–1902.
25. A. Rouault, 'Precise estimates of presence probabilities in the branching random walk', *Stochastic Process. Appl.* 44 (1993) 27–39.
26. K. Sato, *Lévy processes and infinitely divisible distributions* (Cambridge University Press, Cambridge, 1999).
27. K. Uchiyama, 'Spatial growth of a branching process of particles living in R^d', *Ann. Probab.* 10 (1982) 896–918.

Jean Bertoin
Laboratoire de Probabilités et Modèles Aléatoires and Institut Universitaire de France
Université Pierre et Marie Curie
175 rue du Chevaleret
F-75013 Paris
France

jbe@ccr.jussieu.fr

Alain Rouault
LAMA
Bâtiment Fermat
Université de Versailles
45 avenue des Etats-Unis
F-78035 Versailles Cedex
France

rouault@math.uvsq.fr

https://doi.org/10.1112/S0024610705006423 Published online by Cambridge University Press