Canad. Math. Bull. 2026, pp. 1–15
http://dx.doi.org/10.4153/S0008439526102343
© The Author(s), 2026. Published by Cambridge University Press on behalf of Canadian Mathematical Society

# Eigenvalue–eigenvector identity for dual Hermitian matrices

Sarishti Singh and Frédéric Morneau-Guérin

*Abstract.* This article studies an analog of the classical eigenvalue–eigenvector identity for dual Hermitian matrices. The classical identity relates the eigenvalues of a Hermitian matrix, the eigenvalues of its principal submatrices, and the entries of its eigenvectors. Assuming that the standard eigenvalues are simple, we establish the corresponding identity for dual Hermitian matrices and show that it naturally decomposes into a standard component and a dual component. The standard component recovers the classical Hermitian identity, while the dual component yields an explicit first-order correction formula relating eigenvalue and eigenvector corrections to perturbations of the corresponding principal submatrices. Numerical examples are presented to verify the theoretical results.

## 1 Introduction

For more than a century, the mathematical frameworks of dual numbers and dual quaternions, which we briefly review below, including their corresponding vectors and matrices, have been extensively studied and applied in various fields of mathematics and engineering. The theory of dual numbers originated in the late 19th century [5] and was further developed in the early 20th century through the notion of dual angles and related geometric constructions [33]. These early contributions provided a basis for employing dual numbers, along with their vector and matrix representations, in fields, such as kinematics, mechanics, and robotics [1, 11, 13, 19, 20, 27, 28].

In subsequent years [10, 12], dual quaternions, specifically their unit forms, have gained prominence in various modern applications, including neuroscience, hand–eye calibration, cooperative control of multiple agents, and simultaneous localization and mapping (SLAM) [2–4, 6, 14, 16, 23, 25, 29]. More recently, dual complex matrices [17, 32] have also been utilized in brain signal analysis [30]. These developments illustrate the continuing growth and relevance of dual matrix theory in both theoretical and applied settings.

The eigenvector–eigenvalue identity has evolved through a long and complex historical development. It has appeared, sometimes in slightly different or hidden

Received by the editors May 22, 2026 UTC; revised July 4, 2026 UTC; accepted July 14, 2026 UTC.
Published online on Cambridge Core July 20, 2026.
The first author gratefully acknowledges Mitacs Globalink (Canada) for providing partial funding for this research.
AMS subject classification: **15A18**, **15B57**, **93B60**.
Keywords: Dual numbers, dual Hermitian matrices, eigenvalue–eigenvector identity.

https://doi.org/10.4153/S0008439526102343 Published online by Cambridge University Press

forms, and has been rediscovered several times in different research areas, such as random matrix theory, numerical linear algebra, inverse eigenvalue problems, graph theory (including graph reconstruction, chemical graph theory, and walks on graphs), and even neutrino physics [7–9, 18, 26, 31].

While the identity remains valid for all Hermitian matrices and admits extensions to normal and diagonalizable matrices, it is particularly significant in the case of symmetric tridiagonal matrices, such as Jacobi matrices, which play a central role in several core algorithms of numerical linear algebra. In several instances, the identity did not appear explicitly as a direct relation between eigenvalues and eigenvectors, but rather emerged implicitly as an auxiliary result in other analytical developments. The form and notation of the identity also varied widely, making it difficult to locate with normal literature searches.

Interest in the identity was revived following an expository publication by Wolchover in Quanta Magazine [31], which discussed its modern rediscovery by Denton et al. [8, 9]. As reported in that article, Terence Tao remarked that "something this short and simple should have been in textbooks already." This observation reflects the surprising historical trajectory of an identity that had largely escaped attention despite its elementary nature. In subsequent work, Denton et al. [8] presented a concise survey of the identity, including several proofs and generalizations. Their rediscovery arose in the study of numerically stable formulas for the eigenvectors of neutrino oscillation Hamiltonians, where the authors derived the identity explicitly in the case of $3 \times 3$ matrices [9].

The study of eigenvalue problems for dual matrices remains in its early stages and continues to offer opportunities for further investigation. Pennestrì and Stefanelli [19, 20] formulated the problem of determining the eigenvalues and the singular values of matrices defined over dual numbers. Qi and Luo [24] demonstrated that a dual quaternion Hermitian matrix of order $n \times n$ possesses $n$ dual number eigenvalues and further established the singular value decomposition (SVD) for dual quaternion matrices. These results are also applicable to dual number matrices and dual complex matrices [22].

More recently, Qi and Cui [21] introduced a supplementary matrix approach for computing the eigenvalues of dual Hermitian matrices and demonstrated its applicability to problems in multi-agent formation control. Motivated by these developments, we establish in the present work an eigenvalue–eigenvector identity for dual Hermitian matrices.

To the best of our knowledge, no analog of the eigenvalue–eigenvector identity has previously been established for dual Hermitian matrices. The primary contribution of this article is the derivation of an explicit first-order correction to the classical eigenvalue–eigenvector identity in the setting of dual Hermitian matrices. Specifically, we show that the dual eigenvalue–eigenvector identity naturally decomposes into two parts. The standard component recovers the classical identity for Hermitian matrices, while the dual component yields an explicit first-order correction formula relating the corrections to the eigenvalues and eigenvectors to the perturbations of the corresponding principal submatrices. This decomposition extends the classical identity to the dual setting and provides a clear interpretation of the effect of infinitesimal perturbations on the spectral structure of dual Hermitian matrices.

https://doi.org/10.4153/S0008439526102343 Published online by Cambridge University Press

To illustrate the proposed identities, we present two examples and verify the results using the supplementary matrix framework introduced by Qi and Cui [21] for computing the eigenvalues and eigenvectors of dual Hermitian matrices.

## 2 Preliminaries

Let $\mathbb{R}$ and $\mathbb{C}$ denote the fields of real and complex numbers, respectively, and let $\mathbb{K}$ represent either $\mathbb{R}$ or $\mathbb{C}$. The corresponding sets of dual numbers and dual complex numbers are denoted by $\widehat{\mathbb{R}}$ and $\widehat{\mathbb{C}}$, respectively. We write $\widehat{\mathbb{K}}$ to denote either of these two sets. An element of $\widehat{\mathbb{K}}$ will be called a *dual element*.

A dual element is an expression of the form $a = a_s + a_d\varepsilon$, where $a_s, a_d \in \mathbb{K}$, and $\varepsilon$ is an infinitesimal unit satisfying $\varepsilon^2 = 0$ and commuting with all real or complex scalars. The quantities $a_s$ and $a_d$ are called the standard part and dual part of $a$, respectively. The conjugate of $a$ is defined by $a^* = a_s^* + a_d^*\varepsilon$, where $a_s^*$ and $a_d^*$ denote the ordinary conjugates of $a_s$ and $a_d$. A dual element is said to be *appreciable* if $a_s \neq 0$, and *infinitesimal* otherwise. Moreover, a dual element is said to be *invertible* if there exists a dual element $b$ such that $ab = ba = 1$. Equivalently, a dual element is invertible if and only if it is appreciable (see [21]).

A dual matrix $A = A_s + \varepsilon A_d \in \widehat{\mathbb{K}}^{n\times n}$ is said to be *Hermitian* if $A = A^*$. A dual vector $x = x_s + x_d\varepsilon$, where $x_s,\ x_d \in \mathbb{K}^n$, is called a *unit dual vector* if $\|x_s\|_2 = 1$ and $x_s^* x_d + x_d^* x_s = 0$, where $\|\cdot\|_2$ denotes the standard Euclidean norm.

Let $A = A_s + \varepsilon A_d \in \widehat{\mathbb{K}}^{n\times n}$ be a dual Hermitian matrix. Suppose that $Av_i = \lambda_i v_i$, where $\lambda_i = \lambda_{i,s} + \varepsilon\lambda_{i,d}$ is the $i$-th eigenvalue of $A$, and $v_i = u_i + \varepsilon w_i$ is a corresponding unit dual eigenvector, for $i = 1, \ldots, n$. Then, $\|u_i\|_2 = 1$ and $u_i^* w_i + w_i^* u_i = 0$, for $i = 1, \ldots, n$. Throughout the manuscript, $\mathrm{Re}(z)$ denotes the real part of a complex number $z \in \mathbb{C}$.

Let $p_A : \mathbb{C} \to \mathbb{C}$ denote the characteristic polynomial of $A$. Then

$$p_A(\lambda) = \det(\lambda I_n - A) = \prod_{k=1}^{n}(\lambda - \lambda_k),$$

where $I_n$ (or $I$) represents the $n \times n$ identity matrix.

Evaluating the derivative of $p_A$ at $\lambda = \lambda_i$ gives

$$P_i := p_A'(\lambda_i) = \prod_{k\neq i}(\lambda_i - \lambda_k).$$

For $j = 1, \ldots, n$, let $M_j$ denote the $(n-1) \times (n-1)$ principal submatrix of $A$ obtained by deleting the $j$-th row and column. Its characteristic polynomial is

$$p_{M_j}(\lambda) = \det(\lambda I_{n-1} - M_j) = \prod_{k=1}^{n-1}\big(\lambda - \lambda_k(M_j)\big).$$

The following result of Qi and Cui [21] provides the necessary framework for the subsequent results. Theorem 4.1 of [21] states that for a given dual Hermitian matrix $A = A_s + \varepsilon A_d \in \widehat{\mathbb{K}}^{n\times n}$, let $\lambda_{i,s}$ be a simple eigenvalue of $A_s$, and $u_i$ be the corresponding unit eigenvector of $\lambda_{i,s}$ for $i = 1, \ldots, n$. Then, $\lambda_i = \lambda_{i,s} + \varepsilon\lambda_{i,d}$ is a simple eigenvalue of $A$, where $\lambda_{i,d} = u_i^* A_d u_i$, with the corresponding eigenvector $v_i = u_i +$

https://doi.org/10.4153/S0008439526102343 Published online by Cambridge University Press

$\varepsilon w_i$, where $w_i$ satisfies

$$(A_s - \lambda_{i,s} I) w_i = (\lambda_{i,d} I - A_d) u_i,$$

for $i = 1, \ldots, n$.

## 3 Main results

The following proposition establishes the dual eigenvalue–eigenvector identity for dual Hermitian matrices with simple standard eigenvalues. Its proof adapts the adjugate-matrix argument of Denton et al. [8] to the dual setting.

***Proposition 3.1*** *Let $A = A_s + \varepsilon A_d \in \widehat{\mathbb{K}}^{n\times n}$, where $\mathbb{K} = \mathbb{R}$ or $\mathbb{C}$, be a dual Hermitian matrix. Assume that the eigenvalues $\lambda_{1,s}, \ldots, \lambda_{n,s}$ of $A_s$ are simple. Let*

$$A v_i = \lambda_i v_i, \quad \lambda_i = \lambda_{i,s} + \varepsilon \lambda_{i,d},$$

*where $v_i$ is a unit dual eigenvector corresponding to $\lambda_i$. Let $M_j$ be the principal submatrix of $A$ obtained by deleting the $j$-th row and column. Then, for each $i$, $j = 1, \ldots, n$,*

$$p_{M_j}(\lambda_i) = |v_{i,j}|^2 \, p'_A(\lambda_i), \tag{3.1}$$

*where $v_{i,j}$ is the $j$-th component of the unit dual eigenvector $v_i$ associated with $\lambda_i$.*

**Proof** By Theorem 4.1 of Qi and Cui [21], for each simple eigenvalue $\lambda_{i,s}$ of $A_s$, there exists a dual eigenvalue

$$\lambda_i = \lambda_{i,s} + \varepsilon \lambda_{i,d}$$

and an associated appreciable dual eigenvector $v_i = u_i + \varepsilon w_i$. The eigenvectors may be normalized so that $v_i^* v_i = 1$.

We first prove that $v_1, \ldots, v_n$ are mutually orthogonal. Since $\lambda_k = \lambda_{k,s} + \varepsilon \lambda_{k,d}$, we have

$$\lambda_k^* = \lambda_{k,s}^* + \varepsilon \lambda_{k,d}^*.$$

Because $A_s$ is Hermitian, $\lambda_{k,s} \in \mathbb{R}$. Moreover,

$$\lambda_{k,d} = u_k^* A_d u_k.$$

Since $A_d$ is Hermitian, $\lambda_{k,d} \in \mathbb{R}$. Hence,

$$\lambda_k^* = \lambda_k.$$

Since $A$ is dual Hermitian, $A^* = A$. For $i \neq k$, we have

$$A v_i = \lambda_i v_i, \qquad A v_k = \lambda_k v_k.$$

Thus,

$$v_k^* A v_i = \lambda_i v_k^* v_i.$$

https://doi.org/10.4153/S0008439526102343 Published online by Cambridge University Press

On the other hand,

$$v_k^* A v_i = (A v_k)^* v_i = (\lambda_k v_k)^* v_i = \lambda_k^* v_k^* v_i = \lambda_k v_k^* v_i.$$

It follows that

$$(\lambda_i - \lambda_k) v_k^* v_i = 0.$$

Since $\lambda_{i,s} \neq \lambda_{k,s}$, the dual number

$$\lambda_i - \lambda_k = (\lambda_{i,s} - \lambda_{k,s}) + \varepsilon(\lambda_{i,d} - \lambda_{k,d})$$

has nonzero standard part and is therefore invertible. Hence,

$$v_k^* v_i = 0 \qquad (i \neq k).$$

Together with $v_i^* v_i = 1$, this gives

$$V^* V = I, \qquad V = [v_1, \ldots, v_n].$$

Thus, $V$ is dual unitary. Indeed, $V^* V = I$ implies

$$\det(V)^* \det(V) = 1,$$

so $\det(V)$ is invertible in the dual-number algebra. Hence, $V$ is invertible and $V^{-1} = V^*$.

Since $A v_i = \lambda_i v_i$ for all $i$, we have

$$AV = V\Lambda, \qquad \Lambda = \operatorname{diag}(\lambda_1, \ldots, \lambda_n).$$

Multiplying on the right by $V^*$, we obtain

$$A = V \Lambda V^*.$$

Now let $\lambda$ be a dual number. Then

$$\lambda I - A = V(\lambda I - \Lambda) V^*.$$

The determinant and adjugate are well defined over the commutative algebra of dual numbers. Since the entries belong to a commutative algebra, the standard polynomial identity

$$\operatorname{adj}(PXP^{-1}) = P \operatorname{adj}(X) P^{-1}$$

holds for every invertible matrix $P$. This follows, for instance, from

$$\operatorname{adj}(AB) = \operatorname{adj}(B) \operatorname{adj}(A),$$

which is a polynomial identity valid over every commutative ring. Therefore,

$$\operatorname{adj}(\lambda I - A) = V \operatorname{adj}(\lambda I - \Lambda) V^*.$$

Since $\lambda I - \Lambda$ is diagonal, we have

$$\operatorname{adj}(\lambda I - \Lambda) = \operatorname{diag}\left( \prod_{m \neq 1} (\lambda - \lambda_m), \ldots, \prod_{m \neq n} (\lambda - \lambda_m) \right).$$

https://doi.org/10.4153/S0008439526102343 Published online by Cambridge University Press

Putting $\lambda = \lambda_i$, all diagonal entries of $\mathrm{adj}(\lambda_i I - \Lambda)$ vanish except the $i$-th one. Hence,

$$\mathrm{adj}(\lambda_i I - A) = \left(\prod_{m \neq i} (\lambda_i - \lambda_m)\right) v_i v_i^*.$$

Taking the $(j, j)$-entry of both sides gives

$$\det(\lambda_i I_{n-1} - M_j) = \left(\prod_{m \neq i} (\lambda_i - \lambda_m)\right) v_{i,j} v_{i,j}^*.$$

Since $v_{i,j} v_{i,j}^* = |v_{i,j}|^2$, we get

$$p_{M_j}(\lambda_i) = |v_{i,j}|^2 \prod_{m \neq i} (\lambda_i - \lambda_m).$$

It remains to identify the product with $p'_A(\lambda_i)$. Since $A = V \Lambda V^*$ and $V^* = V^{-1}$, we have

$$\begin{aligned} p_A(\lambda) &= \det(\lambda I - A) = \det\left(V(\lambda I - \Lambda) V^*\right) \\ &= \det(V) \det(\lambda I - \Lambda) \det(V^*) = \det(\lambda I - \Lambda) = \prod_{m=1}^{n} (\lambda - \lambda_m). \end{aligned}$$

Therefore,

$$p'_A(\lambda_i) = \prod_{m \neq i} (\lambda_i - \lambda_m).$$

Hence,

$$p_{M_j}(\lambda_i) = |v_{i,j}|^2 p'_A(\lambda_i).$$

This completes the proof. ∎

Assume that the standard eigenvalues $\lambda_{1,s}, \ldots, \lambda_{n,s}$ of $A_s$ are simple. The dual Hermitian principal submatrix $M_j$ of order $(n-1) \times (n-1)$, obtained by removing the $j$-th row and column of $A$, can be written in standard and dual parts as $M_j = M_{j,s} + \varepsilon M_{j,d}$ for $j = 1, \ldots, n$. For $i = 1, \ldots, n$, denote

$$\begin{aligned} p_{M_{j,s}}(\lambda_{i,s}) &= \det(\lambda_{i,s} I - M_{j,s}), \\ p'_{M_{j,s}}(\lambda_{i,s}) &= \frac{d}{d\lambda} (\det(\lambda I - M_{j,s}))\big|_{\lambda = \lambda_{i,s}}, \\ p'_{A_s}(\lambda_{i,s}) 7 &= \prod_{k \neq i} (\lambda_{i,s} - \lambda_{k,s}) := P_{s,i}. \end{aligned}$$

By the Jacobi formula [15], we obtain

$$\mathrm{tr}(\mathrm{adj}(\lambda_{i,s} I - M_{j,s})) = \frac{d}{d\lambda} (\det(\lambda I - M_{j,s}))\big|_{\lambda = \lambda_{i,s}} = p'_{M_{j,s}}(\lambda_{i,s}). \tag{3.2}$$

The following theorem decomposes the dual eigenvalue–eigenvector identity into its standard and dual components. The standard component coincides with the classical identity, whereas the dual component provides an explicit first-order correction formula describing the interaction between the perturbations of the eigenvalues, eigenvectors, and principal submatrices.

https://doi.org/10.4153/S0008439526102343 Published online by Cambridge University Press

***Theorem 3.2*** *Let $A = A_s + \varepsilon A_d \in \widehat{\mathbb{K}}^{n\times n}$ be a dual Hermitian matrix. Assume that the standard eigenvalues $\lambda_{1,s}, \ldots, \lambda_{n,s}$ of $A_s$ are simple. Let $v_i = u_i + \varepsilon w_i$ be the unit dual vector corresponding to $\lambda_i = \lambda_{i,s} + \varepsilon\lambda_{i,d}$, $i = 1, \ldots, n$. Then, for each $j = 1, \ldots, n$, we have the following:*

(1) *Standard identity:*

$$|u_{i,j}|^2\, p'_{A_s}(\lambda_{i,s}) = p_{M_{j,s}}(\lambda_{i,s}). \tag{3.3}$$

(2) *Dual correction identity:*

$$\begin{aligned} &2\,\mathrm{Re}\left(u_{i,j}\overline{w_{i,j}}\right)\, p'_{A_s}(\lambda_{i,s}) + |u_{i,j}|^2\, p'_{A_s}(\lambda_{i,s}) \sum_{m\neq i} \frac{\lambda_{i,d} - \lambda_{m,d}}{\lambda_{i,s} - \lambda_{m,s}} \\ &\qquad = \lambda_{i,d}\, p'_{M_{j,s}}(\lambda_{i,s}) - \mathrm{tr}\big(\mathrm{adj}(\lambda_{i,s}I - M_{j,s})M_{j,d}\big)\,, \end{aligned} \tag{3.4}$$

*where $u_{i,j}$ and $w_{i,j}$ denote the $j$-th components of $u_i$ and $w_i$, respectively.*

**Proof** Let $X_s := \lambda_{i,s}I - M_{j,s}$ and $Y := \lambda_{i,d}I - M_{j,d}$. Then

$$\begin{aligned} p_{M_j}(\lambda_i) &= \det(\lambda_i I - M_j) \\ &= \det\left((\lambda_{i,s} + \varepsilon\lambda_{i,d})I - (M_{j,s} + \varepsilon M_{j,d})\right) \\ &= \det(X_s + \varepsilon Y). \end{aligned}$$

From Jacobi's formula [15], which states that for any differentiable matrix function $X(t)$,

$$\frac{d}{dt}\det(X(t)) = \mathrm{tr}\left(\mathrm{adj}(X(t))\frac{dX(t)}{dt}\right).$$

Consider

$$X(t) = X_s + tY.$$

Then the scalar function $f(t) := \det(X(t))$ has the first-order Taylor expansion about $t = 0$,

$$f(t) = f(0) + tf'(0) + O(t^2).$$

Since $f(0) = \det(X_s)$ and, by Jacobi's formula,

$$f'(0) = \mathrm{tr}(\mathrm{adj}(X_s)Y)\,,$$

we obtain

$$\det(X_s + tY) = \det(X_s) + t\,\mathrm{tr}(\mathrm{adj}(X_s)Y) + O(t^2).$$

Substituting $t = \varepsilon$ and using $\varepsilon^2 = 0$ yields

$$\det(X_s + \varepsilon Y) = \det(X_s) + \varepsilon\,\mathrm{tr}(\mathrm{adj}(X_s)Y)\,.$$

Now

$$\det(X_s) = \det(\lambda_{i,s}I - M_{j,s}) = p_{M_{j,s}}(\lambda_{i,s}),$$

https://doi.org/10.4153/S0008439526102343 Published online by Cambridge University Press

and

$$\begin{aligned}\operatorname{tr}\big(\operatorname{adj}(X_s)Y\big) &= \operatorname{tr}\big(\operatorname{adj}(\lambda_{i,s}I - M_{j,s})(\lambda_{i,d}I - M_{j,d})\big)\\ &= \lambda_{i,d}\operatorname{tr}\big(\operatorname{adj}(\lambda_{i,s}I - M_{j,s})\big) - \operatorname{tr}\big(\operatorname{adj}(\lambda_{i,s}I - M_{j,s})M_{j,d}\big).\end{aligned}$$

By (3.2),

$$\operatorname{tr}\big(\operatorname{adj}(\lambda_{i,s}I - M_{j,s})\big) = \frac{d}{d\lambda}\det(\lambda I - M_{j,s})\Big|_{\lambda=\lambda_{i,s}} = p'_{M_{j,s}}(\lambda_{i,s}).$$

Hence,

$$p_{M_j}(\lambda_i) = p_{M_{j,s}}(\lambda_{i,s}) + \varepsilon\Big(\lambda_{i,d}p'_{M_{j,s}}(\lambda_{i,s}) - \operatorname{tr}\big(\operatorname{adj}(\lambda_{i,s}I - M_{j,s})M_{j,d}\big)\Big). \tag{3.5}$$

Now

$$P_i = \prod_{k\neq i}(\lambda_i - \lambda_k) = \prod_{k\neq i}\big((\lambda_{i,s} - \lambda_{k,s}) + \varepsilon(\lambda_{i,d} - \lambda_{k,d})\big).$$

Since the standard eigenvalues are simple, we have

$$\lambda_{i,s} - \lambda_{k,s} \neq 0, \quad k \neq i.$$

Hence, each dual number

$$(\lambda_{i,s} - \lambda_{k,s}) + \varepsilon(\lambda_{i,d} - \lambda_{k,d})$$

has a nonzero standard part and is therefore invertible in the dual-number algebra. Factoring out the standard part from each factor, we obtain

$$P_i = \prod_{k\neq i}(\lambda_{i,s} - \lambda_{k,s})\prod_{k\neq i}\left(1 + \varepsilon\frac{\lambda_{i,d} - \lambda_{k,d}}{\lambda_{i,s} - \lambda_{k,s}}\right).$$

Setting

$$P_{s,i} = \prod_{k\neq i}(\lambda_{i,s} - \lambda_{k,s}).$$

Since $\varepsilon^2 = 0$, every term in the product containing two or more factors of $\varepsilon$ vanishes. Consequently, only the constant term and the terms containing exactly one factor of $\varepsilon$ contribute to the expansion. Therefore,

$$\prod_{k\neq i}\left(1 + \varepsilon\frac{\lambda_{i,d} - \lambda_{k,d}}{\lambda_{i,s} - \lambda_{k,s}}\right) = 1 + \varepsilon\sum_{k\neq i}\frac{\lambda_{i,d} - \lambda_{k,d}}{\lambda_{i,s} - \lambda_{k,s}}.$$

Hence,

$$P_i = P_{s,i} + \varepsilon P_{s,i}\sum_{k\neq i}\frac{\lambda_{i,d} - \lambda_{k,d}}{\lambda_{i,s} - \lambda_{k,s}}.$$

Setting

$$P_{d,i} := P_{s,i}\sum_{k\neq i}\frac{\lambda_{i,d} - \lambda_{k,d}}{\lambda_{i,s} - \lambda_{k,s}},$$

https://doi.org/10.4153/S0008439526102343 Published online by Cambridge University Press

we obtain

$$P_i = P_{s,i} + \varepsilon P_{d,i}. \tag{3.6}$$

Now, we expand $|v_{i,j}|^2$ as

$$\begin{aligned} |v_{i,j}|^2 &= (u_{i,j} + \varepsilon w_{i,j})(\overline{u_{i,j}} + \varepsilon \overline{w_{i,j}}) \\ &= u_{i,j}\overline{u_{i,j}} + \varepsilon(u_{i,j}\overline{w_{i,j}} + w_{i,j}\overline{u_{i,j}}) + \varepsilon^2(w_{i,j}\overline{w_{i,j}}). \end{aligned}$$

Since $\varepsilon^2 = 0$, the last term vanishes. Therefore,

$$|v_{i,j}|^2 = u_{i,j}\overline{u_{i,j}} + \varepsilon(u_{i,j}\overline{w_{i,j}} + w_{i,j}\overline{u_{i,j}}) = |u_{i,j}|^2 + 2\varepsilon \operatorname{Re}(u_{i,j}\overline{w_{i,j}}). \tag{3.7}$$

Using (3.6) and (3.7), we obtain

$$\begin{aligned} |v_{i,j}|^2 \, p'_A(\lambda_i) &= (P_{s,i} + \varepsilon P_{d,i})(|u_{i,j}|^2 + 2\varepsilon \operatorname{Re}(u_{i,j}\overline{w_{i,j}})) \\ &= P_{s,i}|u_{i,j}|^2 + \varepsilon\big(2\operatorname{Re}(u_{i,j}\overline{w_{i,j}})P_{s,i} + |u_{i,j}|^2 P_{d,i}\big). \end{aligned} \tag{3.8}$$

Using (3.5) and (3.8) in (3.1), we obtain

$$\begin{aligned} &p_{M_{j,s}}(\lambda_{i,s}) + \varepsilon\Big(\lambda_{i,d}\, p'_{M_{j,s}}(\lambda_{i,s}) - \operatorname{tr}\big(\operatorname{adj}(\lambda_{i,s}I - M_{j,s})M_{j,d}\big)\Big) \\ &\quad = P_{s,i}|u_{i,j}|^2 + \varepsilon\Big(2\operatorname{Re}\left(u_{i,j}\overline{w_{i,j}}\right)P_{s,i} + |u_{i,j}|^2 P_{d,i}\Big). \end{aligned} \tag{3.9}$$

Equating the standard (non $\varepsilon$) parts of (3.9) gives the standard identity (3.3). Substituting the value $P_{s,i}$ and $P_{d,i}$, and equating the $\varepsilon$ parts of (3.9) gives the dual correction identity (3.4). ∎

***Corollary 3.3*** *Under the assumptions of Theorem 3.2, identity (3.1) holds if and only if the standard identity (3.3) and the dual correction identity (3.4) are simultaneously satisfied.*

**Proof** From the proof of Theorem 3.2, equation (3.9) is precisely the decomposition of

$$p_{M_j}(\lambda_i) = |v_{i,j}|^2 \, p'_A(\lambda_i)$$

into its standard and dual parts. By equating the standard and dual parts of both sides, we obtain (3.3) and (3.4). Conversely, if (3.3) and (3.4) hold, then the standard and dual parts of both sides coincide, and therefore identity (3.1) holds. ∎

***Remark 3.4*** Equation (3.4) is new and couples the eigenvalue corrections $\lambda_{i,d}$, the eigenvector corrections $w_i$, and the dual perturbations $A_d, M_{j,d}$. It provides an explicit first-order correction to the classical eigenvalue–eigenvector identity. It quantifies how perturbations in the dual part of a Hermitian matrix influence both the eigenvalue corrections and the eigenvector corrections through the corresponding principal submatrices.

https://doi.org/10.4153/S0008439526102343 Published online by Cambridge University Press

## 4 Illustrative examples

The following examples illustrate the main ideas of Theorem 3.2 in both symbolic and numerical settings. We first consider a symbolic $2 \times 2$ example to highlight the structure of the standard and dual correction identities in a simple setting. We then present a numerical $3 \times 3$ example to demonstrate how the identities can be verified concretely for nontrivial dual Hermitian matrices.

***Example 4.1*** Consider the dual Hermitian matrix

$$A = \begin{pmatrix} a & c \\ \overline{c} & d \end{pmatrix} + \varepsilon \begin{pmatrix} \alpha & \gamma \\ \overline{\gamma} & \delta \end{pmatrix},$$

where $a, d, \alpha, \delta \in \mathbb{R}$ and $c, \gamma \in \mathbb{C}$.

Assume that the standard part $A_s$ has distinct eigenvalues

$$\lambda_{1,s},\ \lambda_{2,s},$$

with corresponding orthonormal eigenvectors

$$u_1 = \begin{pmatrix} u_{1,1} \\ u_{1,2} \end{pmatrix}, \text{ and } u_2 = \begin{pmatrix} u_{2,1} \\ u_{2,2} \end{pmatrix}.$$

Let

$$\lambda_i = \lambda_{i,s} + \varepsilon\lambda_{i,d}, \text{ and } v_i = u_i + \varepsilon w_i,$$

for $i = 1, 2$. For $i = j = 1$, the principal submatrix is

$$M_1 = d + \varepsilon\delta,$$

so that

$$p_{M_1}(\lambda_1) = (\lambda_{1,s} - d) + \varepsilon(\lambda_{1,d} - \delta).$$

Since

$$p'_A(\lambda_1) = \lambda_1 - \lambda_2,$$

we have

$$p'_A(\lambda_1) = (\lambda_{1,s} - \lambda_{2,s}) + \varepsilon(\lambda_{1,d} - \lambda_{2,d}).$$

Moreover,

$$|v_{1,1}|^2 = |u_{1,1}|^2 + 2\varepsilon\,\mathrm{Re}(u_{1,1}\overline{w_{1,1}}).$$

Using

$$p_{M_1}(\lambda_1) = |v_{1,1}|^2 p'_A(\lambda_1),$$

and comparing standard and dual parts, we obtain

$$\lambda_{1,s} - d = |u_{1,1}|^2(\lambda_{1,s} - \lambda_{2,s}),$$

https://doi.org/10.4153/S0008439526102343 Published online by Cambridge University Press

Table 1: Computation of eigenvalues and eigenvectors of $A = A_s + \varepsilon A_d$, where $A_s$ and $A_d$ are given in (4.1).

| $i$ | Eigenvalue $\lambda_i = \lambda_{i,s} + \varepsilon\lambda_{i,d}$ | $u_i$ | $w_i$ |
|---|---|---|---|
| 1 | $3.68893048 + \varepsilon\, 0.51805479$ | $\begin{bmatrix} -0.53776387 \\ -0.82747448 \\ -0.16154260 \end{bmatrix}$ | $\begin{bmatrix} -0.07054694 \\ 0.05121676 \\ -0.02750337 \end{bmatrix}$ |
| 2 | $1.54365251 - \varepsilon\, 0.62078441$ | $\begin{bmatrix} -0.70673578 \\ 0.54691183 \\ -0.44878947 \end{bmatrix}$ | $\begin{bmatrix} -0.00574458 \\ 0.08965822 \\ 0.11830724 \end{bmatrix}$ |
| 3 | $0.76741701 + \varepsilon\, 0.15272962$ | $\begin{bmatrix} -0.45971139 \\ 0.12717482 \\ 0.87891524 \end{bmatrix}$ | $\begin{bmatrix} 0.09135623 \\ -0.05232624 \\ 0.05535469 \end{bmatrix}$ |

which is the classical eigenvector–eigenvalue identity, together with the dual correction identity

$$\lambda_{1,d} - \delta = 2\,\mathrm{Re}(u_{1,1}\overline{w_{1,1}})(\lambda_{1,s} - \lambda_{2,s}) + |u_{1,1}|^2(\lambda_{1,d} - \lambda_{2,d}).$$

This example illustrates how the dual identity naturally extends the classical relation through an additional correction term.

***Example 4.2*** Consider a dual Hermitian matrix $A = A_s + \varepsilon A_d \in \widehat{\mathbb{R}}^{3\times 3}$ with $A_s$ and $A_d$ given as

$$A_s = \begin{pmatrix} 2 & 1 & 0.5 \\ 1 & 3 & 0.2 \\ 0.5 & 0.2 & 1 \end{pmatrix} \quad \text{and} \quad A_d = \begin{pmatrix} 0.10 & 0.50 & -0.20 \\ 0.50 & 0 & 0.3 \\ -0.20 & 0.3 & -0.05 \end{pmatrix}. \tag{4.1}$$

For each simple standard eigenvalue $\lambda_{i,s}$, $i = 1, 2, 3$, a unit eigenvector $u_i$ of $A_s$ is computed. The corresponding dual eigenvalue correction is given by

$$\lambda_{i,d} = u_i^* A_d u_i.$$

The eigenvector correction $w_i$ is then obtained by solving

$$(A_s - \lambda_{i,s} I)w_i = (\lambda_{i,d} I - A_d)u_i,$$

as given by Theorem 4.1 of Qi and Cui [21]. The corresponding dual eigenvectors are

$$v_i = u_i + \varepsilon w_i, \quad i = 1, 2, 3,$$

where $v_i$ is chosen so that $v_i^* v_i = 1$. We now verify Theorem 3.2 for the dominant eigenvalue

$$\lambda_1 = 3.68893048 + \varepsilon\, 0.51805479.$$

The purpose of this example is to illustrate explicitly how the standard identity and the dual correction identity interact with the principal submatrices of $A$ (Table 1).

https://doi.org/10.4153/S0008439526102343 Published online by Cambridge University Press

We first consider the case $j = 1$, corresponding to the removal of the first row and column of $A$. In this case, the quantities appearing in equation (3.4) are computed directly from the associated principal submatrices $M_{1,s}$ and $M_{1,d}$.

Corresponding to $i = j = 1$, the matrices $M_{1,s}$ and $M_{1,d}$ are obtained as

$$M_{1,s} = \begin{pmatrix} 3 & 0.2 \\ 0.2 & 1 \end{pmatrix} \text{ and } M_{1,d} = \begin{pmatrix} 0 & 0.3 \\ 0.3 & -0.05 \end{pmatrix},$$

respectively. For the dual correction identity (3.4),

$$\operatorname{adj}(\lambda_{1,s} I - M_{1,s}) = \begin{pmatrix} \lambda_{1,s} - 1 & 0.2 \\ 0.2 & \lambda_{1,s} - 3 \end{pmatrix}.$$

Using

$$p'_{M_{1,s}}(\lambda_{1,s}) = \operatorname{tr}(\operatorname{adj}(\lambda_{1,s} I - M_{1,s})),$$

we obtain

$$p'_{M_{1,s}}(\lambda_{1,s}) = (\lambda_{1,s} - 1) + (\lambda_{1,s} - 3) = 3.377861.$$

Moreover,

$$\operatorname{tr}\left(\operatorname{adj}(\lambda_{1,s} I - M_{1,s}) M_{1,d}\right) = \frac{3}{25} - \frac{\lambda_{1,s} - 3}{20} = 0.085553.$$

- The quantity

$$P_{s,1} = \prod_{k \neq 1} (\lambda_{1,s} - \lambda_{k,s}) = p'_{A_s}(\lambda_{1,s})$$

  measures the spectral separation of the eigenvalue $\lambda_{1,s}$ from the remaining eigenvalues of $A_s$. Numerically,

$$P_{s,1} = 6.267458.$$

- The corresponding dual correction term is

$$P_{d,1} = P_{s,1} \sum_{k \neq 1} \frac{\lambda_{1,d} - \lambda_{k,d}}{\lambda_{1,s} - \lambda_{k,s}},$$

  which captures the perturbative contribution of the dual eigenvalue corrections. In this example,

$$P_{d,1} = 4.110858.$$

- The left-hand side of the dual correction identity (3.4) is therefore

$$\mathrm{LHS}_\varepsilon(1,1) := \lambda_{1,d} p'_{M_{1,s}}(\lambda_{1,s}) - \operatorname{tr}(\operatorname{adj}(\lambda_{1,s} I - M_{1,s}) M_{1,d}) = 1.664364.$$

- The right-hand side becomes

$$\mathrm{RHS}_\varepsilon(1,1) := 2\,\mathrm{Re}(u_{1,1}\overline{w_{1,1}}) P_{s,1} + |u_{1,1}|^2 P_{d,1} = 1.664364.$$

Since $\mathrm{LHS}_\varepsilon(1,1) = \mathrm{RHS}_\varepsilon(1,1) = 1.664364$, the dual identity is verified for $i = j = 1$.

https://doi.org/10.4153/S0008439526102343 Published online by Cambridge University Press

For the standard identity (3.3), we compute

$$p_{M_{1,s}}(\lambda_{1,s}) = \det(\lambda_{1,s}I - M_{1,s}) = 1.812486,$$

while

$$P_{s,1}|u_{1,1}|^2 = 1.812486.$$

Thus, the standard identity is also verified. Since $p_{M_{1,s}}(\lambda_{1,s}) = P_{s,1}|u_{1,1}|^2 = 1.812486$, the standard identity is verified for $i = j = 1$.

The same procedure may be repeated for $j = 2$. In this case,

$$M_{2,s} = \begin{pmatrix} 2 & 0.5 \\ 0.5 & 1 \end{pmatrix}, \text{ and } M_{2,d} = \begin{pmatrix} 0.10 & -0.20 \\ -0.20 & -0.05 \end{pmatrix}.$$

Substituting these matrices into (3.4) gives

$$\mathrm{LHS}_\varepsilon(1,2) = 2.283519,$$

and

$$\mathrm{RHS}_\varepsilon(1,2) = 2.283525,$$

which agree up to rounding error, thereby confirming the dual correction identity.

For standard identity (3.3),

- $p_{M_{2,s}}(\lambda_{1,s}) = \det(\lambda_{1,s}I - M_{2,s}) = (\lambda_{1,s} - 3)(\lambda_{1,s} - 1) - 0.04 = 1.812486$;
- $P_{s,1}|u_{1,2}|^2 = 1.812486$.

Since $p_{M_{2,s}}(\lambda_{1,s}) = P_{s,1}|u_{1,2}|^2 = 1.812486$, the standard identity is verified for $i = 1$, $j = 2$. Similarly, the results can be verified for other values of $i, j = 1, 2, 3$.

This example shows how the classical eigenvector–eigenvalue identity for the standard part $A_s$ extends to the dual setting through an additional correction term involving both the dual perturbation matrix and the dual eigenvector corrections.

## 5 Conclusion

In this work, we established an eigenvalue–eigenvector identity for dual Hermitian matrices. The classical identity was decomposed into standard and dual components, leading to an explicit dual correction formula involving eigenvalue perturbations, eigenvector corrections, and principal submatrices.

Several directions remain open for future investigation. These include extensions to dual normal matrices, dual quaternion Hermitian matrices, and more general structured eigenvalue problems. It would also be of interest to investigate whether analogous correction identities arise in higher-order nilpotent or perturbative algebraic settings.

**Data availability statement** No datasets generated or analyzed during this work.

https://doi.org/10.4153/S0008439526102343 Published online by Cambridge University Press

**Acknowledgements** hors are grateful to the anonymous referee for their helpful comments and suggestions, which have significantly improved the presentation and overall quality of this manuscript.

**Competing interests** The authors declare that they have no possible competing interests.

## References

[1] J. Angeles, *The dual generalized inverses and their applications in kinematic synthesis*. In: J. Lenarcic and M. Husty (eds.), *Latest advances in robot kinematics*, Springer Netherlands, Dordrecht, 2012, pp. 1–10.
[2] G. Brambley and J. Kim, *Unit dual quaternion-based pose optimisation for visual runway observations*. IET Cyber-Syst. Robot. **2**(2020), no. 4, 181–189.
[3] S. Bultmann, K. Li, and U. D. Hanebeck, *Stereo visual SLAM based on unscented dual quaternion filtering*. In: *2019 22th international conference on information fusion (FUSION)*, IEEE, Ottawa, ON, 2019, pp. 1–8.
[4] J. Cheng, J. Kim, Z. Jiang, and W. Che, *Dual quaternion-based graphical SLAM*. Robot. Auton. Syst. **77**(2016), 15–24.
[5] Clifford, *Preliminary sketch of biquaternions*. Proc. Lond. Math. Soc. **s1-4**(1871), no. 1, 381–395.
[6] K. Daniilidis, *Hand-eye calibration using dual quaternions*. Int. J. Robot. Res. **18**(1999), no. 3, 286–298.
[7] J. W. Demmel, *Applied numerical linear algebra*, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1997.
[8] P. B. Denton, S. J. Parke, T. Tao, and X. Zhang, *Eigenvectors from eigenvalues: a survey of a basic identity in linear algebra*. Bull. Amer. Math. Soc. **59**(2021), no. 1, 2.
[9] P. B. Denton, S. J. Parke, and X. Zhang, *Neutrino oscillations in matter via eigenvalues*. Phys. Rev. D **101**(2020), 93001.
[10] W. Ding, Y. Li, and M. Wei, *A new structure-preserving method for dual quaternion Hermitian eigenvalue problems*. Appl. Math. Lett. **163**(2025), 109432.
[11] I. Fischer, *Dual-number methods in kinematics, statics and dynamics*, CRC Press, Boca Raton, Florida, 2017.
[12] Z.-H. Gao, Q.-W. Wang, and L.-M. Xie, *A novel Moore–Penrose inverse to dual quaternion matrices with applications*. Appl. Math. Lett. **172**(2025), 109727.
[13] Y.-L. Gu and J. Luh, *Dual-number transformation and its applications to robotics*. IEEE J. Robot. Autom. **3**(1987), no. 6, 615–623.
[14] I. Hadi, T. Bendory, and N. Sharon, *SE(3) synchronization by eigenvectors of dual quaternion matrices*. Inf. Inference J. IMA **13**(2024), no. 3, iaae014, 7 pp.
[15] R. A. Horn and C. R. Johnson, *Matrix analysis*, Cambridge University Press, Cambridge, 2012.
[16] G. Leclercq, P. Lefèvre, and G. Blohm, *3D kinematics using dual quaternions: Theory and applications in neuroscience*. Front. Behav. Neurosci. **7**(2013), 7.
[17] C. Liu, Y. Wei, and P. Xie, *On the relation between the exponential of real matrices and that of dual matrices*. Appl. Math. Lett. **163**(2025), 109466.
[18] P. V. Mieghem, *Graph spectra for complex networks*, Cambridge University Press, Cambridge, 2010.
[19] E. Pennestrì and R. Stefanelli, *Linear algebra and numerical algorithms using dual numbers*. Multibody Syst. Dyn. **18**(2007), no. 3, 323–344.
[20] E. Pennestrì and P. P. Valentini, *Linear dual algebra algorithms and their application to kinematics*. In: Edited by C. L. Bottasso. *Multibody dynamics: Computational methods and applications*, Springer, Dordrecht, The Netherlands, 2009, pp. 207–229.
[21] L. Qi and C. Cui, *Eigenvalues of dual Hermitian matrices with application information control*. SIAM J. Matrix Anal. Appl. **45**(2024), no. 4, 2135–2154.
[22] L. Qi and C. Cui, *Eigenvalues and Jordan forms of dual complex matrices*. Commun. Appl. Math. Comput. **7**(2025), no. 4, 1225–1241.
[23] L. Qi, C. Cui, and C. Ouyang, *Unit dual quaternion directed graphs, formation control and general weighted directed graphs*. Discret. Appl. Math. **372**(2025), 197–209.
[24] L. Qi and Z. Luo, *Eigenvalues and singular values of dual quaternion matrices*. Pac. J. Optim. **19**(2023), 257–272.

https://doi.org/10.4153/S0008439526102343 Published online by Cambridge University Press

[25] L. Qi, X. Wang, and Z. Luo, *Dual quaternion matrices in multi-agent formation control*. Commun. Math. Sci. **21**(2023), no. 7, 1865–1874.
[26] M. Stawiska, *A generalized eigenvector–eigenvalue identity from the viewpoint of exterior algebra*. Adv. Appl. Clifford Algebr. **35**(2025), no. 2, 1–12.
[27] F. E. Udwadia, *Dual generalized inverses and their use in solving systems of linear dual equations*. Mech. Mach. Theory **156**(2021), 104158.
[28] H. Wang, *Characterizations and properties of the MPDGI and DMPGI*. Mech. Mach. Theory **158**(2021), 104212.
[29] X. Wang, C. Yu, and Z. Lin, *A dual quaternion solution to attitude and position control for rigid-body coordination*. IEEE Trans. Robot. **28**(2012), no. 5, 1162–1170.
[30] T. Wei, W. Ding, and Y. Wei, *Singular value decomposition of dual matrices and its application to traveling wave identification in the brain*. SIAM J. Matrix Anal. Appl. **45**(2024), no. 1, 634–660.
[31] N. Wolchover, *Neutrinos lead to unexpected discovery in basic math*. Quanta Magazine, 2019.
[32] R. Xu, T. Wei, Y. Wei, and P. Xie, *QR decomposition of dual matrices and its application*. Appl. Math. Lett. **156**(2024), 109144.
[33] K. Zindler, *Geometrie der Dynamen: Von E. Study. Leipzig, bei BG Teubner, 1903. XIII und 603 S. Preis 21 M.(geb. 23 M.)*. Monatsh. Math. Phys. **14**(1903), no. 1, A70–A75.

*Département de mathématiques et de statistique, Université Laval, Québec, QC, G1V 0A6, Canada;*
*and*
*Département Éducation, Université TÉLUQ, Québec, QC, G1K 9H6, Canada*
*e-mail*: sarishtisingh2510@gmail.com

*Département Éducation, Université TÉLUQ, Québec, QC, G1K 9H6, Canada*
*e-mail*: frederic.morneau-guerin@teluq.ca

https://doi.org/10.4153/S0008439526102343 Published online by Cambridge University Press