Laser and Particle Beams

cambridge.org/lpb

Direct electron acceleration with a linearly polarized laser beam in a two-dimensional magnetized plasma channel

Mohammad Ghorbanalilu and Nasim Nozarnejad

Department of Physics, Shahid Beheshti University, G. C., Tehran, Iran

Research Article

Cite this article: Ghorbanalilu M, Nozarnejad N (2019). Direct electron acceleration with a linearly polarized laser beam in a two-dimensional magnetized plasma channel. *Laser and Particle Beams* **37**, 428–434. https://doi.org/10.1017/S0263034619000703

Received: 23 July 2019
Revised: 21 September 2019
Accepted: 21 September 2019
First published online: 18 October 2019

Keywords:
Electron beams; filamentation instability; magnetized plasma; two-fluid model

Author for correspondence:
M. Ghorbanalilu, Department of Physics, Shahid Beheshti University, G. C., Evin, Tehran, Iran. E-mail: m_alilu@sbu.ac.ir

Abstract

We examine the electron acceleration induced by an ultra-relativistic intensity laser–plasma interaction in a two-dimensional plasma channel in the presence of a self-generated transverse magnetic field. We find that the electron dynamics is strongly affected by the laser pulse polarization angle, plasma density, and magnetic field strength. We investigate in detail, the dependencies of the electron acceleration in terms of different parameters and find excellent agreement with non-magnetized plasma in the absence of the magnetic field. The numerical results show that the self-generated magnetic field plays a constructive role in the electron acceleration process. It is shown that electron acceleration is more affected by self-generated magnetic field for the laser radiations with large polarization angles. The numerical results show the maximum enhancement for electron acceleration for a laser radiation with polarization angle $\theta = \pi/2$.

Introduction

A broad range of applications, such as GeV electrons sources (Wang *et al.*, 2013), multi-MeV protons, ion, and gamma ray sources (Fuchs *et al.*, 2006; Kneip *et al.*, 2008; Flippo *et al.*, 2010; Cipiccia *et al.*, 2011), have been employed in relativistic laser target interaction in a properly chosen experimental setup. These applications reveal the importance of investigation of the laser interaction with plasmas and matters (Wilks *et al.*, 1992; Brandl *et al.*, 2009; Arefiev *et al.*, 2012; Kemp and Divol, 2012; Breuer and Hommelhoff, 2013; Chen *et al.*, 2013; Liu *et al.*, 2013; Braenzel *et al.*, 2017; Cheng *et al.*, 2017). The most important question is how we can transfer the enormous fraction of laser energy into the generation of relativistic electrons. In the laser interaction with underdense plasma, the electron acceleration mechanism strongly depends on the laser pulse duration. In the wakefield acceleration, the laser pulse duration should be shorter or comparable to the plasma wave period (Esarey *et al.*, 2009), and this mechanism becomes less efficient by long laser pulses (Malka *et al.*, 2001; Mangles *et al.*, 2005).

For generation more energetic electrons by making use of a laser beam, different additional concept have been used, such as fast electron beams generation from plasma mirrors (Bocoum *et al.*, 2016), vacuum laser acceleration of relativistic electron (MeLikian, 2014; Thevenet *et al.*, 2016; Zhang *et al.*, 2017; Pandari *et al.*, 2019), relativistic electron mirrors from nanoscale foils (Kiefer *et al.*, 2013), boosted high harmonic pulse from a double-sided relativistic mirror (Esirkepov *et al.*, 2009), and electron acceleration by relativistic surface plasmons (Fedeli *et al.*, 2016).

We can avoid deceleration when the light field is reflected out of the electron trajectory (Singh, 2004; Mulser *et al.*, 2008; Wu *et al.*, 2010) and or by electron injection into the light field with a specific initial momentum (Dodin and Fisch, 2003; Andreev and Platonov, 2013; Bocoum *et al.*, 2016; Thevenet *et al.*, 2016). Moreover, the electron acceleration has already been enhanced by making use of an additional external electric field (Plettner *et al.*, 2005; Dimant and Oppenheim, 2010; He *et al.*, 2013; Cheng *et al.*, 2016).

It is well known that the non-linear plasma properties are intensified by magnetic field due to the Lorentz force, so it is reasonable the plasma dynamics affected by magnetic field. Therefore, generation of magnetic fields is one of the more interesting debatable topics among the scientists in laser–plasma interaction. This issue has a wide range of applications in fast ignition scheme for inertial confinement fusion (Tabak *et al.*, 1994), laboratory astrophysics (Ripin *et al.*, 1990), and particle acceleration (Schmitz and Kull, 2002; Yu *et al.*, 2003; Liu *et al.*, 2004; Qiao *et al.*, 2005). The experiments have been reported the magnetic field strength in the order of about tens of mega-gauss (MG) (Fuchs *et al.*, 1998; Najmudin *et al.*, 2001); however, the recent 3D particle-in-cell (PIC) simulation and analytical investigations predicted generation of the super strong magnetic field up to 100 MG in laser interaction with an overdense plasma channel (Pukhov and Meyer-ter-Vehn, 1996; Qiao *et al.*, 2006; Wang *et al.*, 2011; Stark *et al.*, 2016).

© Cambridge University Press 2019

CAMBRIDGE UNIVERSITY PRESS

The ponderomotive force plays an important role in a long laser pulse interaction with plasma. In this case, the ponderomotive pressure repels the electrons away from the axis of the beam in the transverse direction, while the heavy ions remain at rest. As a result, an ion channel is created with a quasi-static transverse electric field due to the charge separation. The electron acceleration in an ion channel has been experimentally demonstrated in the past decades (Gahn *et al.*, 1999, 2002; Kitagawa *et al.*, 2004; Walton *et al.*, 2006; Kneip *et al.*, 2008, 2009); however, the test electron acceleration by strong laser field in an ion channel has already been studied theoretically by Arefiev *et al.* (2014).

In this paper, we will follow the method is used by Arefiev *et al.*, by taking into account the fact that the plasma channel should be magnetized (Pukhov and Meyer-ter-Vehn, 1996; Stark *et al.*, 2016). Therefore, we would like to examine the electron direct laser acceleration (DLA) in a two-dimensional magnetized plasma channel. We will focus our attention on the role played by the self-generated quasi-static magnetic field on the electron acceleration process.

The paper is organized in the following fashion. In the "Model description" section, a single electron model is formulated for an electron irradiated by an incoming laser pulse in a steady-state two-dimensional transversely magnetized ion channel. The purpose of this section is to justify the necessity of including the self-generated magnetic field during the plasma channel creation, to explain how such a plasma channel can be modeled. We try to explain the effect of the self-generated magnetic field on the dephasing rate in this section. In the "Numerical results and discussion" section, the closed set of electron motion of equations in the presence of the laser pulse and self-generated magnetic field is presented and solved with proper initial condition. Finally, the summary and conclusion are given in the last section.

Model description

In this section, we describe a single electron dynamics irradiated by a laser field in a homogenous transversely magnetized plasma channel. The origin of the self-generated magnetic field is the current induced in the channel. The PIC simulations show that the quasi-static magnetic field increases proportional at distance from axis and modulates in the direction of laser field propagation with spatial variation in the order of the laser wavelength λ (Stark *et al.*, 2016; Pukhov and Meyer-ter-Vehn, 1996).

It is well known in the interaction of high intensity laser field with a plasma, since the longitudinal momentum dominates the kinetic energy of electrons, we can assume that all of the electrons moving forward with relativistic velocity. However, a few number of electrons move under the angle with respect to the channel axis; therefore, if we assume the radius of channel is R, it is easy to show that the magnetic field for $y < R$ is given by $B_x = 2\pi J_z y/c$, where $J_z = en_e c$. On the other hand, if we include the magnetic field modulation in z direction, we can present the quasi-static magnetic field in the form as follows:

$$B_x/B_0 = (n_e/2n_c)(2\pi \mid y \mid /\lambda)\sin 2\pi z/\lambda, \tag{1}$$

where $B_0 = mc\omega/e$ and $n_c = \pi mc^2/(e\lambda)^2$ are the magnetic field of the laser field and critical density, respectively. Figure 1 depicts a two-dimensional spatial setup, with a Cartesian system of coordinates (x, y, z). The figure shows the channel is a slab of heavy immobile ions with density n_0. The static electric field $E_c = \mathbf{e}_y \omega_{pe}^2 m_e y/|e|$ induced due to the charge separation, where $\mathbf{e}_y$ is a unit vector in y direction and $\omega_{pe} = (4\pi n_0 e^2/m_e)^{1/2}$ refers to the plasma frequency (m_e and e are electron mass and charge). The laser field is linearly polarized by polarization angle θ and propagates in the z direction. In addition, the self-generated magnetic field is perpendicular to y–z plane.

It is worthwhile to note that the plasma channel generally is cylindrical; however, because of azimuthal symmetry and with no loss of generality, we can use a two-dimensional (2D) plasma channel. On the other hand, the 2D plasma channel has some advantages: first for simplicity and the second is that for a given laser amplitude, the amplitude of laser field that responsible to direct acceleration of betatron oscillation is changed by changing the polarization angle. Therefore, we can examine the interplay between the oscillating electric field and the electrostatic field of the channel.

At this stage, we consider a single electron dynamics placed in an ion channel (Fig. 1) irradiated by a laser field. We assume that the laser field has the same parameters with the laser which produced the channel. We introduced the laser field by normalized vector potential as follows:

$$\mathbf{a}(z, t) = a(\xi)[\mathbf{e}_x \cos\theta + \mathbf{e}_y \sin\theta], \tag{2}$$

where $\xi = \omega(t - z/c)$ is the wave phase in which ω is wave frequency and c is the speed of light. Also, t is the time in the channel frame of reference, and $\mathbf{e}_x$ is a unit vector in x direction. We consider pulse amplitude in the form of $a(\xi) = a_*(\xi)\sin\xi$, where $0 \le a_*(\xi) \le a_0$ is a given slowly varying envelope. The total electric and magnetic fields inside the plasma channel are as follows:

$$\mathbf{E} = \mathbf{E}_c + \mathbf{E}_L, \quad \mathbf{B} = \mathbf{B}_L + \mathbf{B}_x, \tag{3}$$

where $\mathbf{B}_x$ is the quasi-static self-generated magnetic field directed along the x-axis, and the electric and magnetic fields corresponding to the laser field can be expressed as follows:

$$\mathbf{E}_L = -\frac{m_e c}{|e|}\frac{\partial \mathbf{a}}{\partial t}, \quad \mathbf{B}_L = \frac{m_e c^2}{|e|}\nabla \times \mathbf{a}. \tag{4}$$

The equation of motion for an electron inside the plasma channel is given by the following equation:

$$\frac{d\mathbf{p}}{dt} = \frac{-|e|\mathbf{E}}{m_e c} - \frac{|e|}{\gamma m_e c}\mathbf{p} \times \mathbf{B}, \tag{5}$$

where $\mathbf{p}$ is the dimensionless electron momentum normalized to $m_e c$, and $\gamma = \sqrt{1 + \mathbf{p}^2}$ is the relativistic factor. By making use of Eqs (1)–(5), we obtain the closed set of equations to describe the electron dynamics in the plasma channel as follows:

$$\begin{aligned}
&\frac{d}{d\tau}[p_y - a\sin\theta] = -\gamma\frac{\omega_{pe}^2}{\omega^2}Y_B - \frac{\omega_{ce}}{\omega}p_z,\\
&\frac{dp_z}{d\tau} = p_x\left(\cos\theta\frac{\partial a}{\partial \xi}\right) + p_y\left(\sin\theta\frac{\partial a}{\partial \xi} + \frac{\omega_{ce}}{\omega}\right),\\
&\frac{d\xi}{d\tau} = \gamma - p_z,\\
&\frac{dY_B}{d\tau} = p_y,\\
&p_x = a\cos\theta,
\end{aligned} \tag{6}$$

https://doi.org/10.1017/S0263034619000703 Published online by Cambridge University Press




Fig. 1. Schematic setup of a single electron model in an ion channel.

where τ is a dimensionless proper time defined by the relation $d\tau/dt = \omega/\gamma$, and $Y_B = \omega y/c$ is dimensionless displacement across the channel. Moreover, $\omega_{ce} = eB_x/m_e c$ and $\omega_{pe} = (4\pi n_e^2/m_e)^{1/2}$ are electron cyclotron and plasma frequencies, where n_e shows the electron density. The third relation in Eq. (6) represents a dephasing rate and usually defines as $R_B = \gamma - p_z$. Using Eq. (6), we can show that

$$\frac{d}{d\tau}\left(\gamma - p_z + \frac{1}{2}\frac{\omega_{pe}^2}{\omega^2}Y_B^2 + Y_B\frac{\omega_{ce}}{\omega}\right) = 0. \tag{7}$$

Integrating Eq. (7), we get

$$R_B = \gamma - p_z = I_B - \frac{1}{2}\Omega_B^2 Y_B^2 - \Omega_{ce} Y_B, \tag{8}$$

where I_B is a constant and determined by initial conditions, and $\Omega_B = \omega_{pe}/\omega$ and $\Omega_{ce} = \omega_{ce}/\omega$ are normalized plasma and cyclotron frequencies, respectively. For example, if we assume the electron is initially at rest (i.e., $R \to 1$) and is placed on Y_{0B}, we can find I_B as follows:

$$I_B = 1 + \frac{1}{2}\Omega_B^2 Y_{0B}^2 + \Omega'_{ce} Y_{0B}, \tag{9}$$

where Ω'_{ce} refers to normalized cyclotron frequency at $Y = Y_{0B}$. It should be noted that the last term on the right-hand side of Eq. (8) shows the effect of the self-generated magnetic field on the dephasing rate. It is well known that the axial part of the electron momentum larger than the transverse part at ultra-relativistic wave amplitude ($a \gg 1$); therefore, we expect that $R_B \to 0$ when the electron efficiently accelerated by the laser field.

In order to determine the electron maximum displacement across to the channel, the dephasing rate should becomes small vanishingly in Eq. (8). Therefore, substituting $R_B = 0$ in Eq. (8), we get

$$Y_{B*} = \frac{-\Omega_{ce} \pm \sqrt{\Omega_{ce}^2 + 2I_B\Omega_B^2}}{\Omega_B^2}. \tag{10}$$

Equation (10) is in complete agreement with the previous knowledge (Arefiev *et al.*, 2014) in the limit of $\Omega_{ce} = 0$, for a non-magnetized plasma channel.

Numerical results and discussion

In this section, we investigate the electron dynamics in the plasma channel for a laser irradiated with polarization angle θ in the presence of the self-generated static magnetic field in the form of Eq. (1).

The electron dynamics is studied by the numerical solution of the coupled relations in Eq. (6) for a Gaussian profile pulse laser $a = a_*(\xi)\sin(\xi)$ is defined as follows:

$$a_*(\xi) = a_0 \exp\left[-\frac{(\xi - \xi_0)^2}{2\sigma^2}\right], \tag{11}$$

where $a_0 = 10$, $\xi_0 = 400$, and $\sigma = 100$. Here, a_0 is the normalized laser field amplitude, and parameters ξ_0 and σ characterize the laser beam initial position with respect to the electron and the beam duration. In the course of this paper, we consider the initial condition for electron at $\tau = 0$, with $p_y(0) = 0$, $p_z(0) = 0$, $Y_0 = 0.05$, and $\xi = 0$.

Figure 2 shows the variation of $\gamma_{max}/\gamma_{vac}$ (where γ_{max} is the maximum γ-factor and $\gamma_{vac} = 1 + a_0^2/2$) as a function of normalized plasma frequency in the magnetized and non-magnetized plasma channels. Figure 2a and 2b are plotted for polarization angles $\theta = \pi/2$ and $\theta = 0$, respectively. In addition, Figure 2c and 2d indicate the maximum strength of magnetic field would feel by the test electron for polarization angles $\theta = \pi/2$ and $\theta = 0$. The figure shows the strength of self-generated magnetic field rises as the plasma density increases. We find that the role of magnetic field is more effective in the enhancement of electron acceleration for polarization angle $\theta = \pi/2$.

Actually, the electron acceleration is affected by plasma density, laser field amplitude and polarization angle, and self-generated magnetic field strength. The previous investigation have been revealed the efficient electron acceleration by low density plasma channels and also the maximum enhancement when laser electric field and the field of the channel are collinear (Arefiev *et al.*, 2014). Figure 2 shows when the effect of self-generated magnetic field is included not only the maximum γ-factor increases but also the electron acceleration enhanced for a relatively high-density plasma channel.

The electron has two oscillating motions: (a) oscillation with laser field frequency in polarization plane and (b) betatron oscillating motion around magnetic field with frequency ω_c. Two oscillatory motions are in y–z plane for polarization angle $\theta = \pi/2$. Therefore, we expect the resonance takes place when the electron betatron oscillation closes to the laser field frequency. For example, the resonant absorption for polarization angle $\theta = \pi/2$ is clearly shown in Figure 2a and 2c for $\omega_{pe}/\omega = 0.356$ when the

https://doi.org/10.1017/S0263034619000703 Published online by Cambridge University Press




Fig. 2. Maximum normalized γ-factor as a function of ω_{pe}/ω for a magnetized (red line) and non-magnetized plasma channel. (a) Polarization angle $\theta = \pi/2$ and (b) polarization angle $\theta = 0$. (c,d) Indicate the normalized maximum self-generated magnetic field variation as a function of ω_{pe}/ω for polarization angles $\theta = \pi/2$ and $\theta = 0$, respectively.

$(B_x)_{max} \approx 0.9B_0$. This process is governed mechanism for the enhancement of electron acceleration for polarization angle in the vicinity of $\theta \approx \pi/2$.

Figure 3a and 3b demonstrate the normalized γ-factor variation as a function of $\xi/2\pi$ for polarization angle $\theta = \pi/2$ in the magnetized and non-magnetized plasma channels, respectively. The figure compares the electron acceleration in the presence of the plasma channel and vacuum. Thus, even for $\gamma < \gamma_{vac}$, the electron is accelerated ($\gamma_{vac} = 51$). It is understood from Figure 3a that γ




Fig. 3. Relative γ-factor at $\theta = \pi/2$ and $\omega_p/\omega = 0.353$ (a) for a magnetized and (b) for a non-magnetized plasma channel as a function of $\xi/2\pi$.

$> \gamma_{vac}$ for wide range of $\xi/2\pi$, means that electron acceleration is enhanced in the presence of the magnetized plasma channel. This is not true for a non-magnetized plasma channel in Figure 3b. The maximum γ-factor for electron driven by laser beam is around $\gamma_{max}/\gamma_{vac} \approx 5.7$ which is in good agreement with Figure 2a.

In order to determine simultaneous effects of the polarization angle and channel density, we plotted the variation of maximum γ-factor with respect to the θ and ω_{pe}/ω in Figures 4 and 5. In Fig. 4, the influence of self-generated magnetic field is included, while Fig. 5 are plotted for a non-magnetized plasma channel. It seems for a magnetized plasma channel and for polarization angles $0.39\pi \leq \theta \leq 0.5\pi$ electron acceleration enhanced up to $\gamma_{max} \approx 12\gamma_{vac}$ and the required threshold density for electron acceleration decreases to $n_e = 0.0036n_c$. While for non-magnetized plasma, the maximum acceleration for polarization angles $0.32\pi \leq \theta \leq 0.5\pi$ is around $\gamma_{max} \approx 8\gamma_{vac}$ for threshold density $n_e = 0.0064n_c$. Therefore, for polarization angles near to $\theta \approx \pi/2$, resonant absorption plays an important role in acceleration enhancement. As we mentioned, the enhancement occurs when the frequency of betatron oscillation closes to the laser field frequency. For a magnetized plasma channel, the resonance condition is satisfied for different plasma densities at different polarization angles in the range $0.39\pi \leq \theta \leq 0.5\pi$. Figure 4 indicates that the electron acceleration enhanced in magnetized plasma specially for the relatively dense plasma channel.

Summary and conclusion

In summary, we have analyzed the dynamics of an electron irradiated by a linearly polarized laser pulse in a two-dimensional

https://doi.org/10.1017/S0263034619000703 Published online by Cambridge University Press




Fig. 4. Variation of maximum γ-factor as a function of ω_{pe}/ω and polarization angle θ for a magnetized plasma channel.




Fig. 5. Variation of maximum γ-factor as a function of ω_{pe}/ω and polarization angle θ for a non-magnetized plasma channel.

transversely magnetized ion channel. Taking into account that a real plasma channel is magnetized, we expect magnetic field makes affect the test electron acceleration through plasma channel. Therefore, the outcomes of the present paper help to the reader to study a real physical system. We supposed that the magnetic field is generated due the current induced inside the channel and presented a simple model for a magnetized plasma channel. The results revealed that the self-generated magnetic field sustains the electron acceleration for polarization angle near to $\theta = \pi/2$. For small polarization angle, the effect of magnetic field was no perceptible. It is shown that for polarization angle near to $\theta \approx \pi/2$, when the electron betatron oscillation frequency closes to the laser field frequency, the resonance condition is achieved and electron acceleration enhanced.

References

Andreev AA and Platonov KY (2013) Double relativistic electron-accelerating mirror. *Quantum Electronics* **43**, 435.

Arefiev AV, Breizman BN, Schollmeier M and Khudik VN (2012) Parametric amplification of laser-driven electron acceleration in underdense plasma. *Physical Review Letters* **108**, 145004.

Arefiev AV, Khudik VN and Schollmeier M (2014) Enhancement of laser-driven electron acceleration in an ion channel. *Physics of Plasmas* **21**, 033104.

Bocoum M, Thévenet M, Böhle F, Beaurepaire B, Vernier A, Jullien A, Faure J and Lopez-Martens R (2016) Anticorrelated emission of high harmonics and fast electron beams from plasma mirrors. *Physical Review Letters* **116**, 185001.

Braenzel J, Andreev AA, Abicht F, Ehrentraut L, Platonov K and Schnürer M (2017) Amplification of relativistic electron bunches by acceleration in laser fields. *Physical Review Letters* **118**, 014801.

Brandl F, Hidding B, Osterholz J, Hemmers D, Karmakar A, Pukhov A and Pretzler G (2009) Directed acceleration of electrons from a solid surface by sub-10-fs laser pulses. *Physical Review Letters* **102**, 195001.

Breuer J and Hommelhoff P (2013) Laser-based acceleration of nonrelativistic electrons at a dielectric structure. *Physical Review Letters* **111**, 134803.

Chen H, Nakai M, Sentoku Y, Arikawa Y, Azechi H, Fujioka S, Keane C, Kojima S, Goldstein W, Maddox BR, Miyanaga N, Morita T, Nagai T, Nishimura H, Ozaki T, Park J, Sakawa Y, Takabe H, Williams G and

https://doi.org/10.1017/S0263034619000703 Published online by Cambridge University Press

Zhang Z (2013) New insights into the laser produced electron-positron pairs. *New Journal of Physics* **15**, 065010.

Cheng LH, Xue JK and Liu J (2016) Laser-driven electron acceleration in a plasma channel with an additional electric field. *Physics of Plasmas* **23**, 053102.

Cheng LH, Yao ZW, Zhang XB and Xue JK (2017) Net electron energy gain induced by superluminal phase velocity and subluminal group velocity of a laser in a plasma channel. *Physics of Plasmas* **24**, 082114.

Cipiccia S, Islam MR, Ersfeld B, Shanks RP, Brunetti E, Vieux G, Yang X, Issac RC, Wiggins SM, Welsh GH, Anania M-P, Maneuski D, Montgomery R, Smith G, Hoek M, Hamilton DJ, Lemos NRC, Symes D, Rajeev PP, Shea VO, Dias JM and Jaroszynski DA (2011) Gamma-rays from harmonically resonant betatron oscillations in a plasma wake. *Nature Physics* **7**, 867.

Dimant YS and Oppenheim MM (2010) Interaction of plasma cloud with external electric field in lower ionosphere. *Annales Geophysicae* **28**, 719.

Dodin IY and Fisch NJ (2003) Relativistic electron acceleration in focused laser fields after above-threshold ionization. *Physical Review E* **68**, 056402.

Esarey E, Schroeder CB and Leemans WP (2009) Physics of laser-driven plasma-based electron accelerators. *Reviews of Modern Physics* **81**, 1229.

Esirkepov TZ, Bulanov SV, Kando M, Pirozhkov AS and Zhidkov AG (2009) Boosted high-harmonics pulse from a double-sided relativistic mirror. *Physical Review Letters* **103**, 025002.

Fedeli L, Sgattoni A, Cantono G, Garzella D, Réau F, Prencipe I, Passoni M, Raynaud M, Květoň M, Proska J, Macchi A and Ceccotti T (2016) Electron acceleration by relativistic surface plasmons in laser-grating interaction. *Physical Review Letters* **116**, 015001.

Flippo K, Bartal T, Beg F, Chawla S, Cobble J, Gaillard S, Hey D, MacKinnon A, Macphee A, Nilson P, Offermann D, Le Pape S and Schmitt MJ (2010) Omega EP, laser scalings and the 60 MeV barrier: first observations of ion acceleration performance in the 10 picosecond kilojoule short-pulse regime. *Journal of Physics. Conference Series* **244**, 022033.

Fuchs J, Malka G, Adam JC, Amiranoff F, Baton SD, Blanchot N, Héron A, Laval G, Miquel JL, Mora P, Pépin H and Rousseaux C (1998) Dynamics of subpicosecond relativistic laser pulse self-channeling in an underdense preformed plasma. *Physical Review Letters* **80**, 1658.

Fuchs J, Antici P, d'Humieres E, Lefebvre E, Borghesi M, Brambrink E, Cecchetti CA, Kaluza M, Malka V, Manclossi M, Meyroneinc S, Mora P, Schreiber J, Toncian T, Pepin H and Audebert P (2006) Laser-driven proton scaling laws and new paths towards energy increase. *Nature Physics* **2**, 48.

Gahn C, Tsakiris GD, Pukhov A, Meyer-ter-Vehn J, Pretzler G, Thirolf P, Habs D and Witte KJ (1999) Multi-MeV electron beam generation by direct laser acceleration in high-density plasma channels. *Physical Review Letters* **83**, 4772.

Gahn C, Tsakiris GD, Pretzler G, Witte KJ, Thirolf P, Habs D, Delfin C and Wahlström CG (2002) Generation of MeV electrons and positrons with femtosecond pulses from a table-top laser system. *Physics of Plasmas* **9**, 987.

He ZH, Hou B, Nees JA, Easter JH, Faure J, Krushelnick K and Thomas AGR (2013) High repetition-rate wakefield electron source generated by few-millijoule, 30 fs laser pulses on a density downramp. *New Journal of Physics* **15**, 053016.

Kemp AJ and Divol L (2012) Interaction physics of multipicosecond petawatt laser pulses with overdense plasma. *Physical Review Letters* **109**, 195005.

Kiefer D, Yeung M, Dzelzainis T, Foster PS, Rykovanov SG, Lewis CL, Marjoribanks RS, Ruhl H, Habs D, Schreiber J, Zepf M and Dromey B (2013) Relativistic electron mirrors from nanoscale foils for coherent frequency upshift to the extreme ultraviolet. *Nature Communications* **4**, 1763.

Kitagawa Y, Sentoku Y, Akamatsu S, Sakamoto W, Kodama R, Tanaka KA, Azumi K, Norimatsu T, Matsuoka T, Fujita H and Yoshida H (2004) Electron acceleration in an ultraintense-laser-illuminated capillary. *Physical Review Letters* **92**, 205002.

Kneip S, Nagel SR, Bellei C, Bourgeois N, Dangor AE, Gopal A, Heathcote R, Mangles SPD, Marquès JR, Maksimchuk A, Nilson PM, Ta Phuoc K, Reed S, Tzoufras M, Tsung FS, Willingale L, Mori WB, Rousse A, Krushelnick K and Najmudin Z (2008) Observation of synchrotron radiation from electrons accelerated in a petawatt-laser-generated plasma cavity. *Physical Review Letters* **100**, 105006.

Kneip S, McGuffey C, Nagel SR, Palmer C, Bellei C, Schreiber J, Huntington C, Dollar F, Matsuoka T, Chvykov V, Kalintchenko G, Yanovsky V, Maksimchuk A, Ta Phuoc K, Mangles SPD, Krushelnick K and Najmudin Z (2009) Comparative study of betatron radiation from laser-wakefield and direct-laser accelerated bunches of relativistic electrons. In Jaroszynski DA and Rousse A (eds), *Proceedings of SPIE on Harnessing Relativistic Plasma Waves as Novel Radiation Sources from Terahertz to X-Rays and Beyond*, Vol. 7359, 73590T (Published on 6 May 2009).

Liu H, He XT and Chen SG (2004) Resonance acceleration of electrons in combined strong magnetic fields and intense laser fields. *Physical Review E* **66**, 066409.

Liu B, Wang HY, Liu J, Fu LB, Xu YJ, Yan XQ and He XT (2013) Generating overcritical dense relativistic electron beams via self-matching resonance acceleration. *Physical Review Letters* **110**, 045002.

Malka V, Faure J, Marques JR, Amiranoff F, Rousseau JP, Ranc S, Chambaret JP, Najmudin Z, Walton B, Mora P and Solodov A (2001) Characterization of electron beams produced by ultrashort (30 fs) laser pulses. *Physics of Plasmas* **8**, 2605.

Mangles SPD, Walton BR, Tzoufras M, Najmudin Z, Clarke RJ, Dangor AE, Evans RG, Fritzler S, Gopal A, Hernandez-Gomez C, Mori WB, Rozmus W, Tatarakis M, Thomas AGR, Tsung FSTsung, Wei MS and Krushelnick K (2005) Electron acceleration in cavitated channels formed by a petawatt laser in low-density plasma. *Physical Review Letters* **24**, 245001.

Melikian R (2014) Acceleration of electrons by high intensity laser radiation in a magnetic field. *Laser and Particle Beams* **2**, 205.

Mulser P, Bauer D and Ruhl H (2008) Collisionless laser-energy conversion by anharmonic resonance. *Physical Review Letters* **101**, 225002.

Najmudin Z, Tatarakis M, Pukhov A, Clark EL, Clarke RJ, Dangor AE, Faure J, Malka V, Neely D, Santala MIK and Krushelnick K (2001) Measurements of the inverse Faraday effect from relativistic laser interactions with an underdense plasma. *Physical Review Letters* **87**, 215004.

Pandari MR, Jahangiri F and Niknam AR (2019) Optimizing the electron acceleration in vacuum by chirped ultrashort laser pulse using particle swarm method. *Laser and Particle Beams* **37**(3), 242–251.

Plettner T, Byer RL, Colby E, Cowan B, Sears CMS, Spencer JE and Siemann RH (2005) Visible-laser acceleration of relativistic electrons in a semi-infinite vacuum. *Physical Review Letters* **95**, 134801.

Pukhov A and Meyer-ter-Vehn J (1996) Relativistic magnetic self-channeling of light in near-critical plasma: three-dimensional particle-in-cell simulation. *Physical Review Letters* **76**, 3975.

Qiao B, He XT, Zhu SP and Zheng CY (2005) Electron acceleration in combined intense laser fields and self-consistent quasistatic fields in plasma. *Physics of Plasmas* **12**, 083102.

Qiao B, He XT and Zhu SP (2006) Fluid theory for quasistatic magnetic field generation in intense laser plasma interaction. *Physics of Plasmas* **13**, 053106.

Ripin BH, Manka CK, Peyser TA, McLean EA, Stamper JA, Mostovych AN, Grun J, Kearney K, Crawford JR and Huba JD (1990) Laboratory laser-produced astrophysical-like plasmas. *Laser and Particle Beams* **8**, 183.

Schmitz M and Kull HJ (2002) Single-electron model of direct laser acceleration in plasma channels. *Laser Physics* **12**, 443.

Singh KP (2004) Electron acceleration by a circularly polarized laser pulse in a plasma. *Physics of Plasmas* **11**, 3992.

Stark DJ, Toncian T and Arefiev AV (2016) Enhanced multi-MeV photon emission by a laser-driven electron beam in a self-generated magnetic field. *Physical Review Letters* **116**, 185003.

Tabak M, Hammer J, Glinsky ME, Kruer WL, Wilks SC, Woodworth J, Campbell EM, Perry MD and Mason RJ (1994) Ignition and high gain with ultrapowerful lasers. *Physics of Plasmas* **1**, 1626.

Thévenet M, Leblanc A, Kahaly S, Vincenti H, Vernier A, Quéré F and Faure J (2016) Vacuum laser acceleration of relativistic electrons using plasma mirror injectors. *Nature Physics* **12**, 355.

Walton BR, Mangles SPD, Najmudin Z, Tatarakis M, Wei MS, Gopal A, Marle C, Dangor AE, Krushelnick K, Fritzler S and Malka V (2006)

https://doi.org/10.1017/S0263034619000703 Published online by Cambridge University Press

Measurements of forward scattered laser radiation from intense sub-ps laser interactions with underdense plasmas. *Physics of Plasmas* **13**, 113103.

Wang HY, Lin C, Sheng ZM, Liu B, Zhao S, Guo ZY, Lu YR, He XT, Chen JE and Yan XQ (2011) Laser shaping of a relativistic intense, short Gaussian pulse by a plasma lens. *Physical Review Letters* **107**, 265002.

Wang X, Zgadzaj R, Fazel N, Li Z, Yi SA, Zhang X, Henderson W, Chang Y-Y, Korzekwa R, Tsai H-E, Pai C-H, Quevedo H, Dyer G, Gaul E, Martinez M, Bernstein AC, Borger T, Spinks M, Donovan M, Khudik V, Shvets G, Ditmire T and Downer MC (2013) Quasi-monoenergetic laser-plasma acceleration of electrons to 2 GeV. *Nature Communications* **4**, 1988.

Wilks SC, Kruer WL, Tabak M and Langdon AB (1992) Absorption of ultra-intense laser pulses. *Physical Review Letters* **69**, 1383.

Wu HC, Meyer-ter-Vehn J, Fernández J and Hegelich BM (2010) Uniform laser-driven relativistic electron layer for coherent Thomson scattering. *Physical Review Letters* **104**, 234801.

Yu MY, Yu W, Chen ZY, Zhang J, Yin Y, Cao LH, Lu PX and Xu ZZ (2003) Electron acceleration by an intense short-pulse laser in underdense plasma. *Physics of Plasmas* **10**, 2468.

Zhang Y, Jiao J-L, Zhang B, Zhang Z-M and Gu Y-Q (2017) GV/cm scale laser-magnetic resonant acceleration in vacuum. *Laser and Particle Beams* **3**, 520.

https://doi.org/10.1017/S0263034619000703 Published online by Cambridge University Press