Environmental Conservation **37** (2): 169–176 © Foundation for Environmental Conservation 2010 doi:10.1017/S0376892910000214

Land-use changes as a critical factor for long-term wild rabbit conservation in the Iberian Peninsula

MIGUEL DELIBES-MATEOS[1,2]*, MIGUEL ÁNGEL FARFÁN[1,3], JESÚS OLIVERO[1] AND JUAN MARIO VARGAS[1]

[1]*Departamento de Biología Animal, Facultad de Ciencias, Universidad de Málaga, 29071 Málaga, Spain* [2]*Instituto de Investigación en Recursos Cinegéticos, IREC (CSIC-UCLM-JCCM), Ronda de Toledo s/n, 13071 Ciudad Real, Spain and* [3]*Biogea Consultores, C/Navarro Ledesma, 243, 29010 Málaga, Spain*

Date submitted: 17 September 2009; Date accepted: 27 January 2010; First published online: 12 August 2010

SUMMARY

European rabbits (*Oryctolagus cuniculus*), a multifunctional keystone species in the Iberian Peninsula, have drastically declined over past decades. Rabbit decline has been frequently attributed to the arrival of two viral diseases. However, decline was apparently ongoing before the arrival of the diseases, as a consequence of habitat loss and fragmentation. In this paper, the effect on rabbit populations of land-use changes during recent decades in Andalusia (southern Spain) is analysed. Areas favourable for rabbits both at present and during the 1960s are identified, and the environmental and land-use factors that determine these areas established. In areas where the favourability for rabbits has changed during recent decades, main land use changes are assessed to identify possible factors explaining rabbit favourability in these areas. Areas favourable to rabbits are currently determined by factors similar to those during the 1960s; these areas have undergone geographic changes in recent decades, apparently as a consequence of land-use changes in Andalusia. The percentages of the variables that were positively associated with rabbit favourability in both models (current and 1960s) have declined in Andalusia as a whole, and in areas where rabbit favourability has decreased; hence environments suitable for rabbits have become impoverished. Conversely, in both models, environments suitable for rabbits increased in municipalities, where rabbit favourability also increased. The preservation of rabbit-friendly habitats should be a priority for the conservation of this key species in the western Mediterranean.

Keywords: agricultural intensification, favourability function, habitat loss, land abandonment, *Oryctolagus cuniculus*

*Correspondence: Dr Miguel Delibes-Mateos Tel: +34 926295450 Fax: +34 926295451 e-mail: mdelibesmateos@gmail.com

INTRODUCTION

European wild rabbits (*Oryctolagus cuniculus*), are considered a multifunctional keystone species in the Iberian Peninsula (Delibes-Mateos *et al.* 2007). Rabbits conspicuously alter landscapes and provide foraging, shelter and nesting habitats for a diverse array of species, their grazing activities alter plant species composition and vegetation structure, especially by creating open areas and preserving plant species diversity, and they are prey for a large number of predators (Delibes-Mateos *et al.* 2008*a*; Gálvez-Bravo *et al.* 2009).

Rabbits have massively declined in the Iberian Peninsula since the first half of the 20th century, and this is frequently attributed to the arrival of the viral diseases myxomatosis from 1953, and rabbit haemorrhagic disease (RHD) from 1988. However, rabbit decline was already ongoing in the first half of the 20th century, apparently as a consequence of habitat loss and fragmentation (Ward 2005). Rabbits populations declined in Spain by 73% between 1973 and 1993 and this decline was by no means restricted to the period 1988–1993 (Virgós *et al.* 2007), when RHD was becoming widely established. Other factors such as habitat loss are evidently also involved in the decline of rabbits. The decrease in rabbit numbers may have had important cascading effects on the functioning of the Iberian Mediterranean ecosystem, with serious ecological and economic consequences (Delibes-Mateos *et al.* 2008*a*).

Researchers have therefore made great efforts to study the main causes of rabbit population decline in the Iberian Peninsula; the number of research papers focused on this species has significantly increased in the Iberian Peninsula following the population decline (Piorno 2006). Most of this research effort has focused on diseases. However, and although many studies have addressed the relationships between habitat characteristics and the distribution and abundance of rabbit populations in the Iberian Peninsula (for example Calvete *et al.* 2004; Fernández 2005), information on the effects of habitat loss on Iberian rabbit populations is scarce. In general, nearly all studies have focused on the association of recent rabbit abundances and/or trends with present landscape features (for example Calvete *et al.* 2006), whereas there are no studies on this relationship several decades ago. In fact, past information on rabbit abundance has never been used in previous studies. Therefore, it is very difficult to know how landscape changes have affected rabbit

https://doi.org/10.1017/S0376892910000214 Published online by Cambridge University Press



Figure 1 Study area. The main mountain ranges (Sierra Morena and the Baetic System, sub-divided into two ranges, Sub-baetic and Penibetic) and the most important valley (Guadalquivir valley) are shown in schematic form. Limits between provinces are also indicated.

populations, and how such changes have contributed to rabbit decline in the Iberian Peninsula.

In Andalusia (southern Spain; Fig. 1), where rabbits have traditionally reached high densities (Villafuerte *et al.* 1998), land use has changed substantially over recent decades (Fernández-Alés *et al.* 1992). Economic growth and the rural exodus have led to the intensification of agriculture and livestock farming in certain areas, and to the under-use of other vast rural areas (Fernández-Alés *et al.* 1992). Both processes have destroyed large areas of traditional agricultural landscapes, where rabbits usually reach their highest densities (Calvete *et al.* 2004). We suspect that the drastic changes in land use (Fernández-Alés *et al.* 1992) must have affected the rabbit distribution and abundance in Andalusia.

In this paper, our main aims were to identify areas in Andalusia that are favourable for rabbits at present and during the 1960s, and to establish the environmental and land use factors that determine these areas in both periods. We define favourable areas as those where the environment increases the probability of presence of rabbits, independently of the proportion of area they occupy ('prevalence'; see Real *et al.* 2006). Using this approach, we evaluated whether rabbit habitat requirements may have changed in recent decades. Alternatively, we identified areas where the favourability for rabbits has markedly changed (either increased or decreased) during recent decades, and evaluated the evolution of the main land use that explains rabbit favourability in these areas, as well as in the whole study area. We also discuss how land-use changes have affected rabbit distribution and abundance in recent decades in the western Mediterranean Basin.

METHODS

Study area

Andalusia covers more than 87 000 km^2 in the southernmost part of mainland Spain, and is administratively divided into 771 municipalities. The main mountain ranges are the Sierra Morena, along the northern fringe of the region, and the Baetic System, sub-divided into two ranges, Sub-baetic and Penibaetic, which are oriented NE–SW and mainly occupy the eastern part of the region. The most important plain is the Guadalquivir valley, which is longitudinally oriented between the Sierra Morena and the Baetic System (Fig. 1). Andalusia has a Mediterranean climate, with mild winters and severe summer droughts (see Delibes-Mateos *et al.* 2009*a* for a wider description of the study area).

Variables

Andalusian municipalities were assigned to groups according to whether rabbits were relatively abundant or not at present and during the 1960s, using information from current hunting yields (1993–2001) and from game species abundance maps available from the Mainland Spanish Fish, Game and National Parks Service, respectively.

We analysed 32 134 annual hunting reports (AHRs) from the period 1993–2001 reported by 6049 game estates to estimate the average hunting yields of the abovementioned species in each Andalusian municipality ($n = 771$), according to the following equation:

$$\mathrm{HY} = \frac{\sum \text{mean annual number of individuals hunted per game estate}}{\sum \text{areas of the game estates}} \times 100$$

where HY is the hunting yield per municipality expressed by the number of individuals captured per 100 ha of game estate where the species is hunted (Vargas *et al.* 2007). The Mainland Spanish Fish, Game and National Parks Service made abundance maps for each game species by using estimated hunting yields in the 1960s. These maps, whose scale is 1:2 000 000, indicate the abundance of the main game species throughout Spain on a 1–6 scale (where 1 = absent, 2 = rare, 3 = scarce, 4 = frequent, 5 = abundant and 6 = very abundant; Ministerio de Agricultura 1968; see also Gortázar *et al.* 2000; Delibes-Mateos *et al.* 2009*a*). Using this information, we extracted the mean value of wild rabbit

https://doi.org/10.1017/S0376892910000214 Published online by Cambridge University Press

Table 1 Variables used to model the potential distribution of wild rabbit abundance in south-Iberian municipalities. Sources: [1]US Geological Survey (1996); [2]derived from GlobDEM50 (Farr & Kobrick, 2000); [3]Mapa de usos y coberturas vegetales del suelo de Andalucía (1956, 1999).

Variables		*Code*
Orographic	Altitude (m) [1]	ALTI
	Slope (%) [1]	SLOP
	Exposure to the south [2]	SE
	Exposure to the west [2]	WE
Natural vegetation	Built land (% area) [3]	BL
	Wetlands (% area) [3]	WETL
	Pasture (% area) [3]	PAST
	Oak wood (% area) [3]	OAKW
	Pasture with oaks (% area) [3]	PWO
	Pasture with conifers (% area) [3]	PWC
	Dense scrub with oaks (% area) [3]	DSWO
	Sparse scrub (% area) [3]	SS
	Sparse scrub with oaks (% area) [3]	SSWO
	Dense scrub with conifers (% area) [3]	DSWC
	Sparse scrub with conifers (% area) [3]	SSWC
	Sparse scrub with diverse trees (% area) [3]	SSWD
	Dense scrub with diverse trees (% area) [3]	DSWD
	Conifer wood (% area) [3]	CW
	Dense scrub (% area) [3]	DS
Crop	Irrigated herbaceous crops (% area) [3]	IHER
	Irrigated woody crops (% area) [3]	IWC
	Dry herbaceous crops (% area) [3]	DHER
	Dry heterogeneous crops (% area) [3]	DHET
	Irrigated heterogeneous crops (% area) [3]	IHET
	Dry wood crops (% area) [3]	DWC
	Mosaic of crops and natural vegetation (% area) [3]	MCNV
	Herbaceous crops with oaks (% area) [3]	HCWO

abundance in each municipality following the procedure described in Delibes-Mateos *et al.* (2009*a*).

As our aim was to detect areas favourable to rabbits, we followed the criterion of Farfán *et al.* (2004) and Vargas *et al.* (2006) to estimate where the abundance of this species was good (index of abundance, IA = 1) or poor (IA = 0). In this way, IA = 1 for the present period, when HY > 20, as well as for the 1960s, when the abundance value was 4 or higher. In contrast, IA = 0 for the present period when HY ≤ 20, as well as for the 1960s, when the abundance value was lower than 4. IA was then used as a target variable in the modelling procedure.

We related the IA to 27 predictor variables that provided information on the environmental characteristics, land use and vegetation in the Andalusian municipalities (Table 1). Orographic variables were derived from US Geological Survey (1996), and GlobDEM50 (Farr & Kobrick, 2000), whereas natural vegetation and crops variables were obtained from Mapa de usos y coberturas vegetales del suelo de Andalucía (Junta de Andalucía 1956, 1999). Exposure to the south and exposure to the west (the SE and WE variables) were derived from GlobDEM50 high-resolution digital elevation data, based on raw data from the Shuttle Radar Topography Mission (SRTM; Farr & Kobrick 2000). To this end, we used the spatial analyst toolbox of ArcMap. We calculated a 90 × 90 m resolution aspect map with surface analysis, and from this we extracted the degree of exposure to the south and west, respectively. Thus, for variable SE, a pixel whose aspect is south was given the value 180, a pixel whose aspect is north was given the value 0, and pixels with intermediate aspects were given intermediate values. The procedure was analogous for variable WE. More details in relation to the process used to obtain the rest of the orographic, natural vegetation and crop variables included in Table 1 are provided in Vargas *et al.* (2007).

Predictive models

To select a subset of significant predictor variables, we performed stepwise logistic regression (Hosmer & Lemeshow 1989) of IA on the predictor variables, using SPSS 14.0 statistical software. We then used the environmental favourability function of Real *et al.* (2006) to eliminate from the model the effect of the uneven proportion of ones and zeros in the dataset. The favourability for a positive IA in each municipality was obtained from the formula:

$$F = (P/(1 - P))/((n1/n0) + (P/(1 - P)))$$

where P is the probability value given by logistic regression, and $n1$ and $n0$ are the number of municipalities with IA equal to 1 and 0, respectively (Real *et al.* 2006). This function provided a description of local deviations from the overall probability of obtaining good abundances. Thus, a value $F > 0.5$ meant that the probability of an IA = 1 (anticipated owing to local environmental conditions) was higher than that expected only according to the IA = 1 / IA = 0 ratio (namely the reported IA = 1 prevalence in the territory).

To obtain an explanatory model, the variables introduced in the final predictive model were grouped into orographic, natural vegetation and crop factors (Table 1), and each group of variables was used to obtain partial orographic, natural vegetation and crop favourability models. To take into account the effect not explained by a single factor, which often results in an overlaid effect in space owing to collinearity between them (Borcard *et al.* 1992; Legendre 1993), we

https://doi.org/10.1017/S0376892910000214 Published online by Cambridge University Press

performed a variation partitioning procedure to specify how much of the variation of the final model was explained by the pure effect of each explanatory factor, which proportion was an indistinguishable effect of more than a single factor (intersection) and how these factors interacted and affected the distribution of the wild rabbit abundances (Legendre 1993; Legendre & Legendre 1998; see an application in Farfán *et al.* 2008). Mathematically, negative intersections between factors can appear, which measure the amount by which the effect of a factor is obscured by another factor through interrelationships between variables (Cartron *et al.* 2000; Bárcena *et al.* 2004).

Municipalities with high, low or intermediate environmental favourability for rabbits were defined according to a classification threshold. We considered favourable all municipalities whose favourability was 0.8 or higher, that is, where the odds of good abundance were at least 4:1 (Rojas *et al.* 2001; Muñoz & Real 2006). Municipalities whose favourability was 0.2 or lower were considered unfavourable (maximum odds 1:4), and favourability values between 0.2 and 0.8 were considered intermediate. We then compared the two models for the two periods that represented those municipalities where favourability was high in the 1960s and now, where this was low in the 1960s and now, and where favourability had changed from high to intermediate (F-I), from intermediate to low (I-U), from low to intermediate (U-I) and from intermediate to high (I-F). Changes in the areas dedicated to the different land uses within every municipality were calculated as percentages (according to the Mapa de usos y coberturas vegetales del suelo de Andalucía; Junta de Andalucía 1956, 1999), and then we quantified the changes in every land use within the F-I, I-U, U-I and I-F areas.

RESULTS

The variables included in the logistic regression models, ranked according to their order of entrance in each model (Table 2) included six variables that were common to both models, representing 66% and 75% of the variables included in the 1960s and the current models, respectively. According to these models, municipalities in Andalusia judged as favourable for rabbits tend to be aggregated (Fig. 2). In the 1960s, the most favourable areas for wild rabbit were mainly located in the western part of the Sierra Morena and Guadalquivir valley. However, currently favourable areas are located along the Guadalquivir slope of the Sub-baetic System and less mountainous regions of the western part of the Baetic System (Fig. 2).

The relationships between the explanatory factors were complex (Fig. 3). In the 1960s, the effect owing to the intersection of orography, natural vegetation and crops was positive, and the characteristics of factors that favour good abundances tended to be present simultaneously. However, the effect owing to the intersection between orography and natural vegetation was negative after excluding the effect of crops; this was also the case regarding both the effect owing to the intersection between natural vegetation and crops after excluding the effect of orography, as well as the effect owing to the intersection between orography and crops after excluding the effect of natural vegetation. At present, the intersection between factors shows the same pattern as in the 1960s, except for the effect due to the intersection between orography and natural vegetation, which has changed from negative to positive (Fig. 3).

Table 2 Favourability models including coefficients of variables in the favourability functions. The Wald parameter indicates the relative importance of each variable. p = statistical significance.

Year	*Variable*	*Coefficient*	*Wald*	*p*
1960s	Slope	−0.187	56.266	<0.001
	Sparse scrub with oaks	5.696	24.274	<0.001
	Dry woody crop	2.442	33.198	<0.001
	Pasture	5.157	10.460	<0.01
	Sparse scrub	1.896	10.008	<0.01
	Exposure to the west	−0.0164	7.364	<0.01
	Dense scrub with diverse trees	7.414	6.194	<0.05
	Wetlands	2.889	4.478	<0.05
	Exposure to the south	0.0102	3.966	<0.05
	Constant	−0.254		
Current	Dry woody crop	2.624	45.004	<0.001
	Pasture	10.697	16.549	<0.001
	Slope	−0.120	15.564	<0.001
	Sparse scrub with oaks	4.048	9.970	<0.01
	Exposure to the south	−0.0148	7.782	<0.01
	Altitude	−0.000967	7.469	<0.01
	Herbaceous crops with oaks	−9.479	4.654	<0.05
	Sparse scrub	1.249	4.649	<0.05
	Constant	0.920		

Figure 4 shows the municipalities where favourable and unfavourable conditions for wild rabbits have remained stable from the 1960s to now, and where these conditions have changed. Some municipalities have changed toward more favourable conditions (Fig. 4*a*). Thus, in the western and central region of Andalusia, favourability increased from intermediate to favourable, whereas in the eastern region this improved from unfavourable to intermediate. Other municipalities have changed toward more unfavourable conditions (Fig. 4*b*). Changes in favourability from favourable to intermediate are concentrated in the middle of the western Andalusia, and changes from intermediate to unfavourable mainly in the eastern region.

Municipalities where favourable and unfavourable conditions for rabbits changed between the 1960s and present day showed substantial changes in the percentage of surface area occupied by vegetation types included in the favourability models (Table 3). Thus, the percentages of surface area occupied by dry woody crops, pasture, sparse scrub with oak and sparse scrub have decreased in Andalusia since the

https://doi.org/10.1017/S0376892910000214 Published online by Cambridge University Press




Figure 2 Favourability values for good wild rabbit abundances in the municipalities of Andalusia, shown on a scale ranging from 0 (white) to 1 (black).




Figure 3 Variation partitioning of the final model. Values shown in the diagrams are the percentages of variation in good abundance explained by the factors orography, natural vegetation and crops and by their interactions.

1960s. Nevertheless, in the municipalities where conditions favourable for rabbits have increased, the percentage of surface area occupied by these vegetation types has increased. The percentage of herbaceous crops with oaks increased in Andalusia since the 1960s. However, in the municipalities where favourable conditions increased, the percentage of surface area occupied by this vegetation type decreased, whereas in the municipalities where favourable conditions decreased, the percentage of surface area occupied by this vegetation type increased.

DISCUSSION

The location of areas favourable for rabbits in Andalusia has changed substantially over recent decades. In fact, we have recorded an increase in favourability (especially from intermediate to favourable) in the areas that are currently favourable for the species, and a favourability decrease in areas that were favourable for the lagomorph during the 1960s. Two hypotheses may explain these changes in favourability. First, rabbit habitat requirements could have changed in recent decades. Second, recent changes in land use in Andalusia (Fernández-Alés *et al.* 1992) could explain the geographical differences in rabbit favourability between the two study periods.

As regards the first hypothesis, our results suggest that favourable areas for rabbits are currently determined by environmental and land-use factors similar to those of the 1960s; this is supported by the fact that six variables were repeated in both favourability models (Table 2). Thus, as in the 1960s, favourable areas for rabbits are currently associated with the main habitat requirements for this small mammal: the presence of suitable food types (for example dry crops and pastures; Calvete *et al.* 2004; Fernández 2005), and the availability of some form of cover for protection against predators (such as sparse Mediterranean scrubland; Moreno & Villafuerte 1995). Moreover, land slope was negatively associated with rabbit favourability during both study periods, which is not surprising given that rabbits typically avoid sloping mountain areas (Farfán *et al.* 2008). In only one of

https://doi.org/10.1017/S0376892910000214 Published online by Cambridge University Press

Table 3 Increase in the percentage of surface area occupied by land use and vegetation types included in the favourability models. I-F: municipalities where the environment has changed from intermediate to favourable for the wild rabbit; U-I: from unfavourable to intermediate; F-I: from favourable to intermediate; I-U: from intermediate to unfavourable.

Included in models		*Andalusia*	*I-F, U-I*	*F-I, I-U*
1960s and current	Dry woody crop	−4.3	17.3	−19.4
	Pasture	−14.9	33.5	−66.7
	Sparse scrub with oaks	−0.6	66.5	−30.6
	Sparse scrub	−15.1	3.1	−17.3
1960s	Dense scrub with diverse trees	7.0	−67.8	−43.1
	Wetlands	−4.7	−3.4	−5.4
Current	Herbaceous crops with oaks	60.0	−45.8	162.7




Figure 4 Municipalities where favourable and unfavourable conditions for a high abundance of wild rabbits have remained stable from the 1960s to now (black and light grey, respectively), and where these conditions have changed (dark grey). (*a*) Dark grey indicates change toward more favourable conditions: either from intermediate to favourable (I-F zone), or from unfavourable to intermediate (U-I zone); (*b*) dark grey indicates change toward less favourable conditions: either from favourable to intermediate (F-I zone), and from intermediate to unfavourable (I-U zone). Thresholds for favourable and unfavourable areas are 0.8 and 0.2, respectively.

the models, a few less statistically significant land use variables were retained. For instance, dense scrub with diverse trees was associated with areas favourable to rabbits during the 1960s, but not at present (Table 2). Small patches of dense scrubland were interspersed with pastures and crops several decades ago in Andalusia (Fernández-Alés *et al.* 1992), and therefore rabbits could find refuge and food in this type of landscape. Similarly, herbaceous crops with oaks were only present in the current favourability model (Table 2). This habitat was negatively associated with favourable areas for rabbits (Table 2), apparently because it offers high food availability but little refuge protection for the species (Lombardi *et al.* 2003). The area devoted to herbaceous crops significantly increased in recent decades (Table 3; see also Fernández-Alés *et al.* 1992), which could explain why this habitat type was only significant in the present day favourability model (Table 2). The possibility that rabbits have changed their habitat requirements in the current landscape configuration of southern Iberia is not totally excluded by coincidences in the model, as the internal relationships between the variables that are common to both models (their coefficients and order of importance according to the Wald's parameter) are not identical.

The partial orographic, natural vegetation and crop models show that orography is the most important factor explaining the distribution of wild rabbit abundance in Andalusia (Fig. 3). In the 1960s, the types of natural vegetation landscape that favoured the presence of rabbits were more widely distributed throughout the study area (Fernández-Alés *et al.* 1992) and, as a consequence, orography may have been a more limiting factor for this species than natural vegetation. Negative intersection between natural vegetation and crops during the 1960s (Fig. 3) means that the favourable natural vegetation conditions tend not to coincide with the favourable crop conditions when orography remains constant. After the 1960s, the partial importance of orography and natural vegetation decreased, whereas the weight of their intersection has increased and become positive (Fig. 3), probably because natural vegetation landscapes favourable for rabbits are presently confined to certain orographic zones. Specifically, the most suitable natural habitats are now mainly situated on the Guadalquivir slope of the Sub-baetic system and the western hills of the Penibaetic system (Fig. 2), which are regions with moderate slopes that are also orographically favourable for rabbits.

As habitat requirements for rabbits do not seem to have changed, the most plausible explanation for the changes observed in rabbit favourability would be the land-use changes in Andalusia in recent decades. In agreement with this, we suggest that the changes in landscape have been mainly detrimental to rabbits during the study period. Thus, the percentages of the four habitat variables that were positively associated with rabbit favourability in both models (current and 1960s; Table 2) have declined in Andalusia as a whole and in areas where rabbit favourability has decreased (Table 3; but see also Fig. 4*b*); hence, habitat suitable for rabbits has

https://doi.org/10.1017/S0376892910000214 Published online by Cambridge University Press

become impoverished. The reduction in woody crops and pastures may have led to a decline in the availability of food for rabbits in these areas (Calvete *et al.* 2004). Similarly, the loss of sparse scrubland, mainly as a consequence of rural abandonment (Fernández-Alés *et al.* 1992), has significantly reduced rabbit numbers in some areas in Andalusia (Moreno & Villafuerte 1995). Moreover, the increase in both dense scrubland and herbaceous crops in Andalusia (Table 3) seems to be detrimental for rabbits, since these are not optimum habitats for the species (Lombardi *et al.* 2003). From this point of view, the large decrease in dense scrubland in municipalities where rabbit favourability has decreased would be positive. However, dense scrubland has apparently been replaced in these areas by other habitats unfavourable to rabbits, such as herbaceous crops with oaks (Table 3). In fact, the area devoted to herbaceous crops with oaks has tremendously increased in those municipalities where rabbit favourability has decreased (Table 3). Not only has the surface area of this type of habitat (typically called *dehesa*) increased in the study area, but it has also undergone structural modifications. The scrub layer traditionally linked to the dehesa has progressively disappeared as a consequence of agriculture and livestock intensification (Fernández-Alés *et al.* 1992). It is known that improvements associated with modern agricultural practices can have long-term detrimental effects on rabbit populations (Boag 1987). Interestingly, land uses suitable for rabbits in both models (Table 2) have increased in municipalities where rabbit favourability has also increased (Table 3; see also Fig. 4*a*). Therefore, an improvement in rabbit habitat has occurred in these locations. In addition, the reduction of the surface area devoted to herbaceous crops with oaks and to dense scrubland has contributed to the habitat improvement recorded in these areas.

We have not addressed specifically the effect of diseases on rabbit population decline. Nevertheless, a recent review has showed that positive rabbit trends, after the initial RHD outbreak, have been recorded in species-friendly habitats (Delibes-Mateos *et al.* 2009*b*). Similarly, a theoretical model also showed that the long-term impact of RHD is conditioned by population dynamics, which are primarily determined by habitat suitability (Calvete 2006). According to these findings, the impoverishment of rabbit preferred habitat observed in the present study could be jeopardizing rabbit recovery in the Iberian Peninsula.

CONCLUSIONS

The availability of hunting records with explicit time and space references allows the development of datasets which can be used to analyse species trends in historic and regional contexts (Fernández & Ruiz de Azua 2009). We have used two sources based on hunting records, which were previously used to assess changes in species abundance and distribution (Gortázar *et al.* 2000; Delibes-Mateos *et al.* 2009*a*), to estimate long-term rabbit population trends. Our aim was to detect areas favourable to rabbits, thus to avoid potential bias owing to the existence of differences in the nature and quality of the data between the two study periods, we transformed the original data into a binomial variable (good and poor areas for rabbit) that was directly comparable between both study dates.

There is a widely-held perception that all lagomorphs are fecund and are sufficiently generalist in their ecology to overcome environmental changes. However, approximately a quarter of all lagomorphs are threatened with extinction and, to a great extent this is owing to land-use changes during the last century (Smith 2008). Although it has been previously suggested that habitat loss and fragmentation has been a major cause of rabbit decline (Ward 2005), our study provides the first empirical evidence showing that habitat changes have been highly detrimental for this keystone species in the Iberian Mediterranean ecosystem. Following the rabbit population decline, conservationists and hunters have applied a great number of management tools to improve rabbit densities (for example Delibes-Mateos *et al.* 2008*b*). Based on our findings, it is to be expected that these strategies would be unsuccessful in areas where favourable habitats for rabbits have disappeared as a consequence of landscape changes. Thus, it would be preferable to conserve and recover the landscapes suitable for the species.

ACKNOWLEDGEMENTS

M. Delibes-Mateos was supported by a postdoctoral grant from the regional government of Castilla la Mancha (JCCM) and the European Social Fund (ESF). Special thanks go to E. Martínez for providing the maps of the abundance of game species during the 1960s. We are indebted to D. Anderson, N. Fernández, J. Fernández de Simón, D. Parish and two anonymous reviewers for helpful comments on previous drafts of the manuscript. The Consejería de Medio Ambiente de la Junta de Andalucía kindly supplied the Annual Hunting Reports.

References

Bárcena, S., Real, R., Olivero, J. & Vargas, J.M. (2004) Latitudinal trends in breeding waterbird species richness in Europe and their environmental correlates. *Biodiversity and Conservation* **13**: 1997–2014.

Boag, B. (1987) Reduction in numbers of the wild rabbit (*Oryctolagus cuniculus*) due to changes in agricultural practices and land use. *Crop Protection* **6**: 347–351.

Borcard, D., Legendre, P. & Drapeau, P. (1992) Partialling out the spatial component of ecological variation. *Ecology* **73**: 1045–1055.

Calvete, C. (2006) Modelling the effect of population dynamics on the impact of rabbit hemorrhagic disease. *Conservation Biology* **20**: 1232–1241.

Calvete, C., Estrada, R., Angulo, E. & Cabezas-Ruiz, S. (2004) Habitat factors related to wild rabbit conservation in an agricultural landscape. *Landscape Ecology* **19**: 531–542.

Calvete, C., Pelayo, E. & Sampietro, J. (2006) Habitat factors related to wild rabbit population trends after the initial impact of rabbit haemorrhagic disease. *Wildlife Research* **33**: 467–474.

https://doi.org/10.1017/S0376892910000214 Published online by Cambridge University Press

Cartron, J.L.E., Kelly, J.F. & Brown, J.H. (2000) Constraints on patterns of covariation: a case study in strigid owls. *Oikos* **90**: 381–389.

Delibes-Mateos, M., Delibes, M., Ferreras, P. & Villafuerte, R. (2008*a*) Key role of European rabbits in the conservation of the western Mediterranean Basin Hotspot. *Conservation Biology* **22**: 1106–1117.

Delibes-Mateos, M., Farfán, M.A., Olivero, J., Márquez, A.L. & Vargas, J.M. (2009*a*) Long-term changes in game species over a long period of transformation in the Iberian Mediterranean landscapes. *Environmental Management* **43**: 1256–1268.

Delibes-Mateos, M., Ferreras, F. & Villafuerte, R. (2009*b*) European rabbit population trends and associated factors: a review of the situation in the Iberian Peninsula. *Mammal Review* **39**: 124–40.

Delibes-Mateos, M., Ramírez, E., Ferreras, P. & Villafuerte, R. (2008*b*) Translocations as a risk for the conservation of European wild rabbit *Oryctolagus cuniculus* lineages. *Oryx* **42**: 259–264.

Delibes-Mateos, M., Redpath, S.M., Angulo, E., Ferreras, P. & Villafuerte, R. (2007) Rabbits as a keystone species in southern Europe. *Biological Conservation* **137**: 149–156.

Farfán, M.A., Guerrero, J.C., Real, R., Barbosa, A.M. & Vargas, J.M. (2004) Caracterización del aprovechamiento cinegético de los mamíferos en Andalucía. *Galemys* **16**: 41–59.

Farfán, M.A., Vargas, J.M., Guerrero, J.C., Barbosa, A.M., Duarte, J. & Real, R. (2008) Distribution modelling of wild rabbit hunting yields in its original area (S Iberian Peninsula). *Italian Journal of Zoology* **75**: 161–172.

Farr, T.G. & Kobrick, M. (2000) Shuttle Radar Topography Mission produces a wealth of data. *EOS Transactions of the American Geophysical Union* **81**: 583–585.

Fernández, J.M. & Ruiz de Azua, N. (2009) Historical dynamics of a declining wolf population: persecution vs. prey reduction. *European Journal of Wildlife Research* (in press).

Fernández, N. (2005) Spatial patterns in European rabbit abundance after a population collapse. *Landscape Ecology* **20**: 897–910.

Fernández-Alés, R., Martín, A., Ortega, F. & Ales, E.E. (1992) Recent changes in landscape structure and function in a Mediterranean region of SW of Spain (1950–1984). *Landscape Ecology* **7**: 3–18.

Gálvez-Bravo, L., Belliure, J. & Rebollo, S. (2009) European rabbits as ecosystem engineers: warrens increase lizard density and diversity. *Biodiversity and Conservation* **18**: 869–885.

Gortázar, C., Herrero, J., Villafuerte, R. & Marco, J. (2000) Historical examination of the status of large mammals in Aragon, Spain. *Mammalia* **64**: 411–422.

Hosmer, D.W. & Lemeshow, S. (1989) *Applied Logistic Regression*. New York, NY, USA: John Wiley & Sons, Inc.

Junta de Andalucía (1956) *Mapa de usos y coberturas vegetales del suelo de Andalucía 1956*. Sevilla, Spain: Consejería de Medio Ambiente.

Junta de Andalucía (1999) *Mapa de usos y coberturas vegetales del suelo de Andalucía 1999*. Sevilla, Spain: Consejería de Medio Ambiente.

Legendre, P. (1993) Spatial autocorrelation: trouble or new paradigm? *Ecology* **74**: 1659–1673.

Legendre, P. & Legendre, L. (1998) *Numerical Ecology*. Second English Edition. Amsterdam, the Netherlands: Elsevier Science.

Lombardi, L., Fernández, N., Moreno, S. & Villafuerte, R. (2003). Habitat-related differences in rabbit (*Oryctolagus cuniculus*) abundance, distribution and activity. *Journal of Mammalogy* **84**: 26–36.

Ministerio de Agricultura (1968) *Mapa Cinegético Nacional*. Madrid, Spain: Ministerio de Agricultura.

Moreno, S. & Villafuerte, R. (1995) Traditional management of scrubland for the conservation of rabbits *Oryctolagus cuniculus* and their predators in Doñana National Park, Spain. *Biological Conservation* **73:** 81–85.

Muñoz, A.R. & Real, R. (2006) Assessing the potential range expansion of the exotic monk parakeet in Spain. *Diversity and Distribution* **12:** 656–665.

Piorno, V. (2006) Gestión cinegética y conservación del conejo de monte. Ph.D. thesis, Universidad de Vigo, Vigo, Spain.

Real, R., Barbosa, A.M. & Vargas, J.M. (2006) Obtaining environmental favourability functions from logistic regression. *Environmental and Ecological Statistics* **13:** 237–245.

Rojas, A.B., Cotilla, I., Real, R., Palomo, L.J. (2001) Determinación de las áreas probables de distribución de los mamíferos terrestres en la provincia de Málaga a partir de las presencias conocidas. *Galemys* **13:** 217–229.

Smith, A.T. (2008) Conservation of endangered lagomorphs. In: *Lagomorph Biology: Evolution, Ecology and Conservation*, ed. P.C. Alves & K. Hackländer, pp. 297–315. Berlin, Germany: Springer-Verlag.

US Geological Survey (1996) *GTOPO30*. Land Processes Distributed Archive Center [www document].URL http://edcdaac.usgs.gov/gtopo30/gtopo30.asp

Vargas, J.M., Farfán, M.A., Guerrero, J.C., Barbosa, A.M. & Real, R. (2007) Geographical and environmental correlates of big and small game in Andalusia (southern Spain). *Wildlife Research* **34**: 498–506.

Vargas, J.M., Guerrero, J.C., Farfán, M.A., Barbosa, A.M. & Real, R. (2006) Land use and environmental factors affecting red-legged partridge (*Alectoris rufa*) hunting yields in Southern Spain. *European Journal of Wildlife Research* **52**: 188–195.

Villafuerte, R., Viñuela, J. & Blanco, J.C. (1998) Extensive predator persecution caused by population crash in a game species: the case of red kites and rabbits in Spain. *Biological Conservation* **84**: 181–188.

Virgós, E., Cabezas-Díaz, S. & Lozano, J. (2007) Is the wild rabbit (*Oryctolagus cuniculus*) a threatened species in Spain? Sociological constraints in the conservation of species. *Biodiversity and Conservation* **16:** 3489–3504.

Ward, D. (2005) Reversing rabbit decline: one of the biggest challenges for nature conservation in Spain and Portugal. Technical report. World Conservation Union, Gland, Switzerland [www document]. URL http://www.ualberta.ca/~dhik/lgs/report_lynx_rabit.pdf

https://doi.org/10.1017/S0376892910000214 Published online by Cambridge University Press