Math. Struct. in Comp. Science (2008), *vol.* 18, *pp.* 975–1004. © 2008 Cambridge University Press
doi:10.1017/S0960129508006683 First published online 15 July 2008 Printed in the United Kingdom

Graph lambda theories[†]

ANTONIO BUCCIARELLI[‡] and ANTONINO SALIBRA[§¶]

[‡]*Equipe PPS (case 7014), Université Paris 7, 2, place Jussieu, 72251 Paris Cedex 05, France*
Email: buccia@pps.jussieu.fr
[§]*Dipartimento di Informatica, Università Ca'Foscari di Venezia,*
Via Torino 155, 30172 Venezia, Italy
Email: salibra@dsi.unive.it

Received 1 November 2006; revised 29 October 2007

A longstanding open problem in lambda calculus is whether there exist continuous models of the untyped lambda calculus whose theory is exactly the $\lambda\beta$ or the least sensible λ-theory $\mathscr{H}$ (which is generated by equating all the unsolvable terms). A related question is whether, given a class of lambda models, there are a minimal λ-theory and a minimal sensible λ-theory represented by it. In this paper, we give a positive answer to this question for the class of graph models *à la* Plotkin, Scott and Engeler. In particular, we build two graph models whose theories are the set of equations satisfied in, respectively, any graph model and any sensible graph model. We conjecture that the least sensible graph theory, where 'graph theory' means 'λ-theory of a graph model', is equal to $\mathscr{H}$, while in one of the main results of the paper we show the non-existence of a graph model whose equational theory is exactly the $\lambda\beta$ theory.
Another related question is whether, given a class of lambda models, there is a maximal sensible λ-theory represented by it. In the main result of the paper, we characterise the greatest sensible graph theory as the λ-theory $\mathscr{B}$ generated by equating λ-terms with the same Böhm tree. This result is a consequence of the main technical theorem of the paper, which says that all the equations between solvable λ-terms that have different Böhm trees fail in every sensible graph model. A further result of the paper is the existence of a continuum of different sensible graph theories strictly included in $\mathscr{B}$.

1. Introduction

The untyped lambda calculus was introduced around 1930 (Church 1933; 1941) as part of an investigation into the formal foundations of mathematics and logic. Although lambda calculus is a very basic language, it is sufficient to express all computable functions. The process of application and evaluation reflects the computational behaviour of many modern functional programming languages, which explains the interest in the lambda calculus among computer scientists.

[†] This article is a revised and expanded version of two earlier papers by the authors, which appeared in the Proceedings of the 28th International Symposium on Mathematical Foundations of Computer Science (MFCS'03) (Bucciarelli and Salibra 2003) and in the Proceedings of the 19th Annual IEEE Symposium on Logic in Computer Science (LICS'04) (Bucciarelli and Salibra 2004).

[¶] Work partially supported by the Equipe PPS of the University Paris 7-Denis Diderot, and by MIUR Cofin'04 FOLLIA Project.

https://doi.org/10.1017/S0960129508006683 Published online by Cambridge University Press

Lambda theories are equational extensions of the untyped lambda calculus that are closed under derivation. They may arise through syntactical or semantic considerations. Indeed, a λ-theory may correspond to a possible operational (observational) semantics of the lambda calculus, as well as being induced by a model of lambda calculus through the kernel congruence relation of the interpretation function. Although researchers have mainly focused on a limited number of λ-theories, the class of λ-theories constitutes a very rich and complex structure, see, for example, Barendregt (1984) and Berline (2000). It is usually difficult to employ syntactical techniques in the study of λ-theories, so semantic methods have been extensively investigated.

Topology has lain at the heart of the approaches taken to giving models of the untyped lambda calculus. The first model, found by Scott in 1969 in the category of complete lattices and Scott continuous functions, was successfully used to show that all unsolvable λ-terms can be consistently equated. After Scott, a large number of mathematical models for lambda calculus, arising from syntax-free constructions, have been introduced in various categories of domains, and were classified into semantics according to the nature of their representable functions, see, for example, Abramsky (1991), Barendregt (1984), Berline (2000) and Plotkin (1993). Scott's continuous semantics (Scott 1972) is given in the category whose objects are complete partial orders and whose morphisms are Scott continuous functions. The stable semantics (Berry 1978) and the strongly stable semantics (Bucciarelli and Ehrhard 1991) are a strengthening of the continuous semantics, which was introduced to capture the notion of 'sequential' Scott continuous function. All these semantics are structurally and equationally rich in the sense that in each of them it is possible to build up $2^{\aleph_0}$ models inducing pairwise distinct λ-theories (Kerth 1998; 2001). Nevertheless, the above denotational semantics are *equationally incomplete*: they do not match all possible operational semantics of lambda calculus. The problem of the equational incompleteness was positively solved by Honsell and Ronchi della Rocca for the continuous semantics (Honsell and Ronchi della Rocca 1992), and by Bastonero and Gouy for the stable semantics (Gouy 1995; Bastonero and Gouy 1999). Salibra has recently shown in a uniform way that all semantics that involve monotonicity with respect to some partial order and have a bottom element are equationally incomplete (Salibra 2001; 2003). From this we get the incompleteness of the strongly stable semantics, which had been conjectured by Bastonero and Gouy and by Berline (Bastonero and Gouy 1999; Berline 2000).

If a semantics is incomplete, there exists a λ-theory T that is not induced by any model in the semantics. In such a case we say that the semantics *omits* the λ-theory T. More generally, a semantics *omits* an equation if the equation fails in every model of the semantics (and *forces* it if it holds in every model of the semantics). The set of equations forced by a semantics $\mathscr{C}$ constitutes a λ-theory. It is the minimal (with respect to the inclusion order) λ-theory of $\mathscr{C}$ if it is induced by a model of $\mathscr{C}$.

The following questions arise naturally (Berline 2000) – given a class $\mathscr{C}$ of models of lambda calculus:

1 Is there a minimal λ-theory represented by $\mathscr{C}$?
2 Is there a minimal sensible λ-theory (that is, one that equates all unsolvable λ-terms) represented by $\mathscr{C}$?

It was shown in Di Gianantonio *et al.* (1995) that the answer to question (1) is positive for Scott's continuous semantics, at least if we restrict to extensional models. Another result in the

https://doi.org/10.1017/S0960129508006683 Published online by Cambridge University Press

same spirit is the construction of a model whose theory is $\lambda\beta\eta$, which is *a fortiori* minimal, in the ω_1-semantics. However, the proofs of Di Gianantonio *et al.* (1995) use logical relations, and since logical relations do not allow us to distinguish terms with the same applicative behaviour, the proofs do not carry across to non-extensional models.

In this paper we show that both questions (1) and (2) yield a positive answer for the *graph semantics*, that is, the semantics of lambda calculus given in terms of *graph models*. These models, which were isolated in the seventies by Scott and Engeler (Barendregt 1984) within the continuous semantics, have proved useful for giving proofs of the consistency of extensions of the lambda calculus and for studying its operational features. For example, the simplest graph model, namely Engeler's model, was used in Berline (2000) to give concise proofs of both the head- and left-normalisation theorems of lambda calculus, while a semantical proof of the 'easiness' of $(\lambda x.xx)(\lambda x.xx)$ was given in Baeten and Boerboom (1979). It is well known that the graph semantics is incomplete, since it trivially omits the axiom of extensionality. The main technical device used in the proof of the existence of the least (sensible) graph theory is the notion of a *weak product* of graph models. Roughly speaking, the weak product of a family of graph models is a new graph model that is the 'canonical completion' of the disjoint union of the models in the family. We show that the theory of a weak product is always semisensible (that is, it does not equate solvable and unsolvable terms), and it is included in the intersection of the theories of its factors (the inclusion is in general strict). The *least graph theory* (where 'graph theory' means 'λ-theory of a graph model') is the theory of the weak product of the family $(D_e : e \in I)$, where I is the set of equations between λ-terms that fail to hold in some graph model, and D_e is a fixed graph model not satisfying the equation e.

Two further questions naturally arise: which equations between λ-terms belong to the minimal graph theory and which to the minimal sensible graph theory? The answer to the second, difficult, question is still unknown: we conjecture that the λ-theory $\mathscr{H}$ generated by equating all unsolvable λ-terms is the least sensible graph theory. The first question is related to a longstanding open problem in lambda calculus, which asks whether there exists a non-syntactic model whose equational theory is equal to the least λ-theory $\lambda\beta$. In this paper we show that this model cannot be found within graph semantics (this result gives us a negative answer to Question 1 in Berline (2000, Section 6.2) for the class of graph models). From this result it follows that the minimal graph theory is not equal to $\lambda\beta$, so graph semantics forces equations between non-β-equivalent λ-terms. In this paper we provide an example of an equation of this kind.

The set of all sensible λ-theories constitutes a bounded lattice. The least sensible λ-theory is the λ-theory $\mathscr{H}$ (generated by equating all the unsolvable terms), while the greatest sensible λ-theory is the λ-theory $\mathscr{H}^*$ (generated by equating terms with the same Böhm tree up to possibly infinite η-equivalence). It was shown in Kerth (1998) that there exists a continuum of different sensible graph theories, so it make sense to ask whether there exists a maximal λ-theory represented by graph semantics. In one of our main results in this paper we show that the λ-theory $\mathscr{B}$ (generated by equating λ-terms with the same Böhm tree) is the greatest sensible graph theory. This result is a consequence of the main technical theorem of the paper: the graph semantics omits all equations $M = N$ between λ-terms satisfying the conditions

$$\begin{aligned} \mathscr{H}^* &\vdash M = N \\ \mathscr{B} &\nvdash M = N\,. \end{aligned} \tag{1}$$

https://doi.org/10.1017/S0960129508006683 Published online by Cambridge University Press

In other words, the graph semantics omits all equations $M = N$ between λ-terms that do not have the same Böhm tree, but have the same Böhm tree up to (possibly infinite) η-equivalence.

The following are also consequences of the main result of the paper:

(i) There exists a continuum of different sensible graph theories strictly included in $\mathscr{B}$ (this result provides a positive answer to Question 2 in Berline (2000, Section 6.3)).
(ii) For every closed term P, the λ-theory generated by $\Omega = P$, where Ω is the paradigmatic unsolvable term $(\lambda x.xx)(\lambda x.xx)$, contains no equation satisfying condition (1).

The paper is organised as follows. In Section 2 we review the basic definitions of lambda calculus and graph models. In particular, we recall the formal definition of the canonical completion of a partial model. The notion of a weak product of graph models is introduced and developed in Section 3. The proof of the existence of a minimal (sensible) graph theory is presented in Section 4, while in Section 5 we show that the least graph theory is not equal to $\lambda\beta$. Sections 6–8 are devoted to characterising the maximal sensible graph theory. Conclusions and ideas for future work are presented in Section 9.

2. Preliminaries

To help keep this article self-contained, we begin with a summary of some definitions and results for lambda calculus and graph models. We will follow the notation and terminology used in Barendregt (1984) for the lambda calculus.

2.1. *Lambda calculus*

The set Λ of λ-terms over an infinite set of variables is constructed as usual: every variable is a λ-term; and if M and N are λ-terms, then so are (MN) and $\lambda x.M$ for each variable x. Λ^o denotes the set of closed λ-terms.

The symbol $\equiv$ denotes syntactic equality. The following are some well-known λ-terms:

$$\begin{aligned} \mathbf{\Omega} &\equiv (\lambda x.xx)(\lambda x.xx) \\ \mathbf{\Omega}_3 &\equiv (\lambda x.xxx)(\lambda x.xxx) \\ \mathbf{i} &\equiv \lambda x.x \\ \mathbf{k} &\equiv \lambda xy.x \\ \mathbf{1} &\equiv \lambda xy.xy\,. \end{aligned}$$

A *compatible λ-relation* T is any set of equations between λ-terms that is closed under the following two rules:

(i) If $M = N \in T$ and $P = Q \in T$, then $MP = NQ \in T$.
(ii) If $M = N \in T$, then $\lambda x.M = \lambda x.N \in T$ for every variable x.

We will write either $T \vdash M = N$ or $M =_T N$ for $M = N \in T$.

A *λ-theory* T is any compatible λ-relation that is an equivalence relation and includes (α)- and (β)-conversion. The set of all λ-theories is naturally equipped with a lattice structure, with meet defined as set theoretical intersection. The join of two λ-theories T and S is the least

https://doi.org/10.1017/S0960129508006683 Published online by Cambridge University Press

equivalence relation including $T \cup S$. $\lambda\beta$ denotes the minimal λ-theory, and $\lambda\beta\eta$ denotes the minimal extensional λ-theory (axiomatised by $\mathbf{i} = \mathbf{1}$).

Solvable λ-terms can be characterised as follows: a λ-term M is solvable if and only if it has a *head normal form*, that is, $M =_{\lambda\beta} \lambda x_1 \dots x_n.yM_1 \dots M_k$ for some $n,k \geqslant 0$ and λ-terms $M_1,\dots,M_k$. $M \in \Lambda$ is *unsolvable* if it is not solvable.

The λ-theory $\mathscr{H}$, generated by equating all unsolvable λ-terms, is consistent (Barendregt 1984, Theorem 16.1.3) and admits a unique maximal consistent extension $\mathscr{H}^*$ (Barendregt 1984, Theorem 16.2.6). A λ-theory T is said to be *sensible* if it is consistent and $\mathscr{H} \subseteq T$. The set of all sensible λ-theories is naturally equipped with a bounded lattice structure. $\mathscr{H}$ is the least sensible λ-theory, and $\mathscr{H}^*$ is the greatest one. $\mathscr{H}^*$ is an extensional λ-theory. A λ-theory is *semisensible* if no solvable term is equivalent to an unsolvable term. It is easy to prove that sensible theories are semisensible. It is also possible to characterise semisensible λ-theories as follows: a λ-theory T is semisensible if and only if $T \subseteq \mathscr{H}^*$ (Barendregt 1984, Section 16.2).

2.2. *Böhm trees*

A λ-term M is called *a projection term* if $M \equiv \lambda x_1 \dots x_n.y$ ($n \geqslant 0$). A *Böhm-like tree* is a finite branching labelled tree, whose inner nodes are labelled by projection terms, and whose leaves are labelled by either projection terms or $\perp$. The equality of Böhm-like trees is defined up to α-equivalence of the corresponding labels.

The Böhm tree $BT(M)$ of a λ-term M is a finite or infinite Böhm-like tree. If M is unsolvable, then $BT(M) = \perp$, that is, $BT(M)$ is a tree with a unique node labelled by $\perp$. If M is solvable and $\lambda x_1 \dots x_n.yM_1 \dots M_k$ is the principal head normal form of M (Barendregt 1984, Definition 8.3.20), we have

$$BT(M) = \lambda x_1 \dots x_n.y$$

$$BT(M_1) \dots\dots\dots BT(M_k)$$

The λ-theory $\mathscr{B}$ generated by equating λ-terms with the same Böhm tree is sensible and non-extensional. $\mathscr{B}$ is distinct from $\mathscr{H}$ and $\mathscr{H}^*$, so $\mathscr{H} \subset \mathscr{B} \subset \mathscr{H}^*$. Note that Barendregt (1984, Exercise 16.5.5) gives a non-extensional λ-theory T satisfying $\mathscr{B} \subset T \subset \mathscr{H}^*$ (see the remark following Theorem 46 in the current paper).

In the remaining part of this section we characterise the λ-theory $\mathscr{H}^*$ in terms of Böhm trees.

For all λ-terms M and N, we write $M \leqslant_\eta N$ if $BT(N)$ is a (possibly infinite) *η-expansion* of $BT(M)$ (Barendregt 1984, Definition 10.2.10). For example, let $J \equiv \Theta(\lambda jxy.x(jy))$, where Θ is Turing's fixpoint combinator. Hence, $x \leqslant_\eta Jx$ (Barendregt 1984, Example 10.2.9), since

$$\begin{aligned} Jx &=_{\lambda\beta} \lambda z_0.x(Jz_0) \\ &=_{\lambda\beta} \lambda z_0.x(\lambda z_1.z_0(Jz_1)) \\ &=_{\lambda\beta} \lambda z_0.x(\lambda z_1.z_0(\lambda z_2.z_1(Jz_2))) \\ &=_{\lambda\beta} \dots \end{aligned}$$

https://doi.org/10.1017/S0960129508006683 Published online by Cambridge University Press

The Böhm tree of Jx is

$$BT(Jx) = \begin{array}{c} \lambda z_0.x \\ | \\ \lambda z_1.z_0 \\ | \\ \lambda z_2.z_1 \\ | \\ \ldots\ldots \end{array}$$

We write $N \approx_\eta M$ if there exists a Böhm-like tree A such that we have $BT(M) \leqslant_\eta A$ and $BT(N) \leqslant_\eta A$. For further details, see Barendregt (1984, Definition 10.2.25) and the proof of (i $\Rightarrow$ ii) in Barendregt (1984, Theorem 10.2.31). It is well known that

$$M =_{\mathscr{H}^*} N \;\Leftrightarrow\; M \approx_\eta N \quad \text{(Barendregt 1984, Theorem 19.2.9).}$$

2.3. *Graph models*

The class of graph models belongs to Scott's continuous semantics. Historically, the first graph model was Scott's P_ω, which is also known in the literature as 'the graph model'. 'Graph' was used to refer to the fact that the continuous functions were encoded in the model using (a sufficient fragment of) their graph.

Notation: For every set D, we write D^* for the set of all finite subsets of D, and $\mathscr{P}(D)$ for the powerset of D. If C is a complete partial ordering (cpo, for short), then $[C \to C]$ denotes the cpo of all Scott continuous functions from C into C.

Definition 1. A *graph model* D is a pair $(|D|, c_D)$, where $|D|$ is an infinite set, called the *web* of D, and $c_D : |D|^* \times |D| \to |D|$ is an injective total function.

We use the same notation D for the graph model and its web. Thus, for example, $\alpha \in D$ means $\alpha \in |D|$. In the following, we will use $a, b, a_1, \ldots$ to denote finite subsets of $|D|$.

Notation: We write $a \to_D \alpha$, or just $a \to \alpha$, for $c_D(a, \alpha)$. When parenthesis are omitted, association to the right should be assumed. For example, $a \to b \to \alpha$ stands for $c_D(a, c_D(b, \alpha))$. If $\overline{a} = a_1 \ldots a_n$ is a sequence of finite subsets of D, we write $\overline{a} \to \alpha$ (or $\overline{a}_n \to \alpha$ when we want to show the length of the sequence) for $a_1 \to a_2 \to \ldots \to a_n \to \alpha$.

The function c_D is useful for encoding a fragment of the graph of a Scott continuous function $f : \mathscr{P}(D) \to \mathscr{P}(D)$ as a subset $G(f)$ of D:

$$G(f) = \{a \to_D \alpha \;:\; \alpha \in f(a) \text{ and } a \in D^*\}. \tag{2}$$

Any graph model D can be used to define a model of lambda calculus through the reflexive cpo $(\mathscr{P}(D), \subseteq)$ determined by two Scott continuous mappings $G : [\mathscr{P}(D) \to \mathscr{P}(D)] \to \mathscr{P}(D)$ and $F : \mathscr{P}(D) \to [\mathscr{P}(D) \to \mathscr{P}(D)]$. The function G is defined in (2), while F is defined by

$$F(X)(Y) = \{\alpha \in D : (\exists a \subseteq Y)\; a \to_D \alpha \in X\}.$$

We will sometimes write XY for $F(X)(Y)$ and $XY_1 \ldots Y_n$ for $(\ldots((XY_1)Y_2)\ldots)Y_n)$. For more details, see Berline (2000) and Barendregt (1984, Chapter 5).

https://doi.org/10.1017/S0960129508006683 Published online by Cambridge University Press

Let Env_D be the set of D-environments ρ mapping the variables of lambda calculus into $\mathscr{P}(D)$. If $Y \subseteq D$, then the environment $\rho[x := Y]$ is defined by:

$$\begin{aligned}\rho[x := Y](x) &= Y\\ \rho[x := Y](z) &= \rho(z)\end{aligned}$$

for $z \neq x$. The interpretation M^D of a λ-term M in an environment ρ is defined by

$$\begin{aligned}x^D_\rho &= \rho(x)\\ (MN)^D_\rho &= \{\alpha \in D : (\exists a \subseteq N^D_\rho)\ a \to \alpha \in M^D_\rho\}\\ (\lambda x.M)^D_\rho &= \{a \to \alpha : \alpha \in M^D_{\rho[x:=a]}\}.\end{aligned}$$

If $\overline{x} \equiv x_1 \dots x_n$ is a sequence of variables and $\overline{a} = a_1 \dots a_n$ is a sequence of finite subsets of D, we have

$$(\lambda \overline{x}.M)^D_\rho = \{\overline{a} \to \alpha : \alpha \in M^D_{\rho[x_1:=a_1]\dots[x_n:=a_n]}\}.$$

We turn now to the interpretation of Ω in graph models. The following lemma gives a necessary condition and a sufficient condition for α to be in the interpretation of Ω in a graph model.

Lemma 2 (Baeten and Boerboom 1979). Let D be a graph model and $\alpha \in D$. Then we have:

(i) If $\alpha \in \Omega^D$, then there exists a such that $a \to \alpha \in a$.
(ii) If there exists $\beta \in D$ such that $\{\beta\} \to \alpha = \beta$, then $\alpha \in \Omega^D$.

Given a graph model D, we have $M^D = N^D$ if and only if $M^D_\rho = N^D_\rho$ for all environments ρ. The λ-theory $Th(D)$ induced by D is defined by

$$Th(D) = \{M = N : M^D = N^D\}.$$

A λ-theory induced by a graph model will be called a *graph theory*. The graph model D is called *sensible* if $Th(D)$ is a sensible λ-theory. Kerth (1998) showed that there exists a continuum of different (sensible) graph theories. It is well known that the graph theory $Th(D)$ is never extensional because $(\lambda x.x)^D \neq (\lambda xy.xy)^D$.

Di Gianantonio and Honsell (1993) showed that graph models are related to filter models (Coppo and Dezani 1980; Barendregt *et al.* 1984) since the class of graph theories is included within the class of λ-theories induced by non-extensional filter models. This inclusion is strict (Alessi *et al.* 2001), namely, there exists an equation between λ-terms, which is omitted in graph semantics, whilst it is satisfied in some non-extensional filter model.

A graph theory T will be called:

1 *the minimal graph theory* if $T \subseteq Th(D)$ for all graph models D;
2 *the minimal sensible graph theory* if T is sensible and $T \subseteq Th(D)$ for all sensible graph models D;
3 *the maximal sensible graph theory* if T is sensible and $Th(D) \subseteq T$ for all sensible graph models D.

Definition 3. A class $\mathscr{C}$ of graph models *omits* a λ-theory T if $T \neq Th(D)$ for all graph models $D \in \mathscr{C}$. More generally, $\mathscr{C}$ *omits* an equation if it fails in all models of $\mathscr{C}$, and *forces* it if it holds in all models of $\mathscr{C}$.

https://doi.org/10.1017/S0960129508006683 Published online by Cambridge University Press

If a class $\mathscr{C}$ of graph models omits an equation $M = N$, then it omits all λ-theories including $M = N$. It is easy to verify that the set of equations 'forced' by $\mathscr{C}$ constitutes a λ-theory. It is the minimal λ-theory of $\mathscr{C}$ if it is induced by a model of $\mathscr{C}$.

The completion method for building graph models from 'partial pairs' was introduced in Longo (1983) and developed on a wide scale in Kerth (1998; 2001). This method is useful for building models satisfying prescribed constraints, such as domain equations and inequations, and it is particularly convenient for dealing with the equational theories of graph models.

Definition 4. A *partial pair* A is given by a set $|A|$ and by a partial, injective function $c_A : |A|^* \times |A| \to |A|$.

Notice that the underlying set of a partial pair may be finite. As with graph models, we use the same notation A for the partial pair and its underlying set.

A partial pair is a graph model if and only if c_A is total. We always suppose that no element of A is a pair. This is not restrictive because partial pairs can be considered up to isomorphism.

Lambda terms can be interpreted in the obvious way by induction on a partial pair A. For example, we have

$$(MN)^A_\rho = \{\, \alpha \in A \;:\; (\exists a \subseteq N^A_\rho)\, [(a,\alpha) \in dom(c_A) \wedge c_A(a,\alpha) \in M^A_\rho]\}$$

and

$$(\lambda x.M)^A_\rho = \{\, c_A(a,\gamma) \in A \;:\; (a,\gamma) \in dom(c_A) \wedge \gamma \in M^A_{\rho[x:=a]}\,\}\,.$$

Definition 5. Let A be a partial pair. The *canonical completion*[†] of A is the graph model E defined by:

— $E = \bigcup_{n\in\omega} E_n$, where $E_0 = A$, $E_{n+1} = E_n \cup ((E_n^* \times E_n) - dom(c_A))$.
— Given $a \in E^*$, $\alpha \in E$,

$$c_E(a,\alpha) = \begin{cases} c_A(a,\alpha) & \text{if } c_A(a,\alpha) \text{ is defined,} \\ (a,\alpha) & \text{otherwise.} \end{cases}$$

It is easy to check that the canonical completion of a given partial pair A is actually a graph model. The canonical completion of a total pair A is equal to A.

A notion of *rank* can be naturally defined on the canonical completion E of a partial pair A. The elements of A are the elements of rank 0, while an element $\alpha \in E - A$ has rank n if $\alpha \in E_n$ and $\alpha \notin E_{n-1}$.

Classic graph models, such as Scott's P_ω (Barendregt 1984), Park's $\mathscr{P}$ (Berline 2000) and Engeler's $\mathscr{E}_B$ (Berline 2000), where B is an arbitrary non-empty set, can be viewed as the canonical completions of suitable partial pairs. In fact, P_ω, $\mathscr{P}$ and $\mathscr{E}_B$ are isomorphic to the canonical completions of

$$\begin{aligned} A &= (\{0\}, c_A) && \text{(with } c_A(\varnothing, 0) = 0\text{)} \\ D &= (\{p\}, c_D) && \text{(with } c_D(\{p\}, p) = p\text{)} \\ E &= (B, c_E) && \text{(with } c_E \text{ the empty function)}, \end{aligned}$$

respectively.

[†] Canonical completions are termed *free completions* in Berline (2006), and *Engeler completions* in Bucciarelli and Salibra (2003; 2004).

https://doi.org/10.1017/S0960129508006683 Published online by Cambridge University Press

Let $\overline{x} = x_1 \dots x_n$ be a sequence of variables and ρ be a D-environment such that $\rho(x_i)$ is a finite set.

Notation: We write $\rho(\overline{x}) \to \alpha$ for $\rho(x_1) \to \rho(x_2) \to \dots \to \rho(x_n) \to \alpha$.

3. Weak product

In this section we introduce the notion of the *weak product* of graph models, which is the main technical device used in the proof of the existence of the least (sensible) graph theory. The idea of a weak product is as follows. Given two graph models D_1 and D_2, construct the partial pair whose web is the disjoint union of the webs of D_1 and D_2, and whose coding function is the disjoint union of their coding functions. The canonical completion of this partial pair is the weak product of D_1 and D_2.

Notation: Given two sets A_1 and A_2, we write $A_1 \uplus A_2$ for their disjoint union, $in_i : A_i \to A_1 \uplus A_2$ for the canonical injections and $pr_i : 2^{A_1 \uplus A_2} \to 2^{A_i}$ for the canonical projections.

Definition 6. Let D_1 and D_2 be graph models. We define the partial pair $D_1 \uplus D_2$ by

$$\begin{aligned} |D_1 \uplus D_2| &= |D_1| \uplus |D_2| \\ c_{D_1 \uplus D_2}(b, \beta) &= \begin{cases} in_i(c_{D_i}(a, \alpha)) & \text{if } b = \{in_i(\alpha') \mid \alpha' \in a\}\,,\ \beta = in_i(\alpha), \\ \text{undefined} & \text{otherwise.} \end{cases} \end{aligned}$$

Definition 7. Let D_1 and D_2 be graph models. The graph model $D_1 \diamond D_2$, which is called the *weak product* of D_1 and D_2, is the canonical completion of the partial pair $D_1 \uplus D_2$ defined above.

These definitions can be extended to countable products by considering countable disjoint unions of webs. Countable weak products are denoted by $\diamond_{i \in \omega} D_i$.

To simplify the presentation of statements and proofs, we will suppose that when forming weak products, the factors' webs are disjoint, and that the canonical injections are replaced by set inclusions. So, for instance, if M is a λ-term and D_i is a factor of a weak product E, it makes sense to write $M^{D_i} \subseteq M^E$.

The rest of this section is devoted to the proof of the main properties of this construction:

(i) The theory of a weak product is included in the intersection of the theories of its factors (Section 3.1).
(ii) The theory of a weak product is semisensible (Section 3.2).
(iii) The inclusion in (i) is strict in general (Section 3.3).

3.1. *The theory of a weak product and its factors*

In this section we show that the theory $Th(E)$ of a weak product E is included in the theory $Th(D_i)$ of each of its factor D_i. The idea is to prove that for all closed λ-terms M

$$M^{D_i} = M^E \cap D_i\,. \tag{3}$$

This takes a structural induction on M, and hence the analysis of open terms too. Roughly speaking, we are going to show that equation (3) also holds for open terms, provided the environments satisfy a suitable closure property introduced below.

In the rest of this section, D_i will be a factor of a (finite or countable) weak product E.

https://doi.org/10.1017/S0960129508006683 Published online by Cambridge University Press

Definition 8. We define *i-flattening* by the function $f_i : E \to E$ defined by induction on the rank of $\alpha \in E$ as follows:

if $rank(\alpha) = 0$, then $f_i(\alpha) = \alpha$,

if $rank(\alpha) > 0$ and $\alpha = (a, \beta)$, then :

$$f_i(\alpha) = \begin{cases} c_{D_i}(f_i(a) \cap D_i, f_i(\beta)) & \text{if } f_i(\beta) \in D_i, \\ \alpha & \text{otherwise,} \end{cases}$$

where $f_i(a) = \{f_i(\gamma) : \gamma \in a\}$.

The following easy facts will be useful.

Fact 9.

(a) For all $\alpha \in E$, we have $f_i(\alpha) \neq \alpha$ if and only if $f_i(\alpha) \in D_i$ and $\alpha \notin D_i$.
(b) If $a \cup \{\beta\} \subseteq E$ and $f_i(\beta) \in D_i$, then $f_i(c_E(a, \beta)) \in D_i$.

Note that Fact 9(b) holds *a fortiori* if $\beta \in D_i$.

Definition 10. For $a \subseteq E$, let $\hat{a} = a \cup f_i(a)$. We say that a is *i-closed* if $\hat{a} = a$.

In other words, a is i-closed if $f_i(a) \subseteq a$.

Lemma 11. For all $a \subseteq E$, we have $\hat{a} \cap D_i = f_i(a) \cap D_i$.

Proof. By definition, $\hat{a} = a \cup f_i(a)$, hence

$$\hat{a} \cap D_i = (a \cap D_i) \cup (f_i(a) \cap D_i).$$

Since f_i restricted to D_i is the identity function, we have $a \cap D_i \subseteq f_i(a) \cap D_i$, and we are done. □

Definition 12. Let $\rho : Var \to \mathscr{P}(E)$ be an E-environment. We define the *i-restriction* ρ_i of ρ by $\rho_i(x) = \rho(x) \cap D_i$, while we say that ρ is *i-closed* if for every variable x, $\rho(x)$ is i-closed.

The following is the key technical lemma of this section.

Lemma 13. Let M be a λ-term and ρ be an i-closed E-environment. Then:

(a) M^E_ρ is i-closed.
(b) $M^E_\rho \cap D_i \subseteq M^E_{\rho_i}$.

Proof. We will prove (a) and (b) simultaneously by induction on the structure of M.

— Let $M \equiv x$.
Both (a) and (b) are trivially true.
— Let $M \equiv \lambda x.N$.

(a) Given $\alpha \in M^E_\rho$, we have to show that $f_i(\alpha) \in M^E_\rho$. First, not that if $rank(\alpha) = 0$ or if $\alpha = (a, \beta)$ and $f_i(\beta) \notin D_i$, then by Fact 9(a), we have $f_i(\alpha) = \alpha$ and we are done.

https://doi.org/10.1017/S0960129508006683 Published online by Cambridge University Press

Therefore, let $\alpha = (a, \beta)$ and $f_i(\beta) \in D_i$. We have:

$$\begin{aligned}
\alpha \in M^E_\rho &\Rightarrow \beta \in N^E_{\rho[x:=a]} && \text{by the definition of } (_)^E \\
&\Rightarrow \beta \in N^E_{\rho[x:=\hat{a}]} && \text{by monotonicity of } (_)^E \text{ with respect to environments} \\
&\Rightarrow f_i(\beta) \in N^E_{\rho[x:=\hat{a}]} && \text{by the induction hypothesis for } (a) \text{ (note that } \rho[x := \hat{a}] \text{ is } i\text{-closed)} \\
&\Rightarrow f_i(\beta) \in N^E_{(\rho[x:=\hat{a}])_i} && \text{by the induction hypothesis for } (b) \text{ since } f_i(\beta) \in D_i \\
&\Rightarrow f_i(\beta) \in N^E_{\rho_i[x:=f_i(a)\cap D_i]} && \text{by Lemma 11} \\
&\Rightarrow c_E(f_i(a) \cap D_i, f_i(\beta)) \in M^E_{\rho_i} && \text{by the definition of } M^E \\
&\Rightarrow c_{D_i}(f_i(a) \cap D_i, f_i(\beta)) \in M^E_{\rho_i} && \text{by the definition of } c_E \\
&\Rightarrow f_i(\alpha) \in M^E_{\rho_i} && \text{by the definition of } f_i \\
&\Rightarrow f_i(\alpha) \in M^E_\rho && \text{by the monotonicity of } (_)^E.
\end{aligned}$$

(b) We have:

$$\begin{aligned}
\alpha \in M^E_\rho \cap D_i &\Rightarrow (\exists a \cup \{\beta\} \subseteq D_i)\ \alpha = c_{D_i}(a, \beta) \text{ and } \beta \in N^E_{\rho[x:=a]} && \text{by the definition of } M^E \text{ and since } \alpha \in D_i \\
&\Rightarrow \beta \in N^E_{(\rho[x:=a])_i} && \text{by the induction hypothesis for } (b) \text{ (note that } \hat{a} = a) \\
&\Rightarrow \beta \in N^E_{\rho_i[x:=a]} && \text{since } a \subseteq D_i \\
&\Rightarrow \alpha \in M^E_{\rho_i} && \text{by the definition of } (_)^E.
\end{aligned}$$

— $M \equiv PQ$

(a) Let $\beta \in (PQ)^E_\rho$. If $f_i(\beta) = \beta$, we are done. Otherwise, by Fact 9(a) we have that $f_i(\beta) \in D_i$ and $\beta \notin D_i$. Moreover, by definition of interpretation in a graph model $\exists a \subseteq E$ such that $c_E(a, \beta) \in P^E_\rho$ and $a \subseteq Q^E_\rho$. Applying Fact 9(b) and the induction hypothesis, we get

$$f_i(c_E(a, \beta)) = c_{D_i}(f_i(a) \cap D_i, f_i(\beta)) = c_E(f_i(a) \cap D_i, f_i(\beta)) \in P^E_\rho .$$

Applying the induction hypothesis to Q, we get $f_i(a) \subseteq Q^E_\rho$. Hence $f_i(\beta) \in M^E_\rho$.

(b) If $\beta \in (PQ)^E_\rho \cap D_i$, then $\exists a \subseteq E$ such that $c_E(a, \beta) \in P^E_\rho$ and $a \subseteq Q^E_\rho$. Since ρ is i-closed, then by the induction hypothesis for (a), we get $f_i(c_E(a, \beta)) \in P^E_\rho$ and $f_i(a) \subseteq Q^E_\rho$. Since $\beta \in D_i$, then by Fact 9(b) we get $f_i(c_E(a, \beta)) = c_{D_i}(f_i(a) \cap D_i, \beta) \in P^E_\rho$. Now, by the induction hypothesis for (b), we obtain $c_{D_i}(f_i(a) \cap D_i, \beta) \in P^E_{\rho_i}$ and $f_i(a) \cap D_i \subseteq Q^E_{\rho_i}$, and we conclude $\beta \in (PQ)^E_{\rho_i}$. □

Lemma 14. Let M be a λ-term and $\rho : Var \to \mathscr{P}(D_i)$ be a D_i-environment. We have $M^E_\rho \cap D_i = M^{D_i}_\rho$.

Proof. We use induction on the structure of M to prove that $M^E_\rho \cap D_i \subseteq M^{D_i}_\rho$. The converse is ensured by $M^{D_i}_\rho \subseteq M^E_\rho$ and $M^{D_i}_\rho \subseteq D_i$, which are both trivially true.

— Let $M \equiv x$.

The statement holds trivially.

https://doi.org/10.1017/S0960129508006683 Published online by Cambridge University Press

— Let $M \equiv \lambda x N$.

$$\begin{aligned}
\alpha \in M^E_\rho \cap D_i &\Leftrightarrow (\exists a \cup \{\beta\} \subseteq D_i)\ \alpha = c_{D_i}(a,\beta) \text{ and } \beta \in N^E_{\rho[x:=a]} \\
&\qquad \text{by the definition of } M^E \text{ and since } \alpha \in D_i \\
&\Leftrightarrow (\exists a \cup \{\beta\} \subseteq D_i)\ \alpha = c_{D_i}(a,\beta) \text{ and } \beta \in N^{D_i}_{\rho[x:=a])} \\
&\qquad \text{by the induction hypothesis} \\
&\Leftrightarrow \alpha \in M^{D_i}_\rho \qquad \text{by definition of } M^{D_i}.
\end{aligned}$$

— Let $M \equiv PQ$.
If $\alpha \in (PQ)^E_\rho \cap D_i$, then $\exists a \subseteq E$ such that $c_E(a,\alpha) \in P^E_\rho$ and $a \subseteq Q^E_\rho$. Since ρ is i-closed and $\alpha \in D_i$, we can use Fact 9(b) and Lemma 13(a) to obtain

$$f_i(c_E(a,\alpha)) \ = c_{D_i}(f_i(a) \cap D_i, \alpha) \in P^E_\rho\,.$$

Hence we can use the induction hypothesis to get $c_{D_i}(f_i(a) \cap D_i, \alpha) \in P^{D_i}_\rho$. Moreover, using Lemma 13(a) and the induction hypothesis on Q again,

$$f_i(a) \cap D_i \subseteq Q^{D_i}_\rho\,.$$

Hence $\alpha \in (PQ)^{D_i}_\rho$.
The opposite direction is trivial. □

Theorem 15. $Th(E) \subseteq Th(D_i)$.

Proof. Let $M^E = N^E$. By the previous proposition we have

$$M^{D_i} = M^E \cap D_i = N^E \cap D_i = N^{D_i}\,.$$ □

The existence of the least, and least sensible, graph theories will be a consequence of Theorem 15 (see Section 4).

The following easy properties of weak products will be used in Section 4.2.

Proposition 16. Let $E = \diamond_{i\in I} D_i$. For all $\alpha \in E$ there exists a unique $j \in I$ such that $f_j(\alpha) \in D_j$.

Proof. The proof is by induction on the rank of α. □

Proposition 17. Let $E = \diamond_{i\in I} D_i$ and M be a closed λ-term. For all $\alpha \in M^E$ there exists a unique $j \in I$ such that $f_j(\alpha) \in M^{D_j}$.

Proof. By Proposition 16, we know that there is a unique j such that $f_j(\alpha) \in D_j$, while by Lemma 13(a) we have that $f_j(\alpha) \in M^E$. The conclusion then follows from Lemma 14. □

3.2. *The theory of a weak product is semisensible*

In this section we show that *stratified* graph models have semisensible theories. A graph model is stratified if it is the completion of a proper partial pair, that is, one whose coding function is not total. Since weak products are particular stratified graph models, the theory of a weak product is also semisensible.

The semisensibility of the theory of a stratified graph model is proved by case analysis on the order of the unsolvable terms (see Definition 24 for the definition of the order of an unsolvable

https://doi.org/10.1017/S0960129508006683 Published online by Cambridge University Press

term). The fact that unsolvables of order 0 cannot be equated to a solvable in a stratified graph model is shown in Lemma 25 using the approximation theorem below.

For unsolvable terms of finite order, we introduce the notion of the *height* of an element of the model, and then rely on the previous case (Lemma 27).

For unsolvable terms of infinite order, we rely on a general property of graph models, their non-extensionality, to show that such terms cannot be equated to solvables in any graph model (Lemma 29).

3.2.1. *An approximation theorem.* Approximation theorems are an important tool in the analysis of the λ-theories induced by models of lambda calculus. In this section we provide an approximation theorem for the class of stratified graph models: we show that the interpretation of a λ-term in a stratified graph model is the union of the interpretations of its direct approximants. This approximation theorem will be applied in Section 3.2.2 to show that the interpretation of an unsolvable term of order 0 in a stratified graph model is a set of elements of rank 0. We do not claim any particular originality for the approximation theorem we prove in this section, since it is very similar to the one in Honsell and Ronchi della Rocca (1992), and is a particular case of the approximation theorem in Bastonero and Gouy (1999). However, we will provide a proof for completeness.

Let D be a stratified graph model, which is the completion of the partial pair A. Recall that $D_0 = A$ and $D_{n+1} = D_n \cup (D_n^* \times D_n) - \mathrm{dom}\, c_A$. For every $X \subseteq D$, we use $X_{\underline{n}}$ to denote $X \cap D_n$.

The underlined natural numbers $\underline{n}$ are called *labels*. Lambda terms with occurrences of labels are called *labelled-terms*. For example, $(\lambda x.x_{\underline{n}})_{\underline{m}}y$ and $(y_{\underline{n}})_{\underline{m}}$ are labelled-terms. Note that the set of ordinary λ-terms is a proper subset of the set of labelled terms (those without any label). If N is a labelled term, we use $|N|$ to denote the λ-term obtained by erasing all labels of N. For example, we have $|(\lambda x.x_{\underline{n}})_{\underline{m}}y| = (\lambda x.x)y$.

Labelled terms are interpreted in D: the interpretation function of labelled terms is the unique extension of the interpretation function of λ-terms such that, for every labelled term M and label $\underline{n}$, we have $(M_{\underline{n}})^D_\rho = (M^D_\rho)_{\underline{n}}$.

Notation: We write $M =_{D,\rho} N$ for $M^D_\rho = N^D_\rho$ and $M \subseteq_{D,\rho} N$ for $M^D_\rho \subseteq N^D_\rho$.

An easy fact that we will use later is that, for all labelled terms M, N and environments ρ, if N is obtained by erasing some of the labels of M, we have $M \subseteq_{D,\rho} N$. In particular, for every labelled term M and environment ρ, we have $M \subseteq_{D,\rho} |M|$.

Definition 18. The *weak direct approximant* (w.a.) of a λ-term is defined by induction as follows:

$$\begin{aligned} x^w &= x \\ (\lambda x.M)^w &= \lambda x.M^w \\ (MN)^w &= M^w N^w \text{ if } MN \text{ is not a redex} \\ ((\lambda x.M)N)^w &= (\lambda x.M^w)_{\underline{0}} N^w . \end{aligned}$$

The weak direct approximant M^w of a λ-term M is a labelled term such that $|M^w| = M$. Moreover, it is easy to show that $M^w \subseteq_{D,\rho} M$ for every λ-term M and environment ρ, so we

https://doi.org/10.1017/S0960129508006683 Published online by Cambridge University Press

have

$$\bigcup\{(N^w)^D_\rho : M =_{\lambda\beta} N\} \subseteq M^D_\rho .$$

The rest of this section is devoted to proving that this inclusion is actually an equality.

Theorem 19 (The Approximation Theorem). Let D be a stratified graph model. For every λ-term M and environment ρ, we have

$$M^D_\rho = \bigcup\{(N^w)^D_\rho : M =_{\lambda\beta} N\} .$$

Proof. The proof is divided into a number of claims.

We say that a labelled-term N is *completely labelled* if every subterm of N has at least one label. For example, $((\lambda x.x_{\underline{n}})_{\underline{0}} y_{\underline{m}})_{\underline{0}}$ and $((\lambda x.x_{\underline{n}})_{\underline{0}}(y_{\underline{m}})_{\underline{0}})_{\underline{0}}$ are two completely labelled versions of the λ-term $(\lambda x.x)y$.

Claim 20. For every λ-term M and for every environment ρ we have

$$M^D_\rho = \bigcup\{N^D_\rho : N \text{ is a completely labelled term, } |N| = M\} .$$

It is sufficient to show by induction on M that, if $\alpha \in (M)^D_\rho \cap D_n$, then there is a completely labelled term N such that $|N| = M$ and $\alpha \in (N)^D_\rho$.

Claim 21. The rewriting system generated by the rules

$$(\lambda x.P)_{\underline{n+1}} Q \to_{lab} P_{\underline{n}}[x := Q_{\underline{n}}]$$
$$(P_{\underline{n}})_{\underline{m}} \to_{lab} P_{\underline{min(n,m)}}$$

is Church–Rosser and strongly normalising.

The proof is in Barendregt (1984, Section 14.1) – note that:

— if M, N are labelled terms and $M \to^*_{lab} N$, then $|M| \to^*_\beta |N|$;
— every $\to_{lab}$ reduct of a completely labelled term is completely labelled;
— the usual substitution lemma holds for labelled terms: for all labelled terms P and Q and environments ρ, we have $(P[x := Q])^D_\rho = P^D_{\rho[x:=Q^D_\rho]}$.

The next claim shows that the interpretation of a labelled term does not decrease along $\to_{lab}$ reduction paths.

Claim 22. For all labelled λ-terms P and Q and environments ρ,

$$(\lambda x.P)_{\underline{n+1}} Q \subseteq_{D,\rho} P_{\underline{n}}[x := Q_{\underline{n}}] .$$

Let $\alpha \in ((\lambda x.P)_{\underline{n+1}} Q)^D_\rho$. Then there exists $b \cup \{\alpha\} \subseteq D$ such that $b \to \alpha \in ((\lambda x.P)_{\underline{n+1}})^D_\rho$ and $b \subseteq Q^D_\rho$. Hence, $b \cup \{\alpha\} \subseteq |D|_n$, and $\alpha \in P^D_{\rho[x:=b]}$. Using these two last relations and by $b \subseteq (Q_{\underline{n}})^D_\rho$, we get $\alpha \in (P_{\underline{n}})^D_{\rho[x:=(Q_{\underline{n}})^D_\rho]}$. By the substitution lemma, we conclude that $\alpha \in P_{\underline{n}}[x := Q_{\underline{n}}]^D_\rho$.

Finally, we have the approximation theorem.

Claim 23. For all λ-terms M and environments ρ,

$$M^D_\rho = \bigcup\{(N^w)^D_\rho : M =_{\lambda\beta} N\} .$$

https://doi.org/10.1017/S0960129508006683 Published online by Cambridge University Press

Let N be a completely labelled term such that $|N| = M$. By Claim 22, we get $N \subseteq_{D,\rho} N_1$, where N_1 is the normal form of N with respect to the rewriting rules $\rightarrow_{lab}$. Since N_1 has no redexes with respect to $\rightarrow_{lab}$, and it is completely labelled, as noted above, then every redex of the λ-term $|N_1|$ should occur in N_1 as $(\lambda x.P)_{\underline{0}}Q$. Let N_2 be the λ-term obtained from N_1 by erasing all labels $n > 0$; we have $N_1 \subseteq_{D,\rho} N_2$. Finally, we get a new term N_3 by erasing from N_2 all occurrences of the label $\underline{0}$ that are not in the position $(\lambda x.P)_{\underline{0}}Q$. Note that N_3 is the direct approximant of $|N_3|$.

In conclusion, we have

$$N \subseteq_{D,\rho} N_1 \subseteq_{D,\rho} N_2 \subseteq_{D,\rho} N_3; \quad N_3 = |N_3|^w . \tag{4}$$

Moreover, as noted above, we also have

$$M \rightarrow_\beta^* |N_3| . \tag{5}$$

In conclusion,

$$\begin{array}{lll} M = & \bigcup\{N_\rho^D : N \text{ completely labelled}, |N| = M\} & \text{by Claim 20} \\ \subseteq_{D,\rho} & \bigcup\{Q^w : M =_\beta Q\} & \text{by (4) and (5)} \\ \subseteq_{D,\rho} & M & \text{as noted after Definition 18.} \end{array}$$

This concludes the proof of the approximation theorem. □

3.2.2. *The theory of a stratified graph model.* We will now apply the approximation theorem to show that stratified graph models have semisensible theories. We begin by recalling the definition of the *order* of an unsolvable λ-term.

Definition 24. An unsolvable λ-term U has:

1 order 0 if it is not β-equivalent to an abstraction term;
2 order n if $U =_{\lambda\beta} \lambda x_1 \dots x_n.T$ and T has order 0;
3 order ω if it has no finite order.

For example, Ω and Ω_3 are unsolvable of order 0, $\lambda x.\Omega$ has order 1, while $Y\,\mathbf{k}$ has order ω, where Y is any fixpoint combinator.

Lemma 25. Let D be a stratified model and U be an unsolvable term of order 0. Then, for every environment ρ, we have

$$U_\rho^D \subseteq D_0 .$$

Proof. If $N =_{\lambda\beta} U$, then N is also an unsolvable term of order 0. Therefore, we have $N \equiv (\lambda x.P)Q_1 \dots Q_m$, so $N^w \equiv (\lambda x.P^w)_{\underline{0}}Q_1^w \dots Q_m^w$. The conclusion follows from the approximation theorem because $((\lambda x.P^w)_{\underline{0}}Q_1^w \dots Q_m^w)_\rho^D \subseteq D_0$. □

An easy corollary of this lemma is that, in stratified graph models, unsolvable terms of order 0 cannot be equated to solvable terms, since the interpretation of any solvable term contains elements of arbitrary rank (see Lemma 28).

In order to deal with unsolvable terms of arbitrary order, we introduce the notion of *height* in a stratified model.

https://doi.org/10.1017/S0960129508006683 Published online by Cambridge University Press

Definition 26. Let D be a stratified model and $\alpha \in D$. We define the *height* $h(\alpha)$ of α by induction over the rank:

— If $rank(\alpha) = 0$, then $h(\alpha) = 0$;
— If $rank(\alpha) > 0$ and $\alpha = (b, \beta)$, then $h(\alpha) = 1 + h(\beta)$.

Note that whenever $\alpha = a_1 \to \ldots \to a_n \to \beta$ with $rank(\beta) = 0$, we have $h(\alpha) \leqslant n$.

Lemma 27. Let D be a stratified model and U be an unsolvable term of order n. Then for every environment ρ, we have

$$\alpha \in U_\rho^D \;\Rightarrow\; h(\alpha) \leqslant n\,.$$

Proof. By hypothesis, $U =_{\lambda\beta} \lambda x_1 \ldots x_n.T$ with T of order 0. If $\alpha \in U_\rho^D$, then $\alpha = a_1 \to \ldots \to a_n \to \beta$ and $\beta \in T_\sigma^D$, where $\sigma(x_i) = a_i$ and $\sigma(y) = \rho(y)$ for all $y \neq x_i$. By Lemma 25, we have $rank(\beta) = 0$. The conclusion then follows by the remark following Definition 26. □

We have shown that, for any unsolvable term U of finite order, the interpretation of U in a stratified graph model only contains elements whose height is no greater than the order of U.

The next lemma shows that the interpretation of any solvable terms contains elements of arbitrary height.

Lemma 28. Let D be a stratified model and $S \in \Lambda^o$ be a solvable λ-term. Then, for every natural number k, there is $\alpha \in D$ such that $\alpha \in S^D$ and $h(\alpha) \geqslant k$.

Proof. Let $S =_{\lambda\beta} \lambda x_1 \ldots x_n.x_j P_1 \ldots P_m$. It is easy to show that

$$\alpha = \varnothing^{j-1} \to (\varnothing^m \to \beta) \to \varnothing^{n-j} \to \beta \in S^D$$

for all $\beta \in D$. If we choose $h(\beta) = k$, then $h(\alpha) \geqslant h(\beta) = k$, and we get the conclusion. □

So far we have seen that in a stratified model the interpretation of an unsolvable term of finite order is different from the interpretation of any solvable term.

We will now show that unsolvable terms of infinite order cannot be consistently equated to solvable terms in graph models.

Lemma 29. Let D be a graph model, $U \in \Lambda^o$ be an unsolvable λ-term of infinite order and $S \in \Lambda^o$ be a solvable λ-term. Then

$$U^D \neq S^D\,.$$

Proof. In order to show a contradiction, assume that $U^D = S^D$. Since S is solvable, there exist λ-terms $M_1, \ldots, M_k$ such that $SM_1 \ldots M_k =_{\lambda\beta} x$ for an arbitrary variable x. Then, for any environment ρ, we have

$$x =_{D,\rho} UM_1 \ldots M_k\,.$$

Since U is an unsolvable term of infinite order, we have that $UM_1 \ldots M_k$ is also an unsolvable term of infinite order. This implies that $UM_1 \ldots M_k =_{\lambda\beta} \lambda y.T$ for suitable y and T. However, the equation $x = \lambda y.T$ does not hold in any graph model: consider an environment ρ such that $\rho(x) = \{a \to \alpha\}$ for given finite a and $\alpha \in D$. Hence $(\lambda y.T)_\rho^D = \{a \to \alpha\}$. However, this is impossible because, for all finite $b \subseteq D$, we have $(b \cup a) \to \alpha \in (\lambda y.T)_\rho^D$, which gives a contradiction. □

https://doi.org/10.1017/S0960129508006683 Published online by Cambridge University Press

Summing up, we have proved the following result.

Theorem 30. The theory of any stratified graph model is semisensible.

Corollary 31. The theory of any weak product is semisensible.

3.3. *Self weak product*

Theorem 15 states that the theory of a weak product is included in the intersection of those of its factors. In this section we show that this inclusion is strict in general. Moreover, in Theorem 33 below we show that self weak products do not, in general, preserve equations between unsolvable terms. Hence it is not in general true that $Th(D \diamond D) = Th(D)$, whenever $Th(D)$ is semisensible.

Proposition 32. Let D be a graph model satisfying the equation $\Omega = \mathbf{i}$. The model $D \diamond D$, which we call the *self weak product* of D, does not satisfy $\Omega = \mathbf{i}$.

Proof. By Corollary 31, the theory of $D \diamond D$ is semisensible. □

Theorem 33. There exists a graph model D satisfying the following two conditions:

(i) $D \models \Omega = \lambda x.\Omega$.
(ii) $D \diamond D \not\models \Omega = \lambda x.\Omega$.

Proof. The proof is divided into a number of claims. For clarity, we use D_1 to denote the first copy of D in $D \diamond D$ and D_2 to denote the second copy. Moreover, we assume that these (isomorphic) copies are disjoint.

Recall that every weak product is a stratified graph model.

Claim 34. Let D be a graph model and $E \equiv D \diamond D$ be the self weak product of D. Then we have

$$\Omega^D = \varnothing \iff E \models \Omega = \lambda x.\Omega\,.$$

($\Leftarrow$) Assume $\Omega^D \neq \varnothing$. By Lemma 14 we have that $\Omega^{D_1} = \Omega^E \cap D_1$. Thus the hypothesis implies $\Omega^E \neq \varnothing$. Let $\beta \in \Omega^E$ be an arbitrary element and let $b \subseteq E$ be a finite set containing elements of rank 1. Then $b \to \beta \in (\lambda x.\Omega)^E = \{a \to \alpha : \alpha \in \Omega^E\}$. Finally, by Lemma 25, we have that $\Omega^E \subseteq E_0$ is a set of elements of rank 0, while $(\lambda x.\Omega)^E$ contains elements of rank greater than 0. We get the conclusion $E \not\models \Omega = \lambda x.\Omega$.

($\Rightarrow$) The conclusion follows from the following relations:

$$\begin{aligned} \Omega^E &\subseteq E_0 && && \text{(see Lemma 25)} \\ E_0 &= D_1 \cup D_2 \\ \varnothing = \Omega^{D_i} &= \Omega^E \cap D_i && (i = 1, 2) && \text{(see Lemma 14).} \end{aligned}$$

This concludes the proof of Claim 34.

Claim 35. There exists a graph model D satisfying the following two conditions:

1 $D \models \Omega = \lambda x.\Omega$.

2 $\Omega^D \neq \varnothing$.

https://doi.org/10.1017/S0960129508006683 Published online by Cambridge University Press

We construct a graph model by using the technique of forcing introduced in Baeten and Boerboom (1979). In the following proof we follow Berline and Salibra (2006).

Let D be any infinite countable set. We are going to define 'forcing' by the injective total function $c_D : D^* \times D \to D$.

We fix an enumeration of D and an enumeration of $D^* \times D$. Let p be the first element in the enumeration of D.

We are going to build an infinite sequence of elements $\alpha_n \in D \cup \{v\}$ $(n \geqslant 0)$, where v is some new element, and an infinite sequence of partial pairs A_n $(n \geqslant 1)$ such that $|A_n|$ is a finite set and $c_{A_n} \subseteq c_{A_{n+1}}$ (that is, the graph of c_{A_n} is contained within the graph of $c_{A_{n+1}}$). D becomes a graph model by defining $c_D =_{def} \cup_{n\in\omega} c_{A_n}$.

We start from $|A_1| = \{p\}$, $c_{A_1}(\{p\}, p) = p$ and $\alpha_0 = p$ (note that the canonical completion of the partial pair A_1 is Park's model (see Section 2.3)). It is not difficult to verify that $\Omega^{A_1} = \{p\} = (\lambda x.\Omega)^{A_1}$ (recall that the interpretation of a λ-term in a partial pair was defined in Section 2.3).

Assume that the partial pair A_n and $\alpha_0, \ldots, \alpha_{n-1}$ have been built.

Let α_n be the first element of $(\lambda x.\Omega)^{A_n} - \{\alpha_0, \ldots, \alpha_{n-1}\}$ if this set is non-empty, and v otherwise.

Let (b_n, δ_n) be the first element in $D^* \times D \; - \; dom(c_{A_n})$ and γ_n be the first element in $D \; - \; (range(c_{A_n}) \cup b_n)$.

Case 1. $\alpha_n = v$. Then $|A_{n+1}| = |A_n| \cup b_n \cup \{\delta_n, \gamma_n\}$ and $c_{A_{n+1}}$ is a proper extension of c_{A_n} defined as follows in the new pair (b_n, δ_n):

$$c_{A_{n+1}}(b_n, \delta_n) = \gamma_n \,.$$

Case 2. $\alpha_n \in D$. Then $|A_{n+1}| = |A_n| \cup b_n \cup \{\delta_n, \gamma_n, \beta_n, \alpha_n\}$ and $c_{A_{n+1}}$ is a proper extension of c_{A_n} defined as follows in the new pairs (b_n, δ_n) and $(\{\beta_n\}, \alpha_n)$:

$$\begin{aligned} c_{A_{n+1}}(b_n, \delta_n) &= \gamma_n \\ c_{A_{n+1}}(\{\beta_n\}, \alpha_n) &= \beta_n, \end{aligned}$$

where β_n is the first element of D such that

$$\begin{aligned} \big(\{\beta_n\}, \alpha_n\big) \;&\in\; D^* \times D \;-\; (dom(c_{A_n}) \cup \{(b_n, \delta_n)\}) \text{ and} \\ \beta_n \;&\in\; D \;-\; (range(c_{A_n}) \cup \{\gamma_n\}) \,. \end{aligned}$$

It is clear that c_{A_n} is a strictly increasing sequence of well-defined partial injective maps and that $c_D = \cup c_{A_n}$ is total.

We still need to show that the graph model D satisfies the equation $\Omega = \lambda x.\Omega = B$, where $B =_{def} \{\, \alpha_n \;:\; n \in \omega \,\} \cap D$.

— $B \subseteq (\lambda x.\Omega)^D$

This follows from $\alpha_0 = p \in (\lambda x.\Omega)^{A_1}$, the definition of α_n $(n > 0)$, and the fact that $(\lambda x.\Omega)^{A_n} \subseteq (\lambda x.\Omega)^D$.

— $(\lambda x.\Omega)^D \subseteq B$

Suppose $\gamma \in (\lambda x.\Omega)^D$. Then $\gamma \in (\lambda x.\Omega)^{A_m}$ for some m (and for all the larger ones). If $\gamma \notin B$, then, for all $n \geqslant m$, we have $\alpha_n \neq v$ (that is, $\alpha_n \in D$) is smaller than γ in the enumeration of D, which contradicts the fact that there is only a finite number of such elements.

https://doi.org/10.1017/S0960129508006683 Published online by Cambridge University Press

— $B \subseteq \Omega^D$
  $\alpha_n \in \Omega^D$ follows immediately from $\alpha_0 = p \in \Omega^{A_1} \subseteq \Omega^D$, the fact that $c_D(\{\beta_n\}, \alpha_n) = \beta$ and Lemma 2.
— $\Omega^D \subseteq B$
  If $\varepsilon \in \Omega^D$, there is an $a \in D^*$ such that $c_D(a, \varepsilon) \in a$ (by Lemma 2). Since $c_D = \cup c_{A_n}$, we have either $\varepsilon = \gamma_n$ or $\varepsilon = \alpha_n$ for some n. Because of the choices of the γ_n, the first of these is impossible.

This concludes the proof of Claim 35.

The conclusion of the theorem is now a simple corollary of Claims 34 and 35. □

Corollary 36. There exist graph models D satisfying the following condition:

$$Th(D \diamond D) \neq Th(D) \cap T, \quad \text{for every sensible } \lambda\text{-theory } T\,.$$

4. Weak product and graph theories

In this section we show the existence of a minimal graph theory and a minimal sensible graph theory. The main technical device is the weak product introduced in the previous section.

4.1. *The minimal graph theory*

Let I be the set of equations between λ-terms that fail to hold in some graph model. For every equation $e \in I$, we consider a fixed graph model D_e, where the equation e fails to hold.

Then we consider the weak product $E = \diamond_{e \in I} D_e$.

By Theorem 15, $Th(E) \subseteq Th(D_e)$ for all $e \in I$. In particular, $e \notin Th(E)$ for all $e \in I$. So we have the following theorem.

Theorem 37. The theory of the graph model E is the minimal graph theory.

4.2. *The minimal sensible graph theory*

We proceed as before. Let I_s be the set of equations that fail to hold in some sensible graph model. For every $e \in I_s$, let D_e be a sensible graph model where the equation e fails to hold.

Then we consider the weak product $E_s = \diamond_{e \in I} D_e$.

By Theorem 15 the theory $Th(E_s)$ is contained within any sensible graph theory. So if $Th(E_s)$ is sensible, we are done.

In the remaining part of this section we show that $Th(E_s)$ is indeed sensible.

The proof of the following lemma can be found in Kerth (1995, Example 5.3.7).

Lemma 38 (Kerth 1995). Let D be a graph model. If $\alpha \in (\Omega_3)^D$, there exists a natural number $k \geqslant 1$ such that

$$\alpha = b_1 \to \ldots \to b_k \to \alpha$$

for suitable finite subsets b_i contained in the interpretation of $\lambda x.xxx$.

Lemma 39. If all closed unsolvable λ-terms have the same interpretation in a graph model, it must be the empty set.

https://doi.org/10.1017/S0960129508006683 Published online by Cambridge University Press

Proof. Let D be a graph model and let X be a non-empty subset of D that is the common interpretation of all closed unsolvable terms. Since Ω and $\lambda x.\Omega$ are both unsolvables, we have

$$X = (\lambda x.\Omega)^D = \{a \to \alpha : \alpha \in \Omega^D\} = \{a \to \alpha : \alpha \in X\}. \tag{6}$$

It follows that $a \to \alpha \in X$ for all finite subsets a of D and all $\alpha \in X$. Let γ be an element of X. Then $a \to \gamma \in (\Omega_3)^D$ by (6), since Ω_3 is unsolvable and $(\Omega_3)^D = X$. From Lemma 38 it follows that

$$a \to \gamma = b_1 \to \ldots \to b_k \to a \to \gamma\,,$$

where $b_1, \ldots, b_k$ are finite subsets contained in the interpretation of $\lambda x.xxx$. It follows that $b_1 = a$. By the arbitrariness of a we can conclude that $(\lambda x.xxx)^D = D$. But this is impossible, because, for example, $\varnothing \to \beta \notin (\lambda x.xxx)^D$. □

Theorem 40. The theory of E_s is the minimal sensible graph theory.

Proof. By construction, $Th(E_s)$ is contained within any sensible graph theory. In order to prove that $Th(E_s)$ is sensible, let us suppose that a closed unsolvable term U has a non-empty interpretation in E_s, that is, there exists $\alpha \in U^{E_s}$. By Proposition 16, there exists a unique $e \in I_s$ such that $f_e(\alpha) \in D_e$. By Lemma 13(a), we have that $f_e(\alpha) \in U^{E_s}$, and, finally, by Lemma 13(b), that $f_e(\alpha) \in U^{D_e}$. But since D_e is sensible, this is impossible by Lemma 39. Hence $U^{E_s} = \varnothing$ for any closed unsolvable U (and actually for any unsolvable term in any environment). □

5. The minimal graph theory is not $\lambda\beta$

A longstanding open problem is whether there exists a non-syntactic model of lambda calculus whose equational theory is equal to the least λ-theory $\lambda\beta$. In Theorem 42 below we show that this model cannot be found within graph semantics. This result gives a negative answer to Question 1 in Berline (2000, Section 6.2) for the class of graph models.

We start with a lemma.

Lemma 41. All graph models satisfy the inequality $\Omega_3 \leqslant \lambda y.\Omega_3 y$.

Proof. Let D be an arbitrary graph model and $\alpha \in (\Omega_3)^D$. From Lemma 38 it follows that there exists a natural number $k \geqslant 1$ such that $\alpha = b_1 \to b_2 \to \ldots \to b_k \to \alpha$ for suitable finite subsets b_i contained in the interpretation of $\lambda x.xxx$. We have that $\alpha = b_1 \to b_2 \to \ldots \to b_k \to \alpha \in (\lambda y.\Omega_3 y)^D$ if and only if there exists a finite set d such that $d \to b_2 \to \ldots \to b_k \to \alpha \in (\Omega_3)^D$ and $d \subseteq b_1$. This last relation is true by defining $d \equiv b_1$, so $\alpha \in (\lambda y.\Omega_3 y)^D$. In conclusion, we get $(\Omega_3)^D \subseteq (\lambda y.\Omega_3 y)^D$. □

We would like to point out here that Berline *et al.* have recently shown (Berline *et al.* 2007) that the order theory of a graph model is never recursively enumerable.

Theorem 42. There exists no graph model whose equational theory is $\lambda\beta$.

Proof. Assume that there exists a graph model D whose equational theory is $\lambda\beta$. Now, by Selinger (2003, Corollary 2.4), the denotations of two non-$\lambda\beta$-equivalent closed λ-terms must be incomparable in every model of lambda calculus whose equational theory is $\lambda\beta$. Then, for all

https://doi.org/10.1017/S0960129508006683 Published online by Cambridge University Press

closed λ-terms M and N such that $M \neq_{\lambda\beta} N$, we have that neither $M^D \subseteq N^D$ nor $N^D \subseteq M^D$. Hence, we get a contradiction because of Lemma 41. □

In Theorem 37, we have shown that there exists a minimal graph theory. By Theorem 42, we have that $\lambda\beta$ is strictly included within the minimal graph theory. Thus, there exist equations between non-$\lambda\beta$-equivalent terms satisfied by all graph models. In Theorem 44, whose proof is based on technical results that can be found in Selinger (2003), we will characterise an equation of this kind.

Let f be any λ-term satisfying, through a fixpoint combinator, the recursive equation $fxy =_{\lambda\beta} fx(fx(fxy))$ for variables x, y (in other words, any three applications of fx are equivalent to a single application) and let $A \equiv \lambda xyzwv.fx(fy(fz(fwv)))$. The λ-terms f and A were defined in Selinger (2003). Selinger (2003, Proposition 2.1) showed that

$$Axxxy =_{\lambda\beta} Axyyy \tag{7}$$

and

$$Axxxy \neq_{\lambda\beta} Axxyy\,. \tag{8}$$

This inequality has an ingenious proof based on the notion of a finite lambda reduction model.

For completeness, we recall Selinger (2003, Lemma 2.2), which will be used in the proof of Theorem 44.

Lemma 43 (Selinger 2003). Let $P_1, \ldots, P_n$ be λ-terms that are distinct in $\lambda\beta$, and let x be a variable not free in $P_1, \ldots, P_n$. Then for all terms M, N for which x is not free in M and N, and for all variables $y_1, \ldots, y_n$, we have

$$M(xP_1)(xP_2)\ldots(xP_n) =_{\lambda\beta} N(xP_1)(xP_2)\ldots(xP_n) \;\Rightarrow\; My_1y_2\ldots y_n =_{\lambda\beta} Ny_1y_2\ldots y_n\,.$$

Notation: Let $t \equiv \Omega_3$ and $u \equiv \lambda y.\Omega_3 y$ in the following theorem.

Theorem 44. Let T be the minimal graph theory (whose existence was shown in Theorem 37). Then, for a variable x, we have

$$A(xt)(xt)(xt)(xu) =_T A(xt)(xt)(xu)(xu)\,, \tag{9}$$

while

$$A(xt)(xt)(xt)(xu) \neq_{\lambda\beta} A(xt)(xt)(xu)(xu)\,. \tag{10}$$

Proof. By compatibility, the fact that $t \leqslant u$ (see Lemma 41) and (7), we get that the following relations hold in every graph model:

$$\begin{aligned} A(xt)(xt)(xt)(xu) &\leqslant A(xt)(xt)(xu)(xu) \\ &\leqslant A(xt)(xu)(xu)(xu) \\ &=_{\lambda\beta} A(xt)(xt)(xt)(xu)\,. \end{aligned}$$

Equation (9) then follows easily.

We still have to show the inequality (10). In order to show a contradiction, we assume the opposite:

$$A(xt)(xt)(xt)(xu) =_{\lambda\beta} A(xt)(xt)(xu)(xu)\,.$$

https://doi.org/10.1017/S0960129508006683 Published online by Cambridge University Press

We can apply the hypotheses of Lemma 43 to $M \equiv \lambda xy.Axxxy$, $N \equiv \lambda xy.Axxyy$, $P_1 \equiv t$ and $P_2 \equiv u$. Then we get the conclusion of Lemma 43: $My_1y_2 =_{\lambda\beta} Ny_1y_2$, which implies $Ay_1y_1y_1y_2 =_{\lambda\beta} Ay_1y_1y_2y_2$, which contradicts (8). □

6. Omitting equations and theories

In this section we prove the main results of the paper:

— The λ-theory $\mathscr{B}$ of Böhm trees is the greatest sensible graph theory.
— Graph semantics omits all equations $M = N$ between solvable λ-terms that have distinct Böhm trees.

The following two theorems are the main results of the paper. The proof of Theorem 45 is postponed to Section 8.

Theorem 45. The graph semantics omits all equations $M = N$ satisfying the following conditions:

$$M =_{\mathscr{H}^*} N \text{ and } M \neq_{\mathscr{B}} N\,. \tag{11}$$

The above result shows the intrinsic non-extensionality of graph models. In particular, any equation $M = N$ between λ-terms that are $\lambda\beta\eta$-equivalent but do not have the same Böhm tree fails in every graph model. This is no longer true if M and N are $\lambda\beta\eta$-equivalent and the η-redexes in M and N are pushed to infinity. For example, let M be any λ-term such that $Mz =_{\lambda\beta} \lambda x.x(Mz)$. Then the Böhm tree of $M\mathbf{1}$ does not contain any η-redex, because $\mathbf{1}$ is pushed to infinity when we generate its Böhm tree:

$$M\mathbf{1} =_{\lambda\beta} \lambda x.x(M\mathbf{1}) =_{\lambda\beta} \lambda x.x(\lambda x.x(M\mathbf{1})) =_{\lambda\beta} \ldots \lambda x.x(\lambda x.x(\lambda x.x(\lambda x.x(\ldots)))).$$

This is why the equation $M\mathbf{i} = M\mathbf{1}$ is true, for example, in Scott's graph model P_ω (Barendregt 1984, Section 19.1).

Theorem 46. The λ-theory $\mathscr{B}$ is the unique maximal sensible graph theory.

Proof. $\mathscr{B}$ is the equational theory of Scott's graph model P_ω (Barendregt 1984, Section 19.1) and of Engeler's graph model $\mathscr{E}_A$ (Berline 2000). Let T be a sensible graph theory and suppose $M =_T N$. We have $M =_{\mathscr{H}^*} N$ because $\mathscr{H}^*$ is the unique maximal sensible λ-theory. Since graph semantics does not omit the equation $M = N$, we have from $M =_{\mathscr{H}^*} N$ and Theorem 45 that $M =_{\mathscr{B}} N$, so $T \subseteq \mathscr{B}$. □

It is well known that every graph theory is non-extensional. Note that Theorem 46 is non-trivial, because there exist non-extensional sensible λ-theories that strictly include $\mathscr{B}$ (Barendregt 1984, Exercise 16.5.5).

Berline (2000) asked whether there is a non-syntactic sensible model of lambda calculus whose theory is strictly included in $\mathscr{B}$. The answer is yes, as shown by the following corollary.

Theorem 47. There exists a continuum of different sensible graph theories strictly included in $\mathscr{B}$.

https://doi.org/10.1017/S0960129508006683 Published online by Cambridge University Press

Proof. The proof is based on a syntactic difficult result (which was conjectured in Kerth (1998) and proved in David (2001)). Kerth (1998) showed that there exists a continuum of sensible graph theories. The conclusion then follows from Theorem 46. □

It is well known that the λ-term Ω is easy, that is, it can be consistently equated to every other closed λ-term M. We use $(\Omega = M)^+$ to denote the λ-theory generated by the equation $\Omega = M$.

Theorem 48. Let M be an arbitrary closed λ-term. Then we have

$$P =_{\mathscr{H}^*} Q,\ P \neq_{\mathscr{B}} Q \ \Rightarrow\ (\Omega = M)^+ \nvdash P = Q\,.$$

In other words, $(\Omega = M)^+ \cap \mathscr{H}^* \subseteq \mathscr{B}$.

Proof. From Baeten and Boerboom (1979), the λ-theory $(\Omega = M)^+$ is contained within a graph theory. The conclusion then follows from Theorem 45. □

7. Incompatible sets

In the proof of the main theorem we are going to use families of points of a graph mode that are not just pairwise distinct, but also 'incompatible' in the sense expressed by Definition 49 below. Then, in Lemma 53 we show that such families actually exist in all graph models.

Let D be a graph model and $\alpha, \beta \in D$ and $m \geqslant 1$. We say that

— β *m-constructs* α if there exists a sequence $\overline{a} = a_1 \dots a_m$ of finite subsets of D such that $\alpha = a_1 \to \dots \to a_m \to \beta$.
— β *constructs* α if β m-constructs α for some m.

Let $P \equiv yP_1 \dots P_m$ be a λ-term and ρ be a D-environment. Given $\beta \in D$, we want to check whether $\beta \in (yP_1 \dots P_m)^D_\rho = \rho(y)(P_1)^D_\rho \dots (P_m)^D_\rho$. In general, we have to consider all elements $\alpha \in \rho(y)$ such that β m-constructs α. The aim of the definition of an incompatible set is to reduce these elements to just one, say $a_1 \to \dots a_m \to \beta$. In this case, in order to check that $\beta \notin (yP_1 \dots P_m)^D_\rho$, it would be sufficient to find an i such that $a_i \notin (P_i)^D_\rho$.

Definition 49. An infinite subset X of D is called *incompatible* if, for all $\alpha, \beta \in X$, we have that β does not construct α (note that α may be equal to β).

Before proving that incompatible sets do indeed exist, we will explain how they will be used later.

Lemma 50. Let X be an incompatible set. For every $\alpha \in D$, there exists at most one element β of X such that β constructs α.

We use $q_X : D \to X$ to denote the partial function such that $q_X(\alpha)$ is the unique element of X that constructs α when this element exists.

Definition 51. Let X be an incompatible set. A subset Y of D is called *X-good* if the function $q_X : D \to X$ restricted to Y is total and injective.

Lemma 52. Let X be an incompatible set and $P \equiv yP_1 \dots P_m$ be a λ-term. Let ρ be a D-environment. If $\rho(y)$ is X-good, then the following two conditions are equivalent for all $\alpha \in \rho(y)$:

https://doi.org/10.1017/S0960129508006683 Published online by Cambridge University Press

1 $q_X(\alpha) \in \rho(y)(P_1)^D_\rho \dots (P_m)^D_\rho$.
2 $q_X(\alpha) \in \{\alpha\}(P_1)^D_\rho \dots (P_m)^D_\rho$.

Proof. $q_X(\alpha) \in \rho(y)(P_1)^D_\rho \dots (P_m)^D_\rho$ if and only if there exists $\gamma \in \rho(y)$ such that

$$\gamma \equiv a_1 \to \dots \to a_m \to q_X(\alpha) \in \rho(y)$$

and $a_i \subseteq (P_i)^D_\rho$. Since $\rho(y)$ is X-good, by Lemma 50, $\gamma = \alpha$ is the only possibility. □

As a consequence of the above lemma, in order to prove that $q_X(\alpha) \notin (yP_1 \dots P_m)^D_\rho$, it is sufficient to find an i such that $a_i \nsubseteq (P_i)^D_\rho$. This is why we introduced the notion of incompatible sets.

Lemma 53. There exists an incompatible set.

Proof. Let D be a graph model. Given $\alpha \in D$, we define the *degree* of α as the least natural number $k > 0$ such that there exist finite subsets $b_1, \dots, b_k$ of D satisfying $\alpha = b_1 \to \dots \to b_k \to \alpha$. If such a natural number does not exist, we say that the degree of α is infinite.

We now show that there exists an element of D whose degree is infinite.

In order to show a contradiction, assume that every element of D has a finite degree. Let α be an arbitrary element of D. Then there exists $b_1, \dots, b_k$ ($k \geqslant 1$) such that $\alpha = b_1 \to \dots \to b_k \to \alpha$. Define $\beta_i \equiv b_i \to \dots \to b_k \to \alpha$ ($i = 1, \dots, k$). Then $\alpha = b_1 \to \dots \to b_{i-1} \to \beta_i$. In particular, $\alpha = b_1 \to \beta_2$. Since there exist infinite elements of D^* distinct from b_1, we consider the family of elements $c \to \beta_2$ with $c \neq b_1$. Let c be such that $c \to \beta_2 \neq \beta_1, \dots, \beta_k$. Then $c \to \beta_2$ cannot have a finite degree.

To conclude the proof, let $\alpha \in D$ whose degree is infinite. Given a family $\{a_n\}_{n\in\omega}$ of pairwise distinct, finite subsets of D, define $\beta_n = a_n \to \alpha$ ($n \geqslant 0$). We prove that the set $\{\beta_n : n \geqslant 0\}$ is an incompatible set. In order to show a contradiction, assume that $\beta_i = b_1 \to \dots \to b_t \to \beta_j$ for some i and j, that is,

$$a_i \to \alpha = b_1 \to \dots \to b_t \to a_j \to \alpha\,.$$

It follows that $\alpha = b_2 \to \dots \to b_t \to a_j \to \alpha$. We then get a contradiction because the degree of α is infinite. □

7.1. *Incompatible sets at work*

In the following we assume X is a fixed incompatible set.

Notation: We define $\overline{\varnothing}_n \to \alpha$ as follows:

$$\overline{\varnothing}_0 \to \alpha \equiv \alpha; \quad \overline{\varnothing}_{n+1} \to \alpha \equiv \varnothing \to (\overline{\varnothing}_n \to \alpha)\,.$$

Recall that if ρ is an environment and $\overline{x} \equiv x_1 \dots x_n$ is a sequence of variables, we write $\rho(\overline{x}) \to \alpha$ for $\rho(x_1) \to \dots \to \rho(x_n) \to \alpha$. Note that we implicitly assume that $\rho(x_i)$ is a finite set for all i.

In the following lemma we show how to separate two solvable λ-terms in a graph model.

Lemma 54. Let D be a graph model and X be an incompatible set. Also, let:
— $S \equiv \lambda\overline{x}.z\overline{S}$, where $\overline{S} \equiv S_1 \dots S_t$;

https://doi.org/10.1017/S0960129508006683 Published online by Cambridge University Press

— $T \equiv \lambda\overline{x}\lambda\overline{y}.z\,\overline{T}\,\overline{Q}$, where $\overline{y} = y_1 \dots y_u$ ($u \geqslant 0$), $\overline{T} \equiv T_1 \dots T_t$ and $\overline{Q} \equiv Q_1 \dots Q_r$ ($r \geqslant 1$);
— $\delta \equiv \overline{\varnothing}_{t+r} \to \beta$ and $\alpha \equiv \rho(\overline{x}) \to \rho(\overline{y}) \to \beta$ with $\beta \in X$;
— ρ and σ be environments such that $\rho(z), \sigma(z)$ are X-good (z is the head variable of S), $\delta \in \rho(z) \cap \sigma(z)$, and $\rho(y_i) \neq \varnothing$ for some $1 \leqslant i \leqslant u$.

Then we have $\alpha \in T_\sigma^D$ and $\alpha \notin S_\sigma^D$.

Proof. We define $\tau \equiv \sigma[\overline{x} := \rho(\overline{x})][\overline{y} := \rho(\overline{y})]$. Note that $\tau(z) = \rho(z)$ if $z \in \overline{xy}$, and $\tau(z) = \sigma(z)$ otherwise. We have

$$\begin{array}{lll}
\alpha \in T_\sigma^D & \text{iff} & \beta \in z_\tau^D\,\overline{T}_\tau^D\,\overline{Q}_\tau^D \\
& \text{iff} & \beta \in \tau(z)\,\overline{T}_\tau^D\,\overline{Q}_\tau^D \\
& \text{iff} & \beta \in \{\delta\}\overline{T}_\tau^D\overline{Q}_\tau^D \qquad \text{by } \delta \in \rho(z) \cap \sigma(z) \text{ and Lemma 52} \\
& \text{iff} & \beta \in \{\delta\}\overline{\varnothing}_{t+r} \qquad \text{by the definition } \delta.
\end{array}$$

Since this last condition is true, we have $\alpha \in T_\sigma^D$.

Let $\tau' \equiv \sigma[\overline{x} := \rho(\overline{x})]$. We now have

$$\begin{array}{lll}
\alpha \in S_\sigma^D & \text{iff} & \rho(\overline{y}) \to \beta \in z_{\tau'}^D\overline{S}_{\tau'}^D \\
& \text{iff} & \rho(\overline{y}) \to \beta \in \{\delta\}\overline{S}_{\tau'}^D \qquad \text{by } \delta \in \rho(z) \cap \sigma(z) \text{ and Lemma 52} \\
& \text{iff} & \rho(\overline{y}) \to \beta \in \{\delta\}\overline{\varnothing}_t \qquad \text{by the definition of } \delta \text{ and } \overline{S} = S_1 \dots S_t \\
& \text{iff} & \rho(\overline{y}) \to \beta = \overline{\varnothing}_r \to \beta \qquad \text{by the definition of } \delta.
\end{array}$$

Since $\beta \in X$ (and thus β does not construct β), this last condition holds if and only if $u = r$, where $\overline{y} = y_1 \dots y_u$, and $\rho(y_j) = \varnothing$ for all $1 \leqslant j \leqslant r$. By hypothesis, $(\exists i)\ \rho(y_i) \neq \varnothing$, so $\alpha \notin S_\sigma^D$. □

Lemma 55. Let D be a graph model and X be an incompatible set. Also, let:

— $S \equiv \lambda\overline{x}.z\overline{S}$, where $\overline{S} \equiv S_1 \dots S_t$;
— $\delta = \overline{\varnothing}_{i-1} \to \{\alpha'\} \to \overline{\varnothing}_{t-i} \to \beta$ for some $i \leqslant t$ and $\beta \in X$;
— ρ and σ be environments such that $\rho(z)$ and $\sigma(z)$ are X-good (z is the head variable of S) and $\delta \in \rho(z) \cap \sigma(z)$;
— $\alpha = \rho(\overline{x}) \to \beta$.

Then we have

$$\alpha \in S_\sigma^D \;\Leftrightarrow\; \alpha' \in (S_i)^D_{\sigma[\overline{x}:=\rho(\overline{x})]}\,.$$

Proof. Let $\tau \equiv \sigma[\overline{x} := \rho(\overline{x})]$. Note that $\tau(z) = \rho(z)$ if $z \in \overline{x}$, and $\tau(z) = \sigma(z)$ otherwise. So we have

$$\begin{array}{lll}
\alpha \in S_\sigma^D & \text{iff} & \beta \in z_\tau^D\overline{S}_\tau^D \qquad \text{by the definition of } \alpha \\
& \text{iff} & \beta \in \{\delta\}\overline{S}_\tau^D \qquad \text{by } \delta \in \sigma(z) \cap \rho(z) \text{ and Lemma 52} \\
& \text{iff} & \beta \in \{\delta\}\overline{\varnothing}_{i-1}(S_i)_\tau^D\overline{\varnothing}_{t-i} \qquad \text{by the definition of } \delta \\
& \text{iff} & \alpha' \in (S_i)_\tau^D \qquad \text{by the definition of } \delta.
\end{array}$$

The conclusion of the lemma is now immediate. □

8. The proof of the main theorem

In this section we provide the proof of Theorem 45.

https://doi.org/10.1017/S0960129508006683 Published online by Cambridge University Press

Recall that a node of a tree is a sequence of natural numbers and that the level of a node is the length of the sequence. We will use ε to denote the empty sequence.

Let M, N be closed λ-terms such that $M =_{\mathscr{H}^*} N$ and $M \neq_{\mathscr{B}} N$. This last condition expresses the fact that the Böhm tree $BT(M)$ of M is different from the Böhm tree $BT(N)$ of N. The first condition $M =_{\mathscr{H}^*} N$ implies that if a given node has the same label in $BT(M)$ and $BT(N)$, then this node has the same number of sons in $BT(M)$ and $BT(N)$.

We will now introduce some notation and definitions needed in the proof.

Let $u = r_1 \dots r_k$ be a node at least level where the labels of $BT(M)$ and $BT(N)$ are different. The sequence ε, r_1, r_1r_2, $r_1r_2r_3, \dots, r_1 \dots r_k$ is the sequence of nodes that are in the path from the root ε to u. These nodes will be denoted by $u_0, u_1, u_2, \dots, u_k$. Then, for example, $u_0 = \varepsilon$, $u_2 = r_1r_2$ and $u_k = u$. From the hypothesis of minimality of u it follows that:

(i) The label of the node u_j $(0 \leqslant j < k)$ in the Böhm tree of M is equal to the corresponding one in the Böhm tree of N.
(ii) The labels of the node u in $BT(M)$ and $BT(N)$ are different.

We define two sequences M_{u_j} and N_{u_j} $(0 \leqslant j \leqslant k)$ of λ-terms whose Böhm trees $BT(M_{u_j})$ and $BT(N_{u_j})$ are the subtrees of $BT(M)$ and $BT(N)$ at root u_j, respectively. Without loss of generality, we may assume that M and N are in head normal form (hnf, for short). Then we define

$$\begin{aligned} M_{u_0} &\equiv M \\ N_{u_0} &\equiv N\,. \end{aligned}$$

If $k = 0$, we have finished. Otherwise, assume by the induction hypothesis that we have already defined two λ-terms M_{u_j} and N_{u_j} $(j < k)$ in hnf and that the Böhm trees of M_{u_j} and N_{u_j} are, respectively, the subtrees of $BT(M)$ and $BT(N)$ at root u_j. Assume that M_{u_j} and N_{u_j} have the following forms:

$$\begin{aligned} M_{u_j} &\equiv \lambda x_1^j \dots x_{n_j}^j.z_j M_1^j \dots M_{s_j}^j \\ N_{u_j} &\equiv \lambda x_1^j \dots x_{n_j}^j.z_j N_1^j \dots N_{s_j}^j\,. \end{aligned} \tag{12}$$

To abbreviate the notation, we will write M_{u_j} and N_{u_j} as follows:

$$\begin{aligned} M_{u_j} &\equiv \lambda \overline{x}_{n_j}^j.z_j M_1^j \dots M_{s_j}^j \\ N_{u_j} &\equiv \lambda \overline{x}_{n_j}^j.z_j N_1^j \dots N_{s_j}^j\,. \end{aligned}$$

Then the node u_j in the Böhm trees of M and N has s_j sons. Since $u_{j+1} = u_j r_{j+1}$ is a son of u_j in the Böhm trees of M and N, we have $r_{j+1} \leqslant s_j$. Then we define $M_{u_{j+1}}$ and $N_{u_{j+1}}$ as the principal head normal forms (principal hnfs, for short) of $M_{r_{j+1}}^j$ and $N_{r_{j+1}}^j$, respectively (see Barendregt (1984, Definition 8.3.20) for the definition of principal hnf).

The Böhm trees of $M_{u_{j+1}}$ and $N_{u_{j+1}}$ are, respectively, the subtrees of $BT(M)$ and $BT(N)$ at root u_{j+1}. The principal hnfs of M_{u_k} and N_{u_k} (recall that $u_k = u$ is the node where the Böhm trees are different) cannot have the same number of abstractions. Without loss of generality, we assume that

$$M_{u_k} \equiv \lambda \overline{x}_{n_k}^k.z_k M_1^k \dots M_{s_k}^k \tag{13}$$

$$N_{u_k} \equiv \lambda \overline{x}_{n_k}^k \lambda y_1 \dots y_r.z_k N_1^k \dots N_{s_k}^k Q_1 \dots Q_r \ (r \geqslant 1) \tag{14}$$

https://doi.org/10.1017/S0960129508006683 Published online by Cambridge University Press

where $y_i \leqslant_\eta Q_i$ $(1 \leqslant i \leqslant r)$, that is, Q_i is a possibly infinite η-expansion of the fresh variable y_i.

Let D be an arbitrary graph model. First we will show that the terms N_{u_k} and M_{u_k} have different interpretations in D, that is, there exist an element $\alpha_k \in D$ and a D-environment σ_k such that

$$\alpha_k \in (N_{u_k})^D_{\sigma_k} \quad \text{and} \quad \alpha_k \notin (M_{u_k})^D_{\sigma_k}. \tag{15}$$

Now we will show that this difference at level k can be propagated upward, that is, there exists an element $\alpha_0 \in D$ and a D-environment σ_0 such that

$$\alpha_k \in (N_{u_k})^D_{\sigma_k} \text{ iff } \alpha_0 \in (N_{u_0})^D_{\sigma_0}$$

and

$$\alpha_k \in (M_{u_k})^D_{\sigma_k} \text{ iff } \alpha_0 \in (M_{u_0})^D_{\sigma_0}.$$

From $\alpha_k \in (N_{u_k})^D_{\sigma_k}$ and $\alpha_k \notin (M_{u_k})^D_{\sigma_k}$, we get the conclusion: $\alpha_0 \in (N_{u_0})^D_{\sigma_0}$ (that is, $\alpha_0 \in N^D_{\sigma_0}$) and $\alpha_0 \notin (M_{u_0})^D_{\sigma_0}$ (that is, $\alpha_0 \notin M^D_{\sigma_0}$).

8.1. *The definition of the environments*

Let X be an incompatible set. Let $\beta_0, \ldots, \beta_k, \beta_{k+1}$ be distinct elements of X and let $\Delta = \{\delta_0, \delta_1, \ldots, \delta_k\}$ be an X-good set. Also, let:

— $z_0 \ldots z_k$ be the sequence of the head variables of the λ-terms $M_{u_0}, \ldots, M_{u_k}$ (note that not all these head variables are necessarily distinct);
— y_r be the last variable occurring in the external abstractions of N_{u_k} (note that we have $y_r \neq z_0, \ldots, z_k$);
— $\overline{x}^j_{n_j}$ be the sequence of variables occurring in the external abstractions of M_{u_j}.

We now define a sequence of environments ρ_j $(0 \leqslant j \leqslant k)$ and σ_i $(0 \leqslant i \leqslant k+1)$:

$$\rho_j(x) = \begin{cases} \{\delta_i : j \leqslant i \leqslant k \text{ and } z_i = x\} & \text{if } x \neq y_r \\ \{\delta_k\} & \text{if } x = y_r \end{cases}$$

$$\begin{aligned} \sigma_0 &\equiv \rho_0 \\ \sigma_{i+1} &= \sigma_i[\overline{x}^i_{n_i} := \rho_i(\overline{x}^i_{n_i})]. \end{aligned}$$

Notation: If τ and ρ are environments, we write $\tau \leqslant \rho$ for $\tau(x) \subseteq \rho(x)$ for all variables x.

Lemma 56.

(a) $i \leqslant j \Rightarrow \rho_i \geqslant \rho_j$.
(b) $\rho_j \leqslant \sigma_{j+1} \leqslant \rho_0$ for every $0 \leqslant j \leqslant k$.
(c) $\delta_j \in \sigma_{j+1}(z_j)$ for all $0 \leqslant j \leqslant k$.

Proof. (b) By definition we have $\sigma_1 = \rho_0$. Assume as induction hypothesis that $\rho_{j-1} \leqslant \sigma_j$. Then by (a) we have $\rho_j \leqslant \sigma_j$. The conclusion is an immediate consequence of the definition of σ_{j+1}. □

We now define by induction the structure of the elements δ_j (belonging to the X-good set Δ) and of the elements α_j $(0 \leqslant j \leqslant k)$.

https://doi.org/10.1017/S0960129508006683 Published online by Cambridge University Press

(i) $\delta_k \equiv \overline{\varnothing}_{s_k+r} \to \beta_k$.
(ii) $\alpha_k \equiv \rho_k(\overline{x}^k_{n_k}) \to \rho_k(\overline{y}_r) \to \beta_k$.

Assume we have defined δ_{j+1} and α_{j+1} $(j < k)$. We define δ_j and α_j as follows:

(i) $\delta_j \equiv \overline{\varnothing}_{r_j-1} \to \{\alpha_{j+1}\} \to \overline{\varnothing}_{s_j-r_j} \to \beta_j$.
(ii) $\alpha_j \equiv \rho_j(\overline{x}^j_{n_j}) \to \beta_j$.

8.2. *The conclusion of the proof*

In the following lemma we show that N_{u_k} and M_{u_k} have different interpretations.

Lemma 57. We have $\alpha_k \in (N_{u_k})^D_{\sigma_k}$ and $\alpha_k \notin (M_{u_k})^D_{\sigma_k}$.

Proof. We apply Lemma 54 by putting $S \equiv M_{u_k}$, $T \equiv N_{u_k}$, $\delta \equiv \delta_k$, $t \equiv s_k$, $\beta \equiv \beta_k$, $\rho \equiv \rho_k$, $\sigma \equiv \sigma_k$, $u \equiv r$ and $i = r$. □

The different interpretation of N_{u_k} and M_{u_k} can be propagated upward as shown in the following lemma.

Lemma 58. For every $k > j \geqslant 0$ we have

$$\alpha_j \in (N_{u_j})^D_{\sigma_j} \;\Leftrightarrow\; \alpha_{j+1} \in (N_{u_{j+1}})^D_{\sigma_{j+1}}$$

and

$$\alpha_j \in (M_{u_j})^D_{\sigma_j} \;\Leftrightarrow\; \alpha_{j+1} \in (M_{u_{j+1}})^D_{\sigma_{j+1}}.$$

Proof. We apply Lemma 55 to N_{u_j}, by putting $S \equiv N_{u_j}$, $\overline{x} \equiv \overline{x}^i_{n_j}$, $\delta \equiv \delta_j$, $i \equiv r_j$, $\alpha' \equiv \alpha_{j+1}$, $\alpha \equiv \alpha_j$, $\rho \equiv \rho_j$ and $\sigma \equiv \sigma_j$, and by observing that $\sigma_{j+1} \equiv \sigma_j[\overline{x}^j_{n_j} := \rho_j(\overline{x}^i_{n_j})]$. We similarly apply Lemma 55 to M_{u_j}. □

Lemma 59. We have $\alpha_0 \in N^D_{\sigma_0}$, while $\alpha_0 \notin M^D_{\sigma_0}$.

Proof. Recall that $N \equiv N_{u_0}$ and $M \equiv M_{u_0}$. By applying Lemma 58 it is easy to show that $\alpha_0 \in N^D_{\sigma_0} \Leftrightarrow \alpha_k \in (N_{u_k})^D_{\sigma_k}$, and $\alpha_0 \in M^D_{\sigma_0} \Leftrightarrow \alpha_k \in (M_{u_k})^D_{\sigma_k}$. Then the conclusion is immediate, since, by Lemma 57, we have that $\alpha_k \in (N_{u_k})^D_{\sigma_k}$ and $\alpha_k \notin (M_{u_k})^D_{\sigma_k}$. □

9. Conclusion and future work

In this paper, we have collected together and organised several already published results and some new material: the existence of the minimum and minimun sensible graph theories appeared originally in Bucciarelli and Salibra (2003) and Bucciarelli and Salibra (2004), respectively. The new presentation in Section 3 stresses the relevance and generality of the *weak product* construction underlying these results, and adds some new results (for instance, the fact that the theory of weak products is semisensible and is, in general, strictly finer than the intersection of the factors' theories, which was obtained using the notion of a *self weak product*).

Section 6 covered the main result of Bucciarelli and Salibra (2004), namely the fact that the maximal sensible graph theory is $\mathscr{B}$.

https://doi.org/10.1017/S0960129508006683 Published online by Cambridge University Press

The content of Section 5, which provides a negative answer to the question of whether $\lambda\beta$ is the minimal graph theory, also appeared in Bucciarelli and Salibra (2004). Actually, this negative result opens the way to the investigation of the minimal graph theory.

Sections 3.2 and 3.3 presented new results. First, the fact that stratified graph models, which are those obtained by canonical completion of partial pairs, that is, virtually all known graph models apart from those constructed by *forcing* (Baeten and Boerboom 1979; Berline and Salibra 2006), have semisensible theories. Then we showed that the theory of a weak product is in general strictly finer than the intersection of the factors' theories. Finally, we provided equations between unsolvable terms that are not preserved in weak products.

Several questions about graph λ-theories remain open. Among them, we wish to address that concerning the minimal sensible graph theory: Is it $\mathscr{H}$ (the minimum sensible theory) or is it bigger? For the time being, we are able to separate in a graph model some typical examples of $\mathscr{B}$-equivalent, $\mathscr{H}$-distinct λ-terms, like Yx and Θx.

The notion of an *effective* graph model is a natural one: it is enough to ask that the coding function be total recursive with respect to given enumerations of the model's web, finite sets and pairs of natural numbers. Then one recasts classical recursion theory results in the framework of graph models, and this seems particularly compelling since those are models of the λ-calculus. A paper on this subject is Berline *et al.* (2007).

Acknowledgements

The authors would like to thank Chantal Berline and Giulio Manzonetto for helpful discussions and comments.

References

Abramsky, S. (1991) Domain theory in logical form. *Annals of Pure and Applied Logic* **51** 1–77.

Alessi, F., Dezani, M. and Honsell, F. (2001) Filter models and easy terms. ICTCS'01. *Springer-Verlag Lecture Notes in Computer Science* **2202** 17–37.

Baeten, J. and Boerboom, B. (1979) Omega can be anything it should not be. *Indag. Mathematicae* **41** 111–120.

Barendregt, H. P. (1984) *The lambda calculus: Its syntax and semantics*, North-Holland.

Barendregt, H. P., Coppo, M. and Dezani-Ciancaglini, M. (1984) A filter lambda model and the completeness of type assignment. *J. Symbolic Logic* **48** 931–940.

Bastonero, O. and Gouy, X. (1999) Strong stability and the incompleteness of stable models of λ-calculus. *Annals of Pure and Applied Logic* **100** 247–277.

Berline, C. (2000) From computation to foundations via functions and application: The λ-calculus and its webbed models. *Theoretical Computer Science* **249** 81–161.

Berline, C. (2006) Graph models of λ-calculus at work, and variations. *Mathematical Structures in Computer Science* **16** 185–221.

Berline, C. and Salibra, A. (2006) Easiness in graph models. *Theoretical Computer Science* **354** 4–23.

Berline, C., Manzonetto, G. and Salibra, A. (2007) Lambda theories of effective lambda models. In: 16th EACSL Annual Conference on Computer Science and Logic (CSL'07). *Springer-Verlag Lecture Notes in Computer Science* **4646** 298–312.

Berry, G. (1978) Stable models of typed lambda-calculi. In: Proc. 5th Int. Coll. on Automata, Languages and Programming. *Springer-Verlag Lecture Notes in Computer Science* **62**.

https://doi.org/10.1017/S0960129508006683 Published online by Cambridge University Press

Bucciarelli, A. and Ehrhard, T. (1991) Sequentiality and strong stability. *Sixth Annual IEEE Symposium on Logic in Computer Science*, IEEE Computer Society Press 138–145.

Bucciarelli, A. and Salibra, A. (2003) The minimal graph model of lambda calculus. 28th International Symposium on Mathematical Foundations of Computer Science. *Springer-Verlag Lecture Notes in Computer Science* **2747**.

Bucciarelli, A. and Salibra, A. (2004) The sensible graph theories of lambda calculus. *19th Annual IEEE Symposium on Logic in Computer Science (LICS 2004)*, IEEE Computer Society Press.

Church, A. (1933) A set of postulates for the foundation of logic. *Annals of Math.* **2** 346–366.

Church, A. (1941) *The calculi of lambda conversion*, Princeton University Press.

Coppo, M. and Dezani, M. (1980) An extension of the basic functionality theory for the λ-calculus. *Notre Dame J. Formal Logic* **21** 685–693.

David, R. (2001) Computing with Böhm trees. *Fundamenta Informaticae* **45** 53–77.

Di Gianantonio, P. and Honsell, F. (1993) An abstract notion of application. Typed lambda calculi and applications. *Springer-Verlag Lecture Notes in Computer Science* **664** 124–138.

Di Gianantonio, P., Honsell, F. and Plotkin, G. (1995) Uncountable limits and the lambda calculus. *Nordic J. Comput.* **2** 126–145.

Gouy, X. (1995) *Etude des théories équationnelles et des propriétés algébriques des modéles stables du λ-calcul*, Thèse, Université de Paris 7.

Honsell, F. and Ronchi della Rocca, S. (1992) An approximation theorem for topological lambda models and the topological incompleteness of lambda calculus. *Journal of Computer and System Sciences* **45** 49–75.

Kerth, R. (1995) *Isomorphisme et équivalence équationnelle entre modèles du λ-calcul*, Thèse, Université de Paris 7.

Kerth, R. (1998) Isomorphism and equational equivalence of continuous lambda models. *Studia Logica* **61** 403–415.

Kerth, R. (2001) On the construction of stable models of λ-calculus. *Theoretical Computer Science* **269** 23–46.

Longo, G. (1983) Set-theoretical models of λ-calculus: theories, expansions and isomorphisms. *Ann. Pure Applied Logic* **24** 153–188.

Plotkin, G. (1993) Set-theoretical and other elementary models of the λ-calculus. *Theoretical Computer Science* **121** 351–409.

Salibra, A. (2001) A continuum of theories of lambda calculus without semantics. *16th Annual IEEE Symposium on Logic in Computer Science*, IEEE Computer Society Press 334–343.

Salibra, A. (2003) Topological incompleteness and order incompleteness of the lambda calculus. (LICS'01 Special Issue). *ACM Transactions on Computational Logic* **4** 379–401.

Scott, D. S. (1972) Continuous lattices. In: Toposes, Algebraic geometry and Logic. *Springer-Verlag Lecture Notes in Mathematics* **274**.

Selinger, P. (2003) Order-incompleteness and finite lambda reduction models. *Theoretical Computer Science* **309** 43–63.

https://doi.org/10.1017/S0960129508006683 Published online by Cambridge University Press