Reviews in Clinical Gerontology 2014 24; 41–57 First published online 27 November 2013
© Cambridge University Press 2013 doi:10.1017/S095925981300021X

Dysphagia: implications for older people

N Miller[1] and J Patterson[1,2]

[1]Institute of Health and Society, University of Newcastle upon Tyne, UK and [2]Sunderland Royal Hospital/ University of Newcastle upon Tyne, UK

Summary

Dysphagia represents a salient concern in many conditions prevalent in older people. There are direct implications for morbidity and mortality. The importance of recognizing and managing dysphagia in hospital and the community also extends to psychosocial impact and quality of life, as well as health, economic and ethical-legal issues. This review outlines reasons for the importance of recognizing and treating dysphagia. It then proceeds to look at recent developments in our understanding of the nature, assessment and management of dysphagia in older people. Whilst there are well-established practices in assessment and management, ongoing work continues to challenge the validity and reliability of many methods. These concerns are covered and directions for future developments highlighted.

Key words: dysphagia, assessment, management, older people, ageing.

Introduction

Dysphagia and factors directly associated with it have important implications for older people. We first outline some of these implications before proceeding to a review of recent work on the nature, assessment and management of dysphagia in older people. In preparing the review we consulted search engines Web of Knowledge, Scopus, MEDLINE and Ovid using terms dysphagia, swallow(ing) (disorders); combined with older people, stroke, neurological (disorder), pneumonia, dementia; head and neck, oral, pharyngeal, laryngeal cancer; assessment, rehabilitation. The term dysphagia elicits >50,000 references since 2009, and >8000 for dysphagia and older persons/people. This review focuses on works from predominantly 2009 onwards that answer or address key issues and updates in the clinical assessment and management of older people (65+ years) with a diagnosis or risk of swallowing disorder.

Dysphagia represents a central cause of, or significant sequel to, many conditions faced by older individuals. Severe dysphagia poses an immediate risk for choking and possible death. Chronic dysphagia is independently associated with a range of other negative factors, including:

- poor(er) prognosis for survival;[1,2]
- increased co-morbidities;[3–9]
- risk of short- and long-term undernourishment, weight loss, malnutrition and dehydration;[10,11]
- less positive rehabilitation outcomes, later discharge from acute care, less chance of returning home on discharge and increased likelihood of re-admission.

Eating and drinking ability is strongly associated with quality of life. Dysphagia results in a significant social and psychological burden for both patients and their families.[12–15] Some patients report being fearful and not wanting to eat alone in case they choke and consequently avoid eating, reduce or alter their overall oral intake. For others, embarrassment can lead to unwillingness to eat within a social context, severely restricting lifestyle. Depression and frustration at restricted food choices and changes to eating routines is common. In addition, the impact on partners caring for a dysphagic patient can be profound, from additional time spent planning, preparing and selecting food for a special diet, acquiring new skills in caring for a feeding tube and a restriction in their own social activities and interactions. Perceptions of psychological issues around impact can evolve from acute to chronic phases.[14]

Dysphagia constitutes a significant economic burden to health care services.[1,16,17] The American National Hospital Discharge Survey (2005–2006)[2] of over 77 million admissions found the median number of hospital days for patients without dysphagia was 2.4 compared with 4.04 days for

Address for correspondence: Nick Miller, Institute of Health and Society, Speech and Language Sciences, George VI Building, University of Newcastle upon Tyne, Newcastle upon Tyne NE1 7RU, UK.
Email: Nicholas.Miller@ncl.ac.uk

https://doi.org/10.1017/S095925981300021X Published online by Cambridge University Press

those with dysphagia. Furthermore, the incidence of dysphagia-associated admissions amongst the elderly (>75 years) was twice that of all other age groups. There was a 40% increase in length of hospital stay when a patient is admitted with dysphagia. These patients had a significantly higher risk (RR 9.1) of developing aspiration pneumonia.

What is dysphagia?

Dysphagia in its narrow sense indicates a disorder of the triggering of the patterned response that propels a bolus from the oral cavity into the oropharynx and/or of the subsequent train of automatic events that transport the bolus to the stomach whilst simultaneously guarding the airway. This pharyngeal-oesophageal dysphagia may arise from delay, reduction or absence of triggering of any or all of the patterned responses; inefficiencies or failure in transport due to altered muscle tone, power and co-ordination; inefficiencies or anatomical blockage from altered tissue status or narrowing of the food passage.

More often a broader definition is applied. Triggering of responses and whether reduced or delayed operation of automatic stages become a problem or not are intricately bound up with the ability of an individual to form a bolus in the mouth and voluntarily direct this posteriorly to initiate the automatic sequence. This in turn is influenced by bolus viscosity and volume.[18,19] These problems (generally labelled oral dysphagia) can arise from reduced tongue mobility,[20,21] misco-ordination of breathing and swallowing,[22–24] altered oral sensation,[25] poor lip seal, particularly for fluids; poor dentition,[26,27] and oral pathology.[28,29] Problems may also be associated with cognitive changes,[30] given that successfully delivering food or drink to the mouth, forming a bolus and initiating swallowing depends on intact attention, awareness of material in the mouth, persistence with chewing and clearing of debris. Being unaware of or not acknowledging the presence of dysphagia exposes the individual to possible harmful practices around swallowing and impairs ability to exploit possible compensatory tactics.

As drooling is generally associated with reduced clearing of saliva and foodstuffs[31] some include sialorrhoea as part of the dysphagic complex. Several works differentiate between anterior drooling (loss of secretions and food through the lips) *versus* posterior drooling (uncontrolled escape of secretions, food and drink into the oropharynx).

Table 1. *More common conditions associated with dysphagia in older people*

Dysphagia highly likely (at some stage)
• Stroke
• Head injury
• Degenerative neurological diseases
• Advanced head and neck cancer
Dysphagia quite possible (at some stage)
• Neoplastic changes in swallowing tract
• Displacement of swallowing tract by masses/lesions outside swallowing tract
• Dystonia,
• Muscular dystrophies
• Impaired insight and arousal
• Slowed cognitive processing
Dysphagia can be associated with
• Temporal-mandibular joint displacement or arthritis
• Oesophageal diverticula, webs and rings
• Osteophytes
• Cricopharyngeal bar
• Oropharyngeal infections/inflammation
• Connective tissue disorders of the swallowing tract
• Radiotherapy effects
• Surgical effects (e.g. oropharyngeal; laryngeal; cervical spine; thyroid; thorax; intubation)
• Medication effects (e.g. dry mouth, loss of appetite, nausea, dyskinesias)
• Dental/gingival deterioration
• Pain from any source
• Psychiatric disorders (including elective eating disorders)
• Depression

The basic anatomy, physiology and neurology of swallowing are covered in most standard textbooks. Recent developments have sought a deeper understanding of dysphagia, to go beyond mere description of what is seen in order to build explanatory models incorporating sensory as well as motor and cross-modal variables in swallowing control. The effects of taste, aroma and texture are not neutral factors in the patterned sensorimotor control.[32–36]

Cause and course of dysphagia

Table 1 lists conditions with which dysphagia is commonly associated, though it is by no means exhaustive for all possible causes and associations.

https://doi.org/10.1017/S095925981300021X Published online by Cambridge University Press

Table 2. *Indications for possible presence of dysphagia*

Strongly suggestive of definite dysphagia
• Pain on or after swallow
• Poor management of secretions
• Breathlessness on/after swallow
• Cough on/after swallow
• 'Gurgly' voice after swallow
• Nasal regurgitation of food
• Delayed regurgitation of undigested food
Strongly warrant further investigation for possible dysphagia
• Repeated chest infections of no obvious cause
• Repeated (unexplained) pyrexia
• Unexplained weight loss
• Significant change to eating habits (content, conduct, preparation)
• New intolerance/avoidance of certain foodstuffs
• Individual can manage fluids but not solids or vice versa
• Slowed/prolonged swallow
• Fear of choking
• Report difficulty swallowing medication
May indicate (risk of) dysphagia is a consideration
• Need for repeated swallows to clear bolus
• Weak voluntary cough
• Food residue in mouth
• Poor oral/dental health
• Inability to maintain optimum eating posture
• Reduced appetite

Likelihood and prognosis of dysphagia vary markedly by aetiology. Irrespective of origin, dysphagia should be suspected when particular signs and symptoms are observed. Table 2 summarizes the salient ones.

In certain conditions, dysphagia may be transitory (e.g. post-extubation post-operatively though there is a less favourable prognosis post-extubation after mechanical ventilation in critical illness,[37] transient ischaemic attack, recovered upper gastrointestinal ulceration), and some mechanical and anatomical causes, e.g. cricopharyngeal web, may be correctable. In other conditions, full or partial recovery to functionally viable swallowing may be a reasonable expectation (e.g. mild stroke, spinal surgery[38–40]). In some circumstances dysphagia may be slowly or more rapidly progressive[41–43] and result in significant morbidity or mortality.

Swallowing safety may alter rapidly in the presence of secondary morbidities, even those not directly associated with dysphagia. For instance, in an individual with previously undiagnosed dysphagia or where a relatively mild dysphagia was well compensated, onset of delirium, infections, clouding of consciousness, respiratory disorder or pain may be linked to appearance of significant swallowing problems.

In progressive disorders, patients may reach a threshold where hitherto consciously or non-consciously applied compensatory strategies to manage their inefficient or mildly dysphagic swallow are no longer effective. When severity or multiplicity of other impairments, including the stage of underlying degeneration, reaches a critical level, rapid deterioration in swallowing status can ensue.[44]

Dysphagia is noted as a common and serious problem for individuals with mental illness.[45] Estimates indicate they may be up to forty-three times more likely to die of choking asphyxiation than the general population. Presence of dysphagia in dementia is well documented.[46] Even where cognitive changes are not profound, effects of divided attention on chewing and swallow initiation may exacerbate a dysphagic condition.[47]

Depression may be a sequel to the psychosocial impact of having a swallowing disorder. People with disorders such as stroke, head injury and Parkinson's typically also present with depression. The cause–effect direction or relationship between depression, swallowing and underlying disorder remains a topic of investigation. Nevertheless, depression is strongly associated with dysphagia.[42,48,49]

Advancing age brings changes to the swallowing process, including slower transit times, possible loss of muscle mass and tissue elasticity, and possible reduction in strength of muscles. There is debate as to whether these alterations are an independent cause of dysphagia or whether dysphagia, if present in otherwise healthy individuals, represents a manifestation of other conditions associated with older age, or with undiagnosed conditions – e.g. 'silent' stroke, prodromal phase of a degenerative condition, cognitive or mood disorder (see below regarding dysphagia in healthy community-dwelling elders).

Prevalence of dysphagia

Given the variable causes and courses of dysphagia, a definitive statement on overall prevalence is problematic. Comparison across studies is

https://doi.org/10.1017/S095925981300021X Published online by Cambridge University Press

further complicated by the variety of assessment techniques employed and divergent operational definitions of variables applied.[50] Swallowing changes (e.g. delayed transit time, diminished reflexes) can be detected instrumentally at onset in up to 100% of people with some disorders, but this does not necessarily indicate an individual has dysphagia in the sense that there exists an immediate threat of airway compromise or consequences for hydration and nutrition (though identifying such changes does oblige careful monitoring).

Where self-report measures are employed, an additional factor concerns awareness. Individuals may either be unaware, for a variety of reasons, that they have dysphagia,[42,51,52] or they mention behaviours suggestive to clinicians of dysphagia but attribute these to other factors unrelated to swallowing[53] and therefore do not admit to swallowing problems. This implies that asking in assessment or at review only 'do you have any swallowing problems' may elicit an inaccurate answer. Keener insights are gained from questioning around possible changes to types and consistencies of food that an individual may now avoid or experience difficulty with, or has made changes to their preparation (puree, mash, crush), time to eat food; fear of choking; and observation for behaviours (e.g. Table 2) may also be suggestive of dysphagia.

This underlines the need for a broad approach to assessment. It emphasizes the need for studies of prevalence to ideally chart changes over the course of an illness and offer longer term follow-up, not just on instrumental assessments but on functional, nutritional, respiratory health and swallowing-related quality of life measures, to judge whether or when demonstrated changes to swallowing patterns indeed represent a daily living or health-related problem.

Estimates of prevalence of dysphagia in generally healthy, community-dwelling older people range from around 11% to around 40%. Differences reflect population biases, assessment techniques and operational cut-offs for presence or not of dysphagia. Holland *et al.*[49] analysed self-report responses to a postal questionnaire from 634 respondents, with 11% indicating presence of symptoms suggestive of dysphagia. Increasing age correlated directly with dysphagia severity. Serra-Prat *et al.*[11,54] examined 254 community-dwelling people >70 years for swallowing efficiency and safety and aspiration. They employed a screen involving swallowing different viscosities and volumes, alongside checklists for possible signs of dysphagia (cough on swallow, etc.) and pulse oximetry. Dysphagia was present over the whole sample in 27%, impaired efficacy in 20.5% and impaired safety in 15%, with signs of aspiration in 6.7%. Likelihood increased steeply after 80+ years, with dysphagia prevalence of 17% in 70–79 years *versus* 33% in 80+ years. Roy *et al.*[55] interviewed 117 independently living elders and found 38% reported swallowing disorder at some stage, with 33% indicating a current problem. Estimates for undiagnosed dysphagia in residential and nursing care facilities are correspondingly higher, falling in the range of 30–68%.[56]

Assessment of swallowing

The imperative of assessing for dysphagia in at-risk conditions and populations is recommended in key guideline publications.[57,58] Assessment preferably involves instrumental assessments, but at a minimum should entail clinical assessment of swallowing status as well as psychosocial impact and related risk factors. NICE guidelines stipulate, in acute stroke, a swallowing screening test by a trained healthcare professional before administrating any food, fluid or medication. If a problem is identified, a specialist swallowing assessment should follow within 72 hours. Head and neck cancer patients planned for treatment that will affect swallowing, should be assessed (in UK by a speech and language therapist) and a plan for method of feeding devised.[59] Consideration must be given to risk factors that may result in dysphagia (Tables 1 and 2). For instance, not all tracheostomized patients will have dysphagia, but a referral for swallowing assessment would be indicated if there is co-occurring neurological involvement following head and neck surgery, or there are signs of a swallowing problem such as evidence of secretions or ingested material on suctioning.[60] As discussed above, dysphagia may not always be as a result of specific medical conditions. In addition to this, the person may not be aware of a swallowing difficulty, or may attribute their problem to other sources. Therefore, ongoing vigilance is required by health care staff for signs and symptoms of dysphagia, including more

https://doi.org/10.1017/S095925981300021X Published online by Cambridge University Press

Table 3. *A comparison of features of VF vs FEES instrumental assessment*

Feature	FEES	VF
Equipment	Nasoendoscope, camera, monitor, digital recorder	Fluoroscopy, digital recorder
Personnel	Otolaryngologist and/or trained speech and language therapist	Radiologist and/or radiographer, speech and language therapist
Safety	Mild discomfort	Radiation exposure
Accessibility	Portable, bedside use	X-ray department only
Oral stage	Not observed	Well seen
Pharyngeal stage	Inferred assessment before and after swallowing	Well seen
Oesophageal stage	Not observed	Can be observed
Aspiration/penetration	Well seen	Well seen
Pharyngeal sensation	Can be tested. Sensory testing equipment can provide more specific information	Not assessed
Test material	Any type of food and drink, often dyed for visibility	Radio-opaque contrast added, altering viscosity, texture and taste
Length of assessment	Limited according to patient tolerance. Good for assessing effect of fatigue	Limited by radiation exposure

Table adapted from Hirst.[170]

covert features, with a view to timely referral for specialist swallowing assessment.

Instrumental assessments

Instrumental assessments provide detailed information about the anatomy and physiology of the oropharyngeal swallow as well as evaluating risk factors such as the presence, amount and response to pooled or aspirated material. Two main clinical procedures provide complementary but differing information: videofluoroscopy (VF) and Fibreoptic Endoscopic Evaluation of Swallowing (FEES). A summary comparison between the two assessments is given in Table 3.

The selection of procedure depends on patient suitability and which questions need answering. Both assessments are difficult to conduct with patients who have attention and behavioural difficulties and where there are likely to be movement artefacts. VF uses a modified barium swallow, providing a dynamic X-ray image of the three (oral, pharyngeal, oesophageal) stages of swallowing, enabling visualization of the swallowing process from the lips through to the upper oesophageal area. FEES provides an endoscopic view of the laryngopharynx, giving information on secretion management, altered anatomy, laryngeal sensation and vocal fold motility. The introduction of a swallow trial is usually preceded by a functional assessment of laryngeal pharyngeal movements, e.g. pharyngeal squeeze test. FEES does not directly view the oral stage nor cricopharyngeal opening. However, it is ideal for patients who are unable to attend the radiology department and it is highly repeatable. The reliability and validity for the interpretation of both of these assessments has been published.[61–63] Reliability improves when raters routinely use scales such as the MBSImps or the penetration/aspiration scale.[64–66]

Instrumental assessments may be conducted for diagnostic purposes. Some of the compensatory interventions listed below (e.g. chin-tuck, effortful swallow, altered consistency) may be tested for their effectiveness during instrumental assessment.

Clinical assessment

Practices vary across services and countries.[5,67–70] Some services routinely refer all patients with suspected or risk of dysphagia for VF or FEES assessment. More often, initial evaluation is conducted using a clinical or bedside screen that aims to gauge swallowing efficiency and safety and identify candidates with possible penetration or aspiration for more objective examination.[71–73] There are many published procedures.[74–78]

Typically patients are asked to swallow liquids and solids graded for viscosity, constituency/

https://doi.org/10.1017/S095925981300021X Published online by Cambridge University Press

texture and volume. The examiner checks time taken and number of swallows to clear, and observes for signs said to be indicative of aspiration or penetration, such as cough on swallow, oxygen desaturation, altered voice quality or altered swallow sounds on cervical auscultation. None of these methods or observed behaviours has strong support as an isolated indicator of dysphagia, or more specifically risk for aspiration. There are ongoing issues around the sensitivity and specificity achieved with clinical examinations and the validity and reliability of claimed signs of penetration and aspiration.

A review of thirty-five protocols for screening post-stroke dysphagia[79] found that only four[75,76,80,81] met basic psychometric criteria and delivered acceptable levels of sensitivity (>87%) and negative predictive values (>91%). All were considered to need further development. A similar review[82] of bedside swallow screens as predictors of post-stroke pneumonia came to less optimistic conclusions. They highlighted the lack of validation and standardization on unpreselected populations (a criticism shared by others[83]). For them the diagnostic accuracy of bedside swallow tests remains unclear and the evidence is insufficient to support their use in a general older population, and consequently, the predictive value for pneumonia remains unknown – again a conclusion reached by others.[84]

A systematic review of screening items for dysphagia risk post-stroke[85] found many items associated with dysphagia, with water swallow tests[78,84] performing an important role, especially in combination with monitoring for 'wet' voice quality. However, in their opinion, the optimum combination of non-swallowing and swallowing items and the most valid protocol remains to be determined. Use of single measures or single liquid volumes do not distinguish healthy individuals from those with stoke with or without dysphagia.[50] Recent investigations confirm earlier cautions expressed for practices such as cervical auscultation of swallow sounds, cough on swallow and wet voice.

Specificity of perceived 'wetness' of voice quality was found to have acceptable levels (>86% for dysphagia, 94% for penetration-aspiration) but unacceptably low sensitivity (8 and 14%, respectively).[86] Values for overall voice quality were lower still. Standard acoustic voice perturbation measures (pitch and amplitude perturbation and harmonics to noise ratio) also fared poorly as discriminators. The authors concluded that the presence of abnormalities in voice samples associated with swallowing, detected either perceptually or acoustically, can easily be misinterpreted and is not a valid means of determining a patient's swallowing status.

A similar study with individuals treated for oral or oropharyngeal cancer arrived at like findings.[87] By contrast, they established that alterations to voice intensity were significantly associated with penetration and aspiration, whilst raised fundamental frequency (higher pitch) was strongly linked to swallow inefficiency. Chang *et al.*[88] found no differences in voice perturbation measures between aspirators and non-aspirators. However, they did not find any alteration to fundamental frequency in their patients either.

A further study[89] had participants attempt to raise their voice pitch after a barium swallow. In this pilot study perceived and acoustically measured pitch elevation independently predicted penetration-aspiration on thin liquid swallows, but vocal range (average to falsetto) did not. Encouraging findings also point to the possibility of detecting aspiration non-invasively utilizing acoustic analysis of breath sound changes.[90]

Cough on swallow has long been considered an indicator of possible aspiration-penetration and for referral for more objective testing.[91] Inter and intra-rater reliability at recognizing and agreeing on coughs after citric acid inhalation of experienced speech language pathologists who employ cough monitoring in assessment and inexperienced ones has been shown to be only fair to moderate.[92] The same research group[93] investigated whether decisions around cough reflex testing enhanced accuracy and outcomes in relation to prediction and occurrence of pneumonia. Presence of cough on swallow influenced diet recommendations and referral for instrumental assessment, but had no bearing on rate of pneumonia and mortality at 3 months post-evaluation.

Using objective airflow measures on voluntary coughs in people with Parkinson's[94] and cough peak flow (as part of a suit of pulmonary function tests),[95] accurate predictions of respiratory prognosis and risk level in aspiration were achieved. It may therefore be that cough on swallow evaluation needs to be combined with objective measures rather than rely on naked ear perceptual decisions.

https://doi.org/10.1017/S095925981300021X Published online by Cambridge University Press

However, the key issue may not be so much the presence or not of aspiration. From instrumental studies, it is apparent that many people with dysphagia who aspirate do not cough on swallow,[96] and many known aspirators do not develop aspiration pneumonia. Reports indicate that, in certain populations, even when individuals are known aspirators, oral water intake may not adversely affect outcomes.[97–99] This has led several to argue that maybe the key issue is not the presence or degree of aspiration, but whether there is aspiration or not in relation to other risk factors, in particular oral health, as a source of pathogens.[36,100] Ideally then, screening encompasses oral health.

Oral health

Presence of oral pathologies may be a direct cause of swallowing difficulties, or exacerbate an underlying dysphagia where they interfere with chewing, bolus formation or passage. Dental status can affect preparation of the bolus and turn a borderline inefficient swallow or mild dysphagia into a major problem.[26,27] Poor oral health alongside dysphagia significantly increases the risk of pneumonia and other complications.[29,101–103]

Meal time observation

Eating represents not solely a mechanical event but also a social process. Eating-related problems can extend well beyond dysphagia. Important information on eating and swallowing can be derived from systematic observation of mealtime behaviours. This is particularly useful where the patient is unable to engage in an instrumental assessment or swallow trial during a bedside assessment. Observations may include specific eating problems such as drooling or food pocketing, ability to self-feed, caregiver feeding skills, body and head positioning, behavioural issues such as agitation, sensory problems such as visual perceptual disorders, and distraction by other stimuli. A recent systematic review of assessment tools used to record observations during a natural meal activity[104] identified two assessments with reported acceptable psychometric properties, the Minimal-Eating Observation Form-version II[105] and the McGill Ingestive Skills Assessment.[106]

Quality of life and the psychosocial impact of dysphagia

Swallowing difficulties can have a major impact on the quality of life of individuals and their families. Therefore, a comprehensive assessment includes attention to the psychosocial impact of dysphagia. A number of validated swallowing-specific quality of life questionnaires are available.[107] More established ones include SWAL-QOL and SWAL-CARE;[108] the MD Anderson Dysphagia Inventory,[109] a 20-item questionnaire validated on head and neck cancer patients, but which has also been reported with other elderly groups;[1,110] and EAT-10,[111] validated on a mixed dysphagia group of oropharyngeal and oesophageal origin. A number of published head and neck cancer studies have reported on outcomes using these questionnaires, and there is a growing body of papers on neurological conditions.[112,113]

Longer term swallowing monitoring

A disadvantage of instrumental and clinical assessments as described above is that they deliver a snapshot in time of swallowing status. Swallowing ability and efficiency may vary significantly over time (e.g. stage in drug cycle; diurnal variation) and place (e.g. quiet clinic one-to-one situation *vs* eating and talking, eating under time pressure), as well as in association with different meal consistencies. Deeper insights into overall swallowing status and possible risks would be improved if this variability could be captured. Several attempts have been made recently to develop devices that enable ambulant longer term monitoring of swallowing frequency and efficiency through acoustic analysis of swallow sounds, accelerometry or measurement of surface electromyographic activity.[114–116] Promising preliminary results have been attained and the field represents a definite area of fruitful development.

Importance of protocols and systematic screening

Whatever the pathway to assessment, findings stress the importance of employing an agreed protocol for screening and management of dysphagia.[57,58,117] Use of protocols has been shown to add accuracy in predicting pneumonia[118]

https://doi.org/10.1017/S095925981300021X Published online by Cambridge University Press

and reduce the incidence of pneumonia in an intensive care unit[119] and on a stroke unit.[120] The latter demonstrated advantages not just in reduced mortality and morbidity but in cost savings on drugs. Wilson *et al.*[121] found that routine VF screening (rather than bedside screening, or combined VF and bedside screen) enabled timely recognition of dysphagia in acute stroke and instigation of effective treatment. Cost benefits of reduced pneumonia and reduced impairment of quality of life outweighed costs of VF assessment.

Interventions

Management of dysphagia aims to address swallowing function, to impact on nutritional status, reduce risk of chest infections and other medical complications and enhance quality of life. It often requires a combination of interventions and a team approach, including from professionals such as speech-language therapists, dietitians, geriatricians, care home staff, and general practitioners. Management may include a combination of direct work on swallowing, compensatory interventions, environmental modification, and surgical and radiological procedures. Typically, compensatory strategies are applied to effect immediate advances where possible, alongside environmental modifications that might also afford gains. Direct interventions on oral aspects of feeding and swallowing may be instigated to bring about longer term underlying improvement or maintenance. Patient participation in therapy is an issue, with reports of non-compliance varying from 0 to 80%.[122] Common reasons are denial of a swallowing problem and finding texture modifications unpleasant. Accordingly, patient engagement must be a consideration.

The quality of investigations into swallowing therapy has improved over the past few years. However, carefully controlled, large-scale studies are still lacking. Information on critical dose size and intensity remains patchy.[123] Many investigations report outcomes for objective measures such as diameter of upper oesophageal sphincter opening or hyoid bone excursion, but fail to include whether differences are maintained and crucially whether such changes translate into improvements in swallow safety and amelioration of secondary complications. Even where recommendations are available, it is not always apparent that full awareness of them is reflected in practice[68,124] or that they are followed in clinical situations.[70] The following lists more common approaches to therapy that have had some evaluation.

Compensatory interventions address changes to behaviours that can influence swallow safety and/or compensate by altering food content and presentation:[125–128]

- Seating modification and physical therapy to improve trunk and head control and arm-and hand co-ordination. The optimal posture for most is an upright position.
- Certain head positions, e.g. a chin-tuck or head turn may facilitate swallowing efficiency and safety.[129]
- Swallowing manoeuvres that introduce an airway protection technique or a more effortful swallow.[130]
- Feeding equipment such as adapted cups, shallow based spoons, non-slip table mats.
- Texture modification. Food and drink may be softened (mashed, pureed, liquidized), thickened[131] or presented in smaller bites. Content is adjusted to maintain nutritional balance and calorific value. Standardized dysphagia diet food texture descriptors have been published by the Patient Safety Agency.[132] There can be issues around acceptability of pureed diets and added burden on carers.[12,122] Some modifications may improve taste or texture tolerability but not affect swallow parameters.[133] Reviews of nutritional programmes for undernourished older people (not necessarily diagnosed with dysphagia) support their use for improved nutritional and health outcomes, even though further evidence is required.[134–136]

Environmental modifications have proved successful in improving nutrition in residential care settings[137] and are adaptable to other settings. They cover such strategies as:

- Increase concentration for swallowing by reducing distractions such as the television or attempting to eat and talk simultaneously.
- Allowing more time for meals. A little and often approach may be useful with the addition of snacks.
- Heat-retaining plates to keep hot foods at a more palatable temperature.

https://doi.org/10.1017/S095925981300021X Published online by Cambridge University Press

- Optimal food presentation, such as moulds for puree textures.
- Verbal prompts, especially where cognitive, attentional or consciousness level variables impact on swallowing.
- Staff/carer training in feeding assistance and techniques.

Swallowing therapy directly targeting the mechanics of swallowing has taken several forms, including:

- Physical therapy to increase range, force and sustainability of movements of lips and tongue; to increase upper oesophageal opening; and to reduce misco-ordination of swallowing and breathing has some support.[138–143] These may or may not be combined with other techniques such as electrical or sensory stimulation.[144–148]
- Use of varying techniques based on transcranial magnetic stimulation has demonstrated growing evidence for lasting effects on swallowing in different groups.[149–151]

Surgery/radiological procedures

- Cricopharyngeal dysfunction may be treated with a surgical myotomy, botulinum toxin injection or balloon dilatation
- Vocal fold injection to increase glottis closure for patients with a vocal fold paralysis
- Stenting to maintain or dilate narrowed passage[152]

Oral health

Dysphagia combined with poor oral hygiene increases the risk of aspiration pneumonia. In addition, certain medications can reduce salivary flow, creating a more favourable environment for growth of bacteria as well as lack of lubrication for bolus formation and transport. Regular dental examination combined with good oral care has been shown to reduce these risks.[29,101,102] Artificial saliva sprays can be used to aid lubrication.

Medications

Patients with dysphagia are also likely to be taking a number of medications. Drugs that create a dry mouth, bring about motor fluctuations, reduce alertness, depress reflexes, increase reflux or induce nausea may negatively impact on swallowing and require review to optimize the balance between benefits for co-morbidities and side-effects for swallowing.[153]

The greater the number of medications, the higher the risk of aspiration pneumonia. Altering the medication such as crushing tablets or discontinuation of medication is relatively common practice in dysphagic individuals.[153] Furthermore, dysphagia may be induced or exacerbated by tissue injury resulting from unswallowed medication. Certain formulations such as multiparticulates, orodispersible tablets, skin patches and syrups may facilitate delivery of medication.[153]

Feeding tubes

Use of nasogastric tubes or insertion of PEG (percutaneous endoscopic gastrostomy) feeding may be employed as short or longer term solutions to dysphagia.[154–156] However, placement is not without controversy and further, whilst tube feeding may solve problems around nutritional intake, it does not eliminate problems around aspiration and infection. Pathogens may still enter the lungs from oral-nasal secretions and gastric reflux.

European Society of Clinical Nutrition and Metabolism guidelines[157] recommend tube feeding in patients with severe neurological dysphagia, but not in final disease state, including end-stage dementia. Use of enteral feeding in end-stage condition remains controversial (see below, ethical issues), not least because of the uncertainty in predicting or confirming end-stage state. Reservations around enteral feeding in such situations centre on failure to lengthen survival, and increased mortality in hospital, particularly for frail elderly people. However, as highlighted below, the evidence base awaits further development.

PEG tubes may be placed in patients where nutrition is an issue and for prevention of associated problems such as pressure sore, though PEG use itself also carries risks and is not beneficial in late dementia.[155,158] There appears to be no survival advantage to insertion in nursing home residents. Further, tube feeding exacerbates social isolation and is implicated in depression and reduced quality of life.

https://doi.org/10.1017/S095925981300021X Published online by Cambridge University Press

Prognosis

Aside from the more general effects of the underlying physical or neurological aetiology, prognosis for dysphagia is poor where there is insufficient insight, critically low levels of attention and awareness, lack of motivation, limited comprehension, severe memory impairment or rapidly degenerating physical condition. Compliance with interventions is negatively affected amongst other things by non-awareness or denial of a problem, underplaying the severity of a problem, assuming a calculated risk for non-compliance, dissatisfaction with food preparations, and perception of social stigma.

Many patients have limited ability to follow swallowing recommendations or adhere to a therapy programme due to language, cognitive or other physical impairment. They are therefore often reliant on their family or care home staff for feeding. The corollary is that prognosis is better if the skills and needs of carers are attended to. Education and ongoing support for families on discharge is vital.[12,13] Support in how to cope with real or perceived social stigma because of altered eating habits is essential to ensuring that the individual's and carers' roles within the family and social circle are maintained.[159] A high level of non-adherence with swallowing advice has been recorded, which significantly improves with the introduction of education and training of care staff about how feeding and swallowing needs can be incorporated into daily routines of patient and carer alike.

Ethical considerations

Decisions around swallowing management raise considerable ethical and legal issues, especially in relation to end of life care, and especially where capacity for consent is impaired or absent. Decisions around nasogastric or PEG feeding and withdrawal of feeding support are particularly fraught areas. Issues also arise in non-terminally ill patients, for instance where individuals or families decide not to follow recommendations. Detailed coverage of issues and potential solutions is not possible here. Guideline documents and discussions are available around palliative care and people with dementia.[157,158,160–166]

The direction of discussions in this literature is that there remains much to do in clarifying the evidence base around decisions that need to be made – for instance extensive literature is available in relation to outcomes around PEG placement,[158,164] but how this is applied to specific populations or individuals (imminent *vs* non-imminent death; lacking capacity or not; different medical aetiologies; different co-morbidities present) awaits definitive answers.[161,162] There is also evidence suggesting, that, for some professional groups at least, deeper education is necessary.[68,124]

Different groups[167,168] have attempted to formulate algorithms for decision making when patients or family decline to follow recommended treatments. These emphasize the educational and explanation role of clinicians, in assuring that information has not just been heard but also understood and focuses on shared responsibility and close monitoring of outcomes. All guidelines and discussions stress the importance of holistic assessment, shared decision making across all parties, with the autonomy and best interests of the affected individual placed at the centre. This may involve, on occasions, apparently counterintuitive decisions such as continuation of oral feeding despite tube placement or continuation of thin liquids despite demonstration of aspiration.[97–99,169] Careful charting and outcome measurement in such instances helps to further the evidence base around decision making and management.

Conflicts of interest

There are no conflicts of interest to declare.

References

1 Cichero JAY, Altman KW. Definition, prevalence and burden of oropharyngeal dysphagia: a serious problem among older adults worldwide and the impact on prognosis and hospital resources. In Cichero J, Clave P (eds), *Stepping Stones to Living Well with Dysphagia*, pp. 1–11; 2012.
2 Altman KW, Yu G-P, Schaefer SD. Consequence of dysphagia in the hospitalized patient impact on prognosis and hospital resources. *Arch Otolaryngol Head Neck Surg* 2010; **136**: 784–89.
3 Cabre M, Serra-Prat M, Palomera E, Almirall J, Pallares R, Clavé P. Prevalence and prognostic implications of dysphagia in elderly patients with pneumonia. *Age Ageing* 2010; **39**: 39–45.

https://doi.org/10.1017/S095925981300021X Published online by Cambridge University Press

4 Bouchard J, Presse N, Ferland G. Association between aspiration and malnutrition in patients from active geriatric units. *Canadian J Dietetic Practice Res* 2009; **70**: 152–54.

5 Hibberd J, Fraser J, Chapman C, McQueen H, Wilson A. Can we use influencing factors to predict aspiration pneumonia in the United Kingdom? *Multidisciplinary Resp Med* 2013; **8**: 39.

6 Masrur S, Smith EE, Saver JL, Reeves MJ, Bhatt DL, Zhao X *et al.* Dysphagia screening and hospital-acquired pneumonia in patients with acute ischemic stroke: findings from Get with the Guidelines – Stroke. *J Stroke Cerebrovasc Dis* (in press); http://dx.doi.org/10.1016/j.jstrokecerebrovasdis.2012.11.013

7 Robinson DJ, Jerrard-Dunne P, Greene Z, Lawson S, Lane S, O'Neill D. Oropharyngeal dysphagia in exacerbations of chronic obstructive pulmonary disease. *Eur Geriatric Med* 2011; **2**: 201–3.

8 van der Maarel-Wierink CD, Vanobbergen JNO, Bronkhorst EM, Schols JMGA, de Baat C. Meta-analysis of dysphagia and aspiration pneumonia in frail elders. *J Dent Res* 2011; **90**: 1398–404.

9 van der Maarel-Wierink CD, Vanobbergen JNO, Bronkhorst EM, Schols JMGA, de Baat C. Risk factors for aspiration pneumonia in frail older people: a systematic literature review. *J Am Medical Directors Assoc* 2011; **12**: 344–54.

10 Crary M, Humphrey J, Carnaby-Mann G, Sambandam R, Miller L, Silliman S. Dysphagia, nutrition, and hydration in ischemic stroke patients at admission and discharge from acute care. *Dysphagia* 2013; **28**: 69–76.

11 Serra-Prat M, Palomera M, Gomez C, Sar-Shalom D, Saiz A, Montoya JG *et al.* Oropharyngeal dysphagia as a risk factor for malnutrition and lower respiratory tract infection in independently living older persons: a population-based prospective study. *Age Ageing* 2012; **41**: 376–81.

12 Patterson J, Rapley T, Carding P, Wilson J, McColl E. Head and neck cancer and dysphagia; caring for carers. *Psycho-Oncology* 2013 (in press).

13 Penner JL, McClement S, Lobchuk M, Daeninck P. Family members' experiences caring for patients with advanced head and neck cancer receiving tube feeding: a descriptive phenomenological study. *J Pain Symptom Management* 2012; **44**: 563–71.

14 Martino R, Beaton D, Diamant NE. Perceptions of psychological issues related to dysphagia differ in acute and chronic patients. *Dysphagia* 2010; **25**: 26–34.

15 Klinke M, Wilosn M, Hafsteinsdottir T, Jonsdottir H. Recognizing new perspectives in eating difficulties following stroke: a concept analysis. *Disability Rehabilitation* 2013; **35**: 1491–500.

16 Semenov YR, Starmer HM, Gourin CG. The effect of pneumonia on short-term outcomes and cost of care after head and neck cancer surgery. *Laryngoscope* 2012; **122**: 1994–2004.

17 Tian H, Abouzaid S, Sabbagh MN, Chen W, Gabriel S, Kahler KH *et al.* Health care utilization and costs among patients with AD with and without dysphagia. *Alzheimer Dis Assoc Disorders* 2013; **27**: 138–44.

18 Lee SI, Yoo JY, Kim M, Ryu JS. Changes of timing variables in swallowing of boluses with different viscosities in patients with dysphagia. *Archives Physical Med Rehabil* 2013; **94**: 120–26.

19 Choi KH, Ryu JS, Kim MY, Kang JY, Yoo SD. Kinematic analysis of dysphagia: significant parameters of aspiration related to bolus viscosity. *Dysphagia* 2011; **26**: 392–98.

20 Umemoto G, Tsuboi Y, Kitashima A, Furuya H, Kikuta T. Impaired food transportation in Parkinson's disease related to lingual bradykinesia. *Dysphagia* 2011; **26**: 250–55.

21 Hirota N, Konaka K, Ono T, Tamine K, Kondo J, Hori K *et al.* Reduced tongue pressure against the hard palate on the paralyzed side during swallowing predicts dysphagia in patients with acute stroke. *Stroke* 2010; **41**: 2982–84.

22 Gross RD, Atwood CW Jr, Ross SB, Eichhorn KA, Olszewski JW, Doyle PJ. The coordination of breathing and swallowing in Parkinson's disease. *Dysphagia* 2008; **23**: 136–45.

23 Gross RD, Atwood CW Jr, Ross SB, Olszewski JW, Eichhorn KA. The coordination of breathing and swallowing in chronic obstructive pulmonary disease. *Am J Resp Critical Care Med* 2009; **179**: 559–65.

24 Troche M, Huebner I, Rosenbek J, Okun M, Sapienza C. Respiratory-swallowing coordination and swallowing safety in patients with Parkinson's disease. *Dysphagia* 2011; **26**: 218–24.

25 Hammer MJ, Murphy CA, Abrams TM. Airway somatosensory deficits and dysphagia in Parkinson's disease. *J Parkinson's Dis* 2013; **3**: 39–44.

26 Fukai K, Takiguchi T, Ando Y, Aoyama H, Miyakawa Y, Ito G *et al.* Critical tooth number without subjective dysphagia. *Geriatr Gerontol Int* 2011; **11**: 482–87.

27 Furuta M, Komiya-Nonaka M, Akifusa S, Shimazaki Y, Adachi M, Kinoshita T *et al.* Interrelationship of oral health status, swallowing function, nutritional status, and cognitive ability

https://doi.org/10.1017/S095925981300021X Published online by Cambridge University Press

with activities of daily living in Japanese elderly people receiving home care services due to physical disabilities. *Community Dentistry Oral Epidemiol* 2013; **41**: 173–81.
28 Pace CC, McCullough GH. The association between oral microorgansims and aspiration pneumonia in the institutionalized elderly: review and recommendations. *Dysphagia* 2010; **25**: 307–22.
29 Tada A, Miura H. Prevention of aspiration pneumonia (AP) with oral care. *Archives Gerontol Geriatr* 2012; **55**: 16–21.
30 Brodsky MB, Abbott KV, McNeil MR, Palmer CV, Grayhack JP, Martin-Harris B. Effects of divided attention on swallowing in persons with idiopathic Parkinson's disease. *Dysphagia* 2012; **27**: 390–400.
31 Nobrega AC, Rodrigues B, Torres AC, Scarpel RD, Neves CA, Melo A. Is drooling secondary to a swallowing disorder in patients with Parkinson's disease? *Parkinsonism Rel Disorders* 2008; **14**: 243–45.
32 Steele CM. The physiology of deglutition and the pathophysiology and complications of oropharyngeal dysphagia. In Cichero J, Clave P (eds), *Stepping Stones to Living Well with Dysphagia*, pp. 13–17; 2012:
33 Teismann IK, Suntrup S, Warnecke T, Steinstraeter O, Fischer M, Floeel A *et al.* Cortical swallowing processing in early subacute stroke. *BMC Neurol* 2011; doi:10.1186/1471–2377–11–34.
34 Momosaki R, Abo M, Kakuda W, Uruma G. Which cortical area is related to the development of dysphagia after stroke? a single photon emission computed tomography study using novel analytic methods. *Eur Neurol* 2012; **67**: 74–80.
35 Humbert IA, German RZ. New directions for understanding neural control in swallowing: the potential and promise of motor learning. *Dysphagia* 2013; **28**: 1–10.
36 Altman KW. Oropharyngeal dysphagia pathophysiology, complications and science-based interventions. In Cichero J, Clave P (eds), *Stepping Stones to Living Well with Dysphagia*, pp. 119–26; 2012.
37 Macht M, Wimbish T, Clark BJ, Benson AB, Burnham EL, Williams A *et al.* Postextubation dysphagia is persistent and associated with poor outcomes in survivors of critical illness. *Critical Care* 2011; **15**: R231.
38 Siska PA, Ponnappan RK, Hohl JB, Lee JY, Kang JD, Donaldson WF III. Dysphagia after anterior cervical spine surgery a prospective study using the swallowing-quality of life questionnaire and analysis of patient comorbidities. *Spine* 2011; **36**: 1387–91.
39 Kalb S, Reis MT, Cowperthwaite MC, Fox DJ, Lefevre R, Theodore N *et al.* Dysphagia after anterior cervical spine surgery: incidence and risk factors. *World Neurosurgery* 2012; **77**: 183–87.
40 Danto J, DiCapua J, Nardi D, Pekmezaris R, Moise G, Lesser M *et al.* Multiple cervical levels: increased risk of dysphagia and dysphonia during anterior cervical discectomy. *J Neurosurg Anesthesiol* 2012; **24**: 350–55.
41 Ruoppolo G, Schettino I, Frasca V, Giacomelli E, Prosperini L, Cambieri C *et al.* Dysphagia in amyotrophic lateral sclerosis: prevalence and clinical findings. *Acta Neurologica Scandinavica* 2013 (in press). doi: 10.1111/ane.12136
42 Miller N, Allcock L, Hildreth AJ, Jones D, Noble E, Burn DJ. Swallowing problems in Parkinson disease: frequency and clinical correlates. *J Neurol Neurosurg Psychiatry* 2009; **80**: 1047–49.
43 O'Sullivan SS, Massey LA, Williams DR, Silveira-Moriyama L, Kempster PA, Holton JL *et al.* Clinical outcomes of progressive supranuclear palsy and multiple system atrophy. *Brain* 2008; **131**: 1362–72.
44 Suntrup S, Teismann I, Bejer J, Suttrup I, Winkels M, Mehler D *et al.* Evidence for adaptive cortical changes in swallowing in Parkinson's disease. *Brain* 2013; **136**: 726–38.
45 Aldridge K, Taylor N. Dysphagia is a common and serious problem for adults with mental illness: a systematic review. *Dysphagia* 2012; **27**: 124–37.
46 Alagiakrishnan K, Bhanji RA, Kurian M. Evaluation and management of oropharyngeal dysphagia in different types of dementia: A systematic review. *Archives Gerontol Geriatr* 2013; **56**: 1–9.
47 Brodsky M, Abbott K, McNeil M, Palmer C, Grayhack J, Martin-Harris B. Effects of divided attention on swallowing in persons with idiopathic Parkinson's disease. *Dysphagia* 2012; **27**: 390–400.
48 Han M, Ohnishi H, Nonaka M, Yamauchi R, Hozuki T, Hayashi T *et al.* Relationship between dysphagia and depressive states in patients with Parkinson's disease. *Parkinsonism Rel Disorders* 2011; **17**: 437–39.
49 Holland G, Jayasekeran V, Pendleton N, Horan M, Jones M, Hamdy S. Prevalence and symptom profiling of oropharyngeal dysphagia in a community dwelling of an elderly population: a self-reporting questionnaire survey. *Diseases Esophagus* 2011; **24**: 476–80.
50 Daniels SK, Schroeder MF, DeGeorge PC, Corey DM, Foundas AL, Rosenbek JC. Defining and measuring dysphagia following stroke. *Am J Speech Lang Pathol* 2009; **18**: 74–81.

https://doi.org/10.1017/S095925981300021X Published online by Cambridge University Press

51 Ding RY, Logemann JA. Patient self-perceptions of swallowing difficulties as compared to expert ratings of videofluorographic studies. *Folia Phoniatr Logop* 2008; **60**: 142–50.

52 Bayés-Rusiñol À, Forjaz MJ, Ayala A, Crespo MC, Prats A, Valles E *et al.* [Awareness of dysphagia in Parkinson's disease]. *Rev Neurologia* 2011; **53**: 664–72.

53 Cheon S-M, Ha M-S, Park MJ, Kim JW. Non-motor symptoms of Parkinson's disease: prevalence and awareness of patients and families. *Parkinsonism Rel Dis* 2008; **14**: 286–90.

54 Serra-Prat M, Hinojosa G, Lopez D, Juan M, Fabre E, Voss DS *et al.* Prevalence of oropharyngeal dysphagia and impaired safety and efficacy of swallow in independently living older persons. *J Am Geriatr Soc* 2011; **59**: 186–87.

55 Roy N, Stemple J, Merrill RM, Thomas L. Dysphagia in the elderly: Preliminary evidence of prevalence, risk factors, and socioemotional effects. *Annals Otol Rhinol Laryngol* 2007; **116**: 858–65.

56 Nogueira D, Reis E. Swallowing disorders in nursing home residents: how can the problem be explained? *Clin Interventions Aging* 2013; **8**: 221–27.

57 NICE. Stroke rehabilitation: long term rehabilitation after stroke. London, UK: National Institute for Health and Care Excellence, 2013.

58 SIGN. Management of Patients with Stroke: Identification and Management of Dysphagia SIGN Guideline 119. Published by the Scottish Intercollegiate Guidelines Network; 2010. Available at: wwwsignacuk/guidelines/published/indexhtml#CHD

59 Excellence NIfC. Improving Outcomes in Head and Neck Cancers. 2004 [cited 26 July 2013]; available at: www.nice.org.uk

60 Trust SGHN. Tracheostomy Guidelines. 2013 [cited 27 July 2013]; available at: http://www.stgeorges.nhs.uk/trachindex.asp

61 Bryant KN, Finnegan E, Berbaum K. VFS Interjudge reliability using a free and directed search. *Dysphagia* 2012; **27**: 53–63.

62 Tohara H, Nakane A, Murata S, Mikushi S, Ouchi Y, Wakasugi Y *et al.* Inter- and intra-rater reliability in fibroptic endoscopic evaluation of swallowing. *J Oral Rehabil* 2010; **37**: 884–91.

63 Takahashi N, Kikutani T, Tamura F, Groher M, Kuboki T. Videoendoscopic assessment of swallowing function to predict the future incidence of pneumonia of the elderly. *J Oral Rehabil* 2012; **39**: 429–37.

64 Martin-Harris B, Brodsky MB, Michel Y, Castell DO, Schleicher M, Sandidge J *et al.* MBS Measurement tool for swallow impairment – MBSimp: establishing a standard. *Dysphagia* 2008; **23**: 392–405.

65 Murguia M, Corey DM, Daniels SK. Comparison of sequential swallowing in patients with acute stroke and healthy adults. *Archives Physical Med Rehabil* 2009; **90**: 1860–65.

66 Hind JA, Gensler G, Brandt DK, Gardner PJM, Blumenthal L, Gramigna GD *et al.* Comparison of trained clinician ratings with expert ratings of aspiration on videofluoroscopic images from a randomized clinical trial. *Dysphagia* 2009; **24**: 211–17.

67 Macht M, Wimbish T, Clark BJ, Benson AB, Burnham EL, Williams A *et al.* Diagnosis and treatment of post-extubation dysphagia: results from a national survey. *J Critical Care* 2012; **27**: 578–86.

68 Sharp HM, Shega JW. Feeding tube placement in patients with advanced dementia: the beliefs and practice patterns of speech-language pathologists. *Am J Speech Lang Pathol* 2009; **18**: 222–30.

69 Cocks N, Ferreira H. What Information do UK speech and language therapists use when making oral versus nonoral feeding recommendations for adults with oropharyngeal dysphagia? *Dysphagia* 2013; **28**: 43–57.

70 Archer SK, Wellwood I, Smith CH, Newham DJ. Dysphagia therapy in stroke: a survey of speech and language therapists. *Int J Lang Commun Disord* 2013 (in press). doi: 10.1111/1460–6984.12006.

71 Bours GJJW, Speyer R, Lemmens J, Limburg M, De Wit R. Bedside screening tests vs. videofluoroscopy or fibreoptic endoscopic evaluation of swallowing to detect dysphagia in patients with neurological disorders: systematic review. *J Adv Nurs* 2009; **65**: 477–93.

72 Umay E, Unlu E, Saylam G, Cakci A, Korkmaz H. Evaluation of dysphagia in early stroke patients by bedside, endoscopic, and electrophysiological methods. *Dysphagia* 2013 (in press). doi: 10.1007/s00455-013-9447-z.

73 Patterson JM, Hildreth A, McColl E, Carding PN, Hamilton D, Wilson JA. The clinical application of the 100 ml water swallow test in head and neck cancer. *Oral Oncology* 2011; **47**: 180–84.

74 Clavé P, Arreola V, Romea M, Medina L, Palomera E, Serra-Prat M. Accuracy of the volume-viscosity swallow test for clinical screening of oropharyngeal dysphagia and aspiration. *Clinical Nutrition* 2008; **27**: 806–15.

75 Martino R, Silver F, Teasell R, Bayley M, Nicholson G, Streiner DL *et al.* The Toronto Bedside Swallowing Screening Test (TOR-BSST): development and validation of a dysphagia

https://doi.org/10.1017/S095925981300021X Published online by Cambridge University Press

screening tool for patients with stroke. *Stroke* 2009; **40**: 555–61.
76 Antonios N, Carnaby-Mann G, Crary M, Miller L, Hubbard H, Hood K *et al.* Analysis of a physician tool for evaluating dysphagia on an inpatient stroke unit: the modified Mann Assessment of Swallowing Ability. *J Stroke Cerebrovasc Dis* 2010; **19**: 49–57.
77 Gonzalez-Fernandez M, Sein MT, Palmer JB. Clinical experience using the Mann assessment of swallowing ability for identification of patients at risk for aspiration in a mixed-disease population. *Am J Speech Lang Pathol* 2011; **20**: 331–36.
78 Leder SB, Suiter DM, Warner HL, Acton LM, Siegel MD. Safe initiation of oral diets in hospitalized patients based on passing a 3-ounce (90 cc) water swallow challenge protocol. *QJM* 2012; **105**: 257–63.
79 Schepp SK, Tirschwell DL, Miller RM, Longstreth WT. Swallowing screens after acute stroke: a systematic review. *Stroke* 2012; **43**: 869–71.
80 Turner-Lawrence DE, Peebles M, Price MF, Singh SJ, Asimos AW. A feasibility study of the sensitivity of emergency physician dysphagia screening in acute stroke patients. *Annals Emergency Med* 2009; **54**: 344–48.
81 Edmiaston J, Connor LT, Loehr L, Nassief A. Validation of a dysphagia screening tool in acute stroke patients. *Am J Critical Care* 2010; **19**: 357–64.
82 Wilkinson AH, Burns SL, Witham MD. Aspiration in older patients without stroke: A systematic review of bedside diagnostic tests and predictors of pneumonia. *Eur Geriatr Med* 2012; **3**: 145–52.
83 Daniels S. Letter by Daniels regarding article 'Silent aspiration risk is volume-dependent'. *Dysphagia* 2012; **27**: 294.
84 Osawa A, Maeshima S, Tanahashi N. Water-swallowing test: screening for aspiration in stroke patients. *Cerebrovasc Dis* 2013; **35**: 276–81.
85 Daniels SK, Anderson JA, Willson PC. Valid items for screening dysphagia risk in patients with stroke: a systematic review. *Stroke* 2012; **43**: 892–97.
86 Waito A, Bailey G, Molfenter S, Zoratto D, Steele C. Voice-quality abnormalities as a sign of dysphagia: validation against acoustic and videofluoroscopic data. *Dysphagia* 2011; **26**: 125–34.
87 Bruijn M, Rinkel RPM, Cnossen I, Witte B, Langendijk J, Leemans CR *et al.* Associations between voice quality and swallowing function in patients treated for oral or oropharyngeal cancer. *Support Care Cancer* 2013; **21**: 2025–32.
88 Chang H-Y, Torng P-C, Wang T-G, Chang Y-C. Acoustic voice analysis does not identify presence of penetration/aspiration as confirmed by videofluoroscopic swallowing study. *Archives Physical Med Rehabil* 2012; **93**: 1991–94.
89 Malandraki GA, Hind JA, Gangnon R, Logemann JA, Robbins J. The utility of pitch elevation in the evaluation of oropharyngeal dysphagia: preliminary findings. *Am J Speech Lang Pathol* 2011; **20**: 262–68.
90 Sarraf Shirazi S, Buchel C, Daun R, Lenton L, Moussavi Z. Detection of swallows with silent aspiration using swallowing and breath sound analysis. *Med Biol Eng Comput* 2012; **50**: 1261–68.
91 Sato M, Tohara H, Iida T, Wada S, Inoue M, Ueda K. Simplified cough test for screening silent aspiration. *Archives Physical Med Rehabil* 2012; **93**: 1982–86.
92 Miles A, Huckabee M-L. Intra- and inter-rater reliability for judgement of cough following citric acid inhalation. *Int J Speech-Language Pathol* 2013;**15**: 209–15.
93 Miles A, Zeng ISL, McLauchlan H, Huckabee M-L. Cough reflex testing in dysphagia following stroke: a randomized controlled trial. *J Clin Med Res* 2013; **5**: 222–33.
94 Pitts T, Troche M, Mann G, Rosenbek J, Okun MS, Sapienza C. Using voluntary cough to detect penetration and aspiration during oropharyngeal swallowing in patients with Parkinson disease. *Chest* 2010; **138**: 1426–31.
95 Bianchi C, Baiardi P, Khirani S, Cantarella G. Cough peak flow as a predictor of pulmonary morbidity in patients with dysphagia. *Am J Physical Med Rehabil* 2012; **91**: 783–88.
96 Garon BR, Sierzant T, Ormiston C. Silent aspiration: results of 2000 video fluoroscopic evaluations. *J Neurosci Nursing* 2009; **41**: 178–85.
97 Frey KL, Ramsberger G. Comparison of outcomes before and after implementation of a water protocol for patients with cerebrovascular accident and dysphagia. *J Neurosci Nursing* 2011; **43**: 165–71.
98 Karagiannis MJP, Chivers L, Karagiannis TC. Effects of oral intake of water in patients with oropharyngeal dysphagia. *BMC Geriatrics* 2011; **11**: **9**. doi:10.1186/1471-2318-11-9.
99 Carlaw C, Finlayson H, Beggs K, Visser T, Marcoux C, Coney D *et al.* Outcomes of a pilot water protocol project in a rehabilitation setting. *Dysphagia* 2012; **27**: 297–306.
100 Ewan V, Perry JD, Mawson T, McCracken G, Brown AN, Newton J *et al.* Detecting potential respiratory pathogens in the mouths of older people in hospital. *Age Ageing* 2010; **39**: 122–25.

https://doi.org/10.1017/S095925981300021X Published online by Cambridge University Press

101 Pace C, McCullough G. The association between oral microorganisms and aspiration pneumonia in the institutionalized elderly: review and recommendations. *Dysphagia* 2010; **25**: 307–22.

102 van der Maarel-Wierink CD, Vanobbergen JNO, Bronkhorst EM, Schols JMGA, de Baat C. Oral health care and aspiration pneumonia in frail older people: a systematic literature review. *Gerodontology* 2013; **30**: 3–9.

103 Abe S, Ishihara K, Adachi M, Okuda K. Tongue-coating as risk indicator for aspiration pneumonia in edentate elderly. *Archives Gerontol Geriatr* 2008; **47**: 267–75.

104 Hansen T, Kjaersgaard A, Faber J. Measuring elderly dysphagic patients' performance in eating – a review. *Disability Rehabil* 2011; **33**: 1931–40.

105 Westergren A, Lindholm C, Mattsson A, Ulander K. Minimal eating observation form: reliability and validity. *J Nutrition Health Aging* 2009; **13**: 6–11.

106 Lambert HC, Gisel EG, Groher ME, Abrahamowicz M, Wood-Dauphinee S. Psychometric testing of the McGill Ingestive Skills Assessment. *Am J Occupational Therapy* 2006; **60**: 409–19.

107 Silbergleit A, Schultz L, Jacobson B, Beardsley T, Johnson A. The Dysphagia Handicap Index: development and validation. *Dysphagia* 2012; **27**: 46–52.

108 McHorney CA, Robbins J, Lomax K, Rosenbek JC, Chignell K, Kramer AE *et al.* The SWAL-QOL and SWAL-CARE outcomes tool for oropharyngeal dysphagia in adults: III. Documentation of reliability and validity. *Dysphagia* 2002; **17**: 97–114.

109 Chen AY, Frankowski R, Bishop-Leone J, Hebert T, Leyk S, Lewin J *et al.* The development and validation of a dysphagia-specific quality-of-life questionnaire for patients with head and neck cancer: the M.D. Anderson Dysphagia Inventory. *Archives Otolaryngol Head Neck Surg* 2001; **127**: 870–76.

110 Chen P-H, Golub JS, Hapner ER, Johns MM III. Prevalence of perceived dysphagia and quality-of-life impairment in a geriatric population. *Dysphagia* 2009; **24**: 1–6.

111 Belafsky PC, Mouadeb DA, Rees CJ, Pryor JC, Postma GN, Allen J *et al.* Validity and reliability of the Eating Assessment Tool (EAT-10). *Ann Otol Rhinol Laryngol* 2008; **117**: 919–24.

112 Leow LP, Huckabee M-L, Anderson T, Beckert L. The impact of dysphagia on quality of life in ageing and Parkinson's disease as measured by the Swallowing Quality of Life (SWAL-QOL) Questionnaire. *Dysphagia* 2010; **25**: 216–20.

113 Wilson JA, Carding PN, Patterson JM. Dysphagia after nonsurgical head and neck cancer treatment: patients' perspectives. *Otolaryngol Head Neck Surg* 2011; **145**: 767–71.

114 Crary M, Sura L, Carnaby G. Validation and demonstration of an isolated acoustic recording technique to estimate spontaneous swallow frequency. *Dysphagia* 2013; **28**: 86–94.

115 Steele C, Sejdić E, Chau T. Non-invasive detection of thin-liquid aspiration using dual-axis swallowing accelerometry. *Dysphagia* 2013; **28**: 105–12.

116 Vaiman M, Eviatar E. Surface electromyography as a screening method for evaluation of dysphagia and odynophagia. *Head Face Med* 2009; **5**: 9.

117 Altman KW. Dysphagia evaluation and care in the hospital setting: the need for protocolization. *Otolaryngol Head Neck Surg* 2011; **145**: 895–98.

118 Lakshminarayan K, Tsai AW, Tong X, Vazquez G, Peacock JM, George MG *et al.* Utility of dysphagia screening results in predicting poststroke pneumonia. *Stroke* 2010; **41**: 2849–54.

119 Yeh S-J, Huang K-Y, Wang T-G, Chen Y-C, Chen C-H, Tang S-C *et al.* Dysphagia screening decreases pneumonia in acute stroke patients admitted to the stroke intensive care unit. *J Neurol Sci* 2011; **306**: 38–41.

120 Ickenstein GW, Riecker A, Höhlig C, Müller R, Becker U, Reichmann H *et al.* Pneumonia and in-hospital mortality in the context of neurogenic oropharyngeal dysphagia (NOD) in stroke and a new NOD step-wise concept. *J Neurol* 2010; **57**: 1492–99.

121 Wilson RD, Howe EC. A cost-effectiveness analysis of screening methods for dysphagia after stroke. *PM&R* 2012; **4**: 273–82.

122 King J, Ligman K. Patient non-compliance with swallowing recommendations: reports from speech-language pathologists. *Contemporary Issues Commun Sci Disorders* 2011; **38**: 53–60.

123 Foley N, Teasell R, Salter K, Kruger E, Martino R. Dysphagia treatment post stroke: a systematic review of randomised controlled trials. *Age Ageing* 2008; **37**: 258–64.

124 Vitale CA, Berkman CS, Monteleoni C, Ahronheim JC. Tube feeding in patients with advanced dementia: knowledge and practice of speech-language pathologists. *J Pain Symptom Management* 2011; **42**: 366–78.

125 Butler S, Pelletier C, Steele C. Compensatory strategies and techniques. In Shaker R, Easterling C, Belafsky PC, Postma GN (eds), *Manual of Diagnostic and Therapeutic Techniques for Disorders of Deglutition*, pp. 299–316. Springer, New York; 2013.

126 Lazarus C. Mendelson Maneuver and Masako Maneuver. In Shaker R, Easterling C, Belafsky PC, Postma GN (eds), *Manual of Diagnostic and*

https://doi.org/10.1017/S095925981300021X Published online by Cambridge University Press

Therapeutic Techniques for Disorders of Deglutition, pp. 269–80. Springer, New York; 2013.
127 Smith SK, Roddam H, Sheldrick H. Rehabilitation or compensation: time for a fresh perspective on speech and language therapy for dysphagia and Parkinson's disease? *Int J Lang Commun Disord* 2012 (in press). doi: 10.1111/j.1460–6984.2011.00093.x.
128 Wheeler-Hegland K, Ashford J, Frymark T, McCabe D, Mullen R, Musson N *et al.* Evidence-based systematic review: oropharyngeal dysphagia behavioral treatments. Impact of dysphagia treatment on normal swallow function. *J Rehabil Res Dev* 2009; **46**: 185–94.
129 Baylow HE, Goldfarb R, Taveira CH, Steinberg RS. Accuracy of clinical judgment of the chin-down posture for dysphagia during the clinical/bedside assessment as corroborated by videofluoroscopy in adults with acute stroke. *Dysphagia* 2009; **24**: 423–33.
130 Huckabee M-L, Macrae P. Effortful Swallow. In Shaker R, Easterling C, Belafsky PC, Postma GN (eds), *Manual of Diagnostic and Therapeutic Techniques for Disorders of Deglutition*, pp. 281–98. Springer, New York: 2013.
131 Leder S, Judson B, Sliwinski E, Madson L. Promoting safe swallowing when puree is swallowed without aspiration but thin liquid is aspirated: nectar is enough. *Dysphagia* 2013; **28**: 58–62.
132 Agency NPS. Dysphagia diet food textures descriptors. 2012 [cited 31 July 2013]; available at: http://www.bda.uk.com/publications/statements/NationalDescriptorsTexture-ModificationAdults.pdf
133 Sdravou K, Walshe M, Dagdilelis L. Effects of carbonated liquids on oropharyngeal swallowing measures in people with neurogenic dysphagia. *Dysphagia* 2012; **27**: 240–50.
134 Beck AM, Holst M, Rasmussen HH. Oral nutritional support of older (65 years+) medical and surgical patients after discharge from hospital: systematic review and meta-analysis of randomized controlled trials. *Clinical Rehabil* 2013; **27**: 19–27.
135 Beck AM, Kjær S, Hansen BS, Storm RL, Thal-Jantzen K, Bitz C. Follow-up home visits with registered dietitians have a positive effect on the functional and nutritional status of geriatric medical patients after discharge: a randomized controlled trial. *Clinical Rehabil* 2013; **27**: 483–93.
136 Geeganage C, Beavan J, Ellender S, Bath PMW. Interventions for dysphagia and nutritional support in acute and subacute stroke. *Cochrane Database Syst Rev* 2012; **10**.
137 Abbott RA, Whear R, Thompson-Coon J, Ukoumunne OC, Rogers M, Bethel A *et al.* Effectiveness of mealtime interventions on nutritional outcomes for the elderly living in residential care: a systematic review and meta-analysis. *Ageing Res Rev* 2013 (in press). doi: org/10.1016/j.arr.2013.06.002.
138 Logemann JA, Rademaker A, Pauloski BR, Kelly A, Stangl-McBreen C, Antinoja J *et al.* A randomized study comparing the shaker exercise with traditional therapy: a preliminary study. *Dysphagia* 2009; **24**: 403–11.
139 Easterling C. Shaker Exercise. In Shaker R, Easterling C, Belafsky PC, Postma GN (eds), *Manual of Diagnostic and Therapeutic Techniques for Disorders of Deglutition*, pp. 257–68. Springer, New York; 2013.
140 Troche MS, Okun MS, Rosenbek JC, Musson N, Fernandez HH, Rodriguez R *et al.* Aspiration and swallowing in Parkinson disease and rehabilitation with EMST: a randomized trial. *Neurology* 2010; **75**: 1912–19.
141 Steele CM, Bailey GL, Polacco REC, Hori SF, Molfenter SM, Oshalla M *et al.* Outcomes of tongue-pressure strength and accuracy training for dysphagia following acquired brain injury. *Int J Speech-Language Pathol* 2013 (in press). doi: 10.3109/17549507.2012.752864.
142 Kang J-H, Park R-Y, Lee S-J, Kim J-Y, Yoon S-R, Jung K-I. The effect of bedside exercise program on stroke patients with dysphagia. *Ann Rehabil Med* 2012; **36**: 512–20.
143 Hind J, Robbins J. Oropharyngeal strengthening and rehabilitation of deglutitive disorders. In Shaker R, Easterling C, Belafsky PC, Postma GN (eds), *Manual of Diagnostic and Therapeutic Techniques for Disorders of Deglutition*, pp. 237–55. Springer, New York; 2013.
144 Regan J, Walshe M, Tobin WO. Immediate effects of thermal-tactile stimulation on timing of swallow in idiopathic Parkinson's disease. *Dysphagia* 2010; **25**: 207–15.
145 Doeltgen SH, Dalrymple-Alford J, Ridding MC, Huckabee ML. Differential effects of neuromuscular electrical stimulation parameters on submental motor-evoked potentials. *Neurorehabil Neural Repair* 2010; **24**: 519–27.
146 Carnaby-Mann GD, Crary MA. Examining the evidence on neuromuscular electrical stimulation for swallowing – a meta-analysis. *Archives Otolaryngol Head Neck Surg* 2007; **133**: 564–71.
147 Clark H, Lazarus C, Arvedson J, Schooling T, Frymark T. Evidence-based systematic review: effects of neuromuscular electrical stimulation on swallowing and neural activation. *Am J Speech Lang Pathol* 2009; **18**: 361–75.

https://doi.org/10.1017/S095925981300021X Published online by Cambridge University Press

148 Humbert IA, Michou E, MacRae PR, Crujido L. Electrical stimulation and swallowing: how much do we know? *Seminars Speech Language* 2012; **33**: 203–16.
149 Michou E, Mistry S, Rothwell J, Hamdy S. Priming pharyngeal motor cortex by repeated paired associative stimulation: implications for dysphagia neurorehabilitation. *Neurorehabil Neural Repair* 2013; **27**: 355–62.
150 Khedr E, Abo-Elfetoh N, Rothwell J. Treatment of post-stroke dysphagia with repetitive transcranial magnetic stimulation. *Acta Neurologica Scandinavica* 2009; **119**: 155–61.
151 Shigematsu T, Fujishima I, Ohno K. Transcranial direct current stimulation improves swallowing function in stroke patients. *Neurorehabil Neural Repair* 2013; **27**: 363–69.
152 Repici A, Hassan C, Sharma P, Conio M, Siersema P. Systematic review: the role of self-expanding plastic stents for benign oesophageal strictures. *Alimentary Pharmacol Therapeutics* 2010; **31**: 1268–75.
153 Stegemann S, Gosch M, Breitkreutz J. Swallowing dysfunction and dysphagia is an unrecognized challenge for oral drug therapy. *Int J Pharmaceutics* 2012; **430**: 197–206.
154 Leder SB, Lazarus CL, Suiter DM, Acton LM. Effect of orogastric tubes on aspiration status and recommendations for oral feeding. *Otolaryngol Head Neck Surg* 2011; **144**: 372–75.
155 Gomes CAR Jr, Lustosa SAS, Matos D, Andriolo RB, Waisberg DR, Waisberg J. Percutaneous endoscopic gastrostomy versus nasogastric tube feeding for adults with swallowing disturbances. *Cochrane Database Syst Rev* 2012; **3**.
156 Beavan J, Conroy SP, Harwood R, Gladman JRF, Leonardi-Bee J, Sach T *et al.* Does looped nasogastric tube feeding improve nutritional delivery for patients with dysphagia after acute stroke? A randomised controlled trial. *Age Ageing* 2010; **39**: 624–30.
157 Sobotka L, Schneider SM, Berner YN, Cederholm T, Krznaric Z, Shenkin A *et al.* ESPEN Guidelines on parenteral nutrition: Geriatrics. *Clinical Nutrition* 2009; **28**: 461–66.
158 Kurien M, McAlindon ME, Westaby D, Sanders DS. Percutaneous endoscopic gastrostomy (PEG) feeding. *BMJ* 2010; **340**: c2414.
159 Johansson AEM, Johansson U. Relatives' experiences of family members' eating difficulties. *Scand J Occupational Therapy* 2009; **16**: 25–32.
160 Smith HA, Kindell J, Baldwin RC, Waterman D, Makin AJ. Swallowing problems and dementia in acute hospital settings: practical guidance for the management of dysphagia. *Clinical Med* 2009; **9**: 544–48.
161 Regnard C, Leslie P, Crawford H, Matthews D, Gibson L. Gastrostomies in dementia: bad practice or bad evidence? *Age Ageing* 2010; **39**: 282–84.
162 Chen H-L, Shih S-C, Bair M-J, Lin IT, Wu C-H. Percutaneous endoscopic gastrostomy in the enteral feeding of the elderly. *Int J Gerontol* 2011; **5**: 135–38.
163 Candy B, Sampson EL, Jones L. Enteral tube feeding in older people with advanced dementia: findings from a Cochrane systematic review. *Int J Palliative Nursing* 2009; **15**: 396–404.
164 Sampson EL, Candy B, Jones L. Enteral tube feeding for older people with advanced dementia. *Cochrane Database Syst Rev* 2009; **2**.
165 Clarke G, Harrison K, Holland A, Kuhn I, Barclay S. How are treatment decisions made about artificial nutrition for individuals at risk of lacking capacity? A systematic literature review. *Plos One* 2013; **16**: 8.
166 RCP. Oral feeding difficulties and dilemmas: a guide to practical care particularly towards end of life. London, UK: Royal College of Physicians; 2010.
167 Sandman L, Bolmsjo IA, Westergren A. Ethical considerations of refusing nutrition after stroke. *Nursing Ethics* 2008; **15**: 147–59.
168 Kaizer F, Spiridigliozzi A-M, Hunt M. Promoting shared decision-making in rehabilitation: development of a framework for situations when patients with dysphagia refuse diet modification recommended by the treating team. *Dysphagia* 2012; **27**: 81–87.
169 Becker R, Nieczaj R, Egge K, Moll A, Meinhardt M, Schulz R-J. Functional dysphagia therapy and PEG treatment in a clinical geriatric setting. *Dysphagia* 2011; **26**: 108–16.
170 Hirst Lisa J. Table 151.1, Comparison of videofluoroscopy with FEES and manometry, chapter 151, Functional investigation of the upper GI tract. In *Scott-Brown's Otolaryngology, Head and Neck Surgery*, 7th edn, editors: MJ Gleeson, G Browning, M Burton, R Clarke, J Hibbert, N Jones, V Lund, L Luxon, J Watkinson.

https://doi.org/10.1017/S095925981300021X Published online by Cambridge University Press