*Combinatorics, Probability and Computing* (2001) 10, 367–384. © 2001 Cambridge University Press
DOI: 10.1017/S0963548301004795 Printed in the United Kingdom

# *G*-Intersecting Families

TOM BOHMAN[1], ALAN FRIEZE[1]†,
MIKLÓS RUSZINKÓ[2]‡ and LUBOŠ THOMA[1]§

[1] Department of Mathematical Sciences,
Carnegie Mellon University, USA
(e-mail: tbohman@moser.math.cmu.edu,
alan@random.math.cmu.edu,
thoma@andrew.cmu.edu)

[2] Computer and Automation Research Institute of the Hungarian Academy of Sciences,
Budapest, PO Box 63, Hungary 1518
(e-mail: ruszinko@sztaki.hu)

*Received 1 October 1999; revised 30 November 2000*

Let $G$ be a graph on vertex set $[n]$, and for $X \subseteq [n]$ let $N(X)$ be the union of $X$ and its neighbourhood in $G$. A family of sets $\mathscr{F} \subseteq 2^{[n]}$ is *$G$-intersecting* if $N(X) \cap Y \neq \emptyset$ for all $X, Y \in \mathscr{F}$. In this paper we study the cardinality and structure of the largest $k$-uniform $G$-intersecting families on a fixed graph $G$.

## 1. Introduction

An *intersecting family* is a collection of sets $\mathscr{F} \subseteq 2^{[n]}$ such that

$$X, Y \in \mathscr{F} \Rightarrow X \cap Y \neq \emptyset.$$

The classic Erdős–Ko–Rado theorem [5] states that a $k$-uniform intersecting family on a ground set of size $n$ contains at most $\binom{n-1}{k-1}$ sets. Furthermore, a family achieves this extremum if and only if the family consists of all $k$-sets containing some fixed element of $[n]$. For a general introduction to the theory of intersecting families see [3].

We consider a generalization of the notion of an intersecting family (for a survey of other generalizations of the Erdős–Ko–Rado theorem see [4]).

† Supported in part by NSF grant CCR-9530974.
‡ Supported in part by OTKA Grants T 030059 and T 29074, FKFP 0607/1999 and by the Bolyai Foundation.
§ Supported in part by NSF grant DMS-9970622.

https://doi.org/10.1017/S0963548301004795 Published online by Cambridge University Press

First consider the following. In an intersecting family $\mathscr{F}$ we require that $X, Y \in \mathscr{F}$ should intersect. Suppose that we relax this to require simply that $X, Y \in \mathscr{F}$ are 'close' in some sense. Can something interesting be said? We believe that it can. For example, suppose that we take '$X, Y$ are close' to mean that there exist $x \in X, y \in Y$ such that $\mathrm{dist}(x, y) = |x - y| \leqslant 1$ (it will be convenient to have $\mathrm{dist}(n, 1) = 1$ here). What can we say about the maximum size of a $k$-uniform family where every pair of distinct sets are close? A moment's thought suggests that a candidate for a maximal $\mathscr{F}$ is the set of $k$-sets which contain 2 or 3. This is almost correct. Throwing in those sets which contain both 1 and 4 gives the correct answer, at least for $n$ sufficiently large,

$$|\mathscr{F}| \leqslant 2\binom{n-1}{k-1} - \binom{n-2}{k-2} + \binom{n-4}{k-2}$$

with equality only if $\mathscr{F}$ has the described structure (see Corollary 1.2).

Now for our full generalization. Let $G$ be a graph on vertex set $[n]$. (The above example can be considered to be the case where $G$ is a cycle.) For $X \subseteq [n]$ let

$$N(X) = X \cup \{y \in [n] : \exists x \in X \text{ such that } x \sim_G y\}.$$

We say that a family of sets $\mathscr{F} \subseteq 2^{[n]}$ is *$G$-intersecting* if

$$X, Y \in \mathscr{F} \Rightarrow N(X) \cap Y \neq \emptyset.$$

In other words, $\mathscr{F}$ is $G$-intersecting if, for all $X, Y \in \mathscr{F}$, there exist $x \in X$ and $y \in Y$ such that $x = y$ or $x \sim_G y$. For a fixed graph $G$, how large can a $k$-uniform $G$-intersecting family be? For a graph $G$ on vertex set $[n]$ and $k$ a positive integer, let

$$N(G, k) = \max\left\{|\mathscr{F}| : \mathscr{F} \subseteq \binom{[n]}{k} \text{ and } \mathscr{F} \text{ is } G\text{-intersecting}\right\}.$$

This quantity can be easily computed in some cases. For example, for the complete graph we clearly have $N(K_n, k) = \binom{n}{k}$. Let $G = K_{a,n-a}$ be the complete bipartite graph having parts $A, B \subseteq [n]$. A $G$-intersecting family $\mathscr{F}$ can be partitioned into three families, $\mathscr{F} = \mathscr{F}_1 \cup \mathscr{F}_2 \cup \mathscr{F}_3$, where $X \in \mathscr{F}_1$ implies $X$ meets both $A$ and $B$, $\mathscr{F}_2$ is an intersecting family of subsets of $A$, and $\mathscr{F}_3$ is an intersecting family of subset of $B$. It follows that

$$N(K_{a,n-a}, k) = \binom{n}{k} - \binom{a}{k} - \binom{n-a}{k} + \binom{a-1}{k-1} + \binom{n-a-1}{k-1}. \tag{1.1}$$

As another simple example, consider the case of the union $H_p$ of a clique on $\{1, 2, \ldots, p\}$ and an independent set on $\{p+1, p+2, \ldots, n\}$. Here we can show that

$$N(H_p, k) = \begin{cases} \binom{n}{k} - \binom{n-p}{k}, & k \leqslant n/2, \\ \binom{n}{k}, & k > n/2. \end{cases}$$

The case $k > n/2$ is trivial, and so we assume that $k \leqslant n/2$. We use induction on $p$. $H_1$ is an independent set and so $\mathscr{F}$ is $H_1$-intersecting if and only if it is intersecting. Thus $|\mathscr{F}| \leqslant \binom{n-1}{k-1} = \binom{n}{k} - \binom{n-1}{k}$. For $p \geqslant 2$ let $\mathscr{F} = \mathscr{F}_1 \cup \mathscr{F}_2$, where $\mathscr{F}_1 = \{X \in \mathscr{F} : p \notin X\}$. $\mathscr{F}_1$ is $H'_{p-1}$-intersecting where $H'_{p-1} = H_p - \{p\}$ and so, by induction, $|\mathscr{F}_1| \leqslant \binom{n-1}{k} - \binom{n-p}{k}$. Clearly $|\mathscr{F}_2| \leqslant \binom{n-1}{k-1}$ and the result follows. Taking $\mathscr{F} = \{X : X \cap [p] \neq \emptyset\}$ shows the bound is tight.

https://doi.org/10.1017/S0963548301004795 Published online by Cambridge University Press

The extremal $G$-intersecting families are analogous to the extremal intersecting families given by Erdős–Ko–Rado for sparse $G$ and small $k$. To make this statement more precise, we first note that, if $K \subseteq [n]$ is a clique in $G$, then

$$\mathscr{F}(K) := \left\{ X \subseteq \binom{[n]}{k} : X \cap K \neq \emptyset \right\} \tag{1.2}$$

is a $G$-intersecting family. It follows that

$$N(G,k) \geqslant \binom{n}{k} - \binom{n-\omega}{k}, \tag{1.3}$$

where $\omega = \omega(G)$ is the clique number of $G$. Note that if $G$ is the empty graph, then Erdős–Ko–Rado implies $N(G,k) = \binom{n-1}{k-1}$, $N(G,k)$ equals the bound given in (1.3), and any extremal family is of the form given in (1.2). However, for most graphs we cannot replace the inequality in (1.3) with equality. If $K$ is a clique in $G$ and $M_1, M_2, \ldots, M_r \subseteq [n]\backslash K$ satisfy

$$K \subseteq N(M_i) \text{ for } i = 1, \ldots, r \quad \text{and} \quad M_i \cap N(M_j) \neq \emptyset, \quad \text{for } i \neq j, \tag{1.4}$$

then the collection

$$\mathscr{F}(K; M_1, \ldots, M_r) = \left\{ X \subseteq \binom{[n]}{k} : X \cap K \neq \emptyset \text{ or } M_i \subseteq X \text{ for some } i \right\} \tag{1.5}$$

is $G$-intersecting. Thus, if there exists a maximum clique $K$ and $M \subseteq [n]\backslash K$ such that $|M| \leqslant k$ and $K \subseteq N(M)$, then the bound given in (1.3) is not the truth. However, for some graphs we are able to show that, if $\mathscr{F}$ is a large $G$-intersecting family, then $\mathscr{F}$ is of the form given in (1.5).

**Theorem 1.1.** *Let $G$ be a graph on vertex set $[n]$ and $k$ a positive integer such that*

$$n \geqslant \frac{2e(1+\Delta)\Delta'(k-1)^2 k(k+1)}{\omega},$$

*where $\Delta = \Delta(G)$ is the maximum degree of $G$, $\omega = \omega(G)$ is the clique number of $G$, and $\Delta' = \Delta'(G)$ is the maximum size of a second neighbourhood in $G$:*

$$\Delta'(G) := \max_{v \in [n]} |\{i \in [n] : \operatorname{dist}_G(i,v) \leqslant 2\}|.$$

*If $\mathscr{F}$ is a $G$-intersecting family containing more than $\binom{n}{k} - \binom{n-\omega}{k}$ sets, then there exists a clique $K$ in $G$ and $M_1, \ldots, M_r \subseteq [n]\backslash K$ such that* (1.4) *holds and $\mathscr{F} = \mathscr{F}(K; M_1, \ldots, M_r)$.*

In some sense, Theorem 1.1 is analogous to the results of Hilton and Milner [7] and Hajnal and Rothschild [6] on intersecting systems, which state that an intersecting family that is 'nearly' maximum is 'very similar' in structure to the maximum intersecting families.

Note that Theorem 1.1 contains information about neither the clique $K$ (*e.g.*, must it be a clique on $\omega(G)$ vertices?) nor the collection of sets $M_1, \ldots, M_r$. Therefore, when we apply this theorem we must optimize over all possible choices of these parameters. We do this for three classes of graphs. For an arbitrary graph $G$ we define the $p$th power of $G$, denoted $G^p$, to be the graph with vertex set $V(G)$ and an edge between vertices $u$ and $v$ if and only if there is a path of length at most $p$ between $u$ and $v$ in $G$.

https://doi.org/10.1017/S0963548301004795 Published online by Cambridge University Press

**Corollary 1.2.** *Let $G = C_n^p$ be the $p$th power of a cycle on $n$ vertices and let $k$ be a positive integer. If*

$$n \geqslant \frac{2e(2p+1)(4p+1)(k-1)^2k(k+1)}{p+1} \tag{1.6}$$

*then*

$$N(G,k) = \binom{n}{k} - \binom{n-2p-1}{k} - p\binom{n-2p-2}{k-1}. \tag{1.7}$$

*Furthermore, a $G$-intersecting family $\mathscr{F}$ has cardinality $N(G,k)$ if and only if $\mathscr{F} = \mathscr{F}\left(K, M_1, \dots, M_{\binom{p+1}{2}}\right)$, where $K$ is a clique on $p+1$ elements and*

$$\left\{M_1, \dots, M_{\binom{p+1}{2}}\right\} = \left\{\{i,j\} \in \binom{[n]\setminus K}{2} : i \not\sim_G j \text{ and } K \subseteq N(\{i,j\})\right\}.$$

**Corollary 1.3.** *Let $G$ be a $d$-regular graph on vertex set $[n]$ having no cycle on 5 or fewer vertices. If*

$$k < d,$$

$$n \geqslant (k+1) + 2e(k-2)k(d+1)(d-1)^k \quad \text{and} \tag{1.8}$$

$$n \geqslant e(d+1)(d^2+d+1)(k-1)^2k(k+1), \tag{1.9}$$

*then*

$$N(G,k) = \binom{n}{k} - \binom{n-2}{k} + \sum_{i=1}^{k-2}\left[\sum_{j=0}^{i}(-1)^j\binom{i}{j}\binom{n-j(d-1)-i-2}{k-1}\right]. \tag{1.10}$$

*Furthermore, a $G$-intersecting family has cardinality $N(G,k)$ if and only if it has the following form:*

$$\mathscr{F} = \mathscr{G}_0 \cup \mathscr{G}_1 \cup \cdots \cup \mathscr{G}_{k-2},$$

*where*

$$\mathscr{G}_0 = \left\{X \in \binom{[n]}{k} : X \cap \{x,y\} \neq \emptyset\right\}$$

*and, for $1 \leqslant i \leqslant k-2$,*

$$\mathscr{G}_i = \left\{X \in \binom{[n]}{k} : \begin{array}{l} X \cap \{x,y,v_1,v_2,\dots,v_{i-1}\} = \emptyset, \\ v_i \in X, \\ X \cap A_j \neq \emptyset,\ 1 \leqslant j \leqslant i-1, \\ X \cap B \neq \emptyset \end{array}\right\}.$$

Here

$\{x,y\}$ is an edge of $G$, *i.e.*, a maximum clique,
$N(\{x\}) \setminus \{x,y\} = \{v_1, v_2, \dots, v_{d-1}\}$,
$A_j = N(\{v_j\}) \setminus \{v_j, x\},\ 1 \leqslant j \leqslant d-1$,
$B = N(\{y\}) \setminus \{x,y\}$.

https://doi.org/10.1017/S0963548301004795 Published online by Cambridge University Press

**Remark.** Unfortunately, the $p$-cube $Q_p$ on $n = 2^p$ vertices does not fit the framework of Corollary 1.3: it has cycles of length 4. It is possible to determine the maximal families for small values of $k$, but the complexity of their description soon gets out of hand. For example, if $k \leqslant 3$ and

$$n \geqslant e(p+1)\left(p+1+\binom{p+1}{2}\right)(k-1)^2k(k+1),$$

then

$$N(Q_p,k) = \binom{n}{k} - \binom{n-2}{k} + \binom{n-3}{k-1} + \binom{n-4}{k-1} - 2\binom{n-p-2}{k-1}.$$

While we must rely on the structure of individual graphs to determine whether families of the form given in (1.5) are extremal, and to determine the value of $N(G,k)$ these families give, we would expect that, for most graphs (not just sparse graphs) and small $k$, the dominant term in $N(G,k)$ is $\binom{n}{k} - \binom{n-\omega}{k}$. In order to prove a version of this claim, we appeal to the language of random graphs. Let $p, 0 < p < 1$, be a fixed real number. For the random graph $G_{n,p}$ we consider $N(G_{n,p},k)$, where $k$ is fixed and $n$ goes to infinity. Since the clique number of $G_{n,p}$ is w.h.p.† $(2+o(1))\log_{1/p} n$ (see [2]), it follows from (1.3) that

$$N(G_{n,p},k) \geqslant (2+o(1))\log_{1/p} n \cdot \binom{n-(2+o(1))\log_{1/p} n}{k-1}$$

w.h.p. We show that a constant multiple of this lower bound is an upper bound on $N(G_{n,p},k)$.

**Theorem 1.4.** *Let $p, 0 < p < 1$, be a fixed real number and let $k \geqslant 2$ be a positive integer. Then*

$$\Pr\left(N(G_{n,p},k) \geqslant c_{k,p}\binom{n}{k-1}\lambda_n\right) = o(1),$$

*where $\lambda_n = \max\{\ln n, \log_{1/p} n\}$ and*

$$c_{k,p} = \frac{8}{(1-p)^{k-1}}\left[\frac{4k(k-1)}{(1-p)^{(k-1)^2}} + 1\right].$$

Note that, if $k = 1$, then $N(G_{n,p},1)$ is given by the clique number of $G_{n,p}$ for any $0 < p < 1$.

For many graphs $N(G,k)$ is nearly equal to $\binom{n}{k}$ when $k$ is large.

**Theorem 1.5.** *Let $\delta = \delta(G)$ be the minimum degree of the graph $G$. Then*

$$k(\delta+1) > cn\ln n \quad \Rightarrow \quad N(G,k) \geqslant \left(1-n^{1-c}\right)\binom{n}{k}.$$

† A sequence of events $\mathscr{E}_n$ occurs *with high probability*, w.h.p., if $\Pr(\mathscr{E}_n) = 1 - o(1)$.

https://doi.org/10.1017/S0963548301004795 Published online by Cambridge University Press

**Theorem 1.6.** *Let $\delta$ and $c$ be constants satisfying*

$$r := c - (1-c)^{\delta+1} > 0.$$

*If $G$ is a graph on vertex set $[n]$ having minimum degree at least $\delta$ and having maximum degree $\Delta(G) = \Delta < \frac{r}{s}\sqrt{\frac{n}{\log n}}$ and $k \geqslant cn$, then*

$$N(G,k) \geqslant \left(1 - n^{-s^2/2}\right)\binom{n}{k}.$$

Now, it follows from (1.1) that

$$N(K_{\delta,n-\delta}, \lfloor cn \rfloor) \sim \left[1 - (1-c)^{\delta}\right]\binom{n}{k}$$

for $\delta, c$ fixed constants and $n \to \infty$. Hence, it is *not* the case that $N(G,k)$ is nearly $\binom{n}{k}$ whenever $k$ is on the order of $n$.

We have seen that, for most graphs, $N(G,k)$ is determined by extremal families that 'cluster' around cliques when $k$ is small, and is nearly $\binom{n}{k}$ when $k$ is large. As there is a considerable gap between the values of $k$ for which these two types of behaviour have been observed, this leaves the obvious question: What happens for $k$ in between? Is there a sharp phase transition between these two types of extrema? Do other types of extrema exist? As it seems likely that the answers to these questions will depend on the graph in question, we hazard a conjecture for only one graph, the cycle.

**Conjecture 1.7.** *There exists a constant $c$ such that, for any fixed $\epsilon > 0$,*

$$k \leqslant (c-\epsilon)n \quad \Rightarrow \quad N(C_n,k) = 2\binom{n-1}{k-1} - \binom{n-2}{k-2} + \binom{n-4}{k-2},$$

$$k \geqslant (c+\epsilon)n \quad \Rightarrow \quad N(C_n,k) = (1-o(1))\binom{n}{k}.$$

Note that it follows from Theorem 1.6 that $N(C_n,k)$ is nearly $\binom{n}{k}$ for $k > 0.32n$.

The remainder of this paper is organized as follows. Theorems 1.5 and 1.6 are proved in the next section. In Section 3 we first prove Theorem 1.1 and then prove Corollaries 1.2 and 1.3. Finally, in Section 4, we prove Theorem 1.4.

## 2. Large $k$

Both proofs in this section are probabilistic, and for both proofs we restrict our attention to the probability space $\binom{[n]}{k}$ with the uniform measure. Let $X_{n,k}$ denote a random element of this probability space.

**Proof of Theorem 1.5.** Let

$$\mathscr{F} = \left\{X \in \binom{[n]}{k} : N(X) = [n]\right\}.$$

https://doi.org/10.1017/S0963548301004795 Published online by Cambridge University Press

Clearly, $\mathscr{F}$ is a $G$-intersecting family. For a fixed vertex $v \in [n]$, we have

$$\begin{aligned} \mathrm{P}(v \notin N(X_{n,k})) &= \prod_{i=0}^{\deg_G(v)} \left(1 - \frac{k}{n-i}\right) \\ &\leqslant \left(1 - \frac{k}{n}\right)^{\delta+1} \\ &< e^{-k(\delta+1)/n}. \end{aligned}$$

Therefore, by the first moment method,

$$\mathrm{P}(X_{n,k} \notin \mathscr{F}) < ne^{-k(\delta+1)/n}.$$

Hence, $|\mathscr{F}|$ is at least

$$\binom{n}{k}\left(1 - ne^{-k(\delta+1)/n}\right). \qquad \square$$

**Proof of Theorem 1.6.** Let

$$\mathscr{F} = \left\{X \in \binom{[n]}{k} : |N(X)| > n - k\right\}.$$

Clearly, $\mathscr{F}$ is a $G$-intersecting family. For $1 \leqslant i \leqslant n$, let $Y_i$ be the random variable

$$Y_i(X) = \mathbf{E}\left[|\overline{N(X_{n,k})}| : X_{n,k} \cap [i] = X \cap [i]\right]$$

(here $\overline{N(X_{n,k})} = [n] \setminus N(X_{n,k})$), and let $Y_0$ be the (constant) random variable $Y_0 = \mathbf{E}[|\overline{N(X_{n,k})}|]$. $Y_0, \ldots, Y_n$ is a martingale. Since

$$\mathbf{E}\left[|\overline{N(X_{n,k})}|\right] \leqslant \sum_{i=1}^{n} \left(1 - \frac{k}{n}\right)^{\deg_G(i)+1} \leqslant (1-c)^{\delta+1} n$$

and

$$|Y_i - Y_{i-1}| \leqslant \Delta,$$

it follows from Azuma's inequality (see [1, p. 85]) that, for $\lambda > 0$,

$$\mathrm{P}\left(Y_n > (1-c)^{\delta+1} n + \lambda\Delta\sqrt{n}\right) < e^{-\lambda^2/2}.$$

Since $\mathrm{P}(X_{n,k} \notin \mathscr{F}) \leqslant \mathrm{P}(Y_n \geqslant cn)$, the result follows. $\square$

## 3. Sparse graphs, small $k$

In this section we prove Theorem 1.1 and Corollaries 1.2 and 1.3. We first prove two general lemmas from which Theorem 1.1 follows directly, and then go on to prove the corollaries. Throughout this section we assume $G$ is a graph on vertex set $[n]$ and $k$ is a positive integer satisfying

$$n \geqslant \frac{2e(1+\Delta)\Delta'(k-1)^2k(k+1)}{\omega}, \tag{3.1}$$

https://doi.org/10.1017/S0963548301004795 Published online by Cambridge University Press

where $\Delta = \Delta(G)$ is the maximum degree of $G$, $\omega = \omega(G)$ is the clique number of $G$, and $\Delta' = \Delta'(G)$ is the maximum size of a second neighbourhood in $G$:

$$\Delta'(G) := \max_{v \in [n]} |\{i \in [n] : \mathrm{dist}_G(i, v) \leqslant 2\}|.$$

Note that the assumptions of Corollaries 1.2 and 1.3 imply (3.1). We further assume $\mathscr{F}$ is a large $G$-intersecting family; in particular, we assume $\mathscr{F}$ is a $G$-intersecting family satisfying

$$|\mathscr{F}| > \binom{n}{k} - \binom{n-\omega}{k}. \tag{3.2}$$

At this point, we must introduce some notation and make some preliminary observations. For $i \in [n]$ let

$$\mathscr{F}_i = \{X \in \mathscr{F} : i \in X\} \quad \text{and} \quad v_i = |\mathscr{F}_i|.$$

We set $v_{\max}$ to be the largest of the $v_i$s. Since, for $X \in \mathscr{F}$ fixed, every set in $\mathscr{F}$ intersects $N(X)$, we have

$$|\mathscr{F}| \leqslant |N(X)| v_{\max} \leqslant (\Delta + 1) k v_{\max},$$

and so

$$v_{\max} > \frac{1}{k(\Delta+1)} \left[ \binom{n}{k} - \binom{n-\omega}{k} \right]. \tag{3.3}$$

Finally, we say a vertex $i \in [n]$ is *saturated* if

$$v_i > \frac{1}{k(\Delta+1)} \left[ \binom{n}{k} - \binom{n-\omega}{k} \right].$$

Note that it follows from (3.3) that there exists at least one saturated vertex. We will make use of the following inequality:

$$a, b \geqslant 2, m > ab \quad \Rightarrow \quad \binom{m-2}{b-2} < \frac{2e(b-1)}{m} \binom{m-a}{b-1}. \tag{3.4}$$

**Proof of (3.4).** Let $a, b, m$ be integers such that $a, b \geqslant 2$ and $ab < m$.

$$\begin{aligned} \frac{\binom{m-2}{b-2}}{\binom{m-a}{b-1}} &< \frac{b-1}{m-a} \left( \frac{m-b+1}{m-a-b+2} \right)^{b-2} \\ &< \frac{2(b-1)}{m} \left( 1 + \frac{a-1}{m-a-b+2} \right)^{b-2} \\ &< \frac{2(b-1)}{m} e^{\frac{(a-1)(b-2)}{m-a-b+2}} \\ &< \frac{2e(b-1)}{m}. \end{aligned}$$

□

**Lemma 3.1.** *If $v$ is a saturated vertex, then*

$$v \in N(X) \quad \forall X \in \mathscr{F}.$$

https://doi.org/10.1017/S0963548301004795 Published online by Cambridge University Press

**Proof.** Let $\mathscr{F}'_v = \{X\backslash\{v\} : X \in \mathscr{F}_v\}$. We will call a collection $\mathscr{H} \subseteq \mathscr{F}'_v$ *separated* if

$$\forall X, Y \in \mathscr{H} \qquad N(X) \cap N(Y) = \emptyset.$$

Let $\mathscr{H}$ be a maximum separated collection in $\mathscr{F}'_v$. The number of vertices at a distance less than or equal to 2 from an element of $\mathscr{H}$ is at most

$$M = |\mathscr{H}|(k-1)\Delta'.$$

We have

$$\begin{aligned}
|\mathscr{F}_v| &\leqslant \binom{n-1}{k-1} - \binom{n-M-1}{k-1} && \text{(by the maximality of } \mathscr{H}) \\
&\leqslant M\binom{n-2}{k-2} \\
&\leqslant \frac{M2e(k-1)}{n}\binom{n-\omega}{k-1} && \text{(by (3.1) and (3.4))} \\
&< \frac{M2e(k-1)}{n\omega}\left[\binom{n}{k} - \binom{n-\omega}{k}\right] \\
&< \frac{|\mathscr{H}|}{k(k+1)(\Delta+1)}\left[\binom{n}{k} - \binom{n-\omega}{k}\right] && \text{(by (3.1))}.
\end{aligned}$$

Since $v$ is a saturated vertex, we have

$$|\mathscr{H}| \geqslant k+1.$$

Now consider an arbitrary $X \in \mathscr{F}$. Since $\mathscr{F}$ is $G$-intersecting, $X$ intersects $H \cup \{v\}$ for all $H \in \mathscr{H}$. Thus, by the pigeonhole principle, some vertex of $X$ must be adjacent to at least two of the sets of the form $H \cup \{v\}$. Since $\mathscr{H}$ is separated, this vertex is a neighbour of $v$. □

**Lemma 3.2.** *If $u$ and $v$ are saturated vertices, then $u \sim_G v$.*

**Proof.** Let $u$ and $v$ be saturated vertices, and assume for the sake of contradiction that $u \not\sim_G v$. It follows from Lemma 3.1 that every element of $\mathscr{F}_u$ intersects $N(\{v\})$. Therefore,

$$\begin{aligned}
|\mathscr{F}_u| &\leqslant (\Delta+1)\binom{n-2}{k-2} && \text{(by Lemma 3.1)} \\
&\leqslant \frac{(\Delta+1)2e(k-1)}{n}\binom{n-\omega}{k-1} && \text{(by (3.1) and (3.4))} \\
&< \frac{(\Delta+1)2e(k-1)}{n\omega}\left[\binom{n}{k} - \binom{n-\omega}{k}\right] \\
&< \frac{1}{(1+\Delta+\Delta^2)(k-1)k(k+1)}\left[\binom{n}{k} - \binom{n-\omega}{k}\right] && \text{(by (3.1))} \\
&< \frac{1}{k(\Delta+1)}\left[\binom{n}{k} - \binom{n-\omega}{k}\right].
\end{aligned}$$

□

It follows from Lemma 3.2 that the saturated vertices form a clique $K$ in $G$, and it

https://doi.org/10.1017/S0963548301004795 Published online by Cambridge University Press

follows from Lemma 3.1 that any set $X \in \mathscr{F}$ satisfies $K \subseteq N(X)$. Thus, we have proved Theorem 1.1 (note that we can simply take the collection $\{M_1, \dots, M_r\}$ to be those sets in $\mathscr{F}$ that do not intersect $K$).

**Proof of Corollary 1.2.** For notational convenience we identify the vertex set $[n]$ of $G = C_n^p$ with the cyclic group $\mathbf{Z}_n$ in the natural way; to be precise,

$$\alpha \sim_G \beta \Leftrightarrow \alpha - \beta \in \{1, \dots, p, n-p, \dots, n-1\},$$

where the difference is taken in $\mathbf{Z}_n$. Let $\mathscr{F}$ be a maximum $G$-intersecting family. We may assume that $\mathscr{F}$ is of the form given in (1.5).

If $|K| = p+1$ then $K = \{\alpha, \alpha+1, \dots, \alpha+p\}$ for some $\alpha \in \mathbf{Z}_n$. Furthermore, for each set $M \in \{M_1, \dots, M_r\}$,

$$\exists \alpha + p + x, \alpha - y \in M \text{ such that } x, y \in \{1, \dots, p\} \text{ and } x + y \leqslant p+1. \tag{3.5}$$

However, if $M_i$ and $M_j$ satisfy (3.5) then $M_i \cap N(M_j) \neq \emptyset$. So, we can take the family $\{M_1, \dots, M_r\}$ to be the collection of all 2-element sets that satisfy (3.5), and it suffices to show $|K| = p+1$.

Assume for the sake of contradiction that there exists $\alpha \in K$ such that $\alpha + 1 \notin K$ and $K \cap \{\alpha+2, \dots, \alpha+p\} \neq \emptyset$. Since $\mathscr{F}$ is maximum and $\alpha+1$ is not saturated, the set $\mathscr{F}' := \{X \in \mathscr{F} : \alpha + 1 \notin N(X)\}$ is nonempty. Therefore,

$$|\mathscr{F}_{\alpha+1}| \leqslant k(2p+1)\binom{n-2}{k-2}.$$

Since an arbitrary element of $\mathscr{F}'$ must contain $\alpha - p$ and some element of $\{\alpha+p+2, \dots, \alpha+2p\}$, we have

$$|\mathscr{F}'| \leqslant (p-1)\binom{n-2}{k-2}.$$

It follows from (3.4) and (1.6) that

$$|\mathscr{F}_{\alpha+1}| + |\mathscr{F}'| \leqslant [k(2p+1) + p - 1]\binom{n-2}{k-2} < \binom{n-1}{k-1}.$$

Since the family

$$(\mathscr{F} \setminus \mathscr{F}_{\alpha+1} \setminus \mathscr{F}') \cup \left\{ X \in \binom{[n]}{k-1} : \alpha + 1 \in X \right\}$$

is $G$-intersecting, this is a contradiction.

Thus, we may assume $K = \{\alpha, \alpha+1, \dots, \alpha+l\}$ for some $\alpha \in \mathbf{Z}_n$ and $l \leqslant p$. Assume for the sake of contradiction that $l < p$. Let $\mathscr{F}'' = \{X \in \mathscr{F} : \alpha - 1 \notin N(X)\}$. Since an arbitrary element of $\mathscr{F}''$ covers $\alpha$ without covering $\alpha - 1$, we have $\mathscr{F}'' \subseteq \mathscr{F}_{\alpha+p}$. Since neither $\alpha - 1$ nor $\alpha + p$ is saturated, we have

$$|\mathscr{F}_{\alpha-1}| + |\mathscr{F}_{\alpha+p}| \leqslant \frac{2}{k(2p+1)}\left[\binom{n}{k} - \binom{n-p-1}{k}\right] < \frac{2}{k}\binom{n-1}{k-1}.$$

https://doi.org/10.1017/S0963548301004795 Published online by Cambridge University Press

This yields a contradiction because the family

$$(\mathscr{F}\setminus\mathscr{F}_{\alpha-1}\setminus\mathscr{F}_{\alpha+p})\cup\left\{X\in\binom{[n]}{k}:\alpha-1\in X\right\}$$

is $G$-intersecting and is larger than $\mathscr{F}$.

To establish (1.7) we assume that $K=\{1,2,\dots,p+1\}$. Then we can take

$$\mathscr{F}=\binom{[n]}{k}\setminus\left(\bigcup_{i=0}^{p}\mathscr{G}_i\right),$$

where

$$\mathscr{G}_0=\left\{X\in\binom{[n]}{k}:X\cap(K\cup\{n-p+1,\dots,n\})=\emptyset\right\}$$

and

$$\mathscr{G}_i=\left\{X\in\binom{[n]}{k}:n-i+1\in X,X\cap\{n-i+2,\dots,2p+3-i\}=\emptyset\right\}.$$

Equation (1.7) follows immediately. □

**Proof of Corollary 1.3.** Assume $\mathscr{F}$ is a maximum $G$-intersecting family. We may assume $\mathscr{F}$ is of the form given in (1.5). Let $\mathscr{M}$ be the collection $\{M_1,\dots,M_r\}$ defined in (1.5). We first note that $|K|=2$. If $x$ is the unique saturated vertex, then

$$\begin{aligned}|\mathscr{F}|&\leqslant|\mathscr{F}_x|+\sum_{v:v\sim x}|\mathscr{F}_v|\\&<\binom{n-1}{k-1}+\frac{d}{k(d+1)}\left[\binom{n}{k}-\binom{n-2}{k}\right]\\&=\left(\frac{n-1}{2n-k-1}+\frac{d}{k(d+1)}\right)\left[\binom{n}{k}-\binom{n-2}{k}\right]\\&<\binom{n}{k}-\binom{n-2}{k},\end{aligned}$$

which is a contradiction.

Now, let $K=\{x,y\}$,

$$A=N(\{x\})\setminus\{x,y\}\text{ and }B=N(\{y\})\setminus\{x,y\}.$$

For $v\in A\cup B$, let

$$A_v=N(\{v\})\setminus\{x,y,v\}.$$

Note that, since $G$ contains no small cycle, these sets are pairwise disjoint. Furthermore, the only edges between these sets are those connecting $v\in A\cup B$ with $A_v$, and possibly edges connecting $A_v$ and $A_u$, where $u\in A$ and $v\in B$.

It remains to determine the size and structure of

$$\mathscr{F}_0:=\{X\in\mathscr{F}:X\cap K=\emptyset\}.$$

It follows from Theorem 1.1 that all elements of $\mathscr{F}_0$ intersect both $A$ and $B$ (it follows immediately that $\mathscr{F}_0$ is empty for $k=1$; we henceforth assume $k\geqslant 2$). We proceed in

https://doi.org/10.1017/S0963548301004795 Published online by Cambridge University Press

stages. In the first stage we show that there exists a vertex $v_1 \in A \cup B$ (we assume without loss of generality that $v_1 \in A$) such that

$$\mathscr{M}_1 := \{\{v_1, b\} : b \in B\}$$

is a subset of $\mathscr{M}$ *and* that all sets in

$$\mathscr{F}_1 := \{X \in \mathscr{F}_0 : \nexists Y \in \mathscr{M}_1 \text{ such that } Y \subseteq X\}$$

must intersect $A_1 := A_{v_1}$. Since $\mathscr{F}_1 \subseteq \mathscr{F}_0$, all sets in $\mathscr{F}_1$ must intersect $A_1$, $A \backslash \{v_1\}$ and $B$ (note that any set in $\mathscr{F}_0$ that contains $v_1$ must contain some set in $\mathscr{M}_1$). In stage $2 \leqslant i \leqslant k-1$ we show that there exists a vertex $v_i \in A \backslash \{v_1, \ldots, v_{i-1}\}$ such that

$$\mathscr{M}_i := \left\{ Y \in \binom{[n]}{i+1} : v_i \in Y \text{ and } Y \text{ intersects } B, A_1, \ldots, A_{i-1} \right\}$$

is a subset of $\mathscr{M}$ *and* that all sets in

$$\mathscr{F}_i := \{X \in \mathscr{F}_{i-1} : \nexists Y \in \mathscr{M}_i \text{ such that } Y \subseteq X\}$$

must intersect $A_i$. Thus, all sets in $\mathscr{F}_i$ must intersect $A \backslash \{v_1, \ldots, v_i\}$, $B$, $A_1, \ldots, A_i$. We have a complete characterization of $\mathscr{F}$ after the $(k-1)$st stage because $\mathscr{F}_{k-1}$ is empty: an element of $\mathscr{F}_{k-1}$ has $k$ elements and intersects $A \backslash \{v_1, \ldots, v_{k-1}\}$, $B$, $A_1, \ldots, A_{k-1}$.

We begin with the first stage. As noted above, every element of $\mathscr{F}_0$ must intersect both $A$ and $B$. For $(a, b) \in A \times B$, let

$$\mathscr{F}_{a,b} = \{X \in \mathscr{F}_0 : \{a, b\} \subseteq X\}.$$

Let $\mathscr{P}$ be the set of pairs $(a, b)$ for which $\mathscr{F}_{a,b}$ is nonempty. We say that the pair $(a, b) \in \mathscr{P}$ *covers* $(a', b') \in \mathscr{P}$ if

$$X \in \mathscr{F}_{a,b} \Rightarrow N(X) \cap \{a', b'\} \neq \emptyset.$$

Now, if $(a, b) \in \mathscr{P}$ does not cover $(a', b')$, then there exists $X \in \mathscr{F}_{a,b}$ such that $N(X) \cap \{a', b'\} = \emptyset$, and

$$|\mathscr{F}_{a',b'}| \leqslant |N(X)| \binom{n-5}{k-3} \leqslant (d+1)k \binom{n-5}{k-3}. \tag{3.6}$$

On the other hand, if $(a, b)$ covers $(a', b')$ and $\{a, b\} \cap \{a', b'\} = \emptyset$, then every set in $\mathscr{F}_{a,b}$ contains some element of $A_{a'} \cup A_{b'}$, and

$$|\mathscr{F}_{a,b}| \leqslant 2(d-1) \binom{n-5}{k-3}. \tag{3.7}$$

We call a pair $(a, b) \in \mathscr{P}$ *good* if every pair in $\mathscr{P}$ covers $(a, b)$ and

$$(a, b) \text{ covers } (a', b') \Rightarrow \{a, b\} \cap \{a', b'\} \neq \emptyset.$$

Note that, if $(a, b)$ is good, then we may assume that $\mathscr{F}$ contains all sets that contain $\{a, b\}$. A pair $(a, b) \in \mathscr{P}$ that is not good will be called *bad*. It follows from (3.6) and (3.7) that

$$|\mathscr{F}_{a,b}| \leqslant (d+1)k \binom{n-5}{k-3}$$

https://doi.org/10.1017/S0963548301004795 Published online by Cambridge University Press

for any bad pair $(a, b)$. Thus, the number of sets in $\mathscr{F}$ containing at least one bad pair is at most

$$d^2(d+1)k\binom{n-5}{k-3}.$$

By (1.8) and (3.4) this quantity is less than $\binom{n-2d}{k-2}$, which is a lower bound on the number of sets containing some fixed good pair (and no other good pair). Thus, $\mathscr{F}$ has a maximal number of good pairs.

How many good pairs could $\mathscr{F}$ have? Note that, if $(a, b)$ and $(a', b')$ are good pairs, then $\{a, b\} \cap \{a', b'\} \neq \emptyset$. Clearly, we optimize the number of good pairs by taking all pairs that contain some fixed vertex. Therefore, there exists a vertex $v_1$ such that the collection

$$\mathscr{M}_1 = \{\{v_1, b\} : b \in B\}$$

is a subset of $\mathscr{M}$. Set $\mathscr{F}_1 = \{X \in \mathscr{F}_0 : v_1 \notin X\}$. By the definition of a good pair and the condition $k < d$, any set in $\mathscr{F}_1$ must intersect $A_1$. We have completed the first stage.

We now proceed by induction. Suppose stage $l$ for $1 \leqslant l \leqslant k-2$ has been completed as described above. Each set in $\mathscr{F}_l$, the collection of sets in $\mathscr{F}$ that have not yet been characterized (*i.e.*, sets in $\mathscr{F}$ that neither intersect $K$ nor contain a set in $\mathscr{M}_1 \cup \cdots \cup \mathscr{M}_l$), intersects $A \backslash \{v_1, \ldots, v_l\}$, $B$, and each of $A_1, \ldots, A_l$. We proceed as above: for

$$\mu = (a, b, x_1, \ldots, x_l) \in \big(A \backslash \{v_1, \ldots, v_l\}\big) \times B \times A_1 \times \cdots \times A_l,$$

we define

$$\mathscr{F}_\mu = \{X \in \mathscr{F}_l : \{a, b, x_1, \ldots, x_l\} \subseteq X\},$$

and set

$$\mathscr{P}_l = \{\mu \in A \times B \times A_1 \times \cdots \times A_l : \mathscr{F}_\mu \neq \emptyset\}.$$

We say that $\mu \in \mathscr{P}_l$ *covers* $\mu' = (a', b', x_1', \ldots, x_l') \in \mathscr{P}_l$ if

$$X \in \mathscr{F}_\mu \Rightarrow N(X) \cap \{a', b', x_1', \ldots, x_l'\} \neq \emptyset,$$

and that $\mu = (a, b, x_1, \ldots, x_l) \in \mathscr{P}_l$ is *good* if every $\mu' \in \mathscr{P}_l$ covers $\mu$ and

$$\mu \text{ covers } (a', b', x_1', \ldots x_l') \Rightarrow \{a, b, x_1, \ldots, x_l\} \cap \{a', b', x_1', \ldots x_l'\} \neq \emptyset.$$

Following the argument above, we conclude that $\mathscr{F}_l$ must have the maximum number of good vectors. This maximum is achieved by taking the set of good vectors to be all vectors that contain some fixed vertex. As the number of good vectors is then the product of the cardinalities of the sets that do not contain this fixed vertex, we must take the fixed vertex to be in the smallest of the sets: $A \backslash \{v_1, \ldots, v_l\}$.

Now put $\mathscr{G}_0 = \mathscr{F} \setminus \mathscr{F}_0$ and $\mathscr{G}_i = \mathscr{F}_{i-1} \setminus \mathscr{F}_i$ for $1 \leqslant i \leqslant k-2$ to obtain the collection described in the statement of the theorem.

Then, to obtain (1.10) we see that

$$|\mathscr{G}_0| = \binom{n}{k} - \binom{n-2}{k}$$

https://doi.org/10.1017/S0963548301004795 Published online by Cambridge University Press

and

$$|\mathscr{G}_i| = \sum_{j=0}^{i} (-1)^j \binom{i}{j} \binom{n - j(d-1) - i - 2}{k-1}$$

for $1 \leqslant i \leqslant k-2$. □

## 4. Random graphs

In this section we prove Theorem 1.4. We begin with definitions and notation. For finite sets $A, B$, not necessarily disjoint, let $K(A,B)$ be the graph on vertex set $A \cup B$ having edge set

$$\{\{a,b\} : a \in A, b \in B\}.$$

If $A, B \subseteq V(G)$ are disjoint vertex sets in a graph $G$, then we set

$$e_G(A,B) = e(A,B) = |\{\{x,y\} \in E(G) : x \in A, y \in B\}|.$$

For an arbitrary hypergraph $\mathscr{H}$ we set $\nu(\mathscr{H})$ equal to the size of the largest matching in $\mathscr{H}$ (*i.e.*, collection of pairwise disjoint edges) and $\tau(\mathscr{H})$ equal to the size of the smallest vertex cover in $\mathscr{H}$ (*i.e.*, collection of vertices that meets all edges). Clearly,

$$\mathscr{H} \quad k\text{-uniform} \Rightarrow \tau(\mathscr{H}) \leqslant k\nu(\mathscr{H}). \tag{4.1}$$

Let $0 < p < 1$ and let a positive integer $k \geqslant 2$ be fixed. Let $G = G_{n,p}$, and let $\mathscr{F} \subseteq \binom{[n]}{k}$ be a $G$-intersecting family. For $X \subseteq [n]$, let

$$\mathscr{F}_X = \{Y \subseteq [n] \setminus X : Y \cup X \in \mathscr{F}\} \qquad \text{and} \qquad \nu_X = |\mathscr{F}_X|.$$

In other words, $\nu_X$ is the number of sets in $\mathscr{F}$ that contain $X$. If $|X| = k-1$ we will think of $\mathscr{F}_X$ as a set (rather than a collection of one-element sets). We say that $X \in \binom{[n]}{k-1}$ is *saturated* if

$$\nu_X \geqslant 8k(1-p)^{-k+1}\lambda_n$$

(note that this definition of saturated differs from that of Section 3). We form the $(k-1)$-uniform hypergraph

$$\mathscr{H} = \mathscr{H}_{\mathscr{F}} = \left\{ X \in \binom{[n]}{k-1} : X \text{ is saturated} \right\}.$$

We prove Theorem 1.4 by showing that, for an arbitrary $G$-intersecting family $\mathscr{F}$, the hypergraph $\mathscr{H}_{\mathscr{F}}$ is not too 'spread out'. One measure of the degree to which a hypergraph is 'spread out' is the vertex cover number. We first observe that if $\tau(\mathscr{H})$ is small then we have the desired bound on $\mathscr{F}$: if there exists a vertex cover $A$ of $\mathscr{H}$ such that

$$|A| < 32k(k-1)(1-p)^{k-k^2}\lambda_n,$$

https://doi.org/10.1017/S0963548301004795 Published online by Cambridge University Press

then we have

$$\begin{aligned}
|\mathscr{F}| &= \frac{1}{k}\sum_{X:|X|=k-1} v_X \\
&= \frac{1}{k}\left[\sum_{X \text{ saturated}} v_X + \sum_{X \text{ not saturated}} v_X\right] \\
&\leqslant \frac{1}{k}\left[k\sum_{v\in A} v_{\{v\}} + \sum_{X \text{ not saturated}} v_X\right] \\
&\leqslant |A|\binom{n}{k-1} + \frac{1}{k}\binom{n}{k-1}\cdot 8k(1-p)^{-k+1}\lambda_n \\
&< \left[32k(k-1)(1-p)^{k-k^2} + 8(1-p)^{-k+1}\right]\binom{n}{k-1}\lambda_n.
\end{aligned} \tag{4.2}$$

Now, let $\mathscr{F}$ be a $G$-intersecting family such that

$$|\mathscr{F}| \geqslant \left[32k(k-1)(1-p)^{k-k^2} + 8(1-p)^{-k+1}\right]\binom{n}{k-1}\lambda_n.$$

We show that such a family $\mathscr{F}$ exists only with probability $o(1)$. It follows from (4.2) that $\tau(\mathscr{H}) \geqslant 32k(k-1)(1-p)^{k-k^2}\lambda_n$, and by (4.1) there exists a matching $X_1,\ldots,X_r$ in $\mathscr{H} \subseteq \binom{[n]}{k-1}$ where

$$r = \lceil 32k(1-p)^{k-k^2}\cdot\lambda_n\rceil.$$

We show that such a matching exists only with probability $o(1)$.

Why can there not exist a large collection of disjoint saturated sets with high probability? We begin with an observation about pairs of sets in the matching. Consider disjoint $X_i \neq X_j$ for which $e(X_i, X_j) = 0$. For

$$x \in \mathscr{F}_{X_i} \qquad \text{and} \qquad y \in \mathscr{F}_{X_j},$$

it follows from the $G$-intersecting property that one of the following holds:

$$x \in N(X_j), \quad y \in N(X_i) \quad \text{or} \quad x \sim_G y.$$

In more global terms, if $A = \mathscr{F}_{X_i}\backslash N(X_j)$ and $B = \mathscr{F}_{X_j}\backslash N(X_i)$, then

$$K(A,B) \subseteq G. \tag{4.3}$$

However, if $A$ and $B$ are large then w.h.p. such subgraphs do not appear in the random graph $G_{n,p}$.

**Lemma 4.1.** *If $\mathscr{Q}$ is the event that there exists $A, B \subseteq [n]$ such that*

$$|A|, |B| > 2\log_{1/p} n \text{ and } K(A,B) \subseteq E(G),$$

*then* $\Pr(\mathscr{Q}) = o(1)$.

The proof of Lemma 4.1 is deferred to the end of this section. It follows from Lemma 4.1

https://doi.org/10.1017/S0963548301004795 Published online by Cambridge University Press

and (4.3) that we can assume that, for $i \neq j$,

$$e(X_i, X_j) = 0 \Rightarrow \min\{|\mathscr{F}_{X_i} \setminus N(X_j)|, |\mathscr{F}_{X_j} \setminus N(X_i)|\} \leqslant 2\log_{1/p} n. \tag{4.4}$$

We shall see, using (4.4), that there exists a set $X_a$ in the matching for which many of the sets $N(X_j)$ have a large intersection with $\mathscr{F}_{X_a}$. We will then finish the proof by showing that w.h.p. neighbourhood intersections of this kind do not occur in the random graph $G_{n,p}$.

In order to establish the existence of the set $X_a$ mentioned above, we form a digraph $D$ on vertex set $[r]$. We connect $i$ to $j$ with an arc if

$$e(X_i, X_j) = 0 \qquad \text{and} \qquad |\mathscr{F}_{X_i} \setminus N(X_j)| < |\mathscr{F}_{X_j} \setminus N(X_i)|.$$

It follows from (4.4) that

$$(i, j) \in D \Rightarrow |\mathscr{F}_{X_i} \setminus N(X_j)| \leqslant 2\log_{1/p} n. \tag{4.5}$$

Since $G$ is random, the following lemma shows that $D$ has w.h.p. many arcs.

**Lemma 4.2.** *If $\mathscr{R}$ is the event that there exist disjoint sets $Y_1, \ldots, Y_r \in \binom{[n]}{k-1}$ such that*

$$\left|\left\{\{i, j\} \in \binom{[r]}{2} : e(Y_i, Y_j) = 0\right\}\right| \leqslant \frac{1}{2}\binom{r}{2}(1-p)^{(k-1)^2},$$

*then* $\Pr(\mathscr{R}) = o(1)$.

The proof of Lemma 4.2 is deferred to the end of this section. It follows from Lemma 4.2 that there exists $a \in [r]$ having out-degree in $D$ at least

$$s := \left\lceil \frac{1}{r}\left(\frac{1}{2}\binom{r}{2}(1-p)^{(k-1)^2}\right)\right\rceil \geqslant 8k(1-p)^{-k+1}\lambda_n - \frac{1}{4}(1-p)^{(k-1)^2}.$$

It follows from (4.5) and the fact that $X_a$ is saturated that the set $S := \mathscr{F}_{X_a}$ and the collection of disjoint $(k-1)$-sets

$$\{Y_1, \ldots, Y_s\} := \{X_i : (a, i) \in D\}$$

satisfy

(1) $|S| \geqslant s$, and
(2) $|N(Y_i) \cap S| \geqslant s - 2\log_{1/p} n$, for $i = 1, \ldots, s$.

We now complete the proof by showing that w.h.p. such a collection of sets does not appear in $G$.

**Lemma 4.3.** *If $\mathscr{S}$ is the event that there exist $S \subseteq [n]$ and a collection of disjoint sets $Y_1, Y_2, \ldots, Y_s \in \binom{[n]}{k-1}$ satisfying* (1) *and* (2), *then* $\Pr(\mathscr{S}) = o(1)$.

The proof of Lemma 4.3 is deferred to the end of this section.

**Proof of Lemma 4.1.** Let $q = 2\log_{1/p} n$, and for $j = 1, \ldots, q$ let $\mathscr{Q}_j$ be the event that there exists $A, B \subseteq [n]$ such that

$$|A|, |B| = q, |A \cap B| = j \text{ and } K(A, B) \subseteq E(G).$$

https://doi.org/10.1017/S0963548301004795 Published online by Cambridge University Press

Consider a fixed $j$ and set $l = q - j$. An application of the first moment method yields

$$\begin{aligned}\Pr(\mathscr{Q}_j) &\leqslant \binom{n}{2l+j}\binom{2l+j}{j} \cdot p^{l^2+2lj+\binom{j}{2}} \\ &\leqslant \frac{n^{2l+j}}{(2l+j)!}\frac{(2l+j)!}{(2l)!j!} \cdot p^{q^2-j^2/2-j/2} \\ &= \left(n^q p^{q^2}\right) \cdot n^l p^{-j^2/2} \cdot p^{-j/2} \cdot \frac{1}{(2l)!j!} \\ &\leqslant p^{-j} \cdot \frac{1}{(2l)!j!} \\ &= o\left(\frac{1}{\log n}\right). \end{aligned}$$

□

**Proof of Lemma 4.2.** For disjoint sets $Y_1, \ldots, Y_r$, the random variable

$$E := \left|\left\{\{i,j\} \in \binom{[r]}{2} : e(Y_i, Y_j) = 0\right\}\right|$$

is distributed as $\mathrm{B}\left(\binom{r}{2}, (1-p)^{(k-1)^2}\right)$. The Chernoff bound then implies that

$$\Pr\left[E \leqslant \frac{1}{2}\binom{r}{2} \cdot (1-p)^{(k-1)^2}\right] < \exp\left\{-\frac{r(r-1)}{16} \cdot (1-p)^{(k-1)^2}\right\}.$$

Therefore,

$$\begin{aligned}\Pr(\mathscr{R}) &\leqslant \binom{n}{k-1}^r \exp\left\{-\frac{r(r-1)}{16} \cdot (1-p)^{(k-1)^2}\right\} \\ &\leqslant n^{r(k-1)} \exp\left\{-2rk(1-p)^{-k+1}\lambda_n + \frac{r}{16}(1-p)^{(k-1)^2}\right\} \\ &\leqslant n^{r(k-1)} \exp\left\{-2rk(1-p)^{-k+1}\lambda_n + 2k(1-p)^{-k+1}\lambda_n + 1\right\} \\ &\leqslant O\left(n^{r(k-1)-(2rk-2k)(1-p)^{-k+1}}\right) \\ &= o(1). \end{aligned}$$

□

**Proof of Lemma 4.3.** Let $\hat{s} = s - k + 1$. Consider $A, B \subseteq [n]$ such that $|A| = s$ and $|B| = k - 1$. The random variable

$$M_{A,B} = |\{x \in A \backslash B : e(\{x\}, B) = 0\}|$$

is stochastically dominated by $\mathrm{B}(\hat{s}, (1-p)^{k-1})$. It follows from the Chernoff bound that

$$\begin{aligned}\Pr(M_{A,B} \leqslant 2\log_{1/p} n) &\leqslant \Pr\left[\mathrm{B}(\hat{s}, (1-p)^{k-1}) \leqslant \frac{1}{2} \cdot \hat{s}(1-p)^{k-1}\right] \\ &\leqslant \exp\left\{-\frac{1}{8} \cdot \hat{s}(1-p)^{k-1}\right\}. \end{aligned}$$

https://doi.org/10.1017/S0963548301004795 Published online by Cambridge University Press

Therefore, the probability that fixed $S, Y_1, \dots, Y_s$ satisfy (1) and (2) is at most

$$\exp\left\{-\frac{1}{8}\cdot \hat{s}s(1-p)^{k-1}\right\}.$$

Thus,

$$\begin{aligned}
\Pr(\mathscr{S}) &\leqslant \binom{n}{s}\binom{n}{k-1}^s \exp\left\{-\frac{1}{8}\cdot \hat{s}s(1-p)^{k-1}\right\} \\
&< n^{s+(k-1)s}\exp\left\{-\frac{1}{8}\cdot \hat{s}s(1-p)^{k-1}\right\}\left(\frac{1}{s!}\right) \\
&\leqslant \exp\left\{sk\lambda_n - \frac{1}{8}\cdot \hat{s}s(1-p)^{k-1}\right\}\left(\frac{1}{s!}\right) \\
&\leqslant \exp\left\{\frac{s}{8}\left(\frac{(1-p)^{k^2-k}}{4} + (k-1)(1-p)^{k-1}\right)\right\}\left(\frac{1}{s!}\right) \\
&= o(1).
\end{aligned}$$

□

## References


[1] Alon, N. and Spencer, J. H. (1991) *The Probabilistic Method*, Wiley.

[2] Bollobás, B. (1985) *Random Graphs*, Academic Press.

[3] Bollobás, B. (1986) *Combinatorics: Set Systems, Hypergraphs, Families of Vectors and Combinatorial Probability*, Cambridge University Press.

[4] Deza, M. and Frankl, P. (1983) Erdős–Ko–Rado theorem: 22 years later. *SIAM J. Alg. Disc. Meth.* **4** 419–431.

[5] Erdős, P., Ko, C. and Rado, R. (1961) Intersection theorems for systems of finite sets. *Quart. J. Math. Oxford Ser. 2* **12** 313–320.

[6] Hajnal, A. and Rothschild, B. (1989) A generalization of the Erdős–Ko–Rado theorem on finite set systems. *J. Combin. Theory Ser. A* **51** 24–42.

[7] Hilton, A. J. W. and Milner, E. C. (1967) Some intersection theorems for systems of finite sets. *Quart. J. Math. Oxford Ser. 2* **18** 369–384.


https://doi.org/10.1017/S0963548301004795 Published online by Cambridge University Press