Proceedings of the Royal Society of Edinburgh: Section A Mathematics, **156**, 1645–1684, *2026*
DOI:10.1017/prm.2024.95

Minimal isometric dilations and operator models for the polydisc

Sourav Pal
Mathematics Department, Indian Institute of Technology Bombay, Powai, Mumbai, Maharashtra, 400076, India (souravpal@iitb.ac.in, souravmaths@gmail.com) (corresponding author)

Prajakta Sahasrabuddhe
Mathematics Department, Indian Institute of Technology Bombay, Powai, Mumbai, Maharashtra, 400076, India (prajakta@math.iitb.ac.in, praju1093@gmail.com)

(Received 5 February 2024; revised 27 June 2024; accepted 6 July 2024)

For commuting contractions $T_1, \ldots, T_n$ acting on a Hilbert space $\mathscr{H}$ with $T = \prod_{i=1}^n T_i$, we find a necessary and sufficient condition such that $(T_1, \ldots, T_n)$ dilates to a commuting tuple of isometries $(V_1, \ldots, V_n)$ on the minimal isometric dilation space of T with $V = \prod_{i=1}^n V_i$ being the minimal isometric dilation of T. This isometric dilation provides a commutant lifting of $(T_1, \ldots, T_n)$ on the minimal isometric dilation space of T. We construct both Schäffer and Sz. Nagy–Foias-type isometric dilations for $(T_1, \ldots, T_n)$ on the minimal dilation spaces of T. Also, a different dilation is constructed when the product T is a $C_{\cdot 0}$ contraction, that is, $T^{*n} \to 0$ as $n \to \infty$. As a consequence of these dilation theorems, we obtain different functional models for $(T_1, \ldots, T_n)$ in terms of multiplication operators on vectorial Hardy spaces. One notable fact about our models is that the multipliers are all analytic functions in one variable. The dilation when T is a $C_{\cdot 0}$ contraction leads to a conditional factorization of T. Several examples have been constructed.

Keywords: polydisc; commuting contractions; isometric dilation; minimality of dilation; functional model

2010 Mathematics Subject Classification: 47A20; 47A25; 47A45; 47B35; 47B38

1. Introduction

We consider only bounded operators acting on complex Hilbert spaces. A contraction is an operator with norm not greater than 1.

The aim of dilation, roughly speaking, is to realize a given tuple of commuting operators as a compression of an appropriate commuting tuple of normal operators. Let $(T_1, \ldots, T_n)$ be a tuple of commuting contractions acting on a Hilbert space $\mathscr{H}$. One would like to represent $(T_1, \ldots, T_n)$ as a compression of an n-tuple of

*Dedicated to Prof. B. V. Rajarama Bhat with deepest respect.

© The Author(s), 2024. Published by Cambridge University Press on behalf of The Royal Society of Edinburgh.

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

commuting unitaries or more precisely as a compression of an n-tuple of commuting isometries, because, every such tuple of commuting isometries extends naturally to a commuting tuple of unitaries. A commuting tuple of isometries $(V_1, \ldots, V_n)$ acting on a Hilbert space $\mathscr{K}$ is said to be an *isometric dilation* of $(T_1, \ldots, T_n)$ if $\mathscr{H}$ can be identified as a closed linear subspace of $\mathscr{K}$, i.e., $\mathscr{H} \subseteq \mathscr{K}$ and for any non-negative integers, $k_1, \ldots, k_n$

$$T_1^{k_1} \ldots T_n^{k_n} = P_{\mathscr{H}}(V_1^{k_1} \ldots V_n^{k_n})|_{\mathscr{H}},$$

where $P_{\mathscr{H}} : \mathscr{K} \to \mathscr{H}$ is the orthogonal projection. Moreover, such an isometric dilation is called minimal if

$$\mathscr{K} = \overline{Span}\, \{V_1^{k_1} \ldots V_n^{k_n} h \ : \ h \in \mathscr{H},\ k_1, \ldots, k_n \in \mathbb{N} \cup \{0\}\}.$$

If $(V_1, \ldots, V_n)$ dilates $(T_1, \ldots, T_n)$, then each T_i is a compression of V_i, that is, $T_i = P_{\mathscr{H}} V_i|_{\mathscr{H}}$. It is well-known that a contraction admits an isometric dilation (Sz. Nagy, [57]) and that a pair of commuting contractions always dilates to a pair of commuting isometries (Ando, [4]), though a triple of commuting contractions may or may not dilate to a triple of commuting isometries (Parrott, [48]). In other words, rational dilation succeeds on the closed unit disk $\overline{\mathbb{D}}$ and on the closed bidisc $\overline{\mathbb{D}}^2$ and fails on the closed polydisc $\overline{\mathbb{D}}^n$ when $n \geq 3$. Since a commuting tuple of contractions $(T_1, \ldots, T_n)$ does not dilate unconditionally whenever $n \geq 3$, efforts have been made to find classes of contractions that dilate under certain conditions and some remarkable works have been witnessed, e.g., Agler [1], Arveson [8], Ball, Li, Timotin, and Trent [13], Ball, Trent, and Vinnikov [14], Bhat, Bhattacharyya, and Dey [19], Binding, Farenick, and Li [22], Brehmer [23], Crabb and Davie [26], Curto and Vasilescu [27, 28], Dey [31], Grinshpan, Kaliuzhnyi-Verbovetki, Vinnikov, and Woerdeman [36], Muller and Vasilescu [46], Popescu [53], and many others. See the references therein and also see §2 for further details.

In this article, we consider the minimal isometric dilation space $\mathscr{K}$ (which is always unique up to unitary) of the product $T = \prod_{i=1}^n T_i$ of a tuple commuting contractions $(T_1, \ldots, T_n)$ acting on $\mathscr{H}$. We find a necessary and sufficient condition such that $(T_1, \ldots, T_n)$ dilates to a commuting isometric tuple $(V_1, \ldots, V_n)$ on $\mathscr{K}$ with the product $V = \prod_{i=1}^n V_i$ being the minimal isometric dilation of $T = \prod_{i=1}^n T_i$. Note that the space $\mathscr{K}$ is unique in the sense that any two minimal isometric dilation spaces of the product T are unitarily equivalent. This is one of the main results in this article and is stated below.

THEOREM 1.1 *Let $T_1, \ldots, T_n \in \mathscr{B}(\mathscr{H})$ be commuting contractions and let $T = \prod_{i=1}^n T_i$.*

(a) If $\mathscr{K}$ is the minimal isometric dilation space of T, then $(T_1, \ldots, T_n)$ possesses an isometric dilation $(V_1, \ldots, V_n)$ on $\mathscr{K}$ with $V = \Pi_{i=1}^n V_i$ being the minimal isometric dilation of T if and only if there are unique orthogonal projections $P_1, \ldots, P_n$ and unique commuting unitaries $U_1, \ldots, U_n$ in $\mathscr{B}(\mathscr{D}_T)$ with $\prod_{i=1}^n U_i = I_{\mathscr{D}_T}$ such that the following conditions are satisfied for each $i = 1, \ldots, n$:
(1) $D_T T_i = P_i^{\perp} U_i^* D_T + P_i U_i^* D_T T$,
(2) $P_i^{\perp} U_i^* P_j^{\perp} U_j^* = P_j^{\perp} U_j^* P_i^{\perp} U_i^*$,

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

(3) $U_iP_iU_jP_j = U_jP_jU_iP_i$,
(4) $D_TU_iP_iU_i^*D_T = D_{T_i}^2$,
(5) $P_1 + U_1^*P_2U_1 + U_1^*U_2^*P_3U_2U_1 + \ldots + U_1^*U_2^*\ldots U_{n-1}^*P_nU_{n-1}\ldots U_2U_1 = I_{\mathscr{D}_T}$.

(b) Such an isometric dilation is minimal and unique in the sense that if $(W_1, \ldots, W_n)$ *on* $\mathscr{K}_1$ *and* $(Y_1, \ldots, Y_n)$ *on* $\mathscr{K}_2$ *are two isometric dilations of* $(T_1, \ldots, T_n)$ *such that* $W = \prod_{i=1}^n W_i$ *and* $Y = \prod_{i=1}^n Y_i$ *are minimal isometric dilations of* T *on* $\mathscr{K}_1$ *and* $\mathscr{K}_2$*, respectively, then there is a unitary* $\widetilde{U} : \mathscr{K}_1 \to \mathscr{K}_2$ *such that* $(W_1, \ldots, W_n) = (\widetilde{U}^*Y_1\widetilde{U}, \ldots, \widetilde{U}^*Y_n\widetilde{U})$.

This is Theorem 3.5 in this article and will be proved in §3. We show an explicit construction of a Schäffer-type minimal isometric dilation for $(T_1, \ldots, T_n)$ on the space $\mathscr{K} = \mathscr{H} \oplus l^2(\mathscr{D}_T)$, where $\mathscr{D}_T = \overline{Ran}(I - T^*T)^{\frac{1}{2}}$ (see Theorem 3.5). We also show in Theorem 3.9 that such a dilation can be constructed with the conditions (1)–(4) of Theorem 1.1 only, though we do not have an exact converse part then. A special emphasis is given to the case when the product T is a $C_{\cdot 0}$ contraction, i.e., $T^{*n} \to 0$ strongly as $n \to \infty$. We show in Theorem 4.1 that an analogue of Theorem 1.1 can be achieved for the $C_{\cdot 0}$ case with a weaker hypothesis. We explicitly construct an isometric dilation in this case too. This leads to a functional model and a factorization of a $C_{\cdot 0}$ contraction. A notable fact about this model is that the multipliers involved here are linear analytic functions in one variable. In Theorem 6.1, another main result of this article, we construct explicitly a similar isometric dilation for $(T_1, \ldots, T_n)$ on the Sz. Nagy–Foias minimal isometric dilation space of T. In §5, we provide several examples describing different classes of commuting contractions that dilate to commuting isometries conditionally. There we show that our classes of commuting contractions admitting isometric dilations are not properly contained in any of the previously determined classes in the literature. Also, none of such classes from the literature is a proper subclass of our classes; however, there are intersections. Finally, we present a model theory for a class of commuting contractions in §7.

2. A brief history of dilation on the polydisc

An isometric dilation $(V_1, \ldots, V_n)$ of $(T_1, \ldots, T_n)$ naturally extends to a tuple of commuting unitaries $(U_1, \ldots, U_n)$, and consequently, $(U_1, \ldots, U_n)$ becomes a unitary dilation of $(T_1, \ldots, T_n)$. Since $(U_1, \ldots, U_n)$ is a tuple of commuting unitaries having its Taylor joint spectrum on the n-torus $\mathbb{T}^n$, which is the distinguished boundary of the closed polydisc $\overline{\mathbb{D}}^n$, following Arveson's terminology (see [7]), we say that $\overline{\mathbb{D}}^n$ is a complete spectral set for $(T_1, \ldots, T_n)$. So, it follows that $\overline{\mathbb{D}}^n$ is a spectral set for $(T_1, \ldots, T_n)$. Thus, the n-tuples of commuting contractions that dilate to commuting isometries or unitaries must have $\overline{\mathbb{D}}^n$ as a spectral set. In [37], Halmos constructed a unitary U on a certain larger space for a contraction T acting on a Hilbert space $\mathscr{H}$ such that $T = P_{\mathscr{H}}U|_{\mathscr{H}}$, which is to say that T is a compression of a unitary U. Existence of an isometry satisfying such a compression relation was proved before it by Julia, e.g., see [41–43]. The unitary dilation of Halmos was missing the compression-vs-dilation frame for positive integral powers of U and T. Later, Sz. Nagy resolved this issue in [58] with an innovative idea, where he proved

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

that there is a Hilbert space $\mathscr{K}$ containing $\mathscr{H}$ and a unitary U on $\mathscr{K}$ such that $T^n = P_{\mathscr{H}}U^n|_{\mathscr{H}}$ for any non-negative integer n. This is well-known as Sz. Nagy unitary dilation of a contraction. A few years after Sz. Nagy's famous discovery, Douglas [32] and Schäffer [55] produced distinct and explicit constructions of such a unitary dilation (of a contraction). The pioneering works of Sz. Nagy, Douglas, and Schäffer were further generalized by Ando to a pair of commuting contractions. Indeed, in [4], Ando constructed an isometric dilation (V_1, V_2) for a pair of commuting contractions (T_1, T_2). Success of dilation for a pair of commuting contractions led to the natural question, whether an arbitrary n-tuple of commuting contractions dilates to some n-tuple of commuting isometries or unitaries for $n \geq 3$. This question was answered negatively by Parrott in [48] via a counter example. One way of realizing the impact of this dilation result is the celebrated von Neumann inequality.

THEOREM 2.1 [62] *Let T be a contraction on some Hilbert space $\mathscr{H}$. Then, for every polynomial $p \in \mathbb{C}[z]$,*

$$\|p(T\| \leq \sup_{|z|\leq 1} |p(z)|.$$

It was observed that the existence of a unitary dilation is sufficient for a commuting tuple of contractions to satisfy von Neumann inequality. Using this principle, Crabb and Davie in [26] and Varopoulos in [60] produced examples of a triple of commuting contractions, which do not satisfy von Neumann inequality and hence do not admit a unitary dilation. These examples spurred a lot of mathematicians to look into the von Neumann inequality for commuting contractions at least on the finite dimensional spaces, [25, 33, 34, 40, 44]. In their seminal article [3], Agler and McCarthy proved a sharper version of von Neumann inequality for a pair of commuting and strictly contractive matrices. In [29], Das and Sarkar presented a new proof to the result of Agler and McCarthy with a refinement of the class of matrices. The impact of Ando's dilation is eminent even in the 20th century. In [12], Bagchi, Bhattacharyya, and Misra have presented an elementary proof of Ando's theorem in a C^*-algebraic setting, within a restricted class of homomorphisms modelled after Parrott's example. In [54], Sau gave new proofs to Ando's dilation theorem with Schäffer- and Douglas-type constructions.

In [23], Brehmer introduced the concept of regular unitary dilation and systematically studied the existence of such dilation. For any $\alpha \in \mathbb{Z}^n$, let $\alpha_- = (-\min\{o, \alpha_1\}, \ldots, -\min\{0, \alpha_n\})$ and $\alpha_+ = (\max\{0, \alpha_1\}, \ldots, \max\{0, \alpha_n\})$. For a given commuting n-tuple of contractions $(T_1, \ldots, T_n)$ and a tuple of positive integers $m = (m_1, \ldots, m_n)$, the following notation is used in the literature: $T^m := \Pi_{i=1}^n T_i^{m_i}$.

DEFINITION 2.2. § 9, [17] *A commuting n-tuple of unitaries $U = (U_1, \ldots, U_n)$ on a Hilbert space $\mathscr{K}$ is said to be a regular unitary dilation of a commuting n-tuple of contractions $T = (T_1, \ldots, T_n)$ on $\mathscr{H} \subseteq \mathscr{K}$, if, for any $\alpha \in \mathbb{Z}^n$,*

$$T^{*\alpha_-}T^{\alpha_+} = P_{\mathscr{H}}U^{*\alpha_-}U^{\alpha_+}|_{\mathscr{H}}.$$

Brehmer proved in [23] that a tuple of commuting contractions, if admits a regular unitary (or isometric) dilation, can be completely characterized bysome

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

positivity conditions, which is known as Brehmer's positivity. A tuple T is said to satisfy Brehmer's positivity condition if

$$\sum_{F\subseteq G}(-1)^{|F|}T_F^*T_F \geq 0 \tag{2.1}$$

for all $G \subseteq \{1,\dots n\}$. It follows from the definition that the existence of a regular unitary dilation implies the existence of a unitary dilation for a commuting tuple of contractions. The study of Brehmer was further continued by Halperin in [38] and [39]. The positivity condition introduced by Brehmer attracted considerable attention, e.g., see the novel works due to Agler [1] and Curto and Vasilescu [27],[28]. Indeed, Curto and Vasilescu generalized the original theorem of Brehmer together with Agler's results on hypercontractivity by general model theory for multi-operators which satisfy certain positivity conditions. An alternative approach to the results due to Agler, Curto, and Vasilescu was provided by Timotin in [59]. Timotin's approach had thrown some new light on the geometric and combinatorial parts of the model theory of Agler, Curto, and Vasilescu. In [22], Binding, Farenick, and Li proved that for every m-tuple of operators on a Hilbert space, one can simultaneously dilate them to normal operators on the same Hilbert space such that the dilating operators have finite spectrums. On the other hand, there are non-trivial results on dilation of contractive but not necessarily commuting tuples. In [30], Davis started studying such tuples, and then, Bunce [24] and Frazho [35] provided a wider and concrete form to this analysis. An extensive research in the direction of non-commuting dilation has been carried out by Popescu in [49–53] and also in collaboration with Arias in [5, 6]. In [19], Bhat, Bhattacharyya, and Dey proved that for a commuting contractive tuple, the standard commuting dilation is the maximal commuting dilation sitting inside the standard non-commuting dilation.

In [8], Arveson considered a $d-$tuple $(T_1,\dots,T_d)$ of mutually commuting operators acting on a Hilbert space $\mathscr{H}$ such that

$$\|T_1h_1+\ldots+T_dh_d\|^2 \leq \|h_1\|^2+\ldots+\|h_d\|^2.$$

He showed many of the operator-theoretic aspects of function theory of the unit disk generalize to that of the unit ball B_d in complex $d-$space, including von Neumann inequality and the model theory of contractions. Apart from this, the notable works due to Athavale [9–11], Druy [33], and Vasilescu [61] were among the early contributors to the multi-parameter operator theory on the unit ball in $\mathbb{C}^n$. In [46], Muller and Vasilescu analysed some positivity conditions for commuting multi-operators, which ensured the unitary equivalence of these objects to some standard models consisting of backwards multi-shifts. They considered spherical dilation of a tuple of commuting contractions $(T_1,\dots T_d)$ on $\mathscr{H}$. Such a tuple dilates to a tuple of commuting normal operators $(N_1,\dots,N_d)$ on $\mathscr{K}\supseteq\mathscr{H}$ satisfying

$$N_1^*N_1+\cdots+N_d^*N_d = I_{\mathscr{K}}.$$

In [46], Muller and Vasilescu gave a necessary and sufficient condition for a commuting multi-operator to have spherical dilation in terms of positivity of certain operator polynomials involving T and T^*. The dilation results of Sz. Nagy [57] for

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

contractions and Agler [2] for $m-$hypercontractions follow as a special case of the result due to Muller and Vasilescu. It is evident that, unlike unitary dilation, the tuple of contractions that admit regular dilations can be completely characterized by Brehmer's positivity conditions [23]. So, this means that Curto and Vasilescu in [28] have found a bigger class of contractive tuples, which admit commuting unitary dilations. Later Grinshpan, Kaliuzhnyi, Verbovetskyi, Vinnikov, and Woerdeman [36] extended this result to a bigger class, which was denoted by $\mathscr{P}^d_{p,q}$. Recently, Barik, Das, Haria, and Sarkar [15] introduced even a larger class of commuting contractions, denoted by $\mathscr{T}^n_{p,q}(\mathscr{H})$, which dilate to commuting isometries. Also, Barik and Das established a von Neumann inequality for a tuple of commuting contractions belonging to $\mathscr{B}^n_{p,q}$. In the expository essay [45], Levy and Shalit discussed a finite dimensional approach to dilation theory and have answered to some extent how much of the dilation theory can work out within the realm of linear algebra. Also, an interested reader is referred to [47] due to McCarthy and Shalit. In [56], Stochel and Szafraniec proposed a test for a commutative family of operators to have a unitary power dilation. For a detailed study of dilation theory, an interested reader is also referred to the nice survey articles by Bhattacharyya [20] and Shalit [55].

3. Schäffer-type minimal isometric dilation

Let us recall a few notations and terminologies from the literature. For a contraction T on a Hilbert space $\mathscr{H}$, the *defect operator* of T is the unique positive square root of $I - T^*T$, and it is denoted by D_T. Also, the closure of the range of D_T is denoted by $\mathscr{D}_T$, i.e., $\mathscr{D}_T = \overline{Ran}\, D_T$. A contraction $T \in \mathscr{B}(\mathscr{H})$ is called *completely non-unitary* or simply *c.n.u.* if there is no non-zero subspace $\mathscr{H}_1$ of $\mathscr{H}$ that reduces T and on which T acts as a unitary. The classical L^2 space consists of complex-valued functions defined on the unit circle $\mathbb{T}$ that are square integrable with respect to the Lebesgue measure on $\mathbb{T}$. A canonical basis for L^2 is $\{e^{in\theta} : n \in \mathbb{Z}\}$, and the closed subspace of L^2 generated by the basis $\{e^{in\theta} : n = 0, 1, 2, \dots\}$ is denoted by $\widetilde{H}^2$. For any Hilbert space E, the space $L^2(E)$ is defined similarly as L^2, and the only difference is that the functions in $L^2(E)$ are E-valued. It is well-known that the Hilbert spaces $L^2(E)$ and $L^2 \otimes E$ are unitarily equivalent. Under this unitary equivalence, the replica of $\widetilde{H}^2 \otimes E$ in $L^2(E)$ is denoted by $\widetilde{H}^2(E)$. A *multiplication operator* M_ϕ on $L^2(E)$, where $\phi(z)$ is an essentially bounded function from $\mathbb{T}$ to E, i.e. $\phi \in L^\infty(E)$, is defined by $M_\phi f(z) = \phi(z)f(z)$. For any $\phi \in L^\infty(E)$, the *Toeplitz* operator T_ϕ on $\widetilde{H}^2(E)$ is defined by $T_\phi f(z) = P\phi(z)f(z)$, where $P : L^2(E) \to \widetilde{H}^2(E)$ is the orthogonal projection. For any Hilbert space E, the *Hardy space* $H^2(E)$ consists of analytic functions from the unit disk $\mathbb{D}$ to E with square summable coefficients in its power series, i.e.,

$$H^2(E) = \Bigg\{ f : \mathbb{D} \to E \,:\, f(z) = \sum_{i=0}^{\infty} a_n z^n \,,\ a_n \in E \text{ for all } n \in \mathbb{N} \cup \{0\} \times \sum_{i=0}^{\infty} \|a_n\|^2 < \infty \Bigg\}.$$

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

The Hilbert spaces $\widetilde{H}^2(E)$ and $H^2(E)$ are unitarily equivalent. A multiplication operator M_ϕ on $H^2(E)$, where $\phi(z)$ is an analytic multiplier, is defined by $M_\phi f(z) = \phi(z)f(z)$.

To explain the results of this section, we begin with the Berger–Coburn–Lebow model (or, simply the BCL model) for commuting isometries, which will be used in sequel.

THEOREM 3.1 Berger–Coburn–Lebow, [18] *Let $V_1, \ldots, V_n$ be commuting isometries on $\mathscr{H}$ such that $V = \prod_{i=1}^n V_i$ is a pure isometry. Then, there exist projections $P_1, \ldots, P_n$ and unitaries $U_1, \ldots, U_n$ in $\mathscr{B}(\mathscr{D}_{V^*})$ such that*

$$(V_1, \ldots, V_n, V) \equiv (T_{P_1^\perp U_1 + zP_1U_1}, \ldots, T_{P_n^\perp U_n + zP_nU_n}, T_z) \text{ on } H^2(\mathscr{D}_{V^*}).$$

Later, Bercovici, Douglas, and Foias found a refined operator model for commuting c.n.u. isometries in [16]. They introduced the notion of model n-isometries. A *model n-isometry* is a tuple of commuting n-isometries $(V_1, \ldots, V_n)$ such that each V_i is a multiplication operator of the form $M_{U_iP_i^\perp + zU_iP_i}$ and $\prod_{i=1}^n V_i = M_z$, where $P_1, \ldots, P_n$ are orthogonal projections and $U_1, \ldots, U_n$ are unitaries acting on a Hilbert space $\mathscr{H}$. The following characterization theorem for model n-isometries is nothing but a variant of the model due to Bercovici, Douglas, and Foias and a proof follows from lemma 2.2 in [16] and the discussion below it. This will be used in sequel.

THEOREM 3.2 Bercovici, Douglas, and Foias, [16] *Let $U_1, \ldots, U_n$ be unitaries on Hilbert space $\mathscr{H}$ and $P_1, \ldots, P_n$ be orthogonal projections on $\mathscr{H}$. For each $1 \leq i \leq n$, let $V_i = M_{U_iP_i^\perp + zU_iP_i}$. Then, $(V_1, \ldots, V_n)$ defines a commuting n-tuple of isometries with $\Pi_{i=1}^n V_i = M_z$ if and only if the following conditions are satisfied:*

(1) $U_iU_j = U_jU_i$ *for all* $1 \leq i < j \leq n$,
(2) $U_1 \ldots U_n = I_{\mathscr{H}}$,
(3) $P_j + U_j^*P_iU_j = P_i + U_i^*P_jU_i \leq I_{\mathscr{H}}$ *for all* $i \neq j$ *and*
(4) $P_1 + U_1^*P_2U_1 + U_1^*U_2^*P_3U_2U_1 + \ldots + U_1^*U_2^* \ldots U_{n-1}^*P_nU_{n-1} \ldots U_2U_1 = I_{\mathscr{H}}$.

The following result is a corollary of Theorem 3.2. In other words, this can be treated as a variant of Theorem 3.2. We state it here so that we can directly apply it later.

THEOREM 3.3 *Let $U_1, \ldots, U_n$ be unitaries on Hilbert space $\mathscr{H}$ and $P_1, \ldots, P_n$ be orthogonal projections on $\mathscr{H}$. For each $1 \leq i \leq n$, let $V_i = M_{P_i^\perp U_i^* + zP_iU_i^*}$. Then, $(V_1, \ldots, V_n)$ defines a commuting n-tuple of isometries with $\Pi_{i=1}^n V_i = M_z$ if and only if the following conditions are satisfied:*

(1) $U_iU_j = U_jU_i$ *for all* $1 \leq i < j \leq n$,
(2) $U_1 \ldots U_n = I_{\mathscr{H}}$,
(3) $P_j + U_j^*P_iU_j = P_i + U_i^*P_jU_i \leq I_{\mathscr{H}}$ *for all* $i \neq j$, *and*
(4) $P_1 + U_1^*P_2U_1 + U_1^*U_2^*P_3U_2U_1 + \ldots + U_1^*U_2^* \ldots U_{n-1}^*P_nU_{n-1} \ldots U_2U_1 = I_{\mathscr{H}}$.

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

Now we present a twisted version of lemma 2.2 in [16]. This will be used in the proof of the main theorem. For the sake of completeness, we give a proof here, and it goes along with similar arguments as in the proof of lemma 2.2 in [16].

LEMMA 3.4. *Consider unitary operators U, U_1, and U_2 and orthogonal projections P, P_1, and P_2 on Hilbert space $\mathscr{H}$. If $V_{U,P}, V_{U_1,P_1}$ and V_{U_2,P_2} on $H^2(\mathscr{H})$ are defined as $V_{U,P} = M_{P^\perp U^* + zPU^*}$, $V_{U_1,P_1} = M_{P_1^\perp U_1^* + zP_1U_1^*}$, and $V_{U_2,P_2} = M_{P_2^\perp U_2^* + zP_2U_2^*}$, then the following are equivalent:*

(i) $V_{U,P} = V_{U_1,P_1} V_{U_2,P_2}$,
(ii) $U = U_2U_1$ *and* $P = P_1 + U_1^* P_2 U_1$.

Proof. We prove only $(i) \implies (ii)$, the proof of $(ii) \implies (i)$ follows trivially. From (i), we have $V_{U,P} = V_{U_1,P_1} V_{U_2,P_2}$, and thus,

$$P^\perp U^* + zPU^* = (P_1^\perp U_1^* + zP_1U_1^*)(P_2^\perp U_2^* + zP_2U_2^*),$$
$$\text{i.e.,} \quad P^\perp U^* + zPU^* = P_1^\perp U_1^* P_2^\perp U_2^* + z(P_1U_1^* P_2^\perp U_2^* + P_1^\perp U_1^* P_2 U_2^*) + z^2 P_1U_1^* P_2 U_2^*.$$

Hence, we have

(1) $P^\perp U^* = P_1^\perp U_1^* P_2^\perp U_2^*$;
(2) $PU^* = P_1U_1^* P_2^\perp U_2^* + P_1^\perp U_1^* P_2 U_2^*$;
(3) $P_1U_1^* P_2 U_2^* = 0$.

From (2) and (3), by substituting $P_i^\perp = I - P_i$, we obtain

$$PU^* = U_1^* P_2 U_2^* + P_1 U_1^* U_2^*. \tag{3.1}$$

From (1) and (3), by substituting $P_i^\perp = I - P_i$, we obtain

$$U^* - PU^* = U_1^* U_2^* - U_1^* P_2 U_2^* - P_1 U_1^* U_2^*. \tag{3.2}$$

Hence, (3.1) and (3.2) give us $U = U_2U_1$. Again, multiplying (3.1) from right by U_2U_1 and substituting $U = U_2U_1$, we obtain $P = P_1 + U_1^* P_2 U_1$. The proof is now complete. □

We now present a Schäffer-type minimal isometric dilation for a tuple of commuting contractions, and this is one of the main results of this article. However, the proof of this theorem is going to be lengthy, and so, we will split the proof into several parts. We request the readers to kindly bear with us for once.

THEOREM 3.5 *Let $T_1, \ldots, T_n \in \mathscr{B}(\mathscr{H})$ be commuting contractions and let $T = \prod_{i=1}^n T_i$.*

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

(a) If $\mathscr{K}$ is the minimal isometric dilation space of T, then $(T_1, \ldots, T_n)$ possesses an isometric dilation $(V_1, \ldots, V_n)$ on $\mathscr{K}$ with $V = \Pi_{i=1}^n V_i$ being the minimal isometric dilation of T if and only if there are unique orthogonal projections $P_1, \ldots, P_n$ and unique commuting unitaries $U_1, \ldots, U_n$ in $\mathscr{B}(\mathscr{D}_T)$ with $\prod_{i=1}^n U_i = I_{\mathscr{D}_T}$ such that the following conditions are satisfied for each $i = 1, \ldots, n$:

(1) $D_T T_i = P_i^{\perp} U_i^* D_T + P_i U_i^* D_T T$,
(2) $P_i^{\perp} U_i^* P_j^{\perp} U_j^* = P_j^{\perp} U_j^* P_i^{\perp} U_i^*$,
(3) $U_i P_i U_j P_j = U_j P_j U_i P_i$,
(4) $D_T U_i P_i U_i^* D_T = D_{T_i}^2$,
(5) $P_1 + U_1^* P_2 U_1 + U_1^* U_2^* P_3 U_2 U_1 + \ldots + U_1^* U_2^* \ldots U_{n-1}^* P_n U_{n-1} \ldots U_2 U_1 = I_{\mathscr{D}_T}$.

(b) Such an isometric dilation is minimal and unique in the sense that if $(W_1, \ldots, W_n)$ on $\mathscr{K}_1$ and $(Y_1, \ldots, Y_n)$ on $\mathscr{K}_2$ are two isometric dilations of $(T_1, \ldots, T_n)$ such that $W = \prod_{i=1}^n W_i$ and $Y = \prod_{i=1}^n Y_i$ are minimal isometric dilations of T on $\mathscr{K}_1$ and $\mathscr{K}_2$, respectively, then there is a unitary $\widetilde{U} : \mathscr{K}_1 \to \mathscr{K}_2$ such that $(W_1, \ldots, W_n) = (\widetilde{U}^* Y_1 \widetilde{U}, \ldots, \widetilde{U}^* Y_n \widetilde{U})$.

Proof. **(a). (*The* $\Leftarrow$ *part*).** Suppose there are projections $P_1, \ldots, P_n$ and commuting unitaries $U_1, \ldots, U_n$ in $\mathscr{B}(\mathscr{D}_T)$ with $\prod_{i=1}^n U_i = I$ satisfying the operator identities (1)–(5) for $1 \leq i \leq n$. We first show that conditions (1)–(4) guarantee the existence of an isometric dilation of $(T_1, \ldots, T_n)$. In fact, we shall construct a co-isometric extension of $(T_1^*, \ldots, T_n^*)$. It is well-known from Sz. Nagy–Foias theory (see [17]) that any two minimal isometric dilations of a contraction are unitarily equivalent. Thus, without loss of generality, we consider the Schäffer's minimal isometric dilation space $\mathscr{K}_0$ of T, where $\mathscr{K}_0 = \mathscr{H} \oplus l^2(\mathscr{D}_T) = \mathscr{H} \oplus \mathscr{D}_T \oplus \mathscr{D}_T \oplus \ldots$ and construct an isometric dilation on $\mathscr{K}_0$ for $(T_1, \ldots, T_n)$. Define V_i on $\mathscr{K}_0$ as follows:

$$V_i = \begin{bmatrix} T_i & 0 & 0 & 0 & \ldots \\ P_i U_i^* D_T & P_i^{\perp} U_i^* & 0 & 0 & \ldots \\ 0 & P_i U_i^* & P_i^{\perp} U_i^* & 0 & \ldots \\ 0 & 0 & P_i U_i^* & P_i^{\perp} U_i^* & \ldots \\ \ldots & \ldots & \ldots & \ldots & \ldots \end{bmatrix}, \qquad 1 \leq i \leq n. \tag{3.3}$$

It is evident from the block matrix form that $V_i^*|_{\mathscr{H}} = T_i^*$ for each $i = 1, \ldots, n$.

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

Step 1. First we prove that $(V_1, \ldots, V_n)$ is a commuting tuple. For each i, j, we have that

$$V_iV_j = \begin{bmatrix} T_iT_j & 0 & 0 & \cdots \\ P_iU_i^*D_TT_j + P_i^\perp U_i^*P_jU_j^*D_T & P_i^\perp U_i^*P_j^\perp U_j^* & 0 & \cdots \\ P_iU_i^*P_jU_j^*D_T & P_iU_i^*P_j^\perp U_j^* + P_i^\perp U_i^*P_jU_j^* & P_i^\perp U_i^*P_j^\perp U_j^* & \cdots \\ 0 & P_iU_i^*P_jU_j^* & P_iU_i^*P_j^\perp U_j^* + P_i^\perp U_i^*P_jU_j^* & \cdots \\ \cdots & \cdots & \cdots & \ddots \end{bmatrix}$$

and

$$V_iV_j = \begin{bmatrix} T_iT_j & 0 & 0 & \cdots \\ P_iU_i^*D_TT_j + P_i^\perp U_i^*P_jU_j^*D_T & P_i^\perp U_i^*P_j^\perp U_j^* & 0 & \cdots \\ P_iU_i^*P_jU_j^*D_T & P_iU_i^*P_j^\perp U_j^* + P_i^\perp U_i^*P_jU_j^* & P_i^\perp U_i^*P_j^\perp U_j^* & \cdots \\ 0 & P_iU_i^*P_jU_j^* & P_iU_i^*P_j^\perp U_j^* + P_i^\perp U_i^*P_jU_j^* & \cdots \\ \cdots & \cdots & \cdots & \ddots \end{bmatrix}.$$

Using the commutativity of T_i, T_j and U_i, U_j and applying condition (3) of the theorem, we obtain by simplifying the condition (2), i.e., $P_i^\perp U_i^* P_j^\perp U_j^* = P_j^\perp U_j^* P_i^\perp U_i^*$, that

$$P_iU_i^*U_j^* + U_i^*P_jU_j^* = P_jU_j^*U_i^* + U_j^*P_iU_i^*. \tag{3.4}$$

Now we show that

$$P_jU_j^*P_i^\perp U_i^* + P_j^\perp U_j^*P_iU_i^* = P_iU_i^*P_j^\perp U_j^* + P_i^\perp U_i^*P_jU_j^* \tag{3.5}$$

and

$$P_jU_j^*D_TT_i + P_j^\perp U_j^*P_iU_i^*D_T = P_iU_i^*D_TT_j + P_i^\perp U_i^*P_jU_j^*D_T. \tag{3.6}$$

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

For showing (3.5), we see that

$$\begin{aligned} P_jU_j^*P_i^\perp U_i^* + P_j^\perp U_j^*P_iU_i^* &= P_jU_j^*(U_i^* - P_iU_i^*) + (U_j^* - P_jU_j^*)P_iU_i^* \\ &= P_jU_j^*U_i^* - P_jU_j^*P_iU_i^* + U_j^*P_iU_i^* - P_jU_j^*P_iU_i^* \\ &= P_iU_i^*U_j^* - P_iU_i^*P_jU_j^* + U_i^*P_jU_j^* - P_iU_i^*P_jU_j^* \\ &= P_iU_i^*P_j^\perp U_j^* + P_i^\perp U_i^*P_jU_j^*. \end{aligned}$$

Note that the second last equality follows from Eq. (3.4) and condition (3) of Theorem 3.5.

To prove (3.6), we use a few conditions of Theorem 3.5 here. We have

$$\begin{aligned} &P_jU_j^*D_TT_i + P_j^\perp U_j^*P_iU_i^*D_T \\ &\quad= P_jU_j^*(P_i^\perp U_i^*D_T + P_iU_i^*D_TT) + P_j^\perp U_j^*P_iU_i^*D_T \quad \text{[by condition (1)]} \\ &\quad= (P_jU_j^*P_i^\perp U_i^* + P_j^\perp U_j^*P_iU_i^*)D_T + P_jU_j^*P_iU_i^*D_TT \\ &\quad= (P_iU_i^*P_j^\perp U_j^* + P_i^\perp U_i^*P_jU_j^*)D_T + P_iU_i^*P_jU_j^*D_TT \\ &\qquad\qquad\qquad\qquad\qquad\qquad \text{[by (3.5) and condition(3)]} \\ &\quad= P_iU_i^*(P_j^\perp U_j^*D_T + P_jU_j^*D_TT) + P_i^\perp U_i^*P_jU_j^*D_T \\ &\quad= P_iU_i^*D_TT_j + P_i^\perp U_i^*P_jU_j^*D_T. \qquad\qquad \text{[by condition (1)]} \end{aligned}$$

Hence, it follows that $V_jV_i = V_iV_j$ for all i, j, and consequently, $(V_1, \ldots, V_n)$ is a commuting tuple.

Step 2. We now prove that each V_j is an isometry and that $(V_1, \ldots, V_n)$ is an isometric dilation of $(T_1, \ldots, T_n)$. Note that

$$V_j^*V_j = \begin{bmatrix} T_j^*T_j + D_TU_jP_jU_j^*D_T & 0 & 0 & \ldots \\ U_jP_j^\perp P_jU_j^*D_T & U_jP_j^\perp U_j^* + U_jP_jU_j^* & U_jP_jP_j^\perp U_j & \ldots \\ 0 & U_jP_j^\perp P_jU_j^* & U_jP_j^\perp U_j^* + U_jP_jU_j^* & \ldots \\ 0 & 0 & U_jP_j^\perp P_jU_j^* & \ldots \\ \ldots & \ldots & \ldots & \ddots \end{bmatrix}.$$

By condition (4), we have $T_j^*T_j + D_TU_jP_jU_j^*D_T = I$. Also, the identities $U_jP_j^\perp U_j^* + U_jP_jU_j^* = I$ and $U_jP_jP_j^\perp U_j^* = U_jP_j^\perp P_jU_j^* = 0$ follow trivially. Thus, we have that $V_j^*V_j = I$, and hence, V_j is an isometry for $1 \leq j \leq n$. It is evident from the block matrix of V_i that $V_i^*|_{\mathscr{H}} = T_i^*$ and thus $(V_1, \ldots, V_n)$ on $\mathscr{K}_0$ is an isometric dilation of $(T_1, \ldots, T_n)$.

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

Step 3. It remains to show that $\prod_{i=1}^{n} V_i = V$, where V on $\mathscr{K}_0$ is the Schäffer's minimal isometric dilation of T. Note that V has the block matrix $V = \begin{bmatrix} T & 0 \\ C & S \end{bmatrix}$ with respect to the decomposition $\mathscr{K}_0 = \mathscr{H} \oplus l^2(\mathscr{D}_T)$, where

$$C = \begin{bmatrix} D_T \\ 0 \\ 0 \\ \vdots \end{bmatrix} : \mathscr{H} \to l^2(\mathscr{D}_T) \text{ and } S = \begin{bmatrix} 0 & 0 & 0 & \cdots \\ I & 0 & 0 & \cdots \\ 0 & I & 0 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix} : l^2(\mathscr{D}_T) \to l^2(\mathscr{D}_T).$$

Similarly, for all $1 \leq i \leq n$, V_i has the following matrix form with respect to the decomposition $\mathscr{K}_0 = \mathscr{H} \oplus l^2(\mathscr{D}_T)$:

$$V_i = \begin{bmatrix} T_i & 0 \\ \widetilde{C}_i & \widetilde{S}_i \end{bmatrix},$$

where

$$\widetilde{C}_i = \begin{bmatrix} P_i U_i^* D_T \\ 0 \\ 0 \\ \vdots \end{bmatrix} : \mathscr{H} \to l^2(\mathscr{D}_T) \text{ and}$$

$$\widetilde{S}_i = \begin{bmatrix} P_i^{\perp} U_i^* & 0 & 0 & \cdots \\ P_i U_i^* & P_i^{\perp} U_i^* & 0 & \cdots \\ 0 & P_i U_i^* & P_i^{\perp} U_i^* & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix} : l^2(\mathscr{D}_T) \to l^2(\mathscr{D}_T).$$

Up to a unitary $S \equiv M_z$ and $\widetilde{S}_i \equiv M_{P_i^{\perp} U_i^* + P_i U_i^* z}$ for $1 \leq i \leq n$ on $H^2(\mathscr{D}_T)$. Further (3.4) gives $P_i + U_i^* P_j U_j = P_j + U_j^* P_i U_j$. The conditions (1)–(4) of Theorem 3.3 follow from the hypotheses of this theorem. Therefore, we have $\prod_{i=1}^{n} M_{P_i^{\perp} U_i^* + z P_i U_i^*} = M_z$ and consequently $S = \prod_{i=1}^{n} \widetilde{S}_i$. As observed by Bercovici, Douglas, and Foias in [16], the terms involved in condition (5) are all mutually orthogonal projections. This is because the sum of projections Q_1 and Q_2 is again a projection if and only if they are mutually orthogonal. Now, suppose

$$\underline{T_k} = T_1 \dots T_k,\ \underline{U_k} = U_1 \dots U_k,\ \underline{P_k} = P_1 + \underline{U_1}^* P_2 \underline{U_1} + \dots + \underline{U_{k-1}}^* P_k \underline{U_{k-1}}.$$

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

Then, clearly, each $\underline{U_k}$ is a unitary, and $\underline{P_k}$ is a projection. Let us define

$$V_{\underline{T_k},\underline{U_k},\underline{P_k}} = \begin{bmatrix} \underline{T_k} & 0 & 0 & 0 & \dots \\ \underline{P_k}\,\underline{U_k}^* D_T & \underline{P_k}^{\perp}\underline{U_k}^* & 0 & 0 & \dots \\ 0 & \underline{P_k}\,\underline{U_k}^* & \underline{P_k}^{\perp}\underline{U_k}^* & 0 & \dots \\ 0 & 0 & \underline{P_k}\,\underline{U_k}^* & \underline{P_k}^{\perp}\underline{U_k}^* & \dots \\ \dots & \dots & \dots & \dots & \dots \end{bmatrix}, \qquad 1 \leq k \leq n.$$

We prove that $V_{\underline{T_k},\underline{U_k},\underline{P_k}} V_{\underline{T_{k+1}},\underline{U_{k+1}},\underline{P_{k+1}}} = V_{\underline{T_{k+1}},\underline{U_{k+1}},\underline{P_{k+1}}}$ for all $1 \leq k \leq n-1$. Note that for all $1 \leq k \leq n$, $V_{\underline{T_k},\underline{U_k},\underline{P_k}}$ has the following block matrix form with respect to the decomposition $\mathscr{K}_0 = \mathscr{H} \oplus l^2(\mathscr{D}_T)$:

$$V_{\underline{T_k},\underline{U_k},\underline{P_k}} = \begin{bmatrix} \underline{T_k} & 0 \\ \underline{C_k} & \underline{S_k} \end{bmatrix},$$

where

$$\underline{C_k} = \begin{bmatrix} \underline{P_k}\,\underline{U_k}^* D_T \\ 0 \\ 0 \\ \vdots \end{bmatrix} : \mathscr{H} \to l^2(\mathscr{D}_T) \text{ and}$$

$$\underline{S_k} = \begin{bmatrix} \underline{P_k}^{\perp}\underline{U_k}^* & 0 & 0 & \cdots \\ \underline{P_k}\,\underline{U_k}^* & \underline{P_k}^{\perp}\underline{U_k}^* & 0 & \cdots \\ 0 & \underline{P_k}\,\underline{U_k}^* & \underline{P_k}^{\perp}\underline{U_k}^* & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix} : l^2(\mathscr{D}_T) \to l^2(\mathscr{D}_T).$$

It is clear from the definition that $\underline{U_{k+1}} = \underline{U_k} U_{k+1}$ and $\underline{P_{k+1}} = \underline{P_k} + \underline{U_k}^* P_{k+1} \underline{U_k}$. Hence, Lemma 3.4 tells us that $\underline{S_{k+1}} = \underline{S_k} S_{k+1}$. From the construction, it is clear that $\underline{T_{k+1}} = \underline{T_k} T_{k+1}$. Hence, for proving $V_{\underline{T_k},\underline{U_k},\underline{P_k}} V_{\underline{T_{k+1}},\underline{U_{k+1}},\underline{P_{k+1}}} = V_{\underline{T_{k+1}},\underline{U_{k+1}},\underline{P_{k+1}}}$, it suffices to prove $\underline{C_k} T_{k+1} + \underline{S_k} C_{k+1} = \underline{C_{k+1}}$. Here,

$$\underline{C_k} T_{k+1} + \underline{S_k} C_{k+1} = \begin{bmatrix} \underline{P_k}\,\underline{U_k}^* D_T T_{k+1} + \underline{P_k}^{\perp}\underline{U_k}^* P_{k+1} U_{k+1}^* D_T \\ \underline{P_k}\,\underline{U_k}^* P_{k+1} U_{k+1}^* D_T \\ 0 \\ \vdots \end{bmatrix}.$$

Note that $\underline{S_{k+1}} = \underline{S_k} S_{k+1}$ gives

$$\underline{P_k}\,\underline{U_k}^* P_{k+1} U_{k+1}^* = 0 \qquad \underline{P_k}^{\perp}\underline{U_k}^* P_{k+1} U_{k+1}^* + \underline{P_k}\,\underline{U_k}^* P_{k+1}^{\perp} U_{k+1}^* = \underline{P_{k+1}}\,\underline{U_{k+1}^*}.$$

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

Using these two equations and condition (1) of this theorem, we obtain the following:

$$\begin{aligned}
&\underline{P_k}\,\underline{U_k}^* D_T T_{k+1} + \underline{P_k}^{\perp}\underline{U_k}^* P_{k+1} U_{k+1}^* D_T \\
&\quad = \underline{P_k}\,\underline{U_k}^* D_T T_{k+1} + \underline{P_{k+1}}\,\underline{U_{k+1}}^* D_T - \underline{P_k}\,\underline{U_k}^* P_{k+1}^{\perp} U_{k+1}^* D_T \\
&\quad = \underline{P_k}\,\underline{U_k}^* (D_T T_{k+1} - P_{k+1}^{\perp} U_{k+1}^* D_T) + \underline{P_{k+1}}\,\underline{U_{k+1}}^* D_T \\
&\quad = \underline{P_k}\,\underline{U_k}^* P_{k+1} U_{k+1}^* D_T T + \underline{P_{k+1}}\,\underline{U_{k+1}}^* D_T \\
&\quad = \underline{P_{k+1}}\,\underline{U_{k+1}}^* D_T. \qquad (3.7)
\end{aligned}$$

This proves that $\underline{C_k} T_{k+1} + \underline{S_k} C_{k+1} = \underline{C_{k+1}}$, and hence, $V_{\underline{T_k},\underline{U_k},\underline{P_k}} V_{T_{k+1},U_{k+1},P_{k+1}} = V_{\underline{T_{k+1}},\underline{U_{k+1}},\underline{P_{k+1}}}$ for all $1 \leq k \leq n-1$. Therefore, by induction, we have that $V_{\underline{T_n},\underline{U_n},\underline{P_n}} = V_1 \dots V_n$. Note that we have $\underline{T_n} = T$, $\underline{U_n} = U_1 \dots U_n = I$. Also, it follows from condition (5) that $\underline{P_n} = I$. Therefore, $V_{\underline{T_n},\underline{U_n},\underline{P_n}} = V$, where V is the Schäffer's minimal isometric dilation. Hence, $\prod_{i=1}^n V_i = V$.

Step-4. We now show that such an isometric dilation $(V_1, \dots, V_n)$ is minimal. Note that $V = \prod_{i=1}^n V_i$ is a minimal isometric dilation of $T = \prod_{i=1}^n T_i$. Therefore,

$$\mathscr{K} = \overline{Span}\,\{V^k h \,:\, h \in \mathscr{H},\, k \in \mathbb{N} \cup \{0\}\,\} = \overline{Span}\,\{V_1^k \dots V_n^k h \,:\, h \in \mathscr{H},\, k \in \mathbb{N} \cup \{0\}\,\}.$$

Again

$$Span\,\{V_1^{k_1} \dots V_n^{k_n} h \,:\, h \in \mathscr{H},\, k_1, \dots, k_n \in \mathbb{N} \cup \{0\}\,\} \subseteq \mathscr{K}.$$

Therefore,

$$\begin{aligned}
\mathscr{K} &= \overline{Span}\,\{V_1^k \dots V_n^k h \,:\, h \in \mathscr{H},\, k \in \mathbb{N} \cup \{0\}\,\} \\
&= \overline{Span}\,\{V_1^{k_1} \dots V_n^{k_n} h \,:\, h \in \mathscr{H},\, k_1, \dots, k_n \in \mathbb{N} \cup \{0\}\,\},
\end{aligned}$$

and consequently, $(V_1, \dots, V_n)$ is a minimal isometric dilation of $(T_1, \dots, T_n)$.

(The $\Rightarrow$ part). Let $(W_1, \dots, W_n)$ on $\mathscr{K}$ be an isometric dilation of $(T_1, \dots, T_n)$ such that $W = \Pi_{i=1}^n W_i$ is the minimal isometric dilation of T. Then, $\mathscr{K} = \mathscr{H} \oplus \mathscr{H}^{\perp}$. Suppose $W_i' = \prod_{j \neq i} W_j$ for $1 \leq i \leq n$. So,

$$\mathscr{K} = \overline{span}\,\{W^n h \,:\, h \in \mathscr{H},\, n \in \mathbb{N} \cup \{0\}\}.$$

Since V on $\mathscr{K}_0 = \mathscr{H} \oplus l^2(\mathscr{D}_T)$ is the minimal Schäffer's isometric dilation of T, it follows that

$$\mathscr{K}_0 = \overline{span}\{V^n h \,:\, h \in \mathscr{H},\, n \in \mathbb{N} \cup \{0\}\}.$$

Therefore, the map $\tau : \mathscr{K}_0 \to \mathscr{K}$ defined by $\tau(V^n h) = W^n h$ is a unitary which is identity on $\mathscr{H}$. Thus, $\mathscr{H}$is a reducing subspace for τ, and consequently,

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

$\tau = \begin{pmatrix} I & 0 \\ 0 & \tau_1 \end{pmatrix}$ for some unitary τ_1. Then, $W = \tau V \tau^* = \begin{bmatrix} T & 0 \\ \tau_1 C & \tau_1 S \tau_1^* \end{bmatrix}$, where $V = \begin{bmatrix} T & 0 \\ C & S \end{bmatrix}$ with

$$C = \begin{bmatrix} D_T \\ 0 \\ 0 \\ \vdots \end{bmatrix} : \mathscr{H} \to l^2(\mathscr{D}_T) \quad \text{and} \quad S = \begin{bmatrix} 0 & 0 & 0 & \cdots \\ I & 0 & 0 & \cdots \\ 0 & I & 0 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix} : l^2(\mathscr{D}_T) \to l^2(\mathscr{D}_T).$$

Let us consider the commuting tuple of isometries $(\widehat{V}_1, \ldots, \widehat{V}_n) = (\tau^* W_1 \tau, \ldots, \tau^* W_n \tau)$. Note that $\prod_{i=1}^n \widehat{V}_i = \tau^* W \tau = V$. Define $\widehat{V}_i' = \prod_{j \neq i} \widehat{V}_j$ for $1 \leq i \leq n$. Evidently, each $\widehat{V}_i'$ is an isometry and $\widehat{V}_i, \widehat{V}_j', V$ commute for all i, j. Also, $\widehat{V}_i = \widehat{V}_i'^* V$ for $i = 1, \ldots, n$. Suppose the block matrix of $\widehat{V}_i$ with respect to the decomposition $\mathscr{H} \oplus l^2(\mathscr{D}_P)$ be $\widehat{V}_i = \begin{bmatrix} T_i & A_i \\ C_i & S_i \end{bmatrix}$. Now by the commutativity of $\widehat{V}_i$ and $\widehat{V}$, we have

$$\begin{bmatrix} T_i & A_i \\ C_i & S_i \end{bmatrix} \begin{bmatrix} T & 0 \\ C & S \end{bmatrix} = \begin{bmatrix} T & 0 \\ C & S \end{bmatrix} \begin{bmatrix} T_i & A_i \\ C_i & S_i \end{bmatrix}$$

$$i.e. \quad \begin{bmatrix} T_i T + A_i C & A_i S \\ C_i T + S_i C & S_i S \end{bmatrix} = \begin{bmatrix} T T_i & T A_i \\ C T_i + S C_i & C A_i + S S_i \end{bmatrix}. \tag{3.8}$$

Since T_i and T commute, considering the $(1,1)$ position, we have $A_i C = 0$. We now show that $A_i = 0$. Suppose $A_i = (A_{i1}, A_{i2} \ldots)$ on $\mathscr{D}_T \oplus \mathscr{D}_T \oplus \ldots$. Then, the fact that $A_i C = 0$ implies $A_{i1} = 0$ on $\mathscr{D}_T$. Again $A_i S = T A_i$ gives

$$\begin{bmatrix} 0 & A_{i2} & A_{i3} & \cdots \end{bmatrix} \begin{bmatrix} 0 & 0 & \ldots \\ I & 0 & \ldots \\ 0 & I & \ldots \\ \vdots & \vdots & \ldots \end{bmatrix} = \begin{bmatrix} 0 & T A_{i2} & T A_{i3} & \cdots \end{bmatrix},$$

which further implies that $A_{i2} = 0$ and $A_{ik} = T A_{i(k-1)}$ for all $k \geq 3$. Hence, inductively, we have $A_{ik} = 0$. This proves that $A_i = 0$. A similar argument holds if we consider the commutativity of $\widehat{V}_i'$ and V. Thus, with respect to the decomposition $\mathscr{K} = \mathscr{H} \oplus l^2(\mathscr{D}_T)$, $\widehat{V}_i$ and $\widehat{V}_i'$ have the following block matrix forms:

$$\widehat{V}_i = \begin{bmatrix} T_i & 0 \\ C_i & S_i \end{bmatrix}, \qquad \widehat{V}_i' = \begin{bmatrix} T_i' & 0 \\ C_i' & S_i' \end{bmatrix} \qquad \text{with} \quad T_i' = \prod_{i \neq j} T_j \qquad (1 \leq i \leq n), \tag{3.9}$$

for some bounded operators C_i, C_i' and S_i, S_i'. It follows from the commutativity of $\widehat{V}_i, \widehat{V}_i'$ with V that S_i, S_i' commute with S. Evidently, $S = M_z$ on $H^2(\mathscr{D}_T)$ and

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

by being commutants of S, $S_i = M_\phi$ and $S_i' = M_{\phi_i'}$ for some $\phi_i, \phi_i' \in H^\infty(\mathscr{B}(\mathscr{D}_T))$. The relation $\widehat{V}_i = \widehat{V}_i'^* V$ gives

$$\begin{bmatrix} T_i & 0 \\ C_i & S_i \end{bmatrix} = \begin{bmatrix} T_i'^* & C_i'^* \\ 0 & S_i'^* \end{bmatrix} \begin{bmatrix} T & 0 \\ C & S \end{bmatrix} = \begin{bmatrix} T_i'^* T + C_i'^* C & C_i'^* S \\ S_i'^* C & S_i'^* S \end{bmatrix}, \tag{3.10}$$

where T_i' is as in (3.9). Form here, we have the following identities for each $i = 1, \ldots, n$:

(a) $T_i - T_i'^* T = C_i'^* C$,
(b) $C_i = (M_{\phi_i'})^* C$,
(c) $M_{\phi_i} = (M_{\phi_i'})^* M_z$.

Again, $\widehat{V}_i' = \widehat{V}_i^* V$ leads to

$$\begin{bmatrix} T_i' & 0 \\ C_i' & S_i' \end{bmatrix} = \begin{bmatrix} T_i^* & C_i^* \\ 0 & S_i^* \end{bmatrix} \begin{bmatrix} T & 0 \\ C & S \end{bmatrix} = \begin{bmatrix} T_i^* T + C_i^* C & C_i^* S \\ S_i^* C & S_i^* S \end{bmatrix}, \tag{3.11}$$

and hence, we have, for each $i = 1, \ldots, n$,

(a') $T_i' - T_i^* T = C_i^* C$,
(b') $C_i' = (M_{\phi_i})^* C$,
(c') $M_{\phi_i'} = (M_{\phi_i})^* M_z$.

From (c) above and considering the power series expansions of ϕ_i and ϕ_i', we have that $\phi_i(z) = F_i + F_i'^* z$ and $\phi_i'(z) = F_i' + F_i^* z$ for some $F_i, F_i' \in \mathscr{B}(\mathscr{D}_T)$. Therefore,

$$S_i = M_{\phi_i} = \begin{bmatrix} F_i & 0 & 0 & \cdots \\ F_i'^* & F_i & 0 & \cdots \\ 0 & F_i'^* & F_i & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix} \quad \text{and} \quad S_i' = M_{\phi_i'} = \begin{bmatrix} F_i' & 0 & 0 & \cdots \\ F_i^* & F_i' & 0 & \cdots \\ 0 & F_i^* & F_i' & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix}.$$

From (b) and (b'), we have that

$$C_i = S_i'^* C = \begin{bmatrix} F_i'^* D_T \\ 0 \\ 0 \\ \vdots \end{bmatrix} \quad \text{and} \quad C_i' = S_i^* C = \begin{bmatrix} F_i^* D_T \\ 0 \\ 0 \\ \vdots \end{bmatrix}.$$

The fact that $\widehat{V}_i$ and $\widehat{V}_i'$ are isometries gives us $F_i' F_i = F_i F_i' = 0$, $F_i^* F_i + F_i' F_i'^* = F_i'^* F_i' + F_i F_i^* = I$ and $D_T F_i' F_i'^* D_T = D_{T_i}^2$ for $1 \leq i \leq n$. Again, by the commutativity of $\widehat{V}_i, \widehat{V}_j$ and considering the $(2,2)$ entries of their 2×2 block-matrices, we have the commutativity of ϕ_i and ϕ_j. From here, we have $[F_i, F_j] = 0$ and

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

$[F_i^*, F_j'] = [F_j^*, F_i']$, and they hold for all $1 \leq i, j \leq n$. Similarly, by the commutativity of $\widehat{V}_i', \widehat{V}_j'$, we have that $[F_i', F_j'] = 0$ for each i, j. Thus, combining all together, we have the following identities for $1 \leq i, j \leq n$:

$$(i)\ \ F_iF_j = F_jF_i \quad [F_i^*, F_j'] = [F_j^*, F_i'] \tag{3.12}$$

$$(ii)\ \ F_i'F_j' = F_j'F_i' \tag{3.13}$$

$$(iii)\ \ F_i'F_i = F_iF_i' = 0 \tag{3.14}$$

$$(iv)\ \ F_i^*F_i + F_i'F_i'^* = F_i'^*F_i' + F_iF_i^* = I \tag{3.15}$$

$$(v)\ \ D_TF_i'F_i'^*D_T = D_{T_i}^2. \tag{3.16}$$

For $1 \leq i \leq n$, let us define $U_i = F_i^* + F_i'$, $U_i' = F_i^* - F_i'$, and $P_i = \frac{1}{2}(I - U_i'^*U_i)$. Applying the above identities involving F_i and F_i', we have that U_i, U_i' are unitaries. Note that $F_i = (U_i^* + U_i'^*)/2$ and $F_i' = (U_i - U_i')/2$. Hence, $F_i'F_i = 0$ implies that $U_iU_i'^* = U_i'U_i^*$, and $F_iF_i' = 0$ implies that $U_i^*U_i' = U_i'^*U_i$. Thus, $P_i = \dfrac{1}{2}(I - U_i^*U_i')$. It follows from here that P_i is a projection. It can be verified that $F_i' = U_iP_i$ and $F_i = P_i^{\perp}U_i^*$. From (3.8), we have $C_iT + S_iC = CT_i + SC_i$ for each i and substituting the values of C_i, S_i and C we have

$$F_i'^*D_TT + F_iD_T = D_TT_i \tag{3.17}$$

We now show that the unitaries $U_1, \ldots, U_n$ commute. For each i, j we have

$$\begin{aligned} U_iU_j =& (F_i^* + F_i')(F_j^* + F_j') \\ =& F_i^*F_j^* + (F_i^*F_j' + F_i'F_j^*) + F_i'F_j' \\ =& F_j^*F_i^* + (F_j^*F_i' + F_j'F_i^*) + F_j'F_i' \quad \text{[by the second part of (3.12)]} \\ =& U_jU_i. \end{aligned}$$

Thus, substituting $F_i' = U_iP_i$ and $F_i = P_i^{\perp}U_i^*$, we have from (3.17), (3.12)-part-1, (3.13), and (3.16)

(1) $D_TT_i = P_i^{\perp}U_i^*D_T + P_iU_i^*D_TT$,
(2) $P_i^{\perp}U_i^*P_j^{\perp}U_j^* = P_j^{\perp}U_j^*P_i^{\perp}U_i^*$,
(3) $U_iP_iU_jP_j = U_jP_jU_iP_i$,
(4) $D_TU_iP_iU_i^*D_T = D_{T_i}^2$,

respectively, where $U_1, \ldots, U_n \in \mathscr{B}(\mathscr{D}_T)$ are commuting unitaries and $P_1, \ldots, P_n \in \mathscr{B}(\mathscr{D}_T)$ are orthogonal projections. Since $\prod_{i=1}^n \widehat{V}_i = V$, the $(2,2)$ entry of the block matrix gives $\prod_{i=1}^n S_i = S$. Note that $S_i = M_{\phi_i}$ for each i, where $\phi_i = F_i + zF_i'^* = P_i^{\perp}U_i^* + zP_iU_i^*$. So, we have $\prod_{i=1}^n M_{P_i^{\perp}U_i^* + zP_iU_i^*} = M_z$. Thus, by Lemma 3.3, $\prod_{i=1}^n U_i = I$ and (5) holds.

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

Uniqueness. From (a) and (a') we have $D^2_{T'_i}T_i = D_T F'_i D_T$ and $D^2_{T_i}T'_i = D_T F_i D_T$, respectively. If there is $G'_i \in \mathscr{B}(\mathscr{D}_T)$ such that $D^2_{T'_i}T_i = D_T G'_i D_T$, then $D_T(F'_i - G'_i)D_T = 0$, and thus, for any $h, g \in \mathscr{H}$, we have

$$\langle (F'_i - G'_i)D_T h, D_T g\rangle = \langle D_T(F'_i - G'_i)D_T h, g\rangle = 0.$$

This shows that $F'_i = G'_i$ and hence F'_i is unique. Similarly one can show that F_i is unique. Now $F_i = P_i^{\perp}U_i$ and $F'_i = U_i P_i$ and thus $U_i = F_i^* + F'_i$ and $P_i = F_i'^* F'_i$. Evidently the uniqueness of F_i, F'_i gives the uniqueness of U_i, P_i for $1 \leq i \leq n$.
(b). The minimality of the dilation follows from the fact that $V = \prod_{i=1}^n V_i$ is the minimal isometric dilation of the product $T = \prod_{i=1}^n T_i$. Following the proof of the $(\Rightarrow)$ part of **(a)** we see that any commuting isometric dilation $(W_1, \ldots, W_n)$ of $(T_1, \ldots, T_n)$ on a minimal isometric dilation space $\mathscr{K}_1$ of T is unitarily equivalent to the isometric dilation $(V_1, \ldots, V_n)$ on the Schäffer's minimal space $\mathscr{K}_0$. The rest of the argument follows and the proof is complete. □

Remark 3.6. Note that Theorem 3.5 actually provides a commutant lifting in several variables. In Theorem 3.5, the isometric dilation $(V_1, \ldots, V_n)$ of a commuting contractive tuple $(T_1, \ldots, T_n)$ is constructed in such a way that the product $V = \prod_{i=1}^n V_i$ becomes the minimal isometric dilation of the contraction $T = \prod_{i=1}^n T_i$. Also, it is evident from the block matrix form of V_i as in (3.3) that $\mathscr{H}$ is co-invariant under each V_i and $V_i^*|_{\mathscr{H}} = T_i^*$. Thus, each T_i is a commutant of T and is being lifted to V_i, which is a commutant of the minimal isometric dilation V of T.

Note 3.7. *Ando's theorem tells us that every pair of commuting contractions (T_1, T_2) dilates to a pair of commuting isometries (V_1, V_2), but (T_1, T_2) may not have such an isometric dilation (V_1, V_2) such that $V = V_1V_2$ is the minimal isometric dilation of $T = T_1T_2$. The following example shows that there are commuting contractions T_1, T_2 that violate the conditions of Theorem 3.5.*

Example 3.8. Let us consider the following contractions on $\mathbb{C}^3$:

$$T_1 = \begin{pmatrix} 0 & 0 & 0 \\ 1/3 & 0 & 0 \\ 0 & 1/3\sqrt{3} & 0 \end{pmatrix} \qquad T_2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ -1/\sqrt{3} & 0 & 0 \end{pmatrix}.$$

Evidently, $T = T_1T_2 = 0$, and thus, $D_T = I$. We show that (T_1, T_2) does not dilate to a commuting pair of isometries acting on the minimal dilation space of T. Suppose it happens. Then, there exist projections P_1, P_2 and commuting unitaries U_1, U_2 in $\mathscr{B}(\mathscr{D}_T)$ with $U_1U_2 = I$ satisfying conditions (1)–(5) of Theorem 3.5. Following the arguments in the $(\Rightarrow)$ part of the proof of Theorem 3.5, we see that T_1, T_2 satisfy (a) for $i = 1, 2$ (see the first display after (3.10)), i.e.,

$$D_T U_1 P_1 D_T = T_2, \quad D_T U_2 P_2 D_T = T_1.$$

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

Since $D_T = I$, we have that $U_1P_1 = T_2$ and $U_2P_2 = T_1$. Now we have

$$D_TU_1P_1U_1^*D_T = U_1P_1U_1^* = T_2T_2^* = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ -1/\sqrt{3} & 0 & 0 \end{pmatrix} \begin{pmatrix} 0 & 0 & -1/\sqrt{3} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1/3 \end{pmatrix}.$$

Also,

$$D_{T_1}^2 = I - T_1^*T_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} - \begin{pmatrix} 0 & 1/3 & 0 \\ 0 & 0 & 1/3\sqrt{3} \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} 0 & 0 & 0 \\ 1/3 & 0 & 0 \\ 0 & 1/3\sqrt{3} & 0 \end{pmatrix}.$$

Thus,

$$D_{T_1}^2 = \begin{pmatrix} 8/9 & 0 & 0 \\ 0 & 26/27 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

Hence, $D_TU_1P_1U_1^*D_T \neq D_{T_1}^2$, which contradicts condition (4) of Theorem 3.5. Hence (T_1, T_2) does not dilate to a pair of commuting isometries (V_1, V_2) acting on the minimal isometric dilation space of $T = T_1T_2$.

It is evident from the first part of the proof of Theorem 3.5 that such an isometric dilation $(V_1, \ldots, V_n)$ can be constructed for $(T_1, \ldots, T_n)$ with the help of conditions (1)–(4) of Theorem 3.5 and without even assuming that $\prod_{i=1}^n U_i = I$. Condition (5) was to make the product $\prod_{i=1}^n V_i$ the minimal isometric dilation of $\prod_{i=1}^n T_i$. Thus, a different version of Theorem 3.5 can be presented based on a weaker hypothesis in the following way. Needless to mention that a proof to this follows naturally from the proof of Theorem 3.5.

THEOREM 3.9 *Let $T_1, \ldots, T_n \in \mathscr{B}(\mathscr{H})$ be commuting contractions and let $T = \prod_{i=1}^n T_i$. Then, $(T_1, \ldots, T_n)$ possesses an isometric dilation on the minimal isometric dilation space of T if there are projections $P_1, \ldots, P_n$ and commuting unitaries $U_1, \ldots, U_n$ in $\mathscr{B}(\mathscr{D}_T)$ such that the following hold for $i = 1, \ldots, n$:*

(1) $D_TT_i = P_i^\perp U_i^* D_T + P_iU_i^*D_TT$,
(2) $P_i^\perp U_i^* P_j^\perp U_j^* = P_j^\perp U_j^* P_i^\perp U_i^*$,
(3) $U_iP_iU_jP_j = U_jP_jU_iP_i$,
(4) $D_TU_iP_iU_i^*D_T = D_{T_i}^2$.

Conversely, if $(T_1, \ldots, T_n)$ possesses an isometric dilation $(\widehat{V}_1, \ldots, \widehat{V}_n)$ with $V = \prod_{i=1}^n V_i$ being the minimal isometric dilation of T, then there are unique projections $P_1, \ldots, P_n$ and unique commuting unitaries $U_1, \ldots, U_n$ in $\mathscr{B}(\mathscr{D}_T)$ satisfying the conditions (1)–(4) above.

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

4. Minimal isometric dilation and functional model when the product is a $C_{\cdot 0}$ contraction

In this section, we consider a tuple of commuting contractions $(T_1, \ldots, T_n)$ with the product $T = \prod_{i=1}^{n} T_i$ being a $C_{\cdot 0}$ contraction, i.e., $T^{*n} \to 0$ strongly as $n \to \infty$. Such a tuple dilates (on the minimal isometric dilation space of T) with a weaker hypothesis than that of Theorem 3.5 as the following theorem shows. This is another main result of this article.

THEOREM 4.1 *Let $T_1, \ldots, T_n$ be commuting contractions on a Hilbert space $\mathscr{H}$ such that their product $T = \prod_{i=1}^{n} T_i$ is a $C_{\cdot 0}$ contraction. Then, $(T_1, \ldots, T_n)$ possesses an isometric dilation $(V_1, \ldots, V_n)$ with $V = \prod_{i=1}^{n} V_i$ being a minimal isometric dilation of T if and only if there are unique orthogonal projections $P_1, \ldots, P_n$ and unique commuting unitaries $U_1, \ldots, U_n$ in $\mathscr{B}(\mathscr{D}_{T^*})$ with $\prod_{i=1}^{n} U_i = I$ such that the following conditions hold for $i = 1, \ldots, n$:*

(1) $D_{T^*} T_i^* = P_i^{\perp} U_i^* D_{T^*} + P_i U_i^* D_{T^*} T^*$,
(2) $P_i^{\perp} U_i^* P_j^{\perp} U_j^* = P_j^{\perp} U_j^* P_i^{\perp} U_i^*$,
(3) $U_i P_i U_j P_j = U_j P_j U_i P_i$,
(4) $P_1 + U_1^* P_2 U_1 + U_1^* U_2^* P_3 U_2 U_1 + \ldots + U_1^* U_2^* \ldots U_{n-1}^* P_n U_{n-1} \ldots U_2 U_1 = I_{\mathscr{D}_{T^*}}$.

Proof. First, we assume that there exist orthogonal projections $P_1, \ldots, P_n$ and commuting unitaries $U_1, \ldots, U_n$ in $\mathscr{B}(\mathscr{D}_{T^*})$ such that $\prod_{i=1}^{n} U_i = I$ and the above conditions (1)–(4) hold. Since T is a $C_{\cdot 0}$ contraction, $H^2(\mathscr{D}_{T^*})$ is a minimal isometric dilation space for T. We first construct an isometric dilation $(V_1, \ldots, V_n)$ for $(T_1, \ldots, T_n)$ on the minimal isometric dilation space $H^2(\mathscr{D}_{T^*})$ of T with the assumptions $(1) - (3)$ only. Condition (4) will imply that $\prod_{i=1}^{n} V_i = V$, the minimal isometric dilation of T. Let us consider the following multiplication operators acting on $H^2(\mathscr{D}_{T^*})$:

$$V_i = M_{U_i P_i^{\perp} + z U_i P_i} \qquad (1 \leq i \leq n).$$

Since U_i is a unitary and P_i is a projection, it follows that

$$(P_i^{\perp} U_i^* + \bar{z} P_i U_i^*)(U_i P_i^{\perp} + z U_i P_i) = P_i^{\perp} + P_i = I,$$

and thus, V_i is an isometry for each i. Now we show that $(V_1, \ldots, V_n)$ is a commuting tuple. Note that for any $1 \leq i < j \leq n$, $V_i V_j = V_j V_i$ if and only if

$$(U_i P_i^{\perp} + z U_i P_i)(U_j P_j^{\perp} + z U_j P_j) = (U_j P_j^{\perp} + z U_j P_j)(U_i P_i^{\perp} + z U_i P_i),$$

which happens if and only if the given conditions (2) and (3) hold along with

$$U_i P_i U_j P_j^{\perp} + U_i P_i^{\perp} U_j P_j = U_j P_j^{\perp} U_i P_i + U_j P_j U_i P_i^{\perp}, \tag{4.1}$$

which we prove now. From the given condition (2), we have $[U_i P_i^{\perp}, U_j P_j^{\perp}] = 0$, and this implies that

$$U_i U_j - U_i U_j P_j - U_i P_i U_j + U_i P_i U_j P_j - U_j U_i + U_j U_i P_i + U_j P_j U_i - U_j P_j U_i P_i = 0.$$

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

From here we have that

$$U_iP_iU_j + U_iU_jP_j = U_jU_iP_i + U_jP_jU_i \tag{4.2}$$

We now prove that

$$(U_iP_iU_jP_j^{\perp} + U_iP_i^{\perp}U_jP_j) = (U_jP_j^{\perp}U_iP_i + U_jP_jU_iP_i^{\perp}). \tag{4.3}$$

We have that

$$\begin{aligned}
&(U_iP_iU_jP_j^{\perp} + U_iP_i^{\perp}U_jP_j) - (U_jP_j^{\perp}U_iP_i + U_jP_jU_iP_i^{\perp}) \\
&\quad = U_iP_iU_jP_j^{\perp} - U_jP_j^{\perp}U_iP_i - U_jP_jU_iP_i^{\perp} + U_iP_i^{\perp}U_jP_j \\
&\quad = U_iP_iU_j - U_iP_iU_jP_j - U_jU_iP_i + U_jP_jU_iP_i - U_jP_jU_i \\
&\qquad + U_jP_jU_iP_i + U_iU_jP_j - U_iP_iU_jP_j \\
&\quad = U_iP_iU_j - U_jU_iP_i - U_jP_jU_i + U_iU_jP_j \\
&\quad = 0. \qquad [\text{by } (4.2)]
\end{aligned}$$

Therefore, (4.1) is proved, and consequently, $(V_1, \ldots, V_n)$ is a tuple of commuting isometries. We now embed $\mathscr{H}$ inside $H^2(\mathscr{D}_{T^*})$. Let us define $W : \mathscr{H} \to H^2(\mathscr{D}_{T^*})$ by

$$W(h) = \sum_0^{\infty} z^n D_{T^*}T^{*n}h. \tag{4.4}$$

It can be found in the literature (e.g., [17]) that the map W is an isometry. However, we include a proof here for the sake of completeness and for the convenience of a reader.

$$\begin{aligned}
\|Wh\|^2 = \|\sum_{n=0}^{\infty} z^n D_{T^*}T^{*n}h\|^2 &= \left\langle \sum_{n=0}^{\infty} z^n D_{T^*}T^{*n}h, \sum_{m=0}^{\infty} z^m D_{T^*}T^{*m}h \right\rangle \\
&= \sum_{m,n=0}^{\infty} \langle z^n, z^m\rangle \langle D_{T^*}T^{*n}h, D_{T^*}T^{*m}h\rangle \\
&= \sum_{n=0}^{\infty} \langle T^n D_{T^*}^2 T^{*n}h, h\rangle \\
&= \sum_{n=0}^{\infty} \langle T^n (I - TT^*)T^{*n}h, h\rangle \\
&= \sum_{n=0}^{\infty} (\langle T^nT^{*n}h, h\rangle - \langle T^{n+1}T^{*(n+1)}h, h\rangle) \\
&= \lim_{m\to\infty} \sum_{n=0}^{m} (\|T^{*n}h\|^2 - \|T^{*n+1}h\|^2) \\
&= \|h\|^2 - \lim_{m\to\infty} \|T^{*m}h\|^2 \\
&= \|h\|^2.
\end{aligned}$$

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

The second last equality follows from the fact that $\lim_{n\to\infty} \|T^{*n}h\|^2 = 0$ as T is a $C_{\cdot 0}$ contraction. Thus, W is an isometry. We now determine the adjoint of W. For any $n \geq 0$, $\xi \in \mathscr{D}_{T^*}$, we have

$$\langle W^*(z^n\xi), h\rangle = \langle z^n\xi, \sum_{n=0}^{\infty} z^n D_{T^*}T^{*n}h\rangle = \langle \xi, D_{T^*}T^{*n}h\rangle = \langle T^n D_{T^*}\xi, h\rangle.$$

Therefore, $W^*(z^n\xi) = T^n D_{T^*}\xi$. Now for any $1 \leq i \leq n$, for all $k \in \mathbb{N}\cup\{0\}$ and for each $\xi \in \mathscr{D}_{T^*}$, we have

$$\begin{aligned} W^*V_i(z^k\xi) &= W^*M_{U_iP_i^{\perp}+zU_iP_i}(z^k\xi) \\ &= W^*(z^kU_iP_i^{\perp}\xi + z^{k+1}U_iP_i\xi) = T^kD_{T^*}U_iP_i^{\perp}\xi + T^{k+1}D_{T^*}U_iP_i\xi \\ &= T^k(D_{T^*}U_iP_i^{\perp}\xi + TD_{T^*}U_iP_i\xi) \\ &= T^k(T_iD_{T^*}\xi) \ \text{ [by condition (1)]} \\ &= T_i(T^kD_{T^*}\xi) \\ &= T_iW^*(z^k\xi). \end{aligned}$$

Therefore, $W^*V_i = T_iW^*$, i.e., $V_i^*W = WT_i^* = WT_i^*W^*W$, and hence, $V_i^*|_{W(\mathscr{H})} = WT_i^*W^*|_{W(\mathscr{H})}$. This proves that $(V_1, \ldots, V_n)$ is an isometric dilation of $(T_1, \ldots, T_n)$. Now (4.2) gives us $P_i + U_i^*P_jU_i = P_j + U_j^*P_iU_j$ for $1 \leq i < j \leq n$ and condition (4) yields $P_i + U_i^*P_jU_i = P_j + U_j^*P_iU_j \leq I_{\mathscr{D}_{T^*}}$. Hence, by an application of Lemma 3.2, we have that that $\prod_{i=1}^n V_i = M_z$, which is (up to a unitary) the minimal isometric dilation of T.

Conversely, suppose $(Y_1, \ldots, Y_n)$ acting on $\mathscr{K}$ is an isometric dilation of $(T_1, \ldots, T_n)$, where $Y = \prod_{i=1}^n Y_i$ is the minimal isometric dilation of T. Let $Y_i' = \prod_{j\neq i} Y_j$ for $1 \leq i \leq n$. Then,

$$\mathscr{K} = \overline{span}\{Y^nh : h \in \mathscr{H},\, n \in \mathbb{N}\cup\{0\}\}.$$

We first show that $Y_i^*|_{\mathscr{H}} = T_i^*$ for each $i = 1, \ldots, n$. Note that for any $i = 1, \ldots, n$, $k \in \mathbb{N}\cup\{0\}$ and $h \in \mathscr{H}$, $T_iP_{\mathscr{H}}(Y^kh) = T_iT^kh = P_{\mathscr{H}}Y_i(Y^kh)$, and as a consequence, we have $T_iP_{\mathscr{H}} = P_{\mathscr{H}}Y_i$. Now, for any $h \in \mathscr{H}$ and $k \in \mathscr{K}$,

$$\langle Y_i^*h, k\rangle = \langle h, Y_ik\rangle = \langle h, P_{\mathscr{H}}Y_ik\rangle = \langle h, T_iP_{\mathscr{H}}k\rangle = \langle T_i^*h, P_{\mathscr{H}}k\rangle = \langle T_i^*h, k\rangle.$$

Hence, $Y_i|_{\mathscr{H}} = T_i$ for each i. Therefore, the block matrix form of each Y_i with respect to the decomposition $\mathscr{K} = \mathscr{H} \oplus \mathscr{H}^{\perp}$ is $Y_i = \begin{bmatrix} T_i & 0 \\ C_i & S_i \end{bmatrix}$ for some operators C_i, S_i. Also, since $Y = \Pi_{i=1}^n Y_i$, we have $Y = \begin{bmatrix} T & 0 \\ C & S \end{bmatrix}$ for some operators C, S.

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

Again $V = \prod_{i=1}^{n} V_i = M_z$ on $H^2(\mathscr{D}_{T^*})$ is also a minimal isometric dilation of T. Therefore,

$$H^2(\mathscr{D}_{T^*}) = \overline{span}\{V^n(Wh) : h \in \mathscr{H},\, n \in \mathbb{N} \cup \{0\}\}.$$

Now the map $\tau : H^2(\mathscr{D}_{T^*}) \to \mathscr{K}$ defined by $\tau(V^n Wh) = Y^n h$ is a unitary, which maps Wh to h and τ^* maps h to Wh for all $h \in \mathscr{H}$. Therefore, with respect to the decomposition $\mathscr{K} = \mathscr{H} \oplus \mathscr{H}^\perp$ and $H^2(\mathscr{D}_{T^*}) = W(\mathscr{H}) \oplus (W\mathscr{H})^\perp$ the map τ has block matrix form $\tau = \begin{bmatrix} W^* & 0 \\ 0 & \tau_1 \end{bmatrix}$ for some unitary τ_1. Evidently, $V = \tau^* Y \tau$. Now

$$\begin{aligned} \tau^* Y_i \tau &= \begin{bmatrix} W & 0 \\ 0 & \tau_1^* \end{bmatrix} \begin{bmatrix} T_i & 0 \\ C_i & S_i \end{bmatrix} \begin{bmatrix} W^* & 0 \\ 0 & \tau_1 \end{bmatrix} \\ &= \begin{bmatrix} WT_i & 0 \\ \tau_1^* C_i & \tau_1^* S_i \end{bmatrix} \begin{bmatrix} W^* & 0 \\ 0 & \tau_1 \end{bmatrix} = \begin{bmatrix} WT_iW^* & 0 \\ \tau_1^* C_i W^* & \tau_1^* S_i \tau_1 \end{bmatrix}. \end{aligned}$$

Therefore, $\tau^* V_i^* \tau(Wh) = WT_i^* W^*(Wh) = WT_i^* h$. For each $i = 1, \ldots, n$, let us define $\widehat{V}_i := \tau^* V_i \tau$. Therefore,

$$\widehat{V}_i^* Wh = WT_i^* h \quad \text{for all} \;\; h \in \mathscr{H}, \qquad (1 \le i \le n). \tag{4.5}$$

Therefore, $(\widehat{V}_1, \ldots, \widehat{V}_n) = (\tau^* Y_1 \tau, \ldots, \tau^* Y_n \tau)$ on $H^2(\mathscr{D}_{T^*})$ is an isometric dilation of $(T_1, \ldots, T_n)$ such that $\prod_{i=1}^{n} \widehat{V}_i = V$. We now follow the arguments given in the converse part of the proof of Theorem 3.5. Since $\widehat{V}_i$ is a commutant of V $(= M_z)$, $\widehat{V}_i = M_{\phi_i}$, where $\phi_i(z) = F_i'^* + F_i z \in H^\infty(\mathscr{B}(\mathscr{D}_{T^*}))$. Evidently, $U_i = F_i^* + F_i'$ and $U_i' = F_i^* - F_i'$ are commuting unitaries and $P_i = \dfrac{1}{2}(I - U_i'^* U_i)$ is a projection for all $i = 1, \ldots, n$. A simple computation shows that $F_i = U_i P_i$ and $F_i' = P_i^\perp U_i^*$. Also, $[F_i, F_j] = [F_i', F_j'] = 0$ for all i, j. Therefore,

$$\widehat{V}_i = M_{U_i P_i^\perp + U_i P_i z} \qquad (1 \le i \le n).$$

Obviously conditions (2) and (3) follow from the commutativity of F_i, F_j and F_i', F_j', respectively. It remains to show that conditions (1) and (4) hold. From (4.5), we have $\widehat{V}_i^* Wh = WT_i^* h$ for every $h \in \mathscr{H}$. Now for any $h \in \mathscr{H}$, we have

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

$$\begin{aligned}\widehat{V}_i^* W h &= \widehat{V}_i^* \Big(\sum_{k=0}^{\infty} z^k D_{T^*} T^{*k} h\Big) \\ &= P_i^{\perp} U_i^* D_{T^*} h + \sum_{k=1}^{\infty} z^k P_i^{\perp} U_i^* D_{T^*} T^{*k} h + z^{k-1} P_i U_i^* D_{T^*} T^{*k} h \\ &= \sum_{k=0}^{\infty} z^k (P_i^{\perp} U_i^* D_{T^*} T^{*k} + P_i U_i^* D_{T^*} T^{*(k+1)}) h \\ &= \sum_{k=0}^{\infty} z^k (P_i^{\perp} U_i^* D_{T^*} + P_i U_i^* D_{T^*} T^*) T^{*k} h.\end{aligned}$$

Also,

$$W(T_i^* h) = \sum_{k=0}^{\infty} z^k D_{T^*} T^{*k} T_i^* h = \sum_{k=0}^{\infty} z^k D_{T^*} T_i^* T^{*k} h.$$

Comparing the constant terms, we have $D_{T^*} T_i^* = P_i^{\perp} U_i^* D_{T^*} + P_i U_i^* D_{T^*} T^*$. This proves condition (1). Now, since we have $\prod_{i=1}^n \widehat{V}_i = V = M_z$ and $\widehat{V}_i = M_{U_i P_i^{\perp} + z U_i P_i}$, condition (4) follows from Theorem 3.2. The uniqueness of $P_1, \ldots, P_n$ and $U_1, \ldots, U_n$ follows by an argument similar to that in the proof of the ($\Rightarrow$) part of Theorem 3.5. The proof is now complete. □

Now we present an analogue of Theorem 3.9 when the product T is a $C_{\cdot 0}$ contraction, and obviously, a proof follows from Theorem 4.1 and its proof.

THEOREM 4.2 *Let $T_1, \ldots, T_n \in \mathscr{B}(\mathscr{H})$ be commuting contractions such that $T = \prod_{i=1}^n T_i$ is a $C_{\cdot 0}$ contraction. Then, $(T_1, \ldots, T_n)$ possesses an isometric dilation on the minimal isometric dilation space of T if there are projections $P_1, \ldots, P_n$ and commuting unitaries $U_1, \ldots, U_n$ in $\mathscr{B}(\mathscr{D}_{T^*})$ such that the following hold for $i = 1, \ldots, n$:*

(1) $D_T T_i = P_i^{\perp} U_i^* D_T + P_i U_i^* D_T T$,
(2) $P_i^{\perp} U_i^* P_j^{\perp} U_j^* = P_j^{\perp} U_j^* P_i^{\perp} U_i^*$,
(3) $U_i P_i U_j P_j = U_j P_j U_i P_i$.

Conversely, if a commuting tuple of contractions $(T_1, \ldots, T_n)$, with the product $T = \prod_{i=1}^n T_i$ being a $C_{\cdot 0}$ contraction, possesses an isometric dilation $(\widehat{V}_1, \ldots, \widehat{V}_n)$, where $V = \prod_{i=1}^n V_i$ is the minimal isometric dilation of T, then there are unique projections $P_1, \ldots, P_n$ and unique commuting unitaries $U_1, \ldots, U_n$ in $\mathscr{B}(\mathscr{D}_{T^*})$ satisfying the conditions (1)–(3) above.

4.1. A functional model when the product is a $C_{\cdot 0}$ contraction

For a contraction T acting on a Hilbert space $\mathscr{H}$, let Λ_T be the set of all complex numbers for which the operator $I - zT^*$ is invertible. For $z \in \Lambda_T$, the characteristic function of T is defined as

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

$$\Theta_T(z) = [-T + zD_{T^*}(I - zT^*)^{-1}D_T]|_{\mathscr{D}_T}. \tag{4.6}$$

Here, we recall a few definitions and terminologies from the initial part of § 3. The operators D_T and D_{T^*} are the defect operators $(I - T^*T)^{1/2}$ and $(I - TT^*)^{1/2}$, respectively. By virtue of the relation $TD_T = D_{T^*}T$ (section I.3 of [17]), $\Theta_T(z)$ maps $\mathscr{D}_T = \overline{\text{Ran}}D_T$ into $\mathscr{D}_{T^*} = \overline{\text{Ran}}D_{T^*}$ for every z in Λ_T.

In [17], Sz.-Nagy and Foias proved that every $C_{\cdot 0}$ contraction P acting on $\mathscr{H}$ is unitarily equivalent to the operator $\mathbb{T} = P_{\mathbb{H}_T}M_z|_{\mathbb{H}_T}$ on the Hilbert space $\mathbb{H}_T = H^2(\mathscr{D}_{T^*}) \ominus M_{\Theta_T}(H^2(\mathscr{D}_T))$, where M_z is the multiplication operator on $H^2(\mathscr{D}_{T^*})$ and M_{Θ_T} is the multiplication operator from $H^2(\mathscr{D}_T)$ into $H^2(\mathscr{D}_{T^*})$ corresponding to the multiplier Θ_T. This is known as Sz. Nagy–Foias model for a $C_{\cdot 0}$ contraction. Indeed, M_z on $H^2(\mathscr{D}_{T^*})$ dilates $T \in \mathscr{B}(\mathscr{H})$, and $W : \mathscr{H} \to H^2(\mathscr{D}_{T^*})$ as in (4.4) is the concerned isometric embedding. In an analogous manner by an application of Theorem 4.1, we obtain a functional model for a tuple of commuting contractions with $C_{\cdot 0}$ product. A notable fact about this model is that the multiplication operators involved in this model have analytic symbols which are linear functions in one variable.

THEOREM 4.3 *Let $T_1, \ldots, T_n$ be commuting contractions on a Hilbert space $\mathscr{H}$ such that their product $T = \Pi_{i=1}^n T_i$ is a $C_{\cdot 0}$ contraction. If there are projections $P_1, \ldots, P_n$ and commuting unitaries $U_1, \ldots, U_n$ in $\mathscr{B}(\mathscr{D}_{T^*})$ satisfying the following for $1 \leq i \leq n$:*

(1) $D_{T^*}T_i^* = P_i^{\perp}U_i^*D_{T^*} + P_iU_i^*D_{T^*}T^*$,
(2) $P_i^{\perp}U_i^*P_j^{\perp}U_j^* = P_j^{\perp}U_j^*P_i^{\perp}U_i^*$,
(3) $U_iP_iU_jP_j = U_jP_jU_iP_i$,

then $(T_1, \ldots, T_n)$ is unitarily equivalent to $(\widetilde{T}_1, \ldots, \widetilde{T}_n)$ acting on the space $\mathbb{H}_T = H^2(\mathscr{D}_{T^*}) \ominus M_{\Theta_T}(H^2(\mathscr{D}_T))$, where $\widetilde{T}_i = P_{\mathbb{H}_T}(M_{U_iP_i^{\perp}+zU_iP_i})|_{\mathbb{H}_T}$ for $1 \leq i \leq n$.

Proof. For a $C_{\cdot 0}$ contraction T, we have from literature (e.g., [17] or lemma 3.3 in [21]) that

$$WW^* + M_{\Theta_T}M_{\Theta_T}^* = I_{H^2(\mathscr{D}_{P^*})}.$$

It follows from here that $W(\mathscr{H}) = \mathbb{H}_T$, where $W : \mathscr{H} \to H^2(\mathscr{D}_{T^*})$ is as in (4.4). Since $V_i^*|_{W(\mathscr{H})} = T_i^*$, where $V_i = (M_{U_iP_i^{\perp}+zU_iP_i})$, we have that $T_i \cong P_{\mathbb{H}_T}(M_{U_iP_i^{\perp}+zU_iP_i})|_{\mathbb{H}_T}$ for $i = 1, \ldots, n$. □

4.2. A factorization of a $C_{\cdot 0}$ contraction

The model for commuting n-isometries, Theorem 3.2, can be restated in the following way.

THEOREM 4.4 *Let $V_1, \ldots, V_n$ be commuting isometries acting on a Hilbert space $\mathscr{H}$ and let $V = \prod_{i=1}^n V_i$. Then, $V = \prod_{i=1}^n V_i$ is a pure isometry if and only if there are unique orthogonal projections $P_1, \ldots, P_n$ and unique commuting unitaries*

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

$U_1, \dots, U_n$ in $\mathscr{B}(\mathscr{D}_{V^*})$ with $\prod_{i=1}^n U_i = I$ such that the following conditions hold for $i = 1, \dots, n$:

(1) $P_i^{\perp} U_i^* P_j^{\perp} U_j^* = P_j^{\perp} U_j^* P_i^{\perp} U_i^*$,
(2) $U_i P_i U_j P_j = U_j P_j U_i P_i$,
(3) $P_1 + U_1^* P_2 U_1 + U_1^* U_2^* P_3 U_2 U_1 + \dots + U_1^* U_2^* \dots U_{n-1}^* P_n U_{n-1} \dots U_2 U_1 = I_{\mathscr{D}_T}$.

Moreover, $(V_1, \dots, V_n)$ is unitarily equivalent to $(M_{U_1 P_1^{\perp} + z U_1 P_1}, \dots, M_{U_n P_n^{\perp} + z U_n P_n})$ on $H^2(\mathscr{D}_{V^*})$.

Proof. First, suppose $(V_1, \dots, V_n)$ on Hilbert space $\mathscr{H}$ is a commuting n-tuple of isometries with $\prod_{i=1}^n V_i = V$ being a pure isometry. Thus, $\mathscr{H}$ can be identified with $H^2(\mathscr{D}_{V^*})$ via a unitary $\tau : \mathscr{H} \to H^2(\mathscr{D}_{V^*})$, and V can be identified with M_z on $H^2(\mathscr{D}_{V^*})$. Let $\widehat{V}_i = \tau V_i \tau^*$ for $i = 1, 2 \dots, n$. Hence, $(\widehat{V}_1, \dots, \widehat{V}_n)$ is a commuting n-tuple of isometries with $\prod_{i=1}^n \widehat{V}_i = M_z$ on $H^2(\mathscr{D}_{V^*})$. Therefore, $(\widehat{V}_1, \dots, \widehat{V}_n)$ is an isometric dilation of $(V_1, \dots, V_n)$ with M_z being the minimal isometric dilation of V. Therefore, by Theorem 4.1, there are unique commuting unitaries $U_1, \dots, U_n$ and unique orthogonal projections $P_1, \dots P_n$ in $\mathscr{B}(\mathscr{D}_{V^*})$ such that $\prod_{i=1}^n U_i = I$ and that the conditions (1)–(3) are satisfied.

Conversely, suppose there are unique orthogonal projections $P_1, \dots, P_n$ and unique commuting unitaries $U_1, \dots, U_n$ in $\mathscr{B}(\mathscr{D}_{V^*})$ such that $\prod_{i=1}^n U_i = I$ and that the conditions (1)–(3) are satisfied. Let $V_i = M_{U_i P_i^{\perp} + z U_i P_i}$ for each $i = 1, 2 \dots, n$. Then, as seen in the proof of Theorem 4.1, $(V_1, \dots, V_n)$ is a commuting n-tuple of isometries with $\prod_{i=1}^n V_i = M_z$ on $H^2(\mathscr{D}_{V^*})$. □

Also, Theorem 3.2 provides a factorization of a $C_{\cdot 0}$ isometry (i.e., a pure isometry) in terms of n number of commuting isometries. Our result, Theorem 4.1, gives a factorization of a $C_{\cdot 0}$ contraction in the following way:

THEOREM 4.5 *Let $T_1, \dots, T_n$ be commuting contractions on a Hilbert space $\mathscr{H}$ and let their product $T = \prod_{i=1}^n T_i$ be a $C_{\cdot 0}$ contraction. Then, $(T_1^*, \dots, T_n^*) \equiv (V_1^*|_{\mathscr{H}}, \dots, V_n^*|_{\mathscr{H}})$ for a model n-isometry $(V_1, \dots, V_n)$ on $H^2(\mathscr{D}_{T^*})$ if and only if there exist unique orthogonal projections $P_1, \dots, P_n$ and unique commuting unitaries $U_1, \dots, U_n$ in $\mathscr{B}(\mathscr{D}_{T^*})$ such that $\prod_{i=1}^n U_i = I_{\mathscr{D}_{T^*}}$ and the following conditions are satisfied:*

(1) $D_{T^*} T_i^* = P_i^{\perp} U_i^* D_{T^*} + P_i U_i^* D_{T^*} T^*$,
(2) $P_i^{\perp} U_i^* P_j^{\perp} U_j^* = P_j^{\perp} U_j^* P_i^{\perp} U_i^*$,
(3) $U_i P_i U_j P_j = U_j P_j U_i P_i$,
(4) $P_1 + U_1^* P_2 U_1 + U_1^* U_2^* P_3 U_2 U_1 + \dots + U_1^* U_2^* \dots U_{n-1}^* P_n U_{n-1} \dots U_2 U_1 = I_{\mathscr{D}_T}$.

Proof. First suppose there are projections $P_1, \dots, P_n$ and commuting unitaries $U_1, \dots, U_n$ in $\mathscr{B}(\mathscr{D}_{T^*})$ such that $\prod_{i=1}^n U_i = I_{\mathscr{D}_{T^*}}$ and that the conditions (1)–(4) are satisfied. Then, Theorem 4.1 provides commuting n-isometries $V_1, \dots, V_n$ on $H^2(\mathscr{D}_{T^*})$ with $V_i = M_{U_i P_i^{\perp} + z U_i P_i}$ for each i such that

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

$$(T_1^*, \ldots, T_n^*) \equiv (V_1^*|_{W(\mathscr{H})}, \ldots, V_n^*|_{W(\mathscr{H})}),$$

where W is the isometry as in (4.4).

Conversely, suppose $(T_1^*, \ldots, T_n^*)$ is equivalent to $(V_1^*|_{\mathscr{H}}, \ldots, V_n^*|_{\mathscr{H}})$ for some model n-isometry $(V_1, \ldots, V_n)$ on $H^2(\mathscr{D}_{T^*})$. Then, by the $(\Rightarrow)$ part of Theorem 4.1, there are projections $P_1, \ldots, P_n$ and commuting unitaries $U_1, \ldots, U_n$ in $\mathscr{B}(\mathscr{D}_{T^*})$ such that $\prod_{i=1}^n U_i = I_{\mathscr{D}_{T^*}}$ and that the conditions (1)–(4) are satisfied. □

5. Examples

In this section, we present several examples to compare our classes of commuting contractions admitting isometric dilation with the previously determined various classes from the literature. We shall show that neither our classes are properly contained in any of these classes from the literature nor any previously determined classes are contained properly in our classes. However, there are always non-trivial intersections. Note that our theory is one-dimensional in the sense that the operator models that we obtained are all having multiplication operators with multipliers of one complex variable.

Suppose (T_1, T_2) is a commuting pair of contractions admitting isometric dilation (V_1, V_2) on the minimal dilation space of $T = T_1T_2$. If there are unitaries U_1, U_2 and projections P_1, P_2 such that $U_iP_j = U_jP_i$ for $i = 1, 2$ and that the condition (1) of Theorem 3.9 holds, then it can be verified that the conditions (2),(3), and(4) hold as a consequence. Hence, (T_1, T_2) has an isometric dilation on the minimal isometric dilation space of T. Now one may ask a question: if (T_1, T_2) admits isometric dilation on the minimal isometric dilation space of T, then will the corresponding unitaries commute with the projections? The following example gives a negative answer to this:

EXAMPLE 5.1. Let $T_1 = T_2 = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$ be commuting pair of contractions on $\mathbb{C}^2$. Then, clearly, $T = T_1T_2 = 0$, and hence, $D_T = I$. Hence, we need to find commuting U_1, U_2 and projections P_1, P_2 such that $D_TT_i = P_i^{\perp}U_i^*D_T + P_iU_i^*D_TT$ for $i = 1, 2$. Substituting $D_T = I$ and $T = 0$, the above equations are equivalent to $T_1 = P_1^{\perp}U_1^*$ and $T_2 = P_2^{\perp}U_2^*$. One can observe that $P_1 = P_2 = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$ and $U_1 = U_2 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ satisfy the above two equations. It is clear that U_1, U_2 are commuting unitaries and P_1, P_2 are projections. Further as T_1, T_2 commute with each other, it follows that $P_1^{\perp}U_1^*$ commutes with $P_2^{\perp}U_2^*$. Simple calculation shows that $U_1P_1 = T_1$ and $U_2P_2 = T_2$. Therefore, U_1P_1 commutes with U_2P_2. Further $D_TU_iP_iU_i^*D_T = U_iP_iU_i^* = U_iP_iP_iU_i^* = T_iT_i^* = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} = D_{T_i}^2$. Also $P_1 + U_1^*P_2U_1 = I$. Hence, the conditions (1)–(5) of Theorem 3.5 hold. But one can clearly observe that U_i do not commute with P_j as $U_iP_j = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$ and $P_jU_i = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$.

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

Before going to the next example let us note down the following observation from the proofs of the previously stated dilation theorems:

NOTE 5.2. *Let $T_1, \dots, T_n \in \mathscr{B}(\mathscr{H})$ be commuting contractions. Suppose there are projections $P_1, \dots, P_n \in \mathscr{B}(\mathscr{D}_T)$ and commuting unitaries $U_1, \dots, U_n \in \mathscr{B}(\mathscr{D}_T)$ such that $\prod_{i=1}^n U_i = I$ and conditions (1)–(5) of Theorem 3.5 are satisfied. Then, by Theorem 3.5, $(T_1, \dots, T_n)$ possesses an isometric dilation $(V_1, \dots, V_n)$ on the minimal isometric dilation space $\mathscr{K}$ of T such that $\prod_{i=1}^n V_i = V$ is the minimal isometric dilation of T. Without loss of generality, we can assume V_i to be as in (3.3) and V to be the Schäffer's minimal isometric dilation. So, if $V_i' = \prod_{j \neq i} V_j$, then*

$$V_i' = V_i^* V = \begin{bmatrix} T_i' & 0 & 0 & 0 & \dots \\ U_i P_i^\perp D_T & U_i P_i & 0 & 0 & \dots \\ 0 & U_i P_i^\perp & U_i P_i & 0 & \dots \\ 0 & 0 & U_i P_i^\perp & U_i P_i & \dots \\ \dots & \dots & \dots & \dots & \dots \end{bmatrix}.$$

Thus, $(2,1)$ entries of both sides of $V_i' V = V V_i'$ give $D_T T_i' = U_i P_i D_T + U_i P_i^\perp D_T T$ for $1 \leq i \leq n$.

Let $\mathscr{U}^n(\mathscr{H})$ be the class of commuting n-tuples of contractions on $\mathscr{H}$ satisfying conditions (1)–(4) of Theorem 3.9 and let $\mathscr{S}^n(\mathscr{H})$ denote the class satisfying conditions (1)–(5) of Theorem 3.5. The following example shows that $\mathscr{S}^n(\mathscr{H})$ is properly contained in $\mathscr{U}^n(\mathscr{H})$.

EXAMPLE 5.3. Let us consider the following doubly commuting contractions acting on $\mathbb{C}^3$:

$$T_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \; T_2 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \; T_3 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

We show that $(T_1, T_2, T_3) \notin \mathscr{S}^3(\mathbb{C}^3)$ though it belongs to $\mathscr{U}^3(\mathbb{C}^3)$. Note that $T_1' = T_2 T_3 = 0$ and similarly $T_2' = T_3' = 0$. Also, it is clear that $T = T_1 T_2 T_3 = 0$, and hence, $D_T = I$ on $\mathbb{C}^3$. Thus, $\mathscr{D}_T = \mathbb{C}^3$. Now suppose there are commuting unitaries U_1, U_2, U_3 and projections P_1, P_2, P_3 in $\mathscr{B}(\mathbb{C}^3)$ satisfying the hypotheses of Theorem 3.5. Then, for each i, we have that $D_T T_i = P_i^\perp U_i^* D_T + P_i U_i^* D_T T$. Since $T = 0$ and $D_T = I$, it reduces to $T_i = P_i^\perp U_i^*$. Again, from Note 5.2, it is clear that the unitaries and projections satisfying (1)–(5) must also satisfy $D_T T_i' = U_i P_i D_T + U_i^* P_i^\perp D_T T$, which is same as saying that $T_i' = U_i P_i$. Thus, $T_i^* + T_i' = U_i P_i^\perp + U_i P_i = U_i$. Since $T_i' = 0$, we have that $U_i = T_i^* + T_i' = T_i^*$. This contradicts the fact that U_i is a unitary. Hence, $(T_1, T_2, T_3) \notin \mathscr{S}^3(\mathbb{C}^3)$. Now if we take $U_i = I$ and $P_i = I - T_i$ for $i = 1, 2, 3$, then one can easily verify that the conditions (1)–(4) of Theorem 3.9 hold.

In [15], Barik, Das, Haria, and Sarkar introduced a new class of commuting contractions that admit isometric dilation. For each natural number $n \geq 3$ and for every number p, q with $1 \leq p < q \leq n$, let $\mathscr{T}_{p,q}^n(\mathscr{H})$ be defined as follows:

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

$$\mathscr{T}^n_{p,q}(\mathscr{H}) = \{T \in \mathscr{T}^n(\mathscr{H}) : \hat{T}_p, \hat{T}_q \in \mathbb{S}_{n-1}(\mathscr{H}), \text{ and } \hat{T}_p \text{is pure}\}, \tag{5.1}$$

where for any natural number $i \leq n$, $\hat{T}_i = (T_1, \ldots, T_{i-1}, T_{i+1}, \ldots T_n)$, $\mathscr{T}_n(\mathscr{H})$ is a set of commuting n-tuple of contractions on space $\mathscr{H}$ and

$$\mathbb{S}_n(\mathscr{H}) = \{T \in \mathscr{T}^n(\mathscr{H}) : \sum_{k\in\{0,1\}^n} (-1)^{|k|} T^k T^{*k} \geq 0\}.$$

Note that $\mathbb{S}_n(\mathscr{H})$ is the set of all those n-tuples of contractions on $\mathscr{H}$ that satisfy Szego positivity condition, i.e., $\sum_{k\in\{0,1\}^n}(-1)^{|k|}T^kT^{*k} \geq 0$. The class obtained by putting an additional condition $\|T_i\| < 1$ for each i on the elements of $\mathscr{T}^n_{p,q}(\mathscr{H})$ is denoted by $\mathscr{P}^n_{p,q}(\mathscr{H})$. This class has been studied in [36] by Grinshpan, Kaliuzhnyi, Verbovetskyi, Vinnikov, and Woerdeman. In [15], it is shown that $\mathscr{P}^n_{p,q}(\mathscr{H}) \subsetneq \mathscr{T}^n_{p,q}(\mathscr{H})$ for $1 \leq p < q \leq n$. The following example shows that an element of our class may not satisfy the Szego positivity condition.

EXAMPLE 5.4. Let $T_1 = T_2 = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$ and $T_3 = I$. Then, $T = T_1T_2T_3 = 0$, and hence, $D_T = I$. Let $P_1 = P_2 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$, $P_3 = 0$ and $U_1 = U_2 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$, $U_3 = I$. Then U_1, U_2, U_3 are commuting unitaries and P_1, P_2, P_3 are projections satisfying $U_1U_2U_3 = I$ and the conditions (1)–(5) of Theorem 3.5. Hence, (T_1, T_2, T_3) belongs to $\mathscr{S}^3(\mathbb{C}^3)$. Note that

$$I - T_1T_1^* - T_2T_2^* - T_3T_3^* = I - \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} - \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} - I = \begin{bmatrix} 0 & 0 \\ 0 & -2 \end{bmatrix} \not\geq 0.$$

Also,

$$I - T_2T_2^* - T_3T_3^* = I - \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} - I = \begin{bmatrix} 0 & 0 \\ 0 & -1 \end{bmatrix} \not\geq 0$$

and

$$I - T_1T_1^* - T_3T_3^* = I - \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} - I = \begin{bmatrix} 0 & 0 \\ 0 & -1 \end{bmatrix} \not\geq 0.$$

Thus, if we consider $\widehat{T}_1 = (T_2, T_3)$, $\widehat{T}_2 = (T_1, T_3)$ and $\widehat{T}_3 = (T_1, T_2)$, then $\widehat{T}_1$, $\widehat{T}_2$ do not belong to $\mathbb{S}_2(\mathscr{H})$. So, for any p, q satisfying $1 \leq p, q \leq 3$, we have that $(T_1, T_2, T_3) \notin \mathscr{T}^n_{p,q}(\mathscr{H})$.

Note that example 5.4 does not satisfy Brehmer's condition. Recall that $\underline{T} = (T_1, \ldots, T_n)$ satisfies Brehmer's conditions if

$$\sum_{F\subseteq G} (-1)^{|F|} \underline{T}^*_F \underline{T}_F \geq 0$$

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

for all $G \subseteq \{1, \dots n\}$. See (2.1) for the definition. For $\gamma = (\gamma_1, \dots, \gamma_n) \in \mathbb{N}^n$, Curto and Vasilescu [27, 28] introduced a notion of γ contractivity. For $\gamma = (1, \dots, 1) := e$, the notion agrees with Brehmer's condition. The notion of γ contractivity is defined in such a way that for each $\gamma \in \mathbb{N}^n$, if an operator $T = (T_1, \dots, T_n)$ is γ contractive, then it is e contractive. Since example 5.4 does not satisfy Brehmer's condition, it does not have a regular unitary dilation, and it is not e contractive. Hence, it cannot be γ contractive, and consequently, it does not belong to the 'Curto–Vasilescu' class.

Example 5.5. As observed by Barik, Das, Haria, and Sarkar in [15], there is an operator tuple $(M_{z_1}, \dots, M_{z_n})$ in $\mathscr{T}^n_{p,q}(H^2(\mathbb{D}^n))$ that does not belong to $\mathscr{P}^n_{p,q}(H^2(\mathbb{D}^n))$. This class was introduced in [36] by Grinshpan et al. Since $T = M_{z_1} \dots M_{z_n}$ on $H^2(\mathbb{D}^n)$ is an isometry, $\mathscr{D}_T = 0$. So the minimal isometric dilation space of T is $\mathscr{K}_0 = H^2(\mathbb{D}^n)$. Since each $M_{z_i} = T_i$ is an isometry on $H^2(\mathbb{D}^n)$, we have that $(M_{z_1}, \dots, M_{z_n})$ is an isometric dilation of $(T_1, \dots, T_n)$ on the minimal dilation space of T with the product being the minimal isometric dilation of T. Thus, $(M_{z_1}, \dots, M_{z_n}) \in \mathscr{U}^n(\mathscr{K})$ by Theorem 3.5.

Example 5.6. Let us consider the following commuting self-adjoint scalar matrices acting on $\mathbb{C}^3$:

$$T_1 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \; T_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \; T_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

Let $P_i = I - T_i$ and $U_i = I$ for $i = 1, 2, 3$. Note that $D_T = 0$ and the projections P_1, P_2, P_3 satisfy $P_1 + P_2 + P_3 = I$. Thus, the condition (5) holds. Also, it can be easily verified that the conditions (1)–(4) hold. Therefore, this triple of commuting contractions belongs to $\mathscr{S}^3(\mathbb{C}^3)$.

Example 5.7. Let us consider the commuting self-adjoint scalar matrices

$$T_1 = \begin{bmatrix} 0 & 1/3 & 0 \\ 1/3 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \; T_2 = \begin{bmatrix} 1/2 & 1/3 & 0 \\ -1/3 & 1/2 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \; T_3 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

acting on $\mathbb{C}^3$. Evidently, $T = T_1T_2T_3 = 0$, and thus, $D_T = I$. Now suppose there are projections P_1, P_2, P_3 and commuting unitaries U_1, U_2, U_3 satisfying conditions (1)–(4) of Theorem 3.5. So, in particular, we have

$$D_T T_1 = P_1^{\perp} U_1^* D_T + P_1 U_1^* D_T T,$$

which implies that $T_1 = P_1^{\perp} U_1^*$. We also have $D_T U_1 P_1 U_1^* D_T = I - T_1^* T_1$, and this gives $P_1 = U_1^*(I - T_1^* T_1) U_1$, and hence, we have

$$P_1^{\perp} = I - U_1^*(I - T_1^* T_1) U_1 = U_1^* T_1^* T_1 U_1 = U_1^* \begin{bmatrix} 1/9 & 0 & 0 \\ 0 & 1/9 & 0 \\ 0 & 0 & 0 \end{bmatrix} U_1.$$

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

Therefore,

$$T_1 = U_1^* \begin{bmatrix} 1/9 & 0 & 0 \\ 0 & 1/9 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

Again

$$\begin{bmatrix} 3 \\ 0 \\ 0 \end{bmatrix} = T_1 \begin{bmatrix} 0 \\ 9 \\ 0 \end{bmatrix} = U_1^* \begin{bmatrix} 1/9 & 0 & 0 \\ 0 & 1/9 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 0 \\ 9 \\ 0 \end{bmatrix} = U_1^* \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}.$$

This contradicts the fact that U_1 is a unitary. Hence, $(T_1, T_2, T_3) \notin \mathscr{U}^3(\mathbb{C}^3)$.

EXAMPLE 5.8. Let T_1, T_2 be as in example 3.8 and let $T_3 = I$. Then, it is evident that (T_1, T_2, T_3) does not belong to our class $\mathscr{S}^3(\mathbb{C}^3)$. This can be verified using an argument similar to that in example 5.7. However,

$$I - T_2T_2^* - T_3T_3^* + T_2T_3T_2^*T_3^* = I - \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1/3 \end{bmatrix} - I + \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1/3 \end{bmatrix} = 0.$$

Alo,

$$\begin{aligned} I - T_1T_1^* - T_2T_2^* + T_1T_2T_1^*T_2^* &= I - \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1/3 \end{bmatrix} - \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1/9 & 0 \\ 0 & 0 & 1/27 \end{bmatrix} + 0 \\ &= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 8/9 & 0 \\ 0 & 0 & 17/27 \end{bmatrix} \geq 0. \end{aligned}$$

Hence, $\widehat{T}_1$ and $\widehat{T}_3$ belong to $\mathbb{S}_3(\mathbb{C}^3)$. Clearly T_1, T_2 are pure contractions, and hence, $\widehat{T}_3$ is pure. Thus, $(T_1, T_2, T_3) \in \mathscr{T}^3_{1,3}(\mathbb{C}^3)$.

We now consider a few examples of contractions acting on infinite dimensional Hilbert spaces and study their dilations.

EXAMPLE 5.9. Let $\mathscr{H} = l^2$ and let T_1, T_2 be the following weighted shift operators acting on $\mathscr{H}$:

$$T_1(h_0, h_1, \dots) = (0, a_0h_0, 0, a_1h_2, 0, a_2h_4, \dots),$$

$$T_2(h_0, h_1, \dots) = (0, b_0h_0, 0, b_1h_2, 0, b_2h_4, \dots),$$

where $\{a_0, a_1, \dots\}$ and $\{b_0, b_1, \dots\}$ are bounded sequences of non-zero real numbers with $|a_n| < 1$ and $|b_n| < 1$ for every $n \in \mathbb{N}$. Clearly, $T = T_1T_2 = T_2T_1 = 0$. Hence,

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

$D_T = I$ and $\mathscr{D}_T = l^2 = \mathscr{H}$. We show that (T_1, T_2) does not possess isometric dilation (V_1, V_2) on the minimal isometric dilation space of T with $V = V_1V_2$ being the minimal isometric dilation of T. Let us assume the contrary. Then, by Theorem 3.5, there are projections P_1, P_2 commuting unitaries U_1, U_2 in $\mathscr{B}(\mathscr{D}_T) = \mathscr{B}(\mathscr{H})$, with $U_1U_2 = I$ satisfying conditions (1)–(5). Condition (1) gives us $T_i = P_i^{\perp}U_i^*$ for $i = 1, 2$. From condition (4), we have

$$\begin{aligned} & U_1P_1U_1^* = I - T_1^*T_1 \\ \implies & T_1^*T_1 = U_1(I - P_1)U_1^* \\ \implies & T_1^*T_1 = U_1T_1 \quad [\text{as } T_1 = P_1^{\perp}U_1^*]. \end{aligned}$$

Simple calculation shows that $T_1^*T_1 = diag(a_0^2, 0, a_1^2, 0, a_2^2, \dots)$. Hence,

$$\begin{aligned} & U_1T_1(a_0^{-1}, 0, \dots) = T_1^*T_1(a_0^{-1}, 0, \dots) \\ \implies & U_1(0, 1, 0, 0, \dots) = (a_0, 0, 0, \dots). \end{aligned}$$

Since U_1 is a unitary, $\|U_1(0, 1, 0, 0, \dots)\| = \|(0, 1, 0, 0, \dots)\|$, and thus, it follows from here that $\|(a_0, 0, 0, \dots)\| = \|(0, 1, 0, 0, \dots)\|$. This is a contradiction as $|a_0| < 1$. Hence, we are done.

Example 5.10. Let $\mathscr{H} = l^2$ and let $\{e_0, e_1, \dots\}$ be the standard orthonormal basis of l^2. For $k = 1, \dots, n$, let T_k be defined on $\mathscr{H}$ by $T_k(e_m) = 0$ if $m \equiv (k-1) mod\, n$ and $T_k(e_m) = e_m$ otherwise. We show that $(T_1, T_2, \dots, T_n)$ possesses an isometric dilation on the minimal isometric dilation space $\mathscr{K}$ of $T = \prod_{i=1}^{n} T_i$. Evidently, $T = 0$, and thus, $\mathscr{D}_T = \mathscr{H}$. So, we have that $\mathscr{K} = \mathscr{H} \oplus \mathscr{H} \oplus \dots$ is the minimal isometric dilation space of T. Now for each k, define $P_k = I - T_k$ and $U_k = I$ on $\mathscr{D}_T = \mathscr{H}$. Then, it follows that $U_1, \dots, U_n$ are commuting unitaries with $U_1U_2 \dots U_n = I$ and $P_1, \dots, P_n$ are projections. Also, the conditions (1)–(5) of Theorem 3.5 are satisfied straightway. Therefore, Theorem 3.5 guaranties the existence of an isometric dilation of $(T_1, \dots, T_n)$ as desired.

6. Sz. Nagy–Foias-type isometric dilation and functional model

Recall that a c.n.u. contraction T on a Hilbert space $\mathscr{H}$ is a contraction such that there is no non-zero subspace $\mathscr{H}_1$ of $\mathscr{H}$ that reduces T and on $\mathscr{H}_1$ the operator T acts as a unitary. In simple words, a c.n.u. contraction is a contraction without any unitary part. Let T on $\mathscr{H}$ be a c.n.u. contraction and let V on $\mathscr{K}_0$ be the minimal isometric dilation of T. By Wold decomposition, $\mathscr{K}_0$ splits into reducing subspaces $\mathscr{K}_{01}$, $\mathscr{K}_{02}$ of V such that $\mathscr{K}_0 = \mathscr{K}_{01} \oplus \mathscr{K}_{02}$ and that $V|_{\mathscr{K}_{01}}$ is unitarily equivalent to a unilateral shift and $V|_{\mathscr{K}_{02}}$ is a unitary. Then, $\mathscr{K}_{01}$ can be identified with $H^2(\mathscr{D}_{T^*})$ and $\mathscr{K}_{02}$ can be identified with $\overline{\Delta_T(L^2(\mathscr{D}_T))}$, where $\Delta_T(t) = [I_{\mathscr{D}_T} - \Theta_T(e^{it})^*\Theta_T(e^{it})]^{1/2}$ and Θ_T being the characteristic function of the contraction T. For further details, see chapter VI of [17]. Thus,$\mathscr{K}_0 = \mathscr{K}_{01} \oplus \mathscr{K}_{02}$ can be identified

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

with $\mathbb{K}_+ = H^2(\mathscr{D}_{T^*}) \oplus \overline{\Delta_T(L^2(\mathscr{D}_T))}$. Also, the isometry V on $\mathscr{K}_0$ can be realized as $M_z \oplus M_{e^{it}}|_{\overline{\Delta_T(L^2(\mathscr{D}_T))}}$. Therefore, there is a unitary

$$\tau = \tau_1 \oplus \tau_2 : \mathscr{K}_{01} \oplus \mathscr{K}_{02} \to (H^2 \otimes \mathscr{D}_{T^*}) \oplus \overline{\Delta_T(L^2(\mathscr{D}_T))} := \widetilde{\mathbb{K}}_+ \tag{6.1}$$

such that V on $\mathscr{K}_0$ can be realized as $(M_z \otimes I_{\mathscr{D}_{T^*}}) \oplus M_{e^{it}}|_{\overline{\Delta_T(L^2(\mathscr{D}_T))}}$ on $\widetilde{\mathbb{K}}_+$.

If $(T_1, \ldots, T_n)$ is a tuple of commuting contractions acting on $\mathscr{H}$ satisfying the hypotheses of Theorem 3.9, then it possesses an isometric dilation $(V_1, \ldots, V_n)$ on the minimal isometric dilation space $\mathscr{K}_0$ of T. By Wold decomposition of commuting isometries, we have that $\mathscr{K}_{01}$ and $\mathscr{K}_{02}$ are reducing subspaces for each V_i and that

$$V_{i2} = V_i|_{\mathscr{K}_{02}} \tag{6.2}$$

is a unitary for $1 \leq i \leq n$. Now we state our dilation theorem and functional model in the Sz. Nagy–Foias setting. This is another main result of this article.

THEOREM 6.1 *Let $(T_1, \ldots, T_n)$ be a tuple of commuting contractions acting on $\mathscr{H}$ such that $T = \Pi_{i=1}^n T_i$ is a c.n.u. contraction. Suppose there are orthogonal projections $P_1, \ldots, P_n$ and commuting unitaries $U_1, \ldots, U_n$ in $\mathscr{B}(\mathscr{D}_T)$ satisfying*

(1) $D_T T_i = P_i^{\perp} U_i^* D_T + P_i U_i^* D_T T$,
(2) $P_i^{\perp} U_i^* P_j^{\perp} U_j^* = P_j^{\perp} U_j^* P_i^{\perp} U_i^*$,
(3) $U_i P_i U_j P_j = U_j P_j U_i P_i$,
(4) $D_T U_i P_i U_i^* D_T = D_{T_i}^2$

for $1 \leq i < j \leq n$. Then, there are projections $Q_1, \ldots, Q_n$ and commuting unitaries $\widetilde{U}_1, \ldots, \widetilde{U}_n$ in $\mathscr{B}(\mathscr{D}_{T^*})$ such that $(T_1, \ldots, T_n)$ dilates to the tuple of commuting isometries $(\widetilde{V}_{11} \oplus \widetilde{V}_{12}, \ldots, \widetilde{V}_{n1} \oplus \widetilde{V}_{n2})$ on $\widetilde{\mathbb{K}}_+ = H^2 \otimes \mathscr{D}_{T^*} \oplus \overline{\Delta_T(L^2(\mathscr{D}_T))}$, where

$$\widetilde{V}_{i1} = I \otimes \widetilde{U}_i Q_i^{\perp} + M_z \otimes \widetilde{U}_i Q_i \,,$$
$$\widetilde{V}_{i2} = \tau_2 V_{i2} \tau_2^* \,,$$

for unitaries τ_2 and V_{i2} as in (6.1) and (6.2), respectively, for $1 \leq i \leq n$.

Proof. By Theorem 3.9, we have that $(T_1, \ldots, T_n)$ possesses an isometric dilation $(V_1, \ldots, V_n)$ on $\mathscr{K}_0 = \mathscr{H} \oplus l^2(\mathscr{D}_T)$, which in fact satisfies $V_i^*|_{\mathscr{H}} = T_i^*$ for $1 \leq i \leq n$. Now $\mathscr{K}_0$ has an orthogonal decomposition $\mathscr{K}_0 = \mathscr{K}_{01} \oplus \mathscr{K}_{02}$ such that $\mathscr{K}_{01}$ and $\mathscr{K}_{02}$ are common reducing subspaces for $V_1, \ldots, V_n$, $(V_1|_{\mathscr{K}_{01}}, \ldots, V_n|_{\mathscr{K}_{01}})$ is a pure isometric tuple, i.e., $\prod_{i=}^n V_i|_{\mathscr{K}_{01}}$ is a pure isometry and $(V_1|_{\mathscr{K}_{02}}, \ldots, V_n|_{\mathscr{K}_{02}})$ is a unitary tuple. Let $\prod_{i=1}^n V_i = V$, $V_i|_{\mathscr{K}_{01}} = V_{i1}$, and $V_i|_{\mathscr{K}_{02}} = V_{i2}$ for $1 \leq i \leq n$. Also, let $\prod_{i=1}^n V_{i1} = W_1$ and $\prod_{i=1}^n V_{i2} = W_2$. So W_1 on $\mathscr{K}_{01}$ is a pure isometry and W_2 on $\mathscr{K}_{02}$ is a unitary. So,

$$(\tau V_1 \tau^*, \ldots, \tau V_n \tau^*) = (\tau_1 V_{11} \tau_1^* \oplus \tau_2 V_{12} \tau_2^*, \ldots, \tau_1 V_{n1} \tau_1^* \oplus \tau_2 V_{n2} \tau_2^*)$$

is an isometric dilation of $(T_1, \ldots, T_n)$ on $\widetilde{\mathbb{K}}_+$, where $\tau = \tau_1 \oplus \tau_2$ is as in (6.1). Thus, the tuple $(\tau_1 V_{11} \tau_1^*, \ldots \tau_1 V_{n1} \tau_1^*)$ on $H^2 \otimes \mathscr{D}_{T^*}$ is a pure isometric tuple. Hence, by

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

Theorem 3.1, there exist commuting unitaries $\widetilde{U}_1, \ldots, \widetilde{U}_n$ and orthogonal projections $Q_1, \ldots, Q_n$ in $\mathscr{B}(\mathscr{D}_{T^*})$ $(= \mathscr{B}(D_{W_1^*}))$ such that the tuple $(\tau_1 V_{11}\tau_1^*, \ldots \tau_1 V_{n1}\tau_1^*)$ is unitarily equivalent to

$$(I \otimes \widetilde{U}_1 Q_1^\perp + M_z \otimes \widetilde{U}_1 Q_1, \ldots, I \otimes \widetilde{U}_n Q_n^\perp + M_z \otimes \widetilde{U}_n Q_n) \quad \text{on} \quad H^2 \otimes \mathscr{D}_{T^*}$$

via a unitary, say Z. For $1 \leq i \leq n$, let us denote $\widetilde{V}_{i1} = I \otimes \widetilde{U}_i Q_i^\perp + M_z \otimes \widetilde{U}_i Q_i$ and $\widetilde{V}_{i2} = \tau_2 V_{i2}\tau_2^*$. So, $(\tau V_1 \tau^*, \ldots, \tau V_n \tau^*)$ is unitarily equivalent to $(\widetilde{V}_{11} \oplus \widetilde{V}_{12}, \ldots, \widetilde{V}_{n1} \oplus \widetilde{V}_{n2})$ via the unitary $Z \oplus I$. Thus, $(V_1, \ldots, V_n)$ is unitarily equivalent to $(\widetilde{V}_{11} \oplus \widetilde{V}_{12}, \ldots, \widetilde{V}_{n1} \oplus \widetilde{V}_{n2})$ via a unitary

$$Y = Z\tau_1 \oplus \tau_2 : \mathscr{K}_{01} \oplus \mathscr{K}_{02} \to H^2 \otimes \mathscr{D}_{T^*} \oplus \overline{\Delta_T(L^2(\mathscr{D}_T))}.$$

Let $\widetilde{V}_i = \widetilde{V}_{i1} \oplus \widetilde{V}_{i2}$ for $i = 1, \ldots n$. Since for each i, $V_i^*|_{\mathscr{H}} = T_i^*$, we have, for any $h \in \mathscr{H}$,

$$\widetilde{V}_i^*(Yh) = (YV_iY^*)^*Yh = YV_i^*Y^*Yh = YV_i^*h = YT_i^*h.$$

Therefore, $\widetilde{V}_i^*|_{Y(\mathscr{H})} = YT_i^*Y^*|_{Y(\mathscr{H})}$ for $1 \leq i \leq n$. So, we have $T_1^{k_1} \ldots T_n^{k_n} = Y^*\widetilde{V}_1^{k_1} \ldots \widetilde{V}_n^{k_n} Y$. Thus, $(\widetilde{V}_{11} \oplus \widetilde{V}_{12}, \ldots, \widetilde{V}_{n1} \oplus \widetilde{V}_{n2})$ is an isometric dilation of $(T_1, \ldots, T_n)$, where $\widetilde{V}_{i1} = I \otimes \widetilde{U}_i Q_i^\perp + M_z \otimes \widetilde{U}_i Q_i$ and $\widetilde{V}_{i2} = \tau_2 V_{i2}\tau_2^*$ for $1 \leq i \leq n$. This completes the proof. □

7. A model theory for a class of commuting contractions

In this section, we present a model theory for a tuple of commuting contractions satisfying the conditions of Theorems 3.5 and 3.9.

THEOREM 7.1 *Let $(T_1, \ldots, T_n)$ be commuting tuple of contractions on a Hilbert space $\mathscr{H}$ and let $T = \Pi_{i=1}^n T_i$. Suppose there are projections $P_1, \ldots, P_n \in \mathscr{B}(\mathscr{D}_{T^*})$ and commuting unitaries $U_1, \ldots, U_n \in \mathscr{B}(\mathscr{D}_{T^*})$ such that for each $i = 1, \ldots, n$,*

(1) $D_{T^*}T_i^* = P_i^\perp U_i^* D_{T^*} + P_i U_i^* D_{T^*} T^*$,
(2) $P_i^\perp U_i^* P_j^\perp U_j^* = P_j^\perp U_j^* P_i^\perp U_i^*$,
(3) $U_i P_i U_j P_j = U_j P_j U_i P_i$,
(4) $D_{T^*} U_i P_i U_i^* D_{T^*} = D_{T_i^*}^2$.

Let $Z_1, \ldots, Z_n$ on $\mathscr{K} = \mathscr{H} \oplus l^2(\mathscr{D}_{T^*})$ be defined as follows:

$$Z_i = \begin{bmatrix} T_i & D_{T^*}U_iP_i & 0 & 0 & \ldots \\ 0 & U_iP_i^\perp & U_iP_i & 0 & \ldots \\ 0 & 0 & U_iP_i^\perp & U_iP_i & \ldots \\ 0 & 0 & 0 & U_iP_i^\perp & \ldots \\ \ldots & \ldots & \ldots & \ldots & \ldots \end{bmatrix}, \qquad (1 \leq i \leq n).$$

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

Then,

(i) $(Z_1, \ldots, Z_n)$ *is a commuting n-tuple of co-isometries,* $\mathscr{H}$ *is a common invariant subspace of* $Z_1, \ldots, Z_n$*, and* $Z_i|_{\mathscr{H}} = T_i$*.*

(ii) there is an orthogonal decomposition $\mathscr{K} = \mathscr{K}_1 \oplus \mathscr{K}_2$ *into common reducing subspaces of* $Z_1, \ldots, Z_n$ *such that* $(Z_1|_{\mathscr{K}_1}, \ldots, Z_n|_{\mathscr{K}_1})$ *is a pure co-isometric tuple, that is,* $Z|_{\mathscr{K}_1} = \Pi_{i=1}^n Z_i|_{\mathscr{K}_1}$ *is a pure co-isometry and* $(Z_1|_{\mathscr{K}_2}, \ldots, Z_n|_{\mathscr{K}_2})$ *is a unitary tuple.*

Additionally, if

$$(5)\ P_1 + U_1^* P_2 U_1 + U_1^* U_2^* P_3 U_2 U_1 + \ldots + U_1^* U_2^* \ldots U_{n-1}^* P_n U_{n-1} \ldots U_2 U_1 = I_{\mathscr{D}_T},$$

then

(iii) $\mathscr{K}_1$ *can be identified with* $H^2(\mathscr{D}_Z)$*, where* $\mathscr{D}_Z$ *has same dimension as that of* $\mathscr{D}_T$*. Also, there are projections* $\widehat{P_1}, \ldots, \widehat{P_n}$ *and commuting unitaries* $\widehat{U_1}, \ldots, \widehat{U_n}$ *such that the operator tuple* $(Z_1|_{\mathscr{K}_1}, \ldots, Z_n|_{\mathscr{K}_1})$ *is unitarily equivalent to the multiplication operator tuple*

$$\left(M_{\widehat{U_1}\widehat{P_1}^{\perp} + \widehat{U_1}\widehat{P_1}\bar{z}}, \ldots, M_{\widehat{U_n}\widehat{P_n}^{\perp} + \widehat{U_n}\widehat{P_n}\bar{z}}\right)$$

acting on $H^2(\mathscr{D}_T)$*.*

Proof. We apply Theorem 3.9 to the tuple $(T_1^*, \ldots, T_n^*)$ of commuting contractions to have an isometric dilation $(X_1, \ldots, X_n)$ on $\mathscr{K}_0 = \mathscr{H} \oplus l^2(\mathscr{D}_{T^*})$, where

$$X_i = \begin{bmatrix} T_i^* & 0 & 0 & 0 & \ldots \\ P_i U_i^* D_{T^*} & P_i^{\perp} U_i^* & 0 & 0 & \ldots \\ 0 & P_i U_i^* & P_i^{\perp} U_i^* & 0 & \ldots \\ 0 & 0 & P_i U_i^* & P_i^{\perp} U_i^* & \ldots \\ \ldots & \ldots & \ldots & \ldots & \ldots \end{bmatrix}, \qquad (1 \leq i \leq n).$$

Clearly, $Z_i = X_i^*$ for each i, and it is evident from the block matrix that $\mathscr{H}$ is a common invariant subspace for each Z_i and $Z_i|_{\mathscr{H}} = T_i$. This proves (i). Since $(X_1, \ldots, X_n)$ is a commuting tuple of isometry, $X = \Pi_{i=1}^n X_i$ is an isometry. By Wold decomposition, $\mathscr{K} = \mathscr{K}_1 \oplus \mathscr{K}_2$ and that $X|_{\mathscr{K}_1}$ is a pure isometry, $X|_{\mathscr{K}_2}$ is a unitary. Also, they are common reducing subspaces for each X_i. Indeed, if

$$X_i = \begin{pmatrix} A_i & B_i \\ C_i & D_i \end{pmatrix} \qquad X = \begin{pmatrix} X_{K1} & 0 \\ 0 & X_{K2} \end{pmatrix}$$

with respect to the decomposition $\mathscr{K} = \mathscr{K}_1 \oplus \mathscr{K}_2$, then $X_i X = X X_i$ implies that

$$B_i X_{K2} = X_{K1} B_i \qquad C_i X_{K1} = X_{K2} C_i.$$

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

Therefore, for all $k \in \mathbb{N}$, we have

$$X_{K2}^{*k} B_i^* = B_i^* X_{K1}^{*k} \qquad X_{K1}^{*n} C_i^* = C_i^* X_{K2}^{*k}.$$

Now X_{K1} is a pure isometry and X_{K2} is unitary, so on the one hand, we have $\|X_{K2}^{*k} B_i^*\| = \|B_i^*\|$, and on the other hand, $\|B_i^* X_{K1}^{*k}\| \to 0$ as $k \to \infty$. Hence, $B_i = 0$. Similarly, $C_i = 0$ for $1 \leq i \leq n$. Thus, with respect to the decomposition $\mathscr{K} = \mathscr{K}_1 \oplus \mathscr{K}_2$, the operator X_i takes form

$$X_i = \begin{pmatrix} X_{i1} & 0 \\ 0 & X_{i2} \end{pmatrix}, \qquad (1 \leq i \leq n).$$

Now since X_i is an isometry, so will be X_{i1}, $X'_{i1} = \Pi_{j \neq i} X_{j1}$ and X_{i2}, $X'_{i2} = \Pi_{j \neq i} X_{j2}$. Further from the block matrix of X_i and from the fact that $X = \Pi_{i=1}^n X_i$, it is clear that $X_{K2} = \Pi_{i=1}^n X_{i2}$. Again, X_{K2} is a unitary, X'_{i2} is an isometry, and $X_{i2'}^* X_{K2} = X_{i2}$. So, we have

$$X_{i2} X_{i2}^* = X_{i2'}^* X_{K2} X_{K2}^* X_{i2'} = I, \qquad (1 \leq i \leq n).$$

Thus, X_{i2} is a unitary on $\mathscr{K}_2$ for each $i = 1, \ldots, n$. Further, $(X_{11}, \ldots, X_{n1})$ is a pure isometric tuple as $X_{K1} = \Pi_{i=1}^n X_{i1}$ is a pure isometry. Since $Z_i = X_i^*$, we have that $(Z_1|_{\mathscr{K}_1}, \ldots, Z_n|_{\mathscr{K}_1})$ is pure co-isometric tuple and $(Z_1|_{\mathscr{K}_2}, \ldots, Z_n|_{\mathscr{K}_2})$ is a unitary tuple. Hence, (ii) holds.

Additionally, if (5) holds, then the dilation $(X_1, \ldots, X_n)$ satisfies the condition that X is the Schäffer's minimal isometric dilation of T^* by Theorem 3.5. Let us denote $Z = X^*$. Then, Z is a co-isometry as X is an isometry and

$$Z = \begin{bmatrix} T & D_{T^*} & 0 & 0 & \ldots \\ 0 & 0 & I_{\mathscr{D}_{T^*}} & 0 & \ldots \\ 0 & 0 & 0 & I_{\mathscr{D}_{T^*}} & \ldots \\ \vdots & \vdots & \vdots & \vdots & \ldots \end{bmatrix}.$$

Note that the dimensions of $\mathscr{D}_Z$ and $\mathscr{D}_T$ are same. Indeed, if $\tau : \mathscr{D}_T \to \mathscr{D}_Z$ is defined by $\tau D_T h = D_Z h$ for all $h \in \mathscr{H}$ and extended continuously to the closure, then τ is a unitary. We recall the proof here. Since X is the minimal isometric dilation of T^*, we have

$$\mathscr{K} = \overline{span}\{X^k h : k \geq 0,\, h \in \mathscr{H}\} = \overline{span}\{Z^{*k} h : k \geq 0,\, h \in \mathscr{H}\}.$$

Now for $n \in \mathbb{N}$ and $h \in \mathscr{H}$, we have

$$D_Z^2 Z^{*n} h = (I - Z^* Z) Z^{*n} h = Z^{*n} - Z^* Z^n Z^* h = 0.$$

Therefore, $D_Z X^n h = 0$ for any $n \in \mathbb{N}$ and $h \in \mathscr{H}$. So, $\mathscr{D}_Z = \overline{D_Z \mathscr{K}} = \overline{D_Z \mathscr{H}}$. Also,

$$\|D_Z h\|^2 = \{(I - Z^* Z)h, h\} = \|h\|^2 - \|Zh\|^2 = \|h\|^2 - \|Th\|^2 = \|D_T h\|^2.$$

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

Therefore, τ is a unitary. By Theorem 3.1, we have that

$$(X_{11},\dots,X_{n1}) \cong (M_{\widetilde{U}_1Q_1^{\perp}+z\widetilde{U}_1Q_1},\dots,M_{\widetilde{U}_nQ_n^{\perp}+z\widetilde{U}_nQ_n}),$$

where $Q_1,\dots,Q_n$ are projections and $\widetilde{U}_1,\dots\widetilde{U}_n$ are commuting unitaries from $\mathscr{B}(\mathscr{D}_{X^*_{K1}})$ satisfying

$$D^2_{X'^*_{i1}}X^*_{i1} = D_{X^*_{K1}}Q_i^{\perp}\widetilde{U}_i^*D_{X^*_{K1}} \tag{7.1}$$

and

$$D^2_{X^*_{i1}}X'^*_{i1} = D_{X^*_{K1}}\widetilde{U}_iQ_iD_{X^*_{K2}} \tag{7.2}$$

for all $i=1,\dots,n$. Using the fact that $X_{i2},\dots,X_{n2}$ are unitaries on $\mathscr{K}_2$, it follows that

$$\begin{aligned} D^2_{X_i'^*} &= I_{\mathscr{K}} - \begin{bmatrix} X'_{i1} & 0 \\ 0 & X'_{i2}\end{bmatrix}\begin{bmatrix} X'^*_{i1} & 0 \\ 0 & X'^*_{i2}\end{bmatrix} \\ &= \begin{bmatrix} I_{\mathscr{K}_1}-X'_{i1}X'^*_{i1} & 0 \\ 0 & I_{\mathscr{K}_2}-X'_{i2}X'^*_{i2}\end{bmatrix} = \begin{bmatrix} I_{\mathscr{K}_1}-X'_{i1}X'^*_{i1} & 0 \\ 0 & 0\end{bmatrix}. \end{aligned}$$

Therefore, $D_{X_i'^*} = D_{X'^*_{i1}}\oplus 0$. Similarly, we can prove that $D_{X_i^*} = D_{X^*_{i1}}\oplus 0$ for $1\le i\le n$, with respect to the above decomposition of $\mathscr{K}$. So, $D_{X^*} = D_{X^*_{K1}}\oplus 0$. Hence $\mathscr{D}_{X^*_{K1}} = \mathscr{D}_{X^*} = \mathscr{D}_Z$. Let us denote $\widehat{U_i} = \tau^*\widetilde{U}_i\tau$ and $\widehat{Q_i} = \widehat{P_i}$ for $1\le i\le n$. Thus, $\widehat{U_1},\dots,\widehat{U_n}$ are commuting unitaries and $\widehat{P_1},\dots,\widehat{P_n}$ are projections in $\mathscr{B}(\mathscr{D}(T))$ such that $(Z_1|_{\mathscr{K}_1},\dots Z_n|_{\mathscr{K}_1})$ is unitarily equivalent to $\left(M_{\widetilde{U}_1Q_1^{\perp}+z\widetilde{U}_1Q_1},\dots,M_{\widetilde{U}_nQ_n^{\perp}+z\widetilde{U}_nQ_n}\right)$, which can be realized as $\left(M_{\widehat{U_1}\widehat{P_1}^{\perp}+\widehat{U_1}\widehat{P_1}z},\dots,M_{\widehat{U_n}\widehat{P_n}^{\perp}+\widehat{U_n}\widehat{P_n}z}\right)$ on $H^2(\mathscr{D}_T)$ via the unitary τ. This proves (*iii*) and the proof is complete. □

Apart from having the explicit constructions of isometric dilations and functional model for a commuting contractive tuple, another interesting consequence of Theorem 3.5 is that it gives a commutant lifting in several variables as discussed in Remark 3.6. We conclude this article here. There will be two more articles in this direction as sequels. One of them will describe explicit constructions of minimal unitary dilations of commuting contractions $(T_1,\dots,T_n)$ on the minimal unitary dilation spaces of $T=\prod_{i=1}^n T_i$. The other article will deal with dilations when the defect spaces $\mathscr{D}_T,\mathscr{D}_{T^*}$ are finite dimensional and their interplay with distinguished varieties in the polydisc.

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

Acknowledgements
S.P. is supported by the Seed Grant of IIT Bombay, the CPDA, and the 'Core Research Grant' with Award No. CRG/2023/005223 of Science and Engineering Research Board, India. P.S. has been supported by the PhD Fellowship of Council of Scientific and Industrial Research, India.

Data availability statement
(1) Data sharing is not applicable to this article as no datasets were generated or analysed during the current study.
(2) In case any datasets are generated during and/or analysed during the current study, they must be available from the corresponding author on reasonable request.

References

[1] J. Agler. The Arveson extension theorem and coanalytic models. *Integ. Equ. Oper. Theory*. **5** (1982), 608–631.
[2] J. Agler. Hypercontractions and subnormality. *J. Oper. Theory*. **13** (1985), 203–217.
[3] J. Agler and J. McCarthy. Distinguished varieties. *Acta Math*. **194** (2005), 133–153.
[4] T. Ando. On a pair of commutative contractions. *Acta Sci. Math. (Szeged)*. **24** (1963), 88–90.
[5] A. Arias and G. Popescu. Noncommutative interpolation and Poisson transforms II. *Houston J. Math*. **25** (1999), 79–98.
[6] A. Arias and G. Popescu. Noncommutative interpolation and Poisson transforms. *Israel J. Math*. **115** (2000), 205–234.
[7] W. Arveson. Subalgebras of C^*-algebras. II. *Acta Math*. **128** (1972), 271–308.
[8] W. Arveson. Subalgebras of C^*-algebras III: multivariable operator theory. *Acta Math*. **181** (1998), 159–228.
[9] A. Athavale. Holomorphic kernels and commuting operators. *Trans. Amer. Math. Soc*. **304** (1987), 101–110.
[10] A. Athavale. Model theory on the unit ball in $\mathbb{C}^m$. *J. Oper. Theory*. **27** (1992), 347–358.
[11] A. Athavale. On the intertwining of joint isometries. *J. Oper. Theory*. **23** (1990), 339–350.
[12] B. Bagchi, T. Bhattacharyya and G. Misra. Some thoughts on Ando's theorem and Parrott's example. *Linear Algebra Appl*. **341** (2002), 357–367.
[13] J. A. Ball, W. S. Li, D. Timotin and T. T. Trent. A commutant lifting theorem on the polydisc. *Indiana Univ. Math. J*. **48** (1999), 653–675.
[14] J. Ball, T. Trent and V. Vinnikov. *Interpolation and commutant lifting for multipliers on reproducing Kernel Hilbert spaces, operator theory and analysis*, Oper. Theory Adv. Appl.. Vol.122, pp. 89–138, (Birkhäuser, Basel, 2001).
[15] S. Barik, B. K. Das, K. J. Haria and J. Sarkar. Isometric dilations and von Neumann inequality for a class of tuples in the polydisc. *Trans. Amer. Math. Soc*. **372** (2019), 1429–1450.
[16] H. Bercovici, R. G. Douglas and C. Foias. On the classification of multi-isometries. *Acta Sci. Math. (Szeged)*. **72** (2006), 639–661.
[17] H. Bercovici, C. Foias, L. Kerchy and B. Sz.-Nagy. *Harmonic analysis of operators on Hilbert space*, Universitext. (Springer, New York, 2010).
[18] C. A. Berger, L. A. Coburn and A. Lebow. Representation and index theory for C^*-algebras generated by commuting isometries. *J. Funct. Anal*. **27** (1978), 51–99.
[19] B. V. R. Bhat, T. Bhattacharyya and S. Dey. Standard noncommuting and commuting dilations of commuting tuples. *Trans. Amer. Math. Soc*. **356** (2004), 1551–1568.
[20] T. Bhattacharyya. Dilation of contractive tuples: a survey. Surveys in analysis and operator theory (Canberra, 2001), Proc. Centre Math. Appl. Austral. Nat. Univ., Vol. 40, pp. 89–126 (Austral. Nat. Univ., Canberra, 2002).
[21] T. Bhattacharyya and S. Pal. A functional model for pure Γ-contractions. *J. Oper. Theory*. **71** (2014), 327–339.

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

[22] P. Binding, D. R. Farenick and C. -K. Li. A dilation and norm in several variable operator theory. *Canad. J. Math.* **47** (1995), 449–461.

[23] S. Brehmer. Über vetauschbare Kontraktionen des Hilbertschen Raumes. *Acta Sci. Math. (Szeged)*. **22** (1961), 106–111.

[24] J. W. Bunce. Models for n-tuples of noncommuting operators. *J. Funct. Anal*. **57** (1984), 21–30.

[25] M. D. Choi and C. K. Li. Constrained unitary dilations and numerical ranges. *J. Oper. Theory*. **46** (2001), 435–447.

[26] M. Crabb and A. Davie. von Nemann's inequality for Hilbert space operators. *Bull. London Math. Soc*. **7** (1975), 49–50.

[27] R. Curto and F. H. Vasilescu. Standard operator models in the polydisc. *Indiana Univ. Math. J*. **42** (1993), 791–810.

[28] R. Curto and F. H. Vasilescu. Standard operator models in the polydisc II. *Indiana Univ. Math. J*. **44** (1995), 727–746.

[29] B. K. Das and J. Sarkar. Ando dilations, von Neumann inequality, and distinguished varieties. *J. Funct. Anal*. **272** (2017), 2114–2131.

[30] C. Davis. Some dilation and representation theorems, Proceedings of the Second International Symposium in West Africa on Functional Analysis and its Applications (Kumasi, Ghana, 1979), pp. 159–182.

[31] S. Dey. Standard commuting dilations and liftings. *Colloq. Math*. **126** (2012), 87–94.

[32] R. G. Douglas. Structure theory for operators I. *J. Reine Angew. Math*. **232** (1968), 180–193.

[33] S. Drury. A generalization of von Neumann's inequality to the complex ball. *Proc. Amer. Math. Soc*. **68** (1978), 300–304.

[34] S. Drury. Remarks on von Neumann's inequality. *Banach spaces, harmonic analysis and probability theory*, Lecture Notes in Math., Vol.995, pp. 14–32 (Springer, Berlin, 1983).

[35] A. E. Frazho. Models for non-commuting operators. *J. Funct. Anal*. **48** (1982), 1–11.

[36] A. Grinshpan, D. S. Kaliuzhnyi-Verbovetskyi, V. Vinnikov and H. J. Woerdeman. Classes of tuples of commuting contractions satisfying the multivariable von Neumann inequality. *J. Funct. Anal*. **256** (2009), 3035–3054.

[37] P. R. Halmos. Normal dilations and extensions of operators. *Summa Brasil. Math*. **2** (1950), 125–134.

[38] I. Halperin. Sz.-Nagy-Brehmer dilations. *Acta Sci. Math. (Szeged)*. **23** (1962), 279–289.

[39] I. Halperin. Intrinsic description of the Sz.-Nagy–Brehmer unitary dilation. *Studia Math*. **22** (1962/1963), 211–219.

[40] J. A. Holbrook. Schur norms and the multivariate von Neumann inequality. *Recent Advances in Operator Theory and Related Topics (Szeged, 1999)* Oper. Theory Adv. Appl., Vol.127, pp. 375–386 (Birkhäuser, Basel, 2001).

[41] G. Julia. Les projections des systèmes orthonormaux de l'espace Hilbertien. *C. R. Acad. Sci. Paris*. **218** (1944), 892–895.

[42] G. Julia. Les projections des systèmes orthonormaux de l'espace Hilbertien et les opérateurs bornés. *C. R. Acad. Sci. Paris*. **219** (1944), 8–11.

[43] G. Julia. Sur la représentation analytique des opérateurs bornés ou fermés de l'espace Hilbertien. *C. R. Acad. Sci. Paris*. **219** (1944), 225–227.

[44] G. Knese. The von Neumann inequality for 3×3 matrices. *Bull. London Math. Soc*. **48** (2016), 53–57.

[45] E. Levy and O. Moshe Shalit. Dilation theory in finite dimensions: the possible, the impossible and the unknown. *Rocky Mountain J. Math*. **44** (2014), 203–221.

[46] V. Müller and F. H. Vasilescu. Standard models for some commuting multishifts. *Proc. Amer. Math. Soc*. **117** (1993), 979–989.

[47] J. McCarthy and O. Shalit. Unitary n-dilations for tuples of commuting matrices. *Proc. Amer. Math. Soc*. **141** (2013), 563–571.

[48] S. Parrott. Unitary dilations for commuting contractions. *Pacific J. Math*. **34** (1970), 481–490.

[49] G. Popescu. Isometric dilations for infinite sequences of noncommuting operators. *Trans. Amer. Math. Soc*. **316** (1989), 523–536.

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press

[50] G. Popescu. Models for infinite sequences of noncommuting operators. *Acta Sci. Math. (Szeged)*. **53** (1989), 355–368.

[51] G. Popescu. Characteristic functions for infinite sequences of noncommuting operators. *J. Oper. Theory*. **22** (1989), 51–71.

[52] G. Popescu. Poisson transforms on some -algebras generated by isometries. *J. Funct. Anal*. **161** (1999), 27–61.

[53] G. Popescu. Curvature invariant for Hilbert modules over free semigroup algebras. *Adv. Math*. **158** (2001), 264–309.

[54] H. Sau. Andó dilations for a pair of commuting contractions: two explicit constructions and functional models. https://arxiv.org/abs/1710.11368.

[55] O. M. Shalit. Dilation theory: a guided tour. *Operator theory, functional analysis and applications* Oper. Theory Adv. Appl., Vol.282, pp. 551–623 (Birkhäuser/Springer, Cham, 2021).

[56] J. Stochel and F. H. Szafraniec. Unitary dilation of several contractions. *Oper. Theory Adv. Appl*. **127** (2001), 585–598.

[57] B. Sz.-Nagy. Sur les contractions de I'espace de Hilbert. *Acta Sci. Math. (Szeged)*. **15** (1953), 87–92.

[58] B. Sz.-Nagy. Transformations of Hilbert space, positive definite functions on a semigroup. *Usp. Mat. Nauk*. **11** (1956), 173–182.

[59] D. Timotin. Regular dilations and models for multicontractions. *Indiana Univ. Math. J*. **47** (1998), 671–684.

[60] N. Varopoulos. On an inequality of von Neumann and an application of the metric theory of tensor products to operator theory. *J. Funct. Anal*. **16** (1974), 83–100.

[61] F. H. Vasilescu. An operator-valued Poisson kernel. *J. Funct. Anal*. **110** (1992), 47–72.

[62] J. von Neumann. Eine Spektraltheorie für allgemeine Operatoren eines unitären Raumes. *Math. Nachr*. **4** (1951), 258–281.

https://doi.org/10.1017/prm.2024.95 Published online by Cambridge University Press