
108.26 PWW: A property of triangular numbers

$$T_n = \frac{n(n+1)}{2} \Rightarrow T_{3n+1} - T_n = (2n+1)^2$$



10.1017/mag.2024.79 © The Authors, 2024
Published by Cambridge
University Press on behalf of
The Mathematical Association

K. B. SUBRAMANIAM
Ganit Ashram,
135, Fine Avenue Phase-I,
Nayapura, Kolar Road,
Bhopal- 462042. India
e-mail: *kbsubramaniam.1950@gmail.com*
AMARNATH MURTHY
D-503, Sharan Residency,
New C G Road, Chandkheda,
Ahmedabad, Gujarat- 382424. India
e-mail: *amaranth.1960@gmail.com*

108.27 More on the Euler limit for *e*

The well-known *Euler limit* is defined as $\lim_{n\to\infty}\left(\frac{n+1}{n}\right)^n = e = 2.71828\ldots$ (see for example [1]). Recently, in [2], appeared the following generalisation of the Euler limit.

https://doi.org/10.1017/mag.2024.80 Published online by Cambridge University Press

Theorem 1: Let A_n be a strictly increasing sequence of positive reals satisfying $A_{n+1} \sim A_n$. Then

$$\lim_{n \to \infty} \left(\frac{A_{n+1}}{A_n} \right)^{\frac{A_n}{A_{n+1} - A_n}} = e.$$

Note that the symbol "$\sim$" means asymptotic equivalence, i.e., $x_n \sim y_n$ if $\lim_{n \to \infty} \frac{x_n}{y_n} = 1$.

Here, we offer the following generalisation.

Theorem 2: Let A_n be a strictly monotone sequence of positive reals satisfying $A_{n+1} \sim A_n$. Let B_n be any sequence of reals satisfying $B_n \sim \frac{A_n}{A_{n+1} - A_n}$. Then

$$\lim_{n \to \infty} \left(\frac{A_{n+1}}{A_n} \right)^{B_n} = e.$$

Proof: First, we consider the case of A_n monotone increasing. Theorem 1 gives

$$\lim_{n \to \infty} \left(\frac{A_{n+1}}{A_n} \right)^{B_n} = \lim_{n \to \infty} \left(\left(\frac{A_{n+1}}{A_n} \right)^{\frac{A_n}{A_{n+1} - A_n}} \right)^{\frac{B_n (A_{n+1} - A_n)}{A_n}} = e^1 = e.$$

Now we consider the other case, of A_n monotone decreasing. We set $A_n' = \frac{1}{A_n}$ and $B_n' = B_n$ to get

$$\lim_{n \to \infty} \left(\frac{A_{n+1}}{A_n} \right)^{B_n} = \lim_{n \to \infty} \left(\frac{A_n'}{A_{n+1}'} \right)^{B_n'}.$$

We conclude by observing that $B_n \sim \dfrac{A_n}{A_{n+1} - A_n} = -\dfrac{A_{n+1}'}{A_{n+1}' - A_n'} \sim -\dfrac{A_n'}{A_{n+1}' - A_n'}$, and applying the first case to B_n' and the monotone increasing A_n'. Theorem 2 is proved.

Theorem 2 allows us to compare the speed of convergence of $\left(\frac{A_{n+1}}{A_n} \right)^{B_n}$ towards e as n increases by choosing different sequences A_n and B_n. For example, let $A_n = n$, $B_n = n$, $n = 100$. This gives $\left(\frac{A_{n+1}}{A_n} \right)^{B_n} \simeq 2.7048$. If $A_n = n$, $B_n = n + \frac{1}{2}$, $n = 100$, then $\left(\frac{A_{n+1}}{A_n} \right)^{B_n} \simeq 2.7183$, which is a much better estimate. However, for these two examples, it can be seen that when n increases, the speeds of convergence in the two cases approach each other.

By changing A_n and B_n, we can further generalise Theorem 2. We take $A_{n+1} = A_n(1 + \varepsilon_n)$, where $\varepsilon_n \to 0$. Our previous assumptions of monotone increasing (decreasing) A_n now correspond to ε_n positive

https://doi.org/10.1017/mag.2024.80 Published online by Cambridge University Press

(negative). We have $B_n \sim \frac{1}{\varepsilon_n}$. Set r_n to be a positive sequence with $r_n \to 1$. Now, Theorem 2 is equivalent to

$$\lim_{n \to \infty} (1 + \varepsilon_n)^{\frac{r_n}{\varepsilon_n}} = e. \tag{1}$$

The sign of ε_n does not matter for this limit, so we can generalise the left-hand side of (1). For any constant k and δ_n a sequence with $|\delta_n|$ monotone decreasing to 0, we have

$$\lim_{n \to \infty} (1 + \varepsilon_n)^{\delta_n + k} = 1. \tag{2}$$

Multiplying (1) by (2) we obtain

$$\lim_{n \to \infty} (1 + \varepsilon_n)^{\frac{r_n}{\varepsilon_n} + \delta_n + k} = e. \tag{3}$$

This allows the reader to choose parameters to optimise convergence.

Acknowledgements

The authors would like to thank the Editor and the anonymous reviewer for their valuable suggestions.

References

1. A. J. Macintyre, Euler's limit for e^x and the exponential series, *Edinburgh Mathematical Notes* **37** (1949) pp. 26-28.
2. R. Farhadian, A generalization of Euler's limit, *Amer. Math. Monthly* **129** (2022) p. 384.

10.1017/mag.2024.80 © The Authors, 2024
Published by Cambridge University Press on behalf of The Mathematical Association

REZA FARHADIAN
Department of Statistics, Razi University, Kermanshah, Iran
e-mail: *farhadian.reza@yahoo.com*
VADIM PONOMARENKO
Department of Mathematics and Statistics, San Diego State University, San Diego, USA
e-mail: *vponomarenko@sdsu.edu*

108.28 π is a mean of 2 and 4

A series of *Mathematical Gazette* contributions, [1, 2, 3, 4], deals with limits of infinite sequences where the first n entries are specified and where latter entries correspond to a specified type of average of the n preceding entries. To the list of recursively defined averages may be added also the more well-known arithmetic-geometric mean, the arithmetic-harmonic mean and the geometric-harmonic mean. We are not aware of studies of recursions where some property of the index k dictates what average to

https://doi.org/10.1017/mag.2024.80 Published online by Cambridge University Press