CrossMark

*Animal* (2016), **10:12**, pp 1949–1954 © The Animal Consortium 2016
doi:10.1017/S1751731116000768

animal

# Technical note: use of internal transcribed spacer for ruminal yeast identification in dairy cows

E. Vargas-Bello-Pérez[1†], N. Cancino-Padilla[1] and J. Romero[2]

[1]*Departamento de Ciencias Animales, Facultad de Agronomía e Ingeniería Forestal, Pontificia Universidad Católica de Chile, Santiago, Casilla 306, C.P. 6904411, Chile;* [2]*Laboratorio de Biotecnología, Instituto de Nutrición y Tecnología de los Alimentos (INTA), Universidad de Chile, Santiago, C.P. 7810000, Chile*

(Received 20 November 2015; Accepted 14 March 2016; First published online 2 May 2016)

*Molecular techniques are important tools for microbiological studies in different habitats, and the internal transcribed spacer (ITS) has been proved to be useful for analyzing fungal diversity. The aim of this study was to use the ITS region to generate ruminal yeast profile and to identify ruminal yeast. DNA from ruminal digesta was extracted to amplify the ribosomal ITS region. The profile from the PCR products was visualized and the excised bands from the profile were identified as the genera* Millerozyma, Pichia, Rhizomucor *and* Hyphopichia. *Overall, the ITS resulted to be a simple, fast and sensitive approach that allowed profiling and identification of ruminal yeast that have not been previously described (*Millerozyma *and* Hyphopichia*) in the rumen microbial community.*

**Keywords:** rumen, yeast identification, dairy cow, internal transcribed spacer

## Implications

There is scarce information regarding the role, importance and diversity of yeast in the rumen of dairy cattle. The internal transcribed spacer (ITS) approach allows the profiling and identification of ruminal yeast in a simple, fast and sensitive way that could provide new insights into the diversity and ecology of many different groups of fungi. This approach could be an important tool in rumen microbiome studies.

## Introduction

The rumen allows the development and interactions of a wide variety of microorganisms (Mendes *et al.*, 2012), which are able to adapt to the rumen anaerobic conditions, temperature and pH (Fonty and Chaucheyras-Durand, 2006). In addition, the composition and proportion of ruminal microorganisms are influenced by external factors such as diet, feeding frequency, age (Denman and McSweeney, 2006) and have an important function in feed digestion (Oliveira *et al.*, 2014).

In the rumen inhabit yeasts, which are a phylogenetically diverse group of unicellular fungi, and belong to the phylum Ascomycota (de Barros Lopes *et al.*, 1998) and Basidiomycota (Osinska-Jaroszuk *et al.*, 2015); however, these microorganisms have received little attention from researchers working on ruminal microbiology.

† E-mail: evargasb@uc.cl

Clarke and Di Menna (1961) and Lund (1974) were pioneers in the study of these microorganisms in bovine through traditional techniques. Traditional methods for yeast identification are based on cellular morphology and reactions on fermentation protocols (Kurtzman and Robnett, 1998). These procedures are difficult and time-consuming (Villa-Carvajal *et al.*, 2006), and can be ambiguous due to strain variability of the microorganisms (Kurtzman and Robnett, 1998). For these reasons, identification of yeast species by phenotypic characteristics has been largely replaced by DNA-based methods (Kurtzman, 2006) because they are quick, sensitive and specific (Hayashi *et al.*, 2013). The ITS is a PCR approach validated to be useful for analysis of fungal diversity (Bellemain *et al.*, 2010). ITS is a non-coding ribosomal DNA (rDNA) spacer region located between the 18S and 28S ribosomal RNA (rRNA) genes (Sirohi *et al.*, 2013), and comprises ITS1/ITS2 intergenic sequences with highly conserved 5.8 rRNA in between (Zhang *et al.*, 2015). These regions were confirmed to be the most suitable marker genes for all classes of fungi providing complementary phylogenetic information (Koetschan *et al.*, 2014), and even useful for the detection and identification of fungal pathogens (Iwen *et al.*, 2005). Moreover, ITS is a DNA region with high variation among closely related species due to its fast rate of evolution (Schoch *et al.*, 2012) providing a molecular basis for determining phylogenetic relationships (Hayashi *et al.*, 2013).

Due to the scarce information regarding the role, importance and diversity of yeast in the rumen of dairy cattle, the aim of



https://doi.org/10.1017/S1751731116000768 Published online by Cambridge University Press

this study was to use the ITS region to generate ruminal yeast profile and to identify ruminal yeast in dairy cows.

## Material and methods

*Animals, diet and management*

The study was conducted at the Estación Experimental Pirque, Santiago, Chile (33°38'28''S, 70°34'27''W). Animals were handled following approved guidelines of the Animal Care and Use Committee of the Pontificia Universidad Católica de Chile. Three non-lactating, non-pregnant Holstein cows (785.5 ± 37.3 kg of BW; with rumen cannulae no. 3C; Bar Diamond Inc., Boise, ID, USA) were used. Cows were individually fed a total mixed ration (TMR) at a fixed rate once daily (0930 h). Animals were housed in individual stalls (2.4 × 6 m) and had continuous access to water. The diet was formulated according to the NRC (2001) with a 56 : 44 forage : concentrate ratio to meet the requirements of a dry cow in the last trimester of gestation consuming 10 kg dry matter (DM) daily. A mixer wagon was used to mix forage (17% of alfalfa hay and 18% of corn silage of DM) and concentrate (10% of high-moisture corn, 34% of soybean hulls and 19% of wheat bran of DM). Briefly, the chemical composition (g/kg DM) of the diet was as follows: 514 of DM, 173 of CP, 342 of NDF, 200 of ADF and 19 of lignin. Cows were fed the diet for 21 days.

*Samples*

On day 21, whole ruminal contents were hand-collected from the anterior, dorsal and mid-ventral regions of the rumen at 0900 h (2 h after feeding) and were squeezed through three layers of cheesecloth. A quantity of 10 ml of residual ruminal fluid was immediately used to determine pH (PP-201; GOnDO Electronic, Taipei, Taiwan), 10 ml were kept for $NH_3$–N analysis and another 10 ml were preserved for volatile fatty acid (VFA) determination by adding 1 ml of 25% metaphosphoric acid. Samples were frozen (−20°C) for later analysis. Samples of ruminal fluid were analyzed for VFA using a gas chromatograph (GC-2010; Shimadzu Scientific Instruments AOC-20s, Columbia, MD, USA) equipped with a 30 m wall-coated open-tubular fused-silica capillary column (Stabilwax-DA; 30 m × 0.32 mm i.d., 0.25 μm film thickness; Restek, Bellefonte, PA, USA). Oven temperature was programmed for 145°C for 2 min and then increased from 145°C to 220°C at 4°C/min. The injector and flame ionization detector were maintained at 250°C and 300°C, respectively. Following pH determination, the strained ruminal fluid was centrifuged for 10 min at 3000 × **g** at room temperature, discarding the supernatant and conserving the residue on ice – with later storage at 20°C – for subsequent DNA extraction.

*DNA extraction*

Before DNA extraction, samples of ruminal digesta were weighed (240 ± 12 μg) and deposited in 1.5 Eppendorf tubes. For preparation and homogenization, 300 μl of phosphate-buffered saline solution was added to the samples. Subsequently, tubes were incubated for 30 min at 37°C with lysozyme (1 μg/μl) and then for 30 min at 37°C with proteinase K (0.1 mg/ml). DNA was extracted by using the PowerSoil DNA Isolation Kit (MO BIO Laboratories Inc., Carlsbad, CA, USA) following the manufacturer's guidelines.

*PCR amplification of internal transcribed spacer region*

The ribosomal ITS region was amplified from genomic DNA by two consecutive PCR reactions, using primers ITF1 (5'-TCCGTAGGTGAACCTGCGG-3') and ITF4 (5'-TCCTCCGCTTATTGATATGC-3'). For the first amplification, the ITS region was amplified in 40 μl of reaction volume and the PCR conditions included 30 cycles, with initial denaturation at 94°C for 5 min, followed by denaturation at 94°C for 30 s, annealing at 55°C for 30 s and extension at 72°C for 1 min. A blank sample was used to ensure that there were no PCR artifacts. The amplified PCR products were separated by electrophoresis in 5% polyacrylamide gels in 1× TBE (tris-borate-EDTA) buffer at 150 V for 50 min and visualized by 30 min of incubation with SYBR Green at room temperature (Molecular Probes, Eugene, OR, USA). The band sizes were estimated by comparison against 100-bp DNA ladders and excised from the gel on a UV-transilluminator using a sterile scalpel as described by Trabal *et al.* (2012). The resulting slices were placed separately in 0.6 ml Eppendorf tube with 50 μl of nuclease-free water and incubated overnight at 4°C for elution of the DNA. Subsequently, 1 μl of each eluted band was used for re-amplification under the same conditions described above. To confirm the amplification, amplicons were visualized by electrophoresis in acrylamide gel under the same electrophoretic conditions previously described, and by silver staining method. Afterward, the amplicons were sent to Macrogen Inc. (Rockville, MD, USA) for sequencing service with both primers: ITF1 and ITF4.

*Sequence and phylogenetic analysis*

The obtained sequences were examined using the BLAST tool from the UNITE database (https://unite.ut.ee/) in order to find the sequence homology with the fungal sequences available in this database. Four UNITE reference sequences from phylum Ascomycota and Zygomycota were used in the phylogenetic analysis. These sequences included *Millerozyma* (SH201340.07FU), *Rhizomucor* (SH220342.07FU), *Hyphopichia* (SH203994.07FU) and *Pichia* (SH199823.07FU). The phylum Neocallimastigomycota reference sequence was used as an outgroup. The evolutionary distances were computed using the maximum likelihood method. The analysis involved 16 nucleotide sequences and all positions containing gaps were eliminated. Phylogenetic analysis was performed using the Molecular Evolutionary Genetics Analysis (MEGA6) software version 6.0.

## Results and discussion

*Ruminal fermentation parameters*

The ruminal microbial composition and environmental parameters may vary depending on the diet composition



https://doi.org/10.1017/S1751731116000768 Published online by Cambridge University Press

(Monteils *et al.*, 2011). In this study, ruminal fermentation parameters were measured to provide data on the rumen environment at the time when ruminal digesta was collected for molecular analysis. Those parameters averaged 6.9 ± 0.01 for pH, 13.3 ± 1.2 mg/dl for $NH_3$–N, 57.5 ± 6.3 mmol/l for total VFA, whereas molar proportions (mol/100 ml) for VFA were as follows: 63.8 ± 0.1 for acetate, 22.7 ± 0.2 for propionate, 11.2 ± 0.1 for butyrate and 2.4 ± 0.1 for valerate. The effect of the diet on the fermentation pattern is well known and the values obtained in the present study are similar to those reported for non-lactating cows fed corn-silage-based diets (Peyrat *et al.*, 2016) and a diet with a 50 : 50 forage : concentrate ratio based on corn silage, alfalfa hay and soybean meal (Shahmoradi *et al.*, 2016).

*PCR amplification of internal transcribed spacer region*

PCR amplification rendered product sizes between 460 and 750 bp (Figure 1). A total of 11 ITS sequences were obtained from those bands named in Figure 1. Ruminal yeast profile consisted mostly of four to six bands, hence potentially, four to six yeast genera. These profiles can be considered as equivalent to ruminal yeast microbiota profiles, as each ITS band may correspond to a particular yeast genus (Ide *et al.*, 2010).

An important part in this study is the PAGE adaptations that allowed an appropriate separation of the bands. Another comparative advantage of the approach used in this study is that we were able to obtain amplicons avoiding the use of purification and cloning kits. Consequently, this approach becomes an easy and economical alternative technique for ruminal yeast identification.

*Identification of ruminal yeast*

ITS bands were sequenced and the identification of the bands corresponded to the genera *Millerozyma*, *Rhizomucor*, *Hyphopichia* and *Pichia*. A more detailed information about the resulting sequence similarity searches against the records of the database UNITE are shown in Table 1. In this table, each ITS band is described based on the observed size of the band and the percentage of identity according to the UNITE RefSeq (sequences of the eluted amplicons).

In this study, the phylogenetic analysis was performed using MEGA6 software comparing the retrieved sequences with the RefSeq for each yeast genus. The resulting maximum likelihood tree (Figure 2) showed the identity of ITS bands for *Rhizomucor* and *Pichia* RefSeq, in contrast with a more diverse genetic composition observed for *Millerozyma* and *Hyphopichia*.

Fungi are the second largest kingdom of eukaryotic life and several DNA regions were evaluated as potential DNA barcodes for this important biological group. For yeasts, the D1/D2 region of large-subunit rDNA was adopted for characterizing species long before the concept of DNA barcoding was promoted (Schoch *et al.*, 2012). It had superior species resolution in some taxonomic groups, such as the early diverging lineages and the Ascomycete yeasts, but was otherwise slightly inferior to the ITS. Therefore, ITS has been suggested as the primary fungal barcode marker to the Consortium for the Barcode of Life (Schoch *et al.*, 2012).

In previous studies, researchers have reported the predominance of the genus *Pichia* in the rumen. Lund (1974) identified *Pichia kudriavzevii* (previously named *Candida krusei*) as the most frequent species in cows fed with different diets that included grass, grass silage, or a mix of beet and straw plus molasses and concentrates. Abrao *et al.* (2011) described *Pichia membranaefaciens* as the predominant ruminal yeast in goats through mycological cultures and micromorphological analysis. Similarly, Mendes *et al.* (2012) reported *P. kudriavzevii* as the most abundant yeast in isolates from ruminal samples from dairy cattle fed tropical forages. The genus *Pichia* was also isolated from the





**Figure 1** Electrophoretic separation profiles of internal transcribed spacer (ITS1) region of ruminal yeast from dairy cows fed a diet with a 56 : 44 forage : concentrate ratio. Lines 1 to 3 = ruminal fluid from cow 1; Lines 4 to 6 = ruminal fluid from cow 2; Lines 7 to 9 = ruminal fluid from cow 3; M = 100-bp DNA marker.



https://doi.org/10.1017/S1751731116000768 Published online by Cambridge University Press

**Table 1** *UNITE reference sequence results for ruminal yeast*

| Genus | Product size (bp) | Band | Identity (%) | RefSeq | GenBank |
|---|---|---|---|---|---|
| *Millerozyma* | 750 | B1.3; B3.2; B5.2; B7.2; B8.2; B9.2 | 99 to 1·00 | SH201340.07FU | EF568067.1 |
| *Rhizomucor* | 690 to 700 | B1.4; B7.3 | 100 | SH220342.07FU | JN206312.1 |
| *Hyphopichia* | 460 | B3.5 | 99 | SH203994.07FU | EU714323.1 |
| *Pichia* | 480 to 490 | B1.6; B7.5 | 99 to 100 | SH199823.07FU | DQ104710.1 |





**Figure 2** Phylogenetic tree constructed from internal transcribed spacer sequences of ruminal yeast from dairy cows and reference sequence from UNITE. The evolutionary history was inferred using the maximum likelihood method. Evolutionary analyses were performed using MEGA6 software version 6.0.

rumen of dairy cows (Sirisan and Pattarajinda, 2011; Sirisan *et al*., 2013) using a PCR approach to amplify the D1/D2 domain of 26S rDNA.

The knowledge of the function exerted by yeast supplementation in the rumen has been the focus of several studies, for example, Yuan *et al.* (2015) studied the effect of yeast supplementation on milk production, feeding behavior and metabolism in dairy cows. They proposed that yeast removed the oxygen in the rumen, promoting anaerobic microorganisms' development, and therefore enhance fiber digestion. The positive effects of yeast in the ruminal environment could also be due to the interaction of yeast with the rumen microorganisms, stimulating certain microbial population with its consequent effect (Sirisan *et al*., 2013; Marrero *et al*., 2015). Mendes *et al.* (2012) suggested that yeasts are able to degrade simple carbohydrates, which could favor the growth of yeast populations, and also contribute to the regulation of ruminal pH. Both live and dead yeast (*Saccharomyces*) are used as feed additives for dairy cows (Salvati *et al*., 2015); however, their precise role in improving rumen function is still unclear. Because yeast is involved in the oxygen utilization, fiber digestion and pH regulation (Salvati *et al*., 2015) in the rumen, data from this study are relevant for a wide range of scientists, animal nutritionists and enterprises. Despite the fact that the effects of dietary supplementation of ruminants with yeast has been extensively studied, there is little information on the identification, quantification and function of these fungi in the rumen. In this regard, a striking feature of this study was that we were able to identify *Millerozyma*, *Hyphopichia* and *Rhizomucor* (mold), which have not been previously described in the rumen microbial community.

The yeast found in this study may be present in ingested feed; however, we did not evaluate that. A recent publication (Henderson *et al*., 2015) reported that rumen microbial community composition varies with diet and host, but a core microbiome can be found across a wide geographical range. Thus, ruminal yeast could be influenced by both diet and host. Jensen *et al.* (1994) have established that the gastrointestinal tract is the main portal of entry for these fungi to the rumen. For example, *Pichia manshurica* was discovered in silage corn and non-fermented TMR under different air exposure, being predominant in low and high moisture levels (Carvalho *et al*., 2014; Hao *et al*., 2015). The species *Hyphopichia burtonii* have been isolated from corn, wheat and rice (Limtong *et al*., 2010) and also described as food spoilage organism (Groenewald and Smith, 2010). The presence of this yeast in different feedstuffs could explain the presence in the rumen through its ingestion from the diet.



https://doi.org/10.1017/S1751731116000768 Published online by Cambridge University Press

According to Lund's (1974) hypothesis, yeast could be introduced to the rumen from feed and water, more recently, Henderson *et al*. (2015) studied the microbial community composition of the ruminant and proposed that rumen microbes' differences are attributed to diet. This assumption can be important for future studies on rumen microbiology and consider the analysis of yeast profile in diets and feedstuffs to confirm this hypothesis.

Finally, there are some aspects that need to be considered when using the ITS approach, for example, the ITS results can be a preliminary step for the quantification of yeast species in rumen. In future studies, the use of quantitative PCR with the same primers used in this study could provide more accurate results if the objective is to determine the total concentration of ruminal yeast. In addition, findings from this study showed that the ITS approach is not accurate enough to discern among yeast species and therefore this molecular technique is more suitable to study genera. In addition, it is recommended to use reference sequences from GenBank RefSeq or UNITE repositories, which are databases with validated sequences to avoid misleading or incorrect sequences.

## Conclusion

In conclusion, the ITS approach used in this study allowed the identification of ruminal yeast in an easier, faster and economic procedure, and could provide new insights into the diversity and ecology of many different groups of fungi. In this study, the use of ITS allowed the identification of *Millerozyma* and *Hyphopichia* (phylum Ascomycota), which have not been previously described in ruminants.

## Acknowledgments

This study was funded by a research grant from FONDECYT 11121142 and 1140734 (Fondo Nacional de Desarrollo Científico y Tecnológico, Chile). The authors would like to thank Juanita Clavijo, Jorge Manzor and Pamela Alvarez for their technical assistance and Fundación Agro U.C. for the animal facilities and assistance in obtaining research data.

## References

Abrao FO, Freitas CES, Duarte ER, Geraseev LC, Barreto SMP, Medeiros AO and Rosa CA 2011. Leveduras no rúmen de caprinos e bovinos de corte criados em pastagem tropicais. Arquivo Brasileiro de Medicina Veterinária e Zootecnia 63, 526–529.

Bellemain E, Carlsen T, Brochmann C, Coissac E, Taberlet P and Kauserud H 2010. ITS as an environmental DNA barcode for fungi: an in silico approach reveals potential PCR biases. BMC Microbiology 10, 189.

Carvalho B, Ávila C, Miguel M, Pinto J, Santos M and Schwan R 2014. Aerobic stability of sugar-cane silage inoculated with tropical strains of lactic acid bacteria. Grass and Forage Science 70, 308–323.

Clarke R and Di Menna M 1961. Yeasts from the bovine rumen. Journal of General Microbiology 25, 113–117.

de Barros Lopes M, Soden A, Martens AL, Henschke PA and Langridge P 1998. Differentiation and species identification of yeasts using PCR. International Journal of Systematic Bacteriology 48, 279–286.

Denman SE and McSweeney CS 2006. Development of a real-time PCR assay for monitoring anaerobic fungal and cellulolytic bacterial populations within the rumen. FEMS Microbiology Ecology 58, 572–582.

Fonty G and Chaucheyras-Durand F 2006. Effects and modes of action of live yeasts in the rumen. Biologia, Bratislava 61, 741–750.

Groenewald M and Smith M 2010. Re-examination of strains formerly assigned to *Hyphopichia burtonii*, the phylogeny of the genus *Hyphopichia*, and the description of *Hyphopichia pseudoburtonii* sp. nov. International Journal of Systematic and Evolutionary Microbiology 60, 2675–2680.

Hao W, Wang H, Ning T, Yang F and Xu C 2015. Aerobic stability and effects of yeasts during deterioration of non-fermented and fermented total mixed ration with different moisture levels. Asian-Australasian Journal of Animal Sciences 28, 816–826.

Hayashi T, Sugita T, Hata E, Katsuda K, Zhang E, Kihu Y, Sugawara K, Ozawa T, Matsubara T, Ando T, Obayashi T, Itoh T, Yabusaki T, Kudo K, Yamamoto H, Koiwa M, Oshida T, Tagawa Y and Kawai K 2013. Molecular-based identification of yeasts isolated from bovine clinical mastitis in Japan. The Journal of Veterinary Medical Science 75, 387–390.

Henderson G, Cox F, Ganesh S, Jonker A and Young W 2015. Rumen microbial community composition varies with diet and host, but a core microbiome is found across a wide geographical range. Scientific Reports 5, 14567.

Ide S, Miyazaki T, Maki H and Kobayashi T 2010. Abundance of ribosomal RNA gene copies maintains genome integrity. Science 327, 693–696.

Iwen PC, Freifeld AG, Sigler L and Tarantolo SR 2005. Molecular identification of *Rhizomucor pusillus* as a cause of sinus-orbital zygomycosis in a patient with acute myelogenous leukemia. Journal of Clinical Microbiology 43, 5819–5821.

Jensen H, Olsen S and Aalbaek B 1994. Gastrointestinal aspergillosis and zygomycosis of cattle. Veterinary Pathology 31, 28–36.

Koetschan C, Kittelmann S, Lu J, Al-Halbouni D, Jarvis G, Müller T, Wolf M and Janssen P 2014. Internal transcribed spacer 1 secondary structure analysis reveals a common core throughout the anaerobic fungi (Neocallimastigomycota). PLOS ONE 9, e91928.

Kurtzman CP 2006. Yeast species recognition from gene sequence analyses and other molecular methods. Mycoscience 47, 65–71.

Kurtzman CP and Robnett CJ 1998. Identification and phylogeny of ascomycetous yeasts from analysis of nuclear large subunit (26S) ribosomal DNA partial sequences. Antonie Van Leeuwenhoek 73, 331–371.

Limtong S, Kaewwichian R, Am-In S, Boonmak C, Jindamorakot S, Yongmanitchai W, Srisuk N, Kawasaki H and Nakase T 2010. Three anamorphic yeast species *Candida sanitii* sp. nov., *Candida sekii* sp. nov. and *Candida suwanaritii*, three novel yeasts in the Saturnispora clade isolated in Thailand. FEMS Yeast Research 10, 114–122.

Lund A 1974. Yeasts and moulds in the bovine rumen. Journal of General Microbiology 81, 453–462.

Marrero Y, Castillo Y, Ruiz O, Burrola E and Angulo C 2015. Feeding of yeast (*Candida* spp.) improves in vitro ruminal fermentation of fibrous substrates. Journal of Integrative Agriculture 14, 514–519.

Mendes P, Robson E, Flávia O, Cláudio S, Castro L and Rosa C 2012. Aerobic fungi in the rumen fluid from dairy cattle fed different sources of forage. Revista Brasileira de Zootecnia 41, 2336–2342.

Monteils V, Rey M, Cauquil L, Troegeler-Meynadier A, Silberberg M and Combes S. 2011. Random changes in the heifer rumen in bacterial community structure, physico-chemical and fermentation parameters, and in vitro fiber degradation. Livestock Science 141, 104–112.

NRC 2001. Nutrient requirements of dairy cattle, 7th revised edition. National Academy of Sciences, Washington, DC, USA.

Oliveira F, Robson E, Silva C, Alves E, Castro L, Ferreira A, Rosa C and Rodrigues N 2014. Characterization of fungi from ruminal fluid of beef cattle with different ages and raised in tropical lignified pastures. Current Microbiology 69, 649–659.

Osinska-Jaroszuk M, Jarosz-Wilkołazka A, Jaroszuk-Sciseł J, Szałapata K, Nowak A, Jaszek M, Ozimek E and Majewska M 2015. Extracellular polysaccharides from Ascomycota and Basidiomycota: production conditions, biochemical characteristics, and biological properties. World Journal of Microbiology and Biotechnology 31, 1823–1844.

Peyrat J, Baumont R, Le Morvan A and Nozière P 2016. Effect of maturity and hybrid on ruminal and intestinal digestion of corn silage in dry cows. Journal of Dairy Science 99, 258–268.



https://doi.org/10.1017/S1751731116000768 Published online by Cambridge University Press

Salvati GGS, Morais Júnior NN, Melo ACS, Vilela RR, Cardoso FF, Aronovich M, Pereira RAN and Pereira MN 2015. Response of lactating cows to live yeast supplementation during summer. Journal of Dairy Science 98, 4062–4073.

Schoch CL, Seifert KA, Huhndorf S, Robert V, Spouge JL, Levesque CA and Chen W, Fungal Barcoding Consortium 2012. Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for fungi. Proceedings of the National Academy of Sciences of the United States of America 109, 6241–6246.

Shahmoradi A, Alikhani M and Riasi A 2016. Effects of partial replacement of barley grain with beet pulp on performance, ruminal fermentation and plasma concentration of metabolites in transition dairy cows. Journal of Animal Physiology and Animal Nutrition 100, 178–188.

Sirisan V and Pattarajinda V 2011. Screening yeasts from ruminal fluid of dairy heifer fed a different ratio roughage to concentrate diets. Journal of Agricultural Science and Technology A 1, 1155–1158.

Sirisan V, Pattarajinda V, Vichitphan K and Leesing R 2013. Isolation, identification and growth determination of lactic acid-utilizing yeasts from the ruminal fluid of dairy cattle. Letters in Applied Microbiology 57, 102–107.

Sirohi S, Choudhury P, Puniya A, Singh D, Dagar S and Singh N 2013. Ribosomal ITS1 sequence-based diversity analysis of anaerobic rumen fungi in cattle fed on high fiber diet. Annals of Microbiology 63, 1571–1577.

Trabal N, Mazón-Suástegui JM, Vázquez-Juárez R, Asencio-Valle F, Morales-Bojórquez E and Romero J 2012. Molecular analysis of bacterial microbiota associated with oysters (*Crassostrea gigas* and *Crassostrea corteziensis*) in different growth phases at two cultivation sites. Microbial Ecology 64, 555–569.

Villa-Carvajal M, Querol A and Belloch C 2006. Identification of species in the genus *Pichia* by restriction of the internal transcribed spacers (ITS1 and ITS2) and the 5.8S ribosomal DNA gene. Antonie Van Leeuwenhoek 90, 171–181.

Yuan K, Liang T, Muckey M, Mendonça L, Hulbert L, Elrod C and Bradford B 2015. Yeast product supplementation modulated feeding behavior and metabolism in transition dairy cows. Journal of Dairy Science 98, 532–540.

Zhang W, Yuan Y, Yang S, Huang J and Huang L 2015. ITS2 secondary structure improves discrimination between medicinal 'Mu Tong' species when using DNA barcoding. PLOS ONE 10, e0131185.



https://doi.org/10.1017/S1751731116000768 Published online by Cambridge University Press