Econometric Theory, **20**, 2004, 1261–1264. Printed in the United States of America.
DOI: 10.1017/S0266466604206089

PROBLEMS AND SOLUTIONS

SOLUTIONS

03.6.1 The Central Limit Theorem for Student's Distribution—Solution

Karim Abadir
University of York, UK

Jan Magnus (the poser of the problem)
Tilburg University, The Netherlands

Consider the Lindeberg–Feller central limit theorem (CLT), which we state as follows. Let $\{x_n\}$ be a sequence of independent random variables with means $\{\mu_n\}$ and nonzero variances $\{\sigma_n^2\}$ (both existing), and c.d.f.s $\{F_n\}$. Define $\lambda_n > 0$ by $\lambda_n^2 = \sum_{i=1}^n \sigma_i^2$. Then, *Lindeberg's condition*

$$\lim_{n\to\infty} \sum_{i=1}^{n} \int_{|u-\mu_i|\geq\lambda_n\epsilon} \left(\frac{u-\mu_i}{\lambda_n}\right)^2 \mathrm{d}F_i(u) = 0, \qquad \forall\epsilon > 0,$$

is equivalent to

$$z_n := \frac{\sum_{i=1}^{n}(x_i-\mu_i)}{\lambda_n} \overset{a}{\sim} \mathrm{N}(0,1) \quad \text{and} \quad \lim_{n\to\infty} \max_{1\leq i\leq n} \Pr\left(\frac{|x_i-\mu_i|}{\lambda_n} \geq \epsilon\right) = 0,$$

where the latter limit is known as the *uniform asymptotic negligibility* (u.a.n.) condition. One can usually interpret λ_n^2 as the variance of the numerator of z_n. We shall see, however, that there are cases where asymptotic normality holds in spite of $\{x_n\}$ having infinite variances.

Let $\{x_n\}$ be a random sample from Student's $\mathrm{t}(\nu)$. For $\nu < 2$, no λ_n exists that can lead to $z_n := \sum_{i=1}^n x_i/\lambda_n \overset{a}{\sim} \mathrm{N}(0,1)$. This is because the tails of the density of $\mathrm{t}(\nu)$ decay at a rate of $u^{-\nu-1}$ and the stable limit theorem tells us that a nonnormal stable law arises if the tails of the p.d.f. of x_i decay at a rate of u^{-a} where $a < 3$; e.g., see Loève (1977, §25) or Hoffmann-Jørgensen (1994, §5.25). For example, for $\nu = 1$, the average of standard Cauchy variates is standard Cauchy too, so that there exists no λ_n achieving asymptotic normality of z_n.

For $\nu > 2$, both the mean and the variance exist, and the Lindeberg–Feller CLT applies, with $\lambda_n^2 = n\,\mathrm{var}(x_i)$. The interesting part is $\nu = 2$, where we will show that asymptotic normality of z_n holds, in spite of $\mathrm{var}(x_i)$ being infinite, and we will derive the appropriate λ_n. We will require the additional assump-

© 2004 Cambridge University Press 0266-4666/04 $12.00

https://doi.org/10.1017/S0266466604216085 Published online by Cambridge University Press

tion that $\lambda_n^2 \to \infty$ as $n \to \infty$. In the standard CLT, this assumption was unnecessary, as it followed from $\lambda_n^2 = n\,\mathrm{var}(x_i)$. We will see subsequently that λ_n^2 can be interpreted in terms of truncated variances for $\nu = 2$.

To prove the asymptotic normality of z_n, we need to show that the characteristic function $\varphi(t) := \mathrm{E}(\exp(\mathrm{i}tx_i))$ satisfies

$$\lim_{n\to\infty} n \log \varphi\left(\frac{t}{\lambda_n}\right) = -\frac{t^2}{2} \tag{1}$$

for some choice of λ_n, with $\lambda_n \to \infty$ as $n \to \infty$. Because the sequence $\{x_n\}$ is i.i.d., the uniform asymptotic negligibility condition

$$\lim_{n\to\infty} \max_{1\le i\le n} \Pr\left(\frac{|x_i|}{\lambda_n} \ge \epsilon\right) = 0$$

is satisfied for all $\epsilon > 0$, thus implying

$$\lim_{n\to\infty} \left|\varphi\left(\frac{t}{\lambda_n}\right) - 1\right| = 0.$$

This allows us to take the leading term of the logarithmic expansion of the left-hand side of (1) as

$$\begin{aligned}\lim_{n\to\infty} n \log \varphi\left(\frac{t}{\lambda_n}\right) &= \lim_{n\to\infty} n\left(\varphi\left(\frac{t}{\lambda_n}\right) - 1\right) \\ &= -\frac{t^2}{2} \lim_{n\to\infty} n \int_{|u|<\lambda_n\epsilon} \frac{u^2}{\lambda_n^2}\,\mathrm{d}F(u), \quad \lambda_n \to \infty,\end{aligned}$$

where the linear term in t drops out because the sequence $\{x_n\}$ is centered around zero. Asymptotic standard-normality obtains if we can find the appropriate λ_n^2 that makes the latter limit equal to 1 for all $\epsilon > 0$. Notice that this limit is the complement of Lindeberg's condition, where $\sum_{i=1}^n$ is replaced by n because $\{x_n\}$ is an i.i.d. sequence.

From Student's t(2) density,

$$\begin{aligned}\int_{-c\sqrt{2}}^{c\sqrt{2}} \frac{u^2}{\sqrt{8}\left(1 + \frac{u^2}{2}\right)^{3/2}}\,\mathrm{d}u &= 2\log(\sqrt{1+c^2} + c) - \frac{2c}{\sqrt{1+c^2}} \\ &= 2\sinh^{-1}(c) - \frac{2c}{\sqrt{1+c^2}}\end{aligned}$$

https://doi.org/10.1017/S0266466604216085 Published online by Cambridge University Press

tends to infinity as $c \to \infty$. We need to solve

$$1 = \lim_{n\to\infty} \frac{n}{\lambda_n^2} \int_{|u|<\lambda_n \epsilon} u^2 \mathrm{d}F(u) = \lim_{n\to\infty} \frac{2n \sinh^{-1}(\lambda_n \epsilon/\sqrt{2})}{\lambda_n^2}$$

where we have dropped $2c/\sqrt{1+c^2} \to 2$ that is dominated by $\sinh^{-1}(c) \to \infty$. By using the logarithmic representation of the latter and simplifying,

$$1 = \lim_{n\to\infty} \frac{2n \log(\lambda_n)}{\lambda_n^2}$$

is solved by $\lambda_n = \sqrt{n \log(n)}$ or any other function that is asymptotically equivalent to it (such as $\sqrt{n \log(n)} + \sqrt{n}$). Therefore,

$$z_n = \frac{1}{\sqrt{n \log(n)}} \sum_{i=1}^{n} x_i \overset{a}{\sim} \mathrm{N}(0,1).$$

REFERENCES

Hoffmann-Jørgensen, J. (1994) *Probability with a View toward Statistics*, vol. I. Chapman and Hall.
Loève, M. (1977) *Probability Theory I*, 4th ed. Springer-Verlag.

03.6.2. Unbiasedness of the OLS Estimator with Random Regressors—Solution

Michael Jansson (the poser of the problem)
University of California, Berkeley

(a) Suppose $n = 1$ and let X and u be independent standard normal variates. Then X is nonzero with probability one and $E(u|X) \overset{\text{a.s.}}{=} 0$, but $E|\hat{\beta}| = \infty$ because the distribution of $\hat{\beta} - \beta = u/X$ is Cauchy.

(b) The matrix X has full column rank with probability one if and only if

$$\Pr[\lambda_{\min}(X'X) > 0] = 1, \tag{1}$$

where $\lambda_{\min}(\cdot)$ denotes the minimal eigenvalue of the argument.

$E(\hat{\beta})$ exists if (and only if) $E|c'(\hat{\beta} - \beta)| < \infty$ for any k-vector c with $c'c = 1$. In the sequel, let c be an arbitrary k-vector with unit length. Now,

$$|c'(\hat{\beta} - \beta)| = |c'(X'X)^{-1}X'u| \leq \sqrt{c'(X'X)^{-1}c}\sqrt{u'u} \leq \sqrt{\lambda_{\min}^{-1}(X'X)}\sqrt{u'u},$$

where the first inequality uses the Cauchy–Schwarz inequality and the second inequality uses Magnus and Neudecker (1988), Theorem 11.4.

If X and u are independent, $E(u|X) \overset{\text{a.s.}}{=} 0$, and

$$E[\lambda_{\min}^{-1/2}(X'X)] < \infty, \tag{3}$$

https://doi.org/10.1017/S0266466604216085 Published online by Cambridge University Press

then

$$E[\sqrt{\lambda_{\min}^{-1}(X'X)}\sqrt{u'u}] = E[\sqrt{\lambda_{\min}^{-1}(X'X)}]E[\sqrt{u'u}]$$
$$\leq E[\lambda_{\min}^{-1/2}(X'X)]E[\|u\|_1] < \infty,$$

where $\|\cdot\|_1$ denotes the l^1 norm (i.e., $\|u\|_1 = \sum_{i=1}^n |u_i|$ for any $u = (u_1,\ldots,u_n)' \in \mathbb{R}^n$), the first inequality uses Jensen's inequality, and the last inequality uses the fact that existence of $E(u|X)$ requires $E(\|u\|_1) < \infty$.

If $E(u'u) < \infty$, $E(u|X) \overset{a.s.}{=} 0$, and

$$E[\lambda_{\min}^{-1}(X'X)] < \infty, \tag{3}$$

then

$$E|c'(\hat{\beta} - \beta)| \leq E[\sqrt{\lambda_{\min}^{-1}(X'X)}\sqrt{u'u}] \leq \sqrt{E[\lambda_{\min}^{-1}(X'X)]}\sqrt{E(u'u)} < \infty,$$

where the second inequality uses the Cauchy–Schwarz inequality.

If $E(u|X) \overset{a.s.}{=} 0$ and

$$\Pr[\lambda_{\min}(X'X) > \delta] = 1 \quad \text{for some } \delta > 0, \tag{4}$$

then

$$E|c'(\hat{\beta} - \beta)| \leq E[\sqrt{\lambda_{\min}^{-1}(X'X)}\sqrt{u'u}] \leq \delta^{-1/2}E(\|u\|_1) < \infty,$$

where the second inequality uses $u'u \leq \|u\|_1^2$ and the last inequality uses the fact that existence of $E(u|X)$ requires $E(\|u\|_1) < \infty$. (This result contains the standard fixed-regressor result as a special case because $\Pr[\lambda_{\min}(X'X) > \delta] = 1$ for some $\delta > 0$ if X is nonrandom and of full column rank.)

Remark. The following chain of implications holds: (4) $\Rightarrow$ (3) $\Rightarrow$ (2) $\Rightarrow$ (1).

Indeed, (4) $\Rightarrow$ (3) because $E[\lambda_{\min}^{-1}(X'X)] < \delta^{-1}$ if $\Pr[\lambda_{\min}(X'X) > \delta] = 1$. Moreover, (3) $\Rightarrow$ (2) because $E[\lambda_{\min}^{-1/2}(X'X)] \leq \sqrt{E[\lambda_{\min}^{-1}(X'X)]}$ by the Cauchy–Schwarz inequality. Finally, (2) $\Rightarrow$ (1) because $E[\lambda_{\min}^{-1/2}(X'X)] = \infty$ whenever $\Pr[\lambda_{\min}(X'X) = 0] > 0$.

REFERENCE

Magnus, J.R. & H. Neudecker (1988) *Matrix Differential Calculus with Applications in Statistics and Econometrics*. Wiley.

https://doi.org/10.1017/S0266466604216085 Published online by Cambridge University Press