Canad. J. Math. Vol. 77 (6), 2025 pp. 2148–2170
http://dx.doi.org/10.4153/S0008414X24000683
© The Author(s), 2024. Published by Cambridge University Press on behalf of Canadian Mathematical Society

Some inequalities between $M(a, b, c; L; n)$ and the partition function $p(n)$

Bing He, Linpei Li, and Jian Cao

Abstract. Let $p(n)$ and $M(m, L; n)$ be the number of partitions of n and the number of partitions of n with crank congruent to m modulo L, respectively, and let

$$M(a, b, c; L; n) := M(a, L; n) + M(b, L; n) + M(c, L; n).$$

In this paper, we focus on some relations between $M(m, L; n)$ and $p(n)$, which Dyson, Andrews, and Garvan etc. have studied previously. By applying a modification of the circle method to estimate the Fourier coefficients of generating functions, we establish the following inequalities between $M(a, b, c; L; n)$ and $p(n)$: for $n \geq 467$,

$$M(0, 1, 1; 9; n) > \frac{p(n)}{3} \quad \text{when} \quad n \equiv 0, 1, 5, 8 \pmod 9,$$
$$M(0, 1, 1; 9; n) < \frac{p(n)}{3} \quad \text{when} \quad n \equiv 2, 3, 4, 6, 7 \pmod 9,$$
$$M(2, 3, 4; 9; n) < \frac{p(n)}{3} \quad \text{when} \quad n \equiv 0, 1, 5, 8 \pmod 9,$$
$$M(2, 3, 4; 9; n) > \frac{p(n)}{3} \quad \text{when} \quad n \equiv 2, 3, 4, 6, 7 \pmod 9.$$

In the proof of these inequalities, an inequality for the logarithm of the generating function for $p(n)$ is derived and applied. Our method reduces the last possible counterexamples to $467 \leq n \leq 22471$, and it will produce more effective estimates when proving inequalities of such types.

1 Introduction

A partition of a positive integer n is a nonincreasing sequence of positive integers whose sum equals n. Typically, $p(n)$ is used to denote the number of partitions of n. The generating function for $p(n)$ is as follows:

$$\sum_{n=1}^{\infty} p(n)q^n = \frac{1}{(q; q)_\infty}.$$

Here and in what follows, $|q| < 1$, and

$$(a; q)_\infty := \prod_{i=0}^{\infty}(1 - aq^i).$$

Received by the editors January 24, 2024; revised June 23, 2024; accepted July 2, 2024.
AMS subject classification: **11P55**, **11P82**, 11P83.
Key words: Inequality, partition, crank, $M(a, b, c; L; n)$, $p(n)$, circle method.

https://doi.org/10.4153/S0008414X24000683 Published online by Cambridge University Press

In 1944, Dyson [12] defined the rank of a partition to be the largest part minus the number of parts. Let $N(m;n)$ be the number of partitions of n with rank m and $N(m,L;n)$ the number of partitions of n with rank congruent to m modulo L. Dyson conjectured that

$$N(a,5;5n+4) = \frac{p(5n+4)}{5},$$
$$N(a,7;7n+5) = \frac{p(5n+4)}{7},$$

which provide combinatorial proofs for the cases of modulo 5 and 7 in Ramanujan's congruences. In addition, Dyson conjectured that there exists a crank function for partitions which would supply a combinatorial proof of Ramanujan's congruences modulo 11. In 1954, Atkin and Swinnerton-Dyer [5] proved the above identities of Dyson by applying the generating function for $N(m,L;n)$.

Forty years later, Andrews and Garvan [3] defined the crank function for partitions. Let $M(m;n)$ be the number of partitions of n with crank m and let $M(m,L;n)$ be the number of partitions of n with crank congruent to m modulo L. Andrews and Garvan proved that

$$M(m,11;11n+6) = \frac{p(11n+6)}{11}, \quad 0 \le m \le 10,$$

which provides a combinatorial proof for Ramanujan's congruences modulo 11. Recently, many inequalities for $M(m,L;n)$ and $N(m,L;n)$ modulo 11 were deduced by Borozenets [7] and Bringmann and Pandey [8]. For inequalities between the rank counts $N(m,L;n)$ or between the crank counts $M(m,L;n)$, see, for example, [1, 10, 11, 16]. For distributions of rank and crank statistics for integer partitions, see, for example, [20, 21].

For convenience, we adopt the following notations:

$$N(a,b,c;L;n) := N(a,L;n) + N(b,L;n) + N(c,L;n),$$
$$M(a,b,c;L;n) := M(a,L;n) + M(b,L;n) + M(c,L;n).$$

In [15, Theorem 4.1], Kang showed the following relationship between rank and crank:

$$\sum_{m=-\infty}^{\infty}\sum_{n=0}^{\infty}(N(3m-1;n)+N(3m;n)+N(3m+1;n))z^m q^n = \frac{(q^3;q^3)_\infty}{(q;q)_\infty}\sum_{m=-\infty}^{\infty}\sum_{n=0}^{\infty}M(m;n)z^m q^{3n}.$$

Inspired by this relationship, Aygin and Chan [6] provided a series of generating functions for $M(a,b,c;L;n)$. For $L \in \{6,9,12\}$, Aygin and Chan found generating functions of

$$N(3j-1,3j,3j+1;L;n), \quad \text{for } 0 \le j \le L/3-1,$$
$$M(3j-1,3j,3j+1;L;n), \quad \text{for } 0 \le j \le L/3-1,$$
$$M(3j-2,3j-1,3j;L;n), \quad \text{for } 0 \le j \le L/3-1$$

https://doi.org/10.4153/S0008414X24000683 Published online by Cambridge University Press

and used the periodicity of the sign of the Fourier coefficients of these generating functions to prove many inequalities between $M(a,b,c;L;n)$ and the partition function $p(n)$. At the end of the paper, they proposed 18 inequalities of such type as a conjecture. In [13], Fan, Xia, and Zhao established some generating functions for $N(a,12;n)$ and $M(a,12;n)$ with $0 \le a \le 11$, and used them to confirm the first 6 inequalities in the conjecture of Aygin and Chan. Later, Yao [19] proved that the remaining 12 inequalities hold for sufficiently large n. Simultaneously, at the end of their paper, Aygin and Chan said that "Additionally, the referee pointed out that the signs of theta parts of $M(0,1,1;9;n)$, $M(2,3,4;9;n)$, $M(0,1,2;9;n)$, $M(3,4,5;9;n)$, and $M(6,7,8;9;n)$ are periodic modulo 9 when $n \ge 467$. Thus, conjectures similar to Conjecture 8.1 can be stated for $M(0,1,1;9;n)$, $M(2,3,4;9;n)$, $M(0,1,2;9;n)$, $M(3,4,5;9;n)$ and $M(6,7,8;9;n)$." Up to now, as far as we are concerned, no inequality for $M(0,1,1;9;n)$, $M(2,3,4;9;n)$, $M(0,1,2;9;n)$, $M(3,4,5;9;n)$ and $M(6,7,8;9;n)$, which is similar to [6, Conjecture 8.1], has been proposed explicitly and proved.

In this paper, we state some inequalities for $M(0,1,1;9;n)$ and $M(2,3,4;9;n)$ and prove them.

Theorem 1.1 *For $n \ge 467$, we have*

$$M(0,1,1;9;n) > \frac{p(n)}{3} \quad \text{when} \quad n \equiv 0,1,5,8 \pmod 9,$$
$$M(0,1,1;9;n) < \frac{p(n)}{3} \quad \text{when} \quad n \equiv 2,3,4,6,7 \pmod 9,$$
$$M(2,3,4;9;n) < \frac{p(n)}{3} \quad \text{when} \quad n \equiv 0,1,5,8 \pmod 9,$$
$$M(2,3,4;9;n) > \frac{p(n)}{3} \quad \text{when} \quad n \equiv 2,3,4,6,7 \pmod 9.$$

We will employ the modification [14] of the circle method[1] to estimate the Fourier coefficients of generating functions for $M(0,1,1;9;n)$ and $M(2,3,4;9;n)$, obtaining information about the sign of the coefficients of $g(q) - 3h(q)$ (see (3.1) and (3.2) for the definitions of $g(q)$ and $h(q)$), and thereby prove the inequalities in Theorem 1.1. An inequality for the logarithm of the generating function for $p(n)$ will be derived and used to reduce the last possible counterexamples to $467 \le n \le 22471$. This method will produce more effective estimates when proving such types of inequalities.

2 Preliminaries

For $f(s) = \sum_{n=0}^{\infty} \alpha(n) s^n$, we apply the residue theorem to the generating function to get

$$\alpha(n) = \frac{1}{2\pi i} \oint_{|s|=r} \frac{f(s)}{s^{n+1}} ds$$

with r smaller than the convergence radius.

[1] See [17] and [18] for details of the circle method.

https://doi.org/10.4153/S0008414X24000683 Published online by Cambridge University Press

2.1 Splitting integral intervals with Farey fractions

We set $r := e^{-2\pi\rho}$ with $\rho = 1/N^2$ for some $N > 0$. Let h/k be a Farey fraction of order N, and let $\xi_{h,k} := [-\theta'_{h,k}, \theta''_{h,k}]$ with $\theta'_{h,k}, \theta''_{h,k}$ being the distance from h/k to its neighboring mediants. If we set

$$s := re^{2\pi i\theta} = e^{-2\pi\rho}e^{2\pi i\theta},$$

and let $\theta := h/k + \varphi$ on each $\xi_{h,k}$, then

$$ds = 2\pi i e^{-2\pi\rho}e^{2\pi ih/k}e^{2\pi i\varphi}d\varphi,$$

so that

$$\alpha(n) = \sum_{\substack{0\le h<k\le N\\(h,k)=1}} e^{-2\pi inh/k}\int_{\xi_{h,k}} f(e^{2\pi i(h/k+i\rho+\varphi)})e^{2\pi n\rho}e^{-2\pi in\varphi}d\varphi.$$

Set $z := k(\rho - i\varphi)$ and $\tau := (h + iz)/k$ to get

$$\begin{aligned}\alpha(n) &= \sum_{\substack{0\le h<k\le N\\(h,k)=1}} e^{-2\pi inh/k}\int_{\xi_{h,k}} f(e^{2\pi i(h+iz)/k})e^{2\pi n\rho}e^{-2\pi in\varphi}d\varphi \\ &= \sum_{\substack{0\le h<k\le N\\(h,k)=1}} e^{-2\pi inh/k}\int_{\xi_{h,k}} f(e^{2\pi i\tau})e^{2\pi n\rho}e^{-2\pi in\varphi}d\varphi.\end{aligned} \tag{2.1}$$

2.2 Dedekind's eta-function

The Dedekind eta-function is defined by

$$\eta(\tau) := q^{1/24}(q;q)_\infty,$$

where $q = e^{2\pi i\tau}$ with $\mathrm{Im}(\tau) > 0$. A well-known transformation formula for the Dedekind eta-function is as follows [4, p. 52].

For $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma := \mathrm{SL}_2(\mathbb{Z})$, $c > 0$, we have [4, Theorem 3.4]

$$\eta(\gamma\tau) = e^{-\pi is(d,c)}e^{\pi i(a+d)/(12c)}\sqrt{-i(c\tau+d)}\eta(\tau), \tag{2.2}$$

where $s(d,c)$ is the Dedekind sum given by

$$s(d,c) := \sum_{n=0}^{c-1}\frac{n}{c}\left(\frac{dn}{c} - \left\lfloor\frac{dn}{c}\right\rfloor - \frac{1}{2}\right).$$

Set

$$F(e^{2\pi i\tau}) := \frac{1}{(q;q)_\infty} = \frac{e^{\pi i\tau/12}}{\eta(\tau)}.$$

It follows from (2.2) that

$$F(e^{2\pi i\tau}) = e^{\pi i(\tau-\gamma\tau)/12}e^{-\pi is(d,c)}e^{\pi i(a+d)/(12c)}\sqrt{-i(c\tau+d)}F(e^{2\pi i\gamma\tau}).$$

https://doi.org/10.4153/S0008414X24000683 Published online by Cambridge University Press

For the interval $\xi_{h,k}$, if $\gcd(n,k)=1$, then we choose an integer h'_n such that $nhh'_n \equiv -1 \pmod{k}$. Hence, there exists $b_n \in \mathbb{Z}$ such that $nhh'_n - b_n k = -1$. Set

$$\gamma_{(nh,k)} := \begin{pmatrix} h'_n & -b_n \\ k & -nh \end{pmatrix} \in \Gamma.$$

Then,

$$\gamma_{(nh,k)} = \frac{h'_n \frac{nh+inz}{k} - b_n}{k\frac{nh+inz}{k} - nh} = \frac{h'_n nh - b_n k + inh'_n z}{kinz} = \frac{h'_n}{k} + i\frac{1}{knz},$$

so that

$$\mathrm{Im}(\gamma_{(nh,k)}) = \mathrm{Re}\left(\frac{1}{knz}\right) = \frac{1}{kn}\mathrm{Re}\left(\frac{1}{z}\right), \tag{2.3}$$

and

$$F(e^{2\pi i n\tau}) = e^{\frac{\pi}{12k}\left(\frac{1}{nz}-nz\right)} e^{\pi i s(nh,k)} \sqrt{nz}\, F\left(e^{2\pi i \gamma_{(nh,k)}(n\tau)}\right). \tag{2.4}$$

2.3 Some bounds and an integral

It is obvious to observe that

$$\frac{1}{2kN} \le \theta'_{h,k}, \theta''_{h,k} \le \frac{1}{kN}.$$

From this, we have

$$\frac{1}{kN} \le |\xi_{h,k}| \le \frac{2}{kN},$$

and

$$|\varphi| \le (kN)^{-1}.$$

Notice that $z = k(\rho - i\varphi)$. Then,

$$\mathrm{Re}\left(\frac{1}{z}\right) = \mathrm{Re}\left(\frac{1}{k(\rho - i\varphi)}\right) = \frac{1}{k}\frac{\rho}{\rho^2+\varphi^2} \ge \frac{1}{k}\frac{N^{-2}}{N^{-4}+k^{-2}N^{-2}} \ge \frac{k}{2}. \tag{2.5}$$

Also, we have

$$|z|^{-1/2} = \frac{1}{k^{1/2}}\frac{1}{(\rho^2+\varphi^2)^{1/4}} \le \frac{1}{(k^2\rho^2)^{1/4}} = k^{-1/2}N. \tag{2.6}$$

We now estimate the function $\log F(x)$ for $0 < x < 1$. Since $0 < x < 1$, we have

$$mx^{m-1} \le \frac{1-x^m}{1-x} = 1 + x + x^2 + \cdots + x^{m-1} \le m.$$

From this, we deduce that

$$\frac{m(1-x)}{x} \le \frac{1-x^m}{x^m} \le \frac{m(1-x)}{x^m},$$

https://doi.org/10.4153/S0008414X24000683 Published online by Cambridge University Press

so that

$$\frac{x^m}{m^2(1-x)} \le \frac{x^m}{m(1-x^m)} \le \frac{x}{m^2(1-x)}.$$

Then

$$\begin{aligned}\log F(x) &= \sum_{n=1}^{\infty} -\log(1-x^n) = \sum_{n=1}^{\infty}\sum_{m=1}^{\infty}\frac{x^{nm}}{m} = \sum_{m=1}^{\infty}\frac{x^m}{m(1-x^m)} \\ &\le \frac{x}{1-x}\sum_{m=1}^{\infty}\frac{1}{m^2} = \frac{\pi^2}{6}\frac{x}{1-x}.\end{aligned} \tag{2.7}$$

Let $q_m(n)$ denote the number of m-colored partitions of n into an even number of distinct parts minus the number of m-colored partitions of n into an odd number of distinct parts. Then, by [2, eq.(6.1.3)]

$$\sum_{n=0}^{\infty} q_1(n)e^{2\pi i n\tau} = \frac{1}{F(e^{2\pi i\tau})} = (q;q)_\infty.$$

Define $\hat{a}_1(n)$ (resp. $\hat{a}_2(n)$) as the number of partitions of n into an even (resp. odd) number of parts with distinct elements. Then, we have $q_1(n) = \hat{a}_1(n) - \hat{a}_2(n)$. Since $0 \le \hat{a}_1(n) \le p(n)$, $0 \le \hat{a}_2(n) \le p(n)$, we have

$$|q_1(n)| = |\hat{a}_1(n) - \hat{a}_2(n)| \le p(n). \tag{2.8}$$

Then,

$$\begin{aligned}\left|\frac{1}{F(e^{2\pi i\tau})}\right| &\le \sum_{n=1}^{\infty}|q_1(n)||e^{2\pi i n\tau}| \le \sum_{n=1}^{\infty}p(n)|e^{2\pi i n\tau}| \\ &= F(|e^{2\pi i\tau}|) = F(e^{-2\pi\mathrm{Im}(\tau)}) \\ &\le \exp\left(\frac{\pi^2 e^{-2\pi\mathrm{Im}(\tau)}}{6(1-e^{-2\pi\mathrm{Im}(\tau)})}\right),\end{aligned} \tag{2.9}$$

where the last inequality follows from (2.7).

An inequality for $F(e^{2\pi i\tau})$ is given in the following lemma.

Lemma 2.1 *Let $\tau_1, \tau_2, \cdots, \tau_n \in \mathbb{C}$ and let $m_1, m_2, \cdots, m_n \in \mathbb{Z}\backslash\{0\}$. Then,*

$$\left|\prod_{i=1}^{n} F(e^{2\pi i\tau_i})^{m_i} - 1\right| \le \exp\left(\sum_{i=1}^{n}\frac{|m_i|\pi^2 e^{-2\pi\mathrm{Im}(\tau_i)}}{6(1-e^{-2\pi\mathrm{Im}(\tau_i)})}\right) - 1. \tag{2.10}$$

Proof Let

$$F(e^{2\pi i\tau_i})^{m_i} =: \sum_{n=1}^{\infty} c_i(n)e^{2\pi i n\tau_i}, \quad F(e^{2\pi i\tau_i})^{|m_i|} =: \sum_{n=1}^{\infty}\tilde{c}_i(n)e^{2\pi i n\tau_i}.$$

If $m_i \ge 0$, then $|c_i(n)| = \tilde{c}_i(n)$; if $m_i < 0$, then, by (2.8),

$$|c_i(n)| \le \sum_{\substack{1\le j\le|m_i| \\ \sum n_j = n}} |q_1(n_j)| \le \sum_{\substack{1\le j\le|m_i| \\ \sum n_j = n}} p(n_j) = \tilde{c}_i(n).$$

https://doi.org/10.4153/S0008414X24000683 Published online by Cambridge University Press

Hence, by (2.7),

$$\begin{aligned}\left|\prod_{i=1}^{n} F(e^{2\pi i\tau_i})^{m_i}-1\right| &\le \left|\sum_{k_1=0}^{\infty}\cdots\sum_{k_n=0}^{\infty} c_1(k_1)\cdots c_n(k_n)e^{2\pi i k_1\tau_1}\cdots e^{2\pi i k_n\tau_n}-1\right| \\ &\le \sum_{\substack{k_1=0\\(k_1,\cdots,k_n)\neq(0,\cdots,0)}}^{\infty}\cdots\sum_{k_n=0}^{\infty} \tilde{c}_1(k_1)\cdots\tilde{c}_n(k_n)|e^{2\pi i k_1\tau_1}|\cdots|e^{2\pi i k_n\tau_n}| \\ &= \prod_{i=1}^{n} F(|e^{2\pi i\tau_i}|)^{|m_i|}-1 \\ &\le \exp\left(\sum_{i=1}^{n}\frac{|m_i|\pi^2 e^{-2\pi\mathrm{Im}(\tau_i)}}{6(1-e^{-2\pi\mathrm{Im}(\tau_i)})}\right)-1.\end{aligned}$$

This completes the proof. ∎

To facilitate certain bounds in the proof of Theorem 1.1, we examine monotonicity of a function here.

Lemma 2.2 *Let $u_1, u_2, \cdots, u_n \in \mathbb{R}_{\ge 1}$ and let $m_1, m_2, \cdots, m_n \in \mathbb{N}$. Then,*

$$W(x) := \frac{1}{x}\left(\exp\left(\sum_{i=1}^{n}\frac{m_i\pi^2 x^{u_i}}{6(1-x^{u_i})}\right)-1\right)$$

is a nondecreasing function of x on $(0,1)$.

Proof Taking derivative of $W(x)$ yields

$$\begin{aligned}W'(x) = &-\frac{1}{x^2}\left(\exp\left(\sum_{i=1}^{n}\frac{m_i\pi^2 x^{u_i}}{6(1-x^{u_i})}\right)-1\right) \\ &+\frac{1}{x}\exp\left(\sum_{i=1}^{n}\frac{m_i\pi^2 x^{u_i}}{6(1-x^{u_i})}\right)\sum_{i=1}^{n}\frac{\pi^2 m_i}{6}\left(\frac{u_i x^{u_i-1}}{(1-x^{u_i})^2}\right).\end{aligned}$$

Therefore, it suffices to prove

$$\sum_{i=1}^{n}\frac{\pi^2 m_i}{6}\left(\frac{u_i x^{u_i}}{(1-x^{u_i})^2}\right)+\exp\left(-\sum_{i=1}^{n}\frac{\pi^2 m_i}{6}\frac{x^{u_i}}{1-x^{u_i}}\right)\ge 1 \tag{2.11}$$

for $x \in (0,1)$. Since

$$\frac{u_i x^{u_i}}{(1-x^{u_i})^2}-\frac{x^{u_i}}{1-x^{u_i}} = \frac{(u_i-1)x^{u_i}+x^{2u_i}}{(1-x^{u_i})^2}\ge 0$$

for $u_i \ge 1$, we have

$$\begin{aligned}&\sum_{i=1}^{n}\frac{\pi^2 m_i}{6}\left(\frac{u_i x^{u_i}}{(1-x^{u_i})^2}\right)+\exp\left(-\sum_{i=1}^{n}\frac{\pi^2 m_i}{6}\frac{x^{u_i}}{1-x^{u_i}}\right) \\ &\quad\ge \sum_{i=1}^{n}\frac{\pi^2 m_i}{6}\frac{x^{u_i}}{1-x^{u_i}}+\exp\left(-\sum_{i=1}^{n}\frac{\pi^2 m_i}{6}\frac{x^{u_i}}{1-x^{u_i}}\right).\end{aligned} \tag{2.12}$$

https://doi.org/10.4153/S0008414X24000683 Published online by Cambridge University Press

Let

$$y := \sum_{i=1}^{n} \frac{\pi^2 m_i}{6} \frac{x^{u_i}}{1-x^{u_i}}.$$

Then, $y > 0$, and so $y + e^{-y} \geq 1$. From this and (2.12), we easily obtain the inequality (2.11). ∎

An estimate for a useful integral is given in the following form. Even though it contains the result in [9, Lemma 3.2] as a special case, its proof is similar to that of [9, Lemma 3.2], and we omit it here.

Lemma 2.3 *Let $(h,k) = 1$, $b \in \mathbb{R}_{\geq 0}$, and define*

$$I := \int_{\xi_{h,k}} e^{\frac{\pi}{12k}\left(\frac{b}{z}-z\right)} z^{-\frac{1}{2}} e^{2\pi n\rho} e^{-2\pi i n\varphi} d\varphi.$$

Then,

$$I = \sqrt{\frac{2}{k(n-1/24)}}\, b^{1/4} \cosh\left(\frac{\pi}{k}\sqrt{\frac{2b}{3}\left(n-\frac{1}{24}\right)}\right) + E^{(b)}(I),$$

where

$$|E^{(b)}(I)| \leq \sqrt{2}\pi^{-1} e^{b\pi/3} \frac{e^{2\pi(n-1/24)\rho} N^{1/2}}{n-1/24}.$$

The following inequalities are also important in the proof of Theorem 1.1.

$$\sum_{k=1}^{N} k^n \leq \begin{cases} N^{n+1}, & \text{if } n \geq 0, \\ \frac{1}{n+1} N^{n+1}, & \text{if } -1 < n < 0. \end{cases}$$

3 Proof of Theorem 1.1

Let

$$\sum_{n=0}^{\infty} a(n) q^n := \frac{(q^9;q^9)_\infty^3}{(q;q)_\infty (q^{27};q^{27})_\infty} = \frac{F(e^{2\pi i\tau}) F(e^{2\pi i(27\tau)})}{F(e^{2\pi i(9\tau)})^3} =: g(q) \tag{3.1}$$

and

$$\begin{aligned} \sum_{n=0}^{\infty} b(n) q^n &:= q^2 \frac{(q^3;q^3)_\infty (q^{27};q^{27})_\infty^2}{(q;q)_\infty (q^9;q^9)_\infty} \\ &= e^{2\pi i(2\tau)} \frac{F(e^{2\pi i\tau}) F(e^{2\pi i(9\tau)})}{F(e^{2\pi i(3\tau)}) F(e^{2\pi i(27\tau)})^2} =: h(q) \end{aligned} \tag{3.2}$$

with $q = e^{2\pi i\tau}$.

In this section, we first employ the modification [14] of the circle method to estimate the coefficient $a(n)$ of $g(q)$ and the coefficient $b(n)$ of $h(q)$ and then obtain information about the sign of the coefficients $\{a(n) - 3b(n)\}_{n\geq 0}$ of $g(q) - 3h(q)$.

https://doi.org/10.4153/S0008414X24000683 Published online by Cambridge University Press

Finally, by [6, eqs.(2.10) and (2.11)], we know that

$$M(0,1,1;9;n)-\frac{p(n)}{3}=\frac{2}{3}(a(n)-3b(n)),$$
$$M(2,3,4;9;n)-\frac{p(n)}{3}=-\frac{1}{3}(a(n)-3b(n)).$$

Therefore, from the sign information of the coefficients $\{a(n)-3b(n)\}_{n\geq 0}$, we can deduce the inequalities in Theorem 1.1.

Since $\gcd(k,27)=1,3,9$ or 27, we split each of the sums $\sum_{\substack{0\leq h<k\leq N\\(h,k)=1}}$ for $a(n)$ and $b(n)$ into four parts according to the values of $\gcd(k,27)$.

It follows from (2.1) that

$$\begin{aligned}a(n)&=\sum_{\substack{0\leq h<k\leq N\\(h,k)=1}}e^{-2\pi inh/k}\int_{\xi_{h,k}}\frac{F(e^{2\pi i\tau})F(e^{2\pi i(27\tau)})}{F(e^{2\pi i(9\tau)})^3}e^{2\pi n\rho}e^{-2\pi in\varphi}d\varphi\\&=\left(\sum_{\substack{1\leq k\leq N\\(k,27)=1}}+\sum_{\substack{1\leq k\leq N\\(k,27)=3}}+\sum_{\substack{1\leq k\leq N\\(k,27)=9}}+\sum_{\substack{1\leq k\leq N\\(k,27)=27}}\right)\sum_{\substack{0\leq h<k\\(h,k)=1}}e^{-2\pi inh/k}\\&\quad\times\int_{\xi_{h,k}}\frac{F(e^{2\pi i\tau})F(e^{2\pi i(27\tau)})}{F(e^{2\pi i(9\tau)})^3}e^{2\pi n\rho}e^{-2\pi in\varphi}d\varphi\\&=:S_1(A)+S_2(A)+S_3(A)+S_4(A)\end{aligned}\tag{3.3}$$

with

$$A=\frac{F(e^{2\pi i\tau})F(e^{2\pi i(27\tau)})}{F(e^{2\pi i(9\tau)})^3}e^{2\pi n\rho}e^{-2\pi in\varphi}$$

and

$$z=k(\rho-i\varphi),\qquad \tau=\frac{h+iz}{k}.$$

Similarly, we get

$$\begin{aligned}b(n)&=\sum_{\substack{0\leq h<k\leq N\\(h,k)=1}}e^{-2\pi inh/k}\int_{\xi_{h,k}}e^{2\pi i(2\tau)}\frac{F(e^{2\pi i\tau})F(e^{2\pi i(9\tau)})}{F(e^{2\pi i(3\tau)})F(e^{2\pi i(27\tau)})^2}e^{2\pi n\rho}e^{-2\pi in\varphi}d\varphi\\&=\left(\sum_{\substack{1\leq k\leq N\\(k,27)=1}}+\sum_{\substack{1\leq k\leq N\\(k,27)=3}}+\sum_{\substack{1\leq k\leq N\\(k,27)=9}}+\sum_{\substack{1\leq k\leq N\\(k,27)=27}}\right)\sum_{\substack{0\leq h<k\\(h,k)=1}}e^{-2\pi inh/k}\\&\quad\times\int_{\xi_{h,k}}e^{2\pi i(2\tau)}\frac{F(e^{2\pi i\tau})F(e^{2\pi i(9\tau)})}{F(e^{2\pi i(3\tau)})F(e^{2\pi i(27\tau)})^2}e^{2\pi n\rho}e^{-2\pi in\varphi}d\varphi\\&=:S_1(B)+S_2(B)+S_3(B)+S_4(B)\end{aligned}\tag{3.4}$$

https://doi.org/10.4153/S0008414X24000683 Published online by Cambridge University Press

with

$$B = e^{2\pi i(2\tau)}\frac{F(e^{2\pi i\tau})F(e^{2\pi i(9\tau)})}{F(e^{2\pi i(3\tau)})F(e^{2\pi i(27\tau)})^2}e^{2\pi n\rho}e^{-2\pi in\varphi}.$$

3.1 Transformation formulas for g and h

We apply (2.4) to transform g and h according to the values of $\gcd(k,27)$.

Case 1: $\gcd(k,27)=1$. By (2.4), we have

$$\frac{F(e^{2\pi i\tau})F(e^{2\pi i(27\tau)})}{F(e^{2\pi i(9\tau)})^3} = \frac{1}{3\sqrt{3}\sqrt{z}}\frac{F(e^{2\pi i\gamma_{(h,k)}(\tau)})F(e^{2\pi i\gamma_{(27h,k)}(27\tau)})}{F(e^{2\pi i\gamma_{(9h,k)}(9\tau)})^3} \times e^{\pi i(s(h,k)+s(27h,k)-3s(9h,k))}e^{\frac{\pi}{12k}(\frac{19}{27z}-z)} \tag{3.5}$$

and

$$\begin{aligned} &e^{2\pi i(2\tau)}\frac{F(e^{2\pi i\tau})F(e^{2\pi i(9\tau)})}{F(e^{2\pi i(3\tau)})F(e^{2\pi i(27\tau)})^2} \\ &= \frac{1}{9\sqrt{3}\sqrt{z}}\frac{F(e^{2\pi i\gamma_{(h,k)}(\tau)})F(e^{2\pi i\gamma_{(9h,k)}(9\tau)})}{F(e^{2\pi i\gamma_{(3h,k)}(3\tau)})F(e^{2\pi i\gamma_{(27h,k)}(27\tau)})^2} \\ &\quad\times e^{2\pi i(2h/k)}e^{\pi\frac{-48z}{12k}}e^{\pi i(s(h,k)+s(9h,k)-s(3h,k)-2s(27h,k))}e^{\frac{\pi}{12k}(\frac{19}{27z}+47z)} \\ &= \frac{1}{9\sqrt{3}\sqrt{z}}\frac{F(e^{2\pi i\gamma_{(h,k)}(\tau)})F(e^{2\pi i\gamma_{(9h,k)}(9\tau)})}{F(e^{2\pi i\gamma_{(3h,k)}(3\tau)})F(e^{2\pi i\gamma_{(27h,k)}(27\tau)})^2} \\ &\quad\times e^{2\pi i(2h/k)}e^{\pi i(s(h,k)+s(9h,k)-s(3h,k))-2s(27h,k)}e^{\frac{\pi}{12k}(\frac{19}{27z}-z)}. \end{aligned} \tag{3.6}$$

Case 2: $\gcd(k,27)=3$. Let $k=3l_1$. Then,

$$\frac{F(e^{2\pi i\tau})F(e^{2\pi i(27\tau)})}{F(e^{2\pi i(9\tau)})^3} = \frac{1}{\sqrt{3}\sqrt{z}}\frac{F(e^{2\pi i\gamma_{(h,3l_1)}(\tau)})F(e^{2\pi i\gamma_{(9h,l_1)}(27\tau)})}{F(e^{2\pi i\gamma_{(3h,l_1)}(9\tau)})^3} \times e^{\pi i(s(h,3l_1)+s(9h,l_1)-3s(3h,l_1))}e^{\frac{\pi}{12k}(-\frac{5}{3z}-z)} \tag{3.7}$$

and

$$\begin{aligned} &e^{2\pi i(2\tau)}\frac{F(e^{2\pi i\tau})F(e^{2\pi i(9\tau)})}{F(e^{2\pi i(3\tau)})F(e^{2\pi i(27\tau)})^2} \\ &= \frac{1}{3\sqrt{3}\sqrt{z}}\frac{F(e^{2\pi i\gamma_{(h,3l_1)}(\tau)})F(e^{2\pi i\gamma_{(3h,l_1)}(9\tau)})}{F(e^{2\pi i\gamma_{(h,l_1)}(3\tau)})F(e^{2\pi i\gamma_{(9h,l_1)}(27\tau)})^2} \\ &\quad\times e^{2\pi i(2h/k)}e^{\pi i(s(h,3l_1)+s(3h,l_1)-s(h,l_1)-2s(9h,l_1))}e^{\frac{\pi}{12k}(-\frac{5}{3z}-z)}. \end{aligned} \tag{3.8}$$

Case 3: $\gcd(k,27)=9$. Let $k=9l_2$. It follows that

$$\frac{F(e^{2\pi i\tau})F(e^{2\pi i(27\tau)})}{F(e^{2\pi i(9\tau)})^3} = \frac{\sqrt{3}}{\sqrt{z}}\frac{F(e^{2\pi i\gamma_{(h,9l_2)}(\tau)})F(e^{2\pi i\gamma_{(3h,l_2)}(27\tau)})}{F(e^{2\pi i\gamma_{(h,l_2)}(9\tau)})^3} \times e^{\pi i(s(h,9l_2)+s(3h,l_2)-3s(h,l_2))}e^{\frac{\pi}{12k}(-\frac{23}{z}-z)}$$

https://doi.org/10.4153/S0008414X24000683 Published online by Cambridge University Press

and

$$e^{2\pi i(2\tau)}\frac{F(e^{2\pi i\tau})F(e^{2\pi i(9\tau)})}{F(e^{2\pi i(3\tau)})F(e^{2\pi i(27\tau)})^2}$$
$$=\frac{1}{3\sqrt{z}}\frac{F(e^{2\pi i\gamma_{(h,9l_2)}(\tau)})F(e^{2\pi i\gamma_{(h,l_2)}(9\tau)})}{F(e^{2\pi i\gamma_{(h,3l_2)}(3\tau)})F(e^{2\pi i\gamma_{(3h,l_2)}(27\tau)})^2}$$
$$\times e^{2\pi i(2h/k)}e^{\pi i(s(h,9l_2)+s(h,l_2)-s(h,3l_1)-2s(3h,l_2))}e^{\frac{\pi}{12k}(\frac{1}{z}-z)}.$$

Case 4: $\gcd(k,27)=27$. Let $k=27l_3$. We have

$$\frac{F(e^{2\pi i\tau})F(e^{2\pi i(27\tau)})}{F(e^{2\pi i(9\tau)})^3}=\frac{1}{\sqrt{z}}\frac{F(e^{2\pi i\gamma_{(h,27l_3)}(\tau)})F(e^{2\pi i\gamma_{(h,l_3)}(27\tau)})}{F(e^{2\pi i\gamma_{(h,3l_3)}(9\tau)})^3}$$
$$\times e^{\pi i(s(h,27l_3)+s(h,l_3)-3s(h,3l_3))}e^{\frac{\pi}{12k}(\frac{1}{z}-z)} \tag{3.9}$$

and

$$e^{2\pi i(2\tau)}\frac{F(e^{2\pi i\tau})F(e^{2\pi i(9\tau)})}{F(e^{2\pi i(3\tau)})F(e^{2\pi i(27\tau)})^2}$$
$$=\frac{1}{\sqrt{z}}\frac{F(e^{2\pi i\gamma_{(h,27l_3)}(\tau)})F(e^{2\pi i\gamma_{(h,3l_3)}(9\tau)})}{F(e^{2\pi i\gamma_{(h,9l_3)}(3\tau)})F(e^{2\pi i\gamma_{(h,l_3)}(27\tau)})^2}$$
$$\times e^{2\pi i(2h/k)}e^{\pi i(s(h,27l_3)+s(h,3l_3)-s(h,9l_3)-2s(h,l_3))}e^{\frac{\pi}{12k}(-\frac{47}{z}-z)}.$$

From the above transformation formulas for g and h, we see that the integrands of the integrals in $S_2(A)$, $S_2(B)$, $S_3(A)$, and $S_4(B)$ have factors of the forms $e^{\frac{\delta}{z}}$ with $\delta<0$. For these integrals, we use (2.3), (2.5), (2.6), and (2.9) to give bounds. However, since the integrands of the integrals in $S_1(A)$, $S_1(B)$, $S_3(B)$, and $S_4(A)$ contain factors of the forms $e^{\frac{\delta}{z}}$ with $\delta>0$, we split these integrals into two parts and then apply Lemma 2.2, (2.3), (2.5), (2.6), and (2.10) to estimate the second parts. For the first parts, we employ Lemma 2.3 to tackle the integrals. It should be mentioned that our main term $P(n)$ (in Subsection 3.3) originates from $S_3(B)$.

3.2 Bounding $S_2(A)$, $S_2(B)$, $S_3(A)$, and $S_4(B)$

For $S_2(A)$, we apply (3.7) to deduce that

$$A=\frac{1}{\sqrt{3}}z^{-\frac{1}{2}}e^{\pi i(s(h,3l_1)+s(9h,l_1)-3s(3h,l_1))}e^{\frac{\pi}{12k}(-\frac{5}{3z}-z)}$$
$$\times\frac{F(e^{2\pi i\gamma_{(h,3l_1)}(\tau)})F(e^{2\pi i\gamma_{(9h,l_1)}(27\tau)})}{F(e^{2\pi i\gamma_{(3h,l_1)}(9\tau)})^3}e^{2\pi n\rho}e^{-2\pi in\varphi}$$

https://doi.org/10.4153/S0008414X24000683 Published online by Cambridge University Press

with $k = 3l_1$. Using (2.3), (2.5), (2.6), and (2.9), we find that

$$\begin{aligned}|A| \le & \frac{1}{\sqrt{3}} F\big(e^{-2\pi(\frac{1}{k})\mathrm{Re}(\frac{1}{z})}\big) F\big(e^{-2\pi(\frac{3}{k})\mathrm{Re}(\frac{1}{9z})}\big) F\big(e^{-2\pi(\frac{3}{k})\mathrm{Re}(\frac{1}{3z})}\big)^3 \\ & \times e^{-\frac{5\pi}{36k}\mathrm{Re}(\frac{1}{z})} e^{-\frac{\pi}{12k}\mathrm{Re}(z)} |z|^{-\frac{1}{2}} e^{2\pi n\rho} \\ \le & \frac{1}{\sqrt{3}} \exp\left(\frac{2\pi^2 e^{-\pi}}{3(1-e^{-\pi})} + \frac{\pi^2 e^{-\pi/3}}{6(1-e^{-\pi/3})}\right) \times e^{-\frac{5\pi}{72}} e^{2\pi(n-\frac{1}{24})\rho} k^{-\frac{1}{2}} N \\ \le & \frac{1}{\sqrt{3}} e^{0.969} e^{2\pi(n-\frac{1}{24})\rho} k^{-\frac{1}{2}} N.\end{aligned}$$

So we have

$$\begin{aligned}|S_2(A)| \le & \sum_{\substack{1\le k\le N \\ (k,27)=3}} \sum_{\substack{0\le h<k \\ (h,k)=1}} \int_{\xi_{h,k}} \frac{1}{\sqrt{3}} e^{0.969} e^{2\pi(n-\frac{1}{24})\rho} k^{-\frac{1}{2}} N d\varphi \\ \le & \sum_{\substack{1\le k\le N \\ (k,27)=3}} \sum_{\substack{0\le h<k \\ (h,k)=1}} \frac{1}{\sqrt{3}} e^{0.969} e^{2\pi(n-\frac{1}{24})\rho} \frac{2}{k^{\frac{3}{2}}} \\ \le & \frac{2}{\sqrt{3}} e^{0.969} e^{2\pi(n-\frac{1}{24})\rho} \sum_{1\le k\le N} k^{-\frac{1}{2}} \\ \le & \frac{4}{\sqrt{3}} e^{0.969} e^{2\pi(n-\frac{1}{24})\rho} N^{\frac{1}{2}}.\end{aligned}$$

For $S_2(B)$, we apply (3.8) to give

$$\begin{aligned}B = & \frac{1}{3\sqrt{3}} z^{-\frac{1}{2}} e^{2\pi i(2h/k)} e^{\pi i(s(h,3l_1)+s(3h,l_1)-s(h,l_1)-2s(9h,l_1))} e^{\frac{\pi}{12k}(-\frac{5}{3z}-z)} \\ & \times \frac{F\big(e^{2\pi i\gamma_{(h,3l_1)}(\tau)}\big) F\big(e^{2\pi i\gamma_{(3h,l_1)}(9\tau)}\big)}{F\big(e^{2\pi i\gamma_{(h,l_1)}(3\tau)}\big) F\big(e^{2\pi i\gamma_{(9h,l_1)}(27\tau)}\big)^2} e^{2\pi n\rho} e^{-2\pi i n\varphi}\end{aligned}$$

with $k = 3l_1$. Using (2.3), (2.5), (2.6), and (2.9), we find that

$$\begin{aligned}|B| \le & \frac{1}{3\sqrt{3}} F\big(e^{-2\pi(\frac{1}{k})\mathrm{Re}(\frac{1}{z})}\big) F\big(e^{-2\pi(\frac{3}{k})\mathrm{Re}(\frac{1}{3z})}\big) F\big(e^{-2\pi(\frac{3}{k})\mathrm{Re}(\frac{1}{z})}\big) F\big(e^{-2\pi(\frac{3}{k})\mathrm{Re}(\frac{1}{9z})}\big)^2 \\ & \times e^{-\frac{5\pi}{36k}\mathrm{Re}(\frac{1}{z})} e^{-\frac{\pi}{12k}\mathrm{Re}(z)} |z|^{-\frac{1}{2}} e^{2\pi n\rho} \\ \le & \frac{1}{3\sqrt{3}} \exp\left(\frac{\pi^2 e^{-\pi}}{3(1-e^{-\pi})} + \frac{\pi^2 e^{-3\pi}}{6(1-e^{-3\pi})} + \frac{\pi^2 e^{-\pi/3}}{3(1-e^{-\pi/3})}\right) e^{-\frac{5\pi}{72}} e^{2\pi(n-\frac{1}{24})\rho} k^{-\frac{1}{2}} N \\ \le & \frac{1}{3\sqrt{3}} e^{1.710} e^{2\pi(n-\frac{1}{24})\rho} k^{-\frac{1}{2}} N\end{aligned}$$

https://doi.org/10.4153/S0008414X24000683 Published online by Cambridge University Press

and

$$\begin{aligned}|S_2(B)| &\le \sum_{\substack{1\le k\le N\\(k,27)=3}}\sum_{\substack{0\le h<k\\(h,k)=1}}\int_{\xi_{h,k}}\frac{1}{3\sqrt{3}}e^{1.710}e^{2\pi(n-\frac{1}{24})\rho}k^{-\frac{1}{2}}Nd\varphi\\ &\le \sum_{\substack{1\le k\le N\\(k,27)=3}}\sum_{\substack{0\le h<k\\(h,k)=1}}\frac{1}{3\sqrt{3}}e^{1.710}e^{2\pi(n-\frac{1}{24})\rho}\frac{2}{k^{\frac{3}{2}}}\\ &\le \frac{2}{3\sqrt{3}}e^{1.710}e^{2\pi(n-\frac{1}{24})\rho}\sum_{1\le k\le N}k^{-\frac{1}{2}}\\ &\le \frac{4}{3\sqrt{3}}e^{1.710}e^{2\pi(n-\frac{1}{24})\rho}N^{\frac{1}{2}}.\end{aligned}$$

Similarly, for $S_3(A)$ and $S_4(B)$, we get

$$|S_3(A)| \le 4\sqrt{3}e^{-2.047}e^{2\pi(n-\frac{1}{24})\rho}N^{\frac{1}{2}}$$

and

$$|S_4(B)| \le 4e^{-6.077}e^{2\pi(n-\frac{1}{24})\rho}N^{\frac{1}{2}}.$$

3.3 Tackling $S_1(A)$, $S_1(B)$, $S_3(B)$, and $S_4(A)$

For $S_1(A)$, set

$$\omega_{(h,k)}^{(1)} := e^{\pi i(s(h,k)+s(27h,k)-3s(9h,k))}.$$

We apply (3.5) to deduce

$$\begin{aligned}S_1(A) &= \sum_{\substack{1\le k\le N\\(k,27)=1}}\sum_{\substack{0\le h<k\\(h,k)=1}}e^{-2\pi inh/k}\int_{\xi_{h,k}}\frac{F(e^{2\pi i\tau})F(e^{2\pi i(27\tau)})}{F(e^{2\pi i(9\tau)})^3}e^{2\pi n\rho}e^{-2\pi in\varphi}d\varphi\\ &= \sum_{\substack{1\le k\le N\\(k,27)=1}}\sum_{\substack{0\le h<k\\(h,k)=1}}e^{-2\pi inh/k}\omega_{(h,k)}^{(1)}\int_{\xi_{h,k}}\frac{1}{3\sqrt{3}\sqrt{z}}e^{\frac{\pi}{12k}(\frac{19}{27z}-z)}\\ &\quad\times\frac{F(e^{2\pi i\gamma_{(h,k)}(\tau)})F(e^{2\pi i\gamma_{(27h,k)}(27\tau)})}{F(e^{2\pi i\gamma_{(9h,k)}(9\tau)})^3}e^{2\pi n\rho}e^{-2\pi in\varphi}d\varphi\\ &= \sum_{\substack{1\le k\le N\\(k,27)=1}}\sum_{\substack{0\le h<k\\(h,k)=1}}e^{-2\pi inh/k}\omega_{(h,k)}^{(1)}\int_{\xi_{h,k}}\frac{1}{3\sqrt{3}\sqrt{z}}e^{\frac{\pi}{12k}(\frac{19}{27z}-z)}e^{2\pi n\rho}e^{-2\pi in\varphi}d\varphi\\ &\quad+\sum_{\substack{1\le k\le N\\(k,27)=1}}\sum_{\substack{0\le h<k\\(h,k)=1}}e^{-2\pi inh/k}\omega_{(h,k)}^{(1)}\int_{\xi_{h,k}}\frac{1}{3\sqrt{3}\sqrt{z}}e^{\frac{\pi}{12k}(\frac{19}{27z}-z)}\\ &\quad\times\left(\frac{F(e^{2\pi i\gamma_{(h,k)}(\tau)})F(e^{2\pi i\gamma_{(27h,k)}(27\tau)})}{F(e^{2\pi i\gamma_{(9h,k)}(9\tau)})^3}-1\right)e^{2\pi n\rho}e^{-2\pi in\varphi}d\varphi\\ &=: T_1+T_2.\end{aligned}$$

https://doi.org/10.4153/S0008414X24000683 Published online by Cambridge University Press

By (2.9) and (2.10), we have

$$\left|e^{\frac{\pi}{12k}\left(\frac{19}{27z}\right)}\left(\frac{F\left(e^{2\pi i\gamma_{(h,k)}(\tau)}\right)F\left(e^{2\pi i\gamma_{(27h,k)}(27\tau)}\right)}{F\left(e^{2\pi i\gamma_{(9h,k)}(9\tau)}\right)^3}-1\right)\right|\le e^{\frac{19\pi}{27\times12k}\mathrm{Re}\left(\frac{1}{z}\right)}\left(e^{f(k,z)}-1\right),$$

where

$$f(k,z)=\frac{\pi^2e^{-2\pi\frac{1}{k}\mathrm{Re}\left(\frac{1}{z}\right)}}{6\left(1-e^{-2\pi\frac{1}{k}\mathrm{Re}\left(\frac{1}{z}\right)}\right)}+\frac{\pi^2e^{-2\pi\frac{1}{27k}\mathrm{Re}\left(\frac{1}{z}\right)}}{6\left(1-e^{-2\pi\frac{1}{27k}\mathrm{Re}\left(\frac{1}{z}\right)}\right)}+\frac{3\pi^2e^{-2\pi\frac{1}{9k}\mathrm{Re}\left(\frac{1}{z}\right)}}{6\left(1-e^{-2\pi\frac{1}{9k}\mathrm{Re}\left(\frac{1}{z}\right)}\right)}.$$

Let

$$x:=e^{-\frac{19\pi}{27\times12k}\mathrm{Re}\left(\frac{1}{z}\right)}.$$

Then,

$$\begin{aligned}&\left|e^{\frac{\pi}{12k}\left(\frac{19}{27z}\right)}\left(\frac{F\left(e^{2\pi i\gamma_{(h,k)}(\tau)}\right)F\left(e^{2\pi i\gamma_{(27h,k)}(27\tau)}\right)}{F\left(e^{2\pi i\gamma_{(9h,k)}(9\tau)}\right)^3}-1\right)\right|\\&\quad\le\frac{1}{x}\left(\exp\left(\frac{\pi^2x^{\frac{648}{19}}}{6\left(1-x^{\frac{648}{19}}\right)}+\frac{\pi^2x^{\frac{24}{19}}}{6\left(1-x^{\frac{24}{19}}\right)}+\frac{3\pi^2x^{\frac{72}{19}}}{6\left(1-x^{\frac{72}{19}}\right)}\right)-1\right)=:W_1(x).\end{aligned}$$

By Lemma 2.2 and (2.5), we get

$$\begin{aligned}W_1(x)&\le W_1\left(e^{-\frac{19\pi}{648}}\right)=e^{\frac{19\pi}{648}}\left(\exp\left(\frac{\pi^2e^{-\pi}}{6\left(1-e^{-\pi}\right)}+\frac{\pi^2e^{-\pi/27}}{6\left(1-e^{-\pi/27}\right)}+\frac{3\pi^2e^{-\pi/9}}{6\left(1-e^{-\pi/9}\right)}\right)-1\right)\\&\le9.819\times10^{10}.\end{aligned}$$

Then, by (2.6),

$$\begin{aligned}|T_2|&\le\sum_{\substack{1\le k\le N\\(k,27)=1}}\sum_{\substack{0\le h<k\\(h,k)=1}}\int_{\xi_{h,k}}1.890\times10^{10}e^{-\frac{\pi}{12k}\mathrm{Re}(z)}|z|^{-1/2}e^{-2\pi in\rho}d\varphi\\&\le\sum_{\substack{1\le k\le N\\(k,27)=1}}\sum_{\substack{0\le h<k\\(h,k)=1}}1.890\times10^{10}e^{2\pi(n-1/24)\rho}\frac{2}{k^{3/2}}\\&\le\sum_{\substack{1\le k\le N\\(k,27)=1}}1.890\times10^{10}e^{2\pi(n-1/24)\rho}\frac{2}{k^{1/2}}\\&\le7.560\times10^{10}e^{2\pi(n-1/24)\rho}N^{1/2}.\end{aligned}$$

Similarly, we use (3.6) to derive that

$$S_1(B)=R_1+R_2,$$

where

$$R_1:=\sum_{\substack{1\le k\le N\\(k,27)=1}}\sum_{\substack{0\le h<k\\(h,k)=1}}e^{-2\pi i(n-2)h/k}\omega^{(2)}_{(h,k)}\int_{\xi_{h,k}}\frac{1}{9\sqrt{3}\sqrt{z}}e^{\frac{\pi}{12k}\left(\frac{19}{27z}-z\right)}e^{2\pi n\rho}e^{-2\pi in\varphi}d\varphi,$$

and

$$|R_2|\le1.435\times10^{13}e^{2\pi i(n-1/24)\rho}N^{1/2}.$$

https://doi.org/10.4153/S0008414X24000683 Published online by Cambridge University Press

Here, $\omega_{(h,k)}^{(2)} := e^{\pi i(s(h,k)+s(9h,k)-s(3h,k)-2s(27h,k))}$.

Using the software *Mathematica*, we find that[2]

$$4s(9h,k)-s(3h,k)-3s(27h,k)+\frac{4h}{k}\equiv 0 \pmod 2, \tag{3.10}$$

where $0\le h<k\le 17$, $(k,3)=1$, and $(h,k)=1$. From this, we have

$$\sum_{\substack{0\le h<k\\(h,k)=1}} e^{-2\pi inh/k}\omega_{(h,k)}^{(1)} = \sum_{\substack{0\le h<k\\(h,k)=1}} e^{-2\pi i(n-2)h/k}\omega_{(h,k)}^{(2)}$$

with $k\le 17$ and $\gcd(k,27)=1$. Then,

$$\begin{aligned}
|T_1-3R_1| &= \Bigg| \sum_{\substack{1\le k\le N\\(k,27)=1}}\sum_{\substack{0\le h<k\\(h,k)=1}} e^{-2\pi inh/k}\omega_{(h,k)}^{(1)}\int_{\xi_{h,k}} \frac{1}{3\sqrt{3}\sqrt{z}} e^{\frac{\pi}{12k}\left(\frac{19}{27z}-z\right)} e^{2\pi n\rho} e^{-2\pi in\varphi} d\varphi \\
&\quad - \sum_{\substack{1\le k\le N\\(k,27)=1}}\sum_{\substack{0\le h<k\\(h,k)=1}} e^{-2\pi i(n-2)h/k}\omega_{(h,k)}^{(2)} \\
&\quad \int_{\xi_{h,k}} \frac{1}{3\sqrt{3}\sqrt{z}} e^{\frac{\pi}{12k}\left(\frac{19}{27z}-z\right)} e^{2\pi n\rho} e^{-2\pi in\varphi} d\varphi \Bigg| \\
&\le \Bigg| \sum_{\substack{19\le k\le N\\(k,27)=1}}\sum_{\substack{0\le h<k\\(h,k)=1}} e^{-2\pi inh/k}\omega_{(h,k)}^{(1)}\int_{\xi_{h,k}} \frac{1}{3\sqrt{3}\sqrt{z}} e^{\frac{\pi}{12k}\left(\frac{19}{27z}-z\right)} e^{2\pi n\rho} e^{-2\pi in\varphi} d\varphi \Bigg| \\
&\quad + \Bigg| \sum_{\substack{19\le k\le N\\(k,27)=1}}\sum_{\substack{0\le h<k\\(h,k)=1}} e^{-2\pi i(n-2)h/k}\omega_{(h,k)}^{(2)} \\
&\quad \int_{\xi_{h,k}} \frac{1}{3\sqrt{3}\sqrt{z}} e^{\frac{\pi}{12k}\left(\frac{19}{27z}-z\right)} e^{2\pi n\rho} e^{-2\pi in\varphi} d\varphi \Bigg|.
\end{aligned}$$

By Lemma 2.3, we have

$$\begin{aligned}
&\sum_{\substack{19\le k\le N\\(k,27)=1}}\sum_{\substack{0\le h<k\\(h,k)=1}} \left| \int_{\xi_{h,k}} \frac{1}{3\sqrt{3}\sqrt{z}} e^{\frac{\pi}{12k}\left(\frac{19}{27z}-z\right)} e^{2\pi n\rho} e^{-2\pi in\varphi} d\varphi \right| \\
&\le \sum_{\substack{19\le k\le N\\(k,27)=1}}\sum_{\substack{0\le h<k\\(h,k)=1}} \sqrt{\frac{2}{k(n-1/24)}} \left(\frac{19}{27}\right)^{1/4} \cosh\left(\frac{\pi}{k}\sqrt{\frac{19}{27}\times\frac{2}{3}\left(n-\frac{1}{24}\right)}\right) \\
&\quad + \sum_{\substack{19\le k\le N\\(k,27)=1}}\sum_{\substack{0\le h<k\\(h,k)=1}} \sqrt{2}\pi^{-1} e^{19\pi/81} \frac{e^{2\pi(n-1/24)\rho} N^{1/2}}{n-1/24} \\
&\le \sum_{\substack{19\le k\le N\\(k,27)=1}} \sqrt{\frac{2k}{n-1/24}} \cosh\left(\frac{\pi}{18}\sqrt{\frac{2}{3}\left(n-\frac{1}{24}\right)}\right)
\end{aligned}$$

[2] We conjecture here that the congruence (3.10) holds for $0\le h<k$, $(k,3)=1$, and $(h,k)=1$.

https://doi.org/10.4153/S0008414X24000683 Published online by Cambridge University Press

$$+\sum_{\substack{19\le k\le N\\(k,27)=1}} 1.2828\frac{e^{2\pi(n-1/24)\rho}kN^{1/2}}{n-1/24}$$
$$\le N^{3/2}\sqrt{\frac{2}{n-1/24}}\cosh\left(\frac{\pi}{18}\sqrt{\frac{2}{3}\left(n-\frac{1}{24}\right)}\right)+1.2828\frac{e^{2\pi(n-1/24)\rho}N^{5/2}}{n-1/24}.$$

So we have

$$|T_1-3R_1|\le 2N^{3/2}\sqrt{\frac{2}{(n-1/24)}}\cosh\left(\frac{\pi}{18}\sqrt{\frac{2}{3}\left(n-\frac{1}{24}\right)}\right)+2.5656\frac{e^{2\pi(n-1/24)\rho}N^{5/2}}{n-1/24}.$$

Setting $N:=\sqrt{2\pi(n-1/24)}$, we obtain

$$\begin{aligned}|T_1-3R_1| &\le 2(2\pi)^{3/4}\sqrt{2}(n-1/24)^{(1/4)}\left(e^{\frac{\pi}{18}\sqrt{\frac{2}{3}\left(n-\frac{1}{24}\right)}}+1\right)\\ &\quad+2.5656e(2\pi)^{5/4}(n-1/24)^{(1/4)}\\ &\le 11.225(n-1/24)^{(1/4)}e^{\frac{\pi}{18}\sqrt{\frac{2}{3}\left(n-\frac{1}{24}\right)}}+80.601(n-1/24)^{(1/4)}.\end{aligned}$$

For $S_3(B)$, set $\Omega_{(h,k)}:=e^{\pi i(s(h,9l_2)+s(h,l_2)-s(h,3l_1)-2s(3h,l_2))}$. We apply (3.1) to conclude that

$$\begin{aligned}S_3(B) &= \sum_{\substack{1\le k\le N\\(k,27)=9}}\sum_{\substack{0\le h<k\\(h,k)=1}} e^{-2\pi inh/k}\int_{\xi_{h,k}} e^{2\pi i(2\tau)}\frac{F(e^{2\pi i\tau})F(e^{2\pi i(9\tau)})}{F(e^{2\pi i(3\tau)})F(e^{2\pi i(27\tau)})^2}e^{2\pi n\rho}e^{-2\pi in\varphi}d\varphi\\
&= \sum_{\substack{1\le k\le N\\(k,27)=9}}\sum_{\substack{0\le h<k\\(h,k)=1}} e^{-2\pi i(n-2)h/k}\Omega_{(h,k)}\int_{\xi_{h,k}}\frac{1}{3\sqrt{z}}e^{\frac{\pi}{12k}\left(\frac{1}{z}-z\right)}\\
&\quad\times\frac{F(e^{2\pi i\gamma_{(h,9l_2)}(\tau)})F(e^{2\pi i\gamma_{(h,l_2)}(9\tau)})}{F(e^{2\pi i\gamma_{(h,3l_2)}(3\tau)})F(e^{2\pi i\gamma_{(3h,l_2)}(27\tau)})^2}e^{2\pi n\rho}e^{-2\pi in\varphi}d\varphi\\
&= \sum_{\substack{1\le k\le N\\(k,27)=9}}\sum_{\substack{0\le h<k\\(h,k)=1}} e^{-2\pi i(n-2)h/k}\Omega_{(h,k)}\int_{\xi_{h,k}}\frac{1}{3\sqrt{z}}e^{\frac{\pi}{12k}\left(\frac{1}{z}-z\right)}e^{2\pi n\rho}e^{-2\pi in\varphi}d\varphi\\
&\quad+\sum_{\substack{1\le k\le N\\(k,27)=9}}\sum_{\substack{0\le h<k\\(h,k)=1}} e^{-2\pi i(n-2)h/k}\Omega_{(h,k)}\int_{\xi_{h,k}}\frac{1}{3\sqrt{z}}e^{\frac{\pi}{12k}\left(\frac{1}{z}-z\right)}\\
&\quad\times\left(\frac{F(e^{2\pi i\gamma_{(h,9l_2)}(\tau)})F(e^{2\pi i\gamma_{(h,l_2)}(9\tau)})}{F(e^{2\pi i\gamma_{(h,3l_2)}(3\tau)})F(e^{2\pi i\gamma_{(3h,l_2)}(27\tau)})^2}-1\right)e^{2\pi n\rho}e^{-2\pi in\varphi}d\varphi\\
&=: P_1+P_2\end{aligned}$$

with $k=9l_2$.

For P_2, by (2.10), we have

$$\left|e^{\frac{\pi}{12k}\left(\frac{1}{z}\right)}\left(\frac{F(e^{2\pi i\gamma_{(h,9l_2)}(\tau)})F(e^{2\pi i\gamma_{(h,l_2)}(9\tau)})}{F(e^{2\pi i\gamma_{(h,3l_2)}(3\tau)})F(e^{2\pi i\gamma_{(3h,l_2)}(27\tau)})^2}-1\right)\right|\le e^{\frac{\pi}{12k}\mathrm{Re}\left(\frac{1}{z}\right)}\left(e^{g(k,z)}-1\right),$$

https://doi.org/10.4153/S0008414X24000683 Published online by Cambridge University Press

where

$$g(k,z)=\frac{\pi^2 e^{-2\pi\frac{1}{k}\mathrm{Re}(\frac{1}{z})}}{6(1-e^{-2\pi\frac{1}{k}\mathrm{Re}(\frac{1}{z})})}+\frac{\pi^2 e^{-2\pi\frac{9}{k}\mathrm{Re}(\frac{1}{z})}}{6(1-e^{-2\pi\frac{9}{k}\mathrm{Re}(\frac{1}{z})})}+\frac{3\pi^2 e^{-2\pi\frac{3}{k}\mathrm{Re}(\frac{1}{z})}}{6(1-e^{-2\pi\frac{3}{k}\mathrm{Re}(\frac{1}{z})})}.$$

Let

$$x:=e^{-\frac{\pi}{12k}\mathrm{Re}(\frac{1}{z})}.$$

Then,

$$\begin{aligned}&\left|e^{\frac{\pi}{12k}(\frac{1}{z})}\left(\frac{F(e^{2\pi i\gamma_{(h,9l_2)}(\tau)})F(e^{2\pi i\gamma_{(h,l_2)}(9\tau)})}{F(e^{2\pi i\gamma_{(h,3l_2)}(3\tau)})F(e^{2\pi i\gamma_{(3h,l_2)}(27\tau)})^2}-1\right)\right|\\&\quad\le\frac{1}{x}\left(\exp\left(\frac{\pi^2x^{24}}{6(1-x^{24})}+\frac{\pi^2x^{216}}{6(1-x^{216})}+\frac{3\pi^2x^{72}}{6(1-x^{72})}\right)-1\right)=:W_3(x).\end{aligned}$$

By Lemma 2.2 and (2.5), we have

$$\begin{aligned}W_3(x)&\le W_3(e^{-\frac{\pi}{24}})=e^{\frac{\pi}{24}}\left(\exp\left(\frac{\pi^2e^{-\pi}}{6(1-e^{-\pi})}+\frac{\pi^2e^{-9\pi}}{6(1-e^{-9\pi})}+\frac{\pi^2e^{-3\pi}}{2(1-e^{-3\pi})}\right)-1\right)\\&\le 0.089.\end{aligned}$$

Hence,

$$\begin{aligned}|P_2|&\le 0.0297\sum_{\substack{1\le k\le N\\(k,27)=9}}\sum_{\substack{0\le h<k\\(h,k)=1}}\int_{\xi_{h,k}}|z|^{-\frac{1}{2}}e^{\frac{-\pi}{12k}\mathrm{Re}(z)}e^{2\pi n\rho}d\varphi\\&\le 0.0297\sum_{\substack{1\le k\le N\\(k,27)=9}}\sum_{\substack{0\le h<k\\(h,k)=1}}\int_{\xi_{h,k}}k^{-\frac{1}{2}}Ne^{2\pi(n-1/24)\rho}d\varphi\\&\le 0.0594\sum_{\substack{1\le k\le N\\(k,27)=9}}\sum_{\substack{0\le h<k\\(h,k)=1}}k^{-\frac{3}{2}}e^{2\pi(n-1/24)\rho}\\&\le 0.0594\sum_{\substack{1\le k\le N\\(k,27)=9}}k^{-\frac{1}{2}}e^{2\pi(n-1/24)\rho}\\&\le 0.119e^{2\pi(n-1/24)\rho}N^{\frac{1}{2}}.\end{aligned}$$

For P_1, applying Lemma 2.3 with $b=1$, we establish

$$\begin{aligned}P_1&=\frac{1}{3}\sum_{\substack{1\le k\le N\\(k,27)=9}}\sum_{\substack{0\le h<k\\(h,k)=1}}e^{-2\pi i(n-2)h/k}\Omega_{(h,k)}\int_{\xi_{h,k}}e^{\frac{\pi}{12k}(\frac{1}{z}-z)}z^{-\frac{1}{2}}e^{2\pi n\rho}e^{-2\pi in\varphi}d\varphi\\&=\frac{1}{3}\sum_{\substack{0\le h<k\\(h,k)=1}}e^{-2\pi i(n-2)h/9}\Omega_{(h,9)}\sqrt{\frac{2}{9(n-1/24)}}\cosh\left(\frac{\pi}{9}\sqrt{\frac{2}{3}\left(n-\frac{1}{24}\right)}\right)\\&\quad+\frac{1}{3}\sum_{\substack{18\le k\le N\\(k,27)=9}}\sum_{\substack{0\le h<k\\(h,k)=1}}e^{-2\pi i(n-2)h/k}\Omega_{(h,k)}\sqrt{\frac{2}{k(n-1/24)}}\cosh\left(\frac{\pi}{k}\sqrt{\frac{2}{3}\left(n-\frac{1}{24}\right)}\right)\end{aligned}$$

https://doi.org/10.4153/S0008414X24000683 Published online by Cambridge University Press

$$
\begin{aligned}
&+\frac{1}{3}\sum_{\substack{1\le k\le N\\(k,27)=9}}\sum_{\substack{0\le h<k\\(h,k)=1}} e^{-2\pi i(n-2)h/k}\Omega_{(h,k)}E^{(1)}(I)\\
&=\frac{1}{3}\Bigg(\sum_{\substack{0\le h<9\\(h,9)=1}} e^{-2\pi i(n-2)h/9}\Omega_{(h,9)}\Bigg)\sqrt{\frac{2}{9(n-1/24)}}\,e^{\frac{\pi}{9}\sqrt{\frac{2}{3}\left(n-\frac{1}{24}\right)}}\\
&\quad+\frac{1}{3}\Bigg(\sum_{\substack{0\le h<9\\(h,9)=1}} e^{-2\pi i(n-2)h/9}\Omega_{(h,9)}\Bigg)\sqrt{\frac{2}{9(n-1/24)}}\,e^{-\frac{\pi}{9}\sqrt{\frac{2}{3}\left(n-\frac{1}{24}\right)}}\\
&\quad+\frac{1}{3}\sum_{\substack{18\le k\le N\\(k,27)=9}}\sum_{\substack{0\le h<k\\(h,k)=1}} e^{-2\pi i(n-2)h/k}\Omega_{(h,k)}\sqrt{\frac{2}{k(n-1/24)}}\cosh\left(\frac{\pi}{k}\sqrt{\frac{2}{3}\left(n-\frac{1}{24}\right)}\right)\\
&\quad+\frac{1}{3}\sum_{\substack{1\le k\le N\\(k,27)=9}}\sum_{\substack{0\le h<k\\(h,k)=1}} e^{-2\pi i(n-2)h/k}\Omega_{(h,k)}E^{(1)}(I)\\
&=: P(n)+Q
\end{aligned}
$$

with

$$
P(n)=\Bigg(\frac{1}{3}\sum_{\substack{0\le h<9\\(h,9)=1}} e^{-2\pi i(n-2)h/9}\Omega_{(h,9)}\Bigg)\sqrt{\frac{2}{9(n-1/24)}}\,e^{\frac{\pi}{9}\sqrt{\frac{2}{3}\left(n-\frac{1}{24}\right)}}
$$

and

$$
\begin{aligned}
|Q|&\le\sum_{\substack{18\le k\le N\\(k,27)=9}}\Bigg(\frac{1}{3}\sum_{\substack{0\le h<k\\(h,k)=1}} e^{-2\pi i(n-2)h/k}\Omega_{(h,k)}\Bigg)\sqrt{\frac{2}{k(n-1/24)}}\\
&\quad\times\left(e^{\frac{\pi}{k}\sqrt{\frac{2}{3}\left(n-\frac{1}{24}\right)}}+e^{-\frac{\pi}{k}\sqrt{\frac{2}{3}\left(n-\frac{1}{24}\right)}}\right)+\frac{2\sqrt{2}}{3}+11.563(n-1/24)^{(1/4)}\\
&\le\sum_{2\le l_2\le N}^{\lfloor\sqrt{2\pi(n-1/24)}/9\rfloor}\frac{\sqrt{k}}{3}\sqrt{\frac{2}{n-1/24}}\left(e^{\frac{\pi}{18}\sqrt{\frac{2}{3}\left(n-\frac{1}{24}\right)}}+1\right)\\
&\quad+\frac{2\sqrt{2}}{3}+11.563(n-1/24)^{(1/4)}\\
&\le\left(\frac{\sqrt{2\pi(n-1/24)}}{9}\right)^{\frac{3}{2}}\sqrt{\frac{2}{n-1/24}}\left(e^{\frac{\pi}{18}\sqrt{\frac{2}{3}\left(n-\frac{1}{24}\right)}}+1\right)\\
&\quad+\frac{2\sqrt{2}}{3}+11.563(n-1/24)^{(1/4)}\\
&\le 0.208(n-1/24)^{(1/4)}e^{\frac{\pi}{18}\sqrt{\frac{2}{3}\left(n-\frac{1}{24}\right)}}+11.771(n-1/24)^{(1/4)}+0.943.
\end{aligned}
$$

Similarly, applying (3.9), we deduce that

$$
S_4(A)=U_1+U_2,
$$

https://doi.org/10.4153/S0008414X24000683 Published online by Cambridge University Press

with

$$U_1 = \sum_{\substack{1\le k\le N\\(k,27)=27}} \sum_{\substack{0\le h<k\\(h,k)=1}} e^{-2\pi i n h/k}\omega_{(h,k)}^{(3)} \int_{\xi_{h,k}} z^{-\frac{1}{2}} e^{\frac{\pi}{12k}(\frac{1}{z}-z)} e^{2\pi n\rho} e^{-2\pi i n\varphi} d\varphi,$$

and

$$|U_2| \le 0.3536 e^{2\pi(n-1/24)\rho} N^{1/2}.$$

Here, $\omega_{(h,k)}^{(3)} := e^{\pi i(s(h,27l_3)+s(h,l_3)-3s(h,3l_3))}$.

For U_1, we have

$$\begin{aligned}
|U_1| &= \sum_{\substack{1\le k\le N\\(k,27)=27}} \sum_{\substack{0\le h<k\\(h,k)=1}} e^{-2\pi i(n)h/k}\omega_{(h,k)}^{(3)} \int_{\xi_{h,k}} z^{-\frac{1}{2}} e^{\frac{\pi}{12k}(\frac{1}{z}-z)} e^{2\pi n\rho} e^{-2\pi i n\varphi} d\varphi \\
&\le \left| \sum_{\substack{1\le k\le N\\(k,27)=27}} \sum_{\substack{0\le h<k\\(h,k)=1}} e^{-2\pi i(n)h/k}\omega_{(h,k)}^{(3)} \sqrt{\frac{2}{k(n-1/24)}} \cosh\left(\frac{\pi}{k}\sqrt{\frac{2}{3}\left(n-\frac{1}{24}\right)}\right)\right| \\
&\quad + \left| \sum_{\substack{1\le k\le N\\(k,27)=27}} \sum_{\substack{0\le h<k\\(h,k)=1}} e^{-2\pi i(n)h/k}\omega_{(h,k)}^{(3)} E^{(1)}(I)\right| \\
&\le \sum_{\substack{1\le k\le N\\(k,27)=27}} \sum_{\substack{0\le h<k\\(h,k)=1}} \sqrt{\frac{2}{k(n-1/24)}} \left(e^{\frac{\pi}{18}\sqrt{\frac{2}{3}(n-\frac{1}{24})}}+1\right) + 34.698(n-1/24)^{(1/4)} \\
&\le \sum_{l_3=1}^{\lfloor\sqrt{2\pi(n-1/24)}/27\rfloor} \sqrt{\frac{54l_3}{n-1/24}} \left(e^{\frac{\pi}{18}\sqrt{\frac{2}{3}(n-\frac{1}{24})}}+1\right) + 34.698(n-1/24)^{(1/4)} \\
&\le \left(\frac{\sqrt{2\pi(n-\frac{1}{24})}}{27}\right)^{\frac{3}{2}} \sqrt{\frac{54}{n-1/24}} \left(e^{\frac{\pi}{18}\sqrt{\frac{2}{3}(n-\frac{1}{24})}}+1\right) + 34.698(n-1/24)^{(1/4)} \\
&\le 0.208(n-1/24)^{(1/4)} e^{\frac{\pi}{18}\sqrt{\frac{2}{3}(n-\frac{1}{24})}} + 34.897(n-1/24)^{(1/4)}.
\end{aligned}$$

3.4 Signs of the Fourier coefficients of $g(q) - 3h(q)$

It follows from (3.1) and (3.2) that

$$g(q) - 3h(q) = \sum_{n=1}^{\infty} (a(n) - 3b(n)) q^n.$$

From (3.3), (3.4) and the results in Subsections 3.2 and 3.3, we derive that

$$\begin{aligned}
a(n) - 3b(n) &= S_1(A) + S_2(A) + S_3(A) + S_4(A) - 3(S_1(B) + S_2(B) + S_3(B) + S_4(B)) \\
&= T_1 - 3R_1 + T_2 - 3R_2 + S_2(A) + S_3(A) - 3S_2(B) \\
&\quad - 3P(n) - 3P_2 - 3Q + U_1 + U_2 - 3S_4(B) \\
&=: -3P(n) + E(n)
\end{aligned}$$

https://doi.org/10.4153/S0008414X24000683 Published online by Cambridge University Press

with

$$-3P(n) = -\left(\sum_{\substack{0\le h<9\\(h,9)=1}} e^{-2\pi i(n-2)h/9}\Omega_{(h,9)}\right)\sqrt{\frac{2}{9(n-1/24)}}\,e^{\frac{\pi}{9}\sqrt{\frac{2}{3}\left(n-\frac{1}{24}\right)}}.$$

From Subsctions 3.2 and 3.3, we know that

$$\begin{cases}
|T_1-3R_1| &\le 11.225(n-1/24)^{(1/4)}e^{\frac{\pi}{18}\sqrt{\frac{2}{3}\left(n-\frac{1}{24}\right)}}+80.601(n-1/24)^{(1/4)},\\
|T_2| &\le 7.560\times 10^{10}e^{2\pi(n-1/24)\rho}N^{\frac{1}{2}} = 3.254\times 10^{11}(n-1/24)^{(1/4)}\\
|R_2| &\le 1.435\times 10^{13}e^{2\pi i(n-1/24)\rho}N^{\frac{1}{2}} = 6.176\times 10^{13}(n-1/24)^{(1/4)},\\
|S_2(A)| &\le \frac{4}{\sqrt{3}}e^{0.969}e^{2\pi(n-\frac{1}{24})\rho}N^{\frac{1}{2}} = 26.193(n-1/24)^{(1/4)},\\
|S_3(A)| &\le 4\sqrt{3}e^{-2.047}e^{2\pi(n-\frac{1}{24})\rho}N^{\frac{1}{2}} = 3.850(n-1/24)^{(1/4)},\\
|S_2(B)| &\le \frac{4}{3\sqrt{3}}e^{1.710}e^{2\pi(n-\frac{1}{24})\rho}N^{\frac{1}{2}} = 18.318(n-1/24)^{(1/4)},\\
|P_2| &\le 0.119e^{2\pi(n-1/24)\rho}N^{\frac{1}{2}} = 0.513(n-1/24)^{(1/4)},\\
|Q| &\le 0.208(n-1/24)^{(1/4)}e^{\frac{\pi}{18}\sqrt{\frac{2}{3}\left(n-\frac{1}{24}\right)}}+11.771(n-1/24)^{(1/4)}+0.943,\\
|U_1| &\le 0.208(n-1/24)^{(1/4)}e^{\left(\frac{\pi}{18}\sqrt{\frac{2}{3}\left(n-\frac{1}{24}\right)}\right)}+34.897(n-1/24)^{(1/4)},\\
|U_2| &\le 0.3536e^{2\pi(n-1/24)\rho}N^{\frac{1}{2}} = 1.522(n-1/24)^{(1/4)},\\
|S_4(B)| &\le 4e^{-6.077}e^{2\pi(n-\frac{1}{24})\rho}N^{\frac{1}{2}} = 0.040(n-1/24)^{(1/4)},
\end{cases}$$

so that

$$\begin{aligned}
|E(n)| &\le |T_1-3R_1|+|T_2|+3|R_2|+|S_2(A)|+|S_3(A)|+3|S_2(B)|\\
&\quad +3|P_2|+3|Q|+|U_1|+|U_2|+3|S_4(B)|\\
&\le 12.057(n-1/24)^{(1/4)}e^{\frac{\pi}{18}\sqrt{\frac{2}{3}\left(n-\frac{1}{24}\right)}}+3.772\\
&\quad +\big(3.254\times 10^{11}+3\times 6.176\times 10^{13}+26.193+3.850+3\times 18.318+\\
&\qquad 3\times 0.513+3\times 12.969+12.969+1.522+3\times 0.040\big)\left(n-\frac{1}{24}\right)^{\frac{1}{4}}\\
&\le 12.057(n-1/24)^{(1/4)}e^{\frac{\pi}{18}\sqrt{\frac{2}{3}\left(n-\frac{1}{24}\right)}}+1.857\times 10^{14}\left(n-\frac{1}{24}\right)^{\frac{1}{4}}.
\end{aligned}$$

It is observed that the signs of the main term $P(n)$ are determined by

$$\sum_{\substack{0\le h<9\\(h,9)=1}} e^{-2\pi i(n-2)h/9}\Omega_{(h,9)}$$

with $\Omega_{(h,9)} := e^{\pi i(s(h,9l_2)+s(h,l_2)-s(h,3l_1)-2s(3h,l_2))}$. After calculations, we have

$$\begin{cases}
\Omega_{(1,9)} = e^{\frac{25}{54}\pi i}, \quad \Omega_{(2,9)} = e^{\frac{11}{54}\pi i},\\
\Omega_{(4,9)} = e^{-\frac{11}{54}\pi i}, \quad \Omega_{(5,9)} = e^{\frac{11}{54}\pi i},\\
\Omega_{(7,9)} = e^{-\frac{11}{54}\pi i}, \quad \Omega_{(8,9)} = e^{-\frac{25}{54}\pi i}.
\end{cases}$$

https://doi.org/10.4153/S0008414X24000683 Published online by Cambridge University Press

Therefore,

$$\sum_{\substack{0\le h<9\\(h,9)=1}} e^{-2\pi i(n-2)h/9}\Omega_{(h,9)} = \begin{cases} 2\cos(23\pi/54)+4\cos(49\pi/54) & \text{if } n\equiv 0 \pmod 9\\ 2\cos(35\pi/54)+4\cos(37\pi/54) & \text{if } n\equiv 1 \pmod 9\\ 2\cos(25\pi/54)+4\cos(11\pi/54) & \text{if } n\equiv 2 \pmod 9\\ 2\cos(49\pi/54)+4\cos(13\pi/54) & \text{if } n\equiv 3 \pmod 9\\ 2\cos(37\pi/54)+4\cos(\pi/54) & \text{if } n\equiv 4 \pmod 9\\ 2\cos(11\pi/54)+4\cos(47\pi/54) & \text{if } n\equiv 5 \pmod 9\\ 2\cos(13\pi/54)+4\cos(23\pi/54) & \text{if } n\equiv 6 \pmod 9\\ 2\cos(\pi/54)+4\cos(35\pi/54) & \text{if } n\equiv 7 \pmod 9\\ 2\cos(47\pi/54)+4\cos(25\pi/54) & \text{if } n\equiv 8 \pmod 9. \end{cases}$$

It is easily computed that

$$\left|\sum_{\substack{0\le h<9\\(h,9)=1}} e^{-2\pi i(n-2)h/9}\Omega_{(h,9)}\right| \ge 0.20142$$

for $n\in\mathbb{N}$. When $n\ge 22472$,

$$\begin{aligned} |-3P(n)| &= \left|\left(\sum_{\substack{0\le h<9\\(h,9)=1}} e^{-2\pi i(n-2)h/9}\Omega_{(h,9)}\right)\sqrt{\frac{2}{9(n-1/24)}}\,e^{\frac{\pi}{9}\sqrt{\frac{2}{3}\left(n-\frac{1}{24}\right)}}\right| \\ &\ge 0.20142\sqrt{\frac{2}{9(n-1/24)}}\,e^{\frac{\pi}{9}\sqrt{\frac{2}{3}\left(n-\frac{1}{24}\right)}} \\ &> 12.057(n-1/24)^{(1/4)}e^{\frac{\pi}{18}\sqrt{\frac{2}{3}\left(n-\frac{1}{24}\right)}}+1.857\times 10^{14}\left(n-\frac{1}{24}\right)^{\frac{1}{4}} \\ &\ge |E(n)|. \end{aligned}$$

From this, we obtain that

$$\begin{cases} a(n)-3b(n)>0 & \text{if } n\equiv 0,1,5,8 \pmod 9\\ a(n)-3b(n)<0 & \text{if } n\equiv 2,3,4,6,7 \pmod 9 \end{cases} \tag{3.11}$$

holds for $n\ge 22472$. Calculating the first 22471 coefficients of $g(q)-3h(q)$ by using the software *Mathematica*, we find that (3.11) is also true for $467\le n\le 22471$.

3.5 Proof of Theorem 1.1

It follows from the identities in [6, eqs.(2.10) and (2.11)] that

$$\sum_{n=1}^{\infty} M(0,1,1;9;n)q^n - \frac{1}{3(q;q)_\infty} = \frac{2}{3}(g(q)-3h(q)),$$

and

$$\sum_{n=1}^{\infty} M(2,3,4;9;n)q^n - \frac{1}{3(q;q)_\infty} = -\frac{1}{3}(g(q)-3h(q)),$$

https://doi.org/10.4153/S0008414X24000683 Published online by Cambridge University Press

so that

$$M(0,1,1;9;n) - \frac{p(n)}{3} = \frac{2}{3}(a(n) - 3b(n)),$$
$$M(2,3,4;9;n) - \frac{p(n)}{3} = -\frac{1}{3}(a(n) - 3b(n)).$$

It is known from Subsection 3.4 that (3.11) holds for $n \geq 467$. Hence, when $n \geq 467$,

$$\begin{cases} M(0,1,1;9;n) > \frac{p(n)}{3}, & \text{if } n \equiv 0,1,5,8 \pmod 9, \\ M(0,1,1;9;n) < \frac{p(n)}{3}, & \text{if } n \equiv 2,3,4,6,7 \pmod 9, \\ M(2,3,4;9;n) < \frac{p(n)}{3}, & \text{if } n \equiv 0,1,5,8 \pmod 9, \\ M(2,3,4;9;n) > \frac{p(n)}{3}, & \text{if } n \equiv 2,3,4,6,7 \pmod 9. \end{cases}$$

This finishes the proof of Theorem 1.1.

Acknowledgements The authors would like to thank the referee for his/her meticulously thorough reading of this paper and for numerous helpful comments and suggestions which greatly improved the exposition of this paper. This work was partially supported by the Natural Science Foundation of Changsha (Grant No. kq2208251).

References

[1] G. A. Andrews and R. Lewis, *The ranks and cranks of partitions moduli 2, 3, and 4*. J. Number Theory **85**(2000), no. 1, 74–84.
[2] G. E. Andrews and B. C. Berndt, *Ramanujan's lost notebook*. Part III, Springer, New York, 2012.
[3] G. E. Andrews and F. G. Garvan, *Dyson's crank of a partition*. Bull. Amer. Math. Soc. (N.S.) **18**(1988), no. 2, 167–171.
[4] T. M. Apostol, *Modular functions and Dirichlet series in number theory*, Graduate Texts in Mathematics, Vol. 41, second edition, Springer-Verlag, New York, 1990.
[5] A. O. L. Atkin and P. Swinnerton-Dyer, *Some properties of partitions*. Proc. London Math. Soc. (3) **4**(1954), 84–106.
[6] Z. S. Aygin and S. H. Chan, *Combinations of ranks and cranks of partitions moduli 6, 9 and 12 and their comparison with the partition function*. Ann. Comb. **23**(2019), nos. 3–4, 489–509.
[7] N. Borozenets, *Deviation of the rank and crank modulo 11*. Preprint, 2023, arXiv:2305.08751v3.
[8] K. Bringmann and B.V. Pandey, *Biases among classes of rank-crank partitions (mod11)*. J. Math. Anal. Appl. **536**(2024), no. 2, Paper No. 128221, 17 pp.
[9] O.-Y. Chan, *Some asymptotics for cranks*. Acta Arith. **120**(2005), no. 2, 107–143.
[10] S. H. Chan and R. Mao, *Inequalities for ranks of partitions and the first moment of ranks and cranks of partitions*. Adv. Math. **258**(2014), 414–437.
[11] N. Chen, S. Chern, Y. Fan and E. X. W. Xia, *Some generating functions and inequalities for the Andrews-Stanley partition functions*. Proc. Edinb. Math. Soc. (2) **65**(2022), no. 1, 120–135.
[12] F. J. Dyson, *Some guesses in the theory of partitions*. Eureka **8**(1944), 10–15.
[13] Y. Fan, E. X. W. Xia and X. Zhao, *New equalities and inequalities for the ranks and cranks of partitions*. Adv. in Appl. Math. **146**(2023), Paper No. 102486, 37 pp.
[14] D. M. Kane, *Resolution of a conjecture of Andrews and Lewis involving cranks of partitions*. Proc. Amer. Math. Soc. **132**(8)(2004), 2247–2256.
[15] S.-Y. Kang, *Mock Jacobi forms in basic hypergeometric series*. Compos. Math. **145**(2009), no. 3, 553–565.
[16] R. Mao, *Ranks of partitions modulo 10*. J. Number Theory **133**(2013), no. 11, 3678–3702.
[17] H. Rademacher, *On the expansion of the partition function in a series*. Ann. of Math. (2) 44(1943), 416–422.

https://doi.org/10.4153/S0008414X24000683 Published online by Cambridge University Press

[18] M. J. Schlosser and N.H. Zhou, *On the infinite Borwein product raised to a positive real power*. Ramanujan J. **61**(2023), no. 2, 515–543.
[19] O. X. M. Yao, Personal communications, 2023.
[20] N. H. Zhou, *On the distribution of rank and crank statistics for integer partitions*. Res. Number Theory **5**(2019), no. 2, Paper No. 18, 8 pp.
[21] N. H. Zhou, *On the distribution of the rank statistic for strongly concave compositions*. Bull. Aust. Math. Soc. **100**(2019), no. 2, 230–238.

School of Mathematics and Statistics, HNP-LAMA, Central South University, Changsha 410083, Hunan, People's Republic of China
e-mail: yuhe123456@foxmail.com yuhelingyun@foxmail.com

School of Mathematics and Statistics, HNP-LAMA, Central South University, Changsha 410083, Hunan, People's Republic of China
e-mail: 232111035@csu.edu.cn

School of Mathematics, Hangzhou Normal University, Hangzhou 311121, People's Republic of China
e-mail: 21caojian@hznu.edu.cn

https://doi.org/10.4153/S0008414X24000683 Published online by Cambridge University Press