Ergod. Th. & Dynam. Sys. (2006), **26**, 19–44
DOI: 10.1017/S0143385705000386
© *2005 Cambridge University Press*
Printed in the United Kingdom

Dynamics of injective quasi-contractions

JULIEN BRÉMONT

Université Paris 12, Département de Mathématiques,
61, Avenue du Général de Gaulle, 94010 Créteil cedex, France
(e-mail: bremont@univ-paris12.fr)

(*Received* 12 *May* 2005 *and accepted* 21 *July* 2005)

Abstract. We study injective locally contracting maps of the interval. After giving an upper-bound on the number of ergodic components, we show that generically finitely many periodic orbits attract the whole dynamics and that this picture is stable under perturbation. In relation to the problem of maximizing measures for regular maps, we then consider a class of probability measures on the circle invariant by $\times p$ generalizing the family of Sturm measures and show its generic periodic character. We detail the structure of order-preserving locally contracting maps on the circle. The rotation number is shown to be generically rational and the transformations having a given rational rotation number are shown explicitly. We also count the periodic attractors. We then deduce the existence of measurable conjugacies with three-interval exchange transformations in non-periodic cases for a model with three pieces on the interval.

1. *Introduction*

This paper is motivated by the study of maximizing measures. This optimization problem in ergodic theory is naturally formulated in a topological setting. Fix a compact metric space X endowed with a continuous transformation T and the set $\mathcal{M}_T(X)$ of T-invariant Borel probability measures. Given some real continuous function f, one considers the variational problem:

$$\beta(f) = \sup\left\{\int f\, d\mu \;\middle|\; \mu \in \mathcal{M}_T(X)\right\}.$$

One then asks for the measures realizing the maximum and, more specifically, for their stochastic properties. This problem is related, via Birkhoff's theorem, to finding points $x \in X$ ensuring the best growth of the ergodic sums $\left(\sum_{0\le k\le n-1} f(T^k x)\right)_{n\ge 0}$. For a detailed presentation of this topic and an introduction to the general theory, see Bousch and Mairesse [**3**], Conze and Guivarc'h [**8**] or Jenkinson [**13**].

A central question in this field concerns the role of periodic measures for regular maps when the dynamical system is of expansive type. A typical context is $X = \mathbb{R}/\mathbb{Z}$ with $Tx = 2x \bmod(1)$ and a conjecture is that *for at least a dense open set in the set of Lipschitz functions, the maximizing measure is unique and supported by a periodic orbit.*

https://doi.org/10.1017/S0143385705000386 Published online by Cambridge University Press

See Contreras *et al.* [**7**] and Hunt and Yuan [**12**] for weak forms of this conjecture. An important example is given in the work of Bousch [**2**], considering the one-parameter family $(f_t)_{t\in[0,1)}$ with $f_t(x) = \cos 2\pi(x+t)$. The result is that for all t the maximizing measure of f_t is unique and Sturmian; moreover, except for t in a zero-dimensional set, this measure is periodic. The proof fundamentally uses the following reduction lemma (valid in a more general context) due to Conze and Guivarc'h (see [**8**]) and Mané (see [**16**]).

LEMMA 1.1. *Let $f \in \mathrm{Lip}(X)$. There exist Lipschitz maps φ and r such that:*

$$f = \beta(f) + (\varphi \circ T - \varphi) - r \quad \textit{with} \quad r \geq 0 \quad \textit{and} \quad r(\cdot)r(\cdot + 1/2) = 0.$$

The conditions on r provide a characterization of the maximizing measures via their support. More precisely, one checks that μ is maximizing for f if and only if $\mathrm{supp}(\mu) \subset r^{-1}\{0\}$. In the study of the cosine family, Bousch shows that for each f_t there exist only two points, diametrically opposed, verifying $r(x) = r(x+1/2) = 0$ and thus that $r^{-1}\{0\}$ is a semi-circle $[\alpha, \alpha + 1/2]$. Finally the argument is that T-invariant Borel probability measures with support in such domains are known to be exactly the family of Sturm measures (see Bullett and Sentenac [**6**]).

In a dynamical approach to the above conjecture, a first step is to consider the case when r admits only finitely many x satisfying the equalities $r(x) = r(x+1/2) = 0$. Then $r^{-1}\{0\}$ is some *finite flower* F of total length $1/2$ and is not necessarily a semi-circle. Omitting boundary questions, the problem of describing the class of T-invariant measures with support in F can be reformulated using the right-inverse η of T defined by $T \circ \eta = \mathrm{Id}$ and $\mathrm{Im}(\eta) = F$. Since $\mu = T\mu$ and $\mathrm{Supp}(\mu) \subset \bar{F}$ is equivalent to $\mu = \eta\mu$, one is lead to studying the dynamics of η which appears to be locally affine and $1/2$-contracting. This class of maps can be seen as a contracting variant of the family of interval exchange transformations, namely the bijective piecewise translations of the interval, introduced by Keane [**15**]. These considerations are the main reason for the present work.

Enlarging the perspective, η belongs to the more general class of injective and locally contracting maps of the interval $[0, 1)$, or *injective quasi-contractions*. Fixing an integer $m \geq 1$ and $0 = D_0 < D_1 < \cdots < D_m < D_{m+1} = 1$, a quasi-contraction T of the interval is defined on each $[D_i, D_{i+1})$ by $T(x) = \Gamma_i(x)$, where each Γ_i is γ-Lipschitzian for some $0 < \gamma < 1$. These maps occur in various contexts, for instance in differential dynamics (see Alsed *et al.* [**1**] and Gambaudo and Tresser [**11**] when $m = 1$) and in physics, since heuristically contraction properties ensure some form of stability. Extending the work of Feely and Chua [**9**], from signal theory, on the map $x \longmapsto \{\gamma x + \alpha\}$, where $0 < \gamma < 1$, Bugeaud and Conze [**5**] have shown that the dynamics can be described via a concatenation algorithm similar to the Farey tree. Fixing γ, then for almost all α the transformation admits a unique periodic attractor. In the non-periodic cases, topological semi-conjugacies are then deduced with irrational rotations. Another point was to extend these results to general locally contracting and order-preserving maps of the circle.

In this paper, we study general injective quasi-contractions with $m+1$ pieces, $m \geq 1$, on the interval. The injectivity requirement is discussed in the next section. We first provide a general analysis on the structure of an injective quasi-contraction which allows one to bound from above the number of ergodic components by $2m$. We next prove

https://doi.org/10.1017/S0143385705000386 Published online by Cambridge University Press

a degeneracy result: for an open set of full Lebesgue measure in the set of parameters, the corresponding maps admit finitely many periodic orbits and the union of the attraction domains is the whole interval. We mention as a general fact that dilating maps are analyzed via inverse branches considered through contracting transfer operators. In this context we naturally look, in an elementary way, at forward images. Our approach is simple and geometric and seems better suited than the usual coding point of view (adopted when $m = 1$ for instance in [**11**] in terms of Sturm sequences) for situations far from rotations. This allows one to slant the orbital complexity which is that of an interval exchange transformation. In fact we show that generically the complexity stops growing in finite time. We complete this section with a perturbation result.

In §3 we solve the problem of invariant measure discussed above. More generally we consider the Borel probability measures on the circle invariant by $Tx = px \bmod(1)$ and with support in some *finite flower*. We prove that, in general, such measures are periodic and that the number of ergodic measures is less than the number of petals of the flower.

Finally, we consider a class of injective quasi-contractions that are order-preserving on the circle $\mathbb{R}/\mathbb{Z}$. After observing the validity of the Poincaré theory on the rotation number, we first deduce from §2 that this number is generically rational. We then completely describe the transformations having a given rational number explicitly and count the number of periodic attractors. As a corollary, we study a model of injective quasi-contractions on the interval with three pieces and show in the non-periodic cases the existence of measurable conjugacies with interval exchange transformations.

2. *Injective quasi-contractions and generic properties*

Let us define injective quasi-contractions on the interval $[0, 1)$.

Definition 2.1.

(1) Let $m \geq 0$ and $\mathcal{D} = \{(D_i)_{1\leq i\leq m} \mid 0 < D_1 < \cdots < D_m < 1\} \subset \mathbb{R}^m$. To $D \in \mathcal{D}$ associate the partition $\mathcal{P}(D)$ in left-closed and right-open intervals. Set $D_0 = 0$ and $D_{m+1} = 1$.

(2) Fix $0 \leq \gamma < 1$ and suppose that injective γ-Lipschitz maps $\Gamma_i : [0, 1] \longrightarrow [0, 1]$, for $0 \leq i \leq m$, are given. For $D \in \mathcal{D}$ define a right-continuous map $T : [0, 1) \to [0, 1)$ by

$$Tx = \Gamma_i(x) \quad \text{for } x \in [D_i, D_{i+1}),$$

under the open condition on $D \in \mathbb{R}^m$: $\mathcal{C} = \mathcal{D} \cap \{T \text{ is injective and } TD_i^{\pm} \neq TD_j^{\pm}, \text{ for } i \neq j, 1 \leq i \leq m, 0 \leq j \leq m+1\}$, where we set $Tx^- = T(x^-)$ for all $x \in [0, 1)$.

(3) If $n \geq 1$ and $(\eta_j)_{1\leq j\leq n} \in \{0, \ldots, m\}^n$, write $\Gamma_{\eta_n,\ldots,\eta_1}$ for $\Gamma_{\eta_n} \circ \cdots \circ \Gamma_{\eta_1}$.

(4) The sequence $(X_n = T^n[0, 1))_{n\geq 0}$ is non-increasing. Set $X_\infty = \bigcap_{n\geq 0} X_n$.

(5) Set $E = [0, 1)\backslash T[0, 1)$. This set is a disjoint union of intervals: $E = \bigcup_{1\leq i\leq m} F_i$.

In the whole text, we say that $x^{\pm}$ is periodic if, for some $q \geq 1$, one has $T^q x^{\pm} = x^{\pm}$. We show that in a generic sense, measurably and topologically, the dynamics of an injective quasi-contraction is degenerated. Denote by λ_m the Lebesgue measure in $\mathbb{R}^m$ and let $\lambda = \lambda_1$.

https://doi.org/10.1017/S0143385705000386 Published online by Cambridge University Press

THEOREM 2.2.

(1) *For any $D \in \mathcal{C}$, the associated quasi-contraction T admits at most $2m$ ergodic components.*

(2) *There is an open set $\Omega \subset \mathcal{C}$ with $\lambda_m(\mathcal{C}\backslash\Omega) = 0$ such that, for $D \in \Omega$, the corresponding T admits a finite stable partition $\mathcal{P}(D')$ in intervals finer than $\mathcal{P}(D)$: every atom of $\mathcal{P}(D')$ is sent by T into another atom of $\mathcal{P}(D')$. As a result and more precisely we can state the following.*

- *The injective quasi-contraction T has $1 \leq n(T) \leq 2m$ attractive periodic orbits. If all Γ_i are increasing then $n(T) \leq m+1$.*
- *The domain of attraction of each periodic orbit is a finite union of intervals with positive length and the union of the domains is $[0, 1)$.*
- *There exists a constant $C(T)$ such that, for all $x \in [0, 1)$, some periodic point y satisfies, for all $n \geq 0$, $\mathrm{dist}(T^n x, T^n y) \leq C(T)\,\gamma^n$.*

Remark. The upper-bound $2m$ on $n(T)$ is not expected to be optimal, whereas the value $m+1$ is a natural candidate. It is easily reached, for instance, if $\Gamma_i[D_i, D_{i+1}) \subset [D_i, D_{i+1})$ for all $0 \leq i \leq m$.

2.1. *Abstract structure of general injective quasi-contractions.* We prove the first part of Theorem 2.2 by giving a general description of $X_n = T^n[0, 1)$, for $n \geq 1$. We proceed inductively.

- For $n = 1$, let $\mathcal{G}(1)$ be the set of disjoint intervals $(T[D_i, D_{i+1}))_{0\leq i\leq m}$. Denote by $\mathcal{R}_1$ the set of D_i in X_1. For $i \in \mathcal{R}_1$, let U_i^1 be the interval of $\mathcal{G}(1)$ containing D_i and $U_{i,\pm}^1$ its two halves, putting D_i on the right-hand interval (the left-hand interval may be empty). Note that if, for instance, $U_{i,+}^1$ contains D_{i+1}, we replace this interval by $[D_i, D_{i+1})$; similar considerations hold on the left-hand side of D_i. Then the elements of $\mathcal{G}(1)$, with the exception of the $(U_i^1)_{i\in\mathcal{R}_1}$, are sent into one another (without being cut) by T. For $i \in \mathcal{R}_1$, both of the intervals $U_{i,+}^1$ and $U_{i,-}$ have the same properties. Let $V_{i,\pm}^1$ be the two intervals of $\mathcal{G}(1)$ such that $TU_{i,\pm}^1 \subset V_{i,\pm}^1$.
  As a result and when considering the intervals different from the $(U_i^1)_{i\in\mathcal{R}_1}$, together with the $(U_{i,-}^1, U_{i,+}^1)_{i\in\mathcal{R}_1}$, the set $\mathcal{G}(1)$ has a natural graph structure, where an edge corresponds to applying T. Starting from any interval, the sequence of visited intervals is either eventually periodic or reaches some interval (only one by injectivity) which is sent in some U_i^1 and that has a non-empty intersection with both $U_{i,-}^1$ and $U_{i,+}^1$. One then stops the graph. Call next *type I graph* a subgraph consisting of a periodic limit cycle together with its basin of attraction and *type II graph* a subgraph where all intervals end in some interval which is sent in a U_i, with $i \in \mathcal{R}_1$, and has a non-empty intersection with $U_{i,-}^1$ and $U_{i,+}^1$. Denote by $\mathcal{T}_I(1)$ and $\mathcal{T}_{II}(1)$ the set of type I graphs and type II graphs at time 1, respectively.
- For $n \geq 1$, the set of disjoint intervals forming X_{n+1} is denoted by $\mathcal{G}(n+1)$ and it is deduced from $\mathcal{G}(n)$ as follows. First, replace all intervals of $\mathcal{G}(n)$ by their images by T. Observe that the attracting limit cycle of a type I graph remains globally unchanged. Next, let $\mathcal{R}_{n+1} \subset \mathcal{R}_n$ be the subset of discontinuities in X_{n+1}. For $i \in \mathcal{R}_{n+1}$, let $U_i^{n+1} = U_{i,-}^{n+1} \cup U_{i,+}^{n+1}$ be the interval containing D_i, with the

https://doi.org/10.1017/S0143385705000386 Published online by Cambridge University Press

same conventions as above. Let $V_{i,\pm}^{n+1}$ be the two intervals in $\mathcal{G}(n+1)$ such that $TU_{i,\pm}^{n+1} \subset V_{i,\pm}^{n+1}$. Then $\mathcal{G}(n+1)$ is defined as before with graphs of types I and II at time n, respectively, denoted by $\mathcal{T}_I(n+1)$ and $\mathcal{T}_{II}(n+1)$.

We now make the following observations.

- The sequence of sets $(\mathcal{R}_n)_{n\geq 1}$ is non-increasing and is therefore eventually constant to some limit set written as $\mathcal{R}$. Denote also by $\mathcal{R}' \subset \mathcal{R}$ the elements D_i such that $U_{i,-}^n \neq \emptyset$ for all n. Recall that $U_{i,+}^n \neq \emptyset$ for $i \in \mathcal{R}$ and for all n. The set $\mathcal{R}'$ is also eventually constant. Let N be such that these properties hold for $n \geq N$, as well as $U_{i,-}^n = \emptyset$ for $i \in \mathcal{R}\backslash\mathcal{R}'$.
- Next, if some non-empty $U_{i,\epsilon}^n$ with $\epsilon \in \{\pm\}$ and $i \in \mathcal{R}$ branches on a type I graph, then observe that this is true for all $U_{i,\epsilon}^p$ with $p \geq n$, since the branching is made on images of the same type I graph. Thus the number of such (i,ϵ) is non-decreasing and as a consequence is eventually constant. Then denote by J the set of (i,ϵ) with $\epsilon \in \{\pm\}$ and $i \in \mathcal{R}$ such that $U_{i,\epsilon}^n$ do not branch on a type I graph. Let N be such that these properties hold for $n \geq N$.
- Each type I graph contracts to a periodic orbit laying in its closure. Note that if $m \geq 1$ then at least one of the boundaries of any interval of X_n has the form $T^k D_s^{\pm}$ with $s \notin \{0, m+1\}$. Thus the periodic orbit corresponding to each type I graph is exactly the ω-limit set of some $D_s^{\pm}$ with $s \notin \{0, m+1\}$. Such distinct periodic orbits require distinct discontinuities of this form. Also, if distinct type I graphs admit the same limit periodic orbit and since T is injective and each D_i, $1 \leq i \leq m$, is a true discontinuity for T, then one type I graph must contain some $U_{i,-}^n$ and the other one must contain the interval $U_{i,+}^n$. Thus the number of distinct type I graphs is bounded from above by $4m$. For the moment we only use that this number is bounded. Next, suppose that no other 'new' type I graph appears after time N.
- For large enough n we assert that no periodic point is contained in the closure of a type II graph. If not, a periodic orbit (say of length q) would be entirely contained in the closure of the family of type II graphs. Denote by $D_{i_1}, \ldots, D_{i_p}$ the discontinuities such that each corresponding type II graph contains a periodic point. Since the lengths of the intervals in $\mathcal{G}(n)$ containing the D_{i_j} tend to zero, as $n \to +\infty$, for some $(\epsilon_j)_{1\leq j\leq p} \in \{\pm\}^p$ the $D_{i_j}^{\epsilon_j}$ are periodic points in the periodic orbit. More precisely, either no D_{i_j} is a periodic point and then all $D_{i_j}^-$ are periodic, or all D_{i_j} are periodic, since T is right-continuous. As T is injective, all D_{i_j} belong to the closure of the boundary of $T^{N+q}[0,1)$. Then take $n \geq N+q$. In the first case, all $U_{i_1,-}^n$ are non-empty and obviously define a type I graph under iterations. In the second case, either all $U_{i_j,-}^n$ are empty, in which case the same holds with the $U_{i_j,+}^n$, or there exists $U_{i_{j_0},-}^n \neq \emptyset$ and then $U_{i_{j_0},+}^n = \{D_{i_{j_0}}\}$ since $D_{i_{j_0}}$ is a boundary point. However, the sequence of intervals starting at $U_{i_{j_0},+}^n$ defines a type I graph. In any case we get a contradiction. Suppose next that N is large enough so that type II graphs for time $n \geq N$ contain no periodic point.
- For $(i,\epsilon) \in J$, the ω-limit set of D_i^{ϵ} contains no periodic orbit. Indeed, $U_{i,\epsilon}^N$ branches on a type II graph. If the forward orbit of D_i^{ϵ} never falls in some $U_{i',\epsilon'}^N$ that branches in a type I graph, then the result holds since the orbit of D_i^{ϵ} is contained in type II

https://doi.org/10.1017/S0143385705000386 Published online by Cambridge University Press

graphs whose closures do not contain any periodic orbit. If on the other hand the forward orbit of D_i^ϵ falls in an interval $U_{i',\epsilon'}^N$ branching in a type I graph, consider the first time p when this holds. Up to taking forward images of D_i^ϵ (with indices in J), assume also that, between times 1 and p, the images of D_i^ϵ are at a positive distance for all D_j with $j \in \mathcal{R}$. Thus for large enough n and since the lengths of the intervals decrease, the interval $U_{i,\epsilon}^n$ will not be cut by forward iterations and will branch on a type I graph. This contradicts the definition of J.

As a result, the number of periodic orbits is less than or equal to $2m - \mathrm{card}(J)$. Next, Borel diffusive probability measures must be supported by type II graphs. We show that the number of ergodic ones is less than or equal to $\mathrm{card}(J)$. First, only branches starting from $V_{i,\epsilon}^N$ with $(i,\epsilon) \in J$ can be weighted by a T-invariant Borel probability measure. Following an idea of Keane [**15**], let $\mu_1, \ldots, \mu_q$ be q T-invariant ergodic diffusive Borel probability measures. Let $(A_i)_{1\le i\le q}$ be disjoint T-invariant Borel sets such that $\mu_i(A_j) = \delta_{i,j}$. Define $\mu = \frac{1}{q}(\mu_1 + \cdots + \mu_q)$ and fix $0 < \epsilon < \min\{1/q, 1/2\}$. Then introduce

$$B_i^n = \{P \in \mathcal{T}_{II}(n) \text{ with } \mu(A_i \cap P) \ge (1-\varepsilon)\mu(P), \mu(P) > 0\}.$$

By regularity of the diffusive measure μ, if n is large enough, then B_i^n is non-empty. Moreover, setting $Y^n = \left(\bigcup_{P\in\mathcal{T}_{II}(n)} P\right) \cup \setminus \bigcup_{i=1}^q B_i^n$, we have

$$\mu(A_i \Delta B_i^n) < \varepsilon \quad \text{for } 1 \le i \le q \quad \text{and} \quad \mu(Y^n) < \varepsilon.$$

Now let $P \in B_i^n$. Using the T-invariance of A_i, we get $\mu(T^{-1}P \cap A_i) = \mu(T^{-1}(P \cap A_i)) = \mu(P \cap A_i) \ge (1-\varepsilon)\mu(P) = (1-\varepsilon)\mu(T^{-1}P)$. As a result, at least one element of the type II graph where P lays that branches to P also belongs to B_i^n. Repeating this procedure in reverse, at least one element of the form $V_{i',\epsilon'}^n$, with $(i',\epsilon') \in J$, belongs to B_i^n. This element does not belong to any other B_k^n, since otherwise the fact that $A_i \cap A_k = \emptyset$ implies $\mu(V_{i',\epsilon'}^n) \ge 2(1-\varepsilon)\mu(V_{i',\epsilon'}^n)$, whereas $\mu(V_{i',\epsilon'}^n) > 0$ and $\varepsilon < 1/2$. Thus $q \le \mathrm{card}(J)$.

This concludes the proof of the first part of the theorem. □

2.2. *Proof of the second part of Theorem 2.2.* In the context of Theorem 2.2, we begin with an elementary analysis suggesting our strategy. Denote by A^{o} the interior of a set A.

LEMMA 2.3.

(1) *We have* $\lim_{n\to+\infty} \lambda(X_n) = \lambda(X_\infty) = 0$.

(2) *For* $n \ge 0$, $[0,1) = T^{n+1}[0,1) \bigcup_{0\le i\le n} T^i E$ *disjointly. Also* $(T^n F_i)_{n\ge 0, 1\le i\le m+2}$ *are disjoint.*

(3) *Set* $\Omega = \{D \in \mathcal{C} \mid \forall 1 \le i \le m, \exists (k_i, l_i) \in \mathbb{N} \times \{1, \ldots, m+2\}$ *with* $D_i \in (T^{k_i} F_{l_i})^{\mathrm{o}}\}$. *Then* Ω *is open. If* $D \in \Omega$, *the conclusion of Theorem 2.2 holds.*

Proof.

(1) We have $\lambda(T^n[0,1)) \to \lambda(X_\infty)$ and $X_\infty = \bigcap_{n\ge 1} T^n[0,1) = \bigcap_{n\ge 0} T(T^n[0,1)) \subset TX_\infty$. Therefore,

$$\lambda(X_\infty) \le \lambda(TX_\infty) \le \sum_{0\le i\le m} \int_{X_\infty \cap [D_i, D_{i+1})} |\Gamma_i'(t)|\, dt \le \gamma\, \lambda(X_\infty).$$

Consequently, $\lambda(X_\infty) = 0$.

https://doi.org/10.1017/S0143385705000386 Published online by Cambridge University Press

(2) As $[0,1) = T[0,1) \cup E$ disjointly, we get recursively $[0,1) = T^{n+1}[0,1) \bigcup_{0\le i\le n} T^i E$ disjointly, since T is injective. Next, if $T^n F_i \cap T^{n'} F_{i'} \neq \emptyset$ with $n \ge n'$, then $n = n'$; otherwise $T^{n-n'} Fi \cap F_{i'} \neq \emptyset$ (as T is injective) and $F_{i'} \subset [0,1)\backslash T[0,1)$. Then $F_i \cap F_{i'} \neq \emptyset$ and thus $i = i'$.

(3) If $D \in \Omega$, observe that, for $1 \le i \le m$, $D_i \notin \{T^n D_j^{\pm} \mid 0 \le j \le m+1, 1 \le n \le K\}$ with $K = \max_{1\le i\le m} k_i$. Thus the boundaries of all $T^{k_i} F_{l_i}$ are locally continuous functions of D and thus Ω is open. Now fix $D \in \Omega$ with a collection of (k_i, l_i) and set $D' = \{T^{-n} D_i \mid 1 \le i \le m, 0 \le n < k_i\}$. The elements of D' are all distinct. Denote by $\mathcal{P}(D') = (U_k)$ the corresponding finite partition of $[0,1)$. For $1 \le i \le m$ let U_{n_i} and U_{m_i} be the components on the left-hand side and on the right-hand side of D_i, respectively.

Taking any U_k, we now show that $(TU_k)^{\mathrm{o}}$ is contained in another atom of $\mathcal{P}(D')$. First note that $(TU_k)^{\mathrm{o}}$ is an interval, since U_k contains no D_i in its interior. Next, if some $T^{-l} D_j \in (TU_k)^{\mathrm{o}}$ with $0 \le l < k_j$, then $l \le k_j - 2$ as $T^{-k_j} D_j$ does not exist. However, this would imply $T^{-l-1} D_j \in (U_k)^{\mathrm{o}}$, contradicting the definition of D'.

Writing $U_k = [x, y)$, there exist $0 \le i, j \le m+1$, $l_i \ge 0$, $l_j \ge 0$ with $x = T^{-l_i} D_i$ and $y = T^{-l_j} D_j$, setting $l_0 = 0$ and $l_{m+1} = 0$. Except for if $m = 0$, in which case T contracts to a unique fixed point, either i or j is in $\{1, \dots, m\}$. Since T is continuous at any $T^{-l} D_k$ with $l \ge 1$ and since $T D_i \notin D'$ for $0 \le i \le m+1$, U_k is never cut under iterations of T and is a positive distance from the boundary of the atom of $\mathcal{P}(D')$ where it lies for times larger than $\max\{k_i, k_j\} + 1$. Writing U_{i_l} for the atom containing $T^l U_k$, the sequence (i_l) contains a term among the $\{n_i, m_i, \text{ for } 1 \le i \le m\}$ and is necessarily preperiodic. Observe that the periodic part of $(i_l)_{l\ge 0}$ begins at a time $\le K$ and the period is $\le 2mK$, where $K = \max_{1\le i\le m} k_i$. An embedded compact argument gives that U_k is attracted exponentially fast to a periodic orbit. Also the ω-limit set of any x is that of some $D_i^{\pm}$, $1 \le i \le m$. Hence there are at most $2m$ periodic orbits. If all Γ_i are increasing then the ω-limit set of any point $x \in [0,1)$ is that of some D_i, $0 \le i \le m$, and the upper bound on $n(T)$ is $m+1$ in this case. □

Remark. When $m = 1$, the inequality $n(T) \le 2$ is contained in [**10**] or [**11**]. The same upper bounds as the ones we give in the general case would also follow easily.

We begin the proof of Theorem 2.2. We first build a denumerable family $\mathcal{F}$ of zero-Lebesgue-measure subsets in $\mathcal{C}$. We next show that $\lambda_m(\mathcal{C}\backslash(\Omega \cup \mathcal{F})) = 0$, giving $\lambda_m(\mathcal{C}\backslash\Omega) = 0$.

Step 1. Fixing $1 \le i \le m, 0 \le j \le m+1, n \ge 1$ and $(\eta_k)_{1\le k\le n} \in \{0, \dots, m\}^n$, observe that

$$\{D \in \mathcal{C} \mid D_i = \Gamma_{\eta_n,\dots,\eta_1}(D_j)\}$$

has zero Lebesgue measure in $\mathcal{C}$. This is obvious if $i \neq j$, as this is the graph of a Lipschitz map. If $i = j$, then D_i is the fixed point of $\Gamma_{\eta_n,\dots,\eta_1}$ in $[0,1]$, since $n \ge 1$. Define $\mathcal{F}$ as the family of subsets of $\mathcal{C}$ obtained when i, j, n and $(\eta_k)_{1\le k\le n}$ vary.

Step 2. Let $D \in (\Omega \cup \mathcal{F})^c$. We make some preliminary remarks on T.

- Any equality $D_i = T^l D_j^{\pm}$ with $1 \le i \le m$, $0 \le j \le m+1$ and $l \ge 1$ is impossible.
- Setting $I = \{1, \dots, m\}$ there exists $\emptyset \neq J \subset I$ such that, for $i \in J$, $D_i \in \bigcap_{k\ge 0}(T^k[0,1))^{\mathrm{o}}$ and, for $i \in I\backslash J$, there exists $(k_i, l_i) \in \mathbb{N} \times \{1, \dots, m+2\}$ such that $D_i \in (T^{k_i} F_{l_i})^{\mathrm{o}}$.

https://doi.org/10.1017/S0143385705000386 Published online by Cambridge University Press

- Set $K = \max_{i\in I\setminus J} k_i$ and denote by $G_1, \dots, G_p$ the intervals forming $T^K E$. Each G_i is not cut under iterations of T. Also $\lambda\big(\bigcup_{n\geq K} T^n E\big) = 1 - \lambda\big(\bigcup_{0\leq n<K} T^n E\big)$.
- Since the $(T^n G_j)_{1\leq j\leq p, n\geq 0}$ form the remaining mass to be added after time K and since $\operatorname{dist}(D_i, G_k) > 0$ for $i \in J$ and $1 \leq k \leq p$, any D_i with $i \in J$ admits arbitrary close intervals among the $(T^n G_j)_{1\leq j\leq p, n\geq 0}$ on its left-hand side and on its right-hand side.

LEMMA 2.4. *For all $i \in J$ there exist arbitrary large n and $1 \leq j \leq p$ such that the interval $T^n G_j$ is arbitrary close to D_i, lies on its right-hand side and verifies*

$$\frac{\lambda(T^n G_j)}{\operatorname{dist}(D_i, T^n G_j)} \geq \kappa = \frac{1-\gamma}{p} > 0. \tag{1}$$

Proof. Introduce the set $Z \subset \{1, \dots, p\}$ of indices such that, for $j \in Z$, G_j admits iterates converging to D_i on its right-hand side. Set $p' = \operatorname{Card}(Z)$ and note that $1 \leq p' \leq p$. Fix $\varepsilon > 0$ such that all approximations of D_i in $[D_i, D_i + \varepsilon]$ are made by iterates of G_j with $j \in Z$. Suppose now that for some N inequality (1) is false for all $n \geq N$ and $j \in Z$.

Let $T^{v_0} G_{u_0}$ be a best record iterate on the right-hand side of D_i with $u_0 \in Z$ and a record time $v_0 \geq N$. Let $(T^{v_k} G_k)_{k\in Z\setminus\{u_0\}}$ be the first iterates of each G_k, with $k \in Z\setminus\{w_0\}$, standing between D_i and $T^{v_0} G_{u_0}$. For such a k, note that $v_k \geq N$. Finally observe that the possible mass that can be added between D_i and $T^{v_0} G_{u_0}$ by iterates of $(G_j)_{j\in Z}$ is at most

$$\begin{aligned}\sum_{k\in Z\setminus\{u_0\}} \frac{1}{1-\gamma}\lambda(T^{v_k} G_k) + \frac{\gamma}{1-\gamma}\lambda(T^{v_0} G_{u_0}) &\leq \sum_{k\in Z} \frac{1}{1-\gamma}\,\lambda(T^{v_k} G_k)\\ &< \frac{\kappa}{1-\gamma} \sum_{k\in Z} \operatorname{dist}(D_i, T^{v_k} G_k)\\ &< \frac{\kappa p'}{1-\gamma}\ \operatorname{dist}(D_i, T^{v_0} G_{u_0}).\end{aligned}$$

Since $\kappa p'/(1-\gamma) \leq \kappa p/(1-\gamma) \leq 1$, this brings a contradiction. □

Step 3. We prove that $\lambda_m(\mathcal{C}\setminus(\Omega \cup \mathcal{F})) = 0$. Supposing the contrary, there exists $J \subset I$ with $J \neq \emptyset$ such that the following set A satisfies $\lambda_m(A) > 0$:

$$A = \left\{ D \in \mathcal{C}\setminus(\Omega \cup \mathcal{F}) \;\middle|\; \{D_j\}_{j\in I\setminus J} \subset \bigcup_{n\geq 0} (T^n E)^{\circ}, \{D_j\}_{j\in J} \cap \left(\bigcup_{n\geq 0} T^n E\right) = \emptyset \right\}.$$

Then A admits a density point D^0. Denoting by $B_m(a, r)$ the Euclidean ball in $\mathbb{R}^m$ of center a and radius $r > 0$, one has

$$\lim_{r\to 0} \frac{\lambda_m(B(D^0, r) \cap A)}{\lambda_m(B(D^0, r))} = 1. \tag{2}$$

We use for D^0 the quantities defined in Step 2. We add an exponent $(\cdot)^0$ to mark the dependence in D^0, except for T for simplicity. We now make arbitrarily small perturbations of D^0. Note that the corresponding map will remain injective and the images

https://doi.org/10.1017/S0143385705000386 Published online by Cambridge University Press

of the wandering intervals cannot overlap. Also each D_i with $i \in I\backslash J$ remains strictly in the interior of $T^{k_i} F_{l_i}$.

Fix $i \in J$. There exists $1 \le j \le p$ and arbitrarily large n such that $T^n G_j$ is on the right-hand side of D_i^0 with $\lambda(T^n G_j^0) \ge \kappa \operatorname{dist}(D_i^0, T^n G_j^0)$. Set $G = G_j$, $d_n = \operatorname{dist}(D_i^0, T^n G^0)$ and $l_n = \lambda(T^n G^0)$. Without loss of generality, assume that

$$d_n = \min\{\operatorname{dist}(D_k^0, T^l G^0) \mid 1 \le k \le m, 0 \le l \le n\},$$

since the lengths of the $(T^l G^0)_{l \ge 0}$ decrease. Denote by $(e_k)_{1 \le k \le m}$ the canonical basis of $\mathbb{R}^m$ and consider the path $D^\eta = D^0 + \eta e_i$, $\eta \ge 0$. All the corresponding quantities for the transformation get some $(\cdot)^\eta$. Introduce an integer q such that $2\gamma^q(1 + \kappa/4) \le 1/2$. Since the boundaries of G^0 are of the form $T^s(D_u^0)^\pm$, $s \ge 1$, we make the following observations for $0 \le l \le n$ and small η:

$$\operatorname{dist}(D_i^\eta, T^n G^\eta) \le d_n - (1 - \gamma^n)\eta \tag{3}$$

$$\lambda(T^n G^\eta) \ge l_n - 2\gamma^n \eta \tag{4}$$

$$\operatorname{dist}(D_k^\eta, T^l G^\eta) \ge \operatorname{dist}(D_k^0, T^l G^0) - \gamma^l \eta \ge d_n - \gamma^l \eta, \text{ if } k \in \{1, \dots, m\}\backslash\{i\}. \tag{5}$$

Also, for any $T^l G^\eta$ with $0 \le l < n$ and close to D_i we have $\operatorname{dist}(D_i^\eta, T^l G^\eta) \ge d_n + \eta(1 - \gamma)$ if it is on its left-hand side and $\operatorname{dist}(D_i^\eta, T^l G^\eta) \ge \operatorname{dist}(D_i^\eta, T^n G^\eta)$ if it stands on its right-hand side, as $T^l G$ is protected by $T^n G$. Then it results from (3)–(5) and $l_n \ge \kappa d_n$ that

$$\eta_1 = \inf\{\eta \ge 0 \mid D_i^\eta \in T^n G^\eta, \operatorname{dist}(D_i^\eta, \partial(T^n G^\eta)) = \kappa d_n/4\}$$

exists and satisfies $\eta_1 \le (1 - \gamma^n)^{-1} d_n(1 + \kappa/4) \le 2d_n(1 + \kappa/4)$ for large enough n. Also, for $q \le l \le n$ and $k \in I\backslash\{i\}$, we have $\operatorname{dist}(D_k^{\eta_1}, T^l G^{\eta_1}) \ge d_n - \gamma^l \eta_1 \ge d_n/2$. Similarly for $q \le l \le n$ we have $\operatorname{dist}(D_i^{\eta_1}, T^l G^{\eta_1}) \ge \min\{d_n, \operatorname{dist}(D_i^{\eta_1}, \partial(T^n G^{\eta_1}))\} \ge d_n(1 \wedge \kappa/4)$. These inequalities evidently remain true for $1 \le l < q$ if n is large enough since q is fixed and $d_n \to 0$.

Finally all boundaries of the $(T^l G)_{0 \le l \le n}$ are at most $1-$Lipschitz maps of D and thus the ball $B(D^{\eta_1}, \eta_2)$ with $\eta_2 = (d_n/4) \wedge (\kappa d_n/8)$ is entirely contained in A^c as $D_i \in T^n G$ in that case. As a result we get the inequality

$$\frac{\lambda_m(B(D^0, \eta_1 + \eta_2) \cap A^c)}{\lambda_m(B(D^0, \eta_1 + \eta_2))} \ge \left(\frac{\eta_2}{\eta_1 + \eta_2}\right)^m \ge \left(\frac{\kappa \wedge 2}{\kappa \wedge 2 + 16(1 + \kappa/4)}\right)^m > 0. \tag{6}$$

As this lower bound is independent of n and $\eta_1 + \eta_2 \to 0$, as $n \to +\infty$, this contradicts (2) and concludes the proof of the theorem. □

Remark 1. In the proof of Theorem 2.2 the injectivity assumption essentially simplifies the description of $[0, 1)\backslash T^n[0, 1)$, $n \ge 1$, which consists of considering the forward orbits of the wandering intervals defined by E. When suppressing the injectivity hypothesis and among other difficulties, the number of pieces of $E_n = T^n[0, 1)\backslash T^{n+1}[0, 1)$ can increase with n and this corresponds to p becoming large in inequality (1). It is, however, natural to conjecture that without supposing injectivity the dynamics still consists of attracting periodic orbits. For instance, Lemma 2.3 extends with only slight changes in the formulation.

https://doi.org/10.1017/S0143385705000386 Published online by Cambridge University Press

Remark 2. Theorem 2.2 provides an algorithm for concretely computing the dynamics of a generic injective quasi-contraction.

(1) Compute the $(T^n F_i)_{1\le i\le m+2}$, $n \ge 0$, until each D_j is covered, $1 \le j \le m$. Theorem 2.2 implies that this process stops almost surely. Then write $D_j \in T^{k_j} F_{l_j}$ with $1 \le j \le m$.

(2) Reversing the calculations, we get the $T^{-k} D_j$, with $0 \le k \le k_j$ and $1 \le j \le m$. Then $\mathcal{D} = \{T^{-k} D_j \mid 0 \le k \le k_j, 1 \le j \le m\}$ defines a stable partition of $[0, 1)$. Considering the orbit of any point in each atom allows one to locate and count the periodic attractors.

For a non-generic injective quasi-contraction having diffusive ergodic components, the first step in the above algorithm does not stop. It may be possible to extend the first part of the proof of Theorem 2.2 to describe the structure of the transformation via an algorithm generalizing the continued fractions algorithm, giving the sequence of best approximations of an irrational number. This would probably provide natural isomorphisms with maps in the class of interval exchange transformations.

2.3. *Structural stability.* We show that perturbations of a generic injective quasi-contraction still have the same behaviour as that described by Theorem 2.2. For any map $\delta : [0, 1) \to \mathbb{R}$, set $T_\delta(x) = T(x) + \delta(x)$.

THEOREM 2.5. *Let Ω be the open set of Lemma 2.3 and Theorem 2.2. For $D \in \Omega$ denote by T the corresponding transformation and suppose that $T[0, 1) \subset (0, 1)$. Then, for $\varepsilon > 0$, there is an $\eta > 0$ such that any sequence of maps $\delta = (\delta_n)_{n\ge1}$ with $\|\delta_n\|_\infty \le \eta$ for all $n \ge 0$ and $\|\delta_n\|_\infty \to 0$ satisfies:*

(1) *for every $x \in [0, 1)$, the sequence $(x_n)_{n\ge0}$ defined by $x_0 = x$ and $x_{n+1} = T_{\delta_n(x_n)} x_n$ converges to a periodic orbit of T;*

(2) *every interval I in the attraction domain of a periodic orbit $\mathcal{O}$ for T contains a subinterval J verifying $|J| \ge (1-\varepsilon)|I|$ such that for $x \in J$ one has $\operatorname{dist}(x_n, T^n x) \to 0$, as $n \to +\infty$.*

Proof. As in Lemma 2.3 assume that there exist $k_i \ge 0$ and $l_i \in \{1, \dots, m+2\}$ for $1 \le i \le m$ such that $D_i \in \operatorname{int}(T^{k_i} E_{l_i})$. Then the partition $\mathcal{P}(D')$ is stable, where

$$D' = \{T^{-k} D_i \mid 0 \le k \le k_i, 1 \le i \le m\}.$$

Take η small enough so that the image segments by T_{δ_n} remain disjoint and contained in $(0, 1)$. Suppose first that each δ_n is a constant map still written as δ_n. Build D'' by replacing $T^{-k} D_i$ by $T_{\delta_1}^{-1} \cdots T_{\delta_k}^{-1} D_i$ in the definition of D'. These quantities exist if η is small enough and are perturbations. Also, if η is small enough, the condition on the length in the second part of the theorem is satisfied. We show that an atom of $\mathcal{P}(D'')$ converges under iterations of the perturbed transformations to the same periodic orbit for T as the corresponding atom of $\mathcal{P}(D')$ under iterations of T.

Recall that the definition of Ω implies that any D_u cannot be written as $T^k D_v^{\pm}$ with $k \ge 1$. Therefore, if $a = [T^{-l_i} D_i, T^{-l_j} D_j)$ is an atom of $\mathcal{P}(D')$, then for $l \ge \min\{l_i, l_j\} + 1$ at least one boundary point of $T^l a$ is at a positive distance of the boundary of the

https://doi.org/10.1017/S0143385705000386 Published online by Cambridge University Press

atom of $\mathcal{P}(D')$ where $T^l a$ lies. For $l \geq \tau_a := \max\{l_i, l_j\} + 1$, both boundaries share this property. Since the elements of the orbit of $T^l a$ that lie in a given atom of $\mathcal{P}(D')$ form a decreasing sequence of intervals, there finally exists $\eta_a > 0$ such that the distance between a boundary of $T^l a$ and the boundary of the atom where $T^l a$ lies is either 0 or $\geq \eta_a$. The last case is verified for both boundaries when $l \geq \tau_a$. This is then also true for the distance between $T^l a$ and any D_i for $l \geq \tau_a$.

Let a' be the atom of $\mathcal{P}(D'')$ corresponding to a. First choose η small enough such that for all atoms a and all $0 \leq l \leq \tau_a$ the distance between a boundary of $T_{\delta_l} \cdots T_{\delta_1} a'$ and the boundary of the atom of $\mathcal{P}(D'')$ where it lies is either 0 or $\geq \eta_a/4$. Assume also that the boundaries of $T_{\delta_{\tau_a}} \cdots T_{\delta_1} a'$ and $T^{\tau_a} a$ differ by less than $\eta_a/4$. Finally, impose

$$\eta \leq \left(\frac{1-\gamma}{4}\right) \min\{\eta_a \mid a \in \mathcal{P}(D')\}.$$

Now let $y \in T_{\delta_{\tau_a}} \cdots T_{\delta_1} a'$ and $z \in T^{\tau_a} a$ be such that $\operatorname{dist}(y, z) \leq \eta_a/4$. We assert that the images of y by $(T_{\delta_{\tau_a+n}} \cdots T_{\delta_{\tau_a+1}})_{n\geq 1}$ and the iterates of z under $(T^n)_{n\geq 1}$ lie in the same intervals of $\mathcal{P}(D)$ and the distance between the corresponding images tends to zero. Indeed, for $n \geq 1$,

$$\operatorname{dist}(T_{\delta_{\tau_a+n}} \cdots T_{\delta_{\tau_a+1}}, T^n z) \leq \gamma^n \, |y - z| + \sum_{s=0}^{n-1} \gamma^{n-s-1} \, \delta_{s+\tau_a} \tag{7}$$

$$\leq \frac{\eta_a}{4} + \frac{\eta_a}{4} \leq \frac{\eta_a}{2}. \tag{8}$$

Then (8) gives the first assertion and (7) the second. To conclude the proof of the theorem, fix x and apply the result for constant maps with the sequence $(\delta_n)_{n\geq 0} := (\delta_n(x_n))_{n\geq 0}$. □

3. *A class of p-invariant measures on the circle*

We use Theorem 2.2 to describe a class of Borel probability measures on the circle invariant by $\times p$ and with zero entropy. These measures generalize the family of Sturm measures (see Bullett and Sentenac [**6**]) and appear in the study of *maximizing measures*, as mentioned in the introduction.

Recall that Sturm measures are characterized by the fact that their support is contained in a semi-circle (see Bullett and Sentenac [**6**]). We introduce a more general family of domains with symmetry properties.

3.1. *Definition of a p-flower with m petals.*

Consider the circle $S^1 = \mathbb{R}/\mathbb{Z}$ endowed with Lebesgue measure and the transformation $T(x) = px \bmod(1)$, where $p \geq 2$ is a fixed integer.

Definition 3.1.

(1) Fix $m \geq 1$. Then let:

- $S_{p,m} = \{(s_i)_{1\leq i\leq m} \mid s_1 < s_2 < \cdots < s_m < s_1 + 1/p, 0 \leq s_1 < 1/p\} \subset \mathbb{R}^m$;
- $K_{p,m} = \{(k_i)_{1\leq i\leq m} \mid 0 \leq k_i \leq p-1, k_i \neq k_{i+1}, \text{for } 1 \leq i \leq m-1, k_m \neq p - k_1\}$.

https://doi.org/10.1017/S0143385705000386 Published online by Cambridge University Press


Also set $s_{m+1} = s_1 + 1/p$ and $k_{m+1} = k_1 - 1$. For any $(s, k) \in S_{p,m} \times K_{p,m}$, define a '$p$-flower with m petals' as follows:

$$\mathrm{Fl}(s, k) = \bigcup_{1 \le i \le m} \left([s_i, s_{i+1}] + \frac{k_i}{p} \right). \tag{9}$$

(2) For $(s, k) \in S_{p,m} \times K_{p,m}$ and $\varepsilon = (\varepsilon_i)_{1 \le i \le m} \in \{0, 1\}^m$, define $\mathrm{Fl}(s, k, \varepsilon)$ as in (9), but delete $s_{i+1} + k_i/p$ if $\varepsilon_i = 0$ and $s_{i+1} + k_{i+1}/p$ if $\varepsilon_i = 1$, for $1 \le i \le m$. To $\mathrm{Fl}(s, k, \varepsilon)$ associate the right-inverse $\eta_{s,k,\varepsilon}$ of T verifying for all $x \in \mathbb{R}/\mathbb{Z}$ that

$$\eta_{s,k,\varepsilon}(x) \in \mathrm{Fl}(s, k, \varepsilon) \quad \text{and} \quad p\eta_{s,k,\varepsilon}(x) = x \bmod(1).$$

Remark. For $p = 2$, $m = 1$ and $(s, k) \in S_{2,1} \times K_{2,1}$, then $\mathrm{Fl}(s, k)$ is a closed semi-circle. An example of a 2-flower with three petals and $k = (k_1, k_2, k_3) = (0, 1, 0)$ is



We consider below the set of T-invariant measures with support in a p-flower with m petals. These tend to be periodic. Observe that any finite union of periodic orbits is contained in the interior of some $\mathrm{Fl}(s, k)$ for suitable m, s and k. Start with an elementary lemma.

LEMMA 3.2. *Let $(s, k) \in S_{p,m} \times K_{p,m}$ and μ be a Borel probability measure on S^1.*

(1) *Let $\mu = T\mu$. Then for any x there is at most one $0 \le k \le p-1$ with $\mu\{x + k/p\} > 0$.*

(2) *Both $\mu = T\mu$ and $\mathrm{supp}(\mu) \subset \mathrm{Fl}(s, k)$ hold if and only if there exists $\varepsilon \in \{0, 1\}^m$ with $\mu = \eta_{s,k,\varepsilon}\mu$.*

Proof. If $\mu\{x\} > 0$, then $T^{-n-n'}\{x\} \cap T^{-n}\{x\} \neq \emptyset$ for some $n \ge 0$ and $n' \ge 1$, since μ is a finite measure. Thus $T^{n'}x = x$. If for some $1 \le k \le p-1$ one has $\mu\{x + k/p\} > 0$, then $x + k/p$ is also periodic. As $x + k/p$ must be in the same orbit as x, this is a contradiction.

Consider the second part. If $\mu = \eta_{s,k,\varepsilon}\mu$, then $T\mu = (T \circ \eta_{s,k,\varepsilon})\mu = \mu$. Also $\mathrm{supp}(\mu) = \eta^{-1}_{s,k,\varepsilon}\,\mathrm{supp}(\mu) \subset \mathrm{Fl}(s, k)$. Reciprocally and using the first part, either $s_{i+1} + k_i/p$ or $s_{i+1} + k_{i+1}/p$ has μ-measure zero for $1 \le i \le m$. Next, choose $\mathrm{Fl}(s, k, \varepsilon)$ corresponding to the exclusion from $\mathrm{Fl}(s, k)$ of any point with zero mass in each couple mentioned above. Then $\eta_{s,k,\varepsilon}\mu = \mu$. □

Remark. If a T-invariant measure μ satisfies $\mathrm{supp}(\mu) \subset \mathrm{Fl}(s, k)$, then it need not be invariant with respect to all $\eta_{s,k,\varepsilon}$. For instance, taking $m = 1$ and the right-inverse η of T defined with $(0, 1/p]$, observe that δ_0 verifies $T\delta_0 = \delta_0$ but $\eta\delta_0 = \delta_{1/p}$.

3.2. *Invariant measures with support in a p-flower with m petals.*

THEOREM 3.3. *Let $p \ge 2$, $m \ge 1$ and $k \in K_{p,m}$. There is an open set $\Omega \subset S_{p,m}$ of full Lebesgue measure in $S_{p,m}$ such that, for $s \in \Omega$, the ergodic T-invariant Borel probability measures with support in $\mathrm{Fl}(s, k)$ are periodic and their number is $\{1, \ldots, n, m\}$. These measures are locally constant functions of s on Ω.*

https://doi.org/10.1017/S0143385705000386 Published online by Cambridge University Press

Proof. Fixing $s \in S_{p,m}$ Lemma 3.2 implies that a T-invariant Borel probability measure with support in $\mathrm{Fl}(s,k)$ is invariant with respect to some $\eta_{s,k,\varepsilon}$. Then fix $\varepsilon \in \{0,1\}^m$ and write η for $\eta_{s,k,\varepsilon}$. On $[0,1)$ observe that η is an injective quasi-contraction with $\{\Gamma_i\} \subset \{x \longmapsto x/p + i/p \mid 0 \leq i \leq p-1\}$ and $D = \{ps_j \bmod(1) \mid 1 \leq j \leq m\}\backslash\{0\}$. Note that if $ps_j \neq 0 \bmod(1)$ for $1 \leq j \leq m$, then 0 is not a discontinuity of η when seen on the circle. This condition can be assumed to be realized.

Now applying Theorem 2.2 and using the notation of this theorem, there is an open set $\Omega \subset S_{p,m}$ of full Lebesgue measure in $S_{p,m}$, such that for $s \in \Omega$, then for all $1 \leq i \leq m$, $ps_i \in (\eta^{k_i} F_{l_i})^{\mathrm{o}}$ for some $k_i \geq 0$ and $1 \leq l_i \leq m+2$, where $(F_j)_{1 \leq j \leq m+2}$ are the wandering intervals of $[0,1)\backslash\eta([0,1))$.

When η is seen on the circle, the partition defined by $\mathcal{D} = \{\eta^{-k}(ps_i) \mid 1 \leq i \leq m, 0 \leq k \leq k_i\}$ is stable under η. Then each point x under iterates of η converges to a periodic orbit and its ω-limit set verifies $\omega(x) \subset \bigcup_{1 \leq i \leq m} \omega(ps_i)$, as η is order-preserving on each interval of continuity. There are then at most m periodic orbits. Also, since $ps_i \in (\eta^{k_i} F_{l_i})^{\mathrm{o}}$ for $1 \leq i \leq m$, each periodic point does not belong to $\mathcal{D}$. Thus all $\eta_{s,k,\varepsilon}$, when ε vary, have the same periodic orbits. Note also that on Ω the periodic orbits vary continuously with s and the lengths of these orbits are locally constant. Thus those orbits are locally constant. □

Remark. If $p = 2$ and $m = 1$, one obtains that for an open set of full Lebesgue measure in $s \in [0,1)$, the semi-circle $[s, s+1/2]$ is uniquely ergodic with respect to $x \longmapsto 2x \bmod(1)$. This is a well-known result on Sturm measures (see Bousch and Mairesse [**3**] or Bullett and Sentenac [**6**]), which is in fact valid for all s. This last point will, however, follow from the study of the next section, since any right-inverse of $x \longmapsto 2x \bmod(1)$ defined with a semi-circle is an 'order-preserving quasi-contraction on the circle' with one piece, in the sense of Definition 4.4. If τ is its rotation number, the conclusion comes from the remark before Proposition 4.16 if τ is rational and from Proposition 4.3 if τ is irrational.

4. *Order-preserving quasi-contractions on the circle*

We examine Theorem 2.2 for a class of quasi-contractions on the circle that preserve cyclic order. We detail fully the maps with a periodic structure.

As for diffeomorphisms of the circle (see Katok and Hasselblatt [**14**, Ch. 11]), the main properties of the dynamics are contained in the rotation number (the average speed of rotation of the transformation). We provide a short theory adapted from the usual one of Poincaré. Proofs result from simple modifications of those in [**14**] and are omitted (see Brémont [**4**]).

4.1. *Rotation number.*

We begin with a few definitions. The cyclic order of three distinct points a, b and c on the circle $S^1 = \mathbb{R}/\mathbb{Z}$ is (a,b,c) if turning in the trigonometric sense and starting from a one meets b before c. A map $f : S^1 \to S^1$ is order-preserving if the order of any triple is preserved by f.

In this section we fix $f : S^1 \to S^1$, an order-preserving injective and right-continuous map of degree one. By definition, there is an increasing and right-continuous map

https://doi.org/10.1017/S0143385705000386 Published online by Cambridge University Press

$\tilde{f} : \mathbb{R} \to \mathbb{R}$ satisfying $\tilde{f}(x+1) = \tilde{f}(x) + 1$ for $x \in \mathbb{R}$ and $f = \tilde{f} \bmod(1)$. Such an $\tilde{f}$ is called a lift of f.

PROPOSITION 4.1. *The map f admits a rotation number $\tau(f)$ defined by*

$$\tau(f) = \lim_{n \to +\infty} \frac{1}{n}[(\tilde{f})^n(x) - x] \quad \bmod(1), \tag{10}$$

where $x \in \mathbb{R}$ and $\tilde{f}$ is any lift of f. This quantity is independent of x and on the lift $\tilde{f}$. Also the limit is uniform in $x \in \mathbb{R}$.

Next, we distinguish the situations $\tau(f) \in \mathbb{Q}$ and $\tau(f) \notin \mathbb{Q}$. As for diffeomorphisms of the circle, the behaviour is significantly different in each case.

PROPOSITION 4.2.

(1) *We have $\tau(f) \in \mathbb{Q}$ if and only if f admits a periodic point.*
(2) *If $\tau(f) = p/q$ with $p \wedge q = 1$, then all periodic points have period q.*
(3) *Let $\tau(f) = p/q$ with $p \wedge q = 1$. If x is q-periodic, the order on the circle of $\{x, f(x), \dots, f^{q-1}(x)\}$ is that of $\{0, p/q, \dots, (q-1)p/q\}$. In particular, the orbit of x induces intervals on the circle that are sent one into another by f.*

Denote by R_α the rotation of angle α. When $\tau(f) \notin \mathbb{Q}$, then f is closely related to $R_{\tau(f)}$. Recall that for sufficiently regular diffeomorphisms of the circle with an irrational rotation number, this latter one is a complete invariant for topological conjugacy.

PROPOSITION 4.3. *Set $\tau = \tau(f)$ and assume that $\tau \notin \mathbb{Q}$.*

(1) *If $\tilde{f}$ is a lift of f, then for all $(n_1, n_2) \geq 0$ and $(m_1, m_2) \in \mathbb{Z}^2$,*

$$n_1\tau + m_1 < n_2\tau + m_2 \quad \text{if and only if for all } x \in \mathbb{R},\ \tilde{f}^{n_1}(x) + m_1 < \tilde{f}^{n_2}(x) + m_2.$$

(2) *The map f is topologically semi-conjugated to R_τ: there is a non-decreasing, continuous and surjective map $h : [0,1) \to [0,1)$ such that $h \circ f = R_\tau \circ h$.*
(3) *The map f is uniquely ergodic.*

4.2. *Definition and first properties.* We define order-preserving quasi-contractions on the circle with $m+1$ pieces. We modify Definition 2.1 to take into account the topology of the circle: the image of an interval is allowed to cover the point 0. Also and up to a change of origin, 0 will always be a discontinuity.

Definition 4.4. Let $m \geq 0$, $0 \leq \gamma < 1$ and there exist strictly increasing γ-Lipschitzian maps $(\Gamma_i)_{0 \leq i \leq m} : [0,1] \to \mathbb{R}$. Set $\mathcal{D} = \{D = (D_i)_{1 \leq i \leq m} \mid 0 = D_0 < D_1 < \cdots < D_m < D_{m+1} = 1\}$. For $D \in \mathcal{D}$ and $\beta = (\beta_i)_{0 \leq i \leq m}$, define $\tilde{T} : \mathbb{R} \to \mathbb{R}$ by $\tilde{T}(x+1) = \tilde{T}(x) + 1$ and on each $[D_i, D_{i+1})$ for $0 \leq i \leq m$ by $\tilde{T} : \mathbb{R} \to \mathbb{R}$ by

$$\tilde{T}(x) = \Gamma_i(x) + \beta_i,$$

under the conditions $\mathcal{C} = \mathcal{D} \cap \{0 \leq \Gamma_0(0) + \beta_0,\ \Gamma_m(1) + \beta_m < \Gamma_0(0) + \beta_0 + 1,\ \Gamma_i(D_{i+1}) + \beta_i < \Gamma_{i+1}(D_{i+1}) + \beta_{i+1}, \text{for } 0 \leq i \leq m\}$ on $(D, \beta) \in \mathbb{R}^{2m+1}$. Write $\mathcal{C}(\beta)$ for the section of $\mathcal{C}$ corresponding to fixing β. Finally, set

$$T = \tilde{T} \quad \bmod(1).$$

https://doi.org/10.1017/S0143385705000386 Published online by Cambridge University Press

From the previous subsection, T admits a rotation number $\tau(T)$. As T is defined by (D, β), we write $\tau(D, \beta)$ for $\tau(T)$. First, the same strategy as for proving Theorem 2.2 gives that by fixing β generically for $(D, \beta) \in \mathcal{C}$, the transformation T admits a periodic point. Combined with Proposition 4.2 we obtain the following result.

THEOREM 4.5. *For all $\beta \in \mathbb{R}^{m+1}$, there is an open set $\Omega(\beta) \subset \mathcal{C}(\beta)$ with $\lambda_m(\mathcal{C}(\beta)\backslash\Omega(\beta)) = 0$ such that $\tau(D, \beta) \in \mathbb{Q}$ for $D \in \Omega(\beta)$.*

We specify below the transformations with a given rational rotation number. We consider first the continuity properties of the rotation number $\tau(D, \beta)$ in $\mathcal{C}$. Set $u = (1, \ldots, 1) \in \mathbb{R}^{m+1}$.

LEMMA 4.6.

(1) *The map $(D, \beta) \longmapsto \tau(D, \beta)$ is continuous on $\overline{\mathcal{C}}$.*
(2) *For $(D, \beta) \in \overline{\mathcal{C}}$, the map $t \longmapsto \tau(D, \beta + tu)$ is non-decreasing and continuous from $[-\Gamma_0(0) - \beta_0, 1 - \Gamma_0(0) - \beta_0]$ onto $[0, 1]$.*

Proof. (1) The following classic idea can be found in Katok and Hasselblatt [**14**]. Let $(D, \beta) \in \overline{\mathcal{C}}$ and denote by T the associated map. Let p/q and p'/q' be irreducible rational numbers such that $p'/q' < \tau(D, \beta) < p/q$ and let U be the lift of T such that $-1 < U^q - \mathrm{Id} - p \leq 0$. Remark that $U^q - \mathrm{Id} - p < 0$, otherwise $\tau(D, \beta) = p/q$.

Let x be such that $T^l x$ is not a discontinuity for T, $0 \leq l \leq q - 1$ (there is a finite set to avoid). If T' is associated to (D', β') and (D', β') is close to (D, β), the orbits $(T^l x)_{0 \leq l \leq q-1}$ and $((T')^l x)_{0 \leq l \leq q-1}$ are close. Thus the lift U' of T' compatible with U satisfies $U^q(x) < p + x$. This gives $\tau(D', \beta') \leq p/q$. Symmetrically $\tau(D', \beta') \geq p'/q'$ for (D', β') close enough to (D, β). This concludes the proof of (1).

(2) From (1) the map $t \longmapsto \tau(D, \beta + tu)$ is continuous on $[-\Gamma_0(0) - \beta_0, 1 - \Gamma_0(0) - \beta_0]$. For $t \in [-\Gamma_0(0) - \beta_0, 1 - \Gamma_0(0) - \beta_0]$ let U_t be the lift as in Definition 4.4 corresponding to the map of parameters $(D, \beta + tu)$. All U_t are compatible and satisfy $0 \leq U_t(0) \leq 1$. Note that, for $-\Gamma_0(0) - \beta_0 \leq t < t' \leq 1 - \Gamma_0(0) - \beta_0$,

$$0 \leq \lim_{n \to +\infty} \frac{U_t^n(0)}{n} \leq \lim_{n \to +\infty} \frac{U_{t'}^n(0)}{n} \leq 1.$$

This implies that $t \longmapsto \tau(D, \beta + tu)$ is non-decreasing. Next, if $t \in \{-\Gamma_0(0) - \beta_0, 1 - \Gamma_0(0) - \beta_0\}$, then $\tau(D, \beta + tu) = 0 \bmod(1)$. Therefore the map $t \longmapsto \tau(D, \beta + tu)$ is either equal to 0 on $[-\Gamma_0(0) - \beta_0, 1 - \Gamma_0(0) - \beta_0]$ or its image is exactly $[0, 1]$. The first case is impossible as for some values of $t \in (-\Gamma_0(0) - \beta_0, 1 - \Gamma_0(0) - \beta_0)$ the graph of U_t is at a positive distance from $x \longmapsto x$ and $x \longmapsto x + 1$. For such t we get $\tau(D, \beta + tu) \neq 0 \bmod(1)$, using Proposition 4.2. □

Definition 4.7.

(1) For $0 \leq r < 1$ define the closed and non-empty interval $[b^-_{D,\beta}(r), b^+_{D,\beta}(r)]$ as the set of parameters t in $[-\Gamma_0(0) - \beta_0, 1 - \Gamma_0(0) - \beta_0]$ for which $\tau(D, \beta + tu) = r$.
(2) Let $\mathcal{C}_0 = \overline{\mathcal{C}} \cap \{\Gamma_i(D_i)\beta_i = \Gamma_{i-1}(D_i) + \beta_{i-1}\ , 1 \leq i \leq m\}$ be the set of parameters in $\overline{\mathcal{C}}$ for which the transformation makes no jump at D_i, $1 \leq i \leq m$.
(3) Set $\mathcal{C}_0(p/q) = \{(D, \beta, t) \mid (D, \beta) \in \mathcal{C}_0, b^-_{D,\beta}(p/q) \leq t \leq b^+_{D,\beta}(p/q)\}$.

https://doi.org/10.1017/S0143385705000386 Published online by Cambridge University Press

We then get the following.

PROPOSITION 4.8. *Let $0 \leq p/q < 1$ with $p \wedge q = 1$. Then we have the following.*

(1) *We have $\tau(D, \beta) = p/q$ if and only if $\tilde{T}^q D_i^- \leq D_i + p \leq \tilde{T}^q D_i$ for some $0 \leq i \leq m$.*

(2) *The maps $b^{\pm}_{D,\beta}(p/q)$ are continuous functions of $(D, \beta) \in \mathcal{C}_0$.*

(3) *For $(D, \beta) \in \mathcal{C}_0$ we have $b^-_{D,\beta}(p/q) < b^+_{D,\beta}(p/q)$. Let $T_{D,\beta,t}$ be associated with (D, β, t). Then the $T^l_{D,\beta,t}0^{\pm}$ for $1 \leq l \leq q$, are continuous maps of $(D, \beta, t) \in \mathcal{C}_0(p/q)$. When $1 \leq l \leq q-1$, these are distinct from each other and from 0. Also,*

$$\tilde{T}^q_{D,\beta,t}(0^-) \leq p \leq \tilde{T}^q_{D,\beta,t}(0),$$

with $\tilde{T}^q_{D,\beta,t}(0^-) = p$ if and only if $t = b^+_{D,\beta}(p/q)$ and $\tilde{T}^q_{D,\beta,t}(0) = p$ if and only if $t = b^-_{D,\beta}(p/q)$.

Proof. (1) By Proposition 4.2 we have $\tau(D, \beta) = p/q$ if and only if $\tilde{T}^q - p$ admits a fixed point. As the graph of $\tilde{T}^q - p$ is increasing and piecewise contracting, this condition is equivalent to the existence of a discontinuity y for $\tilde{T}^q - p$ such that $\tilde{T}^q(y^-) \leq p + y \leq \tilde{T}^q(y)$.

Next, y can be written as $y = \tilde{T}^{-l} D_k$ with $0 \leq l \leq q-1$ and $0 \leq k \leq m$, up to translating by an integer. Suppose that l is minimal. If $l = 0$, then the result is proved. If $l \geq 1$ then y is not a discontinuity of $\tilde{T}$. Applying $\tilde{T}$ we obtain $\tilde{T}^q((\tilde{T}(y))^-) \leq p + \tilde{T}(y) \leq \tilde{T}^q(\tilde{T}(y))$ and $\tilde{T}(y) = \tilde{T}^{-l+1} D_k$ is a discontinuity of $\tilde{T}^q$. Repeating this operation gives the result.

(2) Observe first that the continuity of the rotation number implies that $b^-_{D,\beta}(p/q)$ and $b^+_{D,\beta}(p/q)$ are respectively lower semi-continuous and upper semi-continuous maps of $(D, \beta) \in \overline{\mathcal{C}}$. Suppose now that 0 is the only discontinuity of $\tilde{T}$ on $[0, 1)$, that is $(D, \beta) \in \mathcal{C}_0$.

Let $\delta > 0$. For $(D', \beta') \in \mathcal{C}_0$ close to (D, β), the graph of $\tilde{T}_{D',\beta'+t'u}$ with $t' = b^+_{D,\beta}(p/q) - \delta$ is below the graph of $\tilde{T}_{D,\beta+tu}$ with $t = b^+_{D,\beta}(p/q)$. Thus $\tau(D', \beta' + t'u) \leq p/q$ and then $b^+_{D',\beta'}(p/q) \geq b^+_{D,\beta}(p/q) - \delta$. This gives the continuity of $b^+_{D,\beta}(p/q)$. The same holds for $b^-_{D,\beta}(p/q)$.

(3) For $(D, \beta, t) \in \mathcal{C}_0(p/q)$ we have $\tau(D, \beta + tu) = p/q$. Observe first that $\operatorname{dist}(\tilde{T}^l_{D,\beta,t}(0^{\pm}), \mathbb{Z}) > 0$, for $1 \leq l \leq q-1$, otherwise $\tau(D, \beta + tu)$ would be of the form r/l, whereas p/q is irreducible. This gives the continuity of $\tilde{T}^l_{D,\beta,t}(0^{\pm})$ on $\mathcal{C}_0(p/q)$ for $1 \leq l \leq q$. Similarly all $T^l_{D,\beta,t}0^{\pm}$ for $1 \leq l \leq q-1$ are distinct from each other.

Using point (1) and the fact that 0 is the only discontinuity of $T_{D,\beta,t}$ we have $\tilde{T}^q_{D,\beta,t}(0^-) \leq p \leq \tilde{T}^q_{D,\beta,t}(0)$. As $\tilde{T}^q_{D,\beta,t}(0^-) < \tilde{T}^q_{D,\beta,t}(0)$, suppose for example that $\tilde{T}^q_{D,\beta,t}(0^-) < p \leq \tilde{T}^q_{D,\beta,t}(0)$. By continuity and monotony, for some small $\delta > 0$ we still have $\tilde{T}^q_{D,\beta,t+\delta}(0^-) < p \leq \tilde{T}^q_{D,\beta,t+\delta}(0)$. Consequently, $\tau(D, \beta + (t+\delta)u) = p/q$ and $b^-_{D,\beta}(p/q) \leq t < t + \delta \leq b^+_{D,\beta}(p/q)$.

This also proves that $t < b^+_{D,\beta}(p/q)$. Thus if $t = b^+_{D,\beta}(p/q)$ then $\tilde{T}^q_{D,\beta,t}(0^-) = p$. Reciprocally, if this equality is verified then, for any small $\delta > 0$, we have $p < \tilde{T}^q_{D,\beta,t+\delta}(0^-) < \tilde{T}^q_{D,\beta,t+\delta}(0)$. Thus $\tau(D, \beta + (t+\delta)u) > p/q$ and this proves the other sense. The same holds for $b^-_{D,\beta}(p/q)$. □

https://doi.org/10.1017/S0143385705000386 Published online by Cambridge University Press

4.3. *Transformations with a rational rotation number.* In the following all D_i, $0 \le i \le m$, play the same role. We introduce the expression of the transformation when D_i is taken as the origin.

Definition 4.9. Let T be defined as in Definition 4.4. For $0 \le i \le m$, let T_i be the map deduced from T when taking D_i as the origin. Denote by $\Gamma^i = (\Gamma^i_j)_{0\le j\le m}$ the reordered family of γ-Lipschitzian maps and by (D^i, β^i) the corresponding parameters. Set $\Gamma = \Gamma^0$.

First consider the case of a zero rotation number.

PROPOSITION 4.10. *For $(D,\beta) \in \overline{\mathcal{C}}$, we have $\tau(D,\beta) = 0$ if and only if $(D,\beta) \in E(0) := \bigcup_{0\le i\le m} F_i(0)$, where*

$$F_i(0) = \{(D,\beta) \in \overline{\mathcal{C}} \mid \Gamma^i_0(0) + \beta^i_0 \ge 0, \Gamma^i_m(1) + \beta^i_m \le 1\}.$$

The number of fixed points of T is the number of $0 \le i \le m$ such that $(D,\beta) \in F_i(0)$.

Proof. According to Proposition 4.8, $\tau(D,\beta) = 0$ if and only if $\tilde{T}(D_i^-) \le D_i \le \tilde{T}(D_i)$ for some $0 \le i \le m$. Taking D_i as the origin, this can be written as $\tilde{T}_i(0^-) \le 0 \le \tilde{T}_i(0)$ and corresponds to parameters in $F_i(0)$. Finally note that if D_i and D_j are consecutive discontinuities with $\tilde{T}(D_i^-) \le D_i \le \tilde{T}(D_i)$ and $\tilde{T}(D_j^-) \le D_j \le \tilde{T}(D_j)$, there is exactly one fixed point for $\tilde{T}$ in $[D_i, D_j)$. □

Next, turn to the case when $\tau(T) = p/q$ with $0 < p/q < 1$ and $p \wedge q = 1$. Observe in such a case that the graph of $\tilde{T}$ is strictly contained in the tube determined by the lines $x \longmapsto x$ and $x \longmapsto x+1$. Then a non-empty part of this graph is *above* the horizontal line $y = 1$. According to the position with respect to this line, we have the following discussion.

Definition 4.11.

(1) Assume that $0 < \tau(T) < 1$. Set $h = 2i$ when $\tilde{T}(D_i) \le 1 < \tilde{T}(D_{i+1}^-)$ for $0 \le i \le m$ and $h = 2i-1$ when $\tilde{T}(D_i^-) \le 1 < \tilde{T}(D_i)$ for $1 \le i \le m$. We have $0 \le h \le 2m$.

(2) If h is odd, then T is an injective quasi-contraction on $[0,1)$ as in Definition 2.1 with $m+1$ pieces, parameters D' with $D' = D$ and the family of γ-Lipschitzian maps $\Gamma' = (\Gamma'_i)_{0\le i\le m+1}$ with $\Gamma'_i = \Gamma_i + \beta_i \bmod(1)$.

(3) If $h = 2h'$, then T is an injective quasi-contraction on $[0,1)$ with $m+2$ pieces and parameters D' with $D' = \{D_1, \dots, D_{h'}, \tilde{T}^{-1}(1), D_{h'+1}, \dots, D_m\}$ and the family of γ-Lipschitzian maps $\Gamma' = (\Gamma'_i)_{0\le i\le m+1}$ such that $\Gamma'_i = \Gamma_i + \beta_i$ for $0 \le i \le h$ and $\Gamma'_i = \Gamma_{i-1} + \beta_{i-1} \bmod(1)$ for $h+1 \le i \le m+1$.

To specify the condition in point (1) of Proposition 4.8 we need to consider the possible codes for $(T^l D_i^{\pm})_{0\le l\le q-1}$. Up to taking T_i one only needs to focus on $(T^l 0^{\pm})_{0\le l\le q-1}$.

LEMMA 4.12. *Let $0 < p/q < 1$ with $p \wedge q = 1$. For $0 \le l \le q-1$ let $0 \le n_l \le q-1$ satisfy $pn_l = l \bmod(q)$. If $\tau(T) = p/q$, then the order on the circle of $(T^l 0^{\pm})_{0\le l\le q-1}$ is*

$$(0, T^{n_1}0^-, T^{n_1}0, \dots, T^{n_i}0^-, T^{n_i}0, \dots, T^{n_{q-1}}0^-, T^{n_{q-1}}0).$$

https://doi.org/10.1017/S0143385705000386 Published online by Cambridge University Press

Proof. As $\tau(T) = p/q$, Proposition 4.2 implies the existence of a q-periodic point x. The order of its orbit on the circle is $(x, T^{n_1}x, \dots, T^{n_i}x, \dots, T^{n_{q-1}}x)$. As the intervals induced by this orbit are stable, the conclusion follows by considering the interval where 0 lies. □

The previous lemma leads to the following definition.

Definition 4.13. Let $0 < p/q < 1$ with $p \wedge q = 1$ and let $\tau(T) = p/q$.

(1) Mark the $2q-1$ intervals in $[0,1)$ induced by $(T^l 0^{\pm})_{0\le l\le q-1}$. Specifically:
- $(T^{n_i}0, T^{n_{i+1}}0^-)$ has number $2i+1$, for $0 \le i \le q-2$;
- $[T^{n_i}0^-, T^{n_i}0]$ has number $2i$, for $1 \le i \le q-1$;
- $(T^{n_{q-1}}0, 1)$ has number $2q-1$.

(2) The interval where D_j lies is denoted by d_j, $1 \le j \le m$.

(3) If $h = 2h'-1$ set $S(p/q,h) = \{(d_j)_{1\le j\le m} \mid 1 \le d_j \le d_{j+1} \le 2q-1, d_{h'} = 2(q-p)\}$, where $d_{m+1} = 2q-1$ if $h' = m$. Set $r_j = n_{[(d_j+1)/2]}$ and $r'_j = n_{[d_j/2]}$, for $1 \le j \le m$. Denote by:
- $(\varepsilon_k)_{0\le k\le q-1}$ the code of $(R^k_{p/q}0)_{0\le k\le q-1}$ in the intervals $[0, r_1/q), \dots, [r_m/q, 1)$;
- $(\varepsilon'_k)_{0\le k\le q-1}$ the code of $(R^k_{p/q}0)_{0\le k\le q-1}$ in the intervals $(0, r'_1/q], \dots, (r'_m/q, 1]$.

(4) If $h = 2h'$ set $S(p/q,h) = \{(d_j)_{1\le j\le m} \mid 1 \le d_j \le d_{j+1} \le 2q-1, d_{h'} \le 2(q-p) \le d_{h'+1}\}$, where $d_{m+1} = 2q-1$ if $h' = m$ and $d_0 = 1$ if $h' = 0$. Set $r_j = n_{[(d_j+1)/2]}$ for $1 \le j \le h'$, $r_{h'+1} = n_{q-p}$ and $r_j = n_{[(d_{j-1}+1)/2]}$ for $h'+2 \le j \le m+1$. Also let $r'_j = n_{[d_j/2]}$ for $1 \le j \le h'$, $r'_{h'+1} = n_{q-p}$ and $r'_j = n_{[d_{j-1}/2]}$ for $h'+2 \le j \le m+1$. Denote by:
- $(\varepsilon_k)_{0\le k\le q-1}$ the code of $(R^k_{p/q}0)_{0\le k\le q-1}$ in the intervals $[0, r_1/q), \dots, [r_{m+1}/q, 1)$;
- $(\varepsilon'_k)_{0\le k\le q-1}$ the code of $(R^l_{p/q}0)_{0\le k\le q-1}$ in the intervals $(0, r'_1/q], \dots, (r'_{m+1}/q, 1]$.

Remark. Points (3) and (4) describe the fact that $\tilde{T}^q(0^-) \le p \le \tilde{T}^q(0)$. Then $S(p/q,h)$ lists the possible positions of D with respect to the $(T^l 0^{\pm})_{0\le l\le q-1}$, noting that $\tilde{T}(T^{q-1}0^-) \le 1 \le \tilde{T}(T^{q-1}0)$ and $[T^{q-1}0^-, T^{q-1}0]$ has number $2(q-p)$ since $n_{q-p} = q-1$. Finally, $(\varepsilon_k)_{0\le k\le q-1}$ and $(\varepsilon'_k)_{0\le k\le q-1}$ are the codes of $(T^k 0)_{0\le k\le q-1}$ and $(T^k 0^-)_{0\le k\le q-1}$, respectively.

We now come to the main result of this section. Recall Definitions 4.9, 4.11 and 4.13.

THEOREM 4.14. *Let $0 < p/q < 1$ with $p \wedge q = 1$. Then $\tau(D,\beta) = p/q$ if and only if*

$$(D,\beta) \in E(p/q) := \bigcup_{0\le i\le m} F_i(p/q),$$

where

$$F_i(p/q) := \bigcup_{\substack{d\in S(p/q,h)\\ 0\le h\le 2m,}} G_{i,\Gamma}(p/q,h,d)$$

and $G_i(p/q,h,d) = \{(D,\beta) \in \overline{\mathcal{C}} \mid (D^i,\beta^i) \in G_{0,\Gamma^i}(p/q,h,d)\}$ and finally

https://doi.org/10.1017/S0143385705000386 Published online by Cambridge University Press

$G_{0,\Gamma}(p/q,h,d) \subset \overline{\mathcal{C}}$ *is defined by the inequalities*

$$\begin{cases} \Gamma'_{\varepsilon_{q-1},\dots,\varepsilon_0}(0) \geq 0 \text{ and } \Gamma'_{\varepsilon'_{q-1},\dots,\varepsilon'_0}(1) \leq 1, \\ \Gamma_{h'-1}(D_{h'}) + \beta_{h'-h''} \leq 1 < \Gamma_{h'}(D_{h'+1-h''}) + \beta_{h'}, \quad \text{where } h = 2h' - h'', h'' \in \{0,1\}, \\ D_j \in [\zeta_{d_j}, \zeta_{d_j+1}] \text{ for even } d_j \text{ and } D_j \in (\zeta_{d_j}, \zeta_{d_j+1}) \text{ for odd } d_j,\ 1 \leq j \leq m, \end{cases} \tag{11}$$

where $\zeta_{2t} = \Gamma_{\varepsilon'_{n_t-1},\dots,\varepsilon'_0}(1)$ *and* $\zeta_{2t+1} = \Gamma_{\varepsilon_{n_t-1},\dots,\varepsilon_0}(0)$.

Moreover, for $0 \leq i \leq m$, $0 \leq h, h' \leq m$ *and* $d, d' \in S(p/q,h)$, *the set* $G_{i,\Gamma}(p/q,h,d)$ *is non-empty and even has non-empty interior. If* $(h,d) \neq (h',d')$, *then* $G_{i,\Gamma}(p/q,h,d) \cap G_{i,\Gamma}(p/q,h',d') = \emptyset$.

Proof.

Step 1. From Proposition 4.8 we have $\tau(D,\beta) = p/q$ if and only if some D_i with $0 \leq i \leq m$ verifies point (1) of this proposition. This means that $\tilde{T}_i^q 0^- \leq p \leq \tilde{T}_i^q 0$. Assuming then that $i = 0$, we show that $(D,\beta) \in G_{0,\Gamma}(p/q,h,d)$.

Recall the definition of h given in Definition 4.11. The position of the graph of $\tilde{T}$ with respect to $y = 1$ is then given by the second condition in (11). Next, the order on the circle of $(T^l 0^{\pm})_{0 \leq l \leq q-1}$ is given by Lemma 4.12. Following Definition 4.13, the positions of the elements of D with respect to this sequence are given by some $d \in S(p/q,h)$. This gives the last set of conditions in (11). Finally the first line of (11) is for $\tilde{T}^q(0^-) \leq p \leq \tilde{T}^q(0)$. Thus $(D,\beta) \in G_{0,\Gamma}(p/q,h,d)$.

Step 2. Reciprocally fix $0 \leq i \leq m$, $0 \leq h \leq 2m$ and $d \in S(p/q,h)$. Suppose, for instance, that $h = 2h' - 1$. From the symmetric roles played by the $(D_i)_{0 \leq i \leq m}$, it is enough to prove that $G_{0,\Gamma}(p/q,h,d)$ has non-empty interior and that for (D,β) in this set we have $\tau(D,\beta) = p/q$.

Recall Definition 4.7, Proposition 4.8 and the notation $\mathcal{C}_0$ and $T_{D,\beta,t}$. When $(D,\beta) \in \mathcal{C}_0$ then, if D and β_0 are fixed, the $(\beta_i)_{1 \leq i \leq m}$ are also fixed. Then write $T_{D,\beta_0,t}$ for $T_{D,\beta,t}$. We use the same convention for $b^{\pm}_{D,\beta}(p/q)$. Next, define $\varphi : C \to C$, where $C = \{(s_i)_{1 \leq i \leq m} \mid 0 \leq s_1 \leq \cdots \leq s_m \leq 1\}$, in the following way. For $s \in C$ consider the transformation $T_{s,0,\eta_s}$ with

$$\eta_s = \tfrac{1}{2}(b^-_{s,0}(p/q) + b^+_{s,0}(p/q)).$$

Proposition 4.8 implies that $s \longmapsto \eta_s$ is continuous on C. Similarly $s \longmapsto (T^k_{s,0,\eta_s} 0^{\pm})_{1 \leq k \leq q-1}$ is continuous on C without collapse and each component remains far away from 0. Write the ordered sequence on $[0,1)$ as follows:

$$0 < B_1(s) < \cdots < B_{2q-2}(s) < 1 \quad \text{and set } B_0 = 0,\ B_{2q-1} = 1.$$

Set $\varphi(s) = (s'_1,\dots,s'_m)$, where $s'_{h'} = \tilde{T}^{-1}_{s,0,\eta_s}(1)$. This clearly is a continuous function of $s \in C$. Take now $i \neq h'$, let $N_i = \text{card}\{1 \leq j \leq m \mid d_j = d_i\}$ and suppose that i is exactly the kth index z such that $d_z = d_i$. If $d_i \neq d_{h'}$ we set

$$s'_i = B_{d_i-1}(s) + k\left(\frac{B_{d_i}(s) - B_{d_i-1}(s)}{N_i + 1}\right). \tag{12}$$

https://doi.org/10.1017/S0143385705000386 Published online by Cambridge University Press

If $d_i = d_{h'}$ then $B_{d_{h'}-1}(s) < s'_{h'} < B_{d_{h'}}(s)$ as $\tilde{T}_{s,0,\eta_s}(T^{q-1}_{s,0,\eta_s}0^-) < 1 < \tilde{T}_{s,0,\eta_s}(T^{q-1}_{s,0,\eta_s}0)$. If, for example, $i < h'$, set $N_i = \mathrm{card}\{1 \le j \le h' \mid d_j = d_i\}$ and define s'_i as in (12), replacing $B_{d_i}(s)$ by $s'_{h'}$. As a result we get a continuous $\varphi : C \to C$ with $\varphi(C) \subset C^{\mathrm{o}}$.

From the Brouwer fixed point theorem, φ admits a fixed point $s \in C$ and thus $s \in C^{\mathrm{o}}$. Then $T_{s,0,\eta_s}$ is such that $\tau(T_{s,0,\eta_s}) = p/q$. The $(T^k_{s,0,\eta_s}0^{\pm})_{0\le k\le q-1}$ are all distinct and $\tilde{T}^q_{s,0,\eta_s}(0^-) < p < \tilde{T}^q_{s,0,\eta_s}(0)$. Also the positions of $(D_i)_{1\le i\le m}$ with respect to $(T^k_{s,0,\eta_s}0^{\pm})_{0\le k\le q-1}$ are exactly specified by d. Those points are distinct from each other and from the boundaries of the interval where they lie. Perturbing $(s, 0, \eta_s)$ in such a way that $\tilde{T}(D_{h'})^- < 1 < \tilde{T}(D_{h'})$ gives parameters $(D, \beta) \in \mathcal{C}$ contained in the interior of $G_{0,\Gamma}(p/q, h, d)$. The case when h is even is treated in the same way.

Step 3. We prove by convexity that, for (D, β) in the interior of $G_{0,\Gamma}(p/q, h, d)$, then $\tau(D, \beta) = p/q$, $\tilde{T}^q(0^-) < p < \tilde{T}^q(0)$, the positions of the $(D_i)_{1\le i\le m}$ with respect to the $(T^k0^{\pm})_{0\le k\le q-1}$ are specified by d and the position of $y = 1$ with respect to the graph of $\tilde{T}$ is given by h. Denote by H the parameters in the interior of $G_{0,\Gamma}(p/q, h, d)$ having this property.

By the previous study, H is non-empty and the same argument of perturbation gives that H is open. We now show that H is closed in the interior of $G_{0,\Gamma}(p/q, h, d)$. If $((D, \beta)_n)_{n\ge0} \subset H$ converges to (D, β) in the interior of $G_{0,\Gamma}(p/q, h, d)$, then the expressions associated to $(D, \beta)_n$ in the inequalities defining H converge to the formal expressions associated to (D, β) which verify strict inequalities. Thus the expressions of D and $(T^k0^{\pm})_{0\le k\le q-1}$ corresponding to $(D, \beta)_n$ are still valid for (D, β). Consequently, H is closed in the interior of $G_{0,\Gamma}(p/q, h, d)$ and then H is exactly the interior of $G_{0,\Gamma}(p/q, h, d)$.

As a corollary we get that if $(h, d) \neq (h', d')$ then $G_{0,\Gamma}(p/q, h, d) \cap G_{0,\Gamma}(p/q, h', d') = \emptyset$. This concludes the proof of the theorem. □

Remark. The case $m = 0$ of Theorem 4.14 and when the maps are linear is due to Bugeaud and Conze [**5**]. The case when $m = 1$ in a linear context is also treated in this article but with $\tilde{T}$ continuous at D_1 and $\tilde{T}(D_1) = 1$. Introducing contraction coefficients $0 \le \gamma_0, \gamma_1 < 1$, the map $\tilde{T}$ can be written as

$$\tilde{T}x = \gamma_i x + \beta_i, \quad \text{on } [D_i, D_{i+1}).$$

The previous conditions can be written as $\gamma_0 D_1 + \beta_0 = \gamma_1 D_1 + \beta_1 = 1$ or equivalently $D_1 = (1 - \beta_0)/\gamma_0$ and $\beta_1 = 1 - (1 - \beta_0)\gamma_1/\gamma_0$. Next, choosing β_0 as a parameter as in [**5**], in (11) only the first line remains and one deduces the following result, correcting the formula of [**5**].

THEOREM 4.15. *Let $m = 1$ and $\tilde{T}$ be as above. Set $T = \tilde{T} \bmod(1)$ and let, for $0 \le p/q < 1$ with $p \wedge q = 1$, $(\varepsilon_k)_{0\le k\le q-1}$ be the code of $(R^k_{p/q}0)_{0\le k\le q-1}$ in the intervals $\{[0, 1 - p/q), [1 - p/q, 1)\}$. Then $\tau(T) = p/q$ if and only if*

$$\beta_0 \in \left[\frac{\sum_{k=0}^{q-1} \varepsilon_k(\gamma_{\varepsilon_k} \cdots \gamma_{\varepsilon_{q-2}})}{\sum_{k=0}^{q-1}(\gamma_{\varepsilon_k} \cdots \gamma_{\varepsilon_{q-2}})}, \frac{1 + \gamma_{\varepsilon_{q-1}} \cdots \gamma_{\varepsilon_1}(1 - \gamma_{\varepsilon_0}) + \sum_{k=1}^{q-2} \varepsilon_k(\gamma_{\varepsilon_k} \cdots \gamma_{\varepsilon_{q-2}})}{1 + \gamma_{\varepsilon_{q-1} \cdots \gamma_{\varepsilon_1}} + \sum_{k=1}^{q-2}(\gamma_{\varepsilon_k} \cdots \gamma_{\varepsilon_{q-2}})} \right].$$

In this case T has a unique periodic orbit and it has period q.

https://doi.org/10.1017/S0143385705000386 Published online by Cambridge University Press

Remark. Consider the question of counting the periodic orbits when $\tau(T) \in \mathbb{Q}$. As in Lemma 2.3 the set $\mathcal{D} = \{T^{-k}D_i \mid 0 \le k \le k_i, 1 \le i \le m\}$ induces a stable partition on the circle. There are then less than $m+1$ periodic orbits. We now give an algorithm for counting the periodic orbits.

PROPOSITION 4.16. *Let $0 < p/q < 1$ with $p \wedge q = 1$. Let $(D,\beta) \in \mathcal{C}$ verify $\tau(D,\beta) = p/q$. Using Theorem 4.14, let $I = \{0 \le i \le m \mid (D,\beta) \in F_i(p/q)\}$. For $i \in I$ let h^i and d^i be such that $(D,\beta) \in G_{i,\Gamma}(p/q, h^i, d^i)$.*

(1) *For $i \in I$ set $A_i = \{j \in I \mid d^i_j$ is even $\}$. If $A_i \neq \emptyset$ there is exactly one $j \in A_i$ such that*

$$n_{d^i_j/2} = \min\{n_{d^i_k/2} \mid k \in A_i\}.$$

Then write $i \to j$. Conversely, if $j \in I$ there is at most one $i \in I$ such that $i \to j$.

(2) *The set I decomposes into disjoint closed cycles of length $\le q$ for the relation '$\to$'. The number of distinct periodic orbits for T is the number of cycles in I.*

Proof. Let us reason modulo 1. Using Proposition 4.2 let x be a fixed point for $\tilde{T}^q - p$. The number of distinct q-periodic orbits is the number of fixed points of $\tilde{T}^q - p$ in the interval $[x, \tilde{T}^{n_1}x)$. This is also the number of discontinuities $w \in [x, T^{n_1}x)$ of $\tilde{T}^q - p$ satisfying

$$\tilde{T}^q(w^-) \le p + w \le \tilde{T}^q(w). \tag{13}$$

Next, let $x < y \le T^{n_1}x$ be the fixed point consecutive to x. Call $w_0 \in [x,y)$ the unique discontinuity of $\tilde{T}^q - p$ satisfying (13). We write $w_0 = \tilde{T}^{-l_0}D_{i_0}$ for some D_{i_0} and a minimal $0 \le l_0 \le q-1$. Then D_{i_0} is uniquely determined as the $(D_j)_{0\le j\le m}$ are distinct. If $l_0 > 0$ then by minimality of l_0, for $0 \le k < l_0$, the point $\tilde{T}^k(\tilde{T}^{-l_0}(D_{i_0}))$ does not belong to the $(D_j)_{0\le j\le m}$. Successive iterations give that, for $0 \le k \le l_0$,

$$\tilde{T}^q((\tilde{T}^k(w_0))^-) \le p + \tilde{T}^k(w_0) \le \tilde{T}^q(\tilde{T}^k(w_0)).$$

Consequently, for $0 \le k \le l_0$ the unique discontinuity of $\tilde{T}^q - p$ in $[\tilde{T}^k(x), \tilde{T}^k(y))$ satisfying (13) is $\tilde{T}^k(w_0)$. For $k = l_0$ we have $\tilde{T}^q(D^-_{i_0}) \le p + D_{i_0} \le \tilde{T}^q(D_{i_0})$ giving that $i_0 \in I$. Another application of $\tilde{T}$ leads to

$$\tilde{T}^q(\tilde{T}(D^-_{i_0})) \le p + \tilde{T}(D^-_{i_0}) < p + \tilde{T}(D_{i_0}) \le \tilde{T}^q(\tilde{T}(D_{i_0})).$$

Consequently, if w_1 is the discontinuity of $\tilde{T}^q - p$ satisfying (13) in $[\tilde{T}^{l_0+1}(x), \tilde{T}^{l_0+1}(y))$ then $\tilde{T}(D^-_{i_0}) \le w_1 \le \tilde{T}(D_{i_0})$. Write $w_1 = \tilde{T}^{-l_1}D_{i_1}$ with a minimal $0 \le l_1 \le q-1$ and some uniquely determined D_{i_1}. As above $i_1 \in I$ and, for $0 \le k \le l_1$, $\tilde{T}^k(w_1)$ is the unique discontinuity of $\tilde{T}^q - p$ verifying (13) in $[\tilde{T}^{k+1+l_0}(x), \tilde{T}^{k+1+l_0}(y))$.

Now note that $D_{i_1} \in [\tilde{T}^{l_1+1}(D^-_{i_0}), \tilde{T}^{l_1+1}(D_{i_0})]$, whose associated number is even, following Definition 4.13. As $[\tilde{T}^u(D^-_{i_0}), \tilde{T}^u(D_{i_0})] \subset [\tilde{T}^{l_0+u}(x), \tilde{T}^{l_0+u}(y))$ for $1 \le u \le l_1+1$, we get that i_1 is the only $j \in J$ such that $D_j \in [T^u(D^-_{i_0}), T^u(D_{i_0}))$, for some $1 \le u \le l_1+1$. Indeed D_i must satisfy (13) and l_1 is minimal. We thus get $i_0 \to i_1$.

Repeating this procedure we obtain a sequence $(i_0 \to i_1 \to \cdots \to i_t \to i_0) \subset I$ which is a cycle since $[T^q(x), T^q(y)) = [x,y)$. This also gives $t+1 \le q$. Proceeding in this way for each pair of consecutive q-periodic points in $[x, T^{n_1}x]$ we obtain non-empty

https://doi.org/10.1017/S0143385705000386 Published online by Cambridge University Press

and disjoint cycles in I in bijection with the periodic orbits. Finally, if $i \in I$ then D_i verifies (13) and belongs to some iterate of the interval determined by two consecutive periodic points in $[x, T^{n_1}x]$. Thus i belongs to a cycle. This concludes the proof of the proposition. □

Remark. The case $\tau(T) \notin \mathbb{Q}$ can be considered using the monotony of the rotation number for fixed D. As in Theorem 4.14 the set of parameters (D, β) verifying $\tau(D, \beta) = r$ can be decomposed into a union of polyhedrons, each polyhedron of codimension at least one. However, this union is not denumerable as soon as $m \geq 1$. More details can be found in [**4**].

4.4. *Pictures.* We provide a numerical illustration of the decomposition in Theorem 4.14 and Proposition 4.16. For the sake of simplicity we restrict ourselves to the linear case. Annotated pictures for the case $m = 0$ can then be found in Bugeaud and Conze [**5**]. In the context of Definition 4.4, we consider here the situation when $m = 1$.

Consider a map $T = \tilde{T} \bmod(1)$, where

$$\tilde{T} = \begin{cases} \dfrac{x}{2} + \beta_0, \text{ on } [0, D_1), \\ \dfrac{x}{2} + \beta_1, \text{ on } [D_1, 1). \end{cases}$$

The parameters are $(D_1, \beta_0, \beta_1) \in \mathbb{R}^3$. In the figures, these are respectively represented on the x-axis, y-axis and the z-axis. We look at the case when $\tau(D_1, \beta_0, \beta_1) = 1/3$. Recall that there are one or two 3-periodic orbits:

(1) $F_0(1/3)$;
(2) $F_1(1/3)$;
(3) $E(1/3) = F_0(1/3) \cup F_1(1/3)$;
(4) parameters giving two distinct 3-periodic orbits.

4.5. *Injective quasi-contractions with three pieces on the interval.* We apply the study on order-preserving quasi-contractions on the circle to the analysis of a class of injective locally increasing quasi-contractions with three pieces on $[0, 1)$, as in Definition 2.1. Writing as $(0, 1, 2)$ the three segments, we restrict our argument to the case when the images segments are ordered as $(2, 1, 0)$. A classical construction shows that such an injective quasi-contraction can be seen as the first return map of an order-preserving quasi-contraction on the circle with four pieces.

Definition 4.17.

(1) Let T be defined as in Definition 2.1 with $m = 2$, $D = (D_i)_{1 \leq i \leq 2} \in \mathcal{D}$ and a set of increasing γ-Lipschitzian maps $\Gamma_T = (\Gamma_i)_{0 \leq i \leq 2}$, where $0 \leq \gamma < 1$. Denote by $\mathcal{C}'$ the component of $\mathcal{C}$ corresponding to the fact that the image segments are ordered as $(2, 1, 0)$.
(2) Set $u_0 = -\sqrt{\Gamma_1(D_2) - \Gamma_1(D_1)}$, $u_1 = 0$, $u_2 = D_1$ and $u_3 = D_2$. Let $R : [u_0, 1) \longmapsto [u_0, 1)$ be the order-preserving transformation with four pieces defined by:

https://doi.org/10.1017/S0143385705000386 Published online by Cambridge University Press




FIGURE 1. $F_0(1/3)$.




FIGURE 2. $F_1(1/3)$.

https://doi.org/10.1017/S0143385705000386 Published online by Cambridge University Press




FIGURE 3. $\tau(D, \beta) = 1/3$.




FIGURE 4. $\tau(D, \beta) = 1/3$ and two distinct 3-periodic attractors.

https://doi.org/10.1017/S0143385705000386 Published online by Cambridge University Press

- $\Gamma_R = (\Gamma', \Gamma_0, \Gamma'', \Gamma_2)$;
- $R_{|[u_1,u_2)\cup[u_3,1)} = T_{|[u_1,u_2)\cup[u_3,1)}$;
- the pair (Γ', Γ'') (not unique) is such that Γ'' contracts $[u_1, u_2)$ on $[u_0, u_1)$, and Γ' contracts $[u_0, u_1)$ on $[\Gamma_1(D_1), \Gamma_1(D_2))$ and $\Gamma' \circ \Gamma'' = \Gamma_1$ on $[u_1, u_2)$.

Then T is the first return map of R in $[0, 1)$: if $x \in [0, 1)$ and $t(x) = \inf\{n \geq 1 \mid R^n(x) \in [0, 1)\}$ then $T(x) = R^{t(x)}(x)$. Note that $t(x) \in \{1, 2\}$, $x \in [0, 1)$.

We then obtain the following theorem.

THEOREM 4.18. *For T defined as in Definition 4.17, only two excluding cases can occur.*

(1) *There exists a periodic orbit. Then T admits at most two distinct periodic orbits and no diffusive T-invariant measure.*

(2) *The map T has no periodic orbit. Then T is uniquely ergodic, measurably conjugated and topologically semi-conjugated to a possibly degenerated interval exchange transformation with three pieces and permutation* $(0, 1, 2) \longrightarrow (2, 1, 0)$ *on* $[0, 1)$.

Moreover, each case is non-empty, but the first one holds for at least an open set of full Lebesgue measure in $\mathcal{C}'$. The subcases with exactly one or two periodic orbits have non-empty interior in $\mathcal{C}'$.

Proof. First, an example with two periodic orbits is when the interval i is sent in the interval $\sigma(i)$, where σ is the permutation $(0, 1, 2) \longrightarrow (2, 1, 0)$. This provides a fixed point and a 2-periodic orbit. If all intervals are sent in interval 2, then there is a unique fixed point. Both situations are clearly realized by open subsets in $\mathcal{C}'$.

Next, observe that T admits a periodic orbit if and only if R admits one, that is if and only if $\tau(R) \in \mathbb{Q}$. We show that in this case there are less than two periodic orbits. Denote by (D', β) the parameters of R. As in Proposition 4.16 and after renormalization let I be the set of integers i such that $(D', \beta) \in F_i(p/q)$.

Note that $0 \in [R(D'_2)^-, R(D'_2)]$ and $D'_1 \in [R(D'_3)^-, R(D'_3)]$. Thus if both 0 and D'_2 lay in I they belong to the same cycle of length ≥ 2. The same holds for D'_1 and D'_3. From Proposition 4.16 the number of periodic orbits of R is ≤ 2. This is also true for T.

Next, the situation when R has no periodic orbit is non-empty. Indeed, for some configurations, R admits a fixed point and for other configurations there is no fixed point. This gives respectively $\tau(R) = 0$ and $0 < \tau(R) < 1$. The continuity of the rotation number implies that irrational values must be taken. This then provides parameters for which R (and thus T) has no periodic point.

In this case, Proposition 4.3 implies that R and therefore T are uniquely ergodic. Then let h be a linearizing map as in Proposition 4.3(2) that topologically semi-conjugates R with $R_{\tau(R)}$. Then T is topologically semi-conjugated and measurably conjugated to the first return map of $R_{\tau(R)}$ into $h([D'_1, 1))$. This map is a possibly degenerated interval exchange transformation with three pieces associated to the permutation $(0, 1, 2) \longrightarrow (2, 1, 0)$. □

Acknowledgements. This paper is a part of my PhD thesis and I would like to thank J.-P. Conze for valuable discussions on the subject.

https://doi.org/10.1017/S0143385705000386 Published online by Cambridge University Press

REFERENCES

[1] L. Alsed, J. M. Gambaudo and P. Mumbr. Stable heteroclinic cycles and symbolic dynamics. *Chaos* **4**(2) (1994), 407–419.

[2] T. Bousch. Le poisson n'a pas d'arêtes. *Ann. Inst. H. Poincaré Probab. Statist.* **36**(4) (2000), 489–508.

[3] T. Bousch and J. Mairesse. Asymptotic height optimization for topical IFS, Tetris heaps, and the finiteness conjecture. *J. Amer. Math. Soc.* **15**(1) (2002), 77–111 (electronic).

[4] J. Brémont. Marches aléatoires en milieu aléatoire sur $\mathbb{Z}$; dynamique d'applications localement contractantes sur le Cercle. *Thèse de doctorat*, Université de Rennes I, 2002.

[5] Y. Bugeaud and J.-P. Conze. Dynamics of some contracting linear functions modulo 1. *Noise, Oscillators and Algebraic Randomness (Chapelle des Bois, 1999)*. Springer, Berlin, 2000, pp. 379–387.

[6] S. Bullett and P. Sentenac. Ordered orbits of the shift, square roots and the devil's staircase. *Math. Proc. Cambridge Philos. Soc.* **115**(3) (1994), 451–481.

[7] G. Contreras, A. Lopes and P. Thieullen. Lyapunov minimizing measures for expanding maps of the Circle. *Ergod. Th. & Dynam. Sys.* **17**(21) (2001), 1379–1409.

[8] J.-P. Conze and Y. Guivarc'h. Croissance des sommes ergodiques et principe variationnel. *Technical Report*, Université de Rennes 1, 1990.

[9] O. Feely and L. Chua. The effect of integrator leak in sigma-delta modulation. *IEEE Transactions on Circuits and Systems* **18**(11) (1991), 1293–1310.

[10] J.-M. Gambaudo, P. Glendinning and C. Tresser. The gluing bifurcation. I. Symbolics dynamics of the closed curves. *Nonlinearity* **1**(1) (1988), 203–214.

[11] J.-M. Gambaudo and C. Tresser. On the dynamics of quasi-contractions. *Bol. Soc. Brasil. Mat* **19**(1) (1988), 61–114.

[12] B. Hunt and G. Yuan. Optimal orbits of hyperbolic systems. *Nonlinearity* **12** (1999), 1207–1224.

[13] O. Jenkinson. Frequency locking on the boundary of the barycentre set. *Experiment. Math.* **9**(2) (2000), 309–317.

[14] A. Katok and B. Hasselblatt. *Introduction to the Modern Theory of Dynamical Systems*. Cambridge University Press, Cambridge, 1995.

[15] M. Keane. Interval exchange transformations. *Math. Z.* **141** (1975), 25–31.

[16] R. Mané. Generic properties and problems of minimizing measures of Lagrangian systems. *Nonlinearity* **9**(2) (1996), 273–310.

https://doi.org/10.1017/S0143385705000386 Published online by Cambridge University Press