Ergod. Th. & Dynam. Sys. (2021), **41**, 846–868 doi:10.1017/etds.2019.103
© *Cambridge University Press, 2019*

Frequent universality criterion and densities

R. ERNST† and A. MOUZE‡

† *Romuald Ernst, LMPA, Centre Universitaire de la Mi-Voix, Maison de la Recherche Blaise-Pascal, 50 rue Ferdinand Buisson, BP 699, 62228 Calais Cedex, France (e-mail: ernst.r@math.cnrs.fr)*

‡ *Augustin Mouze, Laboratoire Paul Painlevé, UMR 8524, Cité Scientifique, 59650 Villeneuve d'Ascq, France (e-mail: Augustin.Mouze@math.univ-lille1.fr)*

(*Received* 4 *July* 2018 *and accepted in revised form* 8 *October* 2019)

Abstract. We improve a recent result by giving the optimal conclusion both to the frequent universality criterion and the frequent hypercyclicity criterion using the notion of A-densities, where A refers to some weighted densities sharper than the natural lower density. Moreover, we construct an operator which is logarithmically frequently hypercyclic but not frequently hypercyclic.

Key words: frequent universality, weighted densities, hypercyclicity
2010 Mathematics Subject Classification: 47A16 (Primary); 37B50 (Secondary)

1. *Introduction and notation*

We denote by $\mathbb{N}$ the set of positive integers. Let (α_k) be a non-negative sequence with $\sum_{k\geq 1}\alpha_k = +\infty$. Let us consider the associated admissible matrix $A = (\alpha_{n,k})$ given by

$$\alpha_{n,k} = \begin{cases} \alpha_k \Big/ \left(\sum_{j=1}^{n} \alpha_j\right) & \text{for } 1 \leq k \leq n, \\ 0 & \text{otherwise.} \end{cases}$$

We know that every regular summability matrix A, so any admissible matrix, gives a density $\underline{d}_A$ on subsets of $\mathbb{N}$, called the lower A-density [**9**].

Definition 1.1. For a regular matrix $A = (\alpha_{n,k})$ with non-negative coefficients and a set $E \subset \mathbb{N}$, the lower A-density of E, denoted by $\underline{d}_A(E)$, is defined as

$$\underline{d}_A(E) = \liminf_n \left(\sum_{k=1}^{+\infty} \alpha_{n,k} \mathbb{1}_E(k)\right),$$

and the associated upper A-density, denoted by $\overline{d}_A(E)$, is given by the equality $\overline{d}_A(E) = 1 - \underline{d}_A(\mathbb{N} \setminus E)$.

https://doi.org/10.1017/etds.2019.103 Published online by Cambridge University Press

Moreover, it is well known [**9**] that the upper A-density of any set $E \subset \mathbb{N}$ is given by $\overline{d}_A(E) = \limsup_n (\sum_{k=1}^{+\infty} \alpha_{n,k} \mathbb{1}_E(k))$.

Let X, Y be Fréchet spaces. In the present paper we are interested in the universality of sequences of operators (T_n), $T_n : X \to Y$, in the following sense: a sequence (T_n) is said to be *universal* if there exists $x \in X$ such that the set $\{T_n x \ : \ n \in \mathbb{N}\}$ is dense in Y. Such a vector x is called a universal vector for (T_n). When the sequence (T_n) is given by the iterates of a single operator T, that is, $(T_n) = (T^n)$ and $Y = X$, the notion of universality reduces to the well-known one of *hypercyclicity*, which is a central notion in linear dynamics. Now the following definition extends that of frequent universality and quantifies how often the orbit of a universal vector visits every non-empty open set. For any $x \in X$ and any subset $U \subset Y$, we set $N(x, U) := \{n \in \mathbb{N} : T_n x \in U\}$.

Definition 1.2. Let $A = (\alpha_k / \sum_{j=1}^{n} \alpha_j)$ be an admissible matrix. A sequence of operators (T_n), $T_n : X \to Y$, is called *A-frequently universal* if there exists $x \in X$ such that, for any non-empty open set $U \subset X$, the set $N(x, U)$ has positive lower A-density.

Notice that if $\alpha_k = 1$, $k = 1, 2, \ldots$, the matrix A corresponds to the Cesàro matrix, the lower A-density $\underline{d}_A$ coincides with the natural lower density $\underline{d}$ and we recover the notion of *frequent universality* (or *frequent hypercyclicity* in the case of sequences (T^n)). Bonilla and Grosse-Erdmann have derived a sufficient condition for a sequence of operators to be frequently universal [**6**, **7**]. Their criterion extends the frequent hypercyclicity criterion given by Bayart and Grivaux [**1**].

THEOREM 1.3. (Frequent universality criterion) *Let X be a Fréchet space, Y a separable Fréchet space and $T_n : X \to Y$, $n \in \mathbb{N}$, continuous linear mappings. Suppose that there are a dense subset Y_0 of Y and mappings $S_n : Y_0 \to X$, $n \in \mathbb{N}$, such that:*

(1) $\sum_{n=1}^{k} T_k S_{k-n} y$ *converges unconditionally in Y, uniformly in $k \in \mathbb{N}$, for all $y \in Y_0$;*

(2) $\sum_{n=1}^{+\infty} T_k S_{k+n} y$ *converges unconditionally in Y, uniformly in $k \in \mathbb{N}$, for all $y \in Y_0$;*

(3) $\sum_{n=1}^{+\infty} S_n y$ *converges unconditionally in X, for all $y \in Y_0$;*

(4) $T_n S_n y \to y$ *for all $y \in Y_0$.*

Then the sequence (T_n) is frequently universal.

It is well known that there exist frequently universal sequences of operators which do not satisfy this criterion [**2**]. A natural question arises: if the sequence (T_n) fulfills the hypotheses of the criterion, for which admissible matrices A can one conclude that (T_n) is A-frequently universal? Before answering, let us introduce some useful notation. We denote by $\log^{(s)}$ the iterated logarithmic function $\log \circ \log \circ \cdots \circ \log$ where log appears s times. In the sequel, we shall need the following admissible matrices:

(1) $A_r = (e^{k^r} / \sum_{j=1}^{n} e^{j^r})$ for $0 \leq r \leq 1$;

(2) $\widetilde{D}_s = (\alpha_k / \sum_{j=1}^{n} \alpha_j)$ given by the coefficients $\alpha_k = e^{k/(\log^{(s)}(k))}$ for k large enough and $s \geq 1$;

(3) $\widetilde{B}_s = (\alpha_k / \sum_{j=1}^{n} \alpha_j)$ given by the coefficients $\alpha_k = e^{k/(\log(k) \log^{(s)}(k))}$ for k large enough and $s \geq 2$;

(4) $B_r = (\alpha_k / \sum_{j=1}^{n} \alpha_j)$ given by the coefficients $\alpha_k = e^{k/\log^r(k)}$ for k large enough and $r \geq 1$;

(5) the matrix $L = (k^{-1} / \sum_{j=1}^{n} j^{-1})$ associated to the logarithmic density $\underline{d}_{\log}$.

https://doi.org/10.1017/etds.2019.103 Published online by Cambridge University Press

Lemma 2.8 of [**8**] ensures that we have, for any $2 \leq s \leq s'$, $1 \leq t \leq t'$, $0 < r < r' < 1$ and for all subsets $E \subset \mathbb{N}$,

$$\underline{\mathbf{d}}_{\mathbf{A_1}}(\mathbf{E}) \leq \underline{d}_{\widetilde{D}_{s'}}(E) \leq \underline{d}_{\widetilde{D}_s}(E) \leq \underline{\mathbf{d}}_{\mathbf{B_1}}(\mathbf{E}) \leq \underline{d}_{\widetilde{B}_{s'}}(E) \leq \underline{d}_{\widetilde{B}_s}(E),$$
$$\underline{d}_{\widetilde{B}_s}(E) \leq \underline{d}_{B_t}(E) \leq \underline{d}_{B_{t'}}(E) \leq \underline{d}_{A_{r'}}(E) \leq \underline{d}_{A_r}(E) \leq \underline{\mathbf{d}}(\mathbf{E}) \leq \underline{d}_{\log}(E).$$

Recently the authors have shown that under the assumptions of the frequent universality criterion a sequence (T_n) is automatically $\widetilde{B}_s$-frequently universal for any positive integer $s \geq 1$ [**8**]. Actually the statement is written in the context of $\widetilde{B}_s$-frequent hypercyclicity, but it is easy to check that the proof works along the same lines in the case of $\widetilde{B}_s$-frequent universality. First we improve this result by showing that a sequence of operators which satisfies the frequent universality criterion is necessarily B_1-frequently universal. Then a technical modification of the proof allows us to show that such a sequence of operators is necessarily $\widetilde{D}_s$-frequently universal for any $s \in \mathbb{N}$. On the other hand, we establish that an operator $T : X \to X$ cannot be A_1-frequently hypercyclic. Notice that this result was already proved whenever X was a Banach space [**8**, Proposition 3.7]. Based on these results, this paper determines exactly what quantifies the frequent universality criterion in terms of weighted densities of the return sets. The proof of this result essentially uses combinatorial arguments. Furthermore, in [**8**] the authors exhibited a frequently universal (hypercyclic) operator which is not A_r-frequently hypercyclic for any $0 \leq r < 1$ (in particular, it does not satisfy the above criterion). Conversely, using similar ideas, we build an operator which is log-frequently hypercyclic (i.e. L-frequently hypercyclic) but not frequently hypercyclic. Hence one can find hypercyclic operators whose orbit of some universal vectors visits rather often, in the sense of suitable lower A-densities, every non-empty set but not sufficiently to be frequently hypercyclic. As far as we know, this is the first example in this direction using lower densities.

The paper is organized as follows. In §§2 and 3 we construct specific sequences of integers which will allow us to establish that the frequent universality criterion gives a stronger result. Then we will show that this new result is the best possible. Finally, in §4, we exhibit an example, inspired by [**4**], of an operator which is L-frequently hypercyclic but not frequently hypercyclic.

2. *Construction of a specific sequence*

Let X be a Fréchet space, Y a separable Fréchet space and $T_n : X \to Y$, $n \in \mathbb{N}$, continuous linear mappings. In this section we will establish the following implication: if the sequence (T_n) satisfies the frequent universality criterion, then (T_n) is B_1-frequently universal. This result will already be an improvement of [**8**, Theorem 4.12]. To do this, we begin with an important remark: a careful examination of the proofs of the frequent universality criterion shows that it suffices to find a sequence (δ_k) of integers and (n_k) an increasing sequence of integers such that

$$|n_k - n_l| \geq \delta_k + \delta_l \quad \text{whenever } k \neq l \text{ and for any } p \geq 1, \ \underline{d}(\{n_k : \delta_k = p\}) > 0.$$

We refer the reader to [**3**, Lemma 6.19 and Theorem 6.18] and [**6**, **7**]. An easy modification of the proof of the criterion allows us to obtain the A-frequent universality provided that $\underline{d}_A(\{n_k : \delta_k = p\}) > 0$ (see [**8**]).

https://doi.org/10.1017/etds.2019.103 Published online by Cambridge University Press

First of all, let us recall the following useful lemma to estimate the lower A-density of a given sequence (n_k) [**8**, Lemma 2.7].

LEMMA 2.1. *Let (α_k) be a non-negative sequence such that $\sum_{k\in\mathbb{N}} \alpha_k = +\infty$. Assume that the sequence $(\alpha_n / \sum_{j=1}^{n} \alpha_j)$ converges to zero as n tends to $+\infty$. Let (n_k) be an increasing sequence of integers. Then we have*

$$\underline{d}_A((n_k)_k) = \liminf_{k\to+\infty}\left(\frac{\sum_{j=1}^{k} \alpha_{n_j}}{\sum_{j=1}^{n_k} \alpha_j}\right),$$

where $\underline{d}_A$ is the A-density given by the admissible matrix $A = (\alpha_k / \sum_{j=1}^{n} \alpha_j)$.

In what follows we will build suitable sequences (δ_k) and (n_k). For every positive integer k, we define δ_k and L_0 as follows: $\delta_k = l$ and $L_0 = l - 1$ where l is the place of the first zero in the dyadic representation of k. For example, for $k = 11$ (i.e. $k = 1.2^0 + 1.2^1 + 0.2^2 + 1.2^3$) we have $\delta_k = 3$ and $L_0 = 2$. Then we construct the following increasing sequence (n_k) of positive integers by setting

$$n_1 = 2 \quad \text{and} \quad n_k = 2\sum_{i=1}^{k-1} \delta_i + \delta_k \quad (k \geq 2). \tag{2.1}$$

LEMMA 2.2. *With the above notation, we have, for every $m \geq 1$,*

$$n_{2^m} = 4.2^m - 3 \quad \textit{and} \quad n_{2^m-1} = 4.2^m - m - 5.$$

Proof. We have $n_1 = n_{2-1} = 2 = 4.2 - 1 - 5$ and $n_2 = 5 = 4.2 - 3$. We set, for $k \geq 2$,

$$n'_k = \sum_{i=1}^{k} \delta_i = \frac{n_k + \delta_k}{2}.$$

Let us consider the sets $\Delta_j^{(m)} = \{1 \leq k \leq 2^m - 1;\ \delta_k = j\}$, for $m \geq 1$ and $j = 1, \ldots, m + 1$. We get, for $m \geq 2$,

$$n'_{2^m-1} = \sum_{i=1}^{2^m-1} \delta_i = \sum_{j=1}^{m+1} j\#\Delta_j^{(m)}.$$

Thus it suffices to compute the cardinality of the sets $\Delta_j^{(m)}$. For $j = m + 1$, there is only one possibility to obtain $\delta_l = m + 1$ given by $l = (1, \ldots, 1, 0, \ldots)$ with m ones (i.e. $l = 2^m - 1$). For $2 \leq j \leq m$, to obtain $\delta_l = j$, l necessarily has dyadic representation

$$(\underbrace{1, 1, \ldots, 1}_{\text{length } j-1}, 0, \underbrace{\star, \star, \ldots, \star, \star}_{\text{length } m-j}, 0, 0, \ldots).$$

Thus we have 2^{m-j} possible choices. Finally, to obtain $\delta_l = 1$, l necessarily has dyadic representation

$$(0, \underbrace{\star, \star, \ldots, \star, \star}_{\text{length } m-1}, 0, 0, \ldots),$$

https://doi.org/10.1017/etds.2019.103 Published online by Cambridge University Press

and taking into account $l \neq 0$, we have $2^{m-1} - 1$ possible choices. Therefore we get, for all $m \geq 2$,

$$n'_{2^m-1} = (2^{m-1} - 1) + \sum_{j=2}^{m} j2^{m-j} + (m+1).$$

An easy calculation gives

$$n'_{2^m-1} = 4.2^{m-1} - 2 \quad \text{for } m \geq 2.$$

Therefore we deduce, for all $m \geq 1$,

$$n_{2^m-1} = 4.2^m - 5 - m.$$

Finally, we obtain, for every integer $m \geq 2$,

$$n_{2^m} = n_{2^m-1} + \delta_{2^m-1} + \delta_{2^m} = n_{2^m-1} + (m+1) + 1,$$

which leads to

$$n_{2^m} = 4.2^m - 3 \quad \text{for } m \geq 1. \qquad \square$$

LEMMA 2.3. *With the above notation, we have, for every positive integer k which has dyadic representation $k := 2^n + \sum_{i=0}^{n-1} \alpha_i 2^i$, $2^n < k < 2^{n+1} - 1$, $\alpha_i \in \{0, 1\}$,*

$$n_{k-2^n} = 2 \sum_{i=1+2^n}^{k-1} \delta_i + \delta_k.$$

Proof. It suffices to observe that every integer i with $1 + 2^n \leq i \leq k < 2^{n+1} - 1$ has dyadic representation

$$(\underbrace{\star, \ldots, \star}_{\text{length } n}, 1, 0 \ldots)$$

with at least a zero between the first and the nth position (since $k \neq 2^{n+1} - 1$). Therefore we have $\delta_i = l$ for some $1 \leq l \leq n$ and

$$i - 2^n = (\underbrace{1, \ldots, 1}_{\text{length } l-1}, 0, \underbrace{\star, \ldots, \star}_{\text{length } n-l}, 0, 0 \ldots).$$

We deduce for $1 + 2^n \leq i \leq k$ that $\delta_{i-2^n} = \delta_i$. Therefore we get, for every positive integer k with dyadic representation $k := 2^n + \sum_{i=0}^{n-1} \alpha_i 2^i$, $2^n < k < 2^{n+1} - 1$,

$$n_{k-2^n} = 2 \sum_{i=1}^{k-1-2^n} \delta_i + \delta_{k-2^n} = 2 \sum_{i=1+2^n}^{k-1} \delta_i + \delta_k. \qquad \square$$

LEMMA 2.4. *With the above notation, we have, for every positive integer k which has dyadic representation $k := \sum_{i=1+L_0}^{n} \alpha_i 2^i + 2^{L_0} - 1$ with $n \geq 1$, $\alpha_n = 1$ and $\alpha_i \in \{0, 1\}$, $i = 1 + L_0, \ldots, n$,*

$$n_k = \begin{cases} \displaystyle\sum_{i=1+L_0}^{n} \alpha_i n_{2^i} + \sum_{i=1+L_0}^{n} \alpha_i + n_{2^{L_0}-1} & \text{if } 1 \leq L_0 \leq n-1, \\ \displaystyle\sum_{i=1}^{n} \alpha_i n_{2^i} + \sum_{i=1}^{n} \alpha_i - 1 & \text{if } L_0 = 0. \end{cases}$$

https://doi.org/10.1017/etds.2019.103 Published online by Cambridge University Press

Proof. For $L_0 = 0$, we have $k = \sum_{i=1}^{n-1} \alpha_i 2^i + 2^n$, with $\alpha_n = 1$ and $\alpha_i \in \{0, 1\}$. For $k = 2^n$, we have $\alpha_i = 0$, $i = 1, \ldots, n-1$ and we can write $n_k = n_{2^n} + \sum_{i=1}^{n} \alpha_i - 1$ and the result holds. Otherwise, we define the sequence $1 \le l_1 < l_2 < \cdots < l_m \le n-1$, satisfying $\alpha_{l_j} = 1$, for $j = 1, \ldots, m$, and $\alpha_i = 0$ for $i \notin \{l_1, \ldots, l_m\}$. Taking into account Lemma 2.3, we write

$$\begin{aligned} n_k &= \left(2 \sum_{i=1}^{2^n-1} \delta_i + \delta_{2^n}\right) + \delta_{2^n} + \left(2 \sum_{i=1+2^n}^{k-1} \delta_i + \delta_k\right) \\ &= n_{2^n} + 1 + n_{k-2^n} \\ &= n_{2^n} + \alpha_n + n_{k-2^n}. \end{aligned} \tag{2.2}$$

We get $k - 2^n = \sum_{j=1}^{m} 2^{l_j}$, and a calculation similar to (2.2) leads to

$$n_k = n_{2^n} + \alpha_n + n_{2^{l_m}} + \alpha_{l_m} + n_{k-2^n-2^{l_m}} = \alpha_n n_{2^n} + \alpha_{l_m} n_{2^{l_m}} + \alpha_n + \alpha_{l_m} + n_{k-2^n-2^{l_m}}.$$

By repeating the reasoning we obtain

$$n_k = \sum_{j=2}^{m} n_{2^{l_j}} + n_{2^n} + \sum_{j=2}^{m} \alpha_{l_j} + \alpha_n + n_{2^{l_1}}.$$

Finally, since we have $n_{2^l_1} = n_{2^l_1} + \alpha_{l_1} - 1$, we can write

$$n_k = \sum_{i=1}^{n} \alpha_i n_{2^i} + \sum_{i=1}^{n} \alpha_i - 1.$$

Turning to the case of $1 \le L_0 \le n-1$, if we have $\alpha_i = 0$, for $i = 1 + L_0, \ldots, n-1$, we get $k - 2^n = 2^{L_0} - 1$ and $n_k = n_{2^n} + \alpha_n + n_{2^{L_0-1}}$ which was to be proved. Otherwise, we set $m = \max(1 + L_0 \le i \le n-1; \alpha_i = 1)$. We have $k - 2^n = 2^m + \sum_{i=1+L_0}^{m-1} \alpha_i 2^i + 2^{L_0} - 1$, and a calculation similar to (2.2) leads to

$$n_k = n_{2^n} + n_{2^m} + \alpha_n + \alpha_m + n_{k-2^n-2^m} = \alpha_n n_{2^n} + \alpha_m n_{2^m} + \alpha_n + \alpha_m + n_{k-2^n-2^m}.$$

By repeating the reasoning we obtain

$$n_k = \sum_{i=1+L_0}^{n} \alpha_i n_{2^i} + \sum_{i=1+L_0}^{n} \alpha_i + n_{2^{L_0-1}}. \qquad \square$$

LEMMA 2.5. *With the above notation, for every positive integer k which has dyadic representation $k := \sum_{i=1+L_0}^{n} \alpha_i 2^i + 2^{L_0} - 1$ with $n \ge 1$, $\alpha_n = 1$, $0 \le L_0 \le n-1$ and $\alpha_i \in \{0, 1\}$, $i = 1 + L_0, \ldots, n$, we have*

$$n_k = 4k - 2 \sum_{i=1+L_0}^{n} \alpha_i - L_0 - 1.$$

On the other hand, for every positive integer k with $L_0 = n + 1$, we have $n_k = 4k - L_0 - 1$ again.

https://doi.org/10.1017/etds.2019.103 Published online by Cambridge University Press

Proof. First we deal with the case $L_0 = n+1$ (i.e. $k = 2^{n+1}-1$). Lemma 2.2 ensures that $n_{2^{n+1}-1} = 4.2^{n+1} - (n+1) - 5 = 4.(2^{n+1}-1) - (n+1) - 1$ and we have the desired conclusion. If $L_0 \neq n+1$, we necessarily have $0 \leq L_0 \leq n-1$. Let us consider the case $L_0 = 0$: we apply Lemma 2.4 to write

$$n_k = \sum_{i=1}^{n} \alpha_i n_{2^i} + \sum_{i=1}^{n} \alpha_i - 1.$$

Using Lemma 2.2, we deduce

$$\begin{aligned} n_k &= \sum_{i=1}^{n} \alpha_i (4.2^i - 3) + \sum_{i=1}^{n} \alpha_i - 1 \\ &= 4 \sum_{i=1}^{n} \alpha_i 2^i - 2 \sum_{i=1}^{n} \alpha_i - 1 \\ &= 4k - 2 \sum_{i=1}^{n} \alpha_i - 1 = 4k - 2 \sum_{i=1}^{n} \alpha_i - L_0 - 1. \end{aligned}$$

Otherwise, we consider the case $1 \leq L_0 \leq n-1$ and we apply Lemma 2.4 again to write

$$n_k = \sum_{i=1+L_0}^{n} \alpha_i n_{2^i} + \sum_{i=1+L_0}^{n} \alpha_i + n_{2^{L_0}-1}.$$

We conclude by using Lemma 2.2:

$$\begin{aligned} n_k &= \sum_{i=1+L_0}^{n} \alpha_i (4.2^i - 3) + \sum_{i=1+L_0}^{n} \alpha_i + 4.2^{L_0} - L_0 - 5 \\ &= 4\left(\sum_{i=1+L_0}^{n} \alpha_i 2^i + 2^{L_0} - 1 \right) - 1 - 2 \sum_{i=1+L_0}^{n} \alpha_i - L_0 \\ &= 4k - 2 \sum_{i=1+L_0}^{n} \alpha_i - L_0 - 1. \end{aligned}$$

□

PROPOSITION 2.6. *The sequence* (n_k) *satisfies the following optimal estimate: for every integer* $k \geq 2$,

$$4k - 2\lfloor \log_2(k) \rfloor - 1 \leq n_k \leq 4k - 3$$

Proof. We begin with the case $k = 2^n - 1$, $n \geq 2$. We have $n_k = 4.2^n - n - 5 = 4k - 2 - \lfloor \log_2(k) \rfloor \geq 4k - 2\lfloor \log_2(k) \rfloor - 1$. Now let us consider $k \geq 2$, with $k \neq 2^n - 1$ (i.e. $\log_2(k+1) \notin \mathbb{N}$). The integer k necessarily has the representation $k = 2^n + \sum_{i=1+L_0}^{n} \alpha_i 2^i + 2^{L_0} - 1$ with $n \geq 1$, and $\alpha_i \in \{0, 1\}$, $i = 1 + L_0, \ldots, n-1$. Lemma 2.5 gives $n_k = 4k - 2\sum_{i=1+L_0}^{n} \alpha_i - L_0 - 1$. Clearly we have $n_k \leq 4k - 3$ and $n_{2^n} = 4.2^n - 3$ by Lemma 2.2. On the other hand, we have

$$n_k \geq 4k - 2(n - L_0) - L_0 - 1 \geq 4k - 2n - 1 = 4k - 2\lfloor \log_2(k) \rfloor - 1.$$

Finally, for $k = 2^{m+1} - 2$, we have $k = \sum_{i=1}^{m} 2^i$, $L_0 = 0$, $\lfloor \log_2(2^{m+1} - 2) \rfloor = m$ and Lemma 2.2 gives $n_{2^{m+1}-2} = 4(2^{m+1} - 2) - 2m - 1 = 4(2^{m+1} - 2) - 2\lfloor \log_2(2^{m+1} - 2) \rfloor - 1 = 4k - 2\lfloor \log_2(k) \rfloor - 1$. □

https://doi.org/10.1017/etds.2019.103 Published online by Cambridge University Press

PROPOSITION 2.7. *The sequence (n_k) defined above satisfies $\underline{d}_{B_1}((n_k)_k) > 0$.*

Proof. Using Lemma 2.1, the equality

$$\underline{d}_{B_1}((n_k)_k) = \liminf_{k\to+\infty}\left(\frac{\sum_{j=2}^{k} e^{n_j/\log(n_j)}}{\sum_{j=2}^{n_k} e^{j/\log(j)}}\right)$$

holds. According to Proposition 2.6, there exists a constant $C > 0$ such that, for N large enough and $k \geq N$,

$$\frac{\sum_{j=N}^{k} e^{n_j/\log(n_j)}}{\log(n_k)e^{n_k/\log(n_k)}} \geq \frac{\sum_{j=N}^{k} e^{(4j-C\log(j))/\log(4j-C\log(j))}}{\log(4k)e^{4k/\log(4k)}}.$$

With summation by parts, we get

$$\sum_{j=N}^{k} e^{(4j-C\log(j))/\log(4j-C\log(j))} \sim \frac{\log(k)}{4} e^{(4k-C\log(k))/\log(4k-C\log(k))} \quad \text{as } k \to +\infty.$$

Finally, a simple computation leads to

$$\frac{\log(k)}{4}\frac{e^{(4k-C\log(k))/\log(4k-C\log(k))}}{\log(4k)e^{4k/\log(4k)}} \to \frac{e^{-C}}{4} \quad \text{as } k \to +\infty.$$

This allows us to conclude $\underline{d}_{B_1}((n_k)_k) \geq e^{-C}/4 > 0$. □

Hence we deduce the following result, which improves [**8**, Theorem 4.12].

PROPOSITION 2.8. *Let X be a Fréchet space, Y a separable Fréchet space and $T_n : X \to Y$, $n \in \mathbb{N}$, continuous linear mappings. If the sequence (T_n) satisfies the frequent universality criterion, then (T_n) is B_1-frequently universal.*

3. *Construction of new sequences to reach the extreme densities*

We are going to construct a new sequence with positive A-density for an admissible matrix A defining a sharper density than the natural density and the B_1-density. This construction is inspired by [**8**, §4]. Let us consider an increasing sequence (a_n) of positive integers with $a_1 = 1$. Then we define the function $f : \mathbb{N} \to \mathbb{N}$ by $f(j) = m$ for all $j \in \{a_m, \ldots, a_{m+1} - 1\}$. We also define the sequence $(n_k(f))$ by induction as in (2.1):

$$n_1(f) = 2 \quad \text{and} \quad n_k(f) = 2\sum_{i=1}^{k-1} f(\delta_i) + f(\delta_k) \quad \text{for } k \geq 2. \tag{3.1}$$

Clearly we obtain the following equality, for all $k \geq 3$

$$n_k(f) = n_{k-1}(f) + f(\delta_{k-1}) + f(\delta_k).$$

Observe that, for all $k \neq l$,

$$|n_k(f) - n_l(f)| \geq f(\delta_k) + f(\delta_l). \tag{3.2}$$

In the previous section, the sequence (n_k) satisfied $|n_k - n_l| \geq \delta_k + \delta_l$ in order to prove the frequent universality criterion. In a previous work [**8**], we proved that this condition

https://doi.org/10.1017/etds.2019.103 Published online by Cambridge University Press

could be relaxed as in (3.2), provided that f increases and tends to infinity. Let us observe that, if we set $a_m = m$ for every $m \geq 1$, then the corresponding sequence $(n_k(f))$ is the sequence (n_k) of §2. Throughout this section, we will omit the argument f in $(n_k(f))$ for the sake of readability. Thus our purpose is to compute an exact formula for the new sequence (n_k) to understand its asymptotic behavior and to obtain sharper estimates for the densities given by the frequent universality criterion.

First of all, we will obtain an expression for $n_{2^{a_m+q}}$, with $0 \leq q < a_{m+1} - a_m$.

LEMMA 3.1. *For every $m \in \mathbb{N}$ and every $0 \leq q < a_{m+1} - a_m$, we have*

$$n_{2^{a_m+q}} = -1 - 2m + 2f(1 + a_m + q) + 2^{a_m+q+1} \sum_{i=1}^{m} \frac{1}{2^{a_i-1}}.$$

Proof. Let us define $\Delta_j^{(m,q)} := \{1 \leq l \leq 2^{a_m+q} - 1 : \delta_l = j\}$ for $j \geq 1$. It is clear that, for every $1 \leq l \leq 2^{a_m+q} - 2$, the first zero in the dyadic decomposition of l appears in position less than $a_m + q$, that is, $\delta_l \leq a_m + q$ and $\delta_{2^{a_m+q}-1} = a_m + q + 1$. Thus, for every $j > a_m + q + 1$ we have $\#\Delta_j^{(m,q)} = 0$ and $\#\Delta_{a_m+q+1}^{(m,q)} = 1$. Let us now compute $\#\Delta_j^{(m,q)}$ for $1 \leq j \leq a_m + q$. First, let us observe that we have $j = \delta_l = 1$ if and only if l is even. From this we deduce that $\#\Delta_1^{(m,q)} = 2^{a_m+q-1} - 1$. On the other hand, if $1 \leq l \leq 2^{a_m+q} - 2$ is such that $j = \delta_l \geq 2$ then its dyadic decomposition is given by $j - 1$ ones followed by a zero, and then we have 2^{a_m+q-j} choices as follows:

$$(\overbrace{\underbrace{1, 1, \ldots, 1, 0}_{\text{length } j}, \star, \star, \ldots, \star, \star}^{\text{length } a_m+q}, 0, 0, \ldots).$$

Thus we obtain that, for every $2 \leq j \leq a_m + q$, $\#\Delta_j^{(m,q)} = 2^{a_m+q-j}$. We use these facts to compute $n_{2^{a_m+q}}$, for q satisfying $a_m \leq a_m + q < a_{m+1}$:

$$\begin{aligned}
n_{2^{a_m+q}} &= 2 \sum_{j=1}^{2^{a_m+q}-1} f(\delta_j) + f(\delta_{2^{a_m+q}}) \\
&= 2 \sum_{j=1}^{a_m+q+1} f(j) \#\Delta_j^{(m,q)} + f(\delta_{2^{a_m+q}}) \\
&= 2\left((2^{a_m+q-1} - 1) f(1) + \sum_{j=2}^{a_m+q} f(j) 2^{a_m+q-j} + f(1 + a_m + q) \right) + 1, \\
&\qquad \text{using } f(\delta_{2^{a_m+q}}) = f(1) = 1 \\
&= 2\left(-1 + \sum_{i=1}^{m-1} \sum_{j=a_i}^{a_{i+1}-1} f(j) 2^{a_m+q-j} + \sum_{j=a_m}^{a_m+q} f(j) 2^{a_m+q-j} + f(1 + a_m + q) \right) + 1.
\end{aligned}$$

Setting $u_m = \sum_{i=1}^{m-1} \sum_{j=a_i}^{a_{i+1}-1} f(j) 2^{a_m+q-j} + \sum_{j=a_m}^{a_m+q} f(j) 2^{a_m+q-j}$, we have

$$u_m = \sum_{i=1}^{m-1} i 2^{a_m+q} \sum_{j=a_i}^{a_{i+1}-1} 2^{-j} + m \sum_{j=a_m}^{a_m+q} 2^{a_m+q-j}$$

https://doi.org/10.1017/etds.2019.103 Published online by Cambridge University Press

$$= 2^{a_m+q}\left(\sum_{i=1}^{m-1}\left(\frac{i}{2^{a_i-1}} - \frac{i}{2^{a_{i+1}-1}}\right) + m\left(\frac{1}{2^{a_m-1}} - \frac{1}{2^{a_m+q}}\right)\right)$$

$$= 2^{a_m+q}\left(\sum_{i=1}^{m}\frac{1}{2^{a_i-1}} - \frac{m}{2^{a_m+q}}\right).$$

Therefore we get

$$n_{2^{a_m+q}} = -1 - 2m + 2f(1 + a_m + q) + 2^{a_m+q+1}\sum_{i=1}^{m}\frac{1}{2^{a_i-1}}. \qquad \square$$

From this lemma we get the following result.

LEMMA 3.2. *For every $m \in \mathbb{N}$ and every $0 \le q < a_{m+1} - a_m$, we have*

$$n_{2^{a_m+q}} = \begin{cases} -1 + 2^{a_m+q+1}\displaystyle\sum_{i=1}^{m}\frac{1}{2^{a_i-1}} & \text{if } 0 \le q < a_{m+1} - a_m - 1, \\ 1 + 2^{a_{m+1}}\displaystyle\sum_{i=1}^{m}\frac{1}{2^{a_i-1}} & \text{if } q = a_{m+1} - a_m - 1, \end{cases}$$

Proof. It suffices to apply Lemma 3.1, taking into account that we have $f(a_m + q + 1) = m$ if $0 \le q \le a_{m+1} - a_m - 1$ and $f(a_m + q + 1) = m + 1$ if $q = a_{m+1} - a_m - 1$. $\square$

We will later need an expression for the integers n_{2^L-1} for $L \ge 1$. This is the goal of the following lemma.

LEMMA 3.3. *Let $L \ge 2$ with $a_l - 1 \le L < a_{l+1} - 1$. Then we have*

$$n_{2^L-1} = \begin{cases} 2^{a_l}\displaystyle\sum_{i=1}^{l-1}\frac{1}{2^{a_i-1}} - l & \text{for } L = a_l - 1, \\ 2^{1+L}\displaystyle\sum_{i=1}^{l}\frac{1}{2^{a_i-1}} - (l+2) & \text{for } L > a_l - 1. \end{cases}$$

Proof. By definition the following equality holds:

$$n_{2^L} = n_{2^L-1} + f(\delta_{2^L-1}) + f(\delta_{2^L}) = n_{2^L-1} + f(1 + L) + 1. \tag{3.3}$$

First, let us consider the case $L = a_l - 1 = a_{l-1} + (a_l - a_{l-1} - 1)$. Lemma 3.2 gives $n_{2^L} = n_{2^{a_l-1}} = 1 + 2^{a_l}\sum_{i=1}^{l-1}(1/2^{a_i-1})$. Moreover, since we have $f(1 + L) = f(a_l) = l$, then formula (3.3) implies

$$n_{2^{a_l-1}-1} = n_{2^{a_l-1}} - l - 1 = 2^{a_l}\sum_{i=1}^{l-1}\frac{1}{2^{a_i-1}} - l.$$

On the other hand, if we have $a_l - 1 < L < a_{l+1} - 1$, Lemma 3.2 yields $n_{2^L} = -1 + 2^{L+1}\sum_{i=1}^{l}(1/2^{a_i-1})$. Thus, formula (3.3) gives

$$n_{2^L-1} = 2^{1+L}\sum_{i=1}^{l}\frac{1}{2^{a_i-1}} - (l+2). \qquad \square$$

https://doi.org/10.1017/etds.2019.103 Published online by Cambridge University Press

LEMMA 3.4. *Let $N = 2^n + \sum_{i=0}^{n-1} \alpha_i 2^i$ with $n \geq 1$ and $\sum_{i=0}^{n-1} \alpha_i 2^i \neq 0$. Then*

$$n_{N-2^n} = \begin{cases} 2 \displaystyle\sum_{i=2^n+1}^{N-1} f(\delta_i) + f(\delta_N) & \text{if } N < 2^{n+1} - 1, \\ 2 \displaystyle\sum_{i=2^n+1}^{N-1} f(\delta_i) + 2f(\delta_N) - f(\delta_{N-2^n}) & \text{if } N = 2^{n+1} - 1. \end{cases}$$

Proof. We easily adapt the proof of Lemma 2.3. □

Using Lemma 3.4 instead of Lemma 2.3, we may also prove that Lemma 2.4 remains valid in this context. We shall use it in what follows.

Definition 3.5. In what follows, we express every positive integer N in the following manner, taking into account the properties of the sequence (a_l):

$$N = 2^{L_0} - 1 + \sum_{q=q_0}^{a_{l_0}-a_{l_0-1}-1} \alpha_{a_{l_0-1}+q} 2^{a_{l_0-1}+q} + \sum_{j=l_0}^{l_N-1} \sum_{q=0}^{a_{j+1}-a_j-1} \alpha_{a_j+q} 2^{a_j+q} + \sum_{q=0}^{q_N} \alpha_{a_{l_N}+q} 2^{a_{l_N}+q}$$

with $a_{l_0-1} \leq 1 + L_0 < a_{l_0}$, $a_{l_N} \leq q_N + a_{l_N} < a_{l_N+1}$, $q_0 + a_{l_0-1} = 1 + L_0$ and $\alpha_{a_{l_N}+q_N} = 1$. We also set $w_N = q_N + a_{l_N}$.

LEMMA 3.6. *For every positive integer N, we have, using the notation of Definition 3.5,*

$$n_N = n_{2^{L_0}-1} + 2\left(\sum_{j=l_0}^{1+l_N} \alpha_{a_j-1}\right) + \sum_{q=q_0}^{a_{l_0}-1-a_{l_0-1}} \alpha_{a_{l_0-1}+q} 2^{a_{l_0-1}+q+1} \left(\sum_{i=1}^{l_0-1} \frac{1}{2^{a_i}-1}\right) + \sum_{j=l_0}^{l_N-1} \sum_{q=0}^{a_{j+1}-a_j-1} \alpha_{a_j+q} 2^{a_j+q+1} \left(\sum_{i=1}^{j} \frac{1}{2^{a_i}-1}\right) + \sum_{q=0}^{q_N} \alpha_{a_{l_N}+q} 2^{a_{l_N}+q+1} \left(\sum_{i=1}^{l_N} \frac{1}{2^{a_i}-1}\right).$$

Proof. Using the notation of Definition 3.5, Lemma 2.4 gives the following equality (with the convention $n_0 = -1$). Since $N = \sum_{i=L_0+1}^{n} \alpha_i 2^i + (2^{L_0} - 1) = \sum_{i=0}^{n} \alpha_i 2^i$,

$$n_N = n_{2^{L_0}-1} + \sum_{q=q_0}^{a_{l_0}-1-a_{l_0-1}} \alpha_{a_{l_0-1}+q} n_{2^{a_{l_0-1}+q}} + \sum_{j=l_0}^{l_N-1} \sum_{q=0}^{a_{j+1}-a_j-1} \alpha_{a_j+q} n_{2^{a_j+q}} + \sum_{q=0}^{q_N} \alpha_{a_{l_N}+q} n_{2^{a_{l_N}+q}} + \sum_{i=1+L_0}^{w_N} \alpha_i. \tag{3.4}$$

We begin by transforming the first sum above. We apply Lemma 3.2 and we get

https://doi.org/10.1017/etds.2019.103 Published online by Cambridge University Press

$$\begin{aligned}
&\sum_{q=q_0}^{a_{l_0}-1-a_{l_0-1}} \alpha_{a_{l_0-1}+q} n_{2^{a_{l_0-1}+q}} \\
&\quad = \sum_{q=q_0}^{a_{l_0}-2-a_{l_0-1}} \alpha_{a_{l_0-1}+q}\left(-1+2^{a_{l_0-1}+q+1}\left(\sum_{i=1}^{l_0-1}\frac{1}{2^{a_i-1}}\right)\right) \\
&\qquad + \alpha_{a_{l_0}-1}\left(1+2^{a_{l_0}}\sum_{i=1}^{l_0-1}\frac{1}{2^{a_i-1}}\right) \\
&\quad = \sum_{q=q_0}^{a_{l_0}-1-a_{l_0-1}} \alpha_{a_{l_0-1}+q} 2^{a_{l_0-1}+q+1}\left(\sum_{i=1}^{l_0-1}\frac{1}{2^{a_i-1}}\right) - \sum_{q=q_0}^{a_{l_0}-2-a_{l_0-1}} \alpha_{a_{l_0-1}+q} + \alpha_{a_{l_0}-1}.
\end{aligned}$$

We proceed in the same way with the second sum of (3.4) and we obtain

$$\begin{aligned}
&\sum_{j=l_0}^{l_N-1}\sum_{q=0}^{a_{j+1}-a_j-1} \alpha_{a_j+q} n_{2^{a_j+q}} \\
&\quad = \sum_{j=l_0}^{l_N-1}\sum_{q=0}^{a_{j+1}-a_j-2} \alpha_{a_j+q}\left(-1+2^{a_j+q+1}\left(\sum_{i=1}^{j}\frac{1}{2^{a_i-1}}\right)\right) \\
&\qquad + \sum_{j=l_0}^{l_N-1} \alpha_{a_{j+1}-1}\left(1+2^{a_{j+1}}\left(\sum_{i=1}^{j}\frac{1}{2^{a_i-1}}\right)\right) \\
&\quad = \sum_{j=l_0}^{l_N-1}\sum_{q=0}^{a_{j+1}-a_j-1} \alpha_{a_j+q} 2^{a_j+q+1}\left(\sum_{i=1}^{j}\frac{1}{2^{a_i-1}}\right) \\
&\qquad - \sum_{j=l_0}^{l_N-1}\sum_{q=0}^{a_{j+1}-a_j-2} \alpha_{a_j+q} + \sum_{j=l_0}^{l_N-1} \alpha_{a_{j+1}-1}.
\end{aligned}$$

Finally, to study the third sum of (3.4), we consider two cases. For $w_N < a_{l_N+1} - 1$,

$$\begin{aligned}
\sum_{q=0}^{q_N} \alpha_{a_{l_N}+q} n_{2^{a_{l_N}+q}} &= \sum_{q=0}^{q_N} \alpha_{a_{l_N}+q}\left(-1+2^{a_{l_N}+q+1}\left(\sum_{i=1}^{l_N}\frac{1}{2^{a_i-1}}\right)\right) \\
&= \sum_{q=0}^{q_N} \alpha_{a_{l_N}+q} 2^{a_{l_N}+q+1}\left(\sum_{i=1}^{l_N}\frac{1}{2^{a_i-1}}\right) - \sum_{q=0}^{q_N} \alpha_{a_{l_N}+q}.
\end{aligned}$$

For $w_N = a_{l_N+1} - 1$,

$$\begin{aligned}
&\sum_{q=0}^{q_N} \alpha_{a_{l_N}+q} n_{2^{a_{l_N}+q}} \\
&\quad = \sum_{q=0}^{a_{l_N+1}-2-a_{l_N}} \alpha_{a_{l_N}+q}\left(-1+2^{a_{l_N}+q+1}\left(\sum_{i=1}^{l_N}\frac{1}{2^{a_i-1}}\right)\right)
\end{aligned}$$

https://doi.org/10.1017/etds.2019.103 Published online by Cambridge University Press

$$+\alpha_{a_{l_N+1}-1}\left(1+2^{a_{l_N+1}}\left(\sum_{i=1}^{l_N}\frac{1}{2^{a_i-1}}\right)\right)$$
$$=\sum_{q=0}^{a_{l_N+1}-1-a_{l_N}}\alpha_{a_{l_N}+q}2^{a_{l_N}+q+1}\left(\sum_{i=1}^{l_N}\frac{1}{2^{a_i-1}}\right)-\sum_{q=0}^{a_{l_N+1}-2-a_{l_N}}\alpha_{a_{l_N}+q}+\alpha_{a_{l_N+1}-1}.$$

We now gather these results together in the case where $w_N < a_{l_N+1}-1$ (the case $w_N = a_{l_N+1}-1$ being similar). We get

$$\begin{aligned}
n_N &= n_{2^{L_0}-1}+\sum_{q=q_0}^{a_{l_0}-1-a_{l_0-1}}\alpha_{a_{l_0-1}+q}n_{2^{a_{l_0-1}+q}}+\sum_{j=l_0}^{l_N-1}\sum_{q=0}^{a_{j+1}-a_j-1}\alpha_{a_j+q}n_{2^{a_j+q}}\\
&\quad+\sum_{q=0}^{q_N}\alpha_{a_{l_N}+q}n_{2^{a_{l_N}+q}}+\sum_{i=1+L_0}^{w_N}\alpha_i\\
&=n_{2^{L_0}-1}+\sum_{q=q_0}^{a_{l_0}-1-a_{l_0-1}}\alpha_{a_{l_0-1}+q}2^{a_{l_0-1}+q+1}\left(\sum_{i=1}^{l_0-1}\frac{1}{2^{a_i-1}}\right)\\
&\quad+\sum_{j=l_0}^{l_N-1}\sum_{q=0}^{a_{j+1}-a_j-1}\alpha_{a_j+q}2^{a_j+q+1}\left(\sum_{i=1}^{j}\frac{1}{2^{a_i-1}}\right)+\sum_{q=0}^{q_N}\alpha_{a_{l_N}+q}2^{a_{l_N}+q+1}\left(\sum_{i=1}^{l_N}\frac{1}{2^{a_i-1}}\right)\\
&\quad-\sum_{q=q_0}^{a_{l_0}-2-a_{l_0-1}}\alpha_{a_{l_0-1}+q}-\sum_{j=l_0}^{l_N-1}\sum_{q=0}^{a_{j+1}-a_j-2}\alpha_{a_j+q}\\
&\quad-\sum_{q=0}^{q_N}\alpha_{a_{l_N}+q}+\alpha_{a_{l_0}-1}+\sum_{j=l_0}^{l_N-1}\alpha_{a_{j+1}-1}+\sum_{i=1+L_0}^{w_N}\alpha_i.
\end{aligned}$$

Moreover, observe that, by definition, if $w_N < a_{l_N+1}-1$,

$$\begin{aligned}
\sum_{i=1+L_0}^{w_N}\alpha_i &= \sum_{q=q_0}^{a_{l_0}-1-a_{l_0-1}}\alpha_{a_{l_0-1}+q}+\sum_{j=l_0}^{l_N-1}\sum_{q=0}^{a_{j+1}-a_j-1}\alpha_{a_j+q}+\sum_{q=0}^{q_N}\alpha_{a_{l_N}+q}\\
&=\sum_{q=q_0}^{a_{l_0}-2-a_{l_0-1}}\alpha_{a_{l_0-1}+q}+\sum_{j=l_0}^{l_N-1}\sum_{q=0}^{a_{j+1}-a_j-2}\alpha_{a_j+q}\\
&\quad+\sum_{q=0}^{q_N}\alpha_{a_{l_N}+q}+\alpha_{a_{l_0}-1}+\sum_{j=l_0}^{l_N-1}\alpha_{a_{j+1}-1}.
\end{aligned}$$

Let us observe also that if $w_N < a_{l_N+1}-1$, then we have $\alpha_{a_{1+l_N}-1}=0$. Thus, we derive the result:

$$\begin{aligned}
n_N &= n_{2^{L_0}-1}+\sum_{q=q_0}^{a_{l_0}-1-a_{l_0-1}}\alpha_{a_{l_0-1}+q}2^{a_{l_0-1}+q+1}\left(\sum_{i=1}^{l_0-1}\frac{1}{2^{a_i-1}}\right)\\
&\quad+\sum_{j=l_0}^{l_N-1}\sum_{q=0}^{a_{j+1}-a_j-1}\alpha_{a_j+q}2^{a_j+q+1}\left(\sum_{i=1}^{j}\frac{1}{2^{a_i-1}}\right)
\end{aligned}$$

https://doi.org/10.1017/etds.2019.103 Published online by Cambridge University Press

$$
\begin{aligned}
&+\sum_{q=0}^{q_N}\alpha_{a_{l_N}+q}2^{a_{l_N}+q+1}\Big(\sum_{i=1}^{l_N}\frac{1}{2^{a_i-1}}\Big)+2\alpha_{a_{l_0}-1}+2\sum_{j=l_0}^{l_N-1}\alpha_{a_{j+1}-1}\\
&=n_{2^{L_0}-1}+\sum_{q=q_0}^{a_{l_0}-1-a_{l_0-1}}\alpha_{a_{l_0-1}+q}2^{a_{l_0-1}+q+1}\Big(\sum_{i=1}^{l_0-1}\frac{1}{2^{a_i-1}}\Big)\\
&+\sum_{j=l_0}^{l_N-1}\sum_{q=0}^{a_{j+1}-a_j-1}\alpha_{a_j+q}2^{a_j+q+1}\Big(\sum_{i=1}^{j}\frac{1}{2^{a_i-1}}\Big)\\
&+\sum_{q=0}^{q_N}\alpha_{a_{l_N}+q}2^{a_{l_N}+q+1}\Big(\sum_{i=1}^{l_N}\frac{1}{2^{a_i-1}}\Big)+2\sum_{j=l_0}^{1+l_N}\alpha_{a_j-1}. \qquad \square
\end{aligned}
$$

LEMMA 3.7. *Using the notation of Definition 3.5, we have, for every positive integer N,*

$$
\begin{aligned}
n_N=\;&2N\Big(\sum_{i=1}^{\infty}\frac{1}{2^{a_i-1}}\Big)+2\Big(\sum_{i=1}^{\infty}\frac{1}{2^{a_i-1}}\Big)+2\Big(\sum_{j=l_0}^{1+l_N}\alpha_{a_j-1}\Big)-2^{1+L_0}\Big(\sum_{i=l_0-\tau_0}^{\infty}\frac{1}{2^{a_i-1}}\Big)\\
&-2\Big(\sum_{i=l_0}^{\infty}\frac{1}{2^{a_i-1}}\Big)\Big(\sum_{q=q_0}^{a_{l_0}-a_{l_0-1}-1}\alpha_{a_{l_0-1}+q}2^{a_{l_0-1}+q}\Big)\\
&-2\sum_{j=l_0}^{l_N-1}\Big(\sum_{i=j+1}^{\infty}\frac{1}{2^{a_i-1}}\Big)\Big(\sum_{q=0}^{a_{j+1}-a_j-1}\alpha_{a_j+q}2^{a_j+q}\Big)\\
&-2\Big(\sum_{i=l_N+1}^{\infty}\frac{1}{2^{a_i-1}}\Big)\Big(\sum_{q=0}^{q_N}\alpha_{l_N+q}2^{a_{l_N}+q}\Big)-l_0-1+2\tau_0,
\end{aligned}
$$

with $\tau_0=0$ *if* $L_0>a_{l_0-1}-1$ *and* $\tau_0=1$ *if* $L_0=a_{l_0-1}-1$.

Proof. Let $N=\sum_{i=L_0+1}^{n}\alpha_i2^i+(2^{L_0}-1)=\sum_{i=0}^{n}\alpha_i2^i$ with $a_{l_0-1}-1\le L_0<a_{l_0}-1$. Lemma 3.6 gives

$$
\begin{aligned}
n_N=\;&n_{2^{L_0}-1}+2\Big(\sum_{j=l_0}^{1+l_N}\alpha_{a_j-1}\Big)+\sum_{q=q_0}^{a_{l_0}-1-a_{l_0-1}}\alpha_{a_{l_0-1}+q}2^{a_{l_0-1}+q+1}\Big(\sum_{i=1}^{l_0-1}\frac{1}{2^{a_i-1}}\Big)\\
&+\sum_{j=l_0}^{l_N-1}\sum_{q=0}^{a_{j+1}-a_j-1}\alpha_{a_j+q}2^{a_j+q+1}\Big(\sum_{i=1}^{j}\frac{1}{2^{a_i-1}}\Big)+\sum_{q=0}^{q_N}\alpha_{a_{l_N}+q}2^{a_{l_N}+q+1}\Big(\sum_{i=1}^{l_N}\frac{1}{2^{a_i-1}}\Big).
\end{aligned}
$$

Let us now express every sum of the form $\sum_{i=1}^{K}(1/2^{a_i-1})$ as $\sum_{i=1}^{\infty}(1/2^{a_i-1})-\sum_{i=K+1}^{\infty}(1/2^{a_i-1})$. This gives

$$
\begin{aligned}
n_N=\;&n_{2^{L_0}-1}+2\Big(\sum_{j=l_0}^{1+l_N}\alpha_{a_j-1}\Big)+2\Big(\sum_{i=1}^{\infty}\frac{1}{2^{a_i-1}}\Big)\Big(\sum_{q=q_0}^{a_{l_0}-a_{l_0-1}-1}\alpha_{a_{l_0-1}+q}2^{a_{l_0-1}+q}\Big)\\
&+2\Big(\sum_{i=1}^{\infty}\frac{1}{2^{a_i-1}}\Big)\Big(\sum_{j=l_0}^{l_N-1}\sum_{q=0}^{a_{j+1}-a_j-1}\alpha_{a_j+q}2^{a_j+q}+\sum_{q=0}^{q_N}\alpha_{a_{l_N}+q}2^{a_{l_N}+q}\Big)
\end{aligned}
$$

https://doi.org/10.1017/etds.2019.103 Published online by Cambridge University Press

$$
\begin{aligned}
&- 2\Bigg(\sum_{i=l_0}^{\infty}\frac{1}{2^{a_i-1}}\Bigg)\Bigg(\sum_{q=q_0}^{a_{l_0}-a_{l_0-1}-1}\alpha_{a_{l_0-1}+q}2^{a_{l_0-1}+q}\Bigg)\\
&- 2\Bigg(\sum_{j=l_0}^{l_N-1}\Bigg(\sum_{i=j+1}^{\infty}\frac{1}{2^{a_i-1}}\Bigg)\sum_{q=0}^{a_{j+1}-a_j-1}\alpha_{a_j+q}2^{a_j+q}\Bigg)\\
&- 2\Bigg(\sum_{i=l_N+1}^{\infty}\frac{1}{2^{a_i-1}}\Bigg)\Bigg(\sum_{q=0}^{q_N}\alpha_{a_{l_N}+q}2^{a_{l_N}+q}\Bigg).
\end{aligned}
$$

Moreover, Lemma 3.3 allows us to express

$$
\begin{aligned}
n_{2^{L_0}-1} &= \begin{cases} 2^{1+L_0}\displaystyle\sum_{i=1}^{l_0-2}\frac{1}{2^{a_i-1}} - (l_0-1) & \text{if } L_0 = a_{l_0-1}-1,\\ 2^{1+L_0}\displaystyle\sum_{i=1}^{l_0-1}\frac{1}{2^{a_i-1}} - (l_0+1) & \text{if } L_0 > a_{l_0-1}-1, \end{cases}\\
&= \begin{cases} 2^{1+L_0}\Big(\displaystyle\sum_{i=1}^{\infty}\frac{1}{2^{a_i-1}}\Big) - 2^{1+L_0}\Big(\displaystyle\sum_{i=l_0-1}^{\infty}\frac{1}{2^{a_i-1}}\Big) - (l_0-1) & \text{if } L_0 = a_{l_0-1}-1,\\ 2^{1+L_0}\Big(\displaystyle\sum_{i=1}^{\infty}\frac{1}{2^{a_i-1}}\Big) - 2^{1+L_0}\Big(\displaystyle\sum_{i=l_0}^{\infty}\frac{1}{2^{a_i-1}}\Big) - (l_0+1) & \text{if } L_0 > a_{l_0-1}-1. \end{cases}
\end{aligned}
$$

From now on, we treat only the case where $L_0 = a_{l_0-1}-1$, the other case being completely similar. Let us replace in the first expression the value of $n_{2^{L_0}-1}$. We obtain

$$
\begin{aligned}
n_N &= -l_0 + 1 - 2^{1+L_0}\Bigg(\sum_{i=l_0-1}^{\infty}\frac{1}{2^{a_i-1}}\Bigg)\\
&\quad + 2\Bigg(\sum_{j=l_0}^{1+l_N}\alpha_{a_j-1}\Bigg) + 2\Bigg(\sum_{i=1}^{\infty}\frac{1}{2^{a_i-1}}\Bigg)\Bigg(2^{L_0} + \sum_{q=q_0}^{a_{l_0}-a_{l_0-1}-1}\alpha_{a_{l_0-1}+q}2^{a_{l_0-1}+q}\Bigg)\\
&\quad + 2\Bigg(\sum_{i=1}^{\infty}\frac{1}{2^{a_i-1}}\Bigg)\Bigg(\sum_{j=l_0}^{l_N-1}\sum_{q=0}^{a_{j+1}-a_j-1}\alpha_{a_j+q}2^{a_j+q} + \sum_{q=0}^{q_N}\alpha_{a_{l_N}+q}2^{a_{l_N}+q}\Bigg)\\
&\quad - 2\Bigg(\sum_{i=l_0}^{\infty}\frac{1}{2^{a_i-1}}\Bigg)\Bigg(\sum_{q=q_0}^{a_{l_0}-a_{l_0-1}-1}\alpha_{a_{l_0-1}+q}2^{a_{l_0-1}+q}\Bigg)\\
&\quad - 2\Bigg(\sum_{j=l_0}^{l_N-1}\Bigg(\sum_{i=j+1}^{\infty}\frac{1}{2^{a_i-1}}\Bigg)\sum_{q=0}^{a_{j+1}-a_j-1}\alpha_{a_j+q}2^{a_j+q}\Bigg)\\
&\quad - 2\Bigg(\sum_{i=l_N+1}^{\infty}\frac{1}{2^{a_i-1}}\Bigg)\Bigg(\sum_{q=0}^{q_N}\alpha_{a_{l_N}+q}2^{a_{l_N}+q}\Bigg).
\end{aligned}
$$

Observe now that

https://doi.org/10.1017/etds.2019.103 Published online by Cambridge University Press

$$2^{L_0} + \sum_{q=q_0}^{a_{l_0}-a_{l_0-1}-1} \alpha_{a_{l_0-1}+q} 2^{a_{l_0-1}+q} + \sum_{j=l_0}^{l_N-1} \sum_{q=0}^{a_{j+1}-a_j-1} \alpha_{a_j+q} 2^{a_j+q} + \sum_{q=0}^{q_N} \alpha_{a_{l_N}+q} 2^{a_{l_N}+q} = N+1,$$

and replace this in the preceding expression of n_N to obtain

$$\begin{aligned} n_N = {} & 2\Big(\sum_{i=1}^{\infty} \frac{1}{2^{a_i-1}}\Big)N + 2\Big(\sum_{i=1}^{\infty} \frac{1}{2^{a_i-1}}\Big) + 2\Big(\sum_{j=l_0}^{1+l_N} \alpha_{a_j-1}\Big) - 2^{1+L_0}\Big(\sum_{i=l_0-1}^{\infty} \frac{1}{2^{a_i-1}}\Big) \\ & -2\Big(\sum_{i=l_0}^{\infty} \frac{1}{2^{a_i-1}}\Big)\Big(\sum_{q=q_0}^{a_{l_0}-a_{l_0-1}-1} \alpha_{a_{l_0-1}+q} 2^{a_{l_0-1}+q}\Big) \\ & -2\Big(\sum_{j=l_0}^{l_N-1}\Big(\sum_{i=j+1}^{\infty} \frac{1}{2^{a_i-1}}\Big) \sum_{q=0}^{a_{j+1}-a_j-1} \alpha_{a_j+q} 2^{a_j+q}\Big) \\ & -2\Big(\sum_{i=l_N+1}^{\infty} \frac{1}{2^{a_i-1}}\Big)\Big(\sum_{q=0}^{q_N} \alpha_{a_{l_N}+q} 2^{a_{l_N}+q}\Big) - l_0 + 1. \end{aligned}$$

□

Now observe that we have the following estimates:

$$2\Big(\sum_{i=l_0}^{\infty} \frac{1}{2^{a_i-1}}\Big)\Big(\sum_{q=q_0}^{a_{l_0}-a_{l_0-1}-1} \alpha_{a_{l_0-1}+q} 2^{a_{l_0-1}+q}\Big) \le 4\Big(\sum_{j=1}^{a_{l_0}-a_{l_0-1}-q_0} 2^{-j}\Big) + \frac{4}{2^{a_{l_0+1}-1}} 2^{a_{l_0}} \le 8,$$

$$\begin{aligned} 2\Big(\sum_{j=l_0}^{l_N-1}\Big(\sum_{i=j+1}^{\infty} \frac{1}{2^{a_i-1}}\Big) \sum_{q=0}^{a_{j+1}-a_j-1} \alpha_{a_j+q} 2^{a_j+q}\Big) &\le 8\Big(\sum_{j=l_0}^{l_N-1} 2^{-(a_{j+1}-a_j)}\Big(\sum_{q=0}^{a_{j+1}-a_j-1} 2^q\Big)\Big) \\ &\quad + 16\Big(\sum_{j=l_0}^{l_N-1} 2^{-(a_{j+2}-a_j)}\Big(\sum_{q=0}^{a_{j+1}-a_j-1} 2^q\Big)\Big) \\ &\le 8(l_N - l_0) + 16, \end{aligned}$$

and

$$2\Big(\sum_{i=l_N+1}^{\infty} \frac{1}{2^{a_i-1}}\Big)\Big(\sum_{q=0}^{q_N} \alpha_{a_{l_N}+q} 2^{a_{l_N}+q}\Big) \le 4.$$

The combination of these estimates with Lemma 3.7 leads to the following statement.

LEMMA 3.8. *Using the notation of Definition 3.5, there exist positive real numbers C_1, C_2, C_3, C_4 such that we have, for every positive integer N,*

$$2N\Big(\sum_{i=1}^{\infty} \frac{1}{2^{a_i-1}}\Big) - C_1 l_N - C_2 \le n_N \le 2N\Big(\sum_{i=1}^{\infty} \frac{1}{2^{a_i-1}}\Big) + C_3 l_N + C_4.$$

https://doi.org/10.1017/etds.2019.103 Published online by Cambridge University Press

Thus assume that $a_m = 2^{2^{\cdot^{\cdot^{\cdot^{2^m}}}}}$, where 2 appears $s-1$ times ($s \geq 2$). Then the associated function $f : \mathbb{N} \to \mathbb{N}$ is given by $f(j) = m$, for $j \in \{a_m, \ldots, a_{m+1} - 1\}$, and the estimate

$$CN - C_1 \log^{(s)}(N) \leq n_N \leq CN + C_2 \log^{(s)}(N)$$

holds with $C, C_1, C_2 > 0$. An easy adaptation of [**8**, Lemma 4.10] gives $\underline{d}_{\widetilde{D}_s}((n_k)) > 0$, and we have proved the following result.

THEOREM 3.9. *Let X be a Fréchet space, Y a separable Fréchet space and $T_n : X \to Y$, $n \in \mathbb{N}$, continuous linear mappings. If the sequence (T_n) satisfies the frequent universality criterion, then (T_n) is $\widetilde{D}_s$-frequently universal for any $s \geq 1$.*

Remark 3.10.

(1) Let h be any real strictly increasing C^1-function satisfying the following conditions: $h(x) \to +\infty$, as $x \to +\infty$, $x \mapsto x/h(\log(x))$ (eventually) increases (to infinity), $h(x) = o(\log(x))$ and $h'(x) = o(h(x))$ as $x \to +\infty$. With a good choice of the sequence (a_n) (i.e. $a_n = h^{-1}(n)$) one can show that the conditions of the frequent universality criterion automatically imply A_h-frequent universality, where A_h is the admissible matrix given by the coefficients $e^{k/h(\log(k))}$. Indeed, under the previous assumptions, Lemma 3.8 ensures that there exist $C, C_1, C_2 > 0$ such that

$$CN - C_1 h(\log(N)) \leq n_N \leq CN + C_2 h(\log(N)).$$

Taking into account Lemma 2.1 and the estimate $\sum_{k=1}^{n} e^{k/h(\log(k))} \sim h(\log(n)) e^{n/h(\log(n))}$ as $n \to +\infty$, an adaptation of [**8**, Lemma 4.10] gives $\underline{d}_{A_h}((n_k)) > 0$ and we have proved the result.

(2) Sophie Grivaux showed that one can find a proof of Theorem 3.9 (or Remark 3.10(1)) in the particular case of hypercyclicity in an elegant way thanks to ergodic theory arguments [**10**]. To do this, she combines [**11**, Theorem 7] with [**12**, Theorem 1].

We end this section by proving that an operator $T : X \to X$, acting on a Fréchet space X, cannot be A_1-frequently hypercyclic. In other words, (1) from Remark 3.10 ensures that we have obtained the sharpest result in the context of hypercyclicity.

PROPOSITION 3.11. *Let X be a Fréchet space. Then there is no A_1-frequently hypercyclic operator on X.*

Proof. Let $T : X \to X$ be a continuous operator. Assume that the topology on X is given by an increasing sequence of semi-norms (p_j). Let us consider $J \geq 1$, $f \in X$ with $p_J(f) \geq 1$. We set $0 < \varepsilon < 1$ and $B_J(f, \varepsilon) := \{z \in X : p_J(z - f) < \varepsilon\}$. Assume that T is A_1-frequently hypercyclic and let us consider $x \in X$ so that x is an A_1-frequently hypercyclic vector for T. According to the proof of [**8**, Proposition 3.7], the set $N(x, B_J(f, \varepsilon))$ has bounded gaps. Hence we denote by M an upper bound for the length of these gaps. Let us define $K_0 = J$ and $\eta_0 = 1 - \varepsilon > 0$. The continuity of the operator T ensures that there exist a sequence of natural numbers $(K_i)_{1 \leq i \leq M}$ and a sequence of positive numbers $(\eta_i)_{1 \leq i \leq M}$ such that:

(1) for $0 \leq i \leq M$, $K_i \geq J$;

(2) for $0 \leq i \leq M$, $\eta_i \leq 1 - \varepsilon$;

https://doi.org/10.1017/etds.2019.103 Published online by Cambridge University Press

(3) for $1 \le i \le M$ and for any $z \in X$,

$$p_{K_i}(z) \le \eta_i \implies p_{K_{i-1}}(T(z)) \le \eta_{i-1}.$$

Now we use the A_1-frequent hypercyclicity of x to find a natural number n such that

$$p_{K_M}(T^n(x)) < \eta_M.$$

It follows from (3) that, for every $0 \le i \le M$,

$$p_{K_{M-i}}(T^{n+i}(x)) < \eta_{M-i}.$$

Therefore, by (1) and (2),

$$p_{K_J}(T^{n+i}(x)) < 1 - \varepsilon.$$

Thus, for every $0 \le i \le M$,

$$p_J(T^{n+i}(x) - f) \ge |p_J(f) - p_J(T^{n+i}(x))| \ge 1 - (1 - \varepsilon) = \varepsilon.$$

We have proved that $[n; n + M] \cap N(x, B_J(f, \varepsilon)) = \emptyset$, which contradicts the definition of M. □

Remark 3.12. An anonymous referee pointed out to us that Proposition 3.11 could also be obtained as a consequence of [**5**, §5, Proposition 2] once one observes that both the lower A_1-density and the lower Banach density are zero on non syndetic sets.

4. *A log-frequently hypercyclic operator which is not frequently hypercyclic*

In their very nice paper, Bayart and Ruzsa [**4**] gave a characterization of frequently hypercyclic weighted shifts on the sequence spaces ℓ^p and c_0. In particular, a straightforward modification of the proof of [**4**, Theorem 13] gives an analogous result with respect to the so-called logarithmic density.

THEOREM 4.1. *Let $w = (\omega_n)_{n\in\mathbb{N}}$ be a bounded sequence of positive integers. Then B_w is log-frequently hypercyclic on $c_0(\mathbb{N})$ if and only if there exist a sequence $(M(p))$ of positive real numbers tending to $+\infty$ and a sequence (E_p) of subsets of $\mathbb{N}$ such that:*

(a) *for any $p \ge 1$, $\underline{d}_{\log}(E_p) > 0$;*

(b) *for any $p, q \ge 1$, $p \ne q$, $(E_p + [0, p]) \cap (E_q + [0, q]) = \emptyset$;*

(c) $\lim_{n\to\infty,\ n\in E_p+[0,p]} \omega_1 \cdots \omega_n = +\infty$;

(d) *for any $p, q \ge 1$, for any $n \in E_p$ and any $m \in E_q$ with $m > n$, for any $t \in \{0, \ldots, q\}$,*

$$\omega_1 \cdots \omega_{m-n+t} \ge M(p)M(q).$$

In the same paper, the authors also provide examples of a $\mathcal{U}$-frequently hypercyclic weighted shift which is not frequently hypercyclic and of a frequently hypercyclic weighted shift which is not distributionally chaotic. In what follows, we modify these constructions as well as those made in [**8**] to construct a log-frequently hypercyclic operator which is not frequently hypercyclic. We begin with the following lemma.

LEMMA 4.2. *There exist $a > 1$ and $\varepsilon > 0$ such that, for any integer $u > v \ge 1$, if we let $I_u^{a,\varepsilon} = [2^{(1-\varepsilon)a^{2u}}, 2^{(1+\varepsilon)a^{2u}}]$, then the following properties hold:*

https://doi.org/10.1017/etds.2019.103 Published online by Cambridge University Press

(1) $I_u^{a,4\varepsilon} \cap I_v^{a,4\varepsilon} = \emptyset$;
(2) $I_u^{a,2\varepsilon} - I_v^{a,2\varepsilon} \subset I_u^{a,4\varepsilon}$;
(3) $a^2(1-4\varepsilon/1+4\varepsilon) > 1$.

Proof. A simple calculation shows, for any $u > v \geq 1$, that $I_u^{a,4\varepsilon} \cap I_v^{a,4\varepsilon} = \emptyset$ if and only if $a^2(1-4\varepsilon/1+4\varepsilon) \geq 1$. Similarly, we have

$$I_u^{a,\varepsilon} - I_v^{a,\varepsilon} \subset I_u^{a,4\varepsilon} \quad \text{if and only if } 2^{a^{2(u-1)}(a^2(1-2\varepsilon)-(1+2\varepsilon)}(1-2^{2\varepsilon(2-a^2)a^{2(u-1)}}) \geq 1.$$

This can be achieved if $a^2(1-2\varepsilon)-(1+2\varepsilon) \geq 1$ and $a^2 \geq 2+1/2\varepsilon$, for example. Observe that these conditions and (3) are satisfied if ε is chosen to be very small and a very large. □

The philosophy of Lemma 4.2 can be summarized as follows: it suffices to choose a very large and at the same time ε very small to obtain the result stated. From now on, we suppose that a and ε are given by the previous lemma. Let (A_p) be any syndetic partition of $\mathbb{N}$ and M_p be the maximum length of the gaps in A_p. We consider an increasing sequence of integers (b_p) such that

$$[b_p - 8p;\, b_p + 4p] \subset \bigcup_{u\geq 2}]2^{(1-\varepsilon)a^{2(u-1)}};\, 2^{(1-\varepsilon)a^{2u}} - 2^{(1+\varepsilon)a^{2(u-1)}}[\tag{4.1}$$

and

$$b_p \geq (8p+1)2^p. \tag{4.2}$$

This construction is possible if a is chosen big enough. Finally, let

$$E_p = \bigcup_{u\in A_p} (I_u^{a,\varepsilon} \cap b_p\mathbb{N}).$$

LEMMA 4.3. *With the above notation, we have* $\underline{d}_{\log}(E_p) > 0$.

Proof. Let $(n_k)_{k\in\mathbb{N}}$ be an increasing enumeration of A_p. We set $[c_k;\, d_k] = I_{n_k}^{a,\varepsilon}$. Then we get

$$\underline{d}_{\log}(E_p) = \liminf_{k\to\infty}\left(\frac{\sum_{l=1}^{k}\sum_{j=c_l,\, j\in b_p\mathbb{N}}^{d_l} 1/j}{\sum_{j=1}^{c_{k+1}} 1/j}\right) \geq \liminf_{k\to\infty}\left(\frac{\sum_{j=c_k,\, j\in b_p\mathbb{N}}^{d_k} 1/j}{\log(c_{k+1})}\right).$$

Since there exist $0 \leq \alpha_k,\, \beta_k < b_p$ such that $[c_k+\alpha_k;\, d_k-\beta_k] \subseteq [c_k;\, d_k]$, $c_k+\alpha_k = \alpha b_p$ and $d_k - \beta_k = \beta b_p$ for some $\alpha,\, \beta \in \mathbb{N}$, we obtain

$$\begin{aligned}
\underline{d}_{\log}(E_p) &\geq \liminf_{k\to\infty} \frac{\sum_{j=0}^{\beta-\alpha}(1/\alpha b_p + jb_p)}{\log(c_{k+1})} \\
&\geq \liminf_{k\to\infty} \frac{\log(\beta/\alpha)}{b_p \log(c_{k+1})} \\
&\geq \liminf_{k\to\infty} \frac{\log(d_k - \beta_k/c_k + \alpha_k)}{b_p \log(c_{k+1})} \\
&\geq \liminf_{k\to\infty} \frac{\log(2^{(1+\varepsilon)a^{2n_k}} - b_p/2^{(1-\varepsilon)a^{2n_k}} + b_p)}{b_p \log(2^{(1-\varepsilon)a^{2n_{k+1}}})}
\end{aligned}$$

https://doi.org/10.1017/etds.2019.103 Published online by Cambridge University Press

$$\geq \liminf_{k\to\infty} \frac{2\varepsilon a^{2n_k}}{b_p(1-\varepsilon)a^{2n_{k+1}}}$$
$$\geq \frac{2\varepsilon}{b_p(1-\varepsilon)a^{2M_p}} > 0. \qquad \square$$

Further, the following lemma is almost the same as [**4**, Lemma 3] and it still holds in our context.

LEMMA 4.4. *Let $p, q \geq 1$, $n \in E_p$, $m \in E_q$ with $n \neq m$. Then $|n-m| > \max(p, q)$.*

In particular, $(E_p + [0, p]) \cap (E_q + [0, q]) = \emptyset$ if $p \neq q$. Thus, the sequence of sets (E_p) satisfies conditions (a) and (b) in Theorem 4.1. Observe also that if we remove a finite number of elements from A_p we may suppose that $A_p \subset [p, \infty[$ and, for every $u \in A_p$, $I_u^{a,\varepsilon} + [-2p, 2p] \subset I_u^{a,2\varepsilon}$.

We now turn to the construction of the weights of the weighted shift we are looking for. For this construction, we also draw our inspiration from constructions made in [**4**]. We set

$$w_1^p \cdots w_k^p = \begin{cases} 1 & \text{if } k \notin b_p\mathbb{N} + [-4p, 4p], \\ 2^p & \text{if } k \in b_p\mathbb{N} + [-2p, 2p], \end{cases}$$

and for every $k \in \mathbb{N}$, $\frac{1}{2} \leq w_k^p \leq 2$. Then, for $p, q \geq 1$, $u \in A_p$ and $v \in A_q$ with $u > v$, we define

$$w_1^{u,v} \cdots w_k^{u,v} \begin{cases} = 1 & \text{if } k \notin I_u^{a,4\varepsilon}, \\ \geq \max(2^p, 2^q) & \text{if } k \in I_u^{a,\varepsilon} - I_v^{a,\varepsilon} + [0, p], \\ = 2^u & \text{if } k \in I_u^{a,\varepsilon} + [0, p], \end{cases}$$

with $\frac{1}{2} \leq w_k^{u,v} \leq 2$. We must stress that this construction is made possible by the following conditions.

- $I_u^{a,\varepsilon} - I_v^{a,\varepsilon} + [0, p] - [0, \max(p, q)] \subset I_u^{a,2\varepsilon} - I_v^{a,2\varepsilon} \subset I_u^{a,4\varepsilon}$, which ensures that we can pass from 1 to $\max(2^p, 2^q)$.
- $I_u^{a,\varepsilon} - [0, u] \subset I_u^{a,4\varepsilon}$ (i.e. $2^{(1-4\varepsilon)a^{2u}} + u \leq 2^{(1-\varepsilon)a^{2u}}$), which is possible if a is sufficiently big and ε is sufficiently small. Thus, it is possible to grow from 1 to 2^u.
- $I_u^{a,\varepsilon} + [0, p] + [0, u] \subset I_u^{a,\varepsilon} + [0, 2u] \subset I_u^{a,4\varepsilon}$, which is again possible if a is sufficiently big and ε is sufficiently small. This allows the weights to decrease from 2^u to 1.

We are now able to give the definition of the weight w. This is constructed in order to satisfy the equality

$$w_1 \cdots w_n = \max_{p,u,v}(w_1^p \cdots w_n^p, w_1^{u,v} \cdots w_n^{u,v}).$$

It is clear by construction that, for every $n \in \mathbb{N}$, $\frac{1}{2} \leq w_n \leq 2$, so the weighted backward shift B_w is bounded and invertible. Moreover, this construction satisfies condition (c) in Theorem 4.1.

Since we want to prove that B_w is log-frequently hypercyclic, the only condition left to prove is condition (d) in Theorem 4.1. Thus let $p, q \geq 1$, $n \in E_p$ and $m \in E_q$ with $m > n$ and $t \in [0, q]$. Then we have two cases.

https://doi.org/10.1017/etds.2019.103 Published online by Cambridge University Press

• If $p = q$, then $m - n + t \in b_q\mathbb{N} + [0, q]$ and the definition of w ensures that $w_1 \cdots w_{m-n+t} \geq 2^q$.

• If $p \neq q$, then there exists $u > v$ such that $n \in I_v^{a,\varepsilon}$ and $m \in I_u^{a,\varepsilon}$. Thus, by definition of w,

$$w_1 \cdots w_{m-n+t} \geq \max(2^p; 2^q) \geq 2^{p+q/2} \geq \lfloor 2^{p/2} \rfloor \cdot \lfloor 2^{q/2} \rfloor.$$

Now, one may define $M(p) := \lfloor 2^{p/2} \rfloor$ and each case above satisfies condition (d) in Theorem 4.1. Thus we have proved that the weighted shift B_w is logarithmically frequently hypercyclic.

We now turn to the frequent hypercyclicity of B_w. We will prove by contradiction that B_w is not frequently hypercyclic. Let us suppose that B_w is frequently hypercyclic. Also let x be a frequent hypercyclic vector and $E = \{n \in \mathbb{N} : \|B_w^n(x) - e_0\| \leq \frac{1}{2}\}$. Thus we have $\underline{d}(E) > 0$ and $\lim_{n\to\infty, n\in E} w_1 \cdots w_n = +\infty$. For every $p \geq 1$, we consider the set

$$F_p = \{n \in E : w_1 \cdots w_n > 2^p\}.$$

This set is a cofinite subset of E, so it has the same lower density. We also consider an increasing enumeration (n_k) of A_p. For readability reasons, we define some notation here for the end of this part. Let $[c_{k,\varepsilon}; d_{k,\varepsilon}] = I_{n_k}^{a,\varepsilon}$. Then we get

$$\underline{d}(F_p) \leq \liminf_{k\to\infty} \left(\frac{\#\{n \in F_p : n \leq c_{k+1,\varepsilon}\}}{c_{k+1,\varepsilon}} \right).$$

We write

$$\begin{aligned} \#\{n \in F_p : n \leq c_{k+1,\varepsilon}\} \leq{}& \#\{n \in F_p : n \leq d_{k,\varepsilon}\} \\ &+ \#\left\{ n \in \bigcup_{q>p} (b_q\mathbb{N} + [-4q; 4q]) : d_{k,\varepsilon} < n \leq c_{k+1,\varepsilon} \right\} \\ &+ \#\left\{ n \in \bigcup_{n_k \leq u \leq n_{k+1}} I_u^{a,4\varepsilon} : d_{k,\varepsilon} < n \leq c_{k+1,\varepsilon} \right\}. \end{aligned}$$

For the first term, observe that

$$\begin{aligned} \frac{\#\{n \in F_p : n \leq d_{k,\varepsilon}\}}{c_{k+1,\varepsilon}} &\leq \frac{d_{k,\varepsilon}}{c_{k+1,\varepsilon}} \\ &\leq 2^{(1-\varepsilon)a^{2n_{k+1}}((1+\varepsilon/1-\varepsilon)a^{2(n_k - n_{k+1})} - 1)} \\ &\leq 2^{(1-\varepsilon)a^{2n_{k+1}}(a^{-2}(1+\varepsilon/1-\varepsilon)-1)}. \end{aligned}$$

Moreover, we can easily check that $a^{-2}(1 + \varepsilon/1 - \varepsilon) - 1 < 0$ by (3). Thus we deduce that this first sum tends to zero. Let us estimate the third term $J_{k,\varepsilon}^{(3)} := (\#\{n \in \bigcup_{n_k \leq u \leq n_{k+1}} I_u^{a,4\varepsilon} : d_{k,\varepsilon} < n \leq c_{k+1,\varepsilon}\}/c_{k+1,\varepsilon})$. We have

$$\begin{aligned} J_{k,\varepsilon}^{(3)} &\leq \frac{\sum_{u=n_k}^{n_{k+1}} \sum_{j=2^{(1-4\varepsilon)a^{2u}}}^{2^{(1+4\varepsilon)a^{2u}}} 1}{c_{k+1,\varepsilon}} \\ &\leq \sum_{u=n_k}^{n_{k+1}} \frac{2^{(1+4\varepsilon)a^{2u}} - 2^{(1-4\varepsilon)a^{2u}} + 1}{2^{(1-\varepsilon)a^{2n_{k+1}}}} \end{aligned}$$

https://doi.org/10.1017/etds.2019.103 Published online by Cambridge University Press

$$\leq \sum_{u=n_k}^{n_{k+1}} 2^{(1-\varepsilon)a^{2n_{k+1}}((1+4\varepsilon/1-\varepsilon)a^{2(u-n_{k+1})}-1)}(1-2^{-8\varepsilon a^{2u}}+2^{-(1+4\varepsilon)a^{2u}})$$

$$\leq 2\sum_{u=n_k}^{n_{k+1}} 2^{(1-\varepsilon)a^{2n_{k+1}}(a^{-2}(1+4\varepsilon/1-4\varepsilon)-1)}$$

$$\leq 2(1+M_p)2^{(1-\varepsilon)a^{2n_{k+1}}(a^{-2}(1+4\varepsilon/1-4\varepsilon)-1)} \to 0.$$

We now focus on the second term, $J_{k,\varepsilon}^{(2)} := (\#\{n \in \bigcup_{q>p}(b_q\mathbb{N}+[-4q;4q]) : d_{k,\varepsilon} < n \leq c_{k+1,\varepsilon}\})/(c_{k+1,\varepsilon})$. Notice that the union over $q > p$ is in fact a finite union. For those $q > p$ such that $c_{k+1,\varepsilon} + 4q - (d_{k,\varepsilon} - 4q) \geq b_q$ we get

$$\begin{aligned}
&\#\{]d_{k,\varepsilon}; c_{k+1,\varepsilon}] \cap (b_q\mathbb{N}+[-4q;4q])\} \\
&\quad \leq (8q+1)\#\{]2^{(1+\varepsilon)a^{2n_k}} - 4q; 2^{(1-\varepsilon)a^{2n_{k+1}}} + 4q] \cap b_q\mathbb{N}\} \\
&\quad \leq 3(8q+1)\frac{2^{(1-\varepsilon)a^{2n_{k+1}}} - 2^{(1+\varepsilon)a^{2n_k}} + 8q}{b_q}.
\end{aligned}$$

Moreover, by condition (4.1), there is no $q > p$ such that $c_{k+1,\varepsilon} - d_{k,\varepsilon} + 8q < b_q < c_{k+1,\varepsilon} + 4q$. Thus, replacing the finite sum by an infinite one, we obtain that

$$\begin{aligned}
J_{k,\varepsilon}^{(2)} &\leq \sum_{q>p} \frac{3(8q+1)}{2^{(1-\varepsilon)a^{2n_{k+1}}}} \frac{2^{(1-\varepsilon)a^{2n_{k+1}}} - 2^{(1+\varepsilon)a^{2n_k}} + 8q}{b_q} \\
&\leq \sum_{q>p} 3(8q+1)\frac{1 - 2^{(1+\varepsilon)a^{2n_k}-(1-\varepsilon)a^{2n_{k+1}}} + 8q}{b_q} \\
&\leq 3\sum_{q>p}(8q+1)\frac{1 - 2^{(1-\varepsilon)a^{2n_{k+1}}}((1+\varepsilon/1-\varepsilon)a^{2(n_k-n_{k+1})} - 1) + 8q}{b_q} \\
&\leq 6\sum_{q>p}\frac{8q+1}{b_q}.
\end{aligned}$$

Since this last inequality does not require any property on p, we can let p tend to infinity which, thanks to (4.2), implies that $\underline{d}(E) = \lim_{p\to\infty} \underline{d}(F_p) = 0$, and we obtain a contradiction. Thus the weighted shift B_w is not frequently hypercyclic. From this construction, we deduce the following result.

THEOREM 4.5. *There exists a log-frequently hypercyclic operator that is not frequently hypercyclic.*

Acknowledgements. The authors were partly supported by grant ANR-17-CE40-0021 of the French National Research Agency ANR (project Front) and by Programme PEPS JC 2018 INSMI.

https://doi.org/10.1017/etds.2019.103 Published online by Cambridge University Press

References

[**1**] F. Bayart and S. Grivaux. Frequently hypercyclic operators. *Trans. Amer. Math. Soc.* **358**(11) (2006), 5083–5117.
[**2**] F. Bayart and S. Grivaux. Invariant Gaussian measures for operators on banach spaces and linear dynamics. *Proc. Lond. Math. Soc. (3)* **94**(1) (2007), 181–210.
[**3**] F. Bayart and É. Matheron. *Dynamics of Linear Operators (Cambridge Tracts in Mathematics)*. Cambridge University Press, Cambridge, 2009.
[**4**] F. Bayart and I. Z. Ruzsa. Difference sets and frequently hypercyclic weighted shifts. *Ergod. Th. & Dynam. Sys.* **35**(3) (2015), 691–709.
[**5**] J. Bès, Q. Menet, A. Peris and Y. Puig. Recurrence properties of hypercyclic operators. *Math. Ann.* **366**(1–2) (2016), 545–572.
[**6**] A. Bonilla and K.-G. Grosse-Erdmann. Frequently hypercyclic operators and vectors. *Ergod. Th. & Dynam. Sys.* **27**(2) (2007), 383–404.
[**7**] A. Bonilla and K.-G. Grosse-Erdmann. Erratum: Frequently hypercyclic operators and vectors. *Ergod. Th. & Dynam. Sys.* **29**(6) (2009), 1993–1994.
[**8**] R. Ernst and A. Mouze. A quantitative interpretation of the frequent hypercyclicity criterion. *Ergod. Th. & Dynam. Sys.* **39**(4) (2019), 898–924.
[**9**] A. R. Freedman and J. J. Sember. Densities and summability. *Pacific J. Math.* **95**(2) (1981), 293–305.
[**10**] S. Grivaux. Personal communication (2018).
[**11**] C. Jardas and N. Sarapa. A summability method in some strong laws of large numbers. *Math. Commun.* **2**(2) (1997), 107–124.
[**12**] M. Murillo-Arcila and A. Peris. Strong mixing measures for linear operators and frequent hypercyclicity. *J. Math. Anal. Appl.* **398**(2) (2013), 462–465.

https://doi.org/10.1017/etds.2019.103 Published online by Cambridge University Press