Euro. Jnl of Applied Mathematics (2005), *vol.* 16, *pp.* 583–599. © 2005 Cambridge University Press
doi:10.1017/S0956792505006418 Printed in the United Kingdom

A new formulation of the dam problem

S. CHALLAL and A. LYAGHFOURI

King Fahd University of Petroleum and Minerals,
P.O. Box 728, Dhahran 31261, Saudi Arabia
email: lyaghfo@kfupm.edu.sa

(*Received 22 July 2004; revised 24 May 2005*)

We study a heterogeneous dam supplied by two reservoirs, for which we propose a new formulation based on the stream function. Without any monotonicity assumption on the permeability matrix, we prove that the free boundary is a continuous curve of the form $x = \phi(y)$. We also prove the uniqueness of the solution.

Introduction

The dam problem consists of studying the filtration of a fluid through a porous medium Ω subject to Darcy's law and to some boundary conditions. We look for the saturated region S (see Figure 1) and the fluid pressure p inside Ω. The interface $\Gamma = \partial[p > 0] \cap \Omega$ which separates the wet and dry regions is called a free boundary.

This problem has been studied by many authors. First, Baiocchi [5] solved the case of a homogeneous rectangular dam supplied by two reservoirs of water, by using the theory of variational inequalities. He also proved that Γ is an analytic curve of the form $y = \Phi(x)$. Benci [6] extended Baiocchi's results to rectangular heterogeneous dams with permeability of the form $k_1(x)k_2(y)I_2$, where I_2 is the 2-by-2 unit matrix. In particular, he proved that Γ is a continuous decreasing curve $y = \Phi(x)$ if $k_2'(y) \geqslant 0$. Caffarelli & Friedman [8] proved that Γ is a curve of the form $x = \Psi(y)$ if $k_1(x) = 1$ and $k_2(y)$ is a non-increasing step function.

Unfortunately, the variational inequalities approach is not possible for dams with general geometry. Later, Alt [1] and Brezis *et al.* [7] introduced a new approach. Moreover, for the homogeneous dam Alt [2] proved that the free boundary is an analytic curve $y = \Phi(x)$. Uniqueness of the S_3-connected solution was proved by Carrillo & Chipot [9] and also by Alt & Gilardi [4].

Friedman & Huang [12] and Stavre & Vernescu [15] showed that if the permeability matrix is given by $a(X) = k(x, y)I_2$ with $\partial k/\partial y \geqslant 0$ in $\mathscr{D}'(\Omega)$, then Γ is a continuous curve $y = \Phi(x)$ and the S_3-connected solution is unique. These results were generalized by Lyaghfouri [14] to the case where the permeability matrix $a(X) = (a_{ij}(X))$ is such that $a_{12}(X) \equiv 0$ and $\partial a_{22}/\partial y \geqslant 0$ in $\mathscr{D}'(\Omega)$. Recently, the authors have considered the case where $a_{12}, a_{22} \in C^1$ and $div(a_{12}, a_{22}) \geqslant 0$. They proved, via a change of variables, that Γ is locally represented by continuous curves.

In this paper, we consider a heterogeneous dam supplied by two reservoirs each containing the same fluid. Our main objective is to study the free boundary, and to

https://doi.org/10.1017/S0956792505006418 Published online by Cambridge University Press

establish some of its properties without assuming any kind of monotonicity on the permeability. This is done by introducing a new formulation using the stream function instead of the classical one based on the pressure. In this formulation the total flux through the left-hand side of the dam is prescribed instead of prescribing the height of the left reservoir and the pressure on the contact zone between the reservoir and the dam. Note that a similar formulation has been introduced [3] to study a homogenous dam supplied by two reservoirs each containing two immiscible fluids.

We first prove a monotonicity result for the function characterizing the wet region. This allows us to define the free boundary as a curve of the form $x = \phi(y)$. Then we prove the continuity of ϕ and the uniqueness of the solution.

1 Statement of the problem

Throughout this paper, we will denote by X the point of coordinates (x, y). Sometimes we will also use the notation (x_P, y_P) to denote the coordinates of the point P. We will denote by $\chi(E)$ the characteristic function of the set E.

We consider a dam which is a porous medium represented by a bounded domain Ω of $\mathbb{R}^2$ with boundary $\partial\Omega = \overset{\frown}{AB} \cup \overset{\frown}{BT} \cup \overset{\frown}{TA}$. We assume that

$$\begin{aligned}
\overset{\frown}{TA} &= \{(a_0(y), y),\ y \in [0, y_T]\} && \text{with } y \mapsto a_0(y) \text{ an increasing } C^1 \text{ function}\\
\overset{\frown}{BT} &= \{(b_0(y), y),\ y \in [y_B, y_T]\} && \text{with } y \mapsto b_0(y) \text{ a decreasing } C^1 \text{ function}\\
\overset{\frown}{AB} &= \{(x, c(x)),\ x \in [x_A, x_B]\} && \text{with } x \mapsto c(x) \text{ a decreasing } C^1 \text{ function.}
\end{aligned}$$

The dam is supplied by two reservoirs of water (see Figure 1). To the left, we assume that the total entering flux through $\overset{\frown}{A_1A}$ is equal to the constant $Q > 0$ and the fluid in the reservoir is at the level $H(Q) = y_{A_1}$ depending on Q, while to the right it is at the level $h = y_{B_1}$ independent on Q, with $H > h > 0$.

We assume that the flow in Ω has reached a steady state and we denote by S the saturated region of the porous medium. The boundary ∂S of S is divided into four parts (see Figure 1): the impervious part $\overset{\frown}{AB}$, the free boundary Γ, $\overset{\frown}{BB_1}$ and $\overset{\frown}{A_1A}$ representing the part in contact with the reservoirs and, finally, $\overset{\frown}{B_1O_1}$, the part where the fluid flows out of Ω and not directly into a reservoir.

The flow is governed by the following Darcy law:

$$v = -a(X)\nabla(p + y) \tag{1.1}$$

where v is the fluid velocity, p is its pressure, and $a : \bar{\Omega} \longrightarrow \mathbb{R}^{2\times 2}$ is the permeability matrix satisfying

$$\begin{cases} i) & a \in C^{0,\alpha}(\Omega), \qquad \max\limits_{i,j} |a_{i,j}|_{0,\Omega} \leqslant M \\ ii) & a(X)\xi.\xi \geqslant m|\xi|^2 \qquad \text{for all } \xi \in \mathbb{R}^2, \quad X \in \Omega, \end{cases} \tag{1.2}$$

for some $m, M > 0$ and $\alpha \in (0, 1)$ and where $|.|_{0,\Omega}$ is the L^∞ norm.

https://doi.org/10.1017/S0956792505006418 Published online by Cambridge University Press




FIGURE 1. Problem statement illustration.

Since $\widehat{AB}$ is assumed to be impervious, we have

$$v.v = 0 \quad \text{on} \quad \widehat{AB}, \tag{1.3}$$

where v is the exterior unit normal vector to ∂S.

We assume that there is no flux through the free boundary and that p agrees with the atmospheric pressure normalized at zero. So we have

$$p = 0 \quad \text{and} \quad v.v = 0 \quad \text{on} \quad \Gamma. \tag{1.4}$$

On the part of $\partial\Omega$ in contact with the reservoirs, we assume that the pressure is given by the exterior fluid pressure, i.e. we have

$$p = h - y \quad \text{on} \quad \widehat{BB_1}, \qquad p = H - y \quad \text{on} \quad \widehat{A_1A}. \tag{1.5}$$

Finally, on $\widehat{B_1O_1}$, p coincides with the atmospheric pressure and the water is free to exit the porous medium. Thus, we have

$$p = 0 \quad \text{and} \quad v.v \geqslant 0 \quad \text{on} \quad \widehat{B_1O_1}. \tag{1.6}$$

We assume the flow to be incompressible, i.e. $div(v) = 0$ in S. Then there exists a stream function ψ such that

$$v = -Rot\psi = -\left(-\frac{\partial\psi}{\partial y}, \frac{\partial\psi}{\partial x}\right) = -a(X)\nabla(p + y) \quad \text{in} \quad S. \tag{1.7}$$

Let $\zeta \in \mathscr{D}(S)$. We have by (1.1) and (1.7)

$$0 = \int_S \nabla(p + y).Rot\zeta = \int_S a^{-1}(X)Rot\psi.Rot\zeta = \int_S b(X)\nabla\psi.\nabla\zeta \tag{1.8}$$

where $b(X) = \frac{{}^t a(X)}{det(a(X))}$, $det(a(X))$ is the determinant of the matrix $a(X)$ and ${}^t a(X)$ is its transpose. Note that we deduce from (1.2) the existence of $\lambda, \Lambda > 0$ such that

$$\begin{cases} i) & b \in C^{0,\alpha}(\Omega), \qquad \max\limits_{i,j} |b_{i,j}|_{0,\Omega} \leqslant \Lambda \\ ii) & b(X)\xi.\xi \geqslant \lambda|\xi|^2 \quad \text{for all } \xi \in \mathbb{R}^2, \quad X \in \Omega. \end{cases} \tag{1.9}$$

https://doi.org/10.1017/S0956792505006418 Published online by Cambridge University Press

We obtain from (1.8)

$$div(b(X)\nabla\psi) = 0 \qquad \text{in } \mathscr{D}'(S). \tag{1.10}$$

Let us set $v = (v_x, v_y)$ and denote the tangent vector by $\tau = (-v_y, v_x)$. Then

$$v.v = -Rot\psi.v = \frac{\partial\psi}{\partial\tau}. \tag{1.11}$$

Moreover, one can verify that

$$b(X)\nabla\psi.v = \frac{\partial(p+y)}{\partial\tau}. \tag{1.12}$$

It follows from (1.3) and (1.11) that ψ is constant along $\widehat{AB}$. So there exists a constant c_1 such that

$$\psi = c_1 \quad \text{ on } \widehat{AB}\,. \tag{1.13}$$

In the same way, ψ is constant along Γ. So there exists a constant c_2 such that

$$\psi = c_2 \quad \text{ on } \Gamma\,. \tag{1.14}$$

Now, since $p = 0$ on Γ, we obtain from (1.12)

$$b(X)\nabla\psi.v = v_x \quad \text{ on } \Gamma\,. \tag{1.15}$$

By (1.5), $p + y$ is constant along $\widehat{A_1A}$ and $\widehat{BB_1}$. It follows then by (1.12)

$$b(X)\nabla\psi.v = 0 \quad \text{ on } \widehat{A_1A} \cup \widehat{BB_1}\,. \tag{1.16}$$

Finally, we deduce from (1.6) and (1.11)–(1.12) that

$$b(X)\nabla\psi.v = v_x \quad \text{ and } \quad \frac{\partial\psi}{\partial\tau} \geqslant 0 \quad \text{ on } \widehat{B_1O_1}\,. \tag{1.17}$$

We choose $c_1 = 0$ and we are going to prove, in a formal way, that $c_2 > 0$. First we claim that $c_2 \geqslant 0$. Indeed, suppose that $c_2 < 0$ (see Figure 2).

To begin with, since $h < H$, we expect that the point O_1 is located above B_1, and in particular $\widehat{O_1B_1} \neq \emptyset$. Next, since by (1.17) ψ increases from B_1 to O_1, it takes its minimum value on $\widehat{A_1A}$ or on $\widehat{BB_1}$. Suppose there exists $m_0 \in \widehat{BB_1}$ at which ψ achieves its minimum value. Then $\frac{\partial\psi}{\partial\tau}(m_0) = 0$, and consequently, we have $0 = b(m_0)\nabla\psi.v(m_0) = \frac{\partial\psi}{\partial v}(m_0)b(m_0)v.v$. From (1.9)*ii*) we deduce that $\frac{\partial\psi}{\partial v}(m_0) = 0$, which contradicts the Hopf maximum principle.

Similarly, we get a contradiction if ψ achieves its minimum on $\widehat{A_1A}$.

Now, we assert that $c_2 > 0$. Indeed if $c_2 = 0$, then we cannot have as before, a minimum neither on $\widehat{A_1A}$ nor on $\widehat{BB_1}$. So the minimum is achieved on $\widehat{O_1A_1} \cup \widehat{AB} \cup \widehat{B_1O_1}$. But since $\frac{\partial\psi}{\partial\tau} \geqslant 0$ on $\widehat{B_1O_1}$ and $\psi = 0$ on $\widehat{O_1A_1} \cup \widehat{AB}$, the minimum will be achieved at a point $m_0 \in \widehat{B_1O_1}$. We necessarily have, as before, $\frac{\partial\psi}{\partial\tau}(m_0) = 0$, which leads to $(b(m_0)v.v)\frac{\partial\psi}{\partial v}(m_0) =$

https://doi.org/10.1017/S0956792505006418 Published online by Cambridge University Press




FIGURE 2. Strong formulation of the problem.

$v_x > 0$. Hence by (1.9)*ii*) $\frac{\partial\psi}{\partial v}(m_0) > 0$ and we get a contradiction to the Hopf maximum principle.

Remark 1.1 (*i*) We have $c_2 = Q$ is the total flux through $\widehat{A_1A}$. Indeed we have by (1.11)

$$Q = \int_A^{A_1} -v.v = \int_A^{A_1} -\frac{\partial\psi}{\partial\tau} = \psi(A_1) - \psi(A) = c_2 - 0 = c_2.$$

Moreover, by the strong maximum principle we have $0 < \psi < Q$ in S.

(*ii*) Since $b(X)\nabla\psi.v = v_x$ and $\psi = Q$ on Γ, one has $\frac{\partial\psi}{\partial v}b(X)v.v = v_x$ on Γ. Moreover by the maximum principle and the positivity of $b(X)$, we deduce that $v_x \geqslant 0$ on Γ. This means that Γ is a curve $x = \phi(y)$. This will be justified more rigourously in § 2.

To derive a weak formulation for our problem, we first extend ψ by Q to $\Omega \setminus S$. Next we choose a test function $\zeta \in H^1(\Omega)$ with $\zeta = 0$ on $\widehat{AB}$. Taking into account (1.10), (1.15)–(1.17), we obtain, in a formal way,

$$\int_\Omega b(X)\nabla\psi.\nabla\zeta = \int_S b(X)\nabla\psi.\nabla\zeta = \int_{\partial S} b(X)\nabla\psi.v\zeta = \int_{\Gamma\cup\widehat{B_1O_1}} v_x\zeta. \tag{1.18}$$

Assuming Γ smooth enough, we have

$$\int_\Omega \chi([\psi = Q])\zeta_x = \int_{\partial\Omega} \zeta v_x - \int_{\partial S} \zeta v_x = \int_{\widehat{O_1T}\cup\widehat{TA_1}} v_x\zeta - \int_\Gamma v_x\zeta$$

which leads to

$$\int_\Gamma v_x\zeta = -\int_\Omega \chi([\psi = Q])\zeta_x + \int_{\widehat{O_1T}\cup\widehat{TA_1}} v_x\zeta. \tag{1.19}$$

Comparing (1.18) and (1.19), we get

$$\int_\Omega b(X)\nabla\psi.\nabla\zeta + \int_\Omega \chi([\psi = Q])\zeta_x = \int_{\widehat{TA_1}} v_x\zeta + \int_{\widehat{B_1T}} v_x\zeta. \tag{1.20}$$

https://doi.org/10.1017/S0956792505006418 Published online by Cambridge University Press

Since we are assuming that $\psi = Q$ above the level $y = H$, we can rewrite (1.20) as

$$\int_\Omega \big(b(X)\nabla\psi + \chi([\psi = Q])e_x\big).\nabla\zeta - \int_{\widehat{TA}} \chi([\psi = Q])v_x\zeta = \int_{\widehat{B_1T}} v_x\zeta,$$

where $e_x = (0, 1)$. Hence we are led to the following weak formulation:

$$(P)\begin{cases} \text{Find } (\psi, \gamma, \widetilde{\gamma}) \in H^1(\Omega) \times L^\infty(\Omega) \times L^\infty(\widehat{TA}) \text{ such that :} \\ (i) \quad \displaystyle\int_\Omega \big(b(X)\nabla\psi + \gamma e_x\big).\nabla\zeta - \int_{\widehat{TA}} \widetilde{\gamma} v_x\zeta = \int_{\widehat{B_1T}} v_x\zeta \\ \qquad\qquad \forall\zeta \in H^1(\Omega), \quad \zeta = 0 \text{ on } \widehat{AB} \\ (ii) \quad \gamma \in H(\psi - Q) \quad \text{a.e. in } \Omega, \qquad \widetilde{\gamma} \in H(\psi - Q) \quad \text{a.e. in } \widehat{TA} \\ (iii) \quad \psi = 0 \quad \text{on } \widehat{AB}, \end{cases}$$

where H is the heaviside graph.

Remark 1.2 As mentioned in Alt et al. [4], in the case of two fluids, the new formulation is given in terms of prescribed flux Q instead of prescribed height of the left reservoir. Indeed in the formulation (P) one is given Q as well as B_1, but not A_1.

The existence of a solution of (P) can be obtained by arguing as in Alt [1] or Brézis *et al.* [7]. In § 2, we give some properties of the solutions. In particular we prove the continuity of ψ. Then we show a monotonicity result of γ with respect to x. This allows us to define the free boundary as a curve $x = \phi(y)$. We also prove that the dam is wet below the line $y = h$. In § 3, we establish the continuity of ϕ and obtain the expression of γ as a characteristic function of the dry part of the dam. Finally, in § 4, we prove the uniqueness of the solution.

2 Some properties of the solutions

Throughout this section we denote a solution of (P) by $(\psi, \gamma, \widetilde{\gamma})$. First we have the following proposition.

Proposition 2.1

$$div\big(b(X)\nabla\psi\big) = -\gamma_x \qquad in \quad \mathscr{D}'(\Omega). \tag{2.1}$$

Proof It suffices to take $\zeta \in \mathscr{D}(\Omega)$ as a test function in $(P)i)$. □

Proposition 2.2

$$0 \leqslant \psi \leqslant Q \qquad in \quad \Omega. \tag{2.2}$$

https://doi.org/10.1017/S0956792505006418 Published online by Cambridge University Press

Proof (i) Using ψ^- as a test function in $(P)i)$, we obtain since $[\psi \leqslant 0] \subset [\psi < Q]$, $\gamma = 0$ a.e. in $[\psi < Q]$ and $\widetilde{\gamma} = 0$ a.e. in $\widehat{TA} \cap [\psi < Q]$,

$$\int_\Omega b(X)\nabla\psi^-.\nabla\psi^- = -\int_{\widehat{B_1T}} \psi^- v_x \leqslant 0 \qquad \text{since } v_x \geqslant 0 \text{ on } \widehat{B_1T}\,.$$

By $(1.9)ii)$, we deduce that $\nabla\psi^- = 0$ a.e. in Ω. Then ψ^- is constant in Ω. Since $\psi = 0$ on $\widehat{AB}$, we obtain $\psi \geqslant 0$ in Ω.

(ii) Taking $(\psi - Q)^+$ as a test function in $(P)i)$, we obtain since $\gamma = 1$ a.e. in $[\psi > Q]$, $\widetilde{\gamma} = 1$ a.e. in $\widehat{TA} \cap [\psi > Q]$, and $v_x \geqslant 0$ on $\widehat{BB_1}$

$$\int_\Omega b(X)\nabla(\psi - Q)^+.\nabla(\psi - Q)^+ = -\int_\Omega (\psi - Q)_x^+ + \int_{\widehat{TA}} (\psi - Q)^+ v_x$$
$$+ \int_{\widehat{B_1T}} (\psi - Q)^+ v_x = -\int_{\widehat{BB_1}} (\psi - Q)^+ v_x \leqslant 0.$$

It follows that $(\psi - Q)^+$ is constant in Ω. Since $\psi = 0$ on $\widehat{AB}$, we obtain $\psi \leqslant Q$ in Ω. □

Proposition 2.3

$$\psi \in C^{0,\alpha}_{loc}(\overline{\Omega} \setminus \{A, B, T\}) \qquad \textit{for some} \quad \alpha \in (0, 1).$$

Proof First, since $\gamma \in L^\infty(\Omega)$, and $b \in L^\infty(\Omega)$ and is strictly elliptic, we deduce from (2.1) and the regularity theory for elliptic equations (see Gilbarg & Trudinger [13, Theorem 8.22, p. 200; Theorem 8.29, p. 205] that $\psi \in C^{0,\alpha}_{loc}(\Omega \cup Int(\widehat{AB}))$ for some $\alpha \in (0, 1)$.

For the regularity on $Int(\widehat{BT}) \cup Int(\widehat{TA})$, we use the reflection method. We will give the proof only for $Int(\widehat{TA})$, as it is the same for $Int(\widehat{BT})$.

Let $X_0 = (x_0, y_0) = (a_0(y_0), y_0) \in Int(\widehat{TA})$, $\delta > 0$ and consider the set

$$Z = \{(x, y) \,/\, a_0(y) - \delta < x < a_0(y) + \delta \quad \text{and} \quad y_0 - \delta < y < y_0 + \delta\}.$$

We assume that $\delta > 0$ is small enough so that $\{(a_0(y), y) \,/\, y_0 - \delta < y < y_0 + \delta\} \subset \widehat{TA}$ and (see Figure 3)

$$Z \cap \Omega = \{(x, y) \,/\, a_0(y) < x < a_0(y) + \delta \quad \text{and} \quad y_0 - \delta < y < y_0 + \delta\}.$$

Then we define the following functions in Z:

$$\overline{\psi}(x, y) = \begin{cases} \psi(x, y) & \text{if } (x, y) \in Z \cap \Omega \\ \psi(2a_0(y) - x, y) & \text{otherwise} \end{cases} \quad \text{and} \quad \overline{b}(x, y) = \begin{cases} b(x, y) & \text{if } (x, y) \in Z \cap \Omega \\ c(2a_0(y) - x, y) & \text{otherwise;} \end{cases}$$

where

$$\begin{cases} c_{11} = b_{11} + 4a_0'^2(y)b_{22} - 2a_0'(y)(b_{12} + b_{21}) \\ c_{12} = 2a_0'(y)b_{22} - b_{12},\ c_{21} = 2a_0'(y)b_{22} - b_{21},\ c_{22} = b_{22}. \end{cases}$$

https://doi.org/10.1017/S0956792505006418 Published online by Cambridge University Press




FIGURE 3. $C^{0,\alpha}$ regularity upto $\mathrm{Int}(\widehat{\mathrm{TA}})$.

Let $f = \overline{\gamma}\chi(Z \setminus \Omega) + (2\widetilde{\gamma}(y) - \gamma)\chi(Z \cap \Omega)$, where $\overline{\gamma}(x, y) = \gamma(2a_0(y) - x, y)$ if $(x, y) \in Z \setminus \Omega$. Then we have

$$div(\overline{b}(x, y)\nabla\overline{\psi}) = f_x \quad \text{in } \mathscr{D}'(Z). \tag{2.3}$$

Indeed we have, for $\zeta \in \mathscr{D}(Z)$ and $\overline{\zeta}(x, y) = \zeta(2a_0(y) - x, y)$,

$$\begin{aligned}\int_{Z\setminus\Omega} c(2a_0(y) - x, y)\nabla\overline{\psi}.\nabla\zeta &= \int_{Z\cap\Omega} c(x, y)\nabla\overline{\psi}(2a_0(y) - x, y)\nabla\zeta(2a_0(y) - x, y) \\ &= \int_{Z\cap\Omega} b(x, y)\nabla\psi(x, y).\nabla\overline{\zeta}(x, y). \end{aligned}\tag{2.4}$$

Using (2.4) and the fact that $\zeta + \overline{\zeta}$ is a test function for (P), we get

$$\begin{aligned}\int_Z \overline{b}(x, y)\nabla\overline{\psi}.\nabla\zeta &= \int_{Z\cap\Omega} b(x, y)\nabla\psi.\nabla(\zeta + \overline{\zeta}) \\ &= -\int_{Z\cap\Omega} \gamma(\zeta + \overline{\zeta})_x + \int_{Z\cap\widehat{TA}} \widetilde{\gamma}(y)(\zeta + \overline{\zeta})\nu_x \\ &= \int_{Z\setminus\Omega} \overline{\gamma}\zeta_x + \int_{Z\cap\Omega} (2\widetilde{\gamma}(y) - \gamma)\zeta_x = \int_Z f\zeta_x.\end{aligned}$$

Now it is clear that the coefficients of $\overline{b}(x, y)$ and f are uniformly bounded in Z. Moreover we claim that $\overline{b}(x, y)$ is strictly elliptic in Z. Indeed it is enough to verify that $c(x, y)$ is strictly elliptic in $Z \cap \Omega$. Let $c^*(x, y)$ be the matrix defined by $c^*_{ii} = c_{ii}$ and $c^*_{ij} = -c_{ij}$ for $i \neq j$. Then we have for each $\xi = (\xi_1, \xi_2) \in \mathbb{R}^2$

$$c(x, y).\xi.\xi = c_{11}\xi_1^2 + (c_{12} + c_{21})\xi_1\xi_2 + c_{22}\xi_2^2 = c^*(x, y).\xi^*.\xi^*, \quad \text{with } \xi^* = (\xi_1, -\xi_2).$$

Since $|\xi| = |\xi^*|$, it is clear that it is enough to show the strict ellipticity of $c^*(x, y)$. For this purpose we remark that $c^*(x, y) =^t d(x, y)b(x, y)d(x, y)$, where $d(x, y) = (d_{ij}(x, y))$ is the matrix defined by $d_{11}(x, y) = d_{22}(x, y) = 1$, $d_{12}(x, y) = 0$, and $d_{21}(x, y) = -2a_0'(y)$. Then we have by (1.9)*ii*) for $\xi = (\xi_1, \xi_2) \in \mathbb{R}^2$

$$c^*(x, y)\xi.\xi =^t d(x, y)b(x, y)d(x, y)\xi.\xi = b(x, y)(d(x, y)\xi).(d(x, y)\xi) \geqslant \lambda|d(x, y)\xi|^2.$$

Since $|d(x, y)\xi|^2 = (1 + 4a_0'^2(y))\xi_1^2 - 4a'(y)\xi_1\xi_2 + \xi_2^2$, it is enough to choose η such that $0 < \eta < \min(\frac{1}{2}, \frac{1}{16\max_{y_0-\delta\leqslant y\leqslant y_0+\delta} a_0'^2(y)})$, to get $|d(x, y)\xi|^2 \geqslant \eta|\xi|^2$.

https://doi.org/10.1017/S0956792505006418 Published online by Cambridge University Press

Now using (2.3) and taking into account the fact that $f \in L^\infty(Z)$, $\overline{b} \in L^\infty(Z)$, and the strict ellipticity of $\overline{b}(x,y)$, we conclude (see Gilbarg & Trudiger [13, Theorem 8.22, p. 200]) that $\overline{\psi} \in C^{0,\alpha}_{loc}(Z)$ for some $\alpha \in (0,1)$. Thus $\psi \in C^{0,\alpha}_{loc}(\Omega \cup Int(\widehat{TA}))$. □

The following monotonicity result plays a major role later.

Proposition 2.4

$$\gamma_x \geqslant 0 \qquad in \quad \mathscr{D}'(\Omega). \tag{2.5}$$

Proof Let $\zeta \in \mathscr{D}(\Omega)$, $\zeta \geqslant 0$. For $\epsilon > 0$, $\xi = \min(\frac{Q-\psi}{\epsilon}, \zeta)$ is a test function for $(P)i)$. Using the fact that $\xi = 0$ on $[\psi = Q]$ and $\gamma = 0$ on $[\psi < Q]$, we get

$$\int_{\Omega \cap [Q-\psi \geqslant \epsilon\zeta]} b(X)\nabla(Q-\psi).\nabla\zeta = -\frac{1}{\epsilon}\int_{\Omega\cap[Q-\psi<\epsilon\zeta]} b(X)\nabla(Q-\psi).\nabla(Q-\psi) \leqslant 0.$$

Letting $\epsilon \to 0$ and using (2.1), we get the desired result. □

As a consequence of Propositions 2.1, 2.2 and (2.4), we obtain the following proposition.

Proposition 2.5

$$\psi > 0 \qquad in \quad \Omega. \tag{2.6}$$

Proof Indeed, using (2.1) and (2.5), we deduce that $div(b(X)\nabla\psi) \leqslant 0$. Since moreover $\psi \geqslant 0$ in Ω, we obtain by the strong maximum principle that either $\psi \equiv 0$ in Ω or $\psi > 0$ in Ω. If we have $\psi \equiv 0$ in Ω, then $\gamma = 0$ a.e. in Ω and $\widetilde{\gamma} = 0$ a.e. in $\widehat{TA}$. It follows from $(P)i)$

$$\int_{\widehat{B_1T}} \zeta v_x = 0 \qquad \forall \zeta \in H^1(\Omega), \quad \zeta = 0 \text{ on } \widehat{AB}$$

which is impossible. Hence $\psi > 0$ in Ω. □

As a consequence of the continuity of ψ and the monotonicity of γ, we deduce the following key proposition.

Proposition 2.6 *Let $(x_0, y_0) \in \Omega$. Then we have*

(i) *If* $\psi(x_0,y_0) = Q$ *then* $\psi(x,y_0) = Q$ *for all* $x \geqslant x_0$.

(ii) *If* $\psi(x_0,y_0) < Q$ *then there exists* $\epsilon > 0$ *such that*

$$\psi(x,y) < Q \qquad \forall (x,y) \in D_\epsilon = \Big(B_\epsilon(x_0,y_0) \cup \big((-\infty,x_0)\times(y_0-\epsilon,y_0+\epsilon)\big)\Big) \cap \Omega.$$

Proof It suffices to prove (ii). Let $(x_0,y_0) \in \Omega$ be such that $\psi(x_0,y_0) < Q$. By continuity, there exists $\epsilon > 0$ such that $\psi < Q$ in $B_\epsilon(x_0,y_0)$. Then $\gamma = 0$ a.e. in $B_\epsilon(x_0,y_0)$. Since $\gamma_x \geqslant 0$ in $\mathscr{D}'(\Omega)$ and $0 \leqslant \gamma \leqslant 1$, we obtain $\gamma = 0$ a.e. in D_ϵ. This leads by (2.1) to $div(b(X)\nabla\psi) = 0$

https://doi.org/10.1017/S0956792505006418 Published online by Cambridge University Press

in D_ϵ. Finally, given that $\psi \leqslant Q$ in Ω and $\psi < Q$ in $B_\epsilon(x_0, y_0)$, we deduce by the strong maximum principle that $\psi < Q$ in D_ϵ. □

The following theorem shows that the dam is wet under the level $y = h$.

Theorem 2.1 *We have*

$$\gamma = 0 \quad a.e.\ in \quad \Omega \cap [y < h] = \Omega_0 \qquad and \qquad \psi < Q \quad in \quad \Omega_0.$$

Proof Let $u = Q - \psi$ and $\chi = 1 - \gamma$. Then we have $\chi = 1$ a.e. in $[u > 0]$ and $\chi \in [0, 1]$ a.e. in $[u = 0]$. Moreover for each $\zeta \in H^1(\Omega_0)$, $\zeta = 0$ on $\partial\Omega_0 \setminus \widehat{BB_1}$, we have

$$\begin{aligned}\int_{\Omega_0} \big(b(X)\nabla u + \chi e_x\big)\nabla\zeta &= \int_{\Omega_0} \big(-b(X)\nabla\psi + (1-\gamma)e_x\big)\nabla\zeta \\ &= -\int_{\Omega_0} \big(b(X)\nabla\psi + \gamma e_x\big)\nabla\zeta + \int_{\Omega_0} \zeta_x = \int_{\widehat{BB_1}} \zeta\nu_x.\end{aligned}$$

Let $x_0 \in (x_A, x_B)$ and $y_1, y_2 \in (y_B, h)$ such that $y_1 < y_2 < h$ and $Z = ((x_0, +\infty) \times (y_1, y_2)) \cap \Omega \subset \Omega_0$. Let $\eta \in \mathscr{D}(y_1, y_2)$ and $\zeta = (x - x_0)^+\eta$. Then for $\epsilon > 0$ and $H_\epsilon(s) = \min(\frac{s^+}{\epsilon}, 1)$, we have $(H_\epsilon(u) - 1)\zeta \in H^1(\Omega_0)$ and $(H_\epsilon(u) - 1)\zeta = 0$ on $\partial\Omega_0 \setminus \widehat{BB_1}$. It follows that

$$\int_Z \big(b(X)\nabla u + \chi e_x\big)\nabla((H_\epsilon(u) - 1)\zeta) = \int_{\partial Z \cap \widehat{BB_1}} (H_\epsilon(u) - 1)\zeta\nu_x$$

which we can rewrite as

$$\begin{aligned}\int_Z (1-\chi)e_x.\nabla\zeta = &\int_Z (1 - H_\epsilon(u))b(X)\nabla u.\nabla\zeta - \int_Z H'_\epsilon(u)\zeta b(X)\nabla u.\nabla u \\ &+ \int_Z e_x.\nabla\zeta - \int_Z \chi e_x.\nabla(H_\epsilon(u)\zeta) + \int_{\partial Z \cap \widehat{BB_1}} (H_\epsilon(u) - 1)\zeta\nu_x.\end{aligned}$$

Since $H'_\epsilon(t) \geqslant 0$, $b(X)$ is positive, and $\chi\nabla(H_\epsilon(u)\zeta) = \nabla(H_\epsilon(u)\zeta)$ a.e. in Z, we get

$$\int_Z (1-\chi)e_x.\nabla\zeta \leqslant \int_Z (1 - H_\epsilon(u))b(X)\nabla u.\nabla\zeta.$$

Letting $\epsilon \to 0$, we obtain $\int_Z (1-\chi)\eta(y) = 0$. So $\chi = 1$ a.e. in Z. This holds for all such sub-domains Z of Ω_0. Hence $\chi = 1$ a.e. in Ω_0 and $\gamma = 0$ a.e. in Ω_0.

Now, we deduce from (2.1) that $div(b(X)\nabla\psi) = 0$ in Ω_0. Moreover, since $\psi \leqslant Q$ in Ω_0, we obtain by the strong maximum principle that either $\psi < Q$ in Ω_0 or $\psi \equiv Q$ in Ω_0. The last situation cannot happen since $\psi \in C^0(\Omega_0 \cup Int\,\widehat{AB})$ and $\psi = 0$ on $\widehat{AB}$. □

We end this section with a series of properties of the set $[\psi = Q]$.

Proposition 2.7 *Let* $B_1^* = (a(h), h)$, $E \in Int(T\widehat{B_1^*})$ *and* $l_E = \Omega \cap [y = y_E]$.

$$If \qquad \psi = Q \quad on \quad l_E, \qquad then \qquad \psi \equiv Q \quad in \quad \Omega_E = \Omega \cap [y > y_E].$$

https://doi.org/10.1017/S0956792505006418 Published online by Cambridge University Press

Proof Note that $\zeta = (Q-\psi)\chi(\Omega_E)$ is a test function for (P) since $\zeta \in H^1(\Omega)$ and $\zeta = 0$ on $\widehat{AB}$. Given that $\gamma = 0$ a.e. in $[\psi < Q]$ and $\widetilde{\gamma} = 0$ a.e. in $\widehat{TA} \cap [\psi < Q]$, we have

$$\int_{\Omega_E} b(X)\nabla(Q-\psi).\nabla(Q-\psi) = -\int_{\widehat{B_1T}} \zeta v_x \leqslant 0.$$

It follows that $Q-\psi$ is constant in Ω_E. But since $\psi = Q$ on l_E, we obtain $\psi = Q$ in Ω_E. □

Proposition 2.8 *Assume that* $E \in Int(\widehat{TA})$, $\psi(E) = Q$, *and that the interior sphere condition is satisfied at the point* E. *Then there exists a sequence* $E_n \in \Omega$ *such that* $E_n \longrightarrow E$ *and* $\psi(E_n) = Q$.

Proof We argue by contradiction and assume that there exists $\epsilon > 0$ such that $\psi < Q$ in $B_{2\epsilon}(E) \cap \Omega$. Let $\Omega_\epsilon = B_\epsilon(E) \cap \Omega$. Then we obtain, by taking into account the fact that $\gamma = 0$ a.e. in Ω_ϵ and $\widetilde{\gamma} v_x \leqslant 0$ on $\widehat{TA}$, that

$$\int_{\Omega_\epsilon} b(X)\nabla\psi.\nabla\zeta = \int_{\widehat{TA}} \widetilde{\gamma}\zeta v_x \leqslant 0 \quad \forall \zeta \in H^1(B_\epsilon(E)),\ \zeta \geqslant 0,\ \zeta = 0 \quad \text{on } \partial\Omega_\epsilon \cap \Omega. \tag{2.7}$$

Now we consider v defined by

$$\begin{cases} v \in H^1(\Omega_\epsilon), \quad v = \psi \quad \text{on } \partial\Omega_\epsilon \cap \Omega \quad \text{and} \\ \displaystyle\int_{\Omega_\epsilon} b(X)\nabla v.\nabla\zeta = 0 \qquad \forall \zeta \in H^1(\Omega_\epsilon), \quad \zeta = 0 \quad \text{on } \partial\Omega_\epsilon \cap \Omega. \end{cases} \tag{2.8}$$

Since $(v-Q)^+$ is a test function for (2.8), one easily verifies that $v \leqslant Q$ in Ω_ϵ. Now by taking $\zeta = (\psi - v)^+$ as a test function in (2.7) and (2.8) and subtracting the two quantities from each other, we obtain

$$\int_{\Omega_\epsilon} b(X)\nabla(\psi - v).\nabla(\psi - v)^+ \leqslant 0,$$

from which we deduce that $(\psi - v)^+$ is constant in Ω_ϵ. But $(\psi - v)^+ = 0$ on $\partial\Omega_\epsilon \cap \Omega$ leads to $\psi \leqslant v$ in Ω_ϵ. So $v(E) = Q$ since $Q = \psi(E) \leqslant v(E) \leqslant Q$. Hence v takes its maximum at the point E. By the strong maximum principle, we have either $v \equiv Q$ or $v < Q$ in Ω_ϵ. The first assumption is impossible, since it leads to $\psi = Q$ on $\partial\Omega_\epsilon \cap \Omega$ which is in contradiction with $\psi < Q$ in $\Omega_{2\epsilon} \supset \partial\Omega_\epsilon \cap \Omega$. Thus $v < Q$ in Ω_ϵ and we get a contradiction with $\psi(E) = Q$ and $b(E)\nabla v.\nu = 0$. □

Proposition 2.9 *Assume that the interior sphere condition is satisfied at each point of* $\widehat{BB_1} \cup \widehat{B_1^*A}$. *Then we have*

$$\psi < Q \quad \text{on} \quad \widehat{BB_1} \cup \widehat{B_1^*A} \qquad \text{and} \qquad \widetilde{\gamma} = 0 \quad \text{a.e. in} \quad \widehat{B_1^*A}\,.$$

https://doi.org/10.1017/S0956792505006418 Published online by Cambridge University Press

Proof First note that by Theorem 2.1 and Proposition 2.8, we have $\psi < Q$ on $\overset{\frown}{B_1^* A}$, which leads to $\widetilde{\gamma} = 0$ a.e. in $\overset{\frown}{B_1^* A}$.

Next let $\zeta \in \mathscr{D}(\Omega_0 \cup \overset{\frown}{BB_1})$. Then ζ is a test function for (P) and we have by taking into account Theorem 2.1 and the fact that $\zeta = 0$ on $\overset{\frown}{B_1 T} \cup \overset{\frown}{TA}$

$$\int_{\Omega_0} b(X)\nabla\psi.\nabla\zeta \;=\; 0.$$

It follows that $\psi < Q$, $div(b(X)\nabla\psi) = 0$ in Ω_0, and $b(X)\nabla\psi.\nu = 0$ on $\overset{\frown}{BB_1}$. Hence ψ cannot achieve its maximum value on $\overset{\frown}{BB_1}$ and therefore $\psi < Q$ on $\overset{\frown}{BB_1}$. □

Proposition 2.10 *Assume that* $E \in Int(\overset{\frown}{TB_1^*})$ *and that the interior sphere condition is satisfied at the point* E.

$$If \qquad \psi(E) = Q \qquad then \qquad \psi \equiv Q \quad in \quad \Omega \cap [y > y_E].$$

Proof By Proposition 2.8, there exists a sequence $E_n \in \Omega$ such that $E_n \longrightarrow E$ and $\psi(E_n) = Q$. By Proposition 2.6, it follows that $\psi(x, y_{E_n}) = Q$ if $x \geqslant x_{E_n}$. Letting $n \to \infty$, we get $\psi(x, y_E) = Q$ for $x \geqslant x_E$, which means that $\psi = Q$ on l_E. We deduce then from Proposition 2.7 that $\psi = Q$ in $\Omega \cap [y > y_E]$. □

3 Study of the free boundary

This section is devoted to the study of the free boundary which is, by Theorem 2.1, located above the level $y = h$. Throughout the section we denote a solution of (P) by $(\psi, \gamma, \widetilde{\gamma})$.

Proposition 2.6 allows us to define the function $\phi : (h, y_T) \longrightarrow \mathbb{R}$ by

$$\phi(y) = \begin{cases} \sup\{x \in (a(y), b(y)) \;/ \psi(x, y) < Q\} & \text{if this set is not empty} \\ a(y) & \text{otherwise.} \end{cases}$$

Then we have the following.

Theorem 3.1 *ϕ is continuous on (h, y_T).*

Proof Let $\Omega_1 = \Omega \cap [y > h]$, $u = Q - \psi$, $\chi = 1 - \gamma$. Arguing as in the beginning of the proof of Theorem 2.1, one can verify that

$$(\widehat{P}) \begin{cases} \displaystyle\int_{\Omega_1} (b(X)\nabla u + \chi e_x)\nabla\zeta = 0 \qquad \forall \zeta \in H^1(\Omega_1), \quad \zeta = 0 \quad \text{on } \partial\Omega_1 \setminus \overset{\frown}{B_1 T} \\ \\ \qquad u \geqslant 0 \quad \text{in } \Omega_1, \quad \chi \in H(u) \text{ a.e. in } \Omega_1. \end{cases}$$

Problem $(\widehat{P})$ belongs to a class of problems studied in [11] where the continuity of the free boundary was established in a more general framework. □

https://doi.org/10.1017/S0956792505006418 Published online by Cambridge University Press

As a consequence of Theorem 3.1, we obtain the following proposition.

Proposition 3.1 $\gamma = \chi([\psi = Q]) \qquad a.e.\ in \quad \Omega.$

Proof Indeed, we have $\Omega = \Omega_0 \cup \Omega_1 \cup [y = h]$ and we know from Theorem 2.1 and the definition of ϕ, that $\gamma = 0$ a.e. in $\Omega_0 \cup (\Omega_1 \cap [x < \phi(y)])$.

Now let $(x_0, y_0) \in [x > \phi(y)]$. By continuity, there exists $B_r(x_0, y_0) \subset [x > \phi(y)]$ such that $\psi = Q$ in B_r. This leads to $\psi = Q$ in $Z = B_r \cup (((x_0, +\infty) \times (y_0 - r, y_0 + r)) \cap \Omega)$. From (2.1), we deduce that $\gamma = \gamma(y)$ a.e. in Z. Now if we take $\zeta \in H^1(Z)$ in $(P)i)$ such that $\zeta = 0$ on $\partial Z \cap \Omega$, we obtain

$$\int_Z \gamma(y)\zeta_x = \int_{\widehat{B_1 T}} \gamma(y)\zeta \nu_x = \int_{\widehat{B_1 T}} \zeta \nu_x \qquad \Rightarrow \qquad \gamma = 1 \quad \text{a.e. in } Z.$$

Hence $\gamma = 1$ a.e. in $[x > \phi(y)]$. Now, since the sets $[y = h]$ and $[x = \phi(y)]$ are of Lebesgue measure zero, we get the result. □

Proposition 3.2 *We have* $\psi < Q \qquad on \quad [y = h] \cap \Omega.$

Proof Assume that there exists $x_0 \in (x_{B_1^*}, x_{B_1})$ such that $\psi(x_0, h) = Q$. Then $\psi(x, h) = Q$ for all $x \geqslant x_0$. We claim that there exists a sequence of points $(x_n, y_n)_n$ converging to (x_0, h) and satisfying

$$\forall n \geqslant 1 \quad x_n \geqslant x_0, \quad y_n > h \qquad \text{and} \qquad \psi(x_n, y_n) = Q.$$

Indeed, otherwise we would have $\psi < Q$ above the segment $(x_0, x_{B_1}) \times \{h\}$. Then if we consider a small ball $B \subset \Omega$ centred at a point of this segment, we get $\gamma = 0$ a.e. in B since we have also $\psi < Q$ in $[y < h]$. So $div(b(X)\nabla\psi) = 0$ in B and we get a contradiction to the maximum principle.

Using Proposition 2.6, we have $\psi(x, y_n) = Q$ for all $x \geqslant x_n$.

Arguing as in the proof of the continuity of ϕ [11], one can prove that for some $\eta > 0$ small enough and some n large enough, we have

$$\psi \equiv Q \qquad \text{in} \quad \big((x_n + \eta, +\infty) \times (h, y_n)\big) \cap \Omega.$$

Consider now a small ball B centred at a point of the segment $(x_n + \eta, x_{B_1}) \times \{h\}$ and such that $B \cap [y > h] \subset ((x_n + \eta, +\infty) \times (h, y_n)) \cap \Omega$. Since $\gamma = 0$ a.e. in $B \cap [y < h]$ and $\gamma = 1$ a.e. in $B \cap [y > h]$, we have $\gamma_x = 0$ in $\mathscr{D}'(B)$. Using (2.1), we obtain $div(b(X)\nabla\psi) = 0$ in $\mathscr{D}'(B)$ and we get a contradiction to the maximum principle. Hence we have $\psi < Q$ on $[y = h] \cap \Omega$. □

Proposition 3.3 *Assume that the interior sphere condition is satisfied at each point of* $\widehat{TB_1^*}$. *Then the set* $[\psi < Q]$ *is connected by arcs.*

Proof Since $\Omega \cap [y \leqslant h] \subset [\psi < Q]$ is connected, it is enough to show that each point of $[\psi < Q] \cap [y > h]$ can be connected to a point of the line segment $l_{B_1} = [y = h] \cap \Omega$.

https://doi.org/10.1017/S0956792505006418 Published online by Cambridge University Press

Let $(x_0, y_0) \in [\psi < Q] \cap [y > h]$. By Proposition 2.6, we have $\psi(x, y_0) < Q \quad \forall x \leqslant x_0$. Moreover by Proposition 2.10, one has $\psi(a_0(y), y) < Q \quad \forall y \in [h, y_0]$. By continuity of ψ, there exists $\epsilon > 0$ small enough such that $\psi(x, y) < Q \quad \forall y \in [h, y_0]$ and $\forall x \in [a(y), a(y)+\epsilon]$. Therefore (x_0, y_0) is connected by the piecewise smooth curve $[a_0(y_0) + \epsilon, x_0] \times \{y_0\} \cup \{(a_0(y) + \epsilon, y) : \quad y \in [h, y_0]\}$ to the point $(a_0(h) + \epsilon, h)$ of l_{B_1}. □

Proposition 3.4 $\lim_{y \to h^+} \phi(y) = x_{B_1} = b_0(h)$.

Proof Let $\epsilon > 0$. Since $\psi(x_{B_1} - \epsilon, h) < Q$, there exists, by continuity of ψ, $\eta_1 > 0$ such that $\psi(x_{B_1} - \epsilon, y) < Q \quad \forall y \in [h, h + \eta_1]$.

By definition of ϕ, we obtain $\phi(y) > x_{B_1} - \epsilon \quad \forall y \in [h, h + \eta_1]$.

On the other hand, we get from the continuity of $b_0(y)$ at h, the existence of $\eta_2 > 0$ such that $b_0(y) < x_{B_1} + \epsilon \quad \forall y \in [h, h + \eta_2]$. Therefore

$$x_{B_1} - \epsilon < \phi(y) \leqslant b_0(y) < x_{B_1} + \epsilon \qquad \forall y \in [h, h + \min(\eta_1, \eta_2)].$$

□

4 Uniqueness of the solution

In this section we establish the uniqueness of the solution by techniques similar to those in Carrillo *et al.* [9] and Lyaghfouri [14].

Theorem 4.1 *The solution of the problem* (P) *is unique.*

To prove Theorem 4.1, we need two lemmas.

Lemma 4.1 *Let* $(\psi_1, \gamma_1, \widetilde{\gamma}_1)$, $(\psi_2, \gamma_2, \widetilde{\gamma}_2)$ *be two solutions of* (P). *Then we have*

$$\mathscr{T}(\zeta) = \int_\Omega (b(X)\nabla(\psi_1 - \psi_2)^+ + (\gamma_1 - \gamma_2)^+ e_x).\nabla\zeta \leqslant \int_J \zeta(\phi_2(y), y)dy$$

$\forall \zeta \in H^1(\Omega) \cap C(\overline{\Omega}), \quad \zeta \geqslant 0 \quad$ *with* $\quad J = \{\, y \in (h, y_T) : \quad \phi_1(y) < \phi_2(y) \,\}$.

Proof Let $\zeta \in H^1(\Omega) \cap C(\overline{\Omega})$, $\zeta \geqslant 0$. For $\epsilon > 0$ small enough, take $\xi = \min(\frac{(\psi_1 - \psi_2)^+}{\epsilon}, \zeta)$ as a test function in $(P)i)$ for $(\psi_1, \gamma_1, \widetilde{\gamma}_1)$ and for $(\psi_2, \gamma_2, \widetilde{\gamma}_2)$. By subtracting the obtained equations, we get, since $(\psi_1 - \psi_2)(\gamma_1 - \gamma_2) \geqslant 0$ a.e. in Ω, $(\psi_1 - \psi_2)(\widetilde{\gamma}_1 - \widetilde{\gamma}_2) \geqslant 0$ and $v_x \leqslant 0$ a.e. on $\widehat{TA}$,

$$\begin{aligned}
&\int_{\Omega \cap [(\psi_1 - \psi_2)^+ \geqslant \epsilon\zeta]} b(X)\nabla(\psi_1 - \psi_2)^+.\nabla\zeta + \int_\Omega (\gamma_1 - \gamma_2)^+ e_x.\nabla\xi \\
&= -\frac{1}{\epsilon} \int_{\Omega \cap [(\psi_1 - \psi_2)^+ < \epsilon\zeta]} b(X)\nabla(\psi_1 - \psi_2)^+.\nabla(\psi_1 - \psi_2)^+ + \int_{\widehat{TA}} (\widetilde{\gamma}_1 - \widetilde{\gamma}_2)^+ \xi v_x \leqslant 0.
\end{aligned}$$

https://doi.org/10.1017/S0956792505006418 Published online by Cambridge University Press

Since $\xi = \zeta - (\zeta - \frac{(\psi_1-\psi_2)^+}{\epsilon})^+$, we obtain

$$\int_{\Omega\cap[(\psi_1-\psi_2)^+\geqslant\epsilon\zeta]} b(X)\nabla(\psi_1-\psi_2)^+.\nabla\zeta + \int_\Omega(\gamma_1-\gamma_2)^+e_x.\nabla\zeta \leqslant \int_\Omega(\gamma_1-\gamma_2)^+\Big(\zeta-\frac{(\psi_1-\psi_2)^+}{\epsilon}\Big)_x^+ = I_\epsilon. \tag{4.1}$$

Using Proposition 3.1, we have

$$I_\epsilon = \int_{[x<\phi_2(y)]\cap[x\geqslant\phi_1(y)]}\Big(\zeta-\frac{(Q-\psi_2)^+}{\epsilon}\Big)_x^+ = \int_J\Big(\int_{\phi_1(y)}^{\phi_2(y)}\Big(\zeta-\frac{(Q-\psi_2)^+}{\epsilon}\Big)_x^+dx\Big)dy \leqslant \int_J\zeta(\phi_2(y),y)dy.$$

The lemma follows by letting $\epsilon\to 0$ in (4.1). □

Lemma 4.2 *Let $(\psi_1,\gamma_1,\widetilde{\gamma}_1)$, $(\psi_2,\gamma_2,\widetilde{\gamma}_2)$ be two solutions of (P). Then we have*

$$\mathscr{T}(\zeta)=0 \qquad \forall\zeta\in H^1(\Omega).$$

Proof Let $\zeta\in C^1(\bar{\Omega})$, $\zeta\geqslant 0$. For $\delta>0$ small enough, we consider

$$\alpha_\delta(X)=\Big(1-\frac{d(X,U)}{\delta}\Big)^+, \qquad \text{with } U=[\psi_1<Q]=[x<\phi_1(y)].$$

We have $\mathscr{T}(\zeta)=\mathscr{T}(\alpha_\delta\zeta)+\mathscr{T}((1-\alpha_\delta)\zeta)$ and by Lemma 4.1

$$\mathscr{T}(\alpha_\delta\zeta)\leqslant\int_J\alpha_\delta(\phi_2(y),y)\zeta(\phi_2(y),y)dy.$$

But for $y\in J$, $\phi_2(y)>\phi_1(y)$ and then $(\phi_2(y),y)\notin\overline{U}=\overline{[x<\phi_1(y)]}$ $\forall y\in J$. So $d((\phi_2(y),y),U)>0$ and $\lim_{\delta\to0}\alpha_\delta(\phi_2(y),y)=0$. We deduce that $\limsup_{\delta\to0}\mathscr{T}(\alpha_\delta\zeta)\leqslant 0$.

Next, we have for $\psi_0=\min(\psi_1,\psi_2)$, $\gamma_0=\min(\gamma_1,\gamma_2)$,

$$\begin{aligned}\mathscr{T}((1-\alpha_\delta)\zeta) &= \int_\Omega(b(X)\nabla(\psi_1-\psi_0)+(\gamma_1-\gamma_0)e_x).\nabla((1-\alpha_\delta)\zeta)\\ &= \int_{\Omega\setminus\overline{U}}(b(X)\nabla\psi_1+\gamma_1e_x).\nabla((1-\alpha_\delta)\zeta)\\ &\quad -\int_{\Omega\setminus\overline{U}}(b(X)\nabla\psi_0+\gamma_0e_x).\nabla((1-\alpha_\delta)\zeta).\end{aligned}$$

https://doi.org/10.1017/S0956792505006418 Published online by Cambridge University Press

Note that on $[x > \phi_1(y)] = \Omega \setminus \overline{U}$, we have $\psi_1 = Q$, $\gamma_1 = 1$, $\psi_0 = \psi_2$ and $\gamma_0 = \gamma_2$. Moreover, $(1-\alpha_\delta)\zeta$ is a test function for (P) since $\widehat{AB} \subset \overline{U}$. Then we obtain

$$\begin{aligned}\mathscr{T}((1-\alpha_\delta)\zeta) &= \int_\Omega ((1-\alpha_\delta)\zeta)_x - \int_\Omega (b(X)\nabla\psi_2 + \gamma_2 e_x).\nabla((1-\alpha_\delta)\zeta) \\ &= \int_{\partial\Omega} (1-\alpha_\delta)\zeta \nu_x - \int_{\widehat{TA}} \widetilde{\gamma}_2 (1-\alpha_\delta)\zeta \nu_x - \int_{\widehat{B_1T}} (1-\alpha_\delta)\zeta \nu_x \\ &= \int_{\widehat{TA}} (1-\widetilde{\gamma}_2)(1-\alpha_\delta)\zeta \nu_x \leqslant 0\end{aligned}$$

since $\alpha_\delta = 1$ on $\widehat{AB} \cup \widehat{BB_1}$. Hence $\mathscr{T}(\zeta) \leqslant 0 \quad \forall \zeta \in C^1(\bar{\Omega}), \quad \zeta \geqslant 0$.

Now let $\zeta \in C^1(\bar{\Omega})$ and $M = \sup_{\bar{\Omega}} |\zeta|$. Note that we have $M \pm \zeta \in C^1(\bar{\Omega})$ and $M \pm \zeta \geqslant 0$ in Ω. It follows that $\mathscr{T}(M \pm \zeta) \leqslant 0$ and therefore $\mathscr{T}(\zeta) = 0 \quad \forall \zeta \in C^1(\bar{\Omega})$. Finally, since $C^1(\bar{\Omega})$ is dense in $H^1(\Omega)$, Lemma 4.2 follows. □

Proof of Theorem 4.1 Starting from Lemma 4.2, we establish as in Lyaghfouri [14] that

$$\psi_1 = \psi_2 \qquad \text{in} \qquad [\psi_1 < Q] \cap [\psi_2 < Q] = [\psi_1 < Q] = [\psi_2 < Q].$$

Therefore, $\psi_1 = \psi_2$ in Ω and by Proposition 3.1, we obtain $\gamma_1 = \gamma_2$ a.e. in Ω. Finally, let $\zeta \in H^1(\Omega)$, $\zeta = 0$ on $\partial\Omega \setminus \widehat{TA}$. Writing the equations satisfied by the two solutions and subtracting them from each other, we obtain $\int_{\widehat{TA}} (\widetilde{\gamma}_1 - \widetilde{\gamma}_2)\zeta \nu_x = 0$ which leads to $\widetilde{\gamma}_1 = \widetilde{\gamma}_2$ a.e. in $\widehat{TA}$.

5 Conclusion

To the best of our knowledge, no regularity result for the free boundary has been obtained from the classical formulation in the absence of monotonicity of the permeability. This is the main motivation for this work in which we established, without any monotonicity of the permeability, that the free boundary is a curve of a continuous function $x = \phi(y)$. Finally, it would be quite interesting to study the equivalence of the two formulations.

Acknowledgements

We would like to thank KFUPM for its continuing support.

References

[1] Alt, H. W. (1979) Strömungen durch inhomogene poröse Medien mit freiem Rand. *J. für die Reine und Angewandte Mathematik*, **305**, 89–115.

[2] Alt, H. W. (1977) The fluid flow Through Porous Media. Regularity of the free Surface. *Manuscripta Math.* **21**, 255–272.

[3] Alt, H. W., Caffarelli, L. A. & Friedman, A. (1984) The dam problem with two fluids. *Comm. Pure & Appl. Math.* **37**, 601–645.

https://doi.org/10.1017/S0956792505006418 Published online by Cambridge University Press

[4] Alt, H. W. & Gilardi, G. (1981) The Behavior of the Free Boundary for the Dam Problem. *Ann. Sc. Norm. Sup. Pisa*, **(IV)**(IX), 571–626.
[5] Baiocchi, C. (1972) Su un problema di frontiera libera connesso a questioni di idraulica. *Ann. Mat. Pura Appl.* **92**, 107–127.
[6] Benci, V. (1974) On a filtration through a porous medium. *Ann. Mat. Pura Appl.* **100**, 191–209.
[7] Brézis, H., Kinderlehrer, D. & Stampacchia, G. (1978) Sur une nouvelle formulation du problème de l'écoulement à travers une digue. *C. R. Acd. Sci Paris Serie A* **287**, 711–714.
[8] Caffarelli, L. A. & Friedman, A. (1978) The Dam problem with two layers. *Arch. Ration. Mech. Anal.* **68**, 125–154.
[9] Carrillo, J. & Chipot, M. (1982) On the Dam Problem. *J. Diff. Eq.* **45**, 234–271.
[10] Challal, M. & Lyaghfour, L. (2004) A Filtration Problem through a Heterogeneous Porous Medium. *Interfaces & Free Boundaries*, **6**, 55–79.
[11] Challal, M. & Lyaghfouri, A. (2005) On the Continuity of the Free Boundary in Problems of type $div(a(x)\nabla u) = -(h\chi)_{x_1}$. *Nonlinear Anal.: Theory, Methods & Applica.* **62**(2), 283–300.
[12] Friedman, A. & Huang, S.-Y. (1987) The inhomogeneous dam problem with discountinuous permeability. *Ann. Scu. Norm. Sup. Pisa, Ser. IV,* **14**, 49–77.
[13] Gilbarg, D. & Trudinger, N. S. (1983) *Elliptic Partial Differential Equations of Second Order*. Springer-Verlag.
[14] Lyaghfouri, A. (1996) The inhomogeneous dam problem with linear Darcy's law and Dirichlet boundary conditions. *Mathematical Models & Methods in Appl. Sci.* **8**(6), 1051–1077.
[15] Stavre, R. & Vernescu, B. (1985) Incompressible fluid flow through a nonhomogeneous and anistropic dam. *Nonlinear Anal.: Theory, Methods & Applic*. **9**, 799–810.

https://doi.org/10.1017/S0956792505006418 Published online by Cambridge University Press