Glasgow Mathematical Journal (2026), **68**, pp. 337–344
doi:10.1017/S0017089525100785

Glasgow Mathematical Journal

RESEARCH ARTICLE

A note on twisted moments of Dirichlet *L*-functions

Julio Andrade[1] and Joseph Macmillan[2]

[1]Department of Mathematics, University of Exeter, Exeter EX4 4QF, UK
[2]Department of Mathematics, Swansea University, Swansea SA1 8EN, UK
Corresponding author: Joseph Macmillan; Email: j.b.macmillan@swansea.ac.uk

Received: 2 July 2024; **Revised:** 23 June 2025; **Accepted:** 23 June 2025; **First published online:** 24 September 2025

Keywords: twisted moments; Dirichlet *L*-functions; function fields

2020 Mathematics Subject Classification: *Primary* - 11M38; *Secondary* - 11M06, 11T06, 11T55

Abstract
In this paper, we establish an asymptotic formula for the twisted second moments of Dirichlet *L*-functions with one twist when averaged over all primitive Dirichlet characters of modulus R, where R is a monic polynomial in $\mathbb{F}_q[T]$.

1. Introduction

It is well known that the study of moments of the Riemann zeta-function and *L*-functions is an important topic in analytic number theory. It can be even argued that a great part of research in analytic number theory in the last century has been guided and motivated by this topic.

Applications of moments of *L*-functions appear more notably in the Lindelöf hypothesis, but also when studying proportions of zeros satisfying the Riemann hypothesis and nonvanishing at the central point of families of *L*-functions. For some of these applications, it is important to understand not only the moments of *L*-functions but also what is known as *twisted moments*.

Let χ be a Dirichlet character modulo p, where p is a prime number. The problem is then to obtain a formula for

$$\mathcal{S}(p,h) := \sideset{}{^*}\sum_{\chi \,(\mathrm{mod}\, p)} \left| L\left(\frac{1}{2}, \chi\right)\right|^2 \chi(h), \tag{1.1}$$

where h is a fixed prime number and the $*$ indicates a summation over all primitive Dirichlet characters modulo p. With this notation, Conrey [3, Theorem 10] proved the following.

Theorem 1.1 (Conrey [3]). *For primes p, h with $2 \le h < p$, we have that*

$$\mathcal{S}(p,h) = \frac{p^{1/2}}{h^{1/2}}\mathcal{S}(h,-p) + \frac{p}{h^{1/2}}\left(\log\frac{p}{h} + \gamma - \log(8\pi)\right) + \zeta\left(\frac{1}{2}\right)^2 p^{1/2} + O\left(h + \log p + \frac{p^{1/2}}{h^{1/2}}\log p\right),$$

where γ is Euler's constant and ζ is the Riemann zeta-function.

© The Author(s), 2025. Published by Cambridge University Press on behalf of Glasgow Mathematical Journal Trust. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (https://creativecommons.org/licenses/by/4.0/), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

https://doi.org/10.1017/S0017089525100785 Published online by Cambridge University Press

In [8], Young extended Conrey's result as follows.

Theorem 1.2 (Young [8]). *For primes p, h with $h < p^{1-\varepsilon}$, we have that*

$$\frac{p^{1/2}}{\varphi(p)}\mathcal{S}(p,h) - \frac{h^{1/2}}{\varphi(h)}\mathcal{S}(h,-p) = \frac{p^{1/2}}{h^{1/2}}\left(\log\frac{p}{h} + \gamma - \log(8\pi)\right) + \zeta\left(\frac{1}{2}\right)^2\left(1 - 2\frac{p^{1/2}}{\varphi(p)}(1-p^{-1/2}) + 2\frac{h^{1/2}}{\varphi(h)}(1-h^{-1/2})\right) + \mathcal{E}(p,h),$$

where $\varphi(p)$ is Euler's totient function and

$$\mathcal{E}(p,h) \ll hp^{-1-\varepsilon} + h^{-C},$$

for all fixed $\varepsilon, C > 0$.

Advancing the study of twisted moments of Dirichlet L-functions, Bettin [2] showed that the error term $\mathcal{E}(p,h)$ can be extended to a continuous function with respect to the real topology. In his work, Bettin extended the known reciprocity results for twisted moments by establishing an exact formula with shifts.

More recently, there have been some interesting developments on the study of twisted second moments of Dirichlet L-functions over rational function fields. Let q be the power of an odd prime number and $\mathbb{A} = \mathbb{F}_q[T]$ the polynomials with coefficients in the finite field $\mathbb{F}_q$. In this setting, Djanković [4] proved the following.

Theorem 1.3 (Djanković [4]). *Let P, H be irreducible polynomials in $\mathbb{F}_q[T]$ and*

$$\mathcal{S}(P,H) := \sum_{\chi \,(\mathrm{mod}\, P)}^{*} \left|L\left(\frac{1}{2},\chi\right)\right|^2 \chi(H).$$

If $H \neq P$ and $\deg(H) \leq \deg(P)$, then

$$\frac{|P|^{1/2}}{\phi(P)}\mathcal{S}(P,H) - \frac{|H|^{1/2}}{\phi(H)}\mathcal{S}(H,-P) = \frac{|P|^{1/2}}{|H|^{1/2}}\left(\deg(P) - \deg(H) - \zeta_{\mathbb{A}}\left(\frac{1}{2}\right)^2\right) + \zeta_{\mathbb{A}}\left(\frac{1}{2}\right)^2\left(1 - 2\frac{|P|^{1/2}}{\phi(P)}\left(1-|P|^{-1/2}\right) + 2\frac{|H|^{1/2}}{\phi(H)}\left(1-|H|^{-1/2}\right)\right),$$

where $L(s,\chi)$ is the Dirichlet L-function in function fields associated with the Dirichlet character χ modulo P, with $\zeta_{\mathbb{A}}(s)$ being the zeta-function for $\mathbb{F}_q[T]$, $\phi(P)$ is the Euler's totient function for polynomials and $|P| = q^{\deg(P)}$ denotes the norm of a polynomial P in $\mathbb{F}_q[T]$.

The aim of this note is to extend the above result of Djanković. In their work, they only consider Dirichlet characters modulo a monic irreducible polynomial, that is, they only prove results for prime moduli. In this note, we establish results for general moduli. In particular, we prove the following.

Theorem 1.4. *Let H and R be monic polynomials in $\mathbb{F}_q[T]$ with $(H,R) = 1$ and $\deg(H) \ll (\frac{1}{3} - \epsilon)\deg(R)$, then*

$$\frac{1}{\phi^*(R)}\sum_{\chi \,(\mathrm{mod}\, R)}^{*}\left|L\left(\frac{1}{2},\chi\right)\right|^2\chi(H) = \frac{1}{|H|^{\frac{1}{2}}}\frac{\phi(R)}{|R|}(\deg(R) - \deg(H)) + O\left(\frac{1}{|H|^{\frac{1}{2}}}\log\omega(R)\right). \tag{1.2}$$

where $\omega(R)$ is the number of distinct prime factors of R, $\phi^*(R)$ denotes the number of primitive Dirichlet characters modulo R and $*$ indicates a summation over all primitive Dirichlet characters modulo R.

https://doi.org/10.1017/S0017089525100785 Published online by Cambridge University Press

2. A short overview of Dirichlet *L*-functions over function fields

In this section, we give a short overview of Dirichlet L-functions in function fields, with most of these facts stated in [6]. Let $\mathbb{F}_q$ denote a finite field with q elements, where q is a power of an odd prime and $\mathbb{A} = \mathbb{F}_q[T]$ be its polynomial ring. Furthermore, we denote by $\mathbb{A}^+$, $\mathbb{A}_n^+$ and $\mathbb{A}_{\leq n}^+$ the set of all monic polynomials in $\mathbb{A}$, the set of all monic polynomials in $\mathbb{A}$ of degree n and the set of all monic polynomials of degree at most n in $\mathbb{A}$, respectively. For $f \in \mathbb{A}$, the norm of f, $|f|$, is defined to be equal to $q^{\deg(f)}$, and $\phi(f)$, $\mu(f)$ and $\omega(f)$ denote the Euler-Totient function for $\mathbb{A}$, the Möbius function for $\mathbb{A}$ and the number of distinct prime factors of f.

For $\Re(s) > 1$, the zeta-function for $\mathbb{A}$ is defined as

$$\zeta_{\mathbb{A}}(s) = \sum_{f \in \mathbb{A}^+} \frac{1}{|f|^s} = \prod_P \left(1 - \frac{1}{|P|^s}\right)^{-1} \tag{2.1}$$

where the product is over all monic irreducible polynomials in $\mathbb{A}$. Since there are q^n monic polynomials of degree n in $\mathbb{A}$, then

$$\zeta_{\mathbb{A}}(s) = \frac{1}{1 - q^{1-s}}.$$

Definition 2.1. *Let $R \in \mathbb{A}^+$. Then, a Dirichlet character modulo R is defined to be a function $\chi : \mathbb{A} \to \mathbb{C}$, which satisfies the following properties:*

(1) $\chi(AB) = \chi(A)\chi(B), \quad \forall A, B \in \mathbb{A},$
(2) $\chi(A + BR) = \chi(A), \quad \forall A, B \in \mathbb{A},$
(3) $\chi(A) \neq 0 \iff (A, R) = 1.$

A Dirichlet character χ is said to be even if $\chi(a) = 1$ for all $a \in \mathbb{F}_q^*$. Otherwise, we say that it is odd.

Definition 2.2. *Let $R \in \mathbb{A}^+$, $S|R$ and χ be a character of modulus R. We say that S is an induced modulus of χ if there exists a character χ_1 of modulus S such that*

$$\chi(A) = \begin{cases} \chi_1(A) & \text{if } (A, R) = 1, \\ 0 & \text{otherwise.} \end{cases}$$

We say χ is primitive if there is no induced modulus of strictly smaller norm than R. Otherwise, χ is said to be non-primitive. Let $\phi^*(R)$ denote the number of primitive characters of modulus R.

Definition 2.3. *Let χ be a Dirichlet character modulo R. Then, the Dirichlet L-function corresponding to χ is defined by*

$$L(s, \chi) := \sum_{f \in \mathbb{A}^+} \frac{\chi(f)}{|f|^s} \tag{2.2}$$

which converges absolutely for $\Re(s) > 1$.

To finish this section, we will state some results about multiplicative functions in function fields, which will be used throughout this paper. Taking Euler products, we see that for all $s \in \mathbb{C}$ and all $R \in \mathbb{A}$, we have

$$\sum_{E|R} \frac{\mu(E)}{|E|^s} = \prod_{P|R} \left(1 - \frac{1}{|P|^s}\right) \tag{2.3}$$

https://doi.org/10.1017/S0017089525100785 Published online by Cambridge University Press

and differentiating (2.3), we see that for all $s \in \mathbb{C}\backslash\{0\}$, we have

$$\sum_{E|R} \frac{\mu(E)\deg(E)}{|E|^s} = -\left(\prod_{P|R} 1 - \frac{1}{|P|^s}\right)\left(\sum_{P|R} \frac{\deg(P)}{|P|^s - 1}\right). \tag{2.4}$$

Lemma 2.4 [1, Lemma 4.5]. *Let $R \in \mathbb{A}^+$. We have that*

$$\sum_{P|R} \frac{\deg(P)}{|P| - 1} \ll \log \omega(R). \tag{2.5}$$

Lemma 2.5 [7, Lemma A.2.3]. *For $\deg(R) > 1$, we have*

$$\omega(R) \ll \frac{\log_q |R|}{\log_q \log_q |R|} \tag{2.6}$$

where the implied constant is independent of q.

Lemma 2.6. *We have*

$$2^{\omega(R)} = \sum_{E|R} |\mu(E)|. \tag{2.7}$$

Also, for any $\epsilon > 0$, we have

$$2^{\omega(R)} \ll_\epsilon |R|^\epsilon. \tag{2.8}$$

Lemma 2.7 [7, Lemma A.2.4]. *For $deg(R) > q$, we have*

$$\phi(R) \gg \frac{|R|}{\log_q \log_q |R|}. \tag{2.9}$$

Lemma 2.8 [7, Lemma A.2.5]. *For $deg(R) > q$, we have*

$$\phi^*(R) \gg \frac{\phi(R)}{\log_q \log_q |R|}. \tag{2.10}$$

Lemma 2.9 [1, Lemma 3.7]. *Let $R \in \mathbb{A}^+$ and $A, B \in \mathbb{A}$. Then,*

$$\sum_{\chi (\mathrm{mod}\, R)}^{*} \chi(A)\bar{\chi}(B) = \begin{cases} \sum_{\substack{EF=R \\ F|(A-B)}} \mu(E)\phi(F) & \text{if } (AB, R) = 1, \\ 0 & \text{otherwise} \end{cases}.$$

As a Corollary, we have the following result.

Corollary 2.10 [1, Corollary 3.8]. *For all $R \in \mathbb{A}^+$, we have that*

$$\phi^*(R) = \sum_{EF=R} \mu(E)\phi(F). \tag{2.11}$$

3. Preliminary lemmas

In this section, we state and prove results that will be needed to prove Theorem 1.4. We start by stating the approximate function equation for $\left|L\left(\frac{1}{2}, \chi\right)\right|^2$.

https://doi.org/10.1017/S0017089525100785 Published online by Cambridge University Press

Lemma 3.1 [5, Lemma 2.5]. *Let χ be a primitive Dirichlet character of modulus R. Then, we have*

$$\left|L\left(\frac{1}{2},\chi\right)\right|^2 = 2\sum_{\substack{A,B\in\mathbb{A}^+\\ \deg(AB)<\deg(R)}}\frac{\chi(A)\bar{\chi}(B)}{|AB|^{\frac{1}{2}}} + O\left(|R|^{-\frac{1}{2}+\epsilon}\right). \tag{3.1}$$

The next lemma will be used to obtain the main term of Theorem 1.4.

Lemma 3.2 [1, Lemma 4.12]. *Let R be a monic polynomial in $\mathbb{F}_q[T]$ and let x be a positive integer. Then,*

$$\sum_{\substack{A\in\mathbb{A}^+_{\leq x}\\ (A,R)=1}}\frac{1}{|A|} = \begin{cases}\dfrac{\phi(R)}{|R|}x + O(\log\omega(R)) & \text{if } x\geq\deg(R)\\[2ex] \dfrac{\phi(R)}{|R|}x + O(\log\omega(R)) + O\left(\dfrac{2^{\omega(R)}x}{q^x}\right) & \text{if } x<\deg(R)\end{cases}.$$

The following lemmas will be used to create a suitable bound for the error term of Theorem 1.4.

Lemma 3.3. *Let F, H and R be fixed monic polynomials in $\mathbb{F}_q[T]$ where $F|R$ and let $z<\deg(R)$. Then,*

$$\sum_{\substack{A,B\in\mathbb{A}^+\\ \deg(AB)=z\\ AH\equiv B(\mathrm{mod}\ F)\\ AH\neq B\\ (ABH,R)=1}}\frac{1}{|AB|^{\frac{1}{2}}} \ll \frac{q^{\frac{z}{2}}(z+1)|H|}{|F|}. \tag{3.2}$$

Proof. We consider three cases, $\deg(AH)>\deg(B)$, $\deg(AH)<\deg(B)$ and $\deg(AH)=\deg(B)$, where $AH\neq B$.

If we first consider the case $\deg(AH)>\deg(B)$ and suppose that $\deg(A)=i$, then since $AH\equiv B(\mathrm{mod}\ F)$ and $AH\neq B$ we have that $AH=LF+B$ for some $L\in\mathbb{A}^+$ with $\deg(L)=i+\deg(H)-\deg(F)$ and $\deg(B)=z-\deg(A)=z-i$. Thus, combining the above, we have

$$\begin{aligned}\sum_{\substack{A,B\in\mathbb{A}^+\\ \deg(AB)=z\\ \deg(AH)>\deg(B)\\ AH\equiv B(\mathrm{mod}\ F)\\ AH\neq B\\ (ABH,R)=1}}\frac{1}{|AB|^{\frac{1}{2}}} &\leq q^{-\frac{z}{2}}\sum_{i=0}^{z}\sum_{\substack{L\in\mathbb{A}^+\\ \deg(L)=i+\deg(H)-\deg(F)}}\sum_{\substack{B\in\mathbb{A}^+\\ \deg(B)=z-i}}1\\ &= q^{\frac{z}{2}}\sum_{i=0}^{z}q^{-i}\sum_{\substack{L\in\mathbb{A}^+\\ \deg(L)=i+\deg(H)-\deg(F)}}1 = \frac{q^{\frac{z}{2}}|H|}{|F|}\sum_{i=0}^{z}1 = \frac{q^{\frac{z}{2}}(z+1)|H|}{|F|}.\end{aligned} \tag{3.3}$$

Similarly, considering the case $\deg(AH)<\deg(B)$ and using similar arguments seen previously, we have

$$\sum_{\substack{A,B\in\mathbb{A}^+\\ \deg(AB)=z\\ \deg(B)>\deg(AH)\\ AH\equiv B(\mathrm{mod}\ F)\\ AH\neq B\\ (ABH,R)=1}}\frac{1}{|AB|^{\frac{1}{2}}} \leq \frac{q^{\frac{z}{2}}(z+1)}{|F|}. \tag{3.4}$$

Finally, if we consider the case where $\deg(AH)=\deg(B)=i$, then $2i=\deg(ABH)=z+\deg(H)$ and so $\deg(B)=i=\frac{z+\deg(H)}{2}$. Furthermore, since $AH\equiv B(\mathrm{mod\ F})$ and $AH\neq B$, then $AH=LF+B$ where $L\in$

https://doi.org/10.1017/S0017089525100785 Published online by Cambridge University Press

$\mathbb{A}$ with $\deg(L) < i - \deg(F) = \frac{z+\deg(H)}{2} - \deg(F)$. Thus, combining the above, we have

$$\sum_{\substack{A,B\in\mathbb{A}^+\\ \deg(AB)=z\\ \deg(AH)=\deg(B)\\ AH\equiv B(\mathrm{mod}\ F)\\ AH\neq B\\ (ABH,R)=1}} \frac{1}{|AB|^{\frac{1}{2}}} \leq q^{-\frac{z}{2}} \sum_{\substack{B\in\mathbb{A}^+\\ \deg(B)=\frac{z+\deg(H)}{2}}} \sum_{\substack{L\in\mathbb{A}\\ \deg(L)<\frac{z+\deg(H)}{2}-\deg(F)}} 1$$

$$\ll \frac{|H|^{\frac{1}{2}}}{|F|} \sum_{\substack{B\in\mathbb{A}^+\\ \deg(B)=\frac{z+\deg(H)}{2}}} 1 = \frac{q^{\frac{z}{2}}|H|}{|F|}. \tag{3.5}$$

Combining all the cases proves the result. □

Lemma 3.4. *For all $R \in \mathbb{A}^+$ and $\epsilon > 0$, we have*

$$\frac{2^{\omega(R)}|R|^{\frac{1}{2}}\deg(R)}{\phi^*(R)} \ll_\epsilon |R|^{\epsilon-\frac{1}{2}}. \tag{3.6}$$

Proof. For $\deg(R) \leq q$, we know, by [7, (A.2.3)], that $\frac{\phi^*(R)}{|R|} \gg 1$. Thus, for $\deg(R) \leq q$, we have

$$\frac{2^{\omega(R)}|R|^{\frac{1}{2}}\deg(R)}{\phi^*(R)} \ll \frac{2^{\omega(R)}\deg(R)}{|R|^{\frac{1}{2}}} \ll \frac{2^{\omega(R)}}{|R|^{\frac{1}{2}-\epsilon}}.$$

From Lemma 2.6, we know that $2^{\omega(R)} \ll |R|^\epsilon$, thus (3.6) holds for $\deg(R) \leq q$.

For $\deg(R) > q$, we know by Lemmas 2.7 and 2.8 that

$$\phi^*(R) \gg \frac{\phi(R)}{\log_q\log_q|R|} \gg \frac{|R|}{(\log_q\log_q|R|)^2}.$$

Thus, if $\deg(R) > q$, then

$$\frac{2^{\omega(R)}|R|^{\frac{1}{2}}\deg(R)}{\phi^*(R)} \ll \frac{2^{\omega(R)}\deg(R)(\log_q\log_q|R|)^2}{|R|^{\frac{1}{2}}} \ll_\epsilon \frac{2^{\omega(R)}}{|R|^{\frac{1}{2}-\epsilon}}.$$

Finally, from Lemma 2.6, we know that $2^{\omega(R)} \ll |R|^\epsilon$, then (3.6) holds for $\deg(R) > q$ and thus completes the proof. □

4. Proof of Theorem 1.4

In this section, we use results stated previously to prove Theorem 1.4.

Proof of Theorem 1.4. Using the approximate function equation Lemma 3.1, we have

$$\frac{1}{\phi^*(R)}\sideset{}{^*}\sum_{\chi(\mathrm{mod}\ R)}\left|L\left(\frac{1}{2},\chi\right)\right|^2\chi(H) = \frac{2}{\phi^*(R)}\sideset{}{^*}\sum_{\chi(\mathrm{mod}\ R)}\sum_{\substack{A,B\in\mathbb{A}^+\\ \deg(AB)<\deg(R)}}\frac{\chi(A)\bar{\chi}(B)\chi(H)}{|AB|^{\frac{1}{2}}} + O\left(|R|^{-\frac{1}{2}+\epsilon}\right). \tag{4.1}$$

Using the orthogonality relation Lemma 2.9, we have

$$\frac{2}{\phi^*(R)}\sideset{}{^*}\sum_{\chi(\mathrm{mod}\ R)}\sum_{\substack{A,B\in\mathbb{A}^+\\ \deg(AB)<\deg(R)}}\frac{\chi(A)\bar{\chi}(B)\chi(H)}{|AB|^{\frac{1}{2}}} = \frac{2}{\phi^*(R)}\sum_{EF=R}\mu(E)\phi(F)\sum_{\substack{A,B\in\mathbb{A}^+\\ \deg(AB)<\deg(R)\\ AH\equiv B(\mathrm{mod}\ F)\\ (ABH,R)=1}}\frac{1}{|AB|^{\frac{1}{2}}}. \tag{4.2}$$

https://doi.org/10.1017/S0017089525100785 Published online by Cambridge University Press

For the second sum on the right-hand side of (4.2), we will consider the contribution of the diagonal, $AH = B$, and the off-diagonal, $AH \neq B$, terms separately. Thus, we write

$$\frac{2}{\phi^*(R)} \sum_{EF=R} \mu(E)\phi(F) \sum_{\substack{A,B\in\mathbb{A}^+ \\ \deg(AB)<\deg(R) \\ AH\equiv B(\mathrm{mod}\ F) \\ (ABH,R)=1}} \frac{1}{|AB|^{\frac{1}{2}}} = \frac{2}{\phi^*(R)} \sum_{EF=R} \mu(E)\phi(F) \sum_{\substack{A,B\in\mathbb{A}^+ \\ \deg(AB)<\deg(R) \\ AH\equiv B(\mathrm{mod}\ F) \\ AH=B \\ (ABH,R)=1}} \frac{1}{|AB|^{\frac{1}{2}}} + \frac{2}{\phi^*(R)} \sum_{EF=R} \mu(E)\phi(F) \sum_{\substack{A,B\in\mathbb{A}^+ \\ \deg(AB)<\deg(R) \\ AH\equiv B(\mathrm{mod}\ F) \\ AH\neq B \\ (ABH,R)=1}} \frac{1}{|AB|^{\frac{1}{2}}}.$$

Considering the contribution of the diagonal, $AH = B$, the double sum over all $A, B \in \mathbb{A}^+$ with $\deg(AB) < \deg(R)$, $AH = B$ and $(ABH, R) = 1$ becomes a single sum over all $A \in \mathbb{A}^+$ with $\deg(A) < \frac{1}{2}(\deg(R) - \deg(H))$ and $(AH, R) = 1$. Therefore, using the arguments stated above and Corollary 2.10, we have

$$\frac{2}{\phi^*(R)} \sum_{EF=R} \mu(E)\phi(F) \sum_{\substack{A,B\in\mathbb{A}^+ \\ \deg(AB)<\deg(R) \\ AH=B \\ (ABH,R)=1}} \frac{1}{|AB|^{\frac{1}{2}}} = \frac{2}{|H|^{\frac{1}{2}}} \sum_{\substack{A\in\mathbb{A}^+ \\ \deg(A)<\frac{\deg(R)-\deg(H)}{2} \\ (AH,R)=1}} \frac{1}{|A|}. \tag{4.3}$$

Since the condition $(AH, R) = 1$ holds if and only if $(A, R) = 1$ and $(H, R) = 1$, then since we have already assumed that $(H, R) = 1$, then, in the sum, we only need to consider the condition $(A, R) = 1$. Thus,

$$\sum_{\substack{A\in\mathbb{A}^+ \\ \deg(A)<\frac{\deg(R)-\deg(H)}{2} \\ (AH,R)=1}} \frac{1}{|A|} = \sum_{\substack{A\in\mathbb{A}^+ \\ \deg(A)<\frac{\deg(R)-\deg(H)}{2} \\ (A,R)=1}} \frac{1}{|A|}. \tag{4.4}$$

Using Lemma 3.2 with $x = \frac{\deg(R)-\deg(H)}{2} - 1$, we have

$$\frac{2}{|H|^{\frac{1}{2}}} \sum_{\substack{A\in\mathbb{A}^+ \\ \deg(A)<\frac{\deg(R)-\deg(H)}{2} \\ (AH,R)=1}} \frac{1}{|A|} = \frac{1}{|H|^{\frac{1}{2}}} \frac{\phi(R)}{|R|}(\deg(R) - \deg(H)) + O\left(\frac{1}{|H|^{\frac{1}{2}}} \log \omega(R)\right). \tag{4.5}$$

For the contribution of the off-diagonal terms, we use Lemma 3.3 to give

$$\sum_{\substack{A,B\in\mathbb{A}^+ \\ \deg(AB)<\deg(R) \\ AH\equiv B(\mathrm{mod}\ F) \\ AH\neq B \\ (ABH,R)=1}} \frac{1}{|AB|^{\frac{1}{2}}} = \sum_{z=0}^{\deg(R)-1} \sum_{\substack{A,B\in\mathbb{A}^+ \\ \deg(AB)=z \\ AH\equiv B(\mathrm{mod}\ F) \\ AH\neq B \\ (ABH,R)=1}} \frac{1}{|AB|^{\frac{1}{2}}} \ll \sum_{z=0}^{\deg(R)-1} \frac{|H|q^{\frac{z}{2}}(z+1)}{|F|} \ll \frac{|H||R|^{\frac{1}{2}}\deg(R)}{|F|}. \tag{4.6}$$

Thus using (4.6), we have

$$\frac{2}{\phi^*(R)} \sum_{EF=R} \mu(E)\phi(F) \sum_{\substack{A,B\in\mathbb{A}^+ \\ \deg(AB)<\deg(R) \\ AH\equiv B(\mathrm{mod}\ F) \\ AH\neq B \\ (ABH,R)=1}} \frac{1}{|AB|^{\frac{1}{2}}} \ll \frac{|H||R|^{\frac{1}{2}}\deg(R)}{\phi^*(R)} \sum_{EF=R} |\mu(E)| \frac{\phi(F)}{|F|}. \tag{4.7}$$

https://doi.org/10.1017/S0017089525100785 Published online by Cambridge University Press

Combining (4.7) with Lemmas 2.6, 3.4 and the fact that $\frac{\phi(R)}{|R|} \leq 1$, we have

$$\frac{2}{\phi^*(R)} \sum_{EF=R} \mu(E)\phi(F) \sum_{\substack{A,B\in\mathbb{A}^+ \\ \deg(AB)<\deg(R) \\ AH\equiv B(\mathrm{mod}\ F) \\ AH\neq B \\ (ABH,R)=1}} \frac{1}{|AB|^{\frac{1}{2}}} \ll \frac{2^{\omega(R)}|H||R|^{\frac{1}{2}}\deg(R)}{\phi^*(R)} \ll |H||R|^{\epsilon-\frac{1}{2}}. \tag{4.8}$$

Since H is a monic polynomial in $\mathbb{F}_q[T]$ with $(H,R)=1$ and $\deg(H) \ll (\frac{1}{3}-\epsilon)\deg(R)$, then $|H||R|^{-\frac{1}{2}+\epsilon} \ll \frac{1}{|H|^{\frac{1}{2}}} \log\omega(R)$ as $deg(R)\to\infty$. Combining the above completes the proof of Theorem 1.4. □

Acknowledgements. The authors are grateful to the Leverhulme Trust (RPG-2017-320) for the support through the research project grant "Moments of L-functions in Function Fields and Random Matrix Theory". The second author is also grateful to Swansea University (MAD1000-100) for sponsoring a research visit to the University of Exeter. The authors would also like to thank Prof. Will Sawin for discussions related to the previous version of this manuscript. They would also like to thank the anonymous referee for a careful reading of the paper and for providing useful suggestions and comments that helped to improve the presentation of the paper and also have given us some ideas for future projects.

References

[1] J. C. Andrade and M. Yiasemides, The fourth power mean of Dirichlet *L-functions in* $\mathbb{F}_q[T]$, *Rev. Mat. Comput.* **34**(1) (2021), 239–296.
[2] S. Bettin, On the reciprocity law for the twisted second moment of Dirichlet L-functions, *Trans. Amer. Math. Soc.* **368**(10) (2016), 6887–6914.
[3] J. B. Conrey, *The mean-square of Dirichlet L-functions* (2007). arXiv:0708.2699.
[4] G. Djanković, The reciprocity law for the twisted second moment of Dirichlet L-functions over rational function fields, *Bull. Aust. Math. Soc.* **98**(3) (2018), 382–388.
[5] P. Gao and L. Zhao, Moments of Dirichlet L-functions to a fixed modulus over function fields, *Int. J. Number Theory* **20**(6) (2024), 1493–1513.
[6] M. Rosen, Number theory in function fields, *Graduate texts in mathematics*, vol. **210** (Springer-Verlag, New York, 2002).
[7] M. Yiasemides, *Dirichlet L-functions and their derivatives in function fields* (PhD thesis, University of Exeter, 2020).
[8] M. P. Young, The reciprocity law for the twisted second moment of Dirichlet L-functions, *Forum Math.* **23**(6) (2011), 1323–1337.

https://doi.org/10.1017/S0017089525100785 Published online by Cambridge University Press