Combinatorics, Probability and Computing (2011) 20, 815–835. © Cambridge University Press 2011
doi:10.1017/S0963548311000435

Computing the Partition Function for Perfect Matchings in a Hypergraph

ALEXANDER BARVINOK[1†] and ALEX SAMORODNITSKY[2‡]

[1]Department of Mathematics, University of Michigan, Ann Arbor, MI 48109-1043, USA
(e-mail: barvinok@umich.edu)
[2]Department of Computer Science, Hebrew University of Jerusalem, Givat Ram Campus, 91904, Israel
(e-mail: salex@cs.huji.ac.il)

Received 30 November 2010; revised 4 September 2011; first published online 17 October 2011

Given non-negative weights w_S on the k-subsets S of a km-element set V, we consider the sum of the products $w_{S_1}\cdots w_{S_m}$ over all partitions $V = S_1 \cup \cdots \cup S_m$ into pairwise disjoint k-subsets S_i. When the weights w_S are positive and within a constant factor of each other, fixed in advance, we present a simple polynomial-time algorithm to approximate the sum within a polynomial in m factor. In the process, we obtain higher-dimensional versions of the van der Waerden and Bregman–Minc bounds for permanents. We also discuss applications to counting of perfect and nearly perfect matchings in hypergraphs.

1. Introduction and main results

Let us fix an integer $k > 1$. A collection $H \subset \binom{V}{k}$ of k-subsets of a finite set V is called a *k-uniform hypergraph* with *vertex* set V, while sets $S \in H$ are called *edges* of H. In particular, a uniform 2-hypergraph is an ordinary undirected graph on V without loops or multiple edges. A set $\{S_1, \ldots, S_m\}$ of pairwise vertex-disjoint edges of H such that $V = S_1 \cup \cdots \cup S_m$ is called a *perfect matching* of hypergraph H. More generally, a *matching* of *size* n is a collection of n pairwise disjoint edges of H.

If a perfect matching exists then the number $|V|$ of vertices of V is divisible by k, so we have $|V| = km$ for some integer m. The hypergraph consisting of all k-subsets of V is called the *complete k-uniform hypergraph* with vertex set V. We denote it by $\binom{V}{k}$. A hypergraph is called a *complete k-partite hypergraph* if the set V of vertices is a union $V = V_1 \cup \cdots \cup V_k$ of pairwise disjoint sets V_i, called *parts*, such that $|V_1| = \cdots = |V_k| = m$ and the edges of the hypergraph are the subsets $S \subset V$ containing exactly one vertex

† Research partially supported by NSF grant DMS 0856640 and by a US–Israel BSF grant 2006377.
‡ Research partially supported by ISF grant 039-7165 and by a US–Israel BSF grant 2006377.

https://doi.org/10.1017/S0963548311000435 Published online by Cambridge University Press

in each part: $|S \cap V_1| = \cdots = |S \cap V_k| = 1$. We denote such a hypergraph by $V_1 \times \cdots \times V_k$.

We introduce the main object of the paper.

1.1. Partition function

Let H be a k-uniform hypergraph with the set V of vertices such that $|V| = km$ for some positive integer m. Suppose that to every edge $S \in H$ a non-negative real number w_S is assigned. Such an assignment $W = \{w_S\}$ we call a *weight* on H. We say that W is *positive* if $w_S > 0$ for all $S \in H$. The polynomial

$$P_H(W) = \sum w_{S_1} \cdots w_{S_m},$$

where the sum is taken over all perfect matchings $\{S_1, \ldots, S_m\}$ of H, is called the *partition function* of perfect matchings in hypergraph H. Sometimes we just write $P(W)$ if the choice of the hypergraph H is clear from the context.

We note that we can obtain the partition function $P_H(W)$ of an arbitrary k-uniform hypergraph $H \subset \binom{V}{k}$ by specializing $w_S = 0$ for $S \notin H$ in the partition function of the complete k-uniform hypergraph $\binom{V}{k}$.

The partition function of $\binom{V}{2}$ with $|V| = 2m$ is known as the *hafnian* of the $2m \times 2m$ symmetric matrix $A = (a_{ij})$, where a_{ij} is the weight of the edge consisting of the ith and jth vertices of V (diagonal elements of A can be chosen arbitrarily); see, for example, Section 8.2 of [19]. If $V_1 \times V_2$ is a complete bipartite graph with $|V_1| = |V_2| = m$, then the corresponding partition function is the *permanent* of the $m \times m$ matrix $B = (b_{ij})$, where b_{ij} is the weight of the edge consisting of the ith vertex of V_1 and jth vertex of V_2. The partition function of the complete k-partite hypergraph gives rise to a version of the permanent of a k-dimensional tensor; see, for example, [7].

In this paper, we address the problem of computing or approximating $P_H(W)$ efficiently. First, we define certain classes of weights W.

1.2. Balanced and k-stochastic weights

We say that a positive weight $W = \{w_S\}$ on a k-uniform hypergraph is α-*balanced* for some $\alpha \geqslant 1$ if

$$\frac{w_{S_1}}{w_{S_2}} \leqslant \alpha \quad \text{for all} \quad S_1, S_2 \in H.$$

Note that an α-balanced weight is also β-balanced for any $\beta > \alpha$.

Weight $Z = \{z_S\}$ is called k-*stochastic* if

$$\sum_{\substack{S \in H \\ S \ni v}} z_S = 1 \quad \text{for all} \quad v \in V.$$

In words: for every vertex, the sum of the weights of the edges containing the vertex is 1.

Now we are ready to state our first main result.

Theorem 1.1. *Let us fix an integer $k > 1$ and a real $\alpha \geqslant 1$. Then there exists a real $\gamma = \gamma(k, \alpha) > 0$ such that if H is a complete k-uniform hypergraph or a complete k-partite*

https://doi.org/10.1017/S0963548311000435 Published online by Cambridge University Press

hypergraph with km vertices and Z is a k-stochastic α-balanced weight on H, then

$$m^{-\gamma}e^{-m(k-1)} \leqslant P_H(Z) \leqslant m^{\gamma}e^{-m(k-1)}$$

provided $m > 1$.

In other words, for fixed k and α, the value of the partition function for a k-stochastic α-balanced weight on a complete k-uniform hypergraph or a complete k-partite hypergraph can vary only within a polynomial in m range.

More precisely, we prove that under conditions of Theorem 1.1 and assuming, additionally, that $\alpha^{k+1} > 2$, we have

$$\epsilon_1 m^{-\gamma_1}e^{-m(k-1)} \leqslant P_H(Z) \leqslant \epsilon_2 m^{\gamma_2}e^{-m(k-1)},$$

where

$$\begin{aligned} &\gamma_1 = \alpha^{3(k+1)}(k^2+k)^2 + (k-1)^2 \quad \text{and} \quad \gamma_2 = \frac{k^2\alpha^{k+1}}{2}, \\ &\epsilon_1 = \alpha^{-(k+1)l}l^l\binom{kl}{k}^{1-l} \quad \text{and} \quad \epsilon_2 = \alpha^{(k+1)l}l^{l-kl+k} \\ &\qquad \text{for} \quad l = \lceil \alpha^{2(k+1)}k^2 \rceil + 1. \end{aligned} \tag{1.1}$$

1.3. Comparison with permanents

The van der Waerden conjecture on permanents proved by Falikman [10] and Egorychev [8] (see also [14] for important new developments) asserts that if $A = (a_{ij})$ is an $m \times m$ doubly stochastic matrix, that is, a non-negative matrix with all row and column sums equal to 1, then

$$\operatorname{per} A \geqslant \frac{m!}{m^m} = \sqrt{2\pi m}e^{-m}\left(1 + O\left(\frac{1}{m}\right)\right).$$

A conjecture by Minc proved by Bregman [3] (see also [21] for a simpler proof) asserts that if $B = (b_{ij})$ is an $m \times m$ matrix with $b_{ij} \in \{0, 1\}$ for all i, j, then

$$\operatorname{per} B \leqslant \prod_{i=1}^{m}(r_i!)^{1/r_i},$$

where r_i is the ith row sum of B. From this inequality one can deduce that if A is an $m \times m$ non-negative matrix with all row sums equal to 1 and all the entries not exceeding α/m for some $\alpha \geqslant 1$, then

$$\operatorname{per} A \leqslant m^{\gamma}e^{-m}$$

for some $\gamma = \gamma(\alpha) > 0$ and all $m > 1$ (one can choose any $\gamma > \alpha/2$ if m is sufficiently large); see [22]. Thus the van der Waerden and Bregman–Minc inequalities together imply that $\operatorname{per} A = e^{-m}m^{O(1)}$ for any $m \times m$ doubly stochastic matrix A whose entries are within a factor of $O(1)$ of each other. Theorem 1.1 presents an extension of this interesting fact to non-bipartite graphs for $k = 2$ and to hypergraphs for $k > 2$. A stronger statement that $\operatorname{per} A = e^{-m}m^{O(1)}$ for an $m \times m$ doubly stochastic matrix whose maximum entry is $O(m^{-1})$

https://doi.org/10.1017/S0963548311000435 Published online by Cambridge University Press

fails to extend to non-bipartite graphs for $k = 2$ or to k-partite hypergraphs for $k > 2$, as the following two examples readily show.

Let $k = 2$ and let H be a graph on a set V of $n = 4r + 2$ vertices, which consists of two vertex-disjoint copies of the complete graph on $2r + 1$ vertices. Let us define a weight $Z = \{z_S\}$ on $\binom{V}{2}$ by letting $z_S = (2r)^{-1}$ if S is an edge of H and $z_S = 0$ otherwise. Then Z is a 2-stochastic weight on $\binom{V}{2}$ and $P(Z) = 0$. That is, the hafnian of an $n \times n$ symmetric doubly stochastic matrix can be zero even when the maximum entry of the matrix is $O(n^{-1})$.

Let $k = 3$, let $m = 4r + 2$ and let us identify each set V_1, V_2 and V_3 with a copy of the integer interval $\{1, 2, \ldots, m\}$. Let us define a weight $Z = \{z_S\}$ on $V_1 \times V_2 \times V_3$ by letting $z_S = ((4r + 2)(2r + 1))^{-1}$ if $S = (a, b, c)$ with $a + b + c$ even and $z_S = 0$ otherwise. Then Z is a 3-stochastic weight on $V_1 \times V_2 \times V_3$ while $P(Z) = 0$, since the sum of all integers in V_1, V_2 and V_3 is odd. Hence the permanent of a 3-stochastic $m \times m \times m$ tensor can be zero even when the maximum entry of the tensor is $O(m^{-2})$. This example was constructed in a conversation with Jeff Kahn.

Summarizing, for general k-stochastic weights Z there is no *a priori* non-zero lower bound for the partition function. If, however, we require Z to be α-balanced for any fixed $\alpha \geqslant 1$, the lower bound jumps to within a polynomial in m factor of the upper bound.

We note that there are extensions of the Bregman–Minc bound to hafnians [1] and to higher-dimensional permanents [6]. Lower bounds appear to be harder to come by; see [9] for the recent proof of the Lovász–Plummer conjecture, which states that the number of perfect matchings in a bridgeless 3-regular graph is exponentially large in the number of vertices of the graph, and [12] and [2] for related developments.

If H is a complete k-uniform hypergraph or a complete k-partite hypergraph, one can estimate $P_H(W)$ for any balanced but not necessarily k-stochastic weight W using scaling.

1.4. Scaling

Let $W = \{w_S\}$ be a weight on the edges of a k-uniform hypergraph H with a vertex set V, where $|V| = km$. Let $\{\lambda_v > 0 : v \in V\}$ be reals. We say that a weight $Z = \{z_S\}$ on the hypergraph H is obtained from W by *scaling* if

$$z_S = \left(\prod_{v \in S} \lambda_v\right) w_S \quad \text{for all} \quad S \in H.$$

It is easy to see that

$$P_H(Z) = \left(\prod_{v \in V} \lambda_v\right) P_H(W).$$

It turns out that any positive weight W on a complete k-uniform hypergraph or a complete k-partite hypergraph can be scaled to a k-stochastic weight Z (see, for example, [11] and Section 3 below). We show that the k-stochastic scaling of an α-balanced weight is α^{k+1}-balanced and obtain the following result.

https://doi.org/10.1017/S0963548311000435 Published online by Cambridge University Press

Theorem 1.2. *Let us fix an integer $k > 1$ and a real $\alpha \geqslant 1$. Then there exists a real $\gamma = \gamma(k, \alpha) > 0$ such that the following holds.*

Let H be a complete k-uniform hypergraph or a complete k-partite hypergraph with km vertices, and let $W = \{w_S : S \in H\}$ be an α-balanced weight on H. Let us consider the function

$$f_W(X) = \sum_{S \in H} x_S \ln \frac{x_S}{w_S}$$

for a weight X on H. Let $\Omega_k(H)$ be the set of all k-stochastic weights on H and let

$$\zeta = \min_{X \in \Omega_k(H)} f_W(X).$$

Then

$$e^{-\zeta - m(k-1)} m^{-\gamma} \leqslant P_H(W) \leqslant e^{-\zeta - m(k-1)} m^{\gamma}.$$

More precisely, we prove that under conditions of Theorem 1.1 and assuming, additionally, that $\alpha^{k+1} > 2$, we have

$$\epsilon_1 m^{-\gamma_1} e^{-\zeta - m(k-1)} \leqslant P_H(W) \leqslant \epsilon_2 m^{\gamma_2} e^{-\zeta - m(k-1)},$$

where $\gamma_1, \gamma_2, \epsilon_1, \epsilon_2$ are defined by (1.1).

The set $\Omega_k(H)$ is naturally identified with a convex polytope in $\mathbb{R}^H$. Function f is strictly convex and hence the optimization problem of computing ζ can be solved efficiently (in polynomial time) by interior point methods; see [20]. Thus Theorem 1.2 allows us to estimate the partition function of an α-balanced weight (for any $\alpha \geqslant 1$, fixed in advance) within a polynomial in m factor.

1.5. A probabilistic interpretation

Let us fix $k > 1$ and let H be either a complete k-uniform hypergraph or a complete k-partite hypergraph with a set V of $|V| = km$ vertices. Let us fix $\alpha \geqslant 1$ and let $W = \{w_S : S \in H\}$ be an α-balanced weight on H. Let $|H|$ denote the number of edges of hypergraph H. Let us assume that

$$\sum_{S \in H} w_S = m,$$

in which case

$$\frac{m}{\alpha |H|} \leqslant w_S \leqslant \frac{\alpha m}{|H|} \quad \text{for all} \quad S \in H.$$

In particular, for all sufficiently large m we have $w_S < 1$ for all $S \in H$, so we can introduce independent random Bernoulli variables X_S indexed by the edges $S \in H$, where

$$\mathbb{P}(X_S = 1) = w_S \quad \text{and} \quad \mathbb{P}(X_S = 0) = 1 - w_S.$$

For each vertex $v \in V$ let us define a random variable

$$Y_v = \sum_{\substack{S \in H \\ S \ni v}} X_S.$$

https://doi.org/10.1017/S0963548311000435 Published online by Cambridge University Press

It is not hard to show that

$$P_H(W) = \exp\left\{m + O\left(\frac{1}{m^{k-2}}\right)\right\}\mathbb{P}(Y_v = 1 \text{ for all } v \in V).$$

For large m, the distribution of each random variable Y_v is approximately Poisson with

$$\mathbb{E}\, Y_v = \mu_v \quad \text{where} \quad \mu_v = \sum_{\substack{S \in H \\ S \ni v}} w_S,$$

so

$$\mathbb{P}(Y_v = 1) \approx \mu_v e^{-\mu_v}.$$

The probability of $Y_v = 1$ is maximized when $\mu_v = 1$, and when W is k-stochastic, the probabilities of $Y_v = 1$ are maximized simultaneously for all $v \in V$, so that

$$\mathbb{P}(Y_v = 1) \approx e^{-1} \quad \text{for all } v \in V.$$

Theorem 1.1 implies that in this case the events $Y_v = 1$ behave as if they were (almost) independent, so that

$$\mathbb{P}(Y_v = 1 \text{ for all } v \in V) \approx e^{-km}$$

up to a polynomial in m factor.

In Section 2, we discuss some combinatorial and algorithmic applications of Theorems 1.1 and 1.2. Namely, we present a simple polynomial-time algorithm to distinguish hypergraphs having sufficiently many perfect matchings from hypergraphs that do not have nearly perfect matchings. We also prove a lower bound for the number of nearly perfect matchings in regular hypergraphs.

In the rest of the paper we prove Theorems 1.1 and 1.2.

In Section 3, we review some results about scaling. The results are not new, but we nevertheless provide proofs for completeness. In Section 4, we prove two crucial lemmas about scaling of α-balanced weights. In Section 5 we complete the proofs of Theorems 1.1 and 1.2.

Scaling was used in [17] to efficiently estimate permanents of non-negative matrices.

1.6. Notation

As usual, for two functions f and g, where g is non-negative, we say that $f = O(g)$ if $|f| \leqslant \gamma g$ for some constant $\gamma > 0$. We will allow our constants γ to depend only on the dimension k of the hypergraph and the parameter $\alpha \geqslant 1$ in the definition of an α-balanced weight in Section 1.2.

2. Combinatorial applications

Let us fix an integer $k > 1$ and let H be a k-uniform hypergraph with km vertices. As is known [23], the problem of counting perfect matchings in H is #P-hard. For $k = 2$ there is a classical polynomial-time algorithm to check whether H has a perfect matching (see [18]) and a fully polynomial randomized approximation scheme is known for counting

https://doi.org/10.1017/S0963548311000435 Published online by Cambridge University Press

perfect matchings if H is bipartite [15]. For $k > 2$, finding if there is a perfect matching in H is an NP-complete problem even when H is k-partite [16].

Theorem 1.2 allows us to distinguish in polynomial time between hypergraphs that have sufficiently many perfect matchings and hypergraphs that do not have nearly perfect matchings.

In this section, we let

$$\Phi_k(m) = \frac{(km)!}{(k!)^m m!}$$

be the number of perfect matchings in a complete k-uniform hypergraph with km vertices.

2.1. Testing hypergraphs

Let us fix integer $k > 1$ and positive real $\delta \leqslant 1$ and $\beta < 1$.

We consider the following algorithm.

Input: A k-uniform hypergraph H, defined by the list of its edges, with a set V of km vertices.

Output: At least one of the following two (not mutually exclusive) conclusions.

(a) The hypergraph H contains a matching with at least βm edges.
(b) The hypergraph H contains at most $\delta^m \Phi_k(m)$ perfect matchings.

Algorithm: Let

$$\epsilon = \frac{1}{2}\delta^{1/(1-\beta)}.$$

Let us define a weight $W = \{w_S\}$ on the complete k-uniform hypergraph $\binom{V}{k}$ as follows:

$$w_S = \begin{cases} 1 & \text{if } S \in H, \\ \epsilon & \text{if } S \notin H. \end{cases} \tag{2.1}$$

The weight W is ϵ^{-1}-balanced and we apply the algorithm of Theorem 1.2 to compute in polynomial in m time a number η such that

$$\eta \cdot m^{-\gamma} \leqslant P(W) \leqslant \eta \cdot m^{\gamma}$$

for some $\gamma = \gamma(\delta, \beta) > 0$.

If $m = 1$ or if

$$\frac{m}{\ln m} \leqslant \frac{2\gamma}{(1-\beta)\ln 2}, \tag{2.2}$$

we check by direct enumeration whether (a) or (b) hold. Since k, β and δ are fixed in advance, this requires only a constant time.

If (2.2) does not hold, we output conclusion (a) if $\eta \cdot m^{\gamma} > \delta^m \Phi_k(m)$ and conclusion (b) if $\eta \cdot m^{\gamma} \leqslant \delta^m \Phi_k(m)$.

It is not hard to see that the algorithm is indeed correct. If $\eta \cdot m^{\gamma} \leqslant \delta^m \Phi_k(m)$ then $P(W) \leqslant \delta^m \Phi_k(m)$ and H necessarily contains no more than $\delta^m \Phi_k(m)$ perfect matchings.

https://doi.org/10.1017/S0963548311000435 Published online by Cambridge University Press

If $\eta \cdot m^{\gamma} > \delta^m \Phi_k(m)$ then, assuming that (2.2) does not hold, we conclude that

$$P(W) \geqslant \frac{\delta^m}{m^{2\gamma}} \Phi_k(m) > \frac{\delta^m}{2^{(1-\beta)m}} \Phi_k(m) = \epsilon^{(1-\beta)m} \Phi_k(m),$$

from which it follows that H contains a matching with no fewer than βm edges.

In particular, confronted with two hypergraphs on km vertices, one of which contains more than $\delta^m \Phi_k(m)$ perfect matchings and the other with no matchings of size βm or bigger, the algorithm will be able to decide which is which. It will necessarily output (a) in the former case and (b) in the latter.

Definition. A k-uniform hypergraph H is called *d-regular* if every vertex of H is contained in exactly d edges of H.

For example, a complete k-uniform hypergraph with km vertices is d-regular for $d = \binom{km-1}{k-1}$. The existence of a perfect or nearly perfect matching in d-regular hypergraphs was extensively studied; see, for example, [24] and references therein. As a corollary of Theorem 1.1, we obtain the following estimate for the number of nearly perfect matchings in a regular hypergraph (see also [4] for some related estimates).

Theorem 2.1. *Let us fix an integer $k > 1$ and $0 < \alpha, \beta < 1$. Then, there exists a positive integer $m_0 = m_0(k, \alpha, \beta)$ such that the following holds.*

Let H be a k-uniform d-regular hypergraph with km vertices where

$$d \geqslant \alpha \binom{km-1}{k-1} \quad \text{and} \quad m \geqslant m_0.$$

Then, for every positive integer $s \leqslant \beta m$ the hypergraph H contains at least

$$\alpha^m \frac{\Phi_k(m)}{\Phi_k(m-s)}$$

matchings of size s.

Proof. All implied constants in the 'O' notation below may depend on k, α and β only. Let V be the set of vertices of a k-uniform d-regular hypergraph, $|V| = km$. Let us choose $0 < \epsilon < 1$ such that

$$\epsilon^{1-\beta} < \alpha + \epsilon(1-\alpha), \tag{2.3}$$

and let us define a weight $W = \{w_S\}$ on the complete k-uniform hypergraph $\binom{V}{k}$ by (2.1). Then

$$\sum_{\substack{S \in \binom{V}{k} \\ S \ni v}} w_S = (1-\epsilon)d + \epsilon \binom{km-1}{k-1} \quad \text{for all} \quad v \in V.$$

https://doi.org/10.1017/S0963548311000435 Published online by Cambridge University Press

It follows from Theorem 1.1 and scaling that

$$P(W) \geqslant \left((1-\epsilon)d + \epsilon\binom{km-1}{k-1}\right)^m e^{-m(k-1)}\frac{1}{m^{O(1)}}$$
$$\geqslant (\alpha+\epsilon(1-\alpha))^m\binom{km-1}{k-1}^m e^{-m(k-1)}\frac{1}{m^{O(1)}}. \tag{2.4}$$

We note that

$$\frac{\binom{km-1}{k-1}^m}{\Phi_k(m)} = \frac{\big((km-1)!\big)^m(k!)^m m!}{\big((k-1)!\big)^m\big((km-k)!\big)^m(km)!} = \frac{\big((km-1)!\big)^m k^m m!}{\big((km-k)!\big)^m(km)!}$$
$$= \frac{\big((km)!\big)^m k^m m!}{(km)^m\big((km-k)!\big)^m(km)!} = \frac{\big((km)!\big)^m m!}{\big((km-k)!\big)^m(km)!m^m}$$
$$= \left(\frac{(km)(km-1)\cdots(km-k+1)}{(km)^k}\right)^m \cdot \frac{(km)^{km}m!}{(km)!m^m}. \tag{2.5}$$

Since $\ln(1-x) \geqslant -2x$ for $0 \leqslant x \leqslant 0.5$, we conclude that

$$\left(\frac{(km)(km-1)\cdots(km-k+1)}{(km)^k}\right)^m = \exp\left\{m\sum_{i=1}^{k-1}\ln\left(1-\frac{i}{km}\right)\right\} \geqslant e^{-k+1}.$$

Using Stirling's formula, we conclude from (2.5) and (2.4) that

$$P(W) \geqslant (\alpha+\epsilon(1-\alpha))^m\Phi_k(m)\frac{1}{m^{O(1)}}. \tag{2.6}$$

If a perfect matching in $\binom{V}{k}$ contains fewer than s edges of H then the contribution of the corresponding term to $P(W)$ is less than ϵ^{m-s}. Since every matching in H of size s can be appended to a perfect matching in $\binom{V}{k}$ in $\Phi_k(m-s)$ ways, we conclude that the number of matchings in H of size s is at least

$$\frac{P(W)-\epsilon^{m-s}\Phi_k(m)}{\Phi_k(m-s)} \geqslant \frac{P(W)-\epsilon^{(1-\beta)m}\Phi_k(m)}{\Phi_k(m-s)}.$$

The proof now follows from (2.6) and (2.3). □

3. General results on scaling

In this section, we summarize some results on scaling which we need for the proofs of Theorems 1.1 and 1.2.

Theorem 3.1. *Let H be a k-uniform hypergraph with a set V of $|V| = km$ vertices and let $\Omega_k(H)$ be the set of all k-stochastic weights on H. Suppose that the set $\Omega_k(H)$ has a non-empty relative interior, that is, it contains a positive weight $Y \in \Omega_k(H)$.*

For a positive weight $W = \{w_S : S \in H\}$ on H, let us define a function

$$f_W : \Omega_k(H) \longrightarrow \mathbb{R}$$

by

$$f_W(X) = \sum_{S\in H} x_S \ln\frac{x_S}{w_S} \quad \text{for} \quad X \in \Omega_k(H),\ X = \{x_S : S \in H\}.$$

https://doi.org/10.1017/S0963548311000435 Published online by Cambridge University Press

Then function f_W attains its minimum on $\Omega_k(H)$ at a unique weight $Z = \{z_S : S \in H\}$. We have $z_S > 0$ for all $S \in H$ and there exist real $\lambda_v > 0 : v \in V$ such that

$$z_S = \left(\prod_{v \in S} \lambda_v\right) w_S \quad \text{for all} \quad S \in H. \tag{3.1}$$

We have

$$f_W(Z) = \sum_{v \in V} \ln \lambda_v.$$

Furthermore, if $\lambda_v > 0 : v \in V$ are reals such that weight Z defined by (3.1) *is k-stochastic, then Z is the minimum point of f_W on $\Omega_k(H)$.*

Proof. First, we observe that function f_W is strictly convex, so its minimum on the convex set $\Omega_k(H)$ is unique. Next,

$$\frac{\partial}{\partial x_S} f_W(X) = \ln \frac{x_S}{w_S} + 1, \tag{3.2}$$

which is finite if $x_S > 0$ and is $-\infty$ if $x_S = 0$ (we consider the right derivative in this case). If $z_S = 0$ for some S, then for a sufficiently small $\epsilon > 0$ we have

$$f_W\big((1-\epsilon)Z + \epsilon Y\big) < f_W(Z),$$

which is a contradiction. Hence $z_S > 0$ for all $S \in H$.

Since the minimum point Z lies in the relative interior of $\Omega_k(H)$, considered as a convex polyhedron in $\mathbb{R}^H$, the Lagrange multiplier condition implies that there exist real $\mu_v : v \in V$ such that

$$\ln \frac{z_S}{w_S} = \sum_{v \in S} \mu_v \quad \text{for all} \quad S \in H. \tag{3.3}$$

Hence, letting $\lambda_v = e^{\mu_v}$ for $v \in V$, we obtain

$$z_S = \left(\prod_{v \in S} \lambda_v\right) w_S \quad \text{for all} \quad S \in H.$$

Now,

$$f_W(Z) = \sum_{S \in H} z_S \left(\sum_{v \in S} \ln \lambda_v\right) = \sum_{v \in V} \ln \lambda_v \left(\sum_{\substack{S \in H \\ S \ni v}} z_S\right) = \sum_{v \in V} \ln \lambda_v,$$

as desired.

If (3.1) holds for some $\lambda_v > 0$ and k-stochastic $Z = \{z_S\}$, then (3.3) holds with $\mu_v = \ln \lambda_v$, and by (3.2) we conclude that Z is a critical point of f_W in the relative interior of $\Omega_k(H)$. Since f_W is strictly convex, Z must be the minimum point of f_W on $\Omega_k(H)$. □

Theorem 3.1 implies that any positive weight W on a hypergraph H having a positive k-stochastic weight can be scaled uniquely to a k-stochastic weight Z, in which case we have $P_H(W) = \exp\{-f_W(Z)\}P_H(Z)$. Scaling factors $\{\lambda_v > 0 : v \in V\}$, however, do not have to be unique, as the example of a complete k-partite hypergraph readily shows (although in

https://doi.org/10.1017/S0963548311000435 Published online by Cambridge University Press

the case of the complete k-uniform hypergraph the scaling factors are unique). We note that if H is the complete k-uniform hypergraph or the complete k-partite hypergraph then there is a positive k-stochastic weight $Y = \{y_S : S \in H\}$ on H. In the former case we can choose

$$y_S = \binom{km-1}{k-1}^{-1} \quad \text{for all} \quad S \in H,$$

while in the latter case we can choose

$$y_S = m^{-k+1} \quad \text{for all} \quad S \in H.$$

We need a dual description of the scaling factors λ_v.

Theorem 3.2. *Let H be a k-uniform hypergraph with a set V of $|V| = km$ vertices and let $W = \{w_S : S \in H\}$ be a positive weight on H. Let $\lambda_v > 0 : v \in V$ be reals such that the weight $Z = \{z_S\}$ defined by*

$$z_S = \left(\prod_{v \in S} \lambda_v\right) w_S \quad \textit{for all} \quad S \in H$$

is k-stochastic.

Let us define a set $C(W) \subset \mathbb{R}^V$ by

$$C(W) = \left\{(x_v,\ v \in V) : \sum_{S \in H} w_S \exp\left\{\sum_{v \in S} x_v\right\} \leqslant m\right\}.$$

Then the point $(\mu_v : v \in V)$, where $\mu_v = \ln \lambda_v$ for all $v \in V$, is a maximum point of the linear function $\sum_{v \in V} x_v$ on $C(W)$.

Proof. Since weight Z is k-stochastic, we have

$$\sum_{\substack{S \in H \\ S \ni u}} w_S \exp\left\{\sum_{v \in S} \mu_v\right\} = 1 \quad \text{for all} \quad u \in V,$$

which means that $(\mu_v : v \in V)$ is a critical point of the linear function $\sum_{v \in V} x_v$ on the smooth surface defined in $\mathbb{R}^V$ by the equation

$$\sum_{S \in H} w_S \exp\left\{\sum_{v \in S} x_v\right\} = m.$$

The set $C(W)$ is convex and hence $(\mu_v : v \in V)$ has to be an extremum point of function $\sum_{v \in V} x_v$ on $C(W)$. Moreover, it has to be a maximum point since the function is unbounded from below on $C(W)$. □

4. Scaling balanced weights

Our proof of Theorem 1.1 is based on two lemmas.

Lemma 4.1. *Let us fix an integer $k > 1$ and a real $\alpha \geqslant 1$, and let H be a complete k-uniform hypergraph or a complete k-partite hypergraph. If $W = \{w_S\}$ is an α-balanced weight on H*

https://doi.org/10.1017/S0963548311000435 Published online by Cambridge University Press

and if $Z = \{z_S\}$ is the k-stochastic weight obtained from W by scaling, then Z is α^{k+1}-balanced.

Proof. Let V be the set of vertices of hypergraph H. Without loss of generality, we assume that $|V| > k$. For a subset $X \subset V$, we let

$$H_X = \{S \in H : S \supset X\}$$

denote the set of edges of H containing X. Let $\{\lambda_v > 0 : v \in V\}$ be scaling factors such that

$$z_S = \left(\prod_{v \in S} \lambda_v\right) w_S \quad \text{for all} \quad S \in H. \tag{4.1}$$

Suppose first that $H = V_1 \times \cdots \times V_k$ is a complete k-partite hypergraph, so $V = V_1 \cup \cdots \cup V_k$ and $|V_1| = \cdots = |V_k|$. For every $i = 1, \ldots, k$ and for every pair of vertices $v, u \in V_i$, we have

$$\sum_{S \in H_{\{v\}}} z_S = \sum_{S \in H_{\{u\}}} z_S = 1. \tag{4.2}$$

Let us consider the bijection $\phi : H_{\{v\}} \longrightarrow H_{\{u\}}$ defined by

$$\phi(S) = S \cup \{u\} \setminus \{v\}. \tag{4.3}$$

By (4.1) we have

$$\frac{z_{\phi(S)}}{z_S} = \frac{\lambda_u}{\lambda_v} \cdot \frac{w_{\phi(S)}}{w_S}. \tag{4.4}$$

Since weight W is α-balanced, in view of (4.2) we conclude that

$$\frac{1}{\alpha} \leqslant \frac{\lambda_u}{\lambda_v} \leqslant \alpha, \tag{4.5}$$

which proves that Z is α^{k+1}-balanced.

Suppose now that $H = \binom{V}{k}$ is a complete k-uniform hypergraph. Then, for any two distinct vertices $u, v \in V$ we have

$$\sum_{S \in H_v \setminus H_{\{u,v\}}} z_S = \sum_{S \in H_u \setminus H_{\{u,v\}}} z_S = 1 - \sum_{S \in H_{\{u,v\}}} z_S > 0. \tag{4.6}$$

Let us consider the bijection $\phi : H_v \setminus H_{\{u,v\}} \longrightarrow H_u \setminus H_{\{u,v\}}$ defined by (4.3). From (4.1) we deduce that (4.4) holds, and in view of (4.6) we conclude that (4.5) follows. Since weight W is α-balanced, (4.5) implies that Z is α^{k+1}-balanced. □

The second lemma asserts that if we scale a weight to a k-stochastic weight, and the weight is already sufficiently close to being k-stochastic, then the product of the scaling factors is close to 1.

Lemma 4.2. *Let us fix an integer $k > 1$ and real $\alpha \geqslant 1$ and $\delta > 0$. Then there exist integer $m_0 = m_0(k, \alpha, \delta) > 0$ and real $\beta = \beta(k, \alpha, \delta) > 0$ such that the following holds.*

https://doi.org/10.1017/S0963548311000435 Published online by Cambridge University Press

Suppose that H is a complete k-uniform hypergraph or a complete k-partite hypergraph with a set V of vertices, where $|V| = km$ and $m \geqslant m_0$. Suppose that $W = \{w_S\}$ is an α-balanced weight on H, that

$$\sum_{S \in H} w_S = m$$

and that

$$\left| 1 - \sum_{\substack{S \in H \\ S \ni v}} w_S \right| \leqslant \frac{\delta}{m} \quad \text{for all} \quad v \in V.$$

Let $\lambda_v > 0 : v \in V$ be reals such that weight $Z = \{z_S\}$ defined by

$$z_S = \left(\prod_{v \in S} \lambda_v \right) w_S \quad \text{for all} \quad S \in H$$

is k-stochastic. Then

$$0 \leqslant \sum_{v \in V} \ln \lambda_v \leqslant \frac{\beta}{m}.$$

One can choose $\beta = \alpha\delta^2(k+1)^2$ and $m_0 = \max\{1 + \lceil \alpha\delta k \rceil,\ k\}$.

Proof. We note that the point $\left(x_v = 0 : v \in V\right)$ belongs to the set $C(W)$ of Theorem 3.2, and so by Theorem 3.2 we have

$$\sum_{v \in V} \ln \lambda_v \geqslant \sum_{v \in V} x_v = 0.$$

Let us define

$$\delta_v = 1 - \sum_{\substack{S \in H \\ S \ni v}} w_S \quad \text{for} \quad v \in V.$$

Then

$$\sum_{v \in V} \delta_v = \sum_{v \in V} \left(1 - \sum_{\substack{S \in H \\ S \ni v}} w_S \right) = km - k \sum_{S \in H} w_S = 0. \tag{4.7}$$

In addition, if $H = V_1 \times \cdots \times V_k$ is the complete k-partite graph, where $V = V_1 \cup \cdots \cup V_k$ and $|V_1| = \cdots = |V_k| = m$, for every $i = 1, \ldots, k$ we have

$$\sum_{v \in V_i} \delta_v = \sum_{v \in V_i} \left(1 - \sum_{\substack{S \in H \\ S \ni v}} w_S \right) = m - \sum_{S \in H} w_S = 0. \tag{4.8}$$

We define numbers $\{\rho_S : S \in H\}$ as follows. If $H = \binom{V}{k}$ is the complete k-uniform hypergraph, we define

$$\rho_S = \binom{km-2}{k-1}^{-1} \sum_{v \in S} \delta_v \quad \text{for all} \quad S \in H.$$

https://doi.org/10.1017/S0963548311000435 Published online by Cambridge University Press

If $H = V_1 \times \cdots \times V_k$ is a complete k-partite graph, we define

$$\rho_S = \frac{1}{m^{k-1}} \sum_{v \in S} \delta_v \quad \text{for all} \quad S \in H.$$

We claim that

$$\sum_{\substack{S \in H \\ S \ni v}} \rho_S = \delta_v \quad \text{for all} \quad v \in V. \tag{4.9}$$

Indeed, if $H = \binom{V}{k}$, then using (4.7) we obtain

$$\begin{aligned} \sum_{\substack{S \in H \\ S \ni v}} \rho_S &= \frac{\binom{km-1}{k-1}}{\binom{km-2}{k-1}} \delta_v + \frac{\binom{km-2}{k-2}}{\binom{km-2}{k-1}} \sum_{u \in V \setminus \{v\}} \delta_u \\ &= \frac{\binom{km-1}{k-1} - \binom{km-2}{k-2}}{\binom{km-2}{k-1}} \delta_v \\ &= \delta_v, \end{aligned}$$

and if $H = V_1 \times \cdots \times V_k$, then using (4.8) we obtain that for all $i = 1, \ldots, k$ and for all $v \in V_i$ we have

$$\sum_{\substack{S \in H \\ S \ni v}} \rho_S = \delta_v + \frac{m^{k-2}}{m^{k-1}} \sum_{u \in V \setminus V_i} \delta_u = \delta_v.$$

In either case, (4.9) holds. In addition, from (4.7),

$$\sum_{S \in H} \rho_S = \frac{1}{k} \sum_{v \in V} \sum_{\substack{S \in H \\ S \ni v}} \rho_S = \frac{1}{k} \sum_{v \in V} \delta_v = 0. \tag{4.10}$$

Let us define

$$x_S = w_S + \rho_S \quad \text{for all} \quad S \in H.$$

Then, from (4.9) we have

$$\sum_{\substack{S \in H \\ S \ni v}} x_S = 1 \quad \text{for all} \quad v \in V.$$

Since weight W is α-balanced, for all $S \in H$ we have

$$w_S \geqslant \binom{km}{k}^{-1} \frac{m}{\alpha} \quad \text{when} \quad H = \binom{V}{k} \tag{4.11}$$

and

$$w_S \geqslant \frac{1}{\alpha m^{k-1}} \quad \text{when} \quad H = V_1 \times \cdots \times V_k. \tag{4.12}$$

On the other hand, for all $S \in H$ we have

$$|\rho_S| \leqslant \binom{km-2}{k-1}^{-1} \frac{\delta k}{m} \quad \text{when} \quad H = \binom{V}{k} \tag{4.13}$$

https://doi.org/10.1017/S0963548311000435 Published online by Cambridge University Press

and

$$|\rho_S| \leqslant \frac{\delta k}{m^k} \quad \text{when} \quad H = V_1 \times \cdots \times V_k. \tag{4.14}$$

From (4.11) and (4.13), we conclude that if $H = \binom{V}{k}$ then

$$x_S \geqslant 0 \quad \text{for all} \quad S \in H \quad \text{provided} \quad \frac{m(m-1)}{km-1} \geqslant \alpha\delta,$$

whereas from (4.12) and (4.14) we conclude that if $H = V_1 \times \cdots \times V_k$ then

$$x_S \geqslant 0 \quad \text{for all} \quad S \in H \quad \text{provided} \quad m \geqslant \alpha\delta k.$$

In either case, $X = \{x_S\}$ is a k-stochastic weight on H provided $m \geqslant \alpha\delta k + 1$. Using (4.10), we conclude from Theorem 3.1 that for $m \geqslant \alpha\delta k + 1$ we have

$$\begin{aligned}\sum_{v\in V} \ln \lambda_v &\leqslant \sum_{S\in H} x_S \ln \frac{x_S}{w_S} = \sum_{S\in H} (w_S + \rho_S) \ln \frac{w_S + \rho_S}{w_S} \\ &= \sum_{S\in H} (w_S + \rho_S) \ln\left(1 + \frac{\rho_S}{w_S}\right) \leqslant \sum_{S\in H} (w_S + \rho_S) \frac{\rho_S}{w_S} \\ &= \sum_{S\in H} \frac{\rho_S^2}{w_S}. \end{aligned} \tag{4.15}$$

From (4.11) and (4.13), we conclude that in the case of $H = \binom{V}{k}$ sum (4.15) does not exceed

$$\frac{\alpha\delta^2 k^2 \binom{km}{k}^2}{m^3 \binom{km-2}{k-1}^2} = \frac{\alpha\delta^2(km-1)^2}{m(m-1)^2},$$

whereas from (4.12) and (4.14) we conclude that in the case of $H = V_1 \times \cdots \times V_k$ sum (4.15) does not exceed

$$\frac{\alpha\delta^2 k^2 m^{k-1}}{m^{2k}} m^k = \frac{\alpha\delta^2 k^2}{m}.$$

In either case, sum (4.15) does not exceed $\alpha\delta^2(k+1)^2/m$ as long as $m \geqslant k$. □

5. Proofs of Theorems 1.1 and 1.2

Our approach is somewhat similar to Bregman's original approach [3], combining scaling and induction to obtain upper bounds on permanents. Before giving a formal proof, we illustrate the idea of the proof by sketching it in the more familiar case of permanents, that is, when $k = 2$ and the underlying graph is bipartite.

All implied constants in the 'O' notation in this section may depend on k and α only.

https://doi.org/10.1017/S0963548311000435 Published online by Cambridge University Press

5.1. The idea of the proof

Let us fix $\alpha \geqslant 1$. Let $A = (a_{ij})$ be an α-balanced $m \times m$ doubly stochastic matrix. Our goal is to prove that $\operatorname{per} A = e^{-m} m^{O(1)}$, or, equivalently, that

$$\operatorname{per} A = \exp\left\{-m + O\left(\sum_{j=1}^{m} \frac{1}{j}\right)\right\}. \tag{5.1}$$

We proceed by induction on m.

Using the first row expansion, we can write

$$\operatorname{per} A = \sum_{j=1}^{m} a_{1j} \operatorname{per} \widehat{A}_j, \tag{5.2}$$

where $\widehat{A}_j$ is the $(m-1) \times (m-1)$ matrix obtained from A by crossing out the first row and jth column. We have $a_{1j} \leqslant \alpha/m$ for all j and, using that A is doubly stochastic, we conclude that the sum of σ_j of all entries of $\widehat{A}_j$ satisfies

$$m - 2 \leqslant \sigma_j \leqslant m - 2 + \frac{\alpha}{m}. \tag{5.3}$$

Let us define $(m-1) \times (m-1)$ matrices B_j by

$$B_j = \frac{m-1}{\sigma_j} \widehat{A}_j.$$

Hence the sum of all entries of B_j is $m-1$, and by (5.3) we have

$$\operatorname{per} \widehat{A}_j = \left(\frac{\sigma_j}{m-1}\right)^{m-1} \operatorname{per} B_j = \exp\left\{-1 + O\left(\frac{1}{m}\right)\right\} \operatorname{per} B_j. \tag{5.4}$$

Matrices B_j are not necessarily doubly stochastic, but they are reasonably close to doubly stochastic, since all the row and column sums of B_j are $1 + O(m^{-1})$. Let C_j be the $(m-1) \times (m-1)$ doubly stochastic matrix obtained from B_j by scaling. By Lemma 4.2 we have

$$\operatorname{per} B_j = \exp\left\{O\left(\frac{1}{m}\right)\right\} \operatorname{per} C_j. \tag{5.5}$$

Combining (5.2) and (5.3)–(5.5), we conclude that

$$\operatorname{per} A = \exp\left\{-1 + O\left(\frac{1}{m}\right)\right\} \sum_{j=1}^{m} a_{1j} \operatorname{per} C_j, \tag{5.6}$$

where

$$\sum_{j=1}^{m} a_{1j} = 1 \quad \text{and} \quad a_{1j} \geqslant 0 \quad \text{for all} \quad j = 1, \ldots, m.$$

Up to this point, we did not really use the condition that A is α-balanced: we used only that the entries of A are uniformly small, of the order of $O(m^{-1})$. To proceed with the induction, we have to show that the entries of the doubly stochastic matrices C_j in (5.6) and all other doubly stochastic matrices obtained by iterating the recursion are also uniformly small. Now we observe that C_j is obtained by scaling of $\widehat{A}_j$ and hence by

https://doi.org/10.1017/S0963548311000435 Published online by Cambridge University Press

Lemma 4.1 is α^3-balanced. Similarly, as we iterate recursion (5.6), the doubly stochastic matrices that we obtain are α^3-balanced, since they are obtained by scaling from some submatrices of an α-balanced matrix A. This allows us to use (5.6) in the induction step to obtain (5.1).

Permanents of α-balanced matrices are studied in [13] and [5].

Proof of Theorem 1.1. Without loss of generality, we assume that $\alpha^{k+1} > 2$. Let H be either a complete k-uniform hypergraph $\binom{V}{k}$ or a complete k-partite hypergraph $V_1 \times \cdots \times V_k$ with a set V of $|V| = km$ vertices. Let $Z = \{z_S\}$ be a k-stochastic α-balanced weight on H.

If $H = \binom{V}{k}$ and $U \subset V$ is a subset such that $|U| = kl$ for some integer $l \geqslant 1$, we consider the induced hypergraph $H|U$ consisting of the edges $S \in H$ such that $S \subset U$. Hence $H|U = \binom{U}{k}$ is the complete k-uniform hypergraph with the set U of vertices.

Similarly, if $H = V_1 \times \cdots \times V_k$ and if $|U \cap V_1| = \cdots = |U \cap V_k| = l$ for some integer $l \geqslant 1$, we consider the restriction $H|U$ consisting of the edges $S \in H$ such that $S \subset U$. In this case, $H|U$ is a uniform k-partite graph with the set U of vertices, $H|U = U_1 \times \cdots \times U_k$ where $U_i = V_i \cap U$ for $i = 1, \ldots, k$.

For a subset $U \subset V$ as above, we define a weight

$$Z^U = \left\{ z_S^U : S \in H|U \right\}$$

on $H|U$ as follows. We consider the restriction of weight Z onto hypergraph $H|U$ and define Z^U to be the scaling of the restriction to a k-stochastic weight. We consider the partition function associated with the hypergraph $H|U$, which we denote by P_U. We want to estimate $P_U(Z^U)$.

Let $A \in H|U$ be an edge. We consider the complement $U \setminus A$, the corresponding hypergraph $H|(U \setminus A)$, weight $Z^{U \setminus A}$ and the partition function $P_{U \setminus A}(Z^{U \setminus A})$.

Our goal is to prove that for some $\gamma_1 = \gamma_1(k, \alpha) > 0$, $\gamma_2 = \gamma_2(k, \alpha) > 0$ and $l_0 = l_0(k, \alpha)$ we have

$$\begin{aligned} P_U(Z^U) &\geqslant \exp\left\{ -k + 1 - \frac{\gamma_1}{l-1} \right\} \min_{A \in H|U} P_{U \setminus A}(Z^{U \setminus A}), \\ P_U(Z^U) &\leqslant \exp\left\{ -k + 1 + \frac{\gamma_2}{l-1} \right\} \max_{A \in H|U} P_{U \setminus A}(Z^{U \setminus A}), \end{aligned} \tag{5.7}$$

provided $l \geqslant l_0$ (recall that $|U| = kl$).

We show that we can choose

$$\gamma_1 = \alpha^{3(k+1)}(k^2 + k)^2 + (k-1)^2, \quad \gamma_2 = \frac{k^2 \alpha^{k+1}}{2} \quad \text{and} \quad l_0 = \lceil \alpha^{2(k+1)} k^2 \rceil + 1.$$

Since the restriction of Z onto $H|U$ is α-balanced, by Lemma 4.1 the weight Z^U is α^{k+1}-balanced. Crude estimates give

$$\alpha^{-(k+1)l_0} l_0^{l_0} \binom{kl_0}{k}^{1-l_0} \leqslant P_U(Z^U) \leqslant \alpha^{(k+1)l_0} l_0^{l_0 - kl_0 + k}$$

if $|U| = kl_0$.

https://doi.org/10.1017/S0963548311000435 Published online by Cambridge University Press

Starting with $U = V$, $l = m$ and $Z^U = Z$, by iterating (5.7) we obtain

$$P(Z) \geqslant \alpha^{-(k+1)l_0} l_0^{l_0} \binom{kl_0}{k}^{1-l_0} \exp\left\{-(k-1)(m-l_0) - \gamma_1 \sum_{j=l_0+1}^{m} \frac{1}{j-1}\right\},$$

$$P(Z) \leqslant \alpha^{(k+1)l_0} l_0^{l_0-kl_0+k} \exp\left\{-(k-1)(m-l_0) + \gamma_2 \sum_{j=l_0+1}^{m} \frac{1}{j-1}\right\}.$$

In particular,

$$P(Z) = \exp\left\{-(k-1)m + O\left(\sum_{j=1}^{m} \frac{1}{j}\right)\right\},$$

which completes the proof of the theorem.

We proceed to prove (5.7), assuming that $l \geqslant \alpha^{2(k+1)}k^2 + 1$. Since weight Z^U is α^{k+1}-balanced, we have

$$z_S^U \leqslant \alpha^{k+1} l \binom{kl}{k}^{-1} \quad \text{for all} \quad S \subset U \quad \text{when} \quad H = \binom{V}{k} \tag{5.8}$$

and

$$z_S^U \leqslant \alpha^{k+1} l^{-k+1} \quad \text{for all} \quad S \subset U \quad \text{when} \quad H = V_1 \times \cdots \times V_k. \tag{5.9}$$

Let us pick an element $u \in U$. Then there is a recursion

$$P_U(Z^U) = \sum_{\substack{A \in H|U \\ A \ni u}} z_A^U \cdot P_{U \setminus A}(Z^U). \tag{5.10}$$

Here $P_{U \setminus A}(Z^U)$ is the partition function computed on the restriction of the weight Z^U onto the hypergraph $H|(U \setminus A)$, which is not the same as $P_{U \setminus A}(Z^{U \setminus A})$, since the weight $Z^{U \setminus A}$ is obtained from Z^U by restricting it onto $H|(U \setminus A)$ and scaling the restriction to a k-stochastic weight.

Since Z^U is a k-stochastic weight on $H|U$, we have

$$\sum_{\substack{A \in H|U \\ A \ni u}} z_A^U = 1 \quad \text{and} \quad z_A^U \geqslant 0 \quad \text{for all} \quad A \in H|U. \tag{5.11}$$

Let

$$\sigma_A^U = \sum_{S \in H|(U \setminus A)} z_S^U$$

be the sum of the weights in the restriction Z^U onto $H|(U \setminus A)$, that is, the sum of the weights z_S^U for the edges $S \subset U$ not intersecting an edge $A \in H|U$. Since weight Z^U is k-stochastic, we have

$$\sigma_A^U = l - k + O\left(\frac{1}{l}\right).$$

https://doi.org/10.1017/S0963548311000435 Published online by Cambridge University Press

More precisely, from (5.8) we have

$$l-k \leqslant \sigma_A^U \leqslant l-k+\binom{k}{2}\binom{kl-2}{k-2}\alpha^{k+1}l\binom{kl}{k}^{-1}$$

if H is a complete k-uniform hypergraph, and from (5.9) we have

$$l-k \leqslant \sigma_A^U \leqslant l-k+\binom{k}{2}\alpha^{k+1}l^{-1},$$

if H is a complete k-partite hypergraph. In either case,

$$l-k \leqslant \sigma_A^U \leqslant l-k+\frac{\alpha^{k+1}k^2}{2(l-1)}. \tag{5.12}$$

Similarly, from (5.8) we have

$$1-k\binom{kl-2}{k-2}\alpha^{k+1}l\binom{kl}{k}^{-1} \leqslant \sum_{\substack{S\in H|(U\setminus A)\\ S\ni v}} z_S^U \leqslant 1 \quad \text{for all} \quad v\in U\setminus A$$

if H is a complete k-uniform hypergraph, and from (5.9) we have

$$1-k\alpha^{k+1}l^{-1} \leqslant \sum_{\substack{S\in H|(U\setminus A)\\ S\ni v}} z_S^U \leqslant 1 \quad \text{for all} \quad v\in U\setminus A$$

if H is a complete k-partite hypergraph. In either case,

$$1-\frac{k\alpha^{k+1}}{l-1} \leqslant \sum_{\substack{S\in H|(U\setminus A)\\ S\ni v}} z_S^U \leqslant 1 \quad \text{for all} \quad v\in U\setminus A. \tag{5.13}$$

Let us define a weight

$$W^{U\setminus A}=\left\{w_S^{U\setminus A} : S\in H|(U\setminus A)\right\}$$

by scaling the restriction of the weight Z^U onto $H|(U\setminus A)$ to the total sum $l-1$, so that

$$w_S^{U\setminus A}=\frac{l-1}{\sigma_A^U}\, z_S^U \quad \text{for all} \quad S\in H|(U\setminus A).$$

Hence

$$P_{U\setminus A}(Z^U)=\left(\frac{\sigma_A^U}{l-1}\right)^{l-1} P_{U\setminus A}(W^{U\setminus A}). \tag{5.14}$$

We have

$$\left(\frac{\sigma_A^U}{l-1}\right)^{l-1}=\exp\left\{-k+1+O\left(\frac{1}{l}\right)\right\}.$$

More precisely, from (5.12) we have

$$\exp\left\{-k+1-\frac{(k-1)^2}{l-1}\right\} \leqslant \left(\frac{\sigma_A^U}{l-1}\right)^{l-1} \leqslant \exp\left\{-k+1+\frac{k^2\alpha^{k+1}}{2(l-1)}\right\}. \tag{5.15}$$

https://doi.org/10.1017/S0963548311000435 Published online by Cambridge University Press

Moreover, from (5.12) and (5.13) we deduce that

$$1-\frac{k\alpha^{k+1}}{l-1}\leqslant\sum_{\substack{S\in H|(U\setminus A)\\ S\ni v}}w_S^{U\setminus A}\leqslant 1+\frac{2(k-1)}{l-1}\quad\text{for all}\quad v\in U\setminus A. \tag{5.16}$$

We intend to apply Lemma 4.2 to weight $W^{U\setminus A}$. We observe that weight $W^{U\setminus A}$ is obtained from weight Z^U by restricting it onto the set $U\setminus A$ and then scaling to the total sum $l-1$ of components. Therefore, the k-stochastic weight on $U\setminus A$ obtained from $W^{U\setminus A}$ by scaling is just $Z^{U\setminus A}$, the k-stochastic weight obtained by restricting the original weight Z onto $U\setminus A$ and scaling.

We have

$$P_{U\setminus A}(W^{U\setminus A})=\exp\left\{O\left(\frac{1}{l}\right)\right\}P_{U\setminus A}(Z^{U\setminus A}).$$

More precisely, since $W^{U\setminus A}$ is α^{k+1}-balanced and (5.16) holds, by Lemma 4.2 we conclude that

$$\begin{aligned}&P_{U\setminus A}(W^{U\setminus A})\geqslant\exp\left\{-\frac{\alpha^{3(k+1)}(k^2+k)^2}{l-1}\right\}P_{U\setminus A}(Z^{U\setminus A}),\\ &P_{U\setminus A}(W^{U\setminus A})\leqslant P_{U\setminus A}(Z^{U\setminus A}).\end{aligned} \tag{5.17}$$

Combining (5.10), (5.11), (5.14), (5.15) and (5.17), we obtain (5.7) with

$$\gamma_1=\alpha^{3(k+1)}(k^2+k)^2+(k-1)^2\quad\text{and}\quad\gamma_2=\frac{k^2\alpha^{k+1}}{2},$$

which completes the proof. □

Proof of Theorem 1.2. Let Z be the k-stochastic weight obtained from weight W by scaling. By Theorem 3.1 we have

$$P_H(Z)=f_W(Z)P_H(W)=e^{\zeta}P_H(W).$$

Moreover, by Lemma 4.1, weight Z is α^{k+1}-balanced and the proof follows by Theorem 1.1 applied to Z. Furthermore, weights Z^U constructed in the proof of Theorem 1.1, being scalings of restrictions of W onto subsets $U\subset V$, are also α^{k+1}-balanced, and hence we can use the same estimates for $P_H(Z)$ as in Theorem 1.1. □

Acknowledgement

The authors are grateful to Jeff Kahn for helpful discussions.

References

[1] Alon, N. and Friedland, S. (2008) The maximum number of perfect matchings in graphs with a given degree sequence. *Electron. J. Combin.* **15** #13.

[2] Barvinok, A. (2011) A bound for the number of vertices of a polytope with applications. `arXiv:1108.2871`.

https://doi.org/10.1017/S0963548311000435 Published online by Cambridge University Press

[3] Bregman, L. M. (1973) Certain properties of nonnegative matrices and their permanents (in Russian). *Dokl. Akad. Nauk SSSR* **211** 27–30.
[4] Colbourn, C., Hoffman, D. G., Phelps, K. T., Rödl, V. and Winkler, P. M. (1991) The number of t-wise balanced designs. *Combinatorica* **11** 207–218.
[5] Costello, K. P. and Vu, V. (2009) Concentration of random determinants and permanent estimators. *SIAM J. Discrete Math.* **23** 1356–1371.
[6] Dow, S. J. and Gibson, P. M. (1987) An upper bound for the permanent of a 3-dimensional $(0,1)$-matrix. *Proc. Amer. Math. Soc.* **99** 29–34.
[7] Dow, S. J. and Gibson, P. M. (1987) Permanents of d-dimensional matrices. *Linear Algebra Appl.* **90** 133–145.
[8] Egorychev, G. P. (1981) The solution of van der Waerden's problem for permanents. *Adv. Math.* **42** 299–305.
[9] Esperet, L., Kardoš, F., King, A. D., Král', D. and Norine, S. (2011) Exponentially many perfect matchings in cubic graphs. Advances in Mathematics **227** 1646–1664.
[10] Falikman, D. I. (1981) Proof of the van der Waerden conjecture on the permanent of a doubly stochastic matrix (in Russian). *Mat. Zametki* **29** 931–938.
[11] Friedland, S. (2011) Positive diagonal scaling of a nonnegative tensor to one with prescribed slice sums. *Linear Algebra Appl.* **434** 1615–1619.
[12] Friedland, S. (2011) Analogs of the van der Waerden and Tverberg conjectures for haffnians. `arXiv:1102.2542`.
[13] Friedland, S., Rider, B. and Zeitouni, O. (2004) Concentration of permanent estimators for certain large matrices. *Ann. Appl. Probab.* **14** 1559–1576.
[14] Gurvits, L. (2008) Van der Waerden/Schrijver-Valiant like conjectures and stable (aka hyperbolic) homogeneous polynomials: One theorem for all. With a corrigendum. *Electron. J. Combin.* **15** #66.
[15] Jerrum, M., Sinclair, A. and Vigoda, E. (2004) A polynomial-time approximation algorithm for the permanent of a matrix with nonnegative entries. *J. Assoc. Comput. Mach.* **51** 671–697.
[16] Karp, R. M. (1972) Reducibility among combinatorial problems. In *Complexity of Computer Computations: Proc. Sympos. IBM Thomas J. Watson Res. Center, 1972*, Plenum, pp. 85–103.
[17] Linial, N., Samorodnitsky, A. and Wigderson, A. (2000) A deterministic strongly polynomial algorithm for matrix scaling and approximate permanents. *Combinatorica* **20** 545–568.
[18] Lovász, L. and Plummer, M. D. (2009) *Matching Theory*, AMS Chelsea Publishing.
[19] Minc, H. (1978) *Permanents*, Vol. 6 of *Encyclopedia of Mathematics and its Applications*, Addison-Wesley.
[20] Nesterov, Y. and Nemirovskii, A. (1994) *Interior-Point Polynomial Algorithms in Convex Programming*, Vol. 13 of *SIAM Studies in Applied Mathematics*, SIAM.
[21] Schrijver, A. (1978) A short proof of Minc's conjecture. *J. Combin. Theory Ser. A* **25** 80–83.
[22] Soules, G. W. (2003) New permanental upper bounds for nonnegative matrices. *Linear Multilinear Algebra* **51** 319–337.
[23] Valiant, L. G. (1979) The complexity of computing the permanent. *Theoret. Comput. Sci.* **8** 189–201.
[24] Vu, V. H. (2000) New bounds on nearly perfect matchings in hypergraphs: Higher codegrees do help. *Random Struct. Alg.* **17** 29–63.

https://doi.org/10.1017/S0963548311000435 Published online by Cambridge University Press