Ergod. Th. & Dynam. Sys. (2005), **25**, 1503–1520
doi:10.1017/S0143385705000143
© *2005 Cambridge University Press*
Printed in the United Kingdom

Subquotients of Hecke C^*-algebras

NATHAN BROWNLOWE†, NADIA S. LARSEN‡‖, IAN F. PUTNAM§ and IAIN RAEBURN†

† *School of Mathematical and Physical Sciences, University of Newcastle, NSW 2308, Australia*
(e-mail: Nathan.Brownlowe@studentmail.newcastle.edu.au, iain@frey.newcastle.edu.au)
‡ *Department of Mathematics, Institute for Mathematical Sciences, University of Copenhagen, Universitetsparken 5, 2100 Copenhagen, Denmark*
(e-mail: nadiasl@math.uio.no)
§ *Department of Mathematics and Statistics, University of Victoria, British Columbia V8W 3P4, Canada*
(e-mail: putnam@math.uvic.ca)

(*Received* 17 *September* 2003 *and accepted in revised form* 28 *April* 2005)

Abstract. We realize the Hecke C^*-algebra $\mathcal{C}_\mathbb{Q}$ of Bost and Connes as a direct limit of Hecke C^*-algebras which are semigroup crossed products by $\mathbb{N}^F$, for F a finite set of primes. For each approximating Hecke C^*-algebra we describe a composition series of ideals. In all cases there is a large type I ideal and a commutative quotient, and the intermediate subquotients are direct sums of simple C^*-algebras. We can describe the simple summands as ordinary crossed products by actions of $\mathbb{Z}^S$ for S a finite set of primes. When $|S| = 1$, these actions are odometers and the crossed products are Bunce–Deddens algebras; when $|S| > 1$, the actions are an apparently new class of higher-rank odometer actions, and the crossed products are an apparently new class of classifiable AT-algebras.

0. *Introduction*

In [**2**], Bost and Connes studied a particular Hecke C^*-algebra $\mathcal{C}_\mathbb{Q}$ arising in number theory. The algebra $\mathcal{C}_\mathbb{Q}$ can be realized as a semigroup crossed product $C^*(\mathbb{Q}/\mathbb{Z}) \rtimes_\alpha \mathbb{N}^*$ by an endomorphic action α of the multiplicative semigroup $\mathbb{N}^*$ on the group C^*-algebra $C^*(\mathbb{Q}/\mathbb{Z})$ (see [**8**]), and this realization has provided useful insight into the analysis of $\mathcal{C}_\mathbb{Q}$ (see [**6, 16**]). Since individual elements of $\mathbb{Q}/\mathbb{Z}$ and $\mathbb{N}^*$ involve only finitely many primes, $C^*(\mathbb{Q}/\mathbb{Z}) \rtimes_\alpha \mathbb{N}^*$ is the direct limit of subalgebras $C^*(G_F) \rtimes_\alpha \mathbb{N}^F$, where F is a finite set of

‖ Current address: Department of Mathematics, University of Oslo, P.O. Box 1053, Blindern, N-0316 Oslo, Norway.

https://doi.org/10.1017/S0143385705000143 Published online by Cambridge University Press

primes, G_F is the subgroup of $\mathbb{Q}/\mathbb{Z}$ in which the denominators have all prime factors in F, and $\mathbb{N}^F$ acts through the embedding $(n_p) \mapsto \prod_{p\in F} p^{n_p}$ of $\mathbb{N}^F$ in $\mathbb{N}^*$ (see §1). One can therefore hope to understand the Hecke algebra $\mathcal{C}_\mathbb{Q}$ in terms of the finite-prime analogues $C^*(G_F) \rtimes_\alpha \mathbb{N}^F$.

Our goal is to analyse the structure of these finite-prime analogues of the Bost–Connes algebra. We started this analysis in [**11**], where we described a composition series for the two-prime analogue and identified the subquotients in familiar terms: there is a large type I ideal, a commutative quotient isomorphic to $C(\mathbb{T}^2)$, and the intermediate subquotient is isomorphic to a direct sum of Bunce–Deddens algebras. Here we describe a composition series for $C^*(G_F) \rtimes_\alpha \mathbb{N}^F$. Again there is a large type I ideal and a commutative quotient, and the intermediate subquotients are direct sums of simple C^*-algebras. We can describe the simple summands as ordinary crossed products by actions of $\mathbb{Z}^S$ for $S \subset F$. When $|S| = 1$, these actions are odometers and the crossed products are Bunce–Deddens algebras; when $|S| > 1$, the actions are an apparently new class of higher-rank odometer actions, and the crossed products are an apparently new class of classifiable AT-algebras.

We begin with a short section in which we describe the algebras we intend to study. In §2, we describe our composition series for the semigroup crossed product $C^*(G_F) \rtimes_\alpha \mathbb{N}^F$. It has $|F| + 1$ subquotients, and all but two of them are direct sums of algebras stably isomorphic to ordinary crossed products of the form $C(\mathcal{U}(\mathbb{Z}_{F\setminus S})) \rtimes \mathbb{Z}^S$, where $S \subset F$ and $\mathcal{U}(\mathbb{Z}_{F\setminus S})$ is the group of units in the ring $\prod_{p\in F\setminus S} \mathbb{Z}_p$. Our main tools are the analysis of invariant ideals in semigroup crossed products from [**10**] and some technical lemmas on sums and intersections of ideals in C^*-algebras. We also use the general results of [**19**] to see that the simple summands are classifiable.

In §3, we show that when $S = \{q\}$ is a singleton, $C(\mathcal{U}(\mathbb{Z}_{F\setminus S})) \rtimes \mathbb{Z}^S$ is a direct sum of finitely many Bunce–Deddens algebras; as in [**11**], the number of summands depends on the orders of q in the finite groups $\prod_{p\neq q} \mathcal{U}(\mathbb{Z}/p^l\mathbb{Z})$ for large $l \in \mathbb{N}$. We then consider the case where $S = \{q, r\}$. By computing the K-theory of $C(\mathcal{U}(\mathbb{Z}_{F\setminus S})) \rtimes \mathbb{Z}^S$, we can see that they are not Bunce–Deddens algebras, for example. We expect these summands to be even harder to recognize when $|S| > 2$.

In §4, we use techniques like those of §2 to identify subquotients of the Bost–Connes algebra $C^*(\mathbb{Q}/\mathbb{Z}) \rtimes_\alpha \mathbb{N}^*$. These include algebras stably isomorphic to $C(\mathcal{U}(\mathbb{Z}_{\mathcal{P}\setminus S})) \rtimes \mathbb{Z}^S$ when S is a cofinite subset of the set $\mathcal{P}$ of all primes; in this case, though, these crossed products are themselves simple, and even though the general theory of [**19**] no longer applies, we can see using results from [**1**] that they are classifiable AT-algebras. We finish with a purely number theoretic Appendix A in which we identify the orders of an odd integer in the groups $\mathcal{U}(\mathbb{Z}/p^l\mathbb{Z})$ and their products. As in [**11**, Theorem 3.1], these are needed when we want to identify the number of simple summands in the various subquotients.

1. *Preliminaries*

We denote by $\mathbb{N}^*$ the semigroup of positive integers under multiplication, and by $\mathbb{N}$ the semigroup of non-negative integers under addition. It was shown in [**8**, Proposition 2.1]

https://doi.org/10.1017/S0143385705000143 Published online by Cambridge University Press

that there is an action α of $\mathbb{N}^*$ by endomorphisms of $C^*(\mathbb{Q}/\mathbb{Z})$ such that

$$\alpha_n(\delta_r) = \frac{1}{n}\sum_{ns=r}\delta_s \quad \text{for } r \in \mathbb{Q}/\mathbb{Z} \text{ and } n \in \mathbb{N}^*.$$

The corresponding semigroup crossed product $C^*(\mathbb{Q}/\mathbb{Z}) \rtimes_\alpha \mathbb{N}^*$ is isomorphic to the Hecke C^*-algebra $\mathcal{C}_\mathbb{Q}$ of Bost and Connes [**8**, Corollary 2.10]. We denote by $(i_A, i_{\mathbb{N}^*})$ the canonical covariant representation of $(C^*(\mathbb{Q}/\mathbb{Z}), \mathbb{N}^*, \alpha)$ in $C^*(\mathbb{Q}/\mathbb{Z}) \rtimes_\alpha \mathbb{N}^*$.

Let F be a set of prime numbers. The rational numbers of the form $k\big(\prod_{p\in F} p^{m_p}\big)^{-1}$ form a subgroup of $\mathbb{Q}$, whose image in $\mathbb{Q}/\mathbb{Z}$ we denote by G_F. The integrated form of the map $r \mapsto \delta_r : G_F \to UC^*(\mathbb{Q}/\mathbb{Z})$ is a homomorphism i_F of $C^*(G_F)$ into $C^*(\mathbb{Q}/\mathbb{Z})$; a standard duality argument shows that i_F is injective, so that we can identify $C^*(G_F)$ with the subalgebra $i_F(C^*(G_F))$ of $C^*(\mathbb{Q}/\mathbb{Z})$. When n has all of its prime factors in F, α_n leaves this subalgebra invariant, and hence composing α with the map $(m_p)_{p\in F} \mapsto \prod_{p\in F} p^{m_p}$ gives an action of $\mathbb{N}^F$ on $C^*(G_F)$, which we also denote by α. The pair $(i_F, i_{\mathbb{N}^*}|_{\mathbb{N}^F})$ is a covariant representation of $(C^*(G_F), \mathbb{N}^F, \alpha)$ in $C^*(\mathbb{Q}/\mathbb{Z}) \rtimes_\alpha \mathbb{N}^*$. Since i_F is injective, we can deduce from the main theorem of [**12**] (or by minor modifications to the argument in [**8**, §3]) that the corresponding homomorphism

$$i_F \times i_{\mathbb{N}^*}|_{\mathbb{N}^F} : C^*(G_F) \rtimes_\alpha \mathbb{N}^F \to C^*(\mathbb{Q}/\mathbb{Z}) \rtimes_\alpha \mathbb{N}^*$$

is also an injection. We use this injection to identify $C^*(G_F) \rtimes_\alpha \mathbb{N}^F$ with a subalgebra of $C^*(\mathbb{Q}/\mathbb{Z}) \rtimes_\alpha \mathbb{N}^*$.

The crossed product $C^*(\mathbb{Q}/\mathbb{Z}) \rtimes_\alpha \mathbb{N}^*$ is spanned by the elements of the form $i_A(\delta_r)i_{\mathbb{N}^*}(m)i_{\mathbb{N}^*}(n)^*$ [**8**, Lemma 3.2]. If F contains all of the prime factors of m, n and the denominator of r, then $i_A(\delta_r)i_{\mathbb{N}^*}(m)i_{\mathbb{N}^*}(n)^*$ lies in $C^*(G_F) \rtimes_\alpha \mathbb{N}^F$. Thus, $C^*(\mathbb{Q}/\mathbb{Z}) \rtimes_\alpha \mathbb{N}^*$ is the direct limit $\bigcup_F C^*(G_F) \rtimes_\alpha \mathbb{N}^F$ over increasing finite subsets F of the set $\mathcal{P}$ of prime numbers.

In the next section, we describe a composition series for $C^*(G_F) \rtimes_\alpha \mathbb{N}^F$ when F is a finite subset of $\mathcal{P}$, and identify the subquotients in terms of ordinary crossed products $C(X_S) \rtimes \mathbb{Z}^S$ associated to subsets S of F. The underlying space X_S is the group of units $\mathcal{U}(\mathbb{Z}_{F\setminus S})$ in the ring $\mathbb{Z}_{F\setminus S} := \prod_{p\in F\setminus S}\mathbb{Z}_p$; as an additive group, $\mathbb{Z}_{F\setminus S}$ is the dual group of $G_{F\setminus S}$. The action of a prime $q \in S$ on

$$C(\mathcal{U}(\mathbb{Z}_{F\setminus S})) \subset C(\mathbb{Z}_{F\setminus S}) \cong C^*(G_{F\setminus S}) \subset C^*(\mathbb{Q}/\mathbb{Z})$$

induced by α_q is multiplication by q on $\mathcal{U}(\mathbb{Z}_{F\setminus S})$ (see [**11**, Lemma 1.1]), which is an automorphism because q is a unit in $\mathbb{Z}_{F\setminus S}$. Thus, the action of $\mathbb{N}^S$ on $C(\mathcal{U}(\mathbb{Z}_{F\setminus S}))$ extends to an action σ of $\mathbb{Z}^S$ such that

$$\sigma_{(m_p)}(f)(x) = f\left(\left(\prod_{p\in S} p^{m_p}\right)^{-1} x\right) \quad \text{for } (m_p) \in \mathbb{N}^S.$$

As a matter of notation, we view a crossed product $A \rtimes_\beta G$ by an action of a group as the universal C^*-algebra generated by a copy of A and a unitary representation $i_G : G \to U(A \rtimes_\beta G)$ satisfying the covariance relation $\beta_s(a) = i_G(s)ai_G(s)^*$.

https://doi.org/10.1017/S0143385705000143 Published online by Cambridge University Press

2. *Finitely many primes*

The object of this section is to prove the following theorem. For the definitions of AT-algebra, real rank zero and stable rank one, see [**23**] and the references therein.

THEOREM 2.1. *Let F be a finite set of primes. Then there is a composition series $\{I_k \mid 1 \le k \le |F|\}$ of ideals in $C^*(G_F) \rtimes_\alpha \mathbb{N}^F$ such that:*

(a) $I_1 \cong C(\mathcal{U}(\mathbb{Z}_F), \mathcal{K}(l^2(\mathbb{N}^F)))$;

(b) $I_{k+1}/I_k \cong \bigoplus_{S\subset F,\,|S|=k}(C(\mathcal{U}(\mathbb{Z}_{F\setminus S})) \rtimes_\sigma \mathbb{Z}^S) \otimes \mathcal{K}(l^2(\mathbb{N}^{F\setminus S}))$ *if* $|F| \ge 2$;

(c) $(C^*(G_F) \rtimes \mathbb{N}^F)/I_{|F|} \cong C(\mathbb{T}^F)$.

If $|F| \ge 2$, each $C(\mathcal{U}(\mathbb{Z}_{F\setminus S})) \rtimes_\sigma \mathbb{Z}^S$ is a finite direct sum of simple AT-algebras with real rank zero and a unique tracial state.

The proof of the theorem will occupy the rest of the section. We need some notation and a number of preliminary results.

Under the Fourier transform $C^*(G_F) \cong C(\mathbb{Z}_F)$ the action α becomes

$$\alpha_{(n_p)}(f)(x) = \begin{cases} f\left(\left(\prod_{p\in F} p^{n_p}\right)^{-1} x\right) & \text{if } x \in \left(\prod_{p\in F} p^{n_p}\right)\mathbb{Z}_F \\ 0 & \text{otherwise} \end{cases}$$

(see [**11**, Lemma 1.1]). For $S \subset F$, we set $\mathcal{Z}_S := \{a \in \mathbb{Z}_F \mid a_p = 0 \text{ for } p \in S\}$, and we write $\mathcal{Z}_p$ for $\mathcal{Z}_{\{p\}}$. The next lemma identifies $C_0(\mathbb{Z}_F \setminus \mathcal{Z}_S)$ as the kind of ideal for which taking crossed products behaves well (see [**10**]).

LEMMA 2.2. *For $S \subset F$, $C_0(\mathbb{Z}_F \setminus \mathcal{Z}_S)$ is an extendibly invariant ideal in $C(\mathbb{Z}_F)$.*

Proof. It suffices by [**10**, Theorem 4.3] to show that for each $n \in \mathbb{N}^F$, the endomorphism $x \mapsto \left(\prod_{p\in F} p^{n_p}\right)x$ of $\mathbb{Z}_F$ leaves both $\mathcal{Z}_S$ and $\mathbb{Z}_F\setminus\mathcal{Z}_S$ invariant. Certainly $\left(\prod_{p\in F} p^{n_p}\right)\mathcal{Z}_S$ is contained in $\mathcal{Z}_S$. If $x \notin \mathcal{Z}_S$, then $x_r \ne 0$ for some $r \in S$, $\prod_{p\in F} p^{n_p} x_r \ne 0$ for this r, and $\left(\prod_{p\in F} p^{n_p}\right)x \notin \mathcal{Z}_S$. □

Theorem 1.7 of [**10**] now allows us to identify $C_0(\mathbb{Z}_F \setminus \mathcal{Z}_S) \rtimes \mathbb{N}^F$ with an ideal J_S in $C(\mathbb{Z}_F) \rtimes_\alpha \mathbb{N}^F$ such that $(C(\mathbb{Z}_F) \rtimes_\alpha \mathbb{N}^F)/J_S = C(\mathcal{Z}_S) \rtimes \mathbb{N}^F$; we write J_p for $J_{\{p\}}$.

LEMMA 2.3. *We have $J_S = \sum_{p\in S} J_p$.*

Proof. Since $\mathcal{Z}_S = \bigcap_{p\in S} \mathcal{Z}_p$, we have $\mathbb{Z}_F \setminus \mathcal{Z}_S = \bigcup_{p\in S} \mathbb{Z}_F \setminus \mathcal{Z}_p$, and $C_0(\mathbb{Z}_F \setminus \mathcal{Z}_S) = \sum_{p\in S} C_0(\mathbb{Z}_F \setminus \mathcal{Z}_p)$. It follows from [**10**, Lemma 1.3] that if I, J and $I + J$ are extendibly invariant ideals in (A, P), then $(I + J) \rtimes P = (I \rtimes P) + (J \rtimes P)$. Thus, the result follows from Lemma 2.2. □

For $1 \le k \le |F|$, we define

$$I_k := \prod_{S\subset F,\,|S|=k} J_S = \bigcap_{S\subset F,\,|S|=k} J_S. \tag{2.1}$$

It follows from [**10**, Lemma 1.3] that if I and J are extendibly invariant ideals in (A, P), then

$$(I \rtimes P)(J \rtimes P) = (IJ) \rtimes P,$$

https://doi.org/10.1017/S0143385705000143 Published online by Cambridge University Press

and hence $I_k = C_0\big(\bigcap_{S\subset F,\,|S|=k}(\mathbb{Z}_F \setminus \mathcal{Z}_S)\big) \rtimes \mathbb{N}^F$. Therefore,

$$I_1 = C_0\Big(\bigcap_{p\in F}(\mathbb{Z}_F \setminus \mathcal{Z}_p)\Big) \rtimes \mathbb{N}^F = C_0\Big(\prod_{p\in F}(\mathbb{Z}_p \setminus \{0\})\Big) \rtimes \mathbb{N}^F;$$

since $\mathbb{Z}_p \setminus \{0\}$ is homeomorphic to $\mathcal{U}(\mathbb{Z}_p) \times \mathbb{N}$ by [**11**, Lemma 2.3], Theorem 2.1(a) follows from an argument similar to that in the last paragraph of [**11**, p. 176]. Similarly, we can prove Theorem 2.1(c) by following the proof of (2.4) of [**11**], because $(C^*(G_F) \rtimes_\alpha \mathbb{N}^F)/I_{|F|} = \mathbb{C} \rtimes \mathbb{N}^F$.

To prove Theorem 2.1(b), we need some lemmas. The first contains some general facts about families of ideals in C^*-algebras.

LEMMA 2.4. *Suppose that $I_1, \ldots, I_n$ are ideals in a C^*-algebra B.*

(a) *With $F_n = \{1, \ldots, n\}$, we have*

$$\prod_{S\subset F_n,\,|S|=k}\Big(\sum_{i\in S} I_i\Big) = \sum_{R\subset F_n,\,|R|=n-k+1}\Big(\prod_{j\in R} I_j\Big) \quad \text{for } 1 \le k \le n. \tag{2.2}$$

(b) *Suppose that K is an ideal such that $I_i I_j \subset K$ for all i, j. Then $\big(\sum_{i=1}^n I_i\big)/K$ is naturally isomorphic to $\bigoplus_{i=1}^n (I_i/I_i \cap K)$.*

Proof. We prove (a) by induction on n. The statement is trivial for $n = 1, 2$. Suppose that it holds for $n-1$. When $k = 1$, both sides of (2.2) are $\prod_{i=1}^n I_i$, so we assume $k \ge 2$. Writing the left-hand side (LHS) of (2.2) as $\big(\prod_{n\in S}\big)\big(\prod_{n\notin S}\big)$ and applying the inductive hypothesis to F_{n-1} shows that

$$\text{LHS} = \Big(\prod_{|S|=k,\,n\in S}\Big(I_n + \sum_{i\in S\setminus\{n\}} I_i\Big)\Big)\Big(\sum_{R\subset F_{n-1},\,|R|=n-k}\Big(\prod_{j\in R} I_j\Big)\Big). \tag{2.3}$$

As I_n is an ideal and $I_n^2 = I_n$, the first term of (2.3) simplifies to give

$$\text{LHS} = \Big(I_n + \prod_{S'\subset F_{n-1},\,|S'|=k-1}\Big(\sum_{i\in S'} I_i\Big)\Big)\Big(\sum_{R\subset F_{n-1},\,|R|=n-k}\Big(\prod_{j\in R} I_j\Big)\Big).$$

We can use the inductive hypothesis on F_{n-1} with k replaced by $k-1$ to get

$$\text{LHS} = \Big(I_n + \sum_{R'\subset F_{n-1},\,|R'|=n-k+1}\Big(\prod_{j\in R'} I_j\Big)\Big)\Big(\sum_{R\subset F_{n-1},\,|R|=n-k}\Big(\prod_{j\in R} I_j\Big)\Big), \tag{2.4}$$

which is contained in

$$\sum_{R\subset F_{n-1},\,|R|=n-k}\Big(\prod_{j\in R\cup\{n\}} I_j\Big) + \sum_{R'\subset F_{n-1},\,|R'|=n-k+1}\Big(\prod_{j\in R'} I_j\Big). \tag{2.5}$$

Since (2.5) is the same as the right-hand side (RHS) of (2.2), LHS $\subset$ RHS. On the other hand, every element of every $\prod_{j\in R'} I_j$ arises in (2.4) because we can pick $R \subset R'$, so RHS $\subset$ LHS.

To prove (b), note that the map $\phi_i : a + I_i \cap K \mapsto a + K$ is an injection of $I_i/(I_i \cap K)$ into $\big(\sum_{i=1}^n I_i\big)/K$, and

$$\phi_i(a + I_i \cap K)\phi_j(b + I_j \cap K) = ab + K = 0 \quad \text{for } i \ne j$$

because $ab \in I_i I_j \subset K$. So the ϕ_j combine to give an injection ϕ of $\bigoplus(I_i/I_i \cap K)$ into $\big(\sum_{i=1}^n I_i\big)/K$, which is clearly surjective. □

https://doi.org/10.1017/S0143385705000143 Published online by Cambridge University Press

LEMMA 2.5. *The ideals I_k of $C^*(G_F) \rtimes_\alpha \mathbb{N}^F$ defined in (2.1) satisfy*

$$I_{k+1}/I_k = \bigoplus_{S\subset F,\,|S|=k} \Big(\bigcap_{p\notin S} J_{S\cup\{p\}}\Big)\Big/ J_S.$$

Proof. Lemma 2.4(a) gives $I_{k+1} = \sum_{R\subset F,\,|R|=n-k}(\prod_{p\in R} J_p)$. The product of any two ideals $\prod_{p\in R} J_p$ with $|R| = n-k$ has at least $n-k+1$ factors J_p, and, hence, is contained in $I_k = \sum_{R\subset F,\,|R|=n-k+1}(\prod_{p\in R} J_p)$. Thus, Lemma 2.4(b) gives

$$I_{k+1}/I_k = \bigoplus_{R\subset F,\,|R|=n-k} \frac{\prod_{p\in R} J_p}{I_k \cap \big(\prod_{p\in R} J_p\big)}. \tag{2.6}$$

Now

$$I_k \cap \Big(\prod_{p\in R} J_p\Big) = \sum_{|T|=n-k+1} \Big(\prod_{q\in T} J_q\Big)\Big(\prod_{p\in R} J_p\Big);$$

each of these summands has at least one factor J_q for $q \notin R$, and is then contained in $J_q\big(\prod_{p\in R} J_p\big)$. Using $I\cap J = IJ$ again gives

$$I_k \cap \Big(\prod_{p\in R} J_p\Big) = \sum_{q\notin R} J_q\Big(\prod_{p\in R} J_p\Big) = \Big(\sum_{q\notin R} J_q\Big)\Big(\prod_{p\in R} J_p\Big),$$

and using the isomorphism $(I+J)/I = J/(I\cap J)$ and Lemma 2.3 gives

$$\frac{\prod_{p\in R} J_p}{I_k \cap \big(\prod_{p\in R} J_p\big)} = \frac{J_{F\setminus R} + \big(\prod_{p\in R} J_p\big)}{J_{F\setminus R}}.$$

Finally we observe that

$$J_{F\setminus R} + \Big(\prod_{p\in R} J_p\Big) = \prod_{p\in R}(J_{F\setminus R} + J_p) = \prod_{p\in R} J_{(F\setminus R)\cup\{p\}}$$

and write $S = F\setminus R$ to deduce the result. □

LEMMA 2.6. *The ideals J_S in $C^*(G_F) \rtimes_\alpha \mathbb{N}^F$ satisfy*

$$\Big(\bigcap_{p\in F\setminus S} J_{S\cup\{p\}}\Big)\Big/ J_S \cong (C(\mathcal{U}(\mathbb{Z}_{F\setminus S})) \rtimes_\sigma \mathbb{Z}^S) \otimes \mathcal{K}(l^2(\mathbb{N}^{F\setminus S})).$$

Proof. We first realize $\big(\bigcap_{p\in F\setminus S} J_{S\cup\{p\}}\big)\big/ J_S$ as a semigroup crossed product:

$$\begin{aligned}\bigcap_{p\in F\setminus S} J_{S\cup\{p\}} &= C_0\Big(\bigcap_{p\in F\setminus S}(\mathbb{Z}_F \setminus \mathcal{Z}_{S\cup\{p\}})\Big) \rtimes \mathbb{N}^F \\ &= C_0\Big(\mathbb{Z}_F \setminus \Big(\bigcup_{p\in F\setminus S} \mathcal{Z}_{S\cup\{p\}}\Big)\Big) \rtimes \mathbb{N}^F.\end{aligned}$$

Thus,

$$\begin{aligned}\Big(\bigcap_{p\in F\setminus S} J_{S\cup\{p\}}\Big)\Big/ J_S &= C_0\Big(\mathcal{Z}_S \setminus \Big(\bigcup_{p\in F\setminus S} \mathcal{Z}_{S\cup\{p\}}\Big)\Big) \rtimes \mathbb{N}^F \\ &= C_0\Big(\Big(\prod_{p\in F\setminus S} \mathbb{Z}_p \setminus \{0\}\Big) \times \Big(\prod_{p\in S}\{0\}\Big)\Big) \rtimes \mathbb{N}^F.\end{aligned}$$

The arguments of Corollary 2.4 and Lemma 2.5 of [**11**] show that this last crossed product is isomorphic to $(C(\mathcal{U}(\mathbb{Z}_{F\setminus S})) \rtimes_\sigma \mathbb{Z}^S) \otimes \mathcal{K}(l^2(\mathbb{N}^{F\setminus S}))$. □

https://doi.org/10.1017/S0143385705000143 Published online by Cambridge University Press

Theorem 2.1(b) follows immediately from Lemmas 2.5 and 2.6.

To finish the proof of Theorem 2.1, it remains to prove the statements about the structure of $C(\mathcal{U}(\mathbb{Z}_{F\setminus S}))\rtimes_\sigma \mathbb{Z}^S$. Corollary A.6 implies that $H := \overline{\mathbb{Z}^S}$ has finite index in $\mathcal{U}(\mathbb{Z}_{F\setminus S})$. The argument at the end of the proof of [**11**, Theorem 3.1] shows that $C(\mathcal{U}(\mathbb{Z}_{F\setminus S}))\rtimes_\sigma \mathbb{Z}^S$ is a finite direct sum of algebras isomorphic to $C(H)\rtimes_\sigma \mathbb{Z}^S$, which is simple because $\mathbb{Z}^S$ acts minimally and freely on H. Since H is an open and closed subset of $\mathcal{U}(\mathbb{Z}_{F\setminus S})$, it is totally disconnected and it follows from [**19**, Theorem 6.11] that $C(H)\rtimes_\sigma \mathbb{Z}^S$ has real rank zero and stable rank one.

The space $\mathcal{U}(\mathbb{Z}_{F\setminus S})$ is the inverse limit of the finite groups $\mathcal{U}\big(\mathbb{Z}\big/\big(\prod_{p\in F\setminus S} p^{l_p}\big)\mathbb{Z}\big)$ over $l=(l_p)\in\mathbb{N}^{F\setminus S}$. The diagonally embedded copy of $\mathbb{N}$ is cofinal in $\mathbb{N}^{F\setminus S}$ and, hence,

$$\mathcal{U}(\mathbb{Z}_{F\setminus S}) = \varprojlim \mathcal{U}\Bigg(\mathbb{Z}\Big/\Big(\prod_{p\in F\setminus S} p^n\Big)\mathbb{Z}\Bigg). \tag{2.7}$$

Let π_n denote the canonical surjection of $\mathcal{U}(\mathbb{Z}_{F\setminus S})$ onto $\mathcal{U}\big(\mathbb{Z}/\big(\prod_{p\in F\setminus S} p^n\big)\mathbb{Z}\big)$.

LEMMA 2.7. *Let $H_n := \pi_n(H)\subset \mathcal{U}\big(\mathbb{Z}/\big(\prod_{p\in F\setminus S} p^n\big)\mathbb{Z}\big)$ and let $\mathbb{Z}^S$ act on H_n via the embedding $(n_q)\mapsto \prod_{q\in S} q^{n_q}$ of $\mathbb{Z}^S$ in $\mathbb{Z}$. Then there are C*-subalgebras A_n of $C(H)\rtimes_\sigma \mathbb{Z}^S$ such that $A_n\cong C(H_n)\rtimes\mathbb{Z}^S$ and $C(H)\rtimes_\sigma\mathbb{Z}^S=\overline{\bigcup A_n}$.*

Proof. The homomorphism π_n induces an injection π_n^* of $C(H_n)$ into $C(H)$, and then $C(H)=\overline{\bigcup_{n\in\mathbb{N}}\pi_n^*(C(H_n))}$. On $\mathbb{Z}^S\subset H$, π_n is reduction modulo $\prod_{p\in F\setminus S}p^n$, so π_n^* converts the action σ into the canonical action of $\mathbb{Z}^S$ by multiplication on H_n. Thus, π_n^* induces a homomorphism $\pi_n^*\rtimes\mathrm{id}$ of $C(H_n)\rtimes\mathbb{Z}^S$ into $C(H)\rtimes_\sigma\mathbb{Z}^S$. The homomorphism π_n^* is faithful on $C(H_n)$ and intertwines the dual actions and, hence, a standard argument shows that $\pi_n^*\rtimes\mathrm{id}$ is faithful on $C(H_n)\rtimes\mathbb{Z}^S$ (see, for example, [**11**, Lemma 4.2]). Since $\bigcup_n\pi_n^*(C(H_n))$ is dense in $C(H)$, we therefore have

$$C(H)\rtimes_\sigma\mathbb{Z}^S=\overline{\bigcup_{n\in\mathbb{N}}\pi_n^*\rtimes\mathrm{id}(C(H_n)\rtimes\mathbb{Z}^S)},$$

as claimed. □

We can identify the subalgebras A_n explicitly.

PROPOSITION 2.8. *Let F be a finite quotient of $\mathbb{Z}^k$. Then $C(F)\rtimes\mathbb{Z}^k$ is isomorphic to $C(\mathbb{T}^k, M_{|F|}(\mathbb{C}))$.*

Proof. Let H be the subgroup of $\mathbb{Z}^k$ with $F=\mathbb{Z}^k/H$. Then H is itself a free abelian group of rank k, and hence has the form $A\mathbb{Z}^k$ for some $A\in M_k(\mathbb{Z})\cap GL_k(\mathbb{Q})$. The matrix A has a Smith normal form: there are matrices $P, Q\in GL_k(\mathbb{Z})$ such that $B:=P^{-1}AQ^{-1}$ is diagonal [**15**, §3.22]. Then $H=A\mathbb{Z}^k=PBQ\mathbb{Z}^k=PB\mathbb{Z}^k\cong B\mathbb{Z}^k=b_{11}\mathbb{Z}\oplus\cdots\oplus b_{kk}\mathbb{Z}$. In other words, multiplying by P^{-1} gives an automorphism of $\mathbb{Z}^k$ which carries H into $\bigoplus b_{ii}\mathbb{Z}$. Thus,

$$C(F)\rtimes\mathbb{Z}^k\cong C\Bigg(\prod_{i=1}^k(\mathbb{Z}/b_{ii}\mathbb{Z})\Bigg)\rtimes\mathbb{Z}^k\cong\bigotimes_{i=1}^k\Big(C(\mathbb{Z}/b_{ii}\mathbb{Z})\rtimes_\tau\mathbb{Z}\Big),$$

where τ is the action of $\mathbb{Z}$ by translation.

https://doi.org/10.1017/S0143385705000143 Published online by Cambridge University Press

By [**17**, Corollary 2.5], $C(\mathbb{Z}/b\mathbb{Z}) \rtimes_\tau \mathbb{Z}$ is isomorphic to the induced algebra

$$\mathrm{Ind}_{(b\mathbb{Z})^\perp}^{\mathbb{T}}(C(\mathbb{Z}/b\mathbb{Z}) \rtimes_\tau (\mathbb{Z}/b\mathbb{Z}), \widehat{\tau}),$$

which is described in terms of a generator β of the dual action $\widehat{\tau}$ as the mapping torus

$$MT(\beta) = \{f : [0,1] \to C(\mathbb{Z}/b\mathbb{Z}) \rtimes_\tau (\mathbb{Z}/b\mathbb{Z}) \mid f(1) = \beta(f(0))\}. \tag{2.8}$$

Since $C(\mathbb{Z}/b\mathbb{Z}) \rtimes_\tau (\mathbb{Z}/b\mathbb{Z}) \cong B(l^2(\mathbb{Z}/b\mathbb{Z})) = M_{|b|}(\mathbb{C})$, the automorphism β is inner and there is a continuous path β_t in $\mathrm{Aut}\, M_{|b|}(\mathbb{C})$ such that $\beta_0 = \mathrm{id}$ and $\beta_1 = \beta$. Now $\phi(f)(t) = \beta_t^{-1}(f(t))$ defines an isomorphism ϕ of (2.8) onto $C(\mathbb{T}, M_{|b|}(\mathbb{C}))$. We therefore have

$$C(F) \rtimes \mathbb{Z}^k \cong \bigotimes_{i=1}^{k} C(\mathbb{T}, M_{|b_{ii}|}(\mathbb{C})) \cong C(\mathbb{T}^k, M_{\prod_i |b_{ii}|}(\mathbb{C})),$$

and the result follows on observing that $\prod_i |b_{ii}| = |\det B| = |\det A| = |F|$. □

It follows from Proposition 2.8 and the decomposition $C(H) \rtimes_\sigma \mathbb{Z}^S = \overline{\bigcup A_n}$ that $C(H) \rtimes_\sigma \mathbb{Z}^S$ is an AH-algebra†, see [**23**]. The K-theory of $C(H_n) \rtimes \mathbb{Z}^S$ is torsion-free and this property is preserved under inductive limits, so $C(H) \rtimes_\sigma \mathbb{Z}^S$ has torsion-free K-theory. Since $C(H) \rtimes_\sigma \mathbb{Z}^S$ is a simple AH-algebra with real rank zero and no dimension growth, it follows from [**20**, Lemma 7.5], using the results of Elliott [**4**, Theorems 8.3 and 4.3] and Lin [**13**, Proposition 2.6, **14**, Theorem 5.2], that it is an AT-algebra.

We also use the decomposition $C(H) \rtimes_\sigma \mathbb{Z}^S = \overline{\bigcup A_n}$ to prove that $C(H) \rtimes_\sigma \mathbb{Z}^S$ has a unique tracial state. Let μ denote the Haar measure on $H \subset \mathcal{U}(\mathbb{Z}_{F\setminus S})$. The action σ permutes the cylinder sets $\{\pi_n^{-1}(m) \mid m \in H_n\}$, so every invariant probability measure agrees with μ on cylinder sets. Since the characteristic functions of such sets span a dense subspace of $C(H)$, it follows that μ is the only invariant probability measure and $C(H) \rtimes \mathbb{Z}^S$ has a unique tracial state by [**3**, Corollary VIII.3.8].

This completes the proof of Theorem 2.1.

3. *The structure of* $C(\mathcal{U}(\mathbb{Z}_{F\setminus S})) \rtimes_\sigma \mathbb{Z}^S$

3.1. *When S contains just one prime.* We consider $C(\mathcal{U}(\mathbb{Z}_{F\setminus S})) \rtimes_\sigma \mathbb{Z}^S$ when $|F| \geq 2$ and $S = \{q\}$. To simplify the notation, we relabel $F \setminus \{q\}$ as F. The following result generalizes [**11**, Theorem 3.1] in two directions: to sets F with $|F| > 1$ and to sets F containing the even prime 2. If $\mathbf{l} = (l_p) \in \mathbb{N}^F$ is a multi-index, we write $o_{\mathbf{l}}(q)$ for the order of q in $\prod_{p\in F} \mathcal{U}(\mathbb{Z}/p^{l_p}\mathbb{Z})$.

THEOREM 3.1. *Suppose that F is a finite set of primes and q is a prime which does not belong to F. Then there are a multi-index $\mathbf{K} = (K_p) \in \mathbb{N}^F$ and $d \in \mathbb{N}$ such that*

$$o_{\mathbf{K}+\mathbf{l}}(q) = d\left(\prod_{p\in F} p^{l_p}\right) \quad \textit{for every } \mathbf{l} \in \mathbb{N}^F, \tag{3.1}$$

† To see that an inductive limit $\overline{\bigcup A_n}$ is an AH-algebra, it suffices to show that each A_n is a corner in a matrix algebra $M_N(C(X))$, or, equivalently, that A_n is a homogeneous algebra with vanishing Dixmier–Douady class. Since the Dixmier–Douady class $\delta(A)$ of an m-homogeneous algebra satisfies $m\delta(A) = 0$ and $H^3(\mathbb{T}^k, \mathbb{Z})$ has no torsion, it suffices to prove that each A_n is a homogeneous algebra with spectrum $\mathbb{T}^k$. In our situation we could prove this in several ways. However, Proposition 2.8 makes the stronger statement that A_n is isomorphic to $M_m(C(\mathbb{T}^k))$.

https://doi.org/10.1017/S0143385705000143 Published online by Cambridge University Press

and $C(\mathcal{U}(\mathbb{Z}_F)) \rtimes_\sigma \mathbb{Z}$ is the direct sum of $\left(\prod_{p\in F}(p-1)p^{K_p-1}\right)/d$ copies of a Bunce–Deddens algebra with supernatural number $d\left(\prod_{p\in F} p^\infty\right)$.

The existence of $\mathbf{K}$ and d satisfying (3.1) is established in Proposition A.5. We saw in §2 that $C(\mathcal{U}(\mathbb{Z}_F)) \rtimes_\sigma \mathbb{Z}$ is the direct sum of copies of the simple algebra $C(H) \rtimes_\sigma \mathbb{Z}$, where H is the closure of $q^{\mathbb{Z}}$ in $\mathcal{U}(\mathbb{Z}_F)$. It remains to prove that $C(H) \rtimes_\sigma \mathbb{Z}$ is a Bunce–Deddens algebra and to calculate the index $|\mathcal{U}(\mathbb{Z}_F) : H|$, which is the number of simple direct summands.

Let $\{n_k\}$ be integers with $n_k \geq 2$, and let $X_k = \{0, 1, \ldots, n_k - 1\}$. Addition by 1 with carry over is a homeomorphism of the totally disconnected space $X := \prod_{k\geq 0} X_k$ called an odometer action, and the resulting crossed product $C(X) \rtimes_\tau \mathbb{Z}$ is a *Bunce–Deddens algebra with supernatural number* $\mathbf{n} := \prod_{k\geq 0} n_k$ (see [**3**, Ch. VIII.4]).

Our claim that $C(H) \rtimes_\sigma \mathbb{Z}$ is a Bunce–Deddens algebra will follow from the next proposition, which generalizes [**11**, Proposition 3.6].

PROPOSITION 3.2. *Suppose $\{G_l \mid \in \mathbb{N}\}$ are finite groups and $G = \varprojlim(G_l, \pi_l)$. Fix $g \in G$ and let L denote the closed subgroup of G generated by g. Consider the action $\sigma : \mathbb{Z} \to \operatorname{Aut} C(G)$ such that $\sigma_n(f)(x) = f(g^{-n}x)$. Let $o_l(g)$ denote the order of $\pi_l(g)$ in G_l and let*

$$d_l := \begin{cases} o_0(g) & \text{if } l = 0 \\ o_l(g)/o_{l-1}(g) & \text{if } l \geq 1. \end{cases} \tag{3.2}$$

Then $C(L) \rtimes_\sigma \mathbb{Z}$ is a Bunce–Deddens algebra with supernatural number $\prod_{l\geq 0} d_l$.

Proof. Let $X := \prod_{l\geq 0}\{0, 1, \ldots, d_l - 1\}$. The argument in the proof of [**11**, Proposition 3.6] shows that the continuous maps $h_l : X \to G_l$ given by

$$h_l(\{a_n\}) = \pi_l(g^{a_0 + a_1 d_0 + \cdots + a_l d_0 d_1 \ldots d_{l-1}}) \tag{3.3}$$

combine to give an equivariant homeomorphism $h : X \to L$, which induces the required isomorphism. □

Our subgroup H of $\mathcal{U}(\mathbb{Z}_F)$ is the inverse limit $\varprojlim \pi_l(H)$, where $\pi_l : \mathcal{U}(\mathbb{Z}_F) \to \mathcal{U}\left(\mathbb{Z}/\left(\prod_{p\in F} p^{K_p+l}\right)\mathbb{Z}\right)$ is the canonical surjection. Then Proposition 3.2 and (3.1) imply that $C(H) \rtimes_\sigma \mathbb{Z}$ is a Bunce–Deddens algebra with supernatural number $d(\prod_{p\in F} p)^\infty$ for $d = o_{\mathbf{K}}(q)$. By Corollary A.6, we have that

$$|\mathcal{U}(\mathbb{Z}_F) : H| = \left(\prod_{p\in F}(p-1)p^{K_p-1}\right)\Big/ d, \tag{3.4}$$

which finishes the proof of Theorem 3.1.

3.2. *When S consists of two primes.* We now analyse $C(\mathcal{U}(\mathbb{Z}_{F\setminus S})) \rtimes \mathbb{Z}^S$ when $S = \{q, r\}$. For simplicity, we consider only the case $F = \{p, q, r\}$, so that we are interested in the crossed product $C(\mathcal{U}(\mathbb{Z}_p)) \rtimes_\sigma \mathbb{Z}^2$, where

$$\sigma_{m,n}(f)(x) = f(q^{-m}r^{-n}x).$$

https://doi.org/10.1017/S0143385705000143 Published online by Cambridge University Press

THEOREM 3.3. *The C^*-algebra $C(\mathcal{U}(\mathbb{Z}_p)) \rtimes_\sigma \mathbb{Z}^2$ is a finite direct sum of copies of a simple AT-algebra A which has real rank zero, a unique tracial state and K-theory satisfying two short exact sequences:*

$$\begin{aligned} 0 &\longrightarrow \mathbb{Z}[p^{-1}] \longrightarrow K_0(A) \longrightarrow \mathbb{Z} \longrightarrow 0 \\ 0 &\longrightarrow \mathbb{Z} \longrightarrow K_1(A) \longrightarrow \mathbb{Z}[p^{-1}] \longrightarrow 0. \end{aligned} \tag{3.5}$$

Everything except the assertion about K-theory was proved in Theorem 2.1; the simple C^*-algebra A is $C(H) \rtimes_\sigma \mathbb{Z}^2$, where H is the closure of $q^{\mathbb{Z}} r^{\mathbb{Z}}$ in $\mathcal{U}(\mathbb{Z}_p)$. We aim to analyse $C(H) \rtimes_\sigma \mathbb{Z}^2$ by writing it as an iterated crossed product $(C(H) \rtimes_{\sigma^q} \mathbb{Z}) \rtimes \mathbb{Z}$. The inside crossed product is not simple unless $q^{\mathbb{Z}}$ is dense in H and it is helpful to reduce to this case using the following lemma.

LEMMA 3.4. *Let H_q denote the closure of $q^{\mathbb{Z}}$ in $\mathcal{U}(\mathbb{Z}_p)$. Then H_q has finite index $I(q)$ in H and, hence, is an open and closed subset of H. The inclusion of $C(H_q)$ in $C(H)$ induces an isomorphism of $C(H_q) \rtimes_\sigma (\mathbb{Z} \times I(q)\mathbb{Z})$ onto the corner $\chi_{H_q}(C(H) \rtimes_\sigma \mathbb{Z}^2)\chi_{H_q}$.*

Proof. Corollary A.6 implies that H_q has finite index in $\mathcal{U}(\mathbb{Z}_p)$, so it certainly has finite index in H. The inclusion of $C(H_q)$ in $C(H)$ and the map

$$(m, I(q)n) \mapsto \chi_{H_q} i_{\mathbb{Z}^2}(m, I(q)n)\chi_{H_q}$$

form a covariant representation of $(C(H_q), \mathbb{Z} \times I(q)\mathbb{Z}, \sigma)$ in $\chi_{H_q}(C(H) \rtimes_\sigma \mathbb{Z}^2)\chi_{H_q}$ and, hence, give a homomorphism

$$\phi : C(H_q) \rtimes_\sigma (\mathbb{Z} \times I(q)\mathbb{Z}) \to \chi_{H_q}(C(H) \rtimes_\sigma \mathbb{Z}^2)\chi_{H_q}.$$

We can identify $(\mathbb{Z} \times I(q)\mathbb{Z})^\wedge$ with $\mathbb{T}^2/(\mathbb{Z} \times I(q)\mathbb{Z})^\perp = \mathbb{T}^2/(1 \times C_{I(q)})$, where C_n denotes the group of nth roots of unity, and then ϕ carries the dual action $\hat{\sigma}_{[w,z]}$ into $\hat{\sigma}_{w,z}$; now a standard argument implies that ϕ is injective (or we could apply [**22**, Corollary 4.3], for example). We have

$$\chi_{H_q}(f i_{\mathbb{Z}^2}(m,n))\chi_{H_q} = (f\chi_{H_q}) i_{\mathbb{Z}^2}(m,n)\chi_{H_q} = i_{\mathbb{Z}^2}(m,n)\sigma^{-1}_{m,n}(f\chi_{H_q})\chi_{H_q}.$$

Since the support of $\sigma^{-1}_{m,n}(f\chi_{H_q})$ is contained in $q^{-m}r^{-n}H_q = r^{-n}H_q$, we have

$$\sigma^{-1}_{m,n}(f\chi_{H_q})\chi_{H_q} = \begin{cases} \sigma^{-1}_{m,n}(f\chi_{H_q}) & \text{if } I(q) \text{ divides } n \\ 0 & \text{otherwise,} \end{cases}$$

and

$$\begin{aligned} \chi_{H_q}(f i_{\mathbb{Z}^2}(m,n))\chi_{H_q} &= \begin{cases} i_{\mathbb{Z}^2}(m,n)\sigma^{-1}_{m,n}(f\chi_{H_q}) & \text{if } I(q) \text{ divides } n \\ 0 & \text{otherwise} \end{cases} \\ &= \begin{cases} \phi((f\chi_{H_q}) i_{\mathbb{Z}^2}(m,n)) & \text{if } I(q) \text{ divides } n \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Thus, every $\chi_{H_q}(f i_{\mathbb{Z}^2}(m,n))\chi_{H_q}$ is in the range of ϕ and ϕ is surjective. □

COROLLARY 3.5. *Define $\gamma : \mathbb{Z} \to \mathrm{Aut}(C(H_q) \rtimes_{\sigma^q} \mathbb{Z})$ by*

$$\gamma_m(f i_{\mathbb{Z}}(n)) = \sigma^r_{I(q)m}(f) i_{\mathbb{Z}}(n). \tag{3.6}$$

Then $(C(H_q) \rtimes_{\sigma^q} \mathbb{Z}) \rtimes_\gamma \mathbb{Z}$ is isomorphic to a full corner in $C(H) \rtimes_\sigma \mathbb{Z}^2$.

https://doi.org/10.1017/S0143385705000143 Published online by Cambridge University Press

Proof. Theorem 4.1 of [**18**] gives $C(H_q) \rtimes_\sigma (\mathbb{Z} \times I(q)\mathbb{Z}) \cong (C(H_q) \rtimes_{\sigma^q} \mathbb{Z}) \rtimes I(q)\mathbb{Z}$, so the result follows from Lemma 3.4 on replacing $I(q)\mathbb{Z}$ by the isomorphic group $\mathbb{Z}$. □

The analysis in §3.1 shows that $C(H_q) \rtimes_{\sigma^q} \mathbb{Z}$ is a Bunce–Deddens algebra. The K-theory of Bunce–Deddens algebras is well known. To state the version we need, recall that if $\mathbf{n} = (n_k)_{k\geq 0}$ is a sequence with $n_k \geq 2$, then $\mathbb{Z}[\mathbf{n}^{-1}]$ denotes the set of rational numbers with denominator $N_k = \prod_{i=0}^{k} n_i$ for some $k \geq 0$.

PROPOSITION 3.6. *Suppose that* $\mathbf{n} = (n_k)_{k\geq 0}$, $X_k = \{0, \ldots, n_k - 1\}$, $X = \prod X_k$ *and* $\tau : \mathbb{Z} \to \operatorname{Aut} C(X)$ *is the associated odometer. Then there are isomorphisms* $\phi_0 : K_0(C(X) \rtimes_\tau \mathbb{Z}) \to \mathbb{Z}[\mathbf{n}^{-1}]$ *such that* $\phi_0([\chi_{J(a_0,\ldots,a_k)}]) = N_k^{-1}$ *for each cylinder set* $J(a_0, \ldots, a_k)$ *and* $\phi_1 : K_1(C(X) \rtimes_\tau \mathbb{Z}) \to \mathbb{Z}$ *such that* $\phi_1(i_{\mathbb{Z}}(1)) = 1$.

Proof. As $K_1(C(X)) = 0$, the Pimsner–Voiculescu sequence for the system $(C(X), \mathbb{Z}, \tau)$ reduces to

$$0 \longrightarrow K_1(C(X) \rtimes_\tau \mathbb{Z}) \xrightarrow{\delta} K_0(C(X)) \xrightarrow{\mathrm{id}-\tau_*} K_0(C(X)) \xrightarrow{\mathrm{id}_*} K_0(C(X) \rtimes_\tau \mathbb{Z}) \longrightarrow 0.$$

Now let $C_k = \{J(a_0, \ldots, a_k)\}$ be the set of cylinder sets of length $k+1$ and note that $C(X) = \overline{\bigcup_{k=1} A_k}$, where $A_k = \operatorname{span}\{\chi_J \mid J \in C_k\}$. Each χ_J for $J \in C_k$ is the sum of n_{k+1} basis elements of A_{k+1}, so the maps $[\chi_{J(a_0,\ldots,a_k)}] \mapsto {N_k}^{-1}$ of A_k into $\mathbb{R}$ combine to give a homomorphism ψ_0 of $K_0(C(X)) = \varinjlim K_0(A_k)$ into $\mathbb{R}$ with range $\mathbb{Z}[\mathbf{n}^{-1}]$. Since the generating automorphism $\tau = \tau_1$ permutes C_k, the kernel of ψ_0 is the range of $\mathrm{id} - \tau_*$ and, hence, ψ_0 induces the required isomorphism ϕ_0 of $K_0(C(X) \rtimes_\phi \mathbb{Z})$ onto $\mathbb{Z}[\mathbf{n}^{-1}]$. To verify the statement about K_1, recall that δ is the coboundary map for the Toeplitz extension of $C(X) \rtimes_\tau \mathbb{Z}$ (see [**21**, §2]) and compute the index of $[i_{\mathbb{Z}}(1)]$ in $K_0(C(X) \otimes \mathcal{K}) \cong K_0(C(X))$. □

Proof of Theorem 3.3. We saw in the proof of Proposition 3.2 and in the paragraph following it that the homeomorphism h of $\prod_{k\geq 0} X_k$ onto the subgroup H_q of $\mathcal{U}(\mathbb{Z}_p)$ satisfies

$$\pi_k(h(\{a_n\})) = \pi_k(q^{a_0 + a_1 o_p(q) + \cdots + a_k o_p(q) p^{k-1}}) \quad \text{for } k \geq 0,$$

and hence carries $J(a_0, \ldots, a_k)$ onto $Z(q^{a_0 + a_1 o_p(q) + \cdots + a_k o_p(q) p^{k-1}})$, where

$$Z_k(n) = \{x \in \mathcal{U}(\mathbb{Z}_p) \mid \pi_k(x) = \pi_k(n)\}.$$

So we deduce from Proposition 3.6 that there is an isomorphism ϕ_0 of $K_0(C(H_q) \rtimes_\sigma \mathbb{Z})$ onto $(1/o_p(q))\mathbb{Z}[p^{-1}]$ such that

$$\phi_0([\chi_{Z_k(m)}]) = \frac{1}{o_p(q)} \frac{1}{p^k}$$

for every integer m which lies in H_q.

Multiplying by the unit $r^{-I(q)l}$ carries $Z_k(m)$ into $Z_k(r^{-I(q)l}m)$ and, hence, $\phi_0 \circ (\gamma_l)_* = \phi_0$. Thus, $(\gamma_l)_*$ is the identity on $K_0(C(H_q) \rtimes_\sigma \mathbb{Z})$. It is also the identity on $K_1(C(H_q) \rtimes_\sigma \mathbb{Z})$ and, hence, the Pimsner–Voiculescu sequence for $((C(H_q) \rtimes_\sigma \mathbb{Z}), \mathbb{Z}, \gamma)$

https://doi.org/10.1017/S0143385705000143 Published online by Cambridge University Press

collapses to the two short exact sequences

$$0 \longrightarrow \frac{1}{o_p(q)}\mathbb{Z}[p^{-1}] \longrightarrow K_0(C(H_q) \rtimes \mathbb{Z}^2) \longrightarrow \mathbb{Z} \longrightarrow 0$$
$$0 \longrightarrow \mathbb{Z} \longrightarrow K_1(C(H_q) \rtimes \mathbb{Z}^2) \longrightarrow \frac{1}{o_p(q)}\mathbb{Z}[p^{-1}] \longrightarrow 0.$$

From this and Corollary 3.5 we can deduce (3.5); since the isomorphism induced by Corollary 3.5 scales the class of [1], we have removed the factor $o_p(q)^{-1}$ by a further scaling to ensure that the final statement does not depend on the order of factors in our decomposition. □

Remark 3.7. The number of simple summands in Theorem 3.3 is $|\mathcal{U}(\mathbb{Z}_p) : H|$ and we can compute this using [**11**, Lemma 3.7]. For example, if p is odd and l is large, we have from (A.1) that

$$\begin{aligned}|\pi_l(H)| &= [o_{p^l}(q), o_{p^l}(r)] = [p^{l-L_p(q)}o_p(q), q^{l-L_p(r)}o_p(r)] \\ &= p^{l-\min(L_p(q),L_p(r))}[o_p(q), o_p(r)];\end{aligned}$$

thus we deduce

$$|\mathcal{U}(\mathbb{Z}_p) : H| = |\mathcal{U}(\mathbb{Z}/p^l\mathbb{Z}) : \pi_l(H)| = \frac{(p-1)p^{\min(L_p(q),L_p(r))-1}}{[o_p(q), o_p(r)]}.$$

We could carry out a similar analysis when $|F| > 1$, although it would not be so easy to work out some of the indices explicitly.

Remark 3.8. Theorem 2.1 implies, in particular, that $C(H) \rtimes_\sigma \mathbb{Z}^2$ satisfies the hypotheses of the classification theorem of Elliott for AT-algebras [**23**, Theorem 3.2.6]. We can tell from the computation of K-theory in Theorem 3.3 that $C(H) \rtimes_\sigma \mathbb{Z}^2$ is not a Bunce–Deddens algebra, but it is still closely related to an odometer. The homeomorphism of $\prod_{k\geq 0} X_k$ onto H_q identifies the action of the first copy of $\mathbb{Z}$ (multiplication by q on H_q) with an odometer (addition of 1 with carry over). The action of the second copy of $\mathbb{Z}$ (multiplication by r on H_q) also acts as a permutation on each X_k: it moves X_0 around in a different order and this action carries over into X_1 when the marker in X_0 returns to the starting point. So we can think of the action of $\mathbb{Z}^2$ as two independent odometers on the same set. We can normalize the scale so that either copy of $\mathbb{Z}$ acts by addition of 1 with carry over, but not so that both act this way at once.

4. *The Bost–Connes algebra*

The Hecke C^*-algebra $\mathcal{C}_\mathbb{Q}$ of Bost and Connes [**2**] is isomorphic to the semigroup crossed product $C^*(\mathbb{Q}/\mathbb{Z}) \rtimes_\alpha \mathbb{N}^*$. The Fourier transform takes $C^*(\mathbb{Q}/\mathbb{Z})$ onto the algebra of continuous functions on the compact group $\mathcal{Z} := \prod_{p\in\mathcal{P}} \mathbb{Z}_p$ and carries α into the action given by (see [**7**, §3.1])

$$\alpha_n(f)(x) = \begin{cases} f(x/n) & \text{if } n \text{ divides } x \\ 0 & \text{otherwise.} \end{cases}$$

https://doi.org/10.1017/S0143385705000143 Published online by Cambridge University Press

Lemma 2.2 is valid with F replaced by $\mathcal{P}$ and $\mathbb{Z}_F$ by $\mathcal{Z}$. Thus, for $S \subset \mathcal{P}$, an application of [**10**, Theorem 1.7] gives that $J_S := C_0(\mathcal{Z} \setminus \mathcal{Z}_S) \rtimes_\alpha \mathbb{N}^*$ is an ideal of $\mathcal{C}_\mathbb{Q} = C(\mathcal{Z}) \rtimes \mathbb{N}^*$, with quotient isomorphic to $C(\mathcal{Z}_S) \rtimes \mathbb{N}^*$. Choose $a \in \mathcal{Z}$ such that $a_p = 0 \iff p \in S$. Then $\{\mathbb{Q}_+^* a \cap \mathcal{Z}\}$ has closure $\mathcal{Z}_S$, so $C_0(\mathcal{Z} \setminus \mathcal{Z}_S)$ is the kernel of the representation π_a considered in [**9**, p. 440], and it follows from [**11**, Lemma 4.2] that J_S is the kernel of the representation $\pi_a \times V$ described in [**9**, p. 440]. We can now deduce that $S \mapsto J_S$, as S runs through the proper subsets of $\mathcal{P}$, is the parametrization of $(\operatorname{Prim} C_\mathbb{Q}) \setminus \widehat{\mathbb{Q}_+^*}$ given in [**9**, Theorem 2.8].

THEOREM 4.1. *Suppose that S is a proper subset of $\mathcal{P}$.*

(a) *If $\mathcal{P} \setminus S$ is infinite, then $J_S = \bigcap_{p \notin S} J_{S \cup \{p\}}$.*

(b) *If $0 < |\mathcal{P} \setminus S| < \infty$, then*

$$\Big(\bigcap_{p \notin S} J_{S\cup\{p\}}\Big) \Big/ J_S \cong (C(\mathcal{U}(\mathbb{Z}_{\mathcal{P}\setminus S})) \rtimes_\sigma \mathbb{Z}^S) \otimes \mathcal{K}(l^2(\mathbb{N}^{\mathcal{P}\setminus S})).$$

(c) *$\mathcal{C}_\mathbb{Q}/J_\mathcal{P}$ is isomorphic to $C^*(\mathbb{Q}_+^*) = C(\widehat{\mathbb{Q}_+^*})$.*

Moreover, $C(\mathcal{U}(\mathbb{Z}_{\mathcal{P}\setminus S})) \rtimes_\sigma \mathbb{Z}^S$ is a simple AT-algebra with real rank zero and a unique tracial state.

It follows from [**9**, §2] that every basic open neighbourhood of J_S has the form

$$U_G = \{J_T \mid T \subset \mathcal{P}, T \cap G = \emptyset\}$$

for some finite subset G of $\mathcal{P} \setminus S$. When $\mathcal{P} \setminus S$ is infinite, there are always lots of $J_{S\cup\{p\}}$ in U_G, and thus $J_S \in \overline{\{J_{S\cup\{p\}} \mid p \notin S\}}$; this states precisely that $\bigcap_{p\notin S} J_{S\cup\{p\}} \subset J_S$. The other inclusion is trivial and (a) follows. Part (c) is true because $\mathcal{Z}_\mathcal{P} = \{0\}$. To prove (b) we just need to replace F by $\mathcal{P}$ and $\mathbb{Z}_F$ by $\mathcal{Z}$ in the proof of Lemma 2.6.

It remains to prove the statements about $C(\mathcal{U}(\mathbb{Z}_{\mathcal{P}\setminus S})) \rtimes_\sigma \mathbb{Z}^S$. The Chinese Remainder Theorem implies that $\mathbb{Z}$ is dense in $\mathbb{Z}_{\mathcal{P}\setminus S}$ and, hence, $\mathbb{Z}^S = \mathbb{Z} \cap \mathcal{U}(\mathbb{Z}_{\mathcal{P}\setminus S})$ is dense in $\mathcal{U}(\mathbb{Z}_{\mathcal{P}\setminus S})$. Thus, $\mathbb{Z}^S$ acts minimally and freely on $\mathcal{U}(\mathbb{Z}_{\mathcal{P}\setminus S})$ and $C(\mathcal{U}(\mathbb{Z}_{\mathcal{P}\setminus S})) \rtimes_\sigma \mathbb{Z}^S$ is simple. However, since $|S| = \infty$, we cannot apply the results of [**19**] to conclude that $C(\mathcal{U}(\mathbb{Z}_{\mathcal{P}\setminus S})) \rtimes_\sigma \mathbb{Z}^S$ has real rank zero and stable rank one, as we did in §2 for $C(H) \rtimes_\sigma \mathbb{Z}^S$. Instead we aim to use Theorems 1 and 2 of [**1**], and to do this we need to show that $C(\mathcal{U}(\mathbb{Z}_{\mathcal{P}\setminus S})) \rtimes_\sigma \mathbb{Z}^S$ is an AH-algebra with the extra property of slow dimension growth.

Since $\mathcal{P} \setminus S$ is finite, we have as in (2.7) that $\mathcal{U}(\mathbb{Z}_{\mathcal{P}\setminus S})$ is the inverse limit of the finite groups $\mathcal{U}\big(\mathbb{Z}/\big(\prod_{p\in\mathcal{P}\setminus S} p^{l_p}\big)\mathbb{Z}\big)$ over $l = (l_p) \in \mathbb{N}^{\mathcal{P}\setminus S}$. Hence, $\mathcal{U}(\mathbb{Z}_{\mathcal{P}\setminus S}) = \varprojlim F_n$, where

$$F_n := \mathcal{U}\Big(\mathbb{Z}\Big/\Big(\prod_{p\in\mathcal{P}\setminus S} p^n\Big)\mathbb{Z}\Big). \tag{4.1}$$

We denote the canonical surjection of $\mathcal{U}(\mathbb{Z}_{\mathcal{P}\setminus S})$ onto F_n by π_n. The analogue of Lemma 2.7 for F_n and the canonical action of $\mathbb{Z}^S$ by multiplication on F_n implies that $C(\mathcal{U}(\mathbb{Z}_{\mathcal{P}\setminus S})) \rtimes_\sigma \mathbb{Z}^S$ is the closed union of C^*-subalgebras isomorphic to $C(F_n) \rtimes \mathbb{Z}^S$.

Towards applying Proposition 2.8, we note that the infinite direct sum $\mathbb{Z}^S$ is the union of the subgroups $\mathbb{Z}^E$ associated to finite subsets E of S. Thus, by using an argument similar to

https://doi.org/10.1017/S0143385705000143 Published online by Cambridge University Press

that in Lemma 2.7 we have that $C(F_n) \rtimes \mathbb{Z}^S$ is the closed union of subalgebras isomorphic to $C(F_n) \rtimes \mathbb{Z}^E$. However, by choosing a particular sequence E_n of finite subsets of S, we can show that $C(\mathcal{U}(\mathbb{Z}_{\mathcal{P}\setminus S})) \rtimes_\sigma \mathbb{Z}^S$ has slow dimension growth.

Indeed, since $\mathbb{Z}/\left(\prod_p p^n\right)\mathbb{Z} \cong \prod_p \mathbb{Z}/p^n\mathbb{Z}$, we have $F_n \cong \prod_{p\in\mathcal{P}\setminus S} \mathcal{U}(\mathbb{Z}/p^n\mathbb{Z})$. Thus, F_n is a product of at most $|\mathcal{P}\setminus S|+1$ cyclic groups (the $+1$ allows for the possibility that $2 \in \mathcal{P}\setminus S$) and, hence, has a generating set $\{x_{n,i}\}$ with at most $|\mathcal{P}\setminus S|+1$ elements. By Dirichlet's Theorem, there are primes $q_{n,i}$ such that

$$q_{n,i} \equiv x_{n,i} \quad \left(\mathrm{mod} \prod_{p\in\mathcal{P}\setminus S} p^n\right),$$

and each $q_{n,i}$ belongs to S because it is a unit modulo $\prod_{p\in\mathcal{P}\setminus S} p^n$. Now let $E_n' := \{q_{n,i}\}$, list the primes in S as $\{r_n \mid n \in \mathbb{N}\}$ and take

$$E_n := \left(\bigcup_{m\le n} E_m'\right) \cup \{r_1, \ldots, r_n\}.$$

We then have $\pi_n(\mathbb{Z}^{E_n}) = F_n$, $E_m \subset E_n$ for $m \le n$, and $\bigcup E_n = S$.

We have now realized $C(\mathcal{U}(\mathbb{Z}_{\mathcal{P}\setminus S})) \rtimes_\sigma \mathbb{Z}^S$ as the closure of an increasing union $\bigcup_{n\in\mathbb{N}} B_n$ in which B_n is isomorphic to the crossed product $C(F_n) \rtimes \mathbb{Z}^{E_n}$ by a transitive action of $\mathbb{Z}^{E_n}$. By an argument identical to that at the end of §2 we conclude that $C(\mathcal{U}(\mathbb{Z}_{\mathcal{P}\setminus S})) \rtimes \mathbb{Z}^S$ has a unique tracial state.

We prove next that $C(\mathcal{U}(\mathbb{Z}_{\mathcal{P}\setminus S})) \rtimes \mathbb{Z}^S$ is an AH-algebra with real rank zero. Proposition 2.8 implies that $B_n \cong C(F_n) \rtimes \mathbb{Z}^{E_n} \cong C(\mathbb{T}^{|E_n|}, M_{|F_n|}(\mathbb{C}))$. However,

$$\frac{|E_n|}{|F_n|} \le \frac{n(|\mathcal{P}\setminus S|+2)}{\prod_p (p-1)p^{n-1}} \to 0 \quad \text{as } n \to \infty,$$

and, thus, the sequence $B_n \cong C(F_n) \rtimes \mathbb{Z}^{E_n}$ of subalgebras of $C(\mathcal{U}(\mathbb{Z}_{\mathcal{P}\setminus S})) \rtimes \mathbb{Z}^S$ has slow dimension growth. It now follows from [**1**, Theorem 1] that $C(\mathcal{U}(\mathbb{Z}_{\mathcal{P}\setminus S})) \rtimes \mathbb{Z}^S$ has topological stable rank one. Since the projections in $C(\mathcal{U}(\mathbb{Z}_{\mathcal{P}\setminus S})) \rtimes \mathbb{Z}^S$ trivially separate the unique tracial state, [**1**, Theorem 2] implies that $C(\mathcal{U}(\mathbb{Z}_{\mathcal{P}\setminus S})) \rtimes \mathbb{Z}^S$ has real rank zero. The K-groups of $C(\mathcal{U}(\mathbb{Z}_{\mathcal{P}\setminus S})) \rtimes \mathbb{Z}^S$ are inductive limits of torsion-free groups and, hence, are themselves torsion-free, so it follows as in §2 that $C(\mathcal{U}(\mathbb{Z}_{\mathcal{P}\setminus S})) \rtimes \mathbb{Z}^S$ is an AT-algebra.

This completes the proof of Theorem 4.1.

A. *Appendix. The orders of a prime in groups of units*

For p prime and $m \in \mathbb{N}$ such that $(m, p) = 1$, we denote by $o_{p^l}(m)$ the order of m in $\mathcal{U}(\mathbb{Z}/p^l\mathbb{Z})$. It was shown in [**11**, Theorem 3.1] that if p is odd, there is a positive integer $L_p(m)$ such that

$$o_{p^l}(m) = \begin{cases} o_p(m) & \text{if } 1 \le l \le L_p(m) \\ p^{l-L_p(m)} o_p(m) & \text{if } l > L_p(m); \end{cases} \tag{A.1}$$

the proof uses that the groups $\mathcal{U}(\mathbb{Z}/p^l\mathbb{Z})$ are cyclic. We now show how to modify the arguments of [**11**, §3] to obtain an analogue of (A.1) for $p = 2$, in which case $\mathcal{U}(\mathbb{Z}/2^l\mathbb{Z})$ are no longer cyclic.

https://doi.org/10.1017/S0143385705000143 Published online by Cambridge University Press

PROPOSITION A.1. *If m is an odd integer and $m \equiv 1 \pmod 4$, then there exists a positive integer $K = L_2(m)$ such that*

$$o_{2^l}(m) = \begin{cases} 1 & \text{if } 1 \le l \le K \\ 2^{l-K} & \text{if } l > K; \end{cases} \tag{A.2}$$

if $m \equiv 3 \pmod 4$, then there exists a positive integer $L = L_2(m)$ such that

$$o_{2^l}(m) = \begin{cases} 1 & \text{if } l = 1 \\ 2 & \text{if } 1 < l \le L \\ 2^{l-(L-1)} & \text{if } l > L. \end{cases} \tag{A.3}$$

To prove Proposition A.1 we use general properties of cyclic groups as in [**11**, §3]. We begin with a lemma.

LEMMA A.2. *Suppose $l \ge 3$. Then the group $\{n \in \mathcal{U}(\mathbb{Z}/2^l\mathbb{Z}) \mid n \equiv 1 \pmod 4\}$ is the cyclic subgroup $\langle 5\rangle_l$ of $\mathcal{U}(\mathbb{Z}/2^l\mathbb{Z})$ generated by* 5.

Proof. Theorem 2′ of [**5**, Ch. 4.1] states that $|\langle 5\rangle_l| = 2^{l-2}$. For $k \ge 0$ we have

$$5^k = (4+1)^k = \sum_{n=0}^{k} \binom{k}{n} 4^n = 4 \sum_{n=1}^{k} \binom{k}{n} 4^{n-1} + 1,$$

so $5^k \equiv 1 \pmod 4$. Hence, if $n \equiv 5^k \pmod{2^l}$ for some $0 \le k < 2^{l-2}$, then $n \equiv 1 \pmod 4$. Since the order of $\{n \in \mathcal{U}(\mathbb{Z}/2^l\mathbb{Z}) \mid n \equiv 1 \pmod 4\}$ is also 2^{l-2}, the result follows. □

COROLLARY A.3. *An element of $\mathcal{U}(\mathbb{Z}/2^l\mathbb{Z})$ is congruent to* 3 (mod 4) *if and only if it is congruent to $-5^k \pmod{2^l}$ for some k satisfying $0 \le k < 2^{l-2}$.*

COROLLARY A.4. *Suppose $m \in \mathbb{Z}$ satisfies $m \equiv 1 \pmod 4$. Then for every $l > 0$ we have*

$$o_{2^l}(m) = \begin{cases} o_{2^{l+1}}(m) & \text{if 2 does not divide } o_{2^{l+1}}(m) \\ o_{2^{l+1}}(m)/2 & \text{if 2 divides } o_{2^{l+1}}(m). \end{cases} \tag{A.4}$$

Proof. Since a number is coprime to 2^l if and only if it is coprime to 2^{l+1}, the reduction map $\pi : \mathcal{U}(\mathbb{Z}/2^{l+1}\mathbb{Z}) \to \mathcal{U}(\mathbb{Z}/2^l\mathbb{Z})$ is a surjective homomorphism. Lemma A.2 implies that $m \equiv 5^r \pmod{2^{l+1}}$, where $r = o_{2^{l+1}}(5)/o_{2^{l+1}}(m) = 2^{l-1}/o_{2^{l+1}}(m)$. Hence, by applying [**11**, Lemma 3.2] to the restriction of π to a homomorphism of $\langle 5\rangle_{l+1}$ onto $\langle 5\rangle_l$, we have

$$\begin{aligned} o_{2^l}(m) &= o(\pi(5^r)) \\ &= \begin{cases} 2^{l-1}/(2^{l-1}/o_{2^{l+1}}(m), 2^{l-1}) & \text{if } 2^{l-1} \text{ divides } 2^{l-1}/o_{2^{l+1}}(m) \\ 2^{l-1}/(2(2^{l-1}/o_{2^{l+1}}(m), 2^{l-1})) & \text{if } 2^{l-1} \text{ does not divide } 2^{l-1}/o_{2^{l+1}}(m), \end{cases} \end{aligned}$$

which simplifies to (A.4). □

https://doi.org/10.1017/S0143385705000143 Published online by Cambridge University Press

Proof of Proposition A.1. Suppose first that $m \equiv 1 \pmod 4$. For fixed N, there exists an $l \in \mathbb{N}$ satisfying $m^N < 2^l$. Then $o_{2^l}(m) > N$ and, hence, the sequence $\{o_{2^l}(m) \mid l \in \mathbb{N}\}$ must be unbounded. In particular, $\{o_{2^l}(m)\}$ is not a constant sequence. Let K be the first integer such that $o_{2^K}(m) < o_{2^{K+1}}(m)$. Then $o_{2^l}(m) = o_2(m) = 1$ for $1 \leq l \leq K$ and by Corollary A.4 we have $o_{2^{K+1}}(m) = 2o_2(m) = 2$. Since $o_{2^{K+1}}(m)$ divides $o_{2^l}(m)$ for all $l > K$, it follows that 2 divides $o_{2^l}(m)$ for all $l > K$. We now apply Corollary A.4 $l - K$ times to deduce that $o_{2^l}(m) = 2^{l-K} o_{2^K}(m) = 2^{l-K}$.

Now suppose that $m \equiv 3 \pmod 4$. Certainly $o_2(m) = 1$. For $l > 1$, Corollary A.3 tells us that $m \equiv -5^k$ for some $0 \leq k < 2^{l-2}$. Thus, $m^2 \equiv 5^{2k} \pmod{2^l}$ and, therefore, $m^2 \in \langle 5 \rangle_l$. Let L be the first integer such that $o_{2^L}(m^2) < o_{2^{L+1}}(m^2)$. Applying Corollary A.4 to m^2 and repeating the argument of the preceding paragraph gives (A.3) because $o_{2^l}(m) = 2o_{2^l}(m^2)$. □

We now need to extend these results to cover actions on $\mathcal{U}(\mathbb{Z}_F)$ for an arbitrary finite set F of primes. We write $F = \{p_1, \ldots, p_n\}$ and fix a prime q which is not in F. We denote by $o_{(l_1,\ldots,l_n)}(q)$ the order of $(q, \ldots, q)$ in $\prod_{i=1}^n \mathcal{U}(\mathbb{Z}/{p_i}^{l_i}\mathbb{Z})$.

PROPOSITION A.5. *There exist positive integers $K_1, \ldots, K_n$ and d such that*

$$o_{(K_1+l_1,\ldots,K_n+l_n)}(q) = d\,{p_1}^{l_1} \ldots {p_n}^{l_n} \tag{A.5}$$

for every $(l_1, \ldots, l_n) \in \mathbb{N}^F$.

Proof. Suppose first that $p_1, \ldots, p_n$ are distinct odd primes, and let $L_{p_i}(q)$ be as in (A.1). Let

$$z_i := \max\{z \mid {p_i}^z \text{ divides } o_{p_j}(q) \text{ for some } j \in \{1, \ldots, n\}\},$$

and define $K_i := L_{p_i}(q) + z_i$ and $d := [o_{p_1}(q), \ldots, o_{p_n}(q)]$, where $[r_1, \ldots, r_n]$ is the least common multiple of the integers r_i. In general, if g_i are elements of order r_i in finite groups G_i, then the order of $(g_1, \ldots, g_n)$ in $G_1 \times \cdots \times G_n$ is $[r_1, \ldots, r_n]$. Thus, from the properties of $L_{p_i}(q)$ we obtain

$$\begin{aligned} o_{(K_1+l_1,\ldots,K_n+l_n)}(q) &= [p_1^{(K_1+l_1)-L_{p_1}(q)} o_{p_1}(q), \ldots, p_n^{(K_n+l_n)-L_{p_n}(q)} o_{p_n}(q)] \\ &= [p_1^{z_1+l_1} o_{p_1}(q), \ldots, p_n^{z_n+l_n} o_{p_n}(q)] \\ &= {p_1}^{l_1} \ldots {p_n}^{l_n} [o_{p_1}(q), \ldots, o_{p_n}(q)], \end{aligned} \tag{A.6}$$

which is (A.5).

Now suppose that $2 \in F$, say $p_1 = 2$. If $q \equiv 1 \pmod 4$, we let

$$z_i := \max\{z \mid {p_i}^z \text{ divides } o_{p_j}(q) \text{ for some } j \in \{2, \ldots, n\}\},$$

and define $K_1 := L_2(q) + z_1$, $K_i := L_{p_i}(q) + z_i$ for $i > 1$ and $d := [o_{p_2}(q), \ldots, o_{p_n}(q)]$. Reasoning as in (A.6) gives (A.5).

If $q \equiv 3 \pmod 4$, we let

$$\begin{aligned} z_1 &= \max(1, \max\{z \mid 2^z \text{ divides } o_{p_j}(q) \text{ for some } j \in \{2, \ldots, n\}\}), \\ z_i &= \max\{z \mid {p_i}^z \text{ divides } o_{p_j}(q) \text{ for some } j \in \{2, \ldots, n\}\} \end{aligned}$$

for $i > 1$, and define $K_1 := L_2(q) + z_1 - 1$, $K_i := L_{p_i}(q) + z_i$ for $i > 1$ and $d := [2, o_{p_2}(q), \ldots, o_{p_n}(q)]$. Again, reasoning as in (A.6) gives (A.5). □

https://doi.org/10.1017/S0143385705000143 Published online by Cambridge University Press

COROLLARY A.6. *The closure H of $q^{\mathbb{Z}}$ in $\mathcal{U}(\mathbb{Z}_F)$ is a subgroup of finite index*

$$|\mathcal{U}(\mathbb{Z}_F) : H| = \left(\prod_{i=1}^{n}(p_i - 1)p_i^{K_i - 1}\right)\Big/ d.$$

Proof. Apply Proposition A.5 to $\mathbf{l} = (l, l, \ldots, l)$ to see that $|\pi_l(H)| = d\big(\prod_{i=1}^{n} p_i^l\big)$ for large l and the result follows from [**11**, Lemma 3.7]. □

Acknowledgements. This research was supported by the Carlsberg Foundation, the Natural Sciences and Engineering Research Council of Canada and the Australian Research Council. We thank Chris Phillips for pointing out that the AH-algebras appearing as simple summands in our structure theorems are in fact AT-algebras.

REFERENCES

[1] B. Blackadar, M. Dadarlat and M. Rørdam. The real rank of inductive limit C^*-algebras. *Math. Scand.* **69** (1991), 211–216.

[2] J.-B. Bost and A. Connes. Hecke algebras, Type III factors and phase transitions with spontaneous symmetry breaking in number theory. *Selecta Math. (N. S.)* **1** (1995), 411–457.

[3] K. R. Davidson. *C^*-Algebras by Example (Fields Inst. Monogr., 6)*. American Mathematical Society, Providence, RI, 1996.

[4] G. A. Elliott. On the classification of C^*-algebras of real rank zero. *J. Reine Angew. Math.* **443** (1993), 179–219.

[5] K. Ireland and M. Rosen. *A Classical Introduction to Modern Number Theory*, 2nd edn *(Graduate Texts in Mathematics, 84)*. Springer, Berlin, 1993.

[6] M. Laca. Semigroups of *-endomorphisms, Dirichlet series, and phase transitions in number theory. *J. Funct. Anal.* **152** (1998), 330–378.

[7] M. Laca. From endomorphisms to automorphisms and back: dilations and full corners. *J. London Math. Soc.* **61** (2000), 893–904.

[8] M. Laca and I. Raeburn. A semigroup crossed product arising in number theory. *J. London Math. Soc.* **59** (1999), 330–344.

[9] M. Laca and I. Raeburn. The ideal structure of the Hecke C^*-algebra of Bost and Connes. *Math. Ann.* **318** (2000), 433–451.

[10] N. S. Larsen. Nonunital semigroup crossed products. *Math. Proc. Royal Irish Acad.* **100A** (2000), 205–218.

[11] N. S. Larsen, I. F. Putnam and I. Raeburn. The two-prime analogue of the Hecke C^*-algebra of Bost and Connes. *Indiana Univ. Math. J.* **51** (2002), 171–186.

[12] N. S. Larsen and I. Raeburn. Faithful representations of crossed products by actions of $\mathbb{N}^k$. *Math. Scand.* **89** (2001), 283–296.

[13] H. Lin. Tracially AF C^*-algebras. *Trans. Amer. Math. Soc.* **353** (2001), 693–722.

[14] H. Lin. Classification of simple C^*-algebras of tracial topological rank zero. *Duke Math. J.* **125** (2004), 91–119.

[15] M. Marcus and H. Minc. *A Survey of Matrix Theory and Matrix Inequalities*. Allyn and Bacon, Boston, MA, 1964.

[16] S. Neshveyev. Ergodicity of the action of the positive rationals on the group of finite adeles and the Bost–Connes phase transition theorem. *Proc. Amer. Math. Soc.* **130** (2002), 2999–3003.

[17] D. Olesen and G. K. Pedersen. Partially inner C^*-dynamical systems. *J. Funct. Anal.* **66** (1986), 262–281.

[18] J. A. Packer and I. Raeburn. Twisted crossed products of C^*-algebras. *Math. Proc. Camb. Phil. Soc.* **160** (1989), 293–311.

[19] N. C. Phillips. Crossed products of the Cantor set by free minimal actions of $\mathbb{Z}^d$. *Comm. Math. Phys.* **256** (2005), 1–42.

[20] N. C. Phillips. Crossed products by finite cyclic group actions with the tracial Rokhlin property. *Preprint*, 2003, math.OA/0306410.

https://doi.org/10.1017/S0143385705000143 Published online by Cambridge University Press

[21] M. Pimsner and D. Voiculescu. Exact sequences for K-groups and Ext-groups of certain crossed product C^*-algebras. *J. Operator Th.* **4** (1980), 93–118.

[22] I. Raeburn. On crossed products by coactions and their representation theory. *Proc. London Math. Soc.* **64** (1992), 625–652.

[23] M. Rørdam. *Classification of Nuclear Simple C^*-algebras (Encyclopaedia Math. Sci.,126)*. Springer, Berlin, 2002, pp. 1–145.

https://doi.org/10.1017/S0143385705000143 Published online by Cambridge University Press