Econometric Theory, **31**, 2015, 294–336.
doi:10.1017/S0266466614000310

SPECIFICATION TESTS FOR LATTICE PROCESSES

JAVIER HIDALGO AND MYUNG HWAN SEO
London School of Economics and Seoul National University

We consider an omnibus test for the correct specification of the dynamics of a sequence $\{x(t)\}_{t\in\mathbb{Z}^d}$ in a lattice. As it happens with causal models and $d = 1$, its asymptotic distribution is not pivotal and depends on the estimator of the unknown parameters of the model under the null hypothesis. One first main goal of the paper is to provide a transformation to obtain an asymptotic distribution that is free of nuisance parameters. Secondly, we propose a bootstrap analog of the transformation and show its validity. Thirdly, we discuss the results when $\{x(t)\}_{t\in\mathbb{Z}^d}$ are the errors of a parametric regression model. As a by product, we also discuss the asymptotic normality of the least squares estimator of the parameters of the regression model under very mild conditions. Finally, we present a small Monte Carlo experiment to shed some light on the finite sample behavior of our test.

1. INTRODUCTION

Random models for space or spatio-temporal data play an important role in many disciplines, and in recent years it has become increasingly important in economics, see e.g., Baltagi et al.'s (2007) special volume on the topic or Cressie (1993). Applications cover various areas like environmental, urban, agricultural economics, as well as economic geography among others. In many circumstances data are actually collected in a regular lattice usually as a consequence of planned experiments or because they are collected based on a systematic sampling scheme. Earlier examples are the celebrated paper by Mercer and Hall (1911) on wheat crop yield data or Batchelor and Reed (1924), which were employed as examples and analyzed in the pioneering paper by Whittle (1954). Other examples are given in Cressie and Huang (1999), see also Fernandez-Casal et al. (2003), or within a view toward environmental economics by Mitchell et al. (2005), who employed a regression model of the type in (1.1) to analyze the effect of CO_2 on crops. The latter might also be of relevance in development economics. See also Genton and Koul (2008) on yield of barley in UK and how the models can be useful when there is evidence that there is spatial movement such as pollutants due to winds or ocean currents. When the spatial dimension is one, we obtain the so-called noncausal models or in Whittle's terminology

We would like to thank the Co-Editor and two referees for helpful comments. The first author gratefully acknowledges the research support by a Catedra of Excellence by the Bank of Santander. Address correspondence to Myung Hwan Seo, London School of Economics, Houghton St., London WC2A 2AE, UK; e-mail: m.seo@lse.ac.uk.; or to: Seoul National University, Kwan-Ak Gu, Seoul, Korea.

 © Cambridge University Press 2014

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

linear trascent models. These models can be regarded as forward looking and have gained some consideration recently in economics, see for instance, Davis et al. (2001) or Lanne and Saikkonen (2011, 2013). In general, we can think of lattice or random field models as very useful and practical models to enable us to capture the spatial or the spatio-temporal dependence, see Cressie (1993), Jenish and Prucha (2009), or the neighborhood structure of the data as in Zhu, Huang, and Reyes (2010) among many others.

In particular, given a spatial process $\{x(t)\}_{t\in\mathbb{Z}^d}$, $d \geq 1$, it is agreed that one of the main purposes is to obtain a correct description of its covariogram $\{\gamma(s)\}_{s\in\mathbb{Z}^d}$, defined as $\gamma(s) = \mathrm{Cov}(x(t), x(t+s))$. The importance of the covariogram relies on the fact that it plays a key role to obtain good and accurate predictions and/or interpolations. For instance, see Cressie (1993) in relation to the Gaussian Markov random fields or conditional autoregressions in a lattice. In regression models it enables either correct inferences on the parameters of the model or efficient estimation. When $X = \{x(t)\}_{t=1}^n$ are the errors in a regression model

$$y(t) = \beta_0' z(t) + x(t), \qquad t = 1, \ldots, n, \tag{1.1}$$

where $Z = \{z(t)\}_{t=1}^n$ is a q-dimensional set of fixed regressors, we have that the asymptotic covariance of the least squares estimator of β_0 depends on $\{\gamma(s)\}_{s\in\mathbb{Z}^d}$. In addition, the predictor of say $y(t^*)$ becomes in this case

$$E\left(y\left(t^*\right) \mid \{y(t)\}_{t=1}^n\right) = \beta_0' z\left(t^*\right) + E\left(x\left(t^*\right) \mid X\right),$$

so that an accurate specification of $\gamma(s)$ is the key to obtain a good predictor of $y(t^*)$ or we are just interested to examine the effect of some variables on $y(t)$, see Mitchell et al. (2005) who study the effect of CO_2 on rice crops in Japan.

More specifically, we are interested to check whether the covariogram $\{\gamma(s)\}_{s\in\mathbb{Z}^d}$ follows a particular parametric family, that is $\{\gamma(s)\}_{s\in\mathbb{Z}^d} = \{\gamma(s;\vartheta)\}_{s\in\mathbb{Z}^d}$, where $\vartheta = \left(\theta', \sigma_\varepsilon^2\right)'$ is a $(p+1)$-dimensional vector of unknown parameters. Observing that for any covariance stationary spatial lattice process $\{x(t)\}_{t\in\mathbb{Z}^d}$, the spectral density function $f(\lambda)$ and the covariogram $\{\gamma(s)\}_{s\in\mathbb{Z}^d}$ are related by the expression

$$\gamma(s) = \int_{\Pi^d} e^{-is'\lambda} f(\lambda)\, d\lambda; \quad s \in \mathbb{Z}^d, \tag{1.2}$$

we might have formulated our interest on whether $f(\lambda) = f(\lambda;\theta)$ for all $\lambda \in \Pi^d$. Herewith "$s'\lambda$" means the inner product of two d-dimensional vectors s and λ and $\Pi = [-\pi, \pi]$.

Thus, one of the aims of the paper is to describe a T_p-type omnibus test for the composite hypothesis that the covariogram of the sequence $\{x(t)\}_{t\in\mathbb{Z}^d}$ follows a specified parametric model. One difference with previous work when $d = 1$ is that we allow for models which are also forward looking, i.e., noncausal models or linear trascent models. In addition, we examine the behavior of the test when

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

$\{x(t)\}_{t\in\mathbb{Z}^d}$ is not observed but they are the errors of a parametric regression model. As a by-product, we obtain the asymptotic distribution of the least squares estimator of β_0 in (1.1) under mild conditions. In particular, we show the asymptotic normality when the regressors are deterministic, without the need to assume that the process $\{x(t)\}_{t\in\mathbb{Z}^d}$ is strong mixing as it was assumed in Bolthausen (1982) or more recently in Jenish and Prucha (2009), although our conditions are quite similar to those in Robinson and Thawornkaiwong (2012). Instead, we assume that the process $\{x(t)\}_{t\in\mathbb{Z}^d}$ is a *Generalized linear* process in the sense put forward by Hannan (1970, p. 210), see (1.3) below. The basic condition that we need to obtain the asymptotic distribution of the least squares estimator of β_0 in (1.1) is that the jump of the spectral distribution function of $\{z(t)\}_{t\in\mathbb{Z}^d}$ does not coincide with the discontinuity of the spectral density function of $\{x(t)\}_{t\in\mathbb{Z}^d}$, so that we allow for strong dependence, see also Yajima and Matsuda (2011).

All throughout the paper we shall assume that the spatial linear process $\{x(t)\}_{t\in\mathbb{Z}^d}$ has a representation by the multilateral model

$$x(t)-\mu=\sum_{s\in\mathbb{Z}^d}\psi(s)\varepsilon(t-s),\quad \sum_{s\in\mathbb{Z}^d}\psi^2(s)<\infty,\quad \psi(0)=1, \tag{1.3}$$

for some sequence $\{\varepsilon(t)\}_{t\in\mathbb{Z}^d}$ satisfying $\mathbb{E}(\varepsilon(t))=0$ and $\mathbb{E}(\varepsilon(0)\varepsilon(t))=\sigma_\varepsilon^2\mathbb{I}(t=0)$, where $\mathbb{I}(\cdot)$ is the indicator function. Under (1.3), we have that

$$f(\lambda)=\frac{\sigma_\varepsilon^2}{(2\pi)^d}|\Psi(\lambda)|^2,$$

where $\Psi(\lambda)=\sum_{s\in\mathbb{Z}^d}\psi(s)e^{-is'\lambda}$, which summarizes the covariogram structure of $\{x(t)\}_{t\in\mathbb{Z}^d}$ as seen in (1.2).

Denoting $[0,\pi]\times\Pi^{d-1}$ as $\widetilde{\Pi}^d$, that is $\lambda\in\widetilde{\Pi}^d$ if $\lambda[1]\in[0,\pi]$ and $\lambda[\ell]\in\Pi$ for $\ell=2,\ldots,d$, where $a[\ell]$ denotes the ℓ-th coordinate of the vector a that belongs to $\mathbb{Z}^d$ (or Π^d), we can write the null hypothesis as follows:

$$H_0:\forall\lambda\in\widetilde{\Pi}^d \text{ and for some } \theta_0\in\Theta,\ |\Psi(\lambda)|^2=\left|\Psi_{\theta_0}(\lambda)\right|^2, \tag{1.4}$$

where $\Theta\subset\mathbb{R}^p$ is a compact parameter space and $\Psi_\theta(\lambda)=\sum_{s\in\mathbb{Z}^d}\psi(s;\theta)e^{-is'\lambda}$. The alternative hypothesis is the negation of H_0.

A particular parameterization of (1.3) is the $ARMA(-k_1,k_2;-\imath_1,\imath_2)$ field model, see Whittle (1954), defined as

$$\sum_{s=-k_1}^{k_2}\alpha(s)(x(t-s)-\mu)=\sum_{s=-\imath_1}^{\imath_2}\beta(s)\varepsilon(t-s)\quad \alpha(0)=\beta(0)=1,$$

whose spectral density function is given by

$$f(\lambda)=\frac{\sigma_\varepsilon^2}{(2\pi)^d}\frac{\left|\sum_{s=-\imath_1}^{\imath_2}\beta(s)e^{is'\lambda}\right|^2}{\left|\sum_{s=-k_1}^{k_2}\alpha(s)e^{is'\lambda}\right|^2}.$$

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

Notice that the latter model is causal if $\imath_1 = k_1 = 0$. It is worth mentioning that Whittle (1954) showed that any given stationary multilateral scheme on a plane lattice has a unilateral autoregression with the same spectral scheme, although not necessarily of finite order as is the case when $d = 1$.

Another parametric model of interest is the extension to the lattice of the classical Bloomfield (1973)'s exponential model. In fact, it was introduced by Whittle (1954, Sect. 6) beforehand and it was also named as the *Cepstrum* model by Solo (1986). These models can be characterized as having a spectral density function defined as

$$f_\vartheta(\lambda) = \sigma_\varepsilon^2 \exp\left\{-\sum_{s \prec 0} a(s;\theta)\cos\left(s'\lambda\right)\right\}, \tag{1.5}$$

where "$\prec$" denotes the lexicographical (dictionary) ordering which is defined as

$$s \prec k \Leftrightarrow (\exists \imath > 0)(\forall i < \imath)(s[i] = k[i] \vee s[\imath] < k[\imath]),$$

that is, if one of the terms $s[\imath] < k[\imath]$ and all the previous ones are equal. Observe that if we allowed s in (1.5) to belong to $\mathbb{Z}^d$, the model would not be identified as $\cos\left(s'\lambda\right) = \cos\left(-s'\lambda\right)$ for all $\lambda \in \Pi^d$ and s. Solo (1986) notes that if $0 < f_\vartheta(\lambda) < M$ the representation of the spectral density in (1.5) exists.

Due to the complicated notation in this paper, we have decided to gather it at this stage for convenience. The numbers $0, 1$ and π can be either scalars or vectors (of dimension d), which should be clear from the context, whereas $\mathring{\pi} = (0, \pi, \ldots, \pi)'$ and e_ℓ denotes a vector in $\mathbb{Z}^d$ whose ℓ-th element is one and the other elements are zero. For two vectors a and b, $a \vee b$ and $a \wedge b$ represent the maximum and minimum of the two, respectively, based on the lexicographical ordering, while $a \geq (\leq) b$ means that $a[\ell] \geq (\leq) b[\ell]$ for all $\ell = 1, \ldots, d$. For $\tilde{n} = [n/2]$, denote

$$\Pi_n^d = \left\{\lambda_k[\ell] = \frac{\pi k[\ell]}{\tilde{n}[\ell]}, \quad k[\ell] = 0, \pm 1, \ldots, \pm\tilde{n}[\ell], \quad \ell = 1, \ldots, d\right\},$$

where λ_k stands for the Fourier frequencies. Similar to $\widetilde{\Pi}^d$, we define $\widetilde{\Pi}_n^d = \left\{\lambda_k \in \Pi_n^d : \lambda_k[1] > 0\right\}$. We use three different summation operators when they are taken over the Fourier frequencies, namely

$$\sum_{\lambda_s} = \sum_{\lambda_s \in \widetilde{\Pi}_n^d}, \quad \sum_{\lambda_s}^{\lambda} = \sum_{\substack{\lambda_s \in \widetilde{\Pi}_n^d \\ \lambda_s \leq \lambda}}, \quad \text{and} \quad \sum_{\lambda_s \prec \lambda} = \sum_{\substack{\lambda_s \in \Pi_n^d; \\ 0 \prec \lambda_s \prec \lambda}}.$$

Note that λ_s's in the last summation are taken from Π_n^d.

The remainder of the paper is organized as follows. In the next section, we present the test and examine its asymptotic properties, showing that it is not pivotal as its asymptotic distribution depends on H_0 and on the estimator employed

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

of the unknown parameters ϑ_0. Because the asymptotic critical values are difficult to obtain, Section 3 describes a transformation in the spirit of Brown, Durbin, and Evans (1975) such that it converges to a functional of the "standard" Brownian sheet in $[0,1]^d$. The transformation mirrors that of Delgado, Hidalgo, and Velasco (2005) to the case when $d > 1$, or when $d = 1$ and the model is not causal. We also describe a bootstrap algorithm, showing its validity, to compute the critical values of the transformation. Section 4 describes the local alternatives and it also examines the consequences, if any, when $\{x(t)\}_{t\in\mathbb{Z}^d}$ are the errors of a parametric regression model. In addition, we show the asymptotic normality of the least squares estimator of the parameters under mild conditions. Section 5 presents the results of a Monte Carlo study to shed some light on the finite sample performance of our test and its bootstrap analog. Section 6 concludes. The proofs of our main results are given in Appendix B, which uses a series of lemmas given in Appendix A.

2. THE TEST AND ITS PROPERTIES

Before we introduce and describe the test, we first observe that we can state the null hypothesis (1.4) as

$$H_0 : \forall \lambda \in \widetilde{\Pi}^d \text{ and some } \theta_0 \in \Theta,\ \frac{G_{\theta_0}(\lambda)}{G_{\theta_0}(\pi)} = \frac{\lambda[1]}{\pi}\prod_{\ell=2}^{d}\left(1+\frac{\lambda[\ell]}{\pi}\right), \tag{2.1}$$

where

$$G_\theta(\lambda) = 2\int_{-\mathring{\pi}}^{\lambda} |\Psi_\theta(\omega)|^{-2} f(\omega)\, d\omega.$$

Under H_0, $G_{\theta_0}(\lambda)/(2\pi)^d$ is the spectral distribution function of the lattice process $\{\varepsilon(t)\}_{t\in\mathbb{Z}^d}$ and $G_{\theta_0}(\pi) = \sigma_\varepsilon^2$. Notice that by symmetry of $f(\lambda)$, it does not matter which coordinate we choose in the interval $[0,\pi]$, as it will not affect the value of $G_\theta(\lambda)$ and hence the test given below. We shall indicate though that for simplicity of arguments, we focus in the case when $d \le 3$. Extensions to $d > 3$ can be adapted easily under suitable modifications.

Let $h_n(t) = 2^{-d}\prod_{\ell=1}^{d} h(t[\ell]/n[\ell])$, where $h(\cdot)$ is a function in $[0,1]$, and define the taper *periodogram* of a generic sequence $\{v(t)\}_{t=1}^n$ by

$$I_v^T(\lambda) = \frac{1}{\sum_{t=1}^n h_n^2(t)}\left|\sum_{t=1}^{n} h_n(t)\, v(t)\, e^{-it'\lambda}\right|^2.$$

The motivation to employ the taper periodogram instead of the standard periodogram, i.e., when $h(\cdot) = 1$, is due to the adverse properties that $\widehat{\vartheta}$ in (2.3) would have with $h(t) = 1$ as Guyon (1982) observed. Recall that tapering is primarily a technique employed to reduce the bias of the "standard" periodogram,

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

although it increases the variance by a factor $P_4^2 = \left(\int_0^1 h^4\right)\left(\int_0^1 h^2\right)^{-2}$. Another possibility is the one described by Robinson and Vidal-Sanz (2006), which would be helpful when $d \geq 4$. However as we only consider explicitly the most common scenario when $d \leq 3$, it then suffices to employ $I_v^T(\lambda_s)$.

Given a record $\{x(t)\}_{t=1}^n$, and denoting henceforth $N = \Pi_{\ell=1}^d n[\ell]$, a natural estimator of $G_{\theta_0}(\lambda)$ is $G_{\widehat{\theta}N}(\lambda)$ for a given estimate $\widehat{\theta}$, where

$$G_{\theta N}(\lambda) = \frac{(2\pi)^d}{N} \sum_{\lambda_s}^{\lambda} \frac{I_x^T(\lambda_s)}{|\Psi_\theta(\lambda_s)|^2}. \tag{2.2}$$

The summation in (2.2) is taken over $\widetilde{\Pi}_n^d$ instead of the half space $\{\lambda_s \succ 0\}$ to ease notation and exposition.

For $\widehat{\theta}$, we employ the Whittle's (1954) estimator of $\vartheta_0 = \left(\theta_0', \sigma_\varepsilon^2\right)'$ defined as

$$\widehat{\vartheta} = \arg \min_{\vartheta \in \Theta \times \mathbb{R}^+} \mathcal{Q}_N(\vartheta), \tag{2.3}$$

where

$$\mathcal{Q}_N(\vartheta) = \frac{1}{N} \sum_{\lambda_s} \left\{ \log f_\vartheta(\lambda_s) + \frac{I_x^T(\lambda_s)}{(2\pi)^d f_\vartheta(\lambda_s)} \right\}$$

with $f_\vartheta(\lambda_s) = \sigma_\varepsilon^2 |\Psi_\theta(\lambda_s)|^2 / (2\pi)^d$. It is worth pointing out that because our model is multilateral, one consequence is that $\{\varepsilon(t)\}_{t \in \mathbb{Z}^d}$ loses its interpretation as the "prediction" error or as the innovations. The implication of the latter is that the standard least squares estimator of the parameters θ, that is

$$\widehat{\theta}^{LSE} = \arg \min_{\theta \in \Theta} \sum_{\lambda_s} \frac{I_x^T(\lambda_s)}{|\Psi_\theta(\lambda_s)|^2},$$

is an inconsistent estimator of θ_0, see Whittle (1954).

The formulation of H_0 given in (2.1) suggests to use the Bartlett's $T_p - process$ $\alpha_{\widehat{\theta}N}(\lambda)$ as the basis to test H_0, where

$$\alpha_{\theta N}(\lambda) = \frac{1}{2^{d/2} P_4} N^{1/2} \left[\frac{G_{\theta N}(\lambda)}{G_{\theta N}(\pi)} - \frac{\lambda[1]}{\pi} \prod_{\ell=2}^d \left(1 + \frac{\lambda[\ell]}{\pi}\right) \right], \quad \lambda \in \widetilde{\Pi}^d, \tag{2.4}$$

with $G_{\theta N}(\lambda)$ given in (2.2). From here, we can base the test for H_0 using $\eta\left(\alpha_{\widehat{\theta}N}\right)$ for some continuous functional $\eta : D\left(\widetilde{\Pi}^d\right) \to \mathbb{R}^+$, where $D\left(\widetilde{\Pi}^d\right)$ is the space of càdlàg functions in $\widetilde{\Pi}^d$.

Let us introduce the following regularity conditions.

***C1*:** (*a*) $\{\varepsilon(t)\}_{t \in \mathbb{Z}^d}$ in (1.3) is a sequence of zero mean independent identically distributed random variables with $\mathbb{E}\left(\varepsilon^2(t)\right) = \sigma_\varepsilon^2 = 1$ and finite $4th$ moments, denoting its fourth cumulant by κ_ε.

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

(*b*) The multilateral *moving average* representation of $\{x(t)\}_{t\in\mathbb{Z}^d}$ in (1.3) can be written as a multilateral *autoregressive* model

$$\sum_{s\in\mathbb{Z}^d}\xi(s)\,x(t-s)=\varepsilon(t)\quad \xi(0)=1,$$

where $\xi(s)$ is the coefficient of z^s in the Fourier expansion of $\wp^{-1}(z)$, where

$$\wp(z)=\wp(z[1],\ldots,z[d])=\sum_{s\in\mathbb{Z}^d}\psi(s)\,z^s$$

using the notation $z^s=\Pi_{\ell=1}^d z[\ell]^{s[\ell]}$ and the convention $0^0=1$.

***C2*:** $n[\ell]\asymp\vec{n}\nearrow\infty$ for $\ell=1,\ldots,d$, where "$a\asymp b$" means that $C^{-1}\le a/b\le C$ for some finite positive constant C.

***C3*:** $h(\cdot)$ is the *cosine-bell* taper, that is,

$$h(z)=(1-\cos(2\pi z)).$$

***C4*:** θ_0 is an interior point of the compact parameter set $\Theta\subset\mathbb{R}^p$.

***C5*:** $|\Psi_\theta(\lambda)|=\left|\sum_{s\in\mathbb{Z}^d}\psi_\theta(s)\,e^{-is'\lambda}\right|$ is a positive and twice continuously differentiable function in θ on $\widetilde{\Pi}^d$ and continuously differentiable in λ for all $\theta\in\Theta$.

***C6*:** If $\theta_1\neq\theta_2$, then $\Psi_{\theta_1}(\lambda)\neq\Psi_{\theta_2}(\lambda)$ in a set $\Delta\subset\widetilde{\Pi}^d$ with positive Lebesgue measure.

Conditions $C1$–$C6$ are similar to those in Hidalgo (2009) and so his comments apply here. Notice that we write explicitly $\mathbb{E}\left(\varepsilon^2(t)\right)=\sigma_\varepsilon^2$ as it is a parameter in itself, although for notational simplicity we have assumed that its true value is 1, cf. Condition $C1(a)$. Also note that the condition $C1(b)$ allows for forwarding looking noncausal models, which draw some attention in econometrics as in e.g., Lanne and Saikonnen (2011, 2013).

Let

$$\phi_\vartheta(\lambda)=\frac{\partial}{\partial\vartheta}\log f_\vartheta(\lambda)=\left(\varphi_\theta'(\lambda),\sigma_\varepsilon^{-2}\right)',\quad \varphi_\theta(\lambda)=\frac{\partial}{\partial\theta}\log|\Psi_\theta(\lambda)|^2 \tag{2.5}$$

and

$$\Phi_\vartheta(\lambda)=(2\pi)^{-d}\int_{-\grave{\pi}}^{\lambda}\phi_\vartheta(\omega)\,d\omega\ \text{ and }\ \Lambda_\vartheta=(2\pi)^{-d}\int_{-\grave{\pi}}^{\pi}\phi_\vartheta(\omega)\,\phi_\vartheta'(\omega)\,d\omega.$$

***C7*:** Λ_ϑ is a continuous positive definite matrix at $\vartheta=\vartheta_0$.

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

Proceeding as in Hidalgo (2009), we have that the Whittle estimator $\widehat{\vartheta}$ in (2.3) satisfies the asymptotic linearization

$$\widehat{\theta}-\theta_0=-\widetilde{\Lambda}_{\theta_0 N}^{-1}\int_{-\mathring{\pi}}^{\pi}\widetilde{\varphi}_{\theta_0}(\lambda)\,\alpha_{\theta_0 N}(d\lambda)+o_p\left(N^{-1/2}\right),$$

where

$$\widetilde{\varphi}_{\theta}(\lambda)=\varphi_{\theta}(\lambda)-\frac{2}{(2\pi)^d}\int_{-\mathring{\pi}}^{\pi}\varphi_{\theta}(\lambda)\,d\lambda,$$

and defining $\widetilde{\varphi}_{\theta N}(\lambda_s)=\varphi_{\theta}(\lambda_s)-\frac{2}{N}\sum_{\lambda_s}\varphi_{\theta}(\lambda_s)$,

$$\widetilde{\Lambda}_{\theta N}=N^{-1}\sum_{\lambda_k}\widetilde{\varphi}_{\theta N}(\lambda_k)\,\widetilde{\varphi}_{\theta N}'(\lambda_k).$$

Let

$$\dot{\boldsymbol{B}}(\lambda)=\boldsymbol{B}\left(\frac{\lambda}{\pi}\right)-\left\{\frac{\lambda[1]}{\pi 2^{d-1}}\prod_{\ell=2}^{d}\left(1+\frac{\lambda[\ell]}{\pi}\right)\right\}\boldsymbol{B}(1)\quad\lambda\in\widetilde{\Pi}^d, \tag{2.6}$$

where $\left\{\boldsymbol{B}(u):u\in[0,1]\times[-1,1]^{d-1}\right\}$ denotes a zero mean Gaussian process such that

$$\mathrm{Cov}\left(\boldsymbol{B}(u),\boldsymbol{B}(v)\right)=2^{1-d}\left|u[1]\wedge v[1]\right|\prod_{\ell=2}^{d}\left|(u[\ell]\wedge v[\ell])+1\right|,$$

that is, $\boldsymbol{B}$ is a time-changed Brownian sheet. Also let

$$\widetilde{\Lambda}_{\theta}=(2\pi)^{-d}\int_{-\mathring{\pi}}^{\pi}\widetilde{\varphi}_{\theta}\left(\bar{\lambda}\right)\widetilde{\varphi}_{\theta}'\left(\bar{\lambda}\right)d\bar{\lambda}$$

and define

$$\alpha_N^0(\lambda)=\frac{N^{1/2}}{2^{d/2}P_4}\left[\frac{G_N^0(\lambda)}{G_N^0(\pi)}-\frac{\lambda[1]}{\pi}\prod_{\ell=2}^{d}\left(1+\frac{\lambda[\ell]}{\pi}\right)\right],\quad\lambda\in\widetilde{\Pi}^d, \tag{2.7}$$

with $G_N^0(\lambda)=\frac{(2\pi)^d}{N}\sum_{\lambda_s}^{\lambda}I_{\varepsilon}^T(\lambda_s)$. Denoting

$$\alpha_{\infty}(\lambda)=\dot{\boldsymbol{B}}(\lambda)-\left(\frac{1}{(2\pi)^d}\int_{-\mathring{\pi}}^{\lambda}\widetilde{\varphi}_{\theta_0}'\left(\bar{\lambda}\right)d\bar{\lambda}\right)\widetilde{\Lambda}_{\theta_0}^{-1}\int_{-\mathring{\pi}}^{\pi}\widetilde{\varphi}_{\theta_0}\left(\bar{\lambda}\right)\dot{\boldsymbol{B}}\left(d\bar{\lambda}\right),$$

we then have the following result.

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

THEOREM 1. *Under H_0 and assuming C1 − C7, uniformly in $\lambda \in \widetilde{\Pi}^d$, we have that*

$$\alpha_{\widehat{\theta}N}(\lambda) = \alpha_N^0(\lambda) - \left(\frac{1}{N}\sum_{\lambda_s}^{\lambda} \widetilde{\varphi}'_{\theta_0 N}(\lambda_s)\right) \widetilde{\Lambda}_{\theta_0 N}^{-1} \frac{1}{N}\sum_{\lambda_s} \widetilde{\varphi}'_{\theta_0 N}(\lambda_s) I_\varepsilon^T(\lambda_s) + o_p(1)$$
$$\Rightarrow \alpha_\infty(\lambda).$$

Proof. See Hidalgo (2009). ■

The main conclusion from Theorem 1 is that the asymptotic distribution of the $T_p - process$ $\alpha_{\widehat{\theta}N}(\lambda)$ depends on the model under H_0 and also on the estimator of θ_0. So, the asymptotic critical values of $\eta\left(\alpha_{\widehat{\theta}N}\right)$, for any continuous functional $\eta(\cdot)$, cannot be easily tabulated. To circumvent this type of problem, several approaches have been described. A first approach proposes to use bootstrap algorithms. This is the route employed, among others, by Chen and Romano (2000) or Hainz and Dahlhaus (2000) using the $U_p - process$ and by Hidalgo and Kreiss (2006) who employed the $T_p - process$. Of course, all those works were for $d = 1$, whereas Hidalgo (2009) extends the previous work when $d \geq 1$. A second alternative compares the parametric and nonparametric fits of the spectral density function. This route was followed, among others, by Hong (1996) or Paparoditis (2000) for $d = 1$ and Crujeiras et al. (2008) when $d > 1$. However, the implementation of the test depends on a bandwidth parameter and they are inefficient compared to tests based on $\eta\left(\alpha_{\widehat{\theta}N}\right)$ due to the loss of local power proportional to the bandwidth parameter. One additional disadvantage is that there is not a clear procedure as to how to choose the bandwidth parameter that, keeping the correct size of the test, conveys good power properties. A third approach is to employ a transformation of $\alpha_{\widehat{\theta}N}$ that converges in distribution to the "standard" Brownian sheet. This is the route that we follow in the next section.

3. MARTINGALE TRANSFORMATION: ITS BOOTSTRAP ANALOG

In this section we shall present and examine a martingale transformation $\mathcal{L}_{\theta N}(\cdot)$ of $\alpha_{\widehat{\theta}N}(\lambda)$, as well as its bootstrap analog, when $d \geq 1$. The transformation resembles ideas introduced by Brown, Durbin, and Evans (1975) and examined in depth by Khmaladze (1981) and Delgado, Hidalgo, and Velasco (2005) when $d = 1$ and the model is causal. Our aim in this section is thus to extend the latter approach to $d > 1$ and/or noncausal models. The approach parallels the existing similarities between Khmaladze's (1981) transformation and the CUSUM of least squares residuals approach followed in Delgado et al. (2005) in that the latter can be considered a discrete version of the former. In our context, as we will see below, we will mirror the transformation given in McKeague et al.'s (1995). More specifically, our aim shall be to show that the transformation $\mathcal{L}_{\widehat{\theta}N}\left(\alpha_{\widehat{\theta}N}\right)$ converges weakly to the time-changed Brownian sheet $\boldsymbol{B}(\cdot)$ defined in (2.6). In addition, we describe a bootstrap analog of $\mathcal{L}_{\widehat{\theta}N}\left(\alpha_{\widehat{\theta}N}\right)$ showing its validity.

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

For that purpose, it is worth first noticing that Theorem 1 part (a) indicates that $\alpha_{\widehat{\theta}N}$ has the uniform asymptotic expansion

$$\sup_{\lambda\in\widetilde{\Pi}^d}\left|\alpha_{\widehat{\theta}N}(\lambda)-\frac{2\pi}{G_N^0(\pi)}\frac{1}{N^{1/2}}\sum_{\lambda_s}^{\lambda}u_N(\lambda_s)\right|=o_p(1),$$

where

$$u_N(\lambda_s)=\check{I}_\varepsilon^T(\lambda_s)-\widetilde{\varphi}'_{\theta_0N}(\lambda_s)\widetilde{\Lambda}_{\theta_0N}^{-1}N^{-1}\sum_{\lambda_k}\widetilde{\varphi}_{\theta_0N}(\lambda_k)\check{I}_\varepsilon^T(\lambda_k)$$

and $\check{I}_\varepsilon^T(\lambda_s)=I_\varepsilon^T(\lambda_s)-\widehat{\sigma}_\varepsilon^2$. Here $\widehat{\sigma}_\varepsilon^2=G_{\widehat{\theta}N}(\pi)$ which is a $N^{1/2}$-consistent estimator of $\sigma_\varepsilon^2=1$, see Hidalgo (2009). Now observing that we can consider $u_N(s)$ as the least squares residuals in the artificial regression model of $I_\varepsilon^T(\lambda_s)$ on $\left(1,\varphi'_{\theta_0N}(\lambda_s)\right)'$, it suggests employing the *CUSUM* of recursive least squares residuals to construct asymptotically pivotal tests as originally proposed by Brown, Durbin, and Evans (1975). In our case, the recursive estimation is based on the lexicographic ordering in $\widetilde{\Pi}_n^d$, whose minimum value is $(\pi/\tilde{n}[1],-\pi,\ldots,-\pi)'$.

Let $\widetilde{\Lambda}_{\theta N}(\lambda)=N^{-1}\sum_{\lambda_k\preceq\lambda}\widetilde{\varphi}_{\theta N}(\lambda_k)\widetilde{\varphi}'_{\theta N}(\lambda_k)$ and assume the following condition.

***C8*:** $\widetilde{\Lambda}_{\theta_0N}\left((\pi/\tilde{n}[1],-\pi,\ldots,-\pi)'\right)$ is nonsingular a.s. (almost surely) for all n.

Condition $C8$ is very mild and satisfied for all common models used with real data. Recall that p is the dimension of the parameter θ_0. Also, notice that we can directly compute from the model the deterministic matrix $\widetilde{\Lambda}_{\theta_0N}(\cdot)$.

The (scaled) CUSUM of recursive least squares residuals is thus defined as

$$\beta_N^0(\lambda)=\frac{1}{G_N^0(\pi)}\frac{2^{1/2}}{N^{1/2}}\sum_{\lambda_s}^{\lambda}e_N(\lambda_s),\quad\lambda\in\widetilde{\Pi}^d,$$

where

$$e_N(\lambda_s)=\check{I}_\varepsilon^T(\lambda_s)-\widetilde{\varphi}'_{\theta_0N}(\lambda_s)b_N(\lambda_s)$$

are the least squares residuals with

$$b_N(\lambda_s)=\widetilde{\Lambda}_{\theta_0N}^{-1}(\lambda_s)\frac{1}{N}\sum_{\lambda_k\prec\lambda_s}\widetilde{\varphi}_{\theta_0N}(\lambda_k)\check{I}_\varepsilon^T(\lambda_k).$$

Of course, we could have used the forward least squares residuals, i.e.,

$$e_N^f(\lambda_s)=\check{I}_\varepsilon^T(\lambda_s)-\widetilde{\varphi}'_{\theta_0N}(\lambda_s)b_N(\lambda_s),\quad s\prec\mathring{p},$$

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

with $\mathring{p} = (\tilde{n}[1], \tilde{n}[2], \ldots, \tilde{n}[d-1], \tilde{n}[d] - p - 1)$ and

$$b_N^f(\lambda_s) = \widetilde{\Lambda}_{\theta_0 N}^{-1}(\lambda_s) \frac{1}{N} \sum_{\lambda_s \prec \lambda_k} \widetilde{\varphi}_{\theta_0 N}(\lambda_k) \check{I}_\varepsilon^T(\lambda_k)$$

being the conclusions the same as with $e_N(\lambda_s)$.

The empirical process β_N^0 is a linear transformation of α_N^0, i.e.,

$$\beta_N^0(\lambda) = \mathcal{L}_{\theta_0 N}\left(\alpha_N^0(\lambda)\right), \quad \lambda \in \widetilde{\Pi}^d,$$

where, for any function $g \in D\left(\widetilde{\Pi}^d\right)$,

$$\mathcal{L}_{\theta N}(g(\lambda)) = g(\lambda) - \frac{1}{N} \sum_{\lambda_s}^{\lambda} \widetilde{\varphi}'_{\theta N}(\lambda_s) \widetilde{\Lambda}_{\theta N}^{-1}(\lambda_s) \frac{1}{N} \sum_{\lambda_k \prec \lambda_s} \widetilde{\varphi}_{\theta N}(\lambda_k) g(\lambda_k).$$

The transformation $\mathcal{L}_{\theta_0 N}$ has the limiting version $\mathcal{L}^0$, defined as

$$\mathcal{L}^0(g(\lambda)) = g(\lambda) - \frac{1}{(2\pi)^d} \int_{-\mathring{\pi}}^{\lambda} \widetilde{\varphi}_{\theta_0}(\overline{\lambda}) \widetilde{\Lambda}_{\theta_0}^{-1}(\overline{\lambda}) \left(\int_{\widetilde{\lambda} \prec \overline{\lambda}} \widetilde{\varphi}_{\theta_0}(\widetilde{\lambda}) g(d\widetilde{\lambda}) \right) d\overline{\lambda}.$$

Notice that for $d = 1$, $\mathcal{L}^0(\alpha_\infty)$ is the martingale innovation of α_∞, see Khmaladze (1981). On the other hand, in our context, $\mathcal{L}^0(\alpha_\infty)$ becomes the transformation examined by McKeague et al. (1995). That is consider

$$\eta(\lambda) = B(\lambda) - K(\lambda)\xi,$$

where $B(\lambda)$ is a Brownian sheet in $[0,1]^2$ and $K(\lambda) = \int_0^{\lambda[1]} \int_0^{\lambda[2]} k(s,x)\, ds dx$ and letting $d = 2$ for simplicity. Then, they show that

$$W(\lambda) = \eta(\lambda) - \int_0^{\lambda[2]} \left[\int_0^{\lambda[1]} k(s,x) \left\{ \frac{\int_x^1 k(s,u)\, d\eta(s,u)}{\int_x^1 k^2(s,r)\, dr} \right\} ds \right] dx$$

follows a Brownian sheet. In this sense, $\mathcal{L}_{\theta N}(g(\lambda))$ becomes the discrete version of the latter. In our context $k(s,x) = \widetilde{\varphi}_\theta(s,x)$ and ξ is the asymptotic distribution of $N^{1/2}\left(\widehat{\theta} - \theta_0\right)$. Also, it is worth mentioning that the transformation is valid whether any other $N^{1/2}$-consistent estimators of θ_0 were employed.

THEOREM 2. *Under H_0 and assuming $C1 - C8$, $\beta_N^0(\lambda) \Rightarrow \boldsymbol{B}(\lambda/\pi)$; $\lambda \in \widetilde{\Pi}^d$.*

Proof. The proof proceeds, if it is not easier, as that of Theorem 4 part (a) and thus it is omitted. ■

Because β_N^0 cannot be computed in practice, as it depends on θ_0, we employ a finite sample analog. Let

$$\mathring{I}_{x,\theta}^T(\lambda_s) = \frac{I_x^T(\lambda_s)}{|\Psi_\theta(\lambda_s)|^2} - \frac{1}{\boldsymbol{J}} \sum_{\lambda_k \prec \lambda_s} \frac{I_x^T(\lambda_k)}{|\Psi_\theta(\lambda_k)|^2},$$

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

where $\boldsymbol{J} = \#\left\{\lambda_k \in \Pi_n^d : 0 \prec \lambda_k \prec \lambda_s\right\}$, and introduce the recursive residuals in the linear projection of $\left\{\mathring{I}_{x,\theta}^T(\lambda_k)\right\}_{0\prec\lambda_k\prec\lambda_s}$ on $\{1, \varphi_\theta(\lambda_k)\}_{0\prec\lambda_k\prec\lambda_s}$, that is,

$$e_{\theta N}(\lambda_s) = \mathring{I}_{x,\theta}^T(\lambda_s) - \widetilde{\varphi}'_{\theta N}(\lambda_s)\, b_{\theta N}(\lambda_s),$$

where $b_{\theta N}(\lambda_s) = \widetilde{\Lambda}_{\theta N}^{-1}(\lambda_s)\frac{1}{N}\sum_{\lambda_k\prec\lambda_s}\widetilde{\varphi}_{\theta N}(\lambda_k)\,\mathring{I}_{x,\theta}^T(\lambda_k)$. Then, we consider $\beta_{\widehat{\theta}N} = \mathcal{L}_{\widehat{\theta}N}\left(\alpha_{\widehat{\theta}N}(\lambda)\right)$ as the finite sample analog of β_N^0, where

$$\beta_{\theta N}(\lambda) = \frac{1}{G_{\theta N}(\pi)\, P_4}\frac{2^{d/2}}{N^{1/2}}\sum_{\lambda_s}^{\lambda} e_{\theta N}(\lambda_s), \quad \lambda \in \widetilde{\Pi}^d.$$

To establish the asymptotic equivalence between $\beta_N^0(\cdot)$ and $\beta_{\widehat{\theta}N}(\cdot)$, we need an extra smoothness condition on the model under H_0.

***C9*:** For all $\lambda \in \widetilde{\Pi}^d$, $\varphi_\theta(\lambda)$ is twice continuously differentiable in θ.

THEOREM 3. *Assuming $C1 - C9$, under H_0,*

$$\sup_{\lambda\in\widetilde{\Pi}^d}\left|\beta_{\widehat{\theta}N}(\lambda) - \beta_N^0(\lambda)\right| = o_p(1).$$

Proof. The proof proceeds, if it is not easier, as that of Theorem 4 part (b) and thus it is omitted. ■

From a computational point of view, it is worth observing that

$$\widetilde{\Lambda}_{\theta N}^{-1}\left(\lambda_{s+1}\right) = \widetilde{\Lambda}_{\theta N}^{-1}(\lambda_s) - \frac{\widetilde{\Lambda}_{\theta N}^{-1}(\lambda_s)\,\widetilde{\varphi}_{\theta N}\left(\lambda_{s+1}\right)\widetilde{\varphi}'_{\theta N}\left(\lambda_{s+1}\right)\widetilde{\Lambda}_{\theta N}^{-1}(\lambda_s)}{N + \widetilde{\varphi}'_{\theta N}\left(\lambda_{s+1}\right)\widetilde{\Lambda}_{\theta N}^{-1}(\lambda_s)\,\widetilde{\varphi}_{\theta N}\left(\lambda_{s+1}\right)}$$

and, proceeding as in Brown, Durbin, and Evans (1975),

$$\begin{aligned} b_{\theta N}\left(\lambda_{s+1}\right) = {} & b_{\theta N}(\lambda_s) + \widetilde{\Lambda}_{\theta N}^{-1}\left(\lambda_{s+1}\right)\widetilde{\varphi}_{\theta N}\left(\lambda_{s+1}\right) \\ & \times\left[\mathring{I}_{x,\theta}^T\left(\lambda_{s+1}\right) - \widetilde{\varphi}'_{\theta N}\left(\lambda_{s+1}\right) b_{\theta N}(\lambda_s)\right], \end{aligned}$$

where $\lambda_{s+1} = \min\left\{\lambda_k \in \widetilde{\Pi}_n^d : \lambda_k \succ \lambda_s\right\}$.

COROLLARY 1. *Let $\eta : D\left(\widetilde{\Pi}^d\right) \to \mathbb{R}^+$ be a continuous functional. Then, under H_0 and the conditions in Theorem 3, we have that*

$$\eta\left(\beta_{\widehat{\theta}N}\right) \xrightarrow{d} \eta(\boldsymbol{B}).$$

Proof. The proof is an immediate consequence of Theorems 2 and 3 and the continuous mapping theorem, so it is omitted. ■

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

Two standard functionals $\eta(\cdot)$ are the Kolmogorov–Smirnov and the Cramer–von Mises defined as

$$\widehat{K}_N = \sup_{\lambda_s \in \widetilde{\Pi}_n^d} \left|\beta_{\widehat{\theta}N}(\lambda_s)\right| \xrightarrow{d} \sup_{\lambda \in \widetilde{\Pi}^d} \left|\boldsymbol{B}\left(\frac{\lambda}{\pi}\right)\right|,$$

$$\widehat{C}_N = \frac{2}{N}\sum_{\lambda_s}^{\lambda} \beta_{\widehat{\theta}N}(\lambda_s)^2 \xrightarrow{d} \frac{2}{(2\pi)^d}\int_{-\mathring{\pi}}^{\pi} \boldsymbol{B}^2\left(\frac{\lambda}{\pi}\right) d\lambda.$$

Note that the limiting random variables can be represented as the supremum and integral of the d-dimensional standard Brownian sheet by an appropriate change-of-variable.

3.1. Bootstrap Approach

As mentioned at the beginning of Section 3, we shall present and examine the bootstrap analog of $\eta\left(\beta_{\widehat{\theta}N}\right)$. To that end, we define for a generic sequence $\{v(t)\}_{t=1}^n$, the *discrete Fourier transform* as

$$w_v(\lambda) = \frac{1}{N^{1/2}}\sum_{t=1}^{n} v(t)\, e^{-it'\lambda}.$$

The bootstrap analog of $\beta_{\theta N}(\lambda)$ is described in the following *3 STEPS*.

STEP 1: We first obtain the residuals $\{\widehat{\varepsilon}(t)\}_{t=1}^n$ as

$$\widehat{\varepsilon}(t) = \frac{1}{N^{1/2}}\sum_{s=-\tilde{n}}^{\tilde{n}} e^{-it'\lambda_s}\Psi_{\widehat{\theta}}^{-1}(\lambda_s)\, w_x(\lambda_s),$$

and we obtain a random sample of size $\mathring{n} = (2n[1], \ldots, 2n[d])$ with replacement from the empirical distribution function of $\{\widehat{\varepsilon}(t)\}_{t=1}^n$. Denote the sample by $\{\varepsilon^*(t)\}_{t=1}^{\mathring{n}}$ and compute $\{\widetilde{x}^*(t)\}_{t=1}^{\mathring{n}}$ by

$$\widetilde{x}^*(t) = \frac{1}{2^{(d+1)/2}N^{1/2}}\sum_{s=-\tilde{n}}^{\tilde{n}} e^{-it'\widetilde{\lambda}_s}\Psi_{\widehat{\theta}}\left(\widetilde{\lambda}_s\right) w_{\varepsilon^*}\left(\widetilde{\lambda}_s\right), \tag{3.1}$$

where $\widetilde{\lambda}_s$ are

$$\widetilde{\lambda}_s[\ell] = \frac{\pi s[\ell]}{n[\ell]}; \quad s[\ell] = 0, \pm 1, \ldots, \pm n[\ell], \quad \ell = 1, \ldots, d.$$

Finally, our bootstrap sample is $\{x^*(t)\}_{t=1}^n = \{\widetilde{x}^*(t+n)\}_{t=n+1}^{\mathring{n}}$.

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

Remark 1. (**a**) Notice that because $\beta_{\widehat{\theta}N} = \mathcal{L}_{\widehat{\theta}N}\left(\alpha_{\widehat{\theta}N}\right)$ is independent of the first two moments of $\{\varepsilon(t)\}_{t\in\mathbb{Z}^d}$, we do not need to standardize $\widehat{\varepsilon}(t)$ to obtain the bootstrap sample. (**b**) The motivation to compute the residuals as in *STEP 1* comes from the observation that, for any generic sequence $\{v(t)\}_{t=1}^{n}$, we have the equality

$$v(t) = \frac{1}{N^{1/2}} \sum_{s=-\tilde{n}}^{\tilde{n}} e^{it'\lambda_s} w_v(\lambda_s),$$

and then that by Lemmas 2 and 3 of Hidalgo (2009), we have that $w_x(\lambda_s) \simeq \Psi_{\theta_0}(\lambda_s) w_\varepsilon(\lambda_s)$, for all s.

STEP 2: The bootstrap analog of $\widehat{\vartheta} = \left(\widehat{\theta}', \widehat{\sigma}_\varepsilon^2\right)'$ is given by

$$\widehat{\vartheta}^* = \widehat{\vartheta} - \left(\sum_{\lambda_s} \phi_{\widehat{\vartheta}}(\lambda_s)\phi'_{\widehat{\vartheta}}(\lambda_s)\right)^{-1} \frac{\partial}{\partial\vartheta}\mathcal{Q}_N^*\left(\widehat{\vartheta}\right), \tag{3.2}$$

where

$$\mathcal{Q}_N^*(\vartheta) = \frac{1}{N}\sum_{\lambda_s}\left\{\log f_\vartheta(\lambda_s) + \frac{I_{x^*}^T(\lambda_s)}{(2\pi)^d f_\vartheta(\lambda_s)}\right\}. \tag{3.3}$$

Remark 2. We can replace the estimator $\widehat{\vartheta}^*$ in (3.2) by

$$\widehat{\vartheta}^* = \arg\min_{\vartheta\in\Theta\times\mathbb{R}^+} \mathcal{Q}_N^*(\vartheta).$$

However, for computational simplicity, see Shao and Tu (1995, p. 228, 336), we keep our definition of $\widehat{\vartheta}^*$ in (3.2).

STEP 3: Compute the bootstrap $T_p - process$ $\alpha_{\widehat{\theta}^*N}^*(\lambda)$, where

$$\alpha_{\theta N}^*(\lambda) = \frac{N^{1/2}}{2^{1/2}P_0}\left[\frac{G_{\theta N}^*(\lambda)}{G_{\theta N}^*(\pi)} - \frac{\lambda[1]}{\pi}\prod_{\ell=2}^{d}\left(1+\frac{\lambda[\ell]}{\pi}\right)\right], \quad \lambda\in\widetilde{\Pi}^d, \tag{3.4}$$

with $G_{\theta N}^*(\lambda) = (2\pi)^d N^{-1}\sum_{\lambda_s}^{\lambda} |\Psi_\theta(\lambda_s)|^{-2} I_{x^*}^T(\lambda_s)$. Finally we compute the bootstrap analog of $\beta_{\widehat{\theta}N}$, $\beta_{\widehat{\theta}^*N}^*$, as

$$\begin{aligned}\beta_{\theta N}^*(\lambda) &= \mathcal{L}_{\theta N}\left(\alpha_{\theta N}^*(\lambda)\right)\\ &= \frac{1}{G_{\theta N}^*(\pi)}\frac{2^{1/2}}{N^{1/2}}\sum_{\lambda_s}^{\lambda} e_{\theta N}^*(\lambda_s), \quad \lambda\in\widetilde{\Pi}^d\end{aligned}$$

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

with

$$e^*_{\theta N}(\lambda_s) = \mathring{I}^T_{x^*,\theta}(\lambda_s) - \widetilde{\varphi}'_\theta(\lambda_s)\, b^*_{\theta N}(\lambda_s),$$
$$b^*_{\theta N}(\lambda_s) = \widetilde{\Lambda}^{-1}_{\theta N}(\lambda_s)\frac{1}{\boldsymbol{J}}\sum_{\lambda_k \prec \lambda_s} \widetilde{\varphi}_{\theta N}(\lambda_k)\, \mathring{I}^T_{x^*,\theta}(\lambda_k),$$

are the recursive residuals in the linear projection of $\left\{\mathring{I}^T_{x^*,\theta}(\lambda_k)\right\}_{0\prec\lambda_k\prec\lambda_s}$ on $\{1, \varphi_\theta(\lambda_k)\}_{0\prec\lambda_k\prec\lambda_s}$ with

$$\mathring{I}^T_{x^*,\theta}(\lambda_s) = \frac{I^T_{x^*}(\lambda_s)}{|\Psi_\theta(\lambda_s)|^2} - \frac{1}{\boldsymbol{J}}\sum_{\lambda_k\prec\lambda_s}\frac{I^T_{x^*}(\lambda_k)}{|\Psi_\theta(\lambda_k)|^2}.$$

With $G^{0*}_N(\lambda) = (2\pi)^d N^{-1}\sum_{\lambda_s}^{\lambda} \check{I}^T_{\varepsilon^*}(\lambda_s)$, $\lambda \in \widetilde{\Pi}^d$, let β^{0*}_N be as

$$\beta^{0*}_N(\lambda) = \frac{1}{G^{0*}_N(\pi)}\frac{2^{1/2}}{N^{1/2}}\sum_{\lambda_s}^{\lambda} e^*_N(\lambda_s), \quad \lambda \in \widetilde{\Pi}^d,$$

where

$$e^*_N(\lambda_s) = \check{I}^T_{\varepsilon^*}(\lambda_s) - \widetilde{\varphi}'_{\widehat{\theta}N}(\lambda_s)\, b^*_N(\lambda_s),$$
$$b^*_N(\lambda_s) = \widetilde{\Lambda}^{-1}_{\widehat{\theta}N}(\lambda_s)\frac{1}{\boldsymbol{J}}\sum_{\lambda_k\prec\lambda_s}\widetilde{\varphi}_{\widehat{\theta}N}(\lambda_k)\,\check{I}^T_{\varepsilon^*}(\lambda_k).$$

Here $\check{I}^T_{\varepsilon^*}(\lambda_s) = I^T_{\varepsilon^*}(\lambda_s) - \left(\sum_{\lambda_k\prec\lambda_s} I^T_{\varepsilon^*}(\lambda_k)\right)/\boldsymbol{J}$. Let the notation $\overset{*}{\Rightarrow}$ (and $o_{p^*}(1)$, $\xrightarrow{d^*}$, etc.) indicate respectively weak convergence (and convergence in probability, distribution, etc.) of a bootstrap statistic conditional on the observed data.

THEOREM 4. *Under the maintained hypothesis and C1 − C9, we have that*

(*a*) $\beta^{0*}_N(\lambda) \overset{*}{\Rightarrow} \boldsymbol{B}(\lambda/\pi), \quad \lambda \in \widetilde{\Pi}^d$ *in probability.*

(*b*) $sup_{\lambda\in\widetilde{\Pi}^d}\left|\beta^*_{\widehat{\theta}^*N}(\lambda) - \beta^{0*}_N(\lambda)\right| = o_{p^*}(1).$

COROLLARY 2. *Let $\eta : D\left(\widetilde{\Pi}^d\right) \to \mathbb{R}^+$ be as in Corollary 1. Then, under the maintained hypothesis and conditions in Theorem 4, we have that*

$$\eta\left(\beta^*_{\widehat{\theta}^*N}\right) \xrightarrow{d^*} \eta(\boldsymbol{B}) \quad \text{in probability.}$$

Proof. The proof is an immediate consequence of Theorem 4 and the continuous mapping theorem, so it is omitted. ■

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

4. EXTENSION TO REGRESSION MODELS AND LOCAL ALTERNATIVES

The aim of this section is twofold. On the one hand, we would like to describe the consequences when the sequence $\{x(t)\}_{t=1}^{n}$ is not observable but they are the errors of a parametric regression model. The second aim of this section is to describe the type of local alternatives that $\eta\left(\beta_{\widehat{\vartheta}N}\right)$ is able to detect.

4.1. Regression Models

With regard to our first aim. Let's consider the model in (1.1), that is

$$y(t) = \beta_0' z(t) + x(t), \qquad t = 1, \ldots, n, \tag{4.1}$$

where $z(t)$ is the q-dimensional regressor. Recall that as we have excluded the frequency $\lambda = 0$ in the computation of $\eta\left(\beta_{\widehat{\vartheta}N}\right)$, we have effectively covered in the previous section the scenario when $z(t) = 1$. In our present context and denoting by $\widehat{\beta}$ the least squares estimator of β_0, the test becomes $\eta\left(\beta_{\widetilde{\vartheta}N}\right)$, where

$$\widetilde{\vartheta} = \arg \min_{\vartheta \in \Theta \times \mathbb{R}^+} \mathcal{Q}_N(\vartheta)$$

with

$$\mathcal{Q}_N(\vartheta) = \frac{1}{N} \sum_{\lambda_s} \left\{ \log f_\vartheta(\lambda_s) + \frac{I_{\widehat{x}}^T(\lambda_s)}{(2\pi)^d f_\vartheta(\lambda_s)} \right\}$$

and $\{\widehat{x}(t)\}_{t=1}^{n} = \left\{y(t) - \widehat{\beta}' z(t)\right\}_{t=1}^{n}$ is the set of the least squares residuals.

Before we state the asymptotic properties of the least squares estimator

$$\widehat{\beta} = \left(\sum_{t=1}^{n} z(t) z'(t) \right)^{-1} \sum_{t=1}^{n} z(t) y(t),$$

let's introduce the following condition denoted as Grenander condition on the deterministic regressors Z, which denotes the $n \times q$ matrix stacking $z(t)$'s.

***Grenander Condition*:** Let $z_s(t)$ denote the s-th element of the vector $z(t)$ and $A_n = diag\left(\sqrt{\sum_{t=1}^{n} z_s^2(t)}\right)_{s=1}^{q}$. Then, for all $s = 1, \ldots, q$, as $n \to \infty$,

(i) $\sum_{t=1}^{n} z_s^2(t) \to \infty$,

(ii) $\max_{1 \le u \le n} \frac{z_s^2(u)}{\sum_{t=1}^{n} z_s^2(t)} \to 0$,

(iii) $A_n^{-1} \sum_{t=s+1}^{n} z(t-s) z'(t) A_n^{-1} \to \mathcal{R}(s) = \int_{-\mathring{\pi}}^{\pi} e^{is'\lambda} \mathcal{M}(d\lambda)$,

where $\mathcal{M}(\lambda_2) - \mathcal{M}(\lambda_1)$ is a Hermitian nonnegative matrix and $\mathcal{R} = \mathcal{R}(0) > 0$ and $t - s = (t[\ell] - s[\ell])_{\ell=1}^{d}$.

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

Examples of deterministic sequences $\{z(t)\}_{t\in\mathbb{Z}^d}$ satisfying the Grenander's conditions are spatial-trend polynomials, see e.g., Section 3.4 of Cressie (1993). That is, in case of $d=2$,

$$z(t)=\left[t[1]^s t[2]^k\right]_{0\le s,k\le r}.$$

If $r=2$,

$$z(t)=\left(1,t[1],t[2],t[1]^2,t[2]^2,t[1]t[2]\right)' \tag{4.2}$$

and hence $q=6$. In this case, using that

$$\frac{1}{m^{\kappa+1}}\sum_{k=1}^{m}k^{\kappa}\underset{m\nearrow\infty}{\to}\frac{1}{\kappa+1},\quad \kappa>-1$$

we obtain

$$\mathcal{R}(s)=\begin{pmatrix}1 & 3^{1/2}/2 & 3^{1/2}/2 & 5^{1/2}/3 & 5^{1/2}/3 & 3/4\\ & 1 & 3/4 & 15^{1/2}/4 & 15^{1/2}/6 & 27^{1/2}/6\\ & & 1 & 15^{1/2}/6 & 15^{1/2}/4 & 27^{1/2}/6\\ & & & 1 & 15^{1/2}/9 & 45^{1/2}/8\\ & & & & 1 & 45^{1/2}/8\\ & & & & & 1\end{pmatrix}.$$

One consequence of $\mathcal{R}(s)$ being independent of s is that $\mathcal{M}(\lambda)$ has a jump at the origin, and the size of the jump is $\mathcal{R}$. That is,

$$\mathcal{M}(\lambda)=\begin{cases}0 \text{ if } \lambda[1]<0 \text{ or } \lambda[2]<0\\ \mathcal{R}(s)=\mathcal{R} \text{ if } \lambda[1]\ge 0 \text{ and } \lambda[2]\ge 0.\end{cases}$$

Let's now introduce a slightly milder condition on the spectral density function of the sequence $\{x(t)\}_{t\in\mathbb{Z}^2}$. Hereafter, we restrict our discussion to $d=2$ for the clarity of our exposition.

***C1'*:** (*a*) The *Generalized Linear process* $\{x(t)\}_{t\in\mathbb{Z}^2}$ in (1.3) has a spectral density function $f(\lambda)$, which is positive and piecewise continuous.

(*b*) The jumps of $\mathcal{M}(\lambda)$ do not coincide with the discontinuities of $f(\lambda)$.

We have then the following proposition.

PROPOSITION 1. *Under C1′, C2, and the Grenander conditions, we have that*

$$A_n\left(\widehat{\beta}-\beta_0\right)\to\mathcal{N}\left(0,\mathcal{R}^{-1}\int_{-\pi}^{\pi}f(\lambda)\mathcal{M}(d\lambda)\mathcal{R}^{-1}\right).$$

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

We now comment on the condition $C1'$ and the results on Proposition 1. First, we observe that $C1'$ indicates that the *Generalized linear process* $\{x(t)\}_{t\in\mathbb{Z}^2}$ does not need to satisfy the standard strong mixing conditions for the central limit theorem of the least squares to hold true. Moreover, the condition that $\sum_{s\in\mathbb{Z}^d}\psi^2(s)<\infty$ implies that it is possible to allow for long memory and still the results of the latter proposition hold. Of course, the conditions in Jenish and Prucha (2009) rule out long memory or jumps in the spectral density function, however they allow for nonlinear processes, say the errors $x(t)=g\left(\check{x}(t)\right)$, where $\check{x}(t)$ is a Generalized linear process. Recall that as we allow for the spectral density function to have jumps, due to results of Ibragimov and Rozanov (1978), it implies that $\{x(t)\}_{t\in\mathbb{Z}^2}$ cannot be strong-mixing. Moreover, our results improve those given in Mardia and Marshall (1984). Finally, the results of Proposition 1 indicate that the fast Fourier transform at λ_0 of $\{x(t)\}_{t=1}^n$ satisfies the central limit theorem if $\lambda_0\neq\breve{\lambda}$ where $\breve{\lambda}$ is a jump/discontinuity point of $f(\lambda)$.

From Proposition 1, we have the following corollary.

COROLLARY 3. *Under $C1'$ and $C1-C7$ and the Grenander conditions, we have that*

$$\widetilde{\vartheta}-\widehat{\vartheta}=O_p\left(N^{-1}\right).$$

So, the first conclusion we have is that the asymptotic distribution of the Whittle estimator of ϑ_0 is unaffected by using the residuals instead of the (un)observable $X=\{x(t)\}_{t=1}^n$ and that the asymptotic distribution of $A_n\left(\widehat{\beta}-\beta_0\right)$ and $N^{1/2}\left(\widetilde{\vartheta}-\vartheta_0\right)$ is independent.

Denote

$$\widehat{\alpha}_{\theta N}(\lambda)=\frac{1}{2^{d/2}P_4}N^{1/2}\left[\frac{\widehat{G}_{\theta N}(\lambda)}{\widehat{G}_{\theta N}(\pi)}-\frac{\lambda[1]}{\pi}\prod_{\ell=2}^{d}\left(1+\frac{\lambda[\ell]}{\pi}\right)\right],\quad\lambda\in\widetilde{\Pi}^d, \tag{4.3}$$

where

$$\widehat{G}_{\theta N}(\lambda)=\frac{(2\pi)^d}{N}\sum_{\lambda_s}^{\lambda}\frac{I_{\widehat{x}}^T(\lambda_s)}{|\Psi_\theta(\lambda_s)|^2} \tag{4.4}$$

are (2.4) and (2.2) but with the residuals $\widehat{x}(t)$ instead of the errors $x(t)$. Similarly, introduce

$$\mathring{I}_{\widehat{x},\theta}^T(\lambda_s)=\frac{I_{\widehat{x}}^T(\lambda_s)}{|\Psi_\theta(\lambda_s)|^2}-\frac{1}{\boldsymbol{J}}\sum_{\lambda_k\prec\lambda_s}\frac{I_{\widehat{x}}^T(\lambda_k)}{|\Psi_\theta(\lambda_k)|^2},$$

and the recursive residuals in the linear projection of $\left\{\mathring{I}_{\widehat{x},\theta}^T(\lambda_k)\right\}_{0\prec\lambda_k\prec\lambda_s}$ on $\{1,\varphi_\theta(\lambda_k)\}_{0\prec\lambda_k\prec\lambda_s}$, that is,

$$\widehat{e}_{\theta N}(\lambda_s)=\mathring{I}_{\widehat{x},\theta}^T(\lambda_s)-\widehat{\varphi}_{\theta N}'(\lambda_s)\widehat{b}_{\theta N}(\lambda_s),$$

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

where $\widehat{b}_{\theta N}(\lambda_s) = \widetilde{\Lambda}_{\theta N}^{-1}(\lambda_s)\frac{1}{N}\sum_{\lambda_k \prec \lambda_s}\widetilde{\varphi}_{\theta N}(\lambda_k)\mathring{I}_{\widehat{x},\theta}^{T}(\lambda_k)$. Then, the martingale transformation becomes

$$\widehat{\beta}_{\theta N}(\lambda) = \frac{1}{\widehat{G}_{\theta N}(\pi) P_4}\frac{2^{d/2}}{N^{1/2}}\sum_{\lambda_s}^{\lambda}\widehat{e}_{\theta N}(\lambda_s), \quad \lambda \in \widetilde{\Pi}^d.$$

With the help of Corollary 3, we have the following theorem:

THEOREM 5. *Under C*1 − *C*9 *and the Grenander conditions, we have that*

(*a*) $\sup_{\lambda\in\widetilde{\Pi}^d}\left|\widehat{\alpha}_{\widehat{\theta}N}(\lambda) - \alpha_{\widehat{\theta}N}(\lambda)\right| = o_p(1)$

(*b*) $\sup_{\lambda\in\widetilde{\Pi}^d}\left|\widehat{\beta}_{\widehat{\theta}N}(\lambda) - \beta_{\widehat{\theta}N}(\lambda)\right| = o_p(1).$

So, the conclusion from Theorem 5 is that, up to first order asymptotics, the behavior of the test based on functionals of $\alpha_{\theta N}(\lambda)$ or $\beta_{\theta N}(\lambda)$ is unaltered. Furthermore, the bootstrap can be performed by applying the same algorithm as described in Section 3.1 to the regression residuals $\{\widehat{x}(t)\}_{t=1}^n$ due to the asymptotic independence implied by Corollary 3. Alternatively, we can add one more step between Steps 1 and 2. That is, do Step 1 with the regression residuals $\{\widehat{x}(t)\}_{t=1}^n$ and obtain $\{x^*(t)\}_{t=1}^n$. Next, generate $\left\{y^*(t) = z(t)'\widehat{\beta} + x^*(t)\right\}_{t=1}^n$ and compute the OLS residuals $\{\widehat{x}^*(t)\}_{t=1}^n$ by the OLS of $\{y^*(t)\}_{t=1}^n$ on $\{z(t)\}_{t=1}^n$. Finally, run Step 2 with $\{\widehat{x}^*(t)\}_{t=1}^n$ there.

4.2. Local Alternatives: Omnibus, Directional and Portmanteau Tests

The aim of this section is twofold. On the one hand, we want to investigate the relationship of our test in Section 3 with those based on, say, a Portmanteau scheme. On the other hand, we would like also to describe the type of local alternatives that $\eta\left(\beta_{\widehat{\theta}N}\right)$ is able to detect. In particular, we will see that $\eta\left(\beta_{\widehat{\theta}N}\right)$ is able to detect local alternatives of the type

$$H_{1N} : |\Psi(\lambda)|^2 = \left|\Psi_{\theta_0}(\lambda)\right|^2\left(1 + \tau\frac{l(\lambda)}{N^{1/2}} + \frac{s_N(\lambda)}{N}\right), \lambda \in \widetilde{\Pi}^d \text{ for some } \theta_0 \in \Theta,$$

where $l(\lambda)$ satisfies the same properties as φ_{θ_0} in *C*9, τ is a constant, possibly unknown, and for some finite N_0, $\sup_{N>N_0}|s_N(\cdot)|$ is an integrable function. Let us consider a couple of examples for $d = 2$.

Example 1

We wish to study departures of total independence (the white noise) hypothesis in the direction of the first-order isotropic conditional autoregressive (*CAR*) scheme

$$\mathbb{E}\{x(t)|\ldots\} = \frac{\theta_0}{N^{1/2}}\left(x(t-e_1) + x(t+e_1) + x(t-e_2) + x(t+e_2)\right).$$

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

In this case, we have that

$$\frac{|\Psi(\lambda)|^2}{\left|\Psi_{\theta_0}(\lambda)\right|^2} = 1 - 2\frac{\theta_0}{N^{1/2}}\{\cos(\lambda[1]) + \cos(\lambda[2])\},$$

so that $l(\lambda) = -2\{\cos(\lambda[1]) + \cos(\lambda[2])\}$ and $\tau = \theta_0$, and the remainder function $s_N(\lambda)$ being equal to zero.

Recall that the general CAR formulation, see Besag (1974), is given by

$$\mathbb{E}\{x(t)|x(r) : r \neq t\} = \sum_{s \in \mathbb{Z}^d \setminus \{0\}} \delta(s)\, x(t-s). \tag{4.5}$$

Example 2
Suppose now that we wish to study departures of total independence (white noise) hypothesis in the direction of a first-order (isotropic) simultaneous autoregressive (SAR) model, see Whittle (1954),

$$x(t) = \frac{\theta_0}{N^{1/2}}(x(t-e_1) + x(t+e_1) + x(t-e_2) + x(t+e_2)) + \varepsilon(t).$$

Then, we obtain that

$$\frac{|\Psi(\lambda)|^2}{\left|\Psi_{\theta_0}(\lambda)\right|^2} = 1 - 2\frac{\theta_0}{N^{1/2}}\{\cos(\lambda[1]) + \cos(\lambda[2])\} + \frac{\theta_0}{N} s_N(\lambda),$$

so that, we have that

$$l(\lambda) = -2\{\cos(\lambda[1]) + \cos(\lambda[2])\} \quad \text{and } \tau = \theta_0,$$

and $s_N(\lambda)$ is a function of $\cos(\lambda[1])$, $\cos(2\lambda[1])$, $\cos(\lambda[2])$, and $\cos(2\lambda[2])$, which satisfies that $|s_N(\lambda)| < C$.

Remark 3. It is worth mentioning that the class of CAR models is more general than the SAR models. In fact, as Cressie (1993, Chap. 6) observed, any SAR model has a CAR representation but not vice versa.

Now, for $\lambda \in \widetilde{\Pi}^d$, let us define

$$\boldsymbol{L}(\lambda) = \frac{1}{(2\pi)^d} \int_{-\mathring{\pi}}^{\lambda} \left\{ l(\overline{\lambda}) - \gamma'_{\theta_0}(\overline{\lambda})\, \Lambda_{\theta_0}^{-1}(\overline{\lambda})\, \frac{1}{(2\pi)^d} \int_{\widetilde{\lambda} \prec \overline{\lambda}} \gamma_{\theta_0}(\widetilde{\lambda})\, l(\widetilde{\lambda})\, d\widetilde{\lambda} \right\} d\overline{\lambda}, \tag{4.6}$$

where $\gamma'_{\theta_0}(\lambda) = \left(1, \varphi'_{\theta_0}(\lambda)\right)$ and $\Lambda_{\theta_0}(\lambda) = \int_{\widetilde{\lambda} \prec \overline{\lambda}} \gamma_{\theta_0}(\widetilde{\lambda})\, \gamma'_{\theta_0}(\widetilde{\lambda})\, d\widetilde{\lambda}$. Also denote

$$\boldsymbol{M}(\lambda) = \boldsymbol{B}(\lambda/\pi) + \tau \cdot \boldsymbol{L}(\lambda), \quad \lambda \in \widetilde{\Pi}^d.$$

Then, we have the following theorem.

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

THEOREM 6. *Assuming the same conditions of Theorem 3, under* H_{1N}, $\beta_{\widehat{\theta}N} \Rightarrow \boldsymbol{M}$.

Proof. The proof follows by Theorem 3 and standard arguments, so it is omitted. ■

As usual, $\boldsymbol{L}(\lambda) \neq 0$ implies that under the alternative our test develops a mean function, $N^{1/2}\boldsymbol{L}(\lambda)$ which clearly increases to infinity in absolute value. It is obvious from (4.6) that $\boldsymbol{L}(\lambda)$ will be different than zero in a set with positive Lebesgue measure unless $l(\lambda)$ is a constant or $\varphi_{\theta_0}(\lambda)$. If so, then $H_{1N} \subseteq H_0$. Indeed, consider the model $\left|\Psi_{\theta_0+1/N^{1/2}}(\lambda)\right|^2$. Then, it is easy to see by Taylor's expansion that

$$\left|\Psi_{\theta_0+1/N^{1/2}}(\lambda)\right|^2 = \left|\Psi_{\theta_0}(\lambda)\right|^2\left(1+\tau\frac{l(\lambda)}{N^{1/2}}+\frac{s_N(\lambda)}{N}\right),$$

so that we can conclude that $H_{1N} \subseteq H_0$.

We now turn our attention to the omnibus and directional tests. When $d=1$, using the fact that $\boldsymbol{M}$ and $\boldsymbol{B}$ are identically distributed, except for the deterministic shift $\tau \cdot \boldsymbol{L}$, and taking into account that $2^{1/2}\sin((j-1/2)\lambda)$ and $1/(j-1/2)^2\pi^2$ are the eigenfunctions and eigenvalues in the Kac-Siegert representation of $\boldsymbol{B}(\lambda/\pi)$, the orthogonal components of $\boldsymbol{M}$

$$m(j) = 2^{1/2}\left(j-\frac{1}{2}\right)\int_0^\pi \sin\left(\left(j-\frac{1}{2}\right)\lambda\right)\boldsymbol{M}(\lambda)\,d\lambda, \quad j=1,2,\ldots,$$

are distributed as independent $\mathcal{N}(\tau \cdot v(j),1)$, where

$$v(j) = 2^{1/2}\left(j-\frac{1}{2}\right)\int_0^\pi \sin\left(\left(j-\frac{1}{2}\right)\lambda\right)\boldsymbol{L}(\lambda)\,d\lambda, \quad j=1,2,\ldots.$$

When $d \geq 1$, the previous formulae become

$$m(j) = 2^{d/2}\prod_{\ell=1}^d\left(j[\ell]-\frac{1}{2}\right) \times \int_{-\mathring{\pi}}^{\pi}\prod_{\ell=1}^d \sin\left(\left(j[\ell]-\frac{1}{2}\right)\lambda\right)\boldsymbol{M}(\lambda)\,d\lambda, \quad j=1,2,\ldots$$

which are distributed as independent $\mathcal{N}(\tau \cdot v(j),1)$, where

$$v(j) = 2^{d/2}\prod_{\ell=1}^d\left(j[\ell]-\frac{1}{2}\right) \times \int_{-\mathring{\pi}}^{\pi}\prod_{\ell=1}^d \sin\left(\left(j[\ell]-\frac{1}{2}\right)\lambda\right)\boldsymbol{L}(\lambda)\,d\lambda, \quad j=1,2,\ldots.$$

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

Using, the (asymptotically) orthogonal components of $\beta_{\widehat{\theta}N}$, for $j = 1, 2, \ldots$,

$$\widetilde{m}_N(j) = 2^{d/2} \prod_{\ell=1}^{d} \left(j[\ell] - \frac{1}{2} \right) \int_{-\vec{\pi}}^{\pi} \prod_{\ell=1}^{d} \sin\left(\left(j[\ell] - \frac{1}{2} \right) \lambda \right) \beta_{\widehat{\theta}N}(\lambda)\, d\lambda,$$

we obtain the spectral representation

$$\beta_{\widehat{\theta}N}(\lambda) = 2^{d/2} \sum_{j=1}^{\infty} \widetilde{m}_N(j) \prod_{\ell=1}^{d} \frac{\sin\left(\left(j[\ell] - \frac{1}{2} \right) \lambda \right)}{\pi \left(j[\ell] - \frac{1}{2} \right)}, \quad \lambda \in \widetilde{\Pi}^d.$$

By Theorem 3 and the continuous mapping theorem, we have that finitely many of the $\widetilde{m}_N(j)'s$ converge in distribution to the corresponding $m(j)'s$ under H_{1N}, for the Cramer–von Mises, using Parseval's Theorem, we obtain

$$\widehat{C}_N \xrightarrow{d} \sum_{j=1}^{\infty} \frac{m^2(j)}{\pi^{2d} \prod_{\ell=1}^{d} \left(j[\ell] - \frac{1}{2} \right)^2}.$$

On the other hand, similar arguments to those in Eubank and LaRicca (1992) imply that for a reasonable choice of $q \geq 1$, tests based on

$$\widetilde{W}_{qN} = \sum_{j=1}^{q} \widetilde{m}_N^2(j)$$

will lead to gains in power, compared to $\widehat{C}_N$, in the direction of alternatives with significant autocorrelations at high lags. These Portmanteau tests are related to Neyman's (1937) smooth tests, a compromise between omnibus and directional tests, and for each $q \geq 1$, under H_{1N}, we have that

$$\widetilde{W}_{qN} \xrightarrow{d} \chi_q^2 \left(\tau^2 \sum_{j=1}^{q} \upsilon^2(j) \right).$$

That is, tests based on $\widetilde{W}_{qN}$ are asymptotically pivotal under H_0 ($\tau = 0$) for each choice of q, and more importantly, they are able to detect local alternatives converging to the null at the parametric rate $N^{-1/2}$, provided that $\upsilon(j) \neq 0$ for some $j = 1, \ldots, n$. The latter is in contrast with the classical Portmanteau tests based on

$$\widetilde{Q}_{q_N N} = N \sum_{j=1}^{q_N} \widetilde{\rho}_N^2(j),$$

where $\widetilde{\rho}_N(j)$ is some estimate of the "$j-th$" autocorrelation of $\{\varepsilon(t)\}_{t \in \mathbb{Z}^d}$. It can be shown (as in the case $d = 1$) that $\widetilde{Q}_{q_N N}$ is approximately distributed as a

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

$\chi^2_{q_N-p}$ under H_0 and assuming that q_N diverges with $\vec{n}$. However, the resulting test is able to detect alternatives converging to the null at the rate $q_N^{1/4}N^{-1/2}$ when $d=1$, which is slower than the rate $N^{-1/2}$ of our tests. Moreover, the performance of the test can be quite sensitive to the choice of q_N as a particular choice of q_N for which the level of the test is close to the nominal one, it turns out that particular choice delivers a test with low power.

In practice, one might recommend to use the discrete version $\widehat{W}_{qN} = \sum_{j=1}^{q} \widehat{m}_N^2(j)$ of $\widetilde{W}_{qN}$, with

$$\widehat{m}_N(j) = 2^{d/2}\left\{\prod_{\ell=1}^{d}\left(j[\ell]-\frac{1}{2}\right)\right\}\frac{\pi^d}{N} \times \sum_{k=-\tilde{n}}^{\tilde{n}}\prod_{\ell=1}^{d}\sin\left(\left(j[\ell]-\frac{1}{2}\right)\frac{\pi k}{\tilde{n}}\right)\beta_{\widehat{\theta}N}\left(\frac{\pi k}{\tilde{n}}\right).$$

Next, optimal tests of H_0 in the direction H_{1N} can be constructed applying results in Grenander (1950), as was suggested by Stute (1997) in the context of goodness-of-fit testing of a regression function. Asymptotically, testing for H_0 in the direction of H_{1N} is equivalent to test $\overline{H}_0 : \mathbb{E}(m(j)) = 0$ for all $j \in \mathbb{Z}^d$, against $\overline{H}_1 : \mathbb{E}(m(j)) = \tau \cdot v(j)$ for all $j \geq 1$ with L known, but maybe with unknown τ. Under $\overline{H}_0$, the distribution of $\{m(j)\}_{j\in\mathbb{Z}^d}$ is completely specified, as is also under $\overline{H}_1$ when the parameter τ is known. Then we can compute an (asymptotically) optimal Neyman–Pearson test in the direction of H_{1N} based on the first q orthogonal components of $\beta_{\widehat{\theta}N}$, using the test statistic

$$\widehat{\psi}_{qN} = \frac{\sum_{j=1}^{q} v(j)\widehat{m}_N(j)}{\left(\sum_{j=1}^{n} v^2(j)\right)^{1/2}}.$$

Arguing as in Schoenfeld's (1977, Thm. 3), it can be shown the convergence in distribution of $\widehat{\psi}_{q_N N}$ when q_N increases with N. Approximately optimal tests for H_0 in the direction of H_{1N} reject H_0 at the α significance level when $\left|\widehat{\psi}_{q_N N}\right| > z_{1-\alpha/2}$ if τ has unknown sign, $\widehat{\psi}_{q_N N} > z_{1-\alpha}$ when $\tau > 0$ and $\widehat{\psi}_{q_N N} < -z_{1-\alpha}$ when $\tau < 0$.

5. MONTE CARLO EXPERIMENT

We examine the finite sample performance of our tests. In particular, we compare Cramer–von Mises tests based on the T_p-process $\alpha_{\widehat{\theta}N}$ and the martingale transformed process $\beta_{\widehat{\theta}N}$. Because the test based on $\alpha_{\widehat{\theta}N}$ is not pivotal, its critical value is computed via bootstrap algorithms. On the other hand, for the test based on the martingale transformation $\beta_{\widehat{\theta}N}(\lambda)$ we employ both the asymptotic critical values as well as those from its bootstrap approach. For all the specifications and sample sizes considered in the experiment, the number of Monte Carlo

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

simulations is 1000. However, to simplify and speed up the computations, we have the bootstrap distribution G_n^* be approximated by the WARP algorithm (Giacomini, Politis, and White, 2013). The WARP algorithm permits to approximate the Monte Carlo distribution of the bootstrap test generating only one additional bootstrap replication for each Monte Carlo sample, $\mathcal{X}_{n,m}^{*(1)}$, $m = 1, \ldots, 1000$. Then the empirical distribution of all 1000 bootstrap resamples of our statistic of interest from every independent replication is used jointly to approximate the distribution of the bootstrap test. The results are denoted by T_p, $\widehat{C}_N^*$ and $\widehat{C}_N$, respectively in the Tables 5.1 through 5.4 below.

Three different models are considered for $\{x(t)\}_{t\in\mathbb{Z}}$ as competing models. These models are the first- and second-order simultaneous autoregressive model and the first-order simultaneous moving average model, denoted by $SAR(1)$, $SAR(2)$, and $SMA(1)$, respectively. Specifically, for $d = 2$, they are

$$SAR(1):$$
$$x(t) = \theta(x(t-e_1) + x(t+e_1) + x(t-e_2) + x(t+e_2)) + \varepsilon(t),$$
$$SAR(2):$$
$$x(t) = \theta(x(t-2e_1) + x(t+2e_1) + x(t-2e_2) + x(t+2e_2)) + \varepsilon(t),$$
$$SMA(1):$$
$$x(t) = \theta(\varepsilon(t-e_1) + \varepsilon(t+e_1) + \varepsilon(t-e_2) + \varepsilon(t+e_2)) + \varepsilon(t),$$

where $\varepsilon(t)$ is an independent and identically distributed (iid) mean zero sequence in $\mathbb{Z}^2$. For all the three specifications, we have considered $\theta = 0, 0.1$, and 0.2 with sample sizes $n = (20, 20), (20, 40)$, and $(40, 40)$. Note that the white noise model is included in our specification by choosing $\theta = 0$. We consider two cases. First, we observe $\{x(t)\}_{t=1}^n$ directly and second, we observe $\{y(t), z(t)\}$ as specified in Section 4.1.

The type I error is examined using three null models, namely the white noise model, $SAR(1)$ and $SMA(1)$ with $\theta = 0.1$ and 0.2. The white noise model is estimated under both $SAR(1)$ and $SMA(1)$ specifications.

Table 1 reports the rejection frequencies of the three tests for three different significance levels, $0.1, 0.05$, and 0.01. The true data generating processes are indicated in each panel and the white noise cases are indicated by $SMA(1)$ and $SAR(1)$, respectively, depending on which model is used in the estimation. The outcome of the Monte-Carlo experiment seems to indicate that our procedure performs reasonable well. All the tests exhibit rejection rates similar to corresponding levels for all the scenarios. The bootstrap test, $\widehat{C}_N^*$, appears to be more conservative than its corresponding asymptotic one $\widehat{C}_N$, while there is some variation in the performance of the T_p test across different scenarios. All the results seem to be within Monte Carlo error band.

Table 2 reports empirical powers of the tests. We considered three scenarios. In the first one, we generated the sample from a $SMA(1)$ process but we wrongly estimated a $SAR(1)$ model. The second scenario we generated a $SAR(1)$ process

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

TABLE 1. Size of the tests

		White noise					
		$SMA(1)$			$SAR(1)$		
n	Level	T_p	$\hat{C}_N^*$	$\hat{C}_N$	T_p	$\hat{C}_N^*$	$\hat{C}_N$
(20, 20)	0.1	0.119	0.098	0.105	0.087	0.097	0.106
	0.05	0.046	0.043	0.065	0.043	0.047	0.065
	0.01	0.016	0.012	0.02	0.015	0.01	0.022
(20, 40)	0.1	0.102	0.103	0.121	0.097	0.108	0.121
	0.05	0.048	0.052	0.074	0.045	0.057	0.082
	0.01	0.015	0.008	0.021	0.006	0.012	0.021
(40, 40)	0.1	0.089	0.111	0.123	0.097	0.107	0.123
	0.05	0.055	0.078	0.076	0.057	0.068	0.078
	0.01	0.005	0.024	0.020	0.012	0.006	0.020
		$SMA(1)$					
		$\theta = 0.1$			$\theta = 0.2$		
n	Level	T_p	$\hat{C}_N^*$	$\hat{C}_N$	T_p	$\hat{C}_N^*$	$\hat{C}_N$
(20, 20)	0.1	0.110	0.095	0.107	0.082	0.079	0.088
	0.05	0.054	0.050	0.064	0.044	0.044	0.052
	0.01	0.019	0.015	0.021	0.005	0.008	0.009
(20, 40)	0.1	0.093	0.104	0.126	0.102	0.109	0.112
	0.05	0.045	0.037	0.066	0.047	0.048	0.061
	0.01	0.008	0.008	0.015	0.005	0.013	0.018
(40, 40)	0.1	0.088	0.082	0.101	0.111	0.103	0.116
	0.05	0.042	0.044	0.052	0.051	0.052	0.066
	0.01	0.011	0.017	0.021	0.011	0.013	0.021
		$SAR(1)$					
		$\theta = 0.1$			$\theta = 0.2$		
n	Level	T_p	$\hat{C}_N^*$	$\hat{C}_N$	T_p	$\hat{C}_N^*$	$\hat{C}_N$
(20, 20)	0.1	0.101	0.099	0.121	0.079	0.104	0.092
	0.05	0.047	0.046	0.066	0.038	0.048	0.048
	0.01	0.009	0.010	0.021	0.008	0.012	0.012
(20, 40)	0.1	0.106	0.092	0.104	0.095	0.092	0.098
	0.05	0.067	0.053	0.055	0.050	0.044	0.059
	0.01	0.015	0.018	0.025	0.010	0.004	0.025
(40, 40)	0.1	0.105	0.112	0.111	0.110	0.103	0.109
	0.05	0.049	0.05	0.055	0.059	0.055	0.064
	0.01	0.01	0.004	0.011	0.011	0.007	0.015

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

TABLE 2. Power of the tests

		$SMA(1)$					
		$\theta = 0.1$			$\theta = 0.2$		
n	Level	T_p	$\hat{C}_N^*$	$\hat{C}_N$	T_p	$\hat{C}_N^*$	$\hat{C}_N$
(20, 20)	0.1	0.119	0.125	0.143	0.270	0.320	0.334
	0.05	0.063	0.054	0.088	0.158	0.211	0.225
	0.01	0.021	0.014	0.033	0.044	0.090	0.100
(20, 40)	0.1	0.165	0.130	0.151	0.394	0.405	0.460
	0.05	0.096	0.079	0.102	0.233	0.294	0.342
	0.01	0.025	0.017	0.046	0.089	0.173	0.205
(40, 40)	0.1	0.154	0.149	0.160	0.493	0.685	0.705
	0.05	0.104	0.083	0.094	0.328	0.569	0.556
	0.01	0.031	0.023	0.033	0.145	0.323	0.345
		$SAR(1)$					
		$\theta = 0.1$			$\theta = 0.2$		
n	Level	T_p	$\hat{C}_N^*$	$\hat{C}_N$	T_p	$\hat{C}_N^*$	$\hat{C}_N$
(20, 20)	0.1	0.109	0.118	0.128	0.431	0.597	0.601
	0.05	0.047	0.066	0.074	0.320	0.443	0.496
	0.01	0.007	0.014	0.022	0.086	0.243	0.334
(20, 40)	0.1	0.106	0.118	0.127	0.704	0.793	0.821
	0.05	0.050	0.071	0.074	0.547	0.695	0.733
	0.01	0.004	0.014	0.022	0.267	0.476	0.592
(40, 40)	0.1	0.088	0.136	0.149	0.917	0.977	0.980
	0.05	0.042	0.077	0.087	0.831	0.955	0.961
	0.01	0.004	0.027	0.027	0.618	0.866	0.891
		$SAR(2)$					
		$\theta = 0.1$			$\theta = 0.2$		
n	Level	T_p	$\hat{C}_N^*$	$\hat{C}_N$	T_p	$\hat{C}_N^*$	$\hat{C}_N$
(20, 20)	0.1	0.323	0.139	0.153	0.994	0.874	0.872
	0.05	0.204	0.070	0.086	0.926	0.662	0.682
	0.01	0.101	0.014	0.025	0.601	0.124	0.308
(20, 40)	0.1	0.509	0.216	0.262	1.000	0.998	0.998
	0.05	0.361	0.090	0.152	1.000	0.980	0.988
	0.01	0.124	0.010	0.036	0.996	0.628	0.858
(40, 40)	0.1	0.811	0.458	0.541	1.000	1.000	1.000
	0.05	0.543	0.231	0.306	1.000	1.000	1.000
	0.01	0.205	0.067	0.095	1.000	1.000	1.000

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

TABLE 3. Size of the tests (from the residuals)

		White noise					
		$SMA(1)$			$SAR(1)$		
n	Level	T_p	$\hat{C}_N^*$	$\hat{C}_N$	T_p	$\hat{C}_N^*$	$\hat{C}_N$
(20, 20)	0.1	0.101	0.102	0.106	0.091	0.093	0.107
	0.05	0.049	0.053	0.062	0.050	0.042	0.065
	0.01	0.018	0.011	0.021	0.014	0.011	0.023
(20, 40)	0.1	0.100	0.098	0.118	0.094	0.110	0.116
	0.05	0.055	0.041	0.064	0.048	0.054	0.062
	0.01	0.009	0.009	0.022	0.007	0.023	0.023
(40, 40)	0.1	0.093	0.090	0.105	0.104	0.084	0.106
	0.05	0.039	0.051	0.055	0.044	0.047	0.056
	0.01	0.013	0.020	0.024	0.015	0.019	0.023
		$SMA(1)$					
		$\theta = 0.1$			$\theta = 0.2$		
n	Level	T_p	$\hat{C}_N^*$	$\hat{C}_N$	T_p	$\hat{C}_N^*$	$\hat{C}_N$
(20, 20)	0.1	0.086	0.086	0.086	0.084	0.060	0.090
	0.05	0.043	0.032	0.046	0.033	0.025	0.039
	0.01	0.005	0.009	0.011	0.003	0.006	0.011
(20, 40)	0.1	0.109	0.093	0.107	0.106	0.094	0.106
	0.05	0.047	0.048	0.064	0.049	0.045	0.064
	0.01	0.009	0.008	0.014	0.007	0.009	0.023
(40, 40)	0.1	0.088	0.108	0.119	0.128	0.093	0.109
	0.05	0.047	0.062	0.077	0.057	0.047	0.056
	0.01	0.014	0.012	0.018	0.012	0.011	0.020
		$SAR(1)$					
		$\theta = 0.1$			$\theta = 0.2$		
n	Level	T_p	$\hat{C}_N^*$	$\hat{C}_N$	T_p	$\hat{C}_N^*$	$\hat{C}_N$
(20, 20)	0.1	0.105	0.097	0.119	0.082	0.093	0.093
	0.05	0.048	0.051	0.067	0.032	0.043	0.054
	0.01	0.009	0.010	0.019	0.008	0.011	0.012
(20, 40)	0.1	0.101	0.097	0.105	0.116	0.098	0.111
	0.05	0.047	0.043	0.063	0.056	0.057	0.068
	0.01	0.007	0.016	0.020	0.014	0.015	0.024
(40, 40)	0.1	0.107	0.083	0.094	0.096	0.099	0.117
	0.05	0.054	0.037	0.047	0.048	0.035	0.072
	0.01	0.010	0.008	0.014	0.011	0.008	0.018

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

TABLE 4. Power of the tests (from the residuals)

		$SMA(1)$					
		$\theta = 0.1$			$\theta = 0.2$		
n	Level	T_p	$\hat{C}_N^*$	$\hat{C}_N$	T_p	$\hat{C}_N^*$	$\hat{C}_N$
(20, 20)	0.1	0.113	0.106	0.116	0.252	0.320	0.323
	0.05	0.061	0.062	0.071	0.167	0.203	0.234
	0.01	0.013	0.014	0.022	0.053	0.056	0.110
(20, 40)	0.1	0.136	0.148	0.151	0.343	0.440	0.466
	0.05	0.060	0.078	0.081	0.246	0.320	0.356
	0.01	0.014	0.010	0.033	0.096	0.126	0.203
(40, 40)	0.1	0.147	0.162	0.175	0.510	0.684	0.721
	0.05	0.073	0.073	0.092	0.387	0.543	0.575
	0.01	0.027	0.022	0.035	0.217	0.286	0.379
		$SAR(1)$					
		$\theta = 0.1$			$\theta = 0.2$		
n	Level	T_p	$\hat{C}_N^*$	$\hat{C}_N$	T_p	$\hat{C}_N^*$	$\hat{C}_N$
(20, 20)	0.1	0.102	0.122	0.124	0.378	0.581	0.581
	0.05	0.043	0.071	0.070	0.249	0.425	0.470
	0.01	0.009	0.014	0.019	0.084	0.194	0.314
(20, 40)	0.1	0.068	0.111	0.124	0.681	0.762	0.811
	0.05	0.029	0.059	0.074	0.542	0.682	0.714
	0.01	0.003	0.008	0.018	0.275	0.436	0.565
(40, 40)	0.1	0.068	0.142	0.142	0.923	0.969	0.975
	0.05	0.027	0.072	0.075	0.857	0.937	0.951
	0.01	0.003	0.009	0.018	0.611	0.882	0.905
		$SAR(2)$					
		$\theta = 0.1$			$\theta = 0.2$		
n	Level	T_p	$\hat{C}_N^*$	$\hat{C}_N$	T_p	$\hat{C}_N^*$	$\hat{C}_N$
(20, 20)	0.1	0.328	0.143	0.146	0.987	0.837	0.857
	0.05	0.180	0.079	0.088	0.918	0.580	0.682
	0.01	0.053	0.013	0.027	0.580	0.199	0.327
(20, 40)	0.1	0.554	0.232	0.278	1.000	0.998	0.999
	0.05	0.365	0.121	0.159	1.000	0.968	0.987
	0.01	0.145	0.021	0.048	0.987	0.673	0.859
(40, 40)	0.1	0.851	0.482	0.530	1.000	1.000	1.000
	0.05	0.687	0.232	0.295	1.000	1.000	1.000
	0.01	0.327	0.030	0.079	1.000	1.000	1.000

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

but we estimated a $SMA(1)$ model; and finally in the third scenario we generated a $SAR(2)$ model but we estimated a $SAR(1)$.

We can signal out some features of the tests. First, the power of each test increases as the sample size increases excluding some exception in the T_p test when $\theta = 0.1$; second, the power also increases as the alternative model deviates more from the null model; and third, it appears that neither of the tests dominates the others. The tests based on the transformed process have more power than the T_p test when the true data generating process is $SMA(1)$ or $SAR(1)$. On the other hand, the latter has more power than the former when it is $SAR(2)$. While the $\widehat{C}_N$ test shows more rejection than the $\widehat{C}_N^*$ test, it seems to be a reflection of the under-rejection tendency of the bootstrap test over the asymptotic test as noted in Table 1.

Finally, Tables 3 and 4 report the empirical sizes and powers of the test when $\{x(t)\}$ is the error sequence of the linear regression model described in Section 4.1. In particular, $z(t)$ is specified as in (4.2) and the true regression coefficients are set as $\beta_0 = (1, \ldots, 1)'$. As predicted by our theory, the error in estimating β_0 does not seem to affect the performance of our test much at least in our simulation design so that the discussion given for the previous tables apply here as well.

6. CONCLUSION

We have described and examined a distribution free test for the correct specification of the dynamics in a lattice model. The methodology employed extends existing one in the situation where the data follows a casual model developed in Delgado et al. (2005). To that end, we present a martingale-type transformation when the dimension of "time" could be greater than one, so that the asymptotic distribution of the test becomes just a functional of a standard Brownian sheet. We also look at a bootstrap analog in the spectral domain of the test showing its asymptotic validity. In addition we demonstrate that the asymptotic behavior of the test remains the same even if the process that we are concerned with is not observed but it is the error term in a linear trending regression model. Both the asymptotic and bootstrap tests seem to work well in small samples as demonstrated by a set of Monte Carlo simulations and, as anticipated from the theoretical results, there is no difference on whether we use the true errors or the residuals of the trending regression model. Moreover, the results are encouraging to practitioners in that the test has reasonable finite sample size when using the critical values from the asymptotic distribution. Finally one issue of interest, which we have not explored in the paper, is to examine the behavior of the least squares estimator of the parameters in the linear trending regression model given in (4.1), where the regressors $z(t)$ might take the form $\left[t[1]^{\alpha}, t[2]^{\beta}\right]$ with $\alpha < -1/2$ and/or $\beta < -1/2$. This would extend results in Robinson (2012). However, in this scenario we will need to develop new results for the asymptotic behavior of the estimators of the parameters and this goes beyond the scope of this paper.

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

REFERENCES

Baltagi, B., H.H. Kelejian, & I.R. Prucha (eds.) (2007) Analysis of spatially dependent data. Annals issue. *Journal of Econometrics*, Annals Issue, 140.

Batchelor, L.I.A. & H.S. Reed (1924) Relation of the variability of yields of fruit trees to the accuracy of field trials. *Journal of Agricultural Research* 12, 245–283.

Besag, J. (1974) Spatial interaction and the statistical analysis of lattice systems (with discussion). *Journal of the Royal Statistical Society, Series B* 36, 192–236.

Bickel, P.J. & M.J. Wichura (1971) Convergence criteria for multiparameter stochastic processes and its applications. *Annals of Mathematical Statistics* 42, 1656–1670.

Billingsley, P. (1968) *Convergence of Probability Measures*. Wiley.

Bloomfield, P. (1973) An exponential model for the spectrum of a scalar time series. *Biometrika* 61, 217–226.

Bolthausen, E. (1982) On the central limit theorem for stationary mixing random fields. *The Annals of Probability* 10, 1047–1050.

Brillinger, D.R. (1981) *Time Series, Data Analysis and Theory*. Holden-Day.

Brown, R.L., J. Durbin, & J.M. Evans (1975) Components of Cramér-von Mises statistics II. *Journal of the Royal Statistical Society, Series B* 37, 216–237.

Chen, H. & J.P. Romano (1999) Bootstrap-assisted goodness-of-fit tests in the frequency domain. *Journal of Time Series Analysis* 20, 619–654.

Cressie, N.A.C. (1993) *Statistics for Spatial Data*. Wiley.

Cressie, N. & H.C. Huang (1999) Classes of nonseparable, spatio-temporal stationary covariance functions. *Journal of the American Statistical Association* 94, 1330–1340.

Crujeiras, R.M., R. Fernandez-Casal, & W. Gonzalez-manteiga (2008) An L_2-test for comparing spatial spectral densities. *Statistics & Probability Letters* 78, 2543–2551.

Dahlhaus, R. & H. Künsch (1986) Edge effect and efficient parameter estimation for stationary random fields. *Biometrika* 74, 877–882.

Davis, R.A., A. Alexandre Trindade, & F.J. Breidt (2001) Least absolute deviation estimation for all-pass time series models. *Annals of Statistics* 29, 919–946.

Delgado, M.A., J. Hidalgo, & C. Velasco (2005) Distribution free goodness-of-fit tests for linear processes. *Annals of Statistics* 33, 2568–2609.

Delgado, M.A., J. Hidalgo, & C. Velasco (2011) Bootstrap assisted specification tests for the Farima model. *Econometric Theory* 27, 1083–1116.

Eubank, R.L. & V. LaRicca (1992) Asymptotic comparison of Cramér-von Mises and nonparametric techniques for testing goodness-of-fit. *Annals of Statistics* 20, 2071–2086.

Fernandez-Casal, R., W. Gonzalez-Manteiga, & M. Febrero-Bande (2003) Flexible spatio-temporal stationary variogram models. *Statistics and Computing* 13, 127–136.

Genton, M.G. & H.L. Koul (2008) Minimum distance inference in unilateral autoregressive lattice processes. *Statistica Sinica* 18, 617–631.

Giacomini, R., D.N. Politis, & H. White (2013) A warp-speed method for conducting Monte Carlo experiments involving bootstrap estimators. *Econometric Theory* 29, 567–609.

Grenander, U. (1950) Stochastic processes and statistical inference. *Arkiv för Matematik* 1, 195–277.

Guyon, X. (1982) Parameter estimation for a stationary process on a d-dimension lattice. *Biometrika* 69, 95–105.

Hainz, G. & R. Dahlhaus (2000) Spectral domain bootstrap tests for stationary time series. Preprint.

Hannan, E.J. (1970) *Multiple Time Series*. Wiley.

Hannan, E.J. (1973) The asymptotic theory of linear time series models. *Journal of Applied Probability* 10, 130–145.

Hidalgo, J. (2009) Goodness of fit for lattice processes. *Journal of Econometrics* 151, 113–128.

Hidalgo, J. & J.P. Kreiss (2006) Bootstrap specification tests for linear covariance stationary processes. *Journal of Econometrics* 133, 807–839.

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

Hong, Y. (1996) Consistent testing for serial-correlation of unknown form. *Econometrica* 64, 837–864.
Ibragimov, I.A. & Y.A. Rozanov (1978) *Gaussian Random Processes*. Springer-Verlag.
Jenish, N. & I.R. Prucha (2009) Central limit theorems and uniform laws of large numbers for arrays of random fields. *Journal of Econometrics* 150, 86–98.
Khmaladze, E.V. (1981) Martingale approach in the theory of goodness-of-fit tests. *Theory of Probability and Applications* 26, 240–257.
Lanne, M. & P. Saikonnen (2011) Noncausal autoregressive for economic time series. *Journal of Time Series Econometrics* 3, 1–30.
Lanne, M. & P. Saikonnen (2013) Noncausal vector autoregression. *Econometric Theory* 29, 447–481.
Mardia, K.V. & R.J. Marshall (1984) Maximum likelihood estimation of models for residual covariance in spatial regression. *Biometrika* 71, 135–146.
McKeague, I.W., A.M. Nikabadze, & Y. Sun (1995) An omnibus test for independence of survival time from a covariate. *Annals of Statistics* 223, 450–475.
Mercer, W.B. & A.D. Hall (1911) The experimental error of field trials. *Journal of Agricultural Science* 4, 107–132.
Mitchell, M.W., M.G. Genton, & M.L. Gumpertz (2005) Testing for separability of space–time covariances. *Environmetrics* 16, 819–831.
Neyman, J. (1937) Smooth test for goodness of fit. *Arkiv for matematik* 20, 150–199.
Paparoditis, E. (2000) Spectral density based goodness-of-fit tests for time series models. *Scandinavian Journal of Statistics* 27, 143–176.
Robinson, P.M. (1988) The stochastic difference between econometric statistics. *Econometrica* 56, 531–548.
Robinson, P.M. (2012) Inference on power law spatial trends. *Bernoulli* 18, 644–677.
Robinson, P.M. & S. Thawornkaiwong (2012) Statistical inference on regressing with spatial dependence. *Journal of Econometrics* 167, 521–542.
Robinson, P.M. & J. Vidal-Sanz (2006) Modified Whittle estimation of multilateral models on a lattice. *Journal of Multivariate Analysis* 60, 234–349.
Schoenfeld, D.A. (1977) Asymptotic properties of tests based on linear combinations of the Cramér-von Mises statistic. *Annals of Statistics* 5, 1017–1026.
Shao, J. & D. Tu (1995) *The Jackknife and Bootstrap*. Springer-Verlag.
Solo, V. (1986) Modeling of two-dimensional random field by parametric Cepstrum. *IEEE Transactions on Information Theory* 32, 743–750.
Stute, W. (1997) Nonparametric model checks for regression. *Annals of Statistics* 25, 613–641.
Whittle, P. (1954) On stationary processes in the plane. *Biometrika* 41, 434–449.
Yajima, Y. & Y. Matsuda (2011) Asymptotic properties of the LSE of a spatial regression in both weakly and strongly dependent stationary random fields. Preprint. University of Tokyo.
Zhu, J., H.C. Huang, & P.E. Reyes (2010) On selection of spatial linear models for lattice data. *Journal of the Royal Statistical Society, Series B* 72, 389–402.

APPENDIX A

We first introduce some more notation. For a generic function $g(\lambda)$, we abbreviate $g(\lambda_s)$ by g_s and C will denote a generic positive and finite constant. Then,

$$\sum_{v \le \lambda_s \le u} g_s = \sum_{\lambda_s \in \widetilde{\Pi}_n^d; v[\ell] \le \lambda_s[\ell] \le u[\ell], \forall \ell} g(\lambda_s),$$

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

for example. We also drop for simplicity any reference to "T" in ω_{ξ}^{T} or I_{ξ}^{T}, and we shall denote $\zeta(\lambda;\theta):\widetilde{\Pi}^{d}\times\Theta\to\mathbb{R}^{p}$ a function twice continuously differentiable in λ and θ, abbreviating $\zeta(\lambda;\theta_0)$ and $\zeta\left(\lambda;\widehat{\theta}\right)$ respectively by $\zeta(\lambda)$ and $\widehat{\zeta}(\lambda)$.

LEMMA 1. *Assume* $C1-C8$. *Then,*

(*a*) $\widehat{\vartheta}^{*}-\widehat{\vartheta}=o_{p^{*}}(1)$

(*b*) $$N^{1/2}\left(\widehat{\theta}^{*}-\widehat{\theta}\right)=\left(\frac{1}{N}\sum_{\lambda_s}\widetilde{\varphi}_{\widehat{\theta}s}\widetilde{\varphi}_{\widehat{\theta}s}'\right)^{-1}\frac{1}{N^{1/2}}\sum_{\lambda_s}\widetilde{\varphi}_{\widehat{\theta}s}I_{\varepsilon^{*}s}+o_{p^{*}}(1).$$

Proof. Part (*a*). The proof is quite immediate. Indeed, (3.3) is

$$\frac{1}{N}\sum_{\lambda_s}\frac{f_{\widehat{\vartheta}s}}{f_{\vartheta s}}\left(\frac{I_{x^{*}s}}{(2\pi)^{d}f_{\widehat{\vartheta}s}}-1\right)+\frac{1}{N}\left\{\sum_{\lambda_s}\frac{f_{\widehat{\vartheta}s}}{f_{\vartheta s}}-\log\frac{f_{\widehat{\vartheta}s}}{f_{\vartheta s}}+\log f_{\widehat{\vartheta}s}\right\}. \tag{A.1}$$

Now, the difference between the second term of $(A.1)$ and

$$\int_{-\mathring{\pi}}^{\pi}\left\{\frac{f_{\widehat{\vartheta}}(\lambda)}{f_{\vartheta}(\lambda)}-\log\left(\frac{f_{\widehat{\vartheta}}(\lambda)}{f_{\vartheta}(\lambda)}\right)\right\}d\lambda+\int_{-\mathring{\pi}}^{\pi}\log f_{\widehat{\vartheta}}(\lambda)\,d\lambda$$

converges to zero in probability using Brillinger (1981, p. 15) and that uniformly in λ, $\left|f_{\widehat{\vartheta}}(\lambda)-f_{\vartheta_0}(\lambda)\right|=o_p(1)$ by the mean value theorem and $C5$. Moreover, the last displayed expression is greater than or equal to $\frac{(2\pi)^{d}}{2}+\int_{-\mathring{\pi}}^{\pi}\log f_{\widehat{\vartheta}}(\lambda)\,d\lambda$ with equality when $f_{\widehat{\vartheta}}(\lambda)=f_{\vartheta}(\lambda)$ for all $\lambda\in\widetilde{\Pi}^{d}$, which is the case only if $\vartheta=\widehat{\vartheta}$ by $C6$. On the other hand, the first term of $(A.1)$ converges to zero uniformly in ϑ by Lemma 15 of Hidalgo (2009) because $f_{\vartheta}^{-1}(\lambda)f_{\widehat{\vartheta}}(\lambda)$ is a twice continuous differentiable function by $C5$. From here the conclusion of the lemma proceeds as in Theorem 1 of Hannan (1973), so we omit its details.

Part (*b*). It follows by an obvious extension of Lemma 14 of Hidalgo (2009), and thus it is omitted. ■

LEMMA 2. *Assume* $C1-C8$. *Under* H_0, *uniform in* $\lambda\in\widetilde{\Pi}^{d}$,

$$\frac{1}{N^{1/2}}\sum_{\lambda_s}^{\lambda}\zeta_s\left(\frac{I_{x^{*}s}}{\left|\Psi_{\widehat{\theta}^{*}s}\right|^{2}}-I_{\varepsilon^{*}s}\right)=-\left(\frac{1}{N}\sum_{\lambda_s}^{\lambda}\zeta_s\varphi_{\widehat{\theta}s}'\right)N^{1/2}\left(\widehat{\theta}^{*}-\widehat{\theta}\right)+o_{p^{*}}(1). \tag{A.2}$$

Proof. See Lemma 17 of Hidalgo (2009). ■

We now introduce the following notation. For $v_1\prec v_2\in\widetilde{\Pi}^{d}$, with $\dot{\varepsilon}^{*}(t)=h(t)\varepsilon^{*}(t)$,

$$\mathcal{E}_{1,N}^{*}(v_1,v_2)=\left(\frac{1}{N}\sum_{v_1\le\lambda_s\le v_2}\zeta_s\right)\left(\frac{N^{1/2}}{\sum_{t=1}^{n}h^{2}(t)}\sum_{t=1}^{n}\left(\dot{\varepsilon}^{*}(t)^{2}-h^{2}(t)\widehat{\sigma}_{\varepsilon}^{2}\right)\right) \tag{A.3}$$

$$\mathcal{E}_{2,N}^{*}(v_1,v_2)=\frac{1}{N}\sum_{v_1\le\lambda_s\le v_2}\zeta_s\frac{N^{1/2}}{\sum_{t=1}^{n}h^{2}(t)}\sum_{t_1\neq t_2=1}^{n}\dot{\varepsilon}^{*}(t_1)\dot{\varepsilon}^{*}(t_2)e^{i(t_1-t_2)'\lambda_s}. \tag{A.4}$$

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

Observe that $\mathcal{E}^*_{1,N}(v_1,v_2)+\mathcal{E}^*_{2,N}(v_1,v_2)=N^{-1/2}\sum_{v_1\le\lambda_s\le v_2}\zeta_s\left(I_{\varepsilon^* s}-\widehat{\sigma}^2_\varepsilon\right)$. Also for $\ell=1,\ldots,d$, we define

$$\mathcal{E}^{*(\ell)}_{1,N}(v_1[\ell],v_2[\ell])=\left(\frac{1}{n[\ell]}\sum_{s[\ell]=[\tilde{n}v_1[\ell]/\pi]}^{[\tilde{n}v_2[\ell]/\pi]}\zeta_{s[\ell]}\right)\times\left(\frac{N^{1/2}}{\sum_{t=1}^n h^2(t)}\sum_{t=1}^n\left(\dot{\varepsilon}^*(t)^2-h^2(t)\widehat{\sigma}^2_\varepsilon\right)\right)$$

$$\mathcal{E}^{*(\ell)}_{2,N}(v_1[\ell],v_2[\ell])=\frac{1}{N}\sum_{\lambda_s;[\tilde{n}v_1[\ell]/\pi]<s[\ell]<[\tilde{n}v_2[\ell]/\pi]}\zeta_s\frac{N^{1/2}}{\sum_{t=1}^n h^2(t)}\times\sum_{t_1\neq t_2=1}^n\dot{\varepsilon}^*(t_1)\dot{\varepsilon}^*(t_2)e^{i(t_1-t_2)'\lambda_s}.$$

We define $H_N(\cdot,\cdot)$ as a $O_p(1)$ sequence of random variables.

Next we prove that the processes $\left(\lambda[1]\prod_{\ell=2}^d(\pi+\lambda[\ell])\right)^{-v}\mathcal{E}^*_{c,N}(-\mathring{\pi}[\ell],\lambda)$, $c=1,2$, are tight for some value of $v>0$. From Bickel and Wichura (1971), it suffices to show the following lemma.

LEMMA 3. *Assuming C*1*, for any* $0\le v<1/4$ *and* $\ell=1,\ldots,d$,

$$(a)\ \mathbb{E}^*\left(\frac{\mathcal{E}^{*(\ell)}_{1,N}(-\mathring{\pi}[\ell],\lambda_1[\ell])}{\lambda_1^v[\ell]}-\frac{\mathcal{E}^{*(\ell)}_{1,N}(-\mathring{\pi}[\ell],\lambda_2[\ell])}{\lambda_2^v[\ell]}\right)^2=H_N(\lambda_1[\ell],\lambda_2[\ell])(\lambda_2[\ell]-\lambda_1[\ell])^{2-2v}$$

$$(b)\ \mathbb{E}^*\left(\frac{\mathcal{E}^{*(\ell)}_{2,N}(-\mathring{\pi}[\ell],\lambda_1[\ell])}{\lambda_1^v[\ell]}-\frac{\mathcal{E}^{*(\ell)}_{2,N}(-\mathring{\pi}[\ell],\lambda_2[\ell])}{\lambda_2^v[\ell]}\right)^4=H_N(\lambda_1[\ell],\lambda_2[\ell])(\lambda_2[\ell]-\lambda_1[\ell])^{2-4v}$$

for all $0<\lambda_1[1]<\lambda_2[1]<\pi$ *and* $-\pi<\lambda_1[\ell]<\lambda_2[\ell]<\pi$ *for* $\ell=2,\ldots,d$.

Proof. The proof proceeds, with standard modifications, as that of Lemma 9 of Hidalgo (2009) and thus it is omitted. ■

In what follows we shall abbreviate $\widetilde{\varphi}'_{\theta q}\widetilde{\Lambda}^{-1}_{\theta N}(\lambda_q)$ by $\Im_{\theta N}(q)$ and we write

$$\varkappa^*_s=\frac{I_{x^* s}}{\left|\Psi_{\widehat{\theta}^* s}\right|^2}-I_{\varepsilon^* s};\quad \jmath^*_s=I_{\varepsilon^* s}-\widehat{\sigma}^2_\varepsilon. \tag{A.5}$$

LEMMA 4. *Assuming C*1 − *C*9*, for all* $\varepsilon>0$*, in probability*

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

$$\lim_{\lambda_0 \to -\mathring{\pi}} \overline{\lim_{\vec{n} \to \infty}} \Pr{}^* \left\{ \sup_{-\mathring{\pi} \prec \lambda \preceq \lambda_0} \left| \frac{1}{N} \sum_{\lambda \preceq \lambda_k \preceq \lambda_0} \frac{\Im_{\widehat{\theta}N}(k)}{N^{1/2}} \sum_{\lambda_s \prec \lambda_k} \widetilde{\varphi}_{\widehat{\theta}N}(\lambda_s) \left(\varkappa_s^* + \jmath_s^*\right) \right| > \epsilon \right\} = 0. \tag{A.6}$$

Proof. Take $\lambda_0 \prec -\mathring{\pi}/2$ without loss of generality. The triangle inequality implies that

$$\begin{aligned} &\sup_{-\mathring{\pi} \preceq \lambda \preceq \lambda_0} \left| \frac{1}{N} \sum_{\lambda \preceq \lambda_k \preceq \lambda_0} \frac{\Im_{\widehat{\theta}N}(k)}{N^{1/2}} \sum_{\lambda_s \prec \lambda_k} \widetilde{\varphi}_{\widehat{\theta}N}(\lambda_s) \left(\varkappa_s^* + \jmath_s^*\right) \right| \\ &\quad \le \frac{C}{N} \sum_{\lambda \preceq \lambda_k \preceq \lambda_0} \left\| \Im_{\widehat{\theta}N}(k) \right\| g_N(k)^{\frac{\delta}{2}} \left\{ \sup_{-\tilde{n} \le k \le [\lambda_0 \tilde{n}]} \left\| \frac{g_N(k)^{-\frac{\delta}{2}}}{N^{1/2}} \sum_{\lambda_s \prec \lambda_k} \widetilde{\varphi}_{\widehat{\theta}N}(\lambda_s) \varkappa_s^* \right\| \right. \\ &\qquad \left. + \sup_{-\tilde{n} < k \le [\lambda_0 \tilde{n}]} \left\| \frac{g_N(k)^{-\frac{\delta}{2}}}{N^{1/2}} \sum_{\lambda_s \prec \lambda_k} \widetilde{\varphi}_{\widehat{\theta}N}(\lambda_s) \jmath_s^* \right\| \right\}, \end{aligned} \tag{A.7}$$

for any $0 < \delta < 1$, where $g_N(k) = N^{-1} (k[1]/n[1]) \prod_{\ell=2}^{d} |1 + k[\ell]/n[\ell]|$. First $C7$ implies that $\left\| \widetilde{\Lambda}_{\theta_0}(\lambda) \right\| \ge C^{-1} |\lambda_0[\ell] + \mathring{\pi}[\ell]|$ and hence because $\widehat{\theta} - \theta_0 = O_p\left(N^{-1/2}\right)$ we have that $\left\| \widetilde{\Lambda}_{\widehat{\theta}N}(\lambda_k) - \widetilde{\Lambda}_{\theta_0}(\lambda_k) \right\| = o_p(1)$. So,

$$\left\| \widetilde{\Lambda}_{\widehat{\theta}N}^{-1}(\lambda_k) \right\| \le C g_N(k)^{-1} \tag{A.8}$$

which implies that the first factor on the right of $(A.7)$ is bounded by

$$C \left| \frac{1}{N} \sum_{\lambda_k}^{\lambda_0} \left\| \widetilde{\varphi}_{\widehat{\theta}k} \right\| g_N(k)^{\frac{\delta}{2}-1} \right| = O_p\left(\prod_{\ell=1}^{d} |\lambda_0[\ell] + \mathring{\pi}[\ell]|^{\frac{\delta}{2}} \right).$$

Next, by Lemma 3, the second term inside the braces on the right of (A.7) is $O_p(1)$ for $\delta > 0$ small enough, whereas Lemmas 3 and 1 imply that the first term on the right of (A.7) is bounded by

$$\begin{aligned} &\sup_{-\tilde{n} < k \le [\lambda_0 \tilde{n}]} \left\| \frac{g_N(k)^{-\frac{\delta}{2}}}{N} \sum_{\lambda_s}^{\lambda_k} \widetilde{\varphi}_{\widehat{\theta}s} \widetilde{\varphi}'_{\widehat{\theta}s} \right\| O_{p^*}(1) + o_{p^*}\left(\sup_{-\tilde{n} < k \le [\lambda_0 \tilde{n}]} \frac{g_N(k)^{-\frac{\delta}{2}}}{N^{\delta}} \right) \\ &\quad = O_{p^*}\left(\prod_{\ell=1}^{d} |\lambda_0[\ell] + \mathring{\pi}[\ell]|^{\frac{\delta}{2}} \right) \end{aligned}$$

because $n^{-1}[\ell] \le \tilde{n}^{-1}[\ell] \le \inf_{-\tilde{n} < k \le [\lambda_0 \tilde{n}]} (k[\ell]/\tilde{n}[\ell])$, $0 < \delta < 1$ and an obvious extension of Brillinger (1981, p. 15). So we conclude that $(A.7) = O_{p^*}\left(\prod_{\ell=1}^{d} |\lambda_0[\ell] + \mathring{\pi}[\ell]|^{\frac{\delta}{2}} \right)$ and hence $(A.6)$ holds true because $\delta > 0$. ∎

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

LEMMA 5. *Assuming* $C1-C8$,

$$\sup_{\lambda\in\widetilde{\Pi}^d}\left\|\sum_{\lambda_s}^{\lambda}\left(\varphi_{\widehat{\theta}^*N}(\lambda_s)-\varphi_{\widehat{\theta}N}(\lambda_s)\right)\left(\varkappa_s^*+\jmath_s^*\right)\right\|=O_{p^*}(1). \tag{A.9}$$

Proof. The expression inside the norm on the left of $(A.9)$ is

$$\sum_{\lambda_s}^{\lambda}\frac{\partial}{\partial\theta}\varphi_{\widehat{\theta}N}(\lambda_s)\varkappa_s^*\left(\widehat{\theta}^*-\widehat{\theta}\right)+\sum_{\lambda_s}^{\lambda}\frac{\partial}{\partial\theta}\varphi_{\widehat{\theta}N}(\lambda_s)\jmath_s^*\left(\widehat{\theta}^*-\widehat{\theta}\right)$$
$$+\sum_{\lambda_s}^{\lambda}\left(\varphi_{\widehat{\theta}^*N}(\lambda_s)-\varphi_{\widehat{\theta}N}(\lambda_s)-\frac{\partial}{\partial\theta}\varphi_{\widehat{\theta}N}(\lambda_s)\left(\widehat{\theta}^*-\widehat{\theta}\right)\right)\left(\varkappa_s^*+\jmath_s^*\right). \tag{A.10}$$

By $C9$ and then noting that $|a-b|\le(a-b)+2b$ for $a>0$ and $b>0$, the norm of the third term of $(A.10)$ is bounded by

$$C\left\|\widehat{\theta}^*-\widehat{\theta}\right\|^2\sum_{\lambda_s}\left|\varkappa_s^*+\jmath_s^*\right|=O_p\left(\left\|\widehat{\theta}^*-\widehat{\theta}\right\|^2\right)\left\{\sum_{\lambda_s}\left(\varkappa_s^*+\jmath_s^*\right)+\frac{\widehat{\sigma}_\varepsilon^2}{\pi}\sum_{\lambda_s}1\right\}$$
$$=O_{p^*}(1)$$

by Lemma 1 and then using Lemmas 3 and 18 of Hidalgo (2009). So, uniformly in λ, the third term of (A.10) is $o_{p^*}(1)$. Likewise, the first term of (A.10) is $O_{p^*}(1)$ uniformly in λ using Lemma 4 with $\widehat{\zeta}(\lambda)=\frac{\partial}{\partial\theta}\varphi_{\widehat{\theta}}(\lambda)$ and Lemma 1. Finally, the second term of (A.10) is $O_{p^*}(1)$ by Lemma 18 of Hidalgo (2009) with $\widehat{\zeta}(\lambda)=\frac{\partial}{\partial\theta}\varphi_{\widehat{\theta}}(\lambda)$. ■

LEMMA 6. *Assuming* $C1-C9$, *for all* $\varepsilon>0$, *in probability*

$$\lim_{\lambda_0\to-\mathring{\pi}}\overline{\lim_{n\to\infty}}\Pr^*\left\{\sup_{-\mathring{\pi}\prec\lambda\preceq\lambda_0}\left|\frac{1}{N}\sum_{\lambda\le\lambda_k\le\lambda_0}\frac{\Im_{\widehat{\theta}^*N}(k)}{N^{1/2}}\sum_{s\prec k}\widetilde{\varphi}_{\widehat{\theta}^*N}(\lambda_s)\left(\varkappa_s^*+\jmath_s^*\right)\right|>\varepsilon\right\}=0. \tag{A.11}$$

Proof. Notice that Lemma 1 implies that it suffices to show $(A.11)$ in the set $\left\{\left\|\widehat{\theta}^*-\widehat{\theta}\right\|<CN^{-1/2}m_N^{-1}\right\}$, where $m_N+m_N^{-1}N^{-1/2}\to0$. On the other hand, Lemmas 2 and 3 imply that, uniformly in k,

$$\frac{1}{N^{1/2}}\sum_{s\prec k}\widetilde{\varphi}_{\widehat{\theta}^*N}(\lambda_s)\varkappa_s^*=-\left(\frac{\widehat{\sigma}_\varepsilon^2}{N}\sum_{s\prec k}\widetilde{\varphi}_{\widehat{\theta}^*N}(\lambda_s)\widetilde{\varphi}'_{\widehat{\theta}^*N}(\lambda_s)\right)N^{1/2}\left(\widehat{\theta}-\widehat{\theta}^*\right)+o_{p^*}(1)$$

$$\frac{1}{N^{1/2}}\sum_{s\prec k}\widetilde{\varphi}_{\widehat{\theta}^*N}(\lambda_s)\jmath_s^*=\frac{1}{N^{1/2}}\sum_{s\prec k}\widetilde{\varphi}_{\widehat{\theta}^*N}(\lambda_s)\jmath_s^*+O_{p^*}\left(n^{-1/2}\right) \tag{A.12}$$

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

proceeding as in the proof of $(A.9)$ but with $\varkappa_s^* + \jmath_s^*$ replaced by $\jmath_s^*$ there. Observe that we can take $\lambda_0 \prec -\mathring{\pi}/2$. Next, $C8$ implies that

$$\sup_{-\tilde{n} < k \leq [\lambda_0 \tilde{n}]} \left\| \widetilde{\Lambda}_{\widehat{\theta}^* N}(k) - \widetilde{\Lambda}_{\widehat{\theta} N}(k) \right\| = O_{p^*}\left(\left\| \widehat{\theta}^* - \widehat{\theta} \right\| \right) \prod_{\ell=1}^{d} |\lambda_0[\ell] + \mathring{\pi}[\ell]|$$

which, together with $(A.8)$, implies that $\left\| \widetilde{\Lambda}^{-1}_{\widehat{\theta}^* N}(k) \right\| = O_p\left(g_N(k)^{-1} \right)$.

So, we have that for $0 < \delta < 1/2$,

$$\begin{aligned}
&\sup_{-\mathring{\pi} \prec \lambda \preceq \lambda_0} \left\| \frac{1}{\widetilde{n}} \sum_{\lambda \leq \lambda_k \leq \lambda_0} \frac{\Im_{\widehat{\theta}^* n}(k)}{N^{1/2}} \sum_{s \prec k} \widetilde{\varphi}_{\widehat{\theta}^* N}(\lambda_s) \left(\varkappa_s^* + \jmath_s^* \right) \right\| \\
&= O_{p^*}(1) \sup_{-\mathring{\pi} \prec \lambda \preceq \lambda_0} \left| \frac{1}{N} \sum_{\lambda \leq \lambda_k \leq \lambda_0} \left\| \widetilde{\varphi}_{\widehat{\theta} N}(k) \right\| g_N(k)^{(-1+\delta/2)} \right| \\
&\quad \times \left\{ \sup_{\lambda_k \leq \lambda_0} \left\| g_N(k)^{-\delta/2} \frac{1}{N^{1/2}} \sum_{s \prec k} \widetilde{\varphi}_{\widehat{\theta} N}(\lambda_s) \jmath_s^* \right\| + O_{p^*}\left(\prod_{\ell=1}^{d} |\lambda_0[\ell] + \mathring{\pi}[\ell]|^{\frac{\delta}{2}} \right) \right\}, \quad \textbf{(A.13)}
\end{aligned}$$

by $(A.12)$ and because $C2$ implies that $\overrightarrow{n} \leq \inf_{\lambda_k \leq \lambda_0} (k[\ell]/\widetilde{n}[\ell])$. But Lemma 4 implies that $\sup_{\lambda_k \leq \lambda_0} \left\| g_N(k)^{-\delta/2} N^{-1/2} \sum_{s \prec k} \widetilde{\varphi}_{\widehat{\theta} N}(\lambda_s) \jmath_s^* \right\| = O_{p^*}(1)$ and $C3$ implies that

$$\sup_{-\mathring{\pi} \prec \lambda \preceq \lambda_0} \frac{1}{N} \sum_{\lambda \leq \lambda_k \leq \lambda_0} \left\| \widetilde{\varphi}_{\widehat{\theta} N}(k) \right\| g_N(k)^{(-1+\delta/2)} = O_p\left(\prod_{\ell=1}^{d} |\lambda[\ell] + \mathring{\pi}[\ell]|^{\frac{\delta}{2}} \right),$$

so it is the left side of $(A.13)$. From here, we conclude because $\delta > 0$. ■

APPENDIX B

B.1. Proof of Theorem 4

We begin with part (a). Using $G_N^*(\pi) = \widehat{\sigma}_\varepsilon^2 + o_{p^*}(1)$ and recalling that $\Im_{\theta N}(\lambda_s) = \widetilde{\varphi}'_{\theta N}(\lambda_s) \widetilde{\Lambda}^{-1}_{\theta N}(\lambda_s)$, we obtain that, uniform in $\lambda \in \widetilde{\Pi}^d$,

$$\widehat{\alpha}^*_{\widehat{\theta}^* N}(\lambda) = \frac{(2\pi)^{2d}}{\widehat{\sigma}_\varepsilon^2} \frac{1}{N^{1/2}} \sum_{\lambda_s}^{\lambda} \jmath_s^* - \frac{(2\pi)^{2d}}{\widehat{\sigma}_\varepsilon^2} \widehat{\Upsilon}_N(\lambda) + o_{p^*}(1), \quad \textbf{(B.1)}$$

where $\widehat{\Upsilon}_N(\lambda) = N^{-3/2} \widehat{\sigma}_\varepsilon^{-2} \sum_{\lambda_s}^{\lambda} \left(\Im_{\widehat{\theta} N}(\lambda_s) \sum_{p^* \prec s}^{[\lambda]} \widetilde{\varphi}_{\widehat{\theta} N}(k) \jmath_k^* \right)$ and $\jmath_k^*$ as defined in (A.5).

Suppose, to be shown later, that the convergence in $\lambda \preceq \lambda_0$ holds true for any $\lambda_0 \in \widetilde{\Pi}^d$. Then, because the Brownian sheet $\boldsymbol{B}(\lambda/\pi)$ and the limit of $N^{-1/2} \sum_{\lambda_s}^{\lambda} \jmath_s^*$ are continuous in $\widetilde{\Pi}^d$, Billingsley's (1968) Theorem 4.2 implies that it suffices to show that for all $\varepsilon > 0$,

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

$$\lim_{\lambda_0 \to -\mathring{\pi}} \overline{\lim_{\vec{n} \to \infty}} \Pr{}^* \left\{ \sup_{-\mathring{\pi} \prec \lambda \preceq \lambda_0} \left| \widehat{\Upsilon}_N(\lambda_0) - \widehat{\Upsilon}_N(\lambda) \right| > \varepsilon \right\} = 0,$$

in probability. But this holds true by Lemma 5, cf. the second term on the right of (A.7).

So, to complete the proof we need to show that, for any $\lambda_0 \in \widetilde{\Pi}^d$, the first two terms on the right of (B.1) converge in bootstrap to $\pi^{-d/2} \boldsymbol{B}(\lambda/\pi)$ in $-\mathring{\pi} \prec \lambda \preceq \lambda_0$ in probability. Fidi's convergence follows by Lemma 18 Hidalgo (2009) part (b) after we write $\widehat{\Upsilon}_N(\lambda)$ as

$$\frac{(2\pi)^{2d}}{\widehat{\sigma}_\varepsilon^2} \frac{1}{N^{1/2}} \sum_{\lambda_k} \left(\frac{1}{N} \sum_{\lambda_s}^{\lambda_k \wedge \lambda} \Im_{\widehat{\theta}N}(s) \right) \widetilde{\varphi}_{\widehat{\theta}N}(k) \jmath_k^*$$

and $\left(N^{-1} \sum_{\lambda_s}^{\lambda_k \wedge \lambda} \Im_{\widehat{\theta}N}(s) \right) \widetilde{\varphi}_{\widehat{\theta}N}(k)$ satisfies the same conditions of Lemma 18 Hidalgo (2009) for $\widehat{\zeta}(\lambda)$. Then, it suffices to prove tightness. Since $N^{-1/2} \sum_{\lambda_s}^{\lambda} \jmath_s^*$ is tight by Lemma 2, we only need to show the tightness condition of $\widehat{\Upsilon}_N(\lambda)$. By Billingsley's (1968) Theorem 15.6, it suffices to show that

$$\mathbb{E}^* \left(\left| \widehat{\Upsilon}_N(\vartheta) - \widehat{\Upsilon}_N(\mu) \right| \left| \widehat{\Upsilon}_N(\lambda) - \widehat{\Upsilon}_N(\vartheta) \right| \right) = O_p(1) \prod_{\ell=1}^{d} |\lambda[\ell] - \mu[\ell]|^{2\delta}$$

for all $-\mathring{\pi}[\ell] \leq \mu[\ell] < \vartheta[\ell] < \lambda[\ell] \leq \pi$ and some $\delta > 1/2$. Observe that we can take $\widetilde{n}^{-1} < |\lambda[\ell] - \mu[\ell]|$ since otherwise the last inequality is trivial. Because $(\lambda - \vartheta)(\vartheta - \mu) < (\lambda - \mu)^2$ by the Cauchy–Schwarz's inequality, it suffices to show the last displayed equality holds for $\mathbb{E}^* \left| \widehat{\Upsilon}_N(\lambda) - \widehat{\Upsilon}_N(\mu) \right|^2$ which is

$$\begin{aligned}
&\frac{1}{\widehat{\sigma}_\varepsilon^4} \frac{1}{N^3} \sum_{\mu \leq \lambda_s, \lambda_k \leq \lambda} \Im_{\widehat{\theta}N}(s) \sum_{\ell_1 \prec s} \sum_{\ell_2 \prec k} \widetilde{\varphi}_{\widehat{\theta}N}(\ell_1) \widetilde{\varphi}'_{\widehat{\theta}N}(\ell_2) \mathbb{E}^* \left(\jmath_{\ell_1}^* \jmath_{\ell_2}^* \right) \Im'_{\widehat{\theta}N}(k) \\
&\quad = H_N(\lambda, \mu) \frac{1}{N^2} \sum_{\mu \leq \lambda_s, \lambda_k \leq \lambda} \left\| \Im_{\widehat{\theta}N}(s) \right\| \left\| \Im_{\widehat{\theta}N}(k) \right\| \\
&\quad = H_N(\lambda, \mu) \left(\left| \widetilde{\Im}_{\widehat{\theta}}(\mu, \lambda) \right|^2 + N^{-2} \right),
\end{aligned}$$

because $\left\| N^{-1} \sum_{\lambda_s}^{\lambda} \left\| \Im_{\widehat{\theta}N}(s) \right\| - \widetilde{\Im}_{\widehat{\theta}}(\lambda) \right\| = O_p\left(N^{-1} \right)$ by Lemma 12 of Hidalgo (2009) with $\widetilde{\Im}_{\widehat{\theta}}(\mu, \lambda) = \pi^{-1} \int_{-\mu}^{\lambda} \Im_{\widehat{\theta}}(w) dw$. From here we conclude the proof of part (a) by Billingsley's (1968) Theorem 15.6, because $\widetilde{\Im}_\theta(\lambda)$ is a monotonic, continuous, and non-decreasing function such that $\left| \widetilde{\Im}_{\widehat{\theta}}(\lambda) - \widetilde{\Im}_{\widehat{\theta}}(\mu) \right| = O_p(1) \prod_{\ell=1}^{d} |\lambda[\ell] - \mu[\ell]|^{\delta}$, $\delta > 1/2$ and $\widetilde{n}^{-1}[\ell] \leq |\lambda[\ell] - \mu[\ell]|$. To show part (b), by definition of $\beta_{\theta N}^*$ and β_N^*, it suffices to show that

$$\left| \frac{1}{N^{1/2}} \sum_{\lambda_k}^{\lambda} \varkappa_k^* - \Im_{\widehat{\theta}N}(k) \frac{1}{N} \sum_{\lambda_s}^{\lambda_k} \widetilde{\varphi}_{\widehat{\theta}N}(s) \varkappa_s^* \right| \tag{B.2}$$

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

$$\frac{1}{G_{\widehat{\theta}^* N}(\pi)}\left(\frac{1}{N}\sum_{\lambda_k}^{\lambda}\Im_{\widehat{\theta}N}(k)\frac{1}{N^{1/2}}\sum_{s\prec k}\widetilde{\varphi}_{\widehat{\theta}N}(s)\left(\frac{I_{x^*s}}{\left|\Psi_{\widehat{\theta}^* s}\right|^2}-\frac{G_{\widehat{\theta}^* N}(\pi)}{2\pi}\right)\right)$$

$$-\frac{1}{G_{\widehat{\theta}^* N}(\pi)}\left(\frac{1}{N}\sum_{\lambda_k}^{\lambda}\Im_{\widehat{\theta}^* N}(k)\frac{1}{N^{1/2}}\sum_{s\prec k}\widetilde{\varphi}_{\widehat{\theta}^* N}(s)\left(\frac{I_{x^*s}}{\left|\Psi_{\widehat{\theta}^* s}\right|^2}-\frac{G_{\widehat{\theta}^* N}(\pi)}{2\pi}\right)\right) \tag{B.3}$$

converge to zero uniformly in $\lambda\in\widetilde{\Pi}^d$. Expression (B.2) is $o_{p^*}(1)$, uniformly in $\lambda\in\widetilde{\Pi}^d$, because as we argued with (57) in Delgado et al. (2011)

$$-\frac{\widetilde{\varphi}'_{\widehat{\theta}^* N}(s)}{G_{\widehat{\theta}^* N}(\pi)}\Im^{-1}_{\widehat{\theta}^* N}\frac{1}{N^{1/2}}\sum_{s\prec k}\widetilde{\varphi}_{\widehat{\theta}^* N}(k)\,I_{\varepsilon^* k}=0.$$

Next, because

$$\frac{1}{N}\sum_{\lambda_k}^{\lambda}\left\|\widetilde{\varphi}_{\widehat{\theta}N}(k)\right\|\left\|\widetilde{\Lambda}^{-1}_{\widehat{\theta}N}(k)\right\|\frac{1}{N}\sum_{s\prec k}\left\|\widetilde{\varphi}_{\widehat{\theta}N}(s)\right\|$$

$$\leq C\frac{1}{N}\sum_{\lambda_k}^{\lambda}\left\|\widetilde{\varphi}_{\widehat{\theta}N}(k)\right\|\left\|\widetilde{\Lambda}^{-1}_{\widehat{\theta}N}(k)\,g_N(k)\right\|\leq C\frac{1}{N}\sum_{\lambda_k}^{\lambda}\left\|\widetilde{\varphi}_{\widehat{\theta}N}(k)\right\|=O_p(1)$$

by integrability of $\varphi_{\widehat{\theta}^*}(\lambda)$ and (A.8), it implies that the contribution into (B.2) due to the term $o_{p^*}(1)$ on part (a) of Theorem 1 is negligible.

Next we examine (B.3). Because $G_{\widehat{\theta}^* N}(\pi)-G_{\widehat{\theta}N}(\pi)=o_{p^*}\left(N^{-1/2}\right)$ by Lemma 3 and $G_{\widehat{\theta}N}(\pi)-G_{\theta_0 N}(\pi)=o_p\left(N^{-1/2}\right)$ by Lemma 15 of Hidalgo (2009), it suffices to show that

$$\frac{1}{N}\sum_{\lambda_k}^{\lambda}\left\{\frac{\Im_{\widehat{\theta}N}(k)}{N^{1/2}}\sum_{\lambda_k\prec\lambda_s}\widetilde{\varphi}_{\widehat{\theta}N}(s)\left(\varkappa^*_s+\jmath^*_s\right)-\frac{\Im_{\widehat{\theta}^* N}(k)}{N^{1/2}}\sum_{s\prec k}\widetilde{\varphi}_{\widehat{\theta}^* N}(s)\left(\varkappa^*_s+\jmath^*_s\right)\right\} \tag{B.4}$$

converges to zero uniformly in $\lambda\in\widetilde{\Pi}^d$ after observing that

$$\sup_{\lambda\in\widetilde{\Pi}^d}\left|\sum_{\lambda_k}^{\lambda}\Im_{\widehat{\theta}^* N}(k)\sum_{s\prec k}\widetilde{\varphi}_{\widehat{\theta}^* N}(s)-\sum_{\lambda_k}^{\lambda}\Im_{\widehat{\theta}N}(k)\sum_{s\prec k}\gamma_{\widehat{\theta}s}\right|=0.$$

First, we observe that Lemmas 3 and 5 imply that it suffices to show the uniform convergence in $-\mathring{\pi}\prec\lambda\preceq\lambda_0$ for any $\lambda_0\prec 0$. But (B.4) is

$$\frac{1}{N}\sum_{\lambda_k}^{\lambda}\Im_{\widehat{\theta}^* N}(k)\frac{1}{N^{1/2}}\sum_{s\prec k}\left(\widetilde{\varphi}_{\widehat{\theta}N}(s)-\widetilde{\varphi}_{\widehat{\theta}^* N}(s)\right)\left(\varkappa^*_s+\jmath^*_s\right) \tag{B.5}$$

$$+\frac{1}{N}\sum_{\lambda_k}^{\lambda}\left(\Im_{\widehat{\theta}N}(k)-\Im_{\widehat{\theta}^* N}(k)\right)\frac{1}{N^{1/2}}\sum_{s\prec k}\widetilde{\varphi}_{\widehat{\theta}N}(s)\left(\varkappa^*_s+\jmath^*_s\right). \tag{B.6}$$

So, the theorem follows if (B.5) and (B.6) are both $o_{p^*}(1)$ uniformly in $-\mathring{\pi}\prec\lambda\preceq\lambda_0$. To that end, we first show that

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

$$\sup_{\lambda\in\widetilde{\Pi}^d}\frac{1}{N}\sum_{\lambda_s}^{\lambda}\left\|\widetilde{\varphi}_{\widehat{\theta}N}(s)-\widetilde{\varphi}_{\widehat{\theta}^*N}(s)\right\|=o_{p^*}(1), \tag{B.7}$$

$$\sup_{-\mathring{\pi}\prec\lambda\preceq\lambda_0}\left\|\widetilde{\Lambda}^{-1}_{\widehat{\theta}N}(\lambda)-\widetilde{\Lambda}^{-1}_{\widehat{\theta}}(\lambda)\right\|=o_p(1), \tag{B.8}$$

$$\sup_{-\mathring{\pi}\prec\lambda\preceq\lambda_0}\left\|\widetilde{\Lambda}^{-1}_{\widehat{\theta}^*N}(\lambda)-\widetilde{\Lambda}^{-1}_{\widehat{\theta}N}(\lambda)\right\|=o_{p^*}(1). \tag{B.9}$$

First, (B.7) follows proceeding as with the proof of (A.9) in Lemma 5 but without the factor $\varkappa_s^*+\jmath_s^*$, (B.8) follows because $C8$ implies that $\widetilde{\Lambda}_{\theta_0}(\lambda_0)>0$ and because by $C3$ $\left\|\widetilde{\varphi}_{\widehat{\theta}}(\lambda)\widetilde{\varphi}'_{\widehat{\theta}}(\lambda)\right\|$ satisfies the same conditions of $\zeta(\lambda)$ in Lemma 12 of Hidalgo (2009) , so that

$$\sup_{-\mathring{\pi}\prec\lambda\preceq\lambda_0}\left\|\widetilde{\Lambda}_{\widehat{\theta}}(\lambda)-\widetilde{\Lambda}_{\widehat{\theta}N}(\lambda)\right\|=O\left(n^{-1}\right),$$

whereas (B.9) follows proceeding as with the proof of (B.7) and (B.8).

Now we show that (B.5) is $o_p(1)$ uniformly in $-\mathring{\pi}\prec\lambda\preceq\lambda_0$, which follows by Lemma 5 and (B.7)–(B.9) noting that $\left(\widetilde{\varphi}'_{\widehat{\theta}N}(s)-\widetilde{\varphi}'_{\widehat{\theta}^*N}(s)\right)=\left(\phi'_{\widehat{\theta}N}(s)-\phi'_{\widehat{\theta}^*N}(s),0\right)$, so does (B.6) by (B.7) and (B.9) and that

$$\sup_{-\mathring{\pi}\prec\lambda\preceq\lambda_0}\left|\frac{1}{N^{1/2}}\sum_{\lambda_s}^{\lambda}\widetilde{\varphi}_{\widehat{\theta}s}\left(\varkappa_s^*+\jmath_s^*\right)\right|=O_{p^*}(1)$$

by Lemmas 1 and 2 with $\widehat{\zeta}(\lambda)=\widetilde{\varphi}_{\widehat{\theta}}(\lambda)$ there and observing Lemma 1 and that Lemma 12 of Hidalgo (2009) implies that $N^{-1}\sum_{\lambda_s}^{\lambda}\widetilde{\varphi}_{\widehat{\theta}N}(s)\widetilde{\varphi}'_{\widehat{\theta}N}(s)\to_P\int_{-\pi}^{\lambda}\widetilde{\varphi}_{\theta_0}(\omega)\widetilde{\varphi}'_{\theta_0}(\omega)d\omega$. ■

B.2. Proof of Proposition 1 and Corollary 3

B.2.1. Proof of Proposition 1. First we notice that it suffices to show that

$$A_n^{-1}\sum_{t=1}^{n}z(t)x(t)\xrightarrow{d}\mathcal{N}\left(0,\int_{-\pi}^{\pi}f(\lambda)\mathcal{M}(d\lambda)\right). \tag{B.10}$$

To that end, we shall show first that

$$A_n^{-1}E\left(\sum_{t=1}^{n}z(t)x(t)\sum_{t=1}^{n}z'(t)x(t)\right)A_n^{-1}\to\int_{-\pi}^{\pi}f(\lambda)\mathcal{M}(d\lambda).$$

For that purpose, we first notice that by Weierstrass approximation theorem, we have that we can find two trigonometric polynomials $f_x^{(1)}(\lambda)$ and $f_x^{(2)}(\lambda)$ such that $f_x^{(2)}(\lambda)-f_x^{(1)}(\lambda)\le\epsilon$ and $f_x^{(1)}(\lambda)\le f(\lambda)\le f_x^{(2)}(\lambda)$. When the spectral density function is not continuous, we can employ the construction given in Hannan (1970, pp. 216–218). Observe that the latter implies that

$$f_x^{(1)}(\lambda)\mathcal{Z}(\lambda)\mathcal{Z}^*(\lambda)\le f(\lambda)\mathcal{Z}(\lambda)\mathcal{Z}^*(\lambda)\le f_x^{(2)}(\lambda)\mathcal{Z}(\lambda)\mathcal{Z}^*(\lambda),$$

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

where $\mathcal{Z}(\lambda) = A_n^{-1} \sum_{t=1}^n z(t) e^{it'\lambda}$ and

$$\int_{-\pi}^{\pi} \left(f_x^{(2)}(\lambda) - f_x^{(1)}(\lambda) \right) \mathcal{Z}(\lambda) \mathcal{Z}^*(\lambda) d\lambda \leq \epsilon \int_{-\pi}^{\pi} \mathcal{Z}(\lambda) \mathcal{Z}^*(\lambda) d\lambda$$
$$= \epsilon A_n^{-1} \sum_{t=1}^n z(t) z'(t) A_n^{-1}.$$
$$\to \epsilon \mathcal{R}. \tag{B.11}$$

So, it suffices to show (B.10) with $x(t)$ being replaced by $\ddot{x}(t)$, where

$$\ddot{x}(t) = \sum_{s \in \mathbb{M}^d} \psi(s) \varepsilon(t-s)$$

and $\mathbb{M}^d = \{s : |s[\ell]| < J, \quad \ell = 1, \ldots, d\}$. This is a moving average of finite order. Now, by standard algebra,

$$\alpha' A_n^{-1} \sum_{t=1}^n z(t) \ddot{x}(t) = \sum_{s \in \mathbb{M}^d} \psi(s) \sum_{t=1}^n \left(\sum_{r=1}^q \alpha[r] \frac{z_r(t)}{A_n[r]} \right) \varepsilon(t-s),$$

where α is a q-dimensional vector with norm 1. Now, for each $s \in \mathbb{M}^d$, the term on right side of the last displayed equation, $\sum_{t=1}^n \left(\sum_{r=1}^q \alpha[r] \frac{z_r(t)}{A_n[r]} \right) \varepsilon(t-s)$, converges in distribution to a normal random variable if the Lindeberg's condition is satisfied. However, this is the case as $\varepsilon(t)$ is iid and thus $\varepsilon^2(t)$ is uniformly integrable, and for any $\delta > 0$

$$\sum_{t=1}^n E \left(\frac{z_s(t)^2}{A_n[s]^2} \right) \varepsilon^2(t-s) \mathbb{I} \left\{ \left(\frac{z_s(t)^2}{A_n[r]^2} \right) \varepsilon^2(t-s) > \delta \right\}$$
$$\leq \mathrm{E}\varepsilon^2(t) \mathbb{I} \left\{ \varepsilon^2(t) > \min_u \frac{A_n[r]^2}{z_{u,r}^2} \delta \right\} \to 0$$

since $\left(\sum_{t=1}^n z_r(t)^2 A_n[r]^{-2} \right) = 1$ and $\max_u z_r^2(u) \left(\sum_{t=1}^n z_r(t)^2 \right)^{-1} \to 0$ for all $r = 1, \ldots, q$, and where we have employed $\mathbb{I}(\cdot)$ for the indicator function.

B.2.2. Proof of Corollary 3. We now show that

$$\widehat{\vartheta} - \widetilde{\vartheta} = O_p\left(N^{-1}\right).$$

To that end, it suffices to check Robinson (1988), that is

$$\frac{1}{N} \sum_{\lambda_s} \phi_{\widehat{\vartheta},s} \left\{ \frac{I_{\widehat{x},s}}{(2\pi)^d f_{\widehat{\vartheta},s}} - 1 \right\} = O_p\left(N^{-1}\right). \tag{B.12}$$

The left side of (B.12), except the multiplicative constant $(2\pi)^{-d}$, is

$$\frac{1}{N} \sum_{\lambda_s} \phi_{\widehat{\vartheta},s} \frac{I_{\widehat{x},s} - I_{x,s}}{f_{\widehat{\vartheta},s}} = \left(\widehat{\beta} - \beta\right)' \frac{1}{N} \sum_{\lambda_s} \phi_{\widehat{\vartheta},s} \frac{I_{z,s}}{f_{\widehat{\vartheta},s}} \left(\widehat{\beta} - \beta\right)$$
$$- 2\left(\widehat{\beta} - \beta\right)' \frac{1}{N} \sum_{\lambda_s} \phi_{\widehat{\vartheta},s} \frac{\mathrm{Re}\left(w_{z,s} \overline{w_{x,s}}\right)}{f_{\widehat{\vartheta},s}}. \tag{B.13}$$

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

First by standard linearization and that $\widehat{\vartheta}-\vartheta_0=O_p\left(N^{-1/2}\right)$, we have that

$$\frac{1}{N}\sum_{\lambda_s}\phi_{\widehat{\vartheta},s}\frac{I_{z,s}}{f_{\widehat{\vartheta},s}}=\frac{1}{N}\sum_{\lambda_s}\phi_{\vartheta_0,s}\frac{I_{z,s}}{f_{\vartheta_0,s}}\left(1+O_p\left(N^{-1/2}\right)\right).$$

Next Proposition 1 implies that the behavior of

$$\left(\widehat{\beta}-\beta\right)'\frac{1}{N}\sum_{\lambda_s}\phi_{\vartheta_0,s}\frac{I_{z,s}}{f_{\vartheta_0,s}}\left(\widehat{\beta}-\beta\right),$$

where $\zeta_\vartheta(\lambda)=f_\vartheta^{-1}(\lambda)\phi_\vartheta(\lambda)$, is that of

$$A_n^{-1}\frac{1}{N}\sum_{\lambda_s}\zeta_{\vartheta_0,s}I_{z,s}A_n^{-1}=\frac{1}{N^2}\sum_{\lambda_s}\sum_{r=-n+1}^{n-1}e^{ir'\lambda_s}\zeta_{\vartheta_0,s}A_n^{-1}\sum_{t=1}^{n-r}z(t)z'(t+r)A_n^{-1}$$

by standard algebra. But by Grenander conditions, the right side of the last displayed expression is

$$\frac{1}{N}\int_{-\mathring{\pi}}^{\pi}\zeta_{\vartheta_0}(\lambda)\mathcal{M}(d\lambda).$$

So, the first term of the right of (B.13) is $O_p\left(N^{-1}\right)$. Next as we have done with the first term on the right of (B.13), the second term is governed by the behavior of

$$\begin{aligned}A_n^{-1}\frac{1}{N}\sum_{\lambda_s}\zeta_{\widehat{\vartheta},s}w_{z,s}\overline{w_{x,s}}&=A_n^{-1}\frac{1}{N}\sum_{\lambda_s}\zeta_{\vartheta_0,s}w_{z,s}\overline{w_{x,s}}\\&\quad+O_p\left(N^{-1/2}\right)A_n^{-1}\frac{1}{N}\sum_{\lambda_s}\left(\frac{\partial}{\partial\vartheta}\zeta_{\vartheta_0,s}\right)w_{z,s}\overline{w_{x,s}}\\&\quad+O_p\left(N^{-1}\right)A_n^{-1}\frac{1}{N}\sum_{\lambda_s}\left\|\left(\frac{\partial^2}{\partial\vartheta^2}\zeta_{\breve{\vartheta},s}\right)w_{z,s}\overline{w_{x,s}}\right\|,\end{aligned}$$

where $\breve{\vartheta}$ is an intermediate point between $\widehat{\vartheta}$ and ϑ_0. From here it is standard to conclude that

$$A_n^{-1}\frac{1}{N}\sum_{\lambda_s}\zeta_{\widehat{\vartheta},s}w_{z,s}\overline{w_{x,s}}=O_p\left(N^{-1}\right)$$

because

$$\mathbb{E}\left|A_n^{-1}\frac{1}{N}\sum_{\lambda_s}\left\{\zeta_{\vartheta_0,s}+\frac{\partial}{\partial\vartheta}\zeta_{\vartheta_0,s}\right\}w_{z,s}\overline{w_{x,s}}\right|^2=O_p\left(N^{-2}\right)\tag{B.14}$$

as we now prove. First, by Lemma 3 of Hidalgo (2009),

$$\mathbb{E}\left|A_n^{-1}\frac{1}{N}\sum_{\lambda_s}\zeta_{\vartheta_0,s}w_{z,s}\overline{w_{x,s}}\right|^2=\mathbb{E}\left|A_n^{-1}\frac{1}{N}\sum_{\lambda_s}\frac{\phi_{\vartheta_0,s}}{f_{\vartheta_0,s}^{1/2}}w_{z,s}\overline{w_{\varepsilon,s}}\right|^2(1+o(1)).$$

Now, from here it is obvious that (B.14) holds true as $\mathbb{E}\left(w_{\varepsilon,s}w_{\varepsilon,k}\right)=\mathbb{I}(s=k)$. So, we have that second term of the right of (B.13) is also $O_p\left(N^{-1}\right)$ and hence (B.12) is shown.

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

B.3. Proof of Theorem 5

We will only show part (a) as part (b) is handled similarly. The proof proceeds very similarly to Corollary 3. Indeed, except multiplicative constants,

$$\sup_{\lambda\in\bar{\Pi}^d}\left|\widehat{\alpha}_{\widetilde{\theta}N}(\lambda)-\alpha_{\widehat{\theta}N}(\lambda)\right|\widehat{G}_{\theta N}(\lambda)=\frac{(2\pi)^d}{N}\sum_{\lambda_s}^{\lambda}\frac{I_{\widehat{x}}^T(\lambda_s)}{|\Psi_\theta(\lambda_s)|^2},$$

where

$$\widehat{\alpha}_{\widetilde{\theta}N}(\lambda)-\alpha_{\widehat{\theta}N}(\lambda)=N^{1/2}\left(\frac{\widehat{G}_{\widetilde{\theta}N}(\lambda)}{\widehat{G}_{\widetilde{\theta}N}(\pi)}-\frac{G_{\widehat{\theta}N}(\lambda)}{G_{\widehat{\theta}N}(\pi)}\right).$$

Now, by standard delta methods, it suffices to show that

$$\sup_{\lambda\in\bar{\Pi}^d}\left|N^{1/2}\left(\widehat{G}_{\widetilde{\theta}N}(\lambda)-G_{\widehat{\theta}N}(\lambda)\right)\right|=o_p(1).$$

But,

$$\begin{aligned}N^{1/2}\left(\widehat{G}_{\widetilde{\theta}N}(\lambda)-G_{\widehat{\theta}N}(\lambda)\right)&=\frac{1}{N^{1/2}}\sum_{\lambda_s}^{\lambda}\left\{\frac{I_{\widehat{x},s}}{\left|\Psi_{\widetilde{\vartheta},s}\right|^2}-\frac{I_{x,s}}{\left|\Psi_{\widehat{\vartheta},s}\right|^2}\right\}\\&=\frac{1}{N^{1/2}}\sum_{\lambda_s}^{\lambda}\frac{I_{\widehat{x},s}-I_{x,s}}{\left|\Psi_{\widehat{\vartheta},s}\right|^2}\\&\quad+\frac{1}{N^{1/2}}\sum_{\lambda_s}^{\lambda}I_{\widehat{x},s}\left\{\frac{1}{\left|\Psi_{\widetilde{\vartheta},s}\right|^2}-\frac{1}{\left|\Psi_{\widehat{\vartheta},s}\right|^2}\right\}.\end{aligned}\tag{B.15}$$

First, it is straightforward to show that

$$\sup_{\lambda\in\bar{\Pi}^d}\left|\frac{1}{N^{1/2}}\sum_{\lambda_s}^{\lambda}I_{\widehat{x},s}\left\{\frac{1}{\left|\Psi_{\widetilde{\vartheta},s}\right|^2}-\frac{1}{\left|\Psi_{\widehat{\vartheta},s}\right|^2}\right\}\right|=o_p(1)$$

because

$$\begin{aligned}\sup_{\lambda\in\bar{\Pi}^d}\left|\left|\Psi_{\widetilde{\vartheta},s}\right|^2-\left|\Psi_{\widetilde{\vartheta},s}\right|^2\right|&=\left(\widetilde{\vartheta}-\widetilde{\vartheta}\right)\sup_{\lambda\in\bar{\Pi}^d}\left|\frac{\partial}{\partial\vartheta}\left|\Psi_{\check{\vartheta}}(\lambda)\right|^2\right|\\&=O_p\left(\left|\widetilde{\vartheta}-\widetilde{\vartheta}\right|\right).\end{aligned}$$

In addition

$$\begin{aligned}\sup_{\lambda\in\bar{\Pi}^d}\left|\frac{1}{N}\sum_{\lambda_s}^{\lambda}I_{\widehat{x},s}\right|&=\frac{1}{N}\sum_{\lambda_s}I_{\widehat{x},s}\\&=\frac{1}{N}\sum_{\lambda_s}\left(I_{\widehat{x},s}-I_{x,s}\right)+\frac{1}{N}\sum_{\lambda_s}I_{x,s}\\&=O_p(1)\end{aligned}$$

as a consequence of Corollary 3.

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press

To conclude the proof it remains to show that the first in the far right of (B.15) satisfies that

$$\sup_{\lambda\in\widetilde{\Pi}^d}\left|\frac{1}{N^{1/2}}\sum_{\lambda_s}^{\lambda}\frac{I_{\widehat{x},s}-I_{x,s}}{\left|\Psi_{\widehat{\vartheta},s}\right|^2}\right|=o_p(1). \tag{B.16}$$

Now as in (B.13),

$$\frac{1}{N^{1/2}}\sum_{\lambda_s}^{\lambda}\frac{I_{\widehat{x},s}-I_{x,s}}{\left|\Psi_{\widehat{\vartheta},s}\right|^2}=\left(\widehat{\beta}-\beta\right)'\frac{1}{N^{1/2}}\sum_{\lambda_s}^{\lambda}\frac{I_{z,s}}{\left|\Psi_{\widehat{\vartheta},s}\right|^2}\left(\widehat{\beta}-\beta\right)$$
$$-2\left(\widehat{\beta}-\beta\right)'\frac{1}{N^{1/2}}\sum_{\lambda_s}^{\lambda}\frac{\mathrm{Re}\left(w_{z,s}\overline{w_{x,s}}\right)}{\left|\Psi_{\widehat{\vartheta},s}\right|^2}. \tag{B.17}$$

The contribution of the first term on the right of (B.17) into the left of (B.16) is bounded by

$$A_n^{-1}\frac{1}{N^{1/2}}\sum_{\lambda_s}\frac{I_{z,s}}{\left|\Psi_{\vartheta_0,s}\right|^2}A_n^{-1}=o_p(1)$$

as we showed in Corollary 3. Finally, the contribution of the second term on the right of (B.17) into the left of (B.16) is given by that of

$$\sup_{\lambda\in\widetilde{\Pi}^d}\left|A_n^{-1}\frac{1}{N^{1/2}}\sum_{\lambda_s}^{\lambda}\frac{w_{z,s}\overline{w_{\varepsilon,s}}}{\left|\Psi_{\vartheta_0,s}\right|}\right|$$

proceeding as in Lemma 3 of Hidalgo (2009). Now,

$$\mathbb{E}\left|A_n^{-1}\frac{1}{N^{1/2}}\sum_{\lambda_s}^{\lambda}\frac{w_{z,s}\overline{w_{\varepsilon,s}}}{\left|\Psi_{\vartheta_0,s}\right|}\right|^2=o(1)$$

because $\mathbb{E}\left(w_{\varepsilon,s}w_{\varepsilon,k}\right)=\mathbb{I}(s=k)$. On the other hand,

$$\mathbb{E}\left|A_n^{-1}\frac{1}{N^{1/2}}\sum_{\lambda_1\le\lambda_s\le\lambda_2}\frac{w_{z,s}\overline{w_{\varepsilon,s}}}{\left|\Psi_{\vartheta_0,s}\right|}\right|^2\le C\left|\lambda_2-\lambda_1\right|^{1+\delta}$$

proceeding as in Lemma 9 of Hidalgo (2009). This concludes the proof of part (a) and the theorem.

https://doi.org/10.1017/S0266466614000310 Published online by Cambridge University Press