The Journal of Symbolic Logic
Volume 80, Number 1, March 2015

STRUCTURE THEORY OF $L(\mathbb{R}, \mu)$ AND ITS APPLICATIONS

NAM TRANG

Abstract. In this paper, we explore the structure theory of $L(\mathbb{R}, \mu)$ under the hypothesis $L(\mathbb{R}, \mu) \vDash$ "$\mathsf{AD} + \mu$ is a normal fine measure on $\mathcal{P}_{\omega_1}(\mathbb{R})$" and give some applications. First we show that "ZFC + there exist ω^2 Woodin cardinals"[1] has the same consistency strength as "$\mathsf{AD} + \omega_1$ is $\mathbb{R}$-supercompact". During this process we show that if $L(\mathbb{R}, \mu) \vDash \mathsf{AD}$ then in fact $L(\mathbb{R}, \mu) \vDash \mathsf{AD}^+$. Next we prove important properties of $L(\mathbb{R}, \mu)$ including Σ_1-reflection and the uniqueness of μ in $L(\mathbb{R}, \mu)$. Then we give the computation of full HOD in $L(\mathbb{R}, \mu)$. Finally, we use Σ_1-reflection and $\mathbb{P}_{\max}$ forcing to construct a certain ideal on $\mathcal{P}_{\omega_1}(\mathbb{R})$ (or equivalently on $\mathcal{P}_{\omega_1}(\omega_2)$ in this situation) that has the same consistency strength as "ZFC + there exist ω^2 Woodin cardinals."

§1. Introduction. Recall that under $\mathsf{ZF} + \mathsf{DC}$, a measure μ on $\mathcal{P}_{\omega_1}(\mathbb{R})$, the set of countable subsets of $\mathbb{R}$, is:

(1) *fine* iff $\{\sigma \in \mathcal{P}_{\omega_1}(\mathbb{R}) \mid x \in \sigma\} \in \mu$ for each $x \in \mathbb{R}$;
(2) *normal* iff for each regressive $F : \mathcal{P}_{\omega_1}(\mathbb{R}) \to \mathcal{P}_{\omega_1}(\mathbb{R})$, that is

$$\{\sigma \in \mathcal{P}_{\omega_1}(\mathbb{R}) \mid F(\sigma) \subseteq \sigma \wedge F(\sigma) \neq \emptyset\} \in \mu$$

then

$$\exists x \in \mathbb{R}\, \{\sigma \in \mathcal{P}_{\omega_1}(\mathbb{R}) \mid x \in F(\sigma)\} \in \mu.$$

It's easy to see that if μ is a fine measure on $\mathcal{P}_{\omega_1}(\mathbb{R})$, ZF proves that normality of μ (condition (2) above) is equivalent to the following "diagonal intersection" property:

(2') If $\langle A_x \mid x \in \mathbb{R}\rangle$ is an $\mathbb{R}$-indexed sequence of μ-measure one sets, then

$$\triangle_{x\in\mathbb{R}} A_x =_{\text{def}} \{\sigma \mid \sigma \in \bigcap_{x\in\sigma} A_x\} \in \mu.$$

We first prove the following (previously unpublished) theorem, due to Woodin, which determines the exact consistency strength of the theory "$\mathsf{AD} + \omega_1$ is $\mathbb{R}$-supercompact".

THEOREM 1.1 (Woodin). *The following are equiconsistent.*

1. ZFC + *there are* ω^2 *Woodin cardinals.*
2. *There is a filter* μ *on* $\mathcal{P}_{\omega_1}(\mathbb{R})$ *such that* $L(\mathbb{R}, \mu) \vDash$ "$\mathsf{ZF} + \mathsf{DC} + \mathsf{AD} + \mu$ *is a normal fine measure on* $\mathcal{P}_{\omega_1}(\mathbb{R})$".

Received September 03, 2012.
Key words and phrases. Descriptive inner model theory, Solovay measure, HOD, derived model, Prikry forcing, $\mathbb{P}_{max}$ forcing.

[1] By this, we mean "ZFC + there is a set W of Woodin cardinals of order type ω^2". We will say "there exist ω^2 Woodin cardinals" for short.

© 2015, Association for Symbolic Logic
0022-4812/15/8001-0002
DOI:10.1017/jsl.2014.65

https://doi.org/10.1017/jsl.2014.65 Published online by Cambridge University Press

The proof of this theorem will occupy part of Section 2. The $(1) \Rightarrow (2)$ direction is proved using the derived model construction. The converse uses a Prikry forcing that forces a model of ZFC with ω^2 Woodin cardinals that realize $L(\mathbb{R}, \mu)$ as its derived model. This also shows that $L(\mathbb{R}, \mu) \vDash \mathsf{AD}$ if and only if $L(\mathbb{R}, \mu) \vDash \mathsf{AD}^+$.

It's worth mentioning that the existence of a normal fine measure on $\mathcal{P}_{\omega_1}(\mathbb{R})$ was first shown by Solovay to follow from $\mathsf{AD}_{\mathbb{R}}$ (see [8]); so $\mathsf{AD}_{\mathbb{R}}$ implies "$\mathsf{AD} + \omega_1$ is $\mathbb{R}$-supercompact". It also follows from [8] that the theory "$\mathsf{AD}+\omega_1$ is $\mathbb{R}$-supercompact" doesn't imply $\mathsf{AD}_{\mathbb{R}}$. Theorem 1.1 determines the exact consistency strength of the former, which is much weaker than that of the latter. It also follows from $\mathsf{AD}_{\mathbb{R}}$ that games on reals of fixed countable length are determined. This gives a hierarchy of normal fine measures extending the Solovay measure in some sense. A sequel to this paper ([16]) gives a construction (due to Woodin) of this hierarchy from $\mathsf{AD}_{\mathbb{R}}$, explores their exact consistency strength, and gives some applications of these measures.

Using the proof of Theorem 1.1, we explore the basic structure theory of $L(\mathbb{R}, \mu)$. We also prove in Section 2 the following theorem, which is also due to Woodin.

THEOREM 1.2 (Woodin). *The following holds in $L(\mathbb{R}, \mu)$ assuming $L(\mathbb{R}, \mu) \vDash$ "$\mathsf{AD}^+ + \mu$ is a normal fine measure on $\mathcal{P}_{\omega_1}(\mathbb{R})$".*

1. $(L_{\delta^2_1}(\mathbb{R})[\mu], \mu) \prec_{\Sigma_1} (L(\mathbb{R}, \mu), \mu)$.
2. *Suppose $L(\mathbb{R}, \mu) \vDash$ "μ_0 and μ_1 are normal fine measures on $\mathcal{P}_{\omega_1}(\mathbb{R})$". Then*
$$L(\mathbb{R}, \mu) \vDash \mu_0 = \mu_1.$$

Using Theorems 1.1, 1.2, and their proofs, we give some applications in Sections 3 and 4. Section 3 is dedicated to the HOD computation in $L(\mathbb{R}, \mu)$. The precise definition of HOD will be given in Section 3. Roughly speaking, HOD of $L(\mathbb{R}, \mu)$ will be shown to be $L(\mathcal{M}_\infty, \Lambda)$ where $\mathcal{M}_\infty \subseteq \mathrm{HOD}$ is a fine-structural premouse that has ω^2 Woodin cardinals cofinal in $o(\mathcal{M}_\infty)$, where $o(\mathcal{M}_\infty)$ is the ordinal height of the transitive structure $\mathcal{M}_\infty$, and agrees with HOD on all bounded subsets of Θ and Λ is a certain strategy that acts on finite stacks of normal trees in $\mathcal{M}_\infty$ based on $\mathcal{M}_\infty|\Theta$. The reader familiar with the HOD analysis in $L(\mathbb{R})$ will not be surprised here. As an application, [16] uses the HOD analysis to prove a "determinacy transfer theorem" which roughly states that the determinacy for real games of length ω^2 with payoff $\underset{\sim}{\Pi}^1_1$ and those with payoff $<\omega^2$-$\underset{\sim}{\Pi}^1_1$ are equivalent.

Finally, in Section 4 we prove the following two theorems. The first one uses $\mathbb{P}_{\max}$ forcing over a model of the form $L(\mathbb{R}, \mu)$ as above and the second one is an application of the core model induction. Woodin's book [19] or Larson's handbook article [5] are good sources for $\mathbb{P}_{\max}$; for details on the core model induction, see [7].

THEOREM 1.3. *Suppose $L(\mathbb{R}, \mu) \vDash$ "$\mathsf{AD}^+ + \mu$ is a normal fine measure on $\mathcal{P}_{\omega_1}(\mathbb{R})$" and let $G \subseteq \mathbb{P}_{max}$ be a generic filter over $L(\mathbb{R}, \mu)$. Then in $L(\mathbb{R}, \mu)[G]$, there is a normal fine ideal $\mathcal{I}$ on $\mathcal{P}_{\omega_1}(\mathbb{R})$ such that*

1. *letting $\mathcal{F}$ be the dual filter of $\mathcal{I}$ and $A \subseteq \mathbb{R}$ such that A is OD_x for some $x \in \mathbb{R}$, either $A \in \mathcal{F}$ or $\mathbb{R} \setminus A \in \mathcal{F}$;*
2. *$\mathcal{I}$ is precipitous;*
3. *for all $s \in \mathrm{OR}^\omega$, for all generics $G_0, G_1 \subseteq \mathcal{I}^+$, letting $j_{G_i} : V \to Ult(V, G_i) = M_i$ for $i \in \{0, 1\}$ be the generic embeddings, then $j_{G_0} \restriction \mathrm{HOD}_{\{\mathcal{I}, s\}} = j_{G_1} \restriction \mathrm{HOD}_{\{\mathcal{I}, s\}}$ and $\mathrm{HOD}^{M_0}_{\mathbb{R}^V \cup \{\mathbb{R}^V\}} = \mathrm{HOD}^{M_1}_{\mathbb{R}^V \cup \{\mathbb{R}^V\}} \in V$.*

https://doi.org/10.1017/jsl.2014.65 Published online by Cambridge University Press

The next theorem establishes the equiconsistency of the conclusion of Theorem 1.3 with the existence of ω^2 Woodin cardinals.

THEOREM 1.4 (ZFC). *Suppose there is a normal fine ideal $\mathcal{I}$ on $\mathcal{P}_{\omega_1}(\mathbb{R})$ such that*

1. *letting $\mathcal{F}$ be the dual filter of $\mathcal{I}$ and $A \subseteq \mathcal{P}_{\omega_1}(\mathbb{R})$ such that A is OD_x for some $x \in \mathbb{R}$, either $A \in \mathcal{F}$ or $\mathbb{R}\backslash A \in \mathcal{F}$;*
2. *$\mathcal{I}$ is precipituous;*
3. *for all $s \in \mathrm{OR}^\omega$, for all generics $G_0, G_1 \subseteq \mathcal{I}^+$, letting $j_{G_i} : V \to Ult(V, G_i) = M_i$ for $i \in \{0,1\}$ be the generic embeddings, then $j_{G_0} \restriction \mathrm{HOD}_{\{\mathcal{I},s\}} = j_{G_1} \restriction \mathrm{HOD}_{\{\mathcal{I},s\}}$ and $\mathrm{HOD}^{M_0}_{\mathbb{R}^V \cup \{\mathbb{R}^V\}} = \mathrm{HOD}^{M_1}_{\mathbb{R}^V \cup \{\mathbb{R}^V\}} \in V$.*

Then in a generic extension $V[G]$ of V, there is a filter μ on $\mathcal{P}_{\omega_1}(\mathbb{R})$ such that

$$L(\mathbb{R},\mu) \vDash \text{“}\mathsf{AD} + \mu \text{ is a normal fine measure on } \mathcal{P}_{\omega_1}(\mathbb{R})\text{”}.$$

Basic notions and notations. For a transitive structure M, we let $o(M)$ denote the ordinal height of M. A transitive $\mathcal{M}$ is a *fine-structural premouse* or simply a *premouse* if $\mathcal{M} = (J_\alpha[E], \in, E, F^{\mathcal{M}})$, where E is a fine-extender sequence in the sense of [12] and $F^{\mathcal{M}}$ is the amenable code for the top extender of M, also in the sense of [12]. We write $\mathcal{M}|\gamma$ for the structure $\mathcal{N} = (J_\gamma[E \restriction \gamma], \in, E \restriction \gamma, F^{\mathcal{N}})$ and $\mathcal{M}||\gamma$ for $\mathcal{N} = (J_\gamma[E \restriction \gamma], \in, E \restriction \gamma, \emptyset)$. Note that $\mathcal{M}|\gamma = \mathcal{M}||\gamma$ if $\mathcal{M}|\gamma$ is passive, that is, its predicate for the top extender is empty. If $\mathcal{P}, \mathcal{Q}$ are premice, we write $\mathcal{P} \triangleleft \mathcal{Q}$ if there is some $\gamma \leq o(\mathcal{Q})$ such that $\mathcal{P} = \mathcal{Q}|\gamma$. For some $k \leq \omega$, a k-sound premouse $\mathcal{M}$ is (k,α,β)-iterable if player II (the good player) has a winning strategy in the game $\mathcal{G}_k(\mathcal{M},\alpha,\beta)$ (see [12], Section 4). We customarily call a k-sound premouse $\mathcal{M}$ that is $(k, 1, \omega_1+1)$-iterable (or (k,ω_1,ω_1+1)-iterable) a *mouse*. When the degree of soundness of $\mathcal{M}$ is clear from the context, we will neglect to mention it in our notations.

The structure $L(\mathbb{R},\mu)$ considered in this paper is a structure of the language $\mathcal{L}^* = \mathcal{L} \cup \{\dot{\mathbb{R}}, \dot{\mu}\}$, where $\mathcal{L}$ is the language of set theory, $\dot{\mu}$ is a unary predicate symbol, and $\dot{\mathbb{R}}$ is a constant symbol, whose intended interpretation is the reals of the model. We sometimes write $L(\mathbb{R})[\mu]$, or $(L(\mathbb{R})[\mu],\mu)$ for the same structure. If μ is a measure on $\mathcal{P}_{\omega_1}(\mathbb{R})$ and $P(v)$ is a property, we often write $\forall^*_\mu \sigma P(\sigma)$ for $\{\sigma \mid P(\sigma)\} \in \mu$. Also, we also say “$\omega_1$ is $\mathbb{R}$-supercompact” to mean “there is a normal fine measure on $\mathcal{P}_{\omega_1}(\mathbb{R})$”.

We use Θ to denote the supremum of α such that there is a surjection from $\mathbb{R}$ onto α. Under ZFC, Θ is simply the successor cardinal of the continuum. Assuming AD^+, which is a technical strengthening of AD (see [9] or [15] for more on AD^+), a *Solovay sequence* is a sequence $\langle \theta_\alpha \mid \alpha \leq \Omega \rangle$ such that: (i) θ_0 is the supremum of ordinals α such that there is an OD surjection from $\mathbb{R}$ onto α; (ii) if $\beta \leq \Omega$ is limit, then $\theta_\beta = \sup_{\gamma<\beta}\theta_\gamma$; (iii) if $\beta = \gamma + 1 \leq \Omega$, then letting $B \subseteq \mathbb{R}$ have Wadge rank θ_γ, θ_β is the supremum of α such that there is an $OD(B)$ surjection from $\mathbb{R}$ onto α. Suppose $\mathsf{AD}^+ + \Theta = \theta_0$. We let $\underset{\sim}{\delta}^2_1$ denote the largest Suslin cardinal. The largest pointclass with the scales property, as shown by Woodin, is $\underset{\sim}{\Sigma}^2_1$.

For cardinals $\alpha \leq \beta$, we write $Col(\alpha, < \beta)$ for the Lévy collapse that adds a surjection from α onto every $\kappa \in [\alpha,\beta)$. If $\beta > \alpha$ is inaccessible then after forcing with $Col(\alpha, < \beta)$, β has cardinality α^+; otherwise, β will have cardinality α.

https://doi.org/10.1017/jsl.2014.65 Published online by Cambridge University Press

Finally, suppose γ is a limit of Woodin cardinals. We let $Hom_{<\gamma}$ denote the collection of $<\gamma$-homogeneously Suslin sets of reals. See [9] for more on the basic theory of $Hom_{<\gamma}$.

§2. The equiconsistency and structure theory of $L(\mathbb{R},\mu)$. We first present a variation of the derived model construction in [9] in the context where we want to construct a model of the form $L(\mathbb{R},\mu)$. See [9] for facts about AD^+ and the derived model construction.

LEMMA 2.1. *Suppose there is a measurable cardinal. Then there is a forcing $\mathbb{P}$ such that in $V^{\mathbb{P}}$, $L(\mathbb{R},\mathcal{C}) \vDash$ "$\mathcal{C}$ is a normal fine measure on $\mathcal{P}_{\omega_1}(\mathbb{R})$" where $\mathcal{C}$ is the club filter on $\mathcal{P}_{\omega_1}(\mathbb{R})$.*

PROOF. Let κ be a measurable cardinal and U be a normal measure on κ. Let $j : V \to M$ be the ultrapower map by U. Let $\mathbb{P}_0$ be $Col(\omega, <\kappa)$. Let $G \subseteq \mathbb{P}_0$ be V-generic. For $\alpha < \kappa$, we write $G \restriction \alpha$ for $G \cap Col(\omega, <\alpha)$. $Col(\omega, <j(\kappa)) = j(\mathbb{P}_0)$ is isomorphic to $\mathbb{P}_0 \Lambda \mathbb{Q}$ for some $\mathbb{Q}$ and whenever $H \subseteq \mathbb{Q}$ is $V[G]$-generic, then j can be lifted to an elementary embedding $j^+ : V[G] \to M[G][H]$ defined by $j^+(\tau_G) = j(\tau)_{G \Lambda H}$. Let $\mathbb{R}^{**} = \cup_{\alpha<\kappa} \mathbb{R}^{V[G|\alpha]}$ be the symmetric reals. Note that since κ is inaccessible, $\mathbb{R}^{**} = \mathbb{R}^{V[G]}$. We define a filter $\mathcal{F}^*$ on $\mathcal{P}_{\omega_1}(\mathbb{R}^{**})$ as follows.

$$A \in \mathcal{F}^* \Leftrightarrow \forall H \subseteq \mathbb{Q}(H \text{ is } V[G]\text{-generic} \Rightarrow \mathbb{R}^{V[G]} \in j^+(A)).$$

It's clear from the definition that $\mathcal{F}^* \in V[G]$.

We first claim that $\mathcal{F}^*$ is a normal fine filter. Fineness is easy; so we just verify normality. To see normality, suppose F is regressive. Then $A := \{\sigma \mid F(\sigma) \subseteq \sigma \wedge F(\sigma) \neq \emptyset\} \in \mathcal{F}$. Then $j^+(F)(\mathbb{R}^*) \subseteq \mathbb{R}^{**} \wedge j^+(F)(\mathbb{R}^*) \neq \emptyset$. Fix some $x \in \mathbb{R}^{**}$ such that $x \in j^+(F)(\mathbb{R}^{**})$. Then $\forall^*_{\mathcal{F}} \sigma\ x \in F(\sigma)$.

We now claim that $L(\mathbb{R}^{**}, \mathcal{F}^*) \vDash \mathcal{F}^*$ is a measure on $\mathcal{P}_{\omega_1}(\mathbb{R}^{**})$. Suppose $A \in L(\mathbb{R}^{**}, \mathcal{F}^*)$ is defined in $V[G]$ by a formula φ from a real $x \in \mathbb{R}^{**}$ (without loss of generality, we suppress parameters $\{U, s\}$, where $s \in \mathrm{OR}^{<\omega}$ that go into the definition of A); so $\sigma \in A \Leftrightarrow V[G] \vDash \varphi[\sigma, x]$. Let $\alpha < \kappa$ be such that $x \in V[G \restriction \alpha]$ and we let U^* be the canonical extension of U in $V[G \restriction \alpha]$. Then either

$$\forall^*_{U^*} \beta V[G \restriction \alpha] \vDash \emptyset \Vdash_{Col(\omega,<\beta)} \emptyset \Vdash_{Col(\omega,<\kappa)} \varphi[\dot{\mathbb{R}}_\beta, x]$$

or

$$\forall^*_{U^*} \beta V[G \restriction \alpha] \vDash \emptyset \Vdash_{Col(\omega,<\beta)} \emptyset \Vdash_{Col(\omega,<\kappa)} \neg\varphi[\dot{\mathbb{R}}_\beta, x].$$

In the above, $\dot{\mathbb{R}}_\beta$ is the canonical $Col(\omega, <\beta)$-name for the symmetric reals in $V^{Col(\omega,<\beta)}$. This easily implies either $A \in \mathcal{F}^*$ or $\neg A \in \mathcal{F}^*$.

Next, note $\mathcal{P}_{\omega_1}(\mathbb{R}^{**})$ has size ω_1 in $V[G]$, so we can use the iterated club shooting construction to turn $\mathcal{F}^*$ into the club filter. We let $\mathbb{P}_1$ be the forcing defined in 17.2 of [1]. By 17.2 of [1], $\mathbb{P}_1$ does not add any ω-sequence of ordinals. In particular, it does not add reals. Letting $H \subseteq \mathbb{P}_1$ be $V[G]$-generic, in $V[G][H]$, we still have $L(\mathbb{R}^{**}, \mathcal{F}^*) \vDash$ "$\mathcal{F}^*$ is a normal fine measure on $\mathcal{P}_{\omega_1}(\mathbb{R})^{**}$" and furthermore, $\mathcal{F}^* \cap L(\mathbb{R}^{**}, \mathcal{F}^*)$ is the restriction of the club filter on $L(\mathbb{R}^{**}, \mathcal{F}^*)$. Our desirable $\mathbb{P}$ is $\mathbb{P}_0 \Lambda \mathbb{P}_1$. ⊣

Suppose there exist ω^2 many Woodin cardinals. Let γ be the sup of the first ω^2 Woodin cardinals and for each $i < \omega$, let η_i be the sup of the first ωi Woodin cardinals. Suppose $G \subseteq Col(\omega, <\gamma)$ is V-generic and for each i, let $\mathbb{R}^* = \cup_{\alpha<\gamma} \mathbb{R}^{V[G\restriction\alpha]}$

https://doi.org/10.1017/jsl.2014.65 Published online by Cambridge University Press

and $\sigma_i = \mathbb{R}^{V[G\restriction Col(\omega,<\eta_i)]}$. We define a filter $\mathcal{F}^*$ as follows: for each $A \subseteq \mathcal{P}_{\omega_1}(\mathbb{R}^*)$ in $V[G]$

$$A \in \mathcal{F}^* \Leftrightarrow \exists n \forall m \geq n (\sigma_m \in A).$$

We call $\mathcal{F}^*$ defined above the **tail filter**.

LEMMA 2.2. *Let $\gamma, \eta_i, \mathbb{R}^*, \mathcal{F}^*$ be as above. Then*

$$L(\mathbb{R}^*, \mathcal{F}^*) \vDash \text{“}\mathcal{F}^* \textit{ is a normal fine measure on } \mathcal{P}_{\omega_1}(\mathbb{R}^*)\text{”}.$$

PROOF. Suppose not. So this statement is forced by the empty condition in $Col(\omega, < \gamma)$ by the homogeneity of $Col(\omega, < \gamma)$. By Lemma 2.1 applied to the first measurable cardinal κ and the fact that the forcing $\mathbb{P}$ used there is of size less than the first Woodin cardinal, by working over $V[g]$, where $g \subseteq \mathbb{P}$ is V-generic, we may assume that in V, the club filter $\mathcal{F}$ on $\mathcal{P}_{\omega_1}(\mathbb{R})$ has the property that $L(\mathbb{R}, \mathcal{F}) \vDash \mathcal{F}$ is a normal fine measure on $\mathcal{P}_{\omega_1}(\mathbb{R})$. Let $\lambda >> \gamma$ be regular and let

$$\begin{aligned} S = \{X \prec V_\lambda \mid\ & X \text{ is countable}, \gamma \in X, \exists \eta \in X \cap \gamma \text{ such that} \\ & \text{for all successor Woodin cardinals } \lambda \in X \cap (\eta, \gamma), \text{ if } D \subseteq \mathbb{Q}_{<\lambda}, \\ & D \in X, \text{ and } D \text{ is predense then } X \text{ captures } D\}. \end{aligned}$$

By Lemma 3.1.14 of [4], S is stationary and furthermore, letting $H \subseteq \mathcal{P}(\mathcal{P}_{\omega_1}(V_\lambda))/\mathcal{I}_{NS}$ [2] be generic such that $S \in H$, then for some $\xi < \gamma$, for all $\xi < \delta < \gamma$ and δ is Woodin, $H \cap \mathbb{Q}_{<\delta}$ is V-generic. We may as well assume ξ is less than the first Woodin cardinal and hence for all $\delta < \gamma$, δ is Woodin, $H \cap \mathbb{Q}_{<\delta}$ is V-generic.

Let $j : V \to (M, E)$ be the induced generic embedding given by H. Of course, (M, E) may not be wellfounded but wellfounded at least up to λ because $j''\lambda \in M$. For each $\alpha < \omega^2$, let $j_\alpha : V \to M_\alpha$ be the induced embedding by $H \cap \mathbb{Q}_{<\delta_\alpha}$, let M^* be the direct limit of the M_α's and $j^* : V \to M^*$ be the direct limit map. Note that j_α, j^* factor into j.

Let $\mathbb{R}^* = \mathbb{R}^{M^*}$ (the $\mathbb{R}^*$ from before is behind us now) and for each $i < \omega$, $\sigma_i = \mathbb{R}^{M_i^*}$ where $M_i^* = lim_n M_{\omega i+n}$. Let $G \subseteq Col(\omega, < \gamma)$ be such that $\cup_{\alpha<\eta_i} \mathbb{R}^{V[G\restriction\alpha]} = \sigma_i$ for all i; so $\mathbb{R}^*$ is the symmetric reals associated to G. Let $\mathcal{F}^*$ be the tail filter defined in $V[G]$. We claim that if $A \in j^*(\mathcal{F})$ then $A \in \mathcal{F}^*$. To see this, let $\pi \in M^*$ witness that A is a club. Let $\alpha < \omega^2$ be such that M_α contains the preimage of π. Then it is clear that $\forall m$ such that $\omega m \geq \alpha$ and $\pi'' \sigma_m \subseteq \sigma_m$. This shows $j^*(\mathcal{F}) \subseteq \mathcal{F}^*$ and hence $L_\lambda(\mathbb{R}^*, j^*(\mathcal{F})) = L_\lambda(\mathbb{R}^*, \mathcal{F}^*) \vDash$ “$\mathcal{F}^*$ is a normal fine measure on $\mathcal{P}_{\omega_1}(\mathbb{R}^*)$”. Since λ can be chosen arbitrarily large, we're done. ⊣

Next, we prove a “reflection phenomenon” analoguous to that in Lemma 6.4 of [9].

LEMMA 2.3. *Let $\gamma, G, \mathbb{R}^*, \mathcal{F}^*$ be defined as above. Suppose $x \in \mathbb{R}^{V[H\restriction\alpha]}$ for some $\alpha < \gamma$, and suppose ψ is a formula in the language of set theory with an additional predicate symbol. Let HC^* be the set of heritarily countable sets (in $V[G]$) coded by $\mathbb{R}^*$. Suppose*

$$\exists B \in \mathcal{P}(\mathbb{R}) \cap L(\mathbb{R}^*, \mathcal{F}^*)((HC^*, \in, B) \vDash \psi[x])$$

then

$$\exists B \in Hom_{<\gamma}^{V[G\restriction\alpha]}((HC^{V[G\restriction\alpha]}, \in, B) \vDash \psi[x]).$$

[2] In this section, $\mathcal{I}_{NS}$ is the nonstationary ideal on $\mathcal{P}_{\omega_1}(V_\lambda)$.

https://doi.org/10.1017/jsl.2014.65 Published online by Cambridge University Press

PROOF. Such a B in the statement of the lemma is called a ψ-witness. To see that Lemma 2.3 holds, pick the least γ_0 such that some $OD(x)^{L(\mathbb{R}^*,\mathcal{F}^*)}$ ψ-witness B is in $L_{\gamma_0}(\mathbb{R}^*,\mathcal{F}^*)$ and by minimizing the sequence of ordinals in the definition of B, we may assume B is definable (over $L_{\gamma_0}(\mathbb{R}^*,\mathcal{F}^*)$) from x without ordinal parameters. We may as well assume $x \in V$. We want to produce an absolute definition of B as in the proof of Lemma 6.4 in [9]. We do this as follows. First let φ be such that

$$u \in B \Leftrightarrow L_{\gamma_0}(\mathbb{R}^*,\mathcal{F}^*) \vDash \varphi[u,x],$$

and

$$\overline{\psi}(v) = \text{“}v \text{ is a } \psi\text{-witness”}.$$

Let $\mathcal{C}$ denote the club filter on $\mathcal{P}_{\omega_1}(\mathbb{R})$ and $\theta(u,v)$ be the natural formula defining B:

$$\begin{aligned}\theta(u,v) = \text{“}&L(\mathbb{R},\mathcal{C}) \vDash \mathcal{C} \text{ is a normal fine measure on } \mathcal{P}_{\omega_1}(\mathbb{R}) \text{ and } L(\mathbb{R},\mathcal{C}) \vDash \exists B \overline{\psi}[B] \\ &\text{and if } \gamma_0 \text{ is the least } \gamma \text{ such that } L_\gamma(\mathbb{R},\mathcal{C}) \vDash \exists B \overline{\psi}[B] \\ &\text{then } L_{\gamma_0}(\mathbb{R},\mathcal{C}) \vDash \varphi[u,v]\text{”}.\end{aligned}$$

We apply the tree production lemma (see [9]) to the definition $\theta(u,v)$ with parameter $x \in \mathbb{R}^V$. It's clear that stationary correctness holds. To verify generic absolutenss, let $\delta < \gamma$ be a Woodin cardinal; let g be $< \delta$ generic over V and h be $< \delta^+$ generic over $V[g]$. We want to show that if $y \in \mathbb{R}^{V[g]}$

$$V[g] \vDash \theta[y,x] \Leftrightarrow V[g][h] \vDash \theta[y,x].$$

There are $G_0, G_1 \subseteq Col(\omega, < \gamma)$ such that G_0 is generic over $V[g]$ and G_1 is generic over $V[g][h]$ with the property that $\mathbb{R}^*_{G_0} = \mathbb{R}^*_{G_1}$ and furthermore, if $\eta < \gamma$ is a limit of Woodin cardinals above δ, then $\mathbb{R}^*_{G_0} \restriction \eta = \mathbb{R}^*_{G_1} \restriction \eta$.[3] Such G_0 and G_1 exist since h is generic over $V[g]$ and $\delta < \gamma$. But this means letting $\mathcal{F}_i$ be the tail filter defined from G_i respectively then $L(\mathbb{R}^*_{G_0},\mathcal{F}_0) = L(\mathbb{R}^*_{G_1},\mathcal{F}_1)$. The proof of Lemma 2.2 implies that $L(\mathbb{R},\mathcal{C})^{V[g]}$ is embeddable into $L(\mathbb{R}^*_{G_0},\mathcal{F}_0)$ and $L(\mathbb{R},\mathcal{C})^{V[g][h]}$ is embeddable into $L(\mathbb{R}^*_{G_1},\mathcal{F}_1)$. This proves generic absoluteness. This gives us that $B \cap \mathbb{R}^V \in Hom^V_{<\gamma}$ and $B \cap \mathbb{R}^V$ is a ψ-witness. Hence we're done. ⊣

LEMMA 2.4. *Let $\gamma, \mathbb{R}^*, \mathcal{F}^*$ be defined as above. Then $L(\mathbb{R}^*,\mathcal{F}^*) \vDash \mathsf{AD}^+$.*

PROOF. Suppose not. Then any failure of AD^+ in $L(\mathbb{R}^*,\mathcal{F}^*)$ can be expressed in the form

$$(HC^*, \in, B) \vDash \psi[x]$$

for some $x \in \mathbb{R}^*$, some $B \in L(\mathbb{R}^*,\mathcal{F}^*) \cap \mathcal{P}(\mathbb{R})$, and some formula ψ. Using Lemma 2.3, we can get a ψ-witness B in $L(\mathbb{R}^*,\mathcal{F}^*)$ such that $B = C^*$, where $C \in Hom^{V[g]}_{<\gamma}$ for some $<\gamma$ generic g such that $x \in V[g]$ and C^* is the canonical blowup of C in the sense of [9]. The lemma then follows verbatim from the proof of Theorem 6.1 from Lemma 6.4 in [9]. ⊣

Now assume $L(\mathbb{R},\mu) \vDash$ “$\mathsf{AD} + \mu$ is a normal fine measure on $\mathcal{P}_{\omega_1}(\mathbb{R})$”. We prove that in a generic extension of $L(\mathbb{R},\mu)$, there is a class model N such that

[3] $\mathbb{R}^*_{G_0}$ is the symmetric reals defined by G_0 and similarly for $\mathbb{R}^*_{G_1}$. $\mathbb{R}^*_{G_0} \restriction \eta = \mathbb{R}^{V[g][G_0 \cap Col(\omega,<\eta)]}$ and $\mathbb{R}^*_{G_1} \restriction \eta = \mathbb{R}^{V[g][h][G_1 \cap Col(\omega,<\eta)]}$.

https://doi.org/10.1017/jsl.2014.65 Published online by Cambridge University Press

1. $N \vDash \mathsf{ZFC}$ + there are ω^2 Woodin cardinals;
2. letting λ be the sup of the Woodin cardinals of N, $\mathbb{R}$ can be realized as the symmetric reals over N via $Col(\omega, < \lambda)$;
3. letting $\mathcal{F}$ be the tail filter on $\mathcal{P}_{\omega_1}(\mathbb{R})$ in $N[G]$ where $G \subseteq Col(\omega, < \lambda)$ is a generic over N such that $\mathbb{R}$ is the symmetric reals induced by G, $L(\mathbb{R}, \mu) = L(\mathbb{R}, \mathcal{F})$ and $\mu \cap L(\mathbb{R}, \mu) = \mathcal{F} \cap L(\mathbb{R}, \mathcal{F})$.

The proof is given in Lemma 2.6. First we introduce some notions. Assume AD^+. Let T be a tree on $\omega \times \mathrm{OR}$ whose projection is a universal Σ^2_1 set. For any real x, by a Σ^2_1 *degree* d_x, we mean the equivalence class of all y such that $L[T, y] = L[T, x]$. Woodin has shown that the notion of Σ^2_1 degrees does not depend on the choice of T. In fact, we can define d_x to be the equivalence class of all y such that $\mathrm{HOD}_y = \mathrm{HOD}_x$. If d_1, d_2 are Σ^2_1 degrees, we say $d_1 \leq d_2$ if for any $x \in d_1$ and $y \in d_2$, $x \in L[T, y]$. $d_1 < d_2$ iff $d_1 \leq d_2$ and $d_1 \neq d_2$. For any reals x, y, we say $d_x = d_y$ or $x \equiv y$ iff $d_x \leq d_y$ and $d_y \leq d_x$. Just like with Turing cones, we define Σ^2_1 cones to be sets of the form $C_d = \{e \mid d \leq e\}$ for some Σ^2_1 degree d.

Theorem 2.5 (Woodin, see [3]). *Assume* AD^+. *Let* R, S *be sets of ordinals. Then for a (Turing or* Σ^2_1*) cone of* x, $\mathrm{HOD}^{L[R,S,x]}_R \vDash \omega_2^{L[R,S,x]}$ *is a Woodin cardinal.*

Lemma 2.6. *There is a forcing notion* $\mathbb{P}$ *in* $L(\mathbb{R}, \mu)$ *and there is an* N *in* $L(\mathbb{R}, \mu)^{\mathbb{P}}$ *satisfying* (1)–(3) *above.*

Proof. First, by arguments from [17], in $L(\mathbb{R}, \mu)$,

$$\Theta = \theta_0 + L(\mathcal{P}(\mathbb{R})) \vDash \Theta = \theta_0 + \mathsf{MC}.^4$$

Hence Σ^2_1 is the largest Suslin pointclass in $L(\mathbb{R}, \mu)$ and by Theorem 17.1 of [10], every set of reals in $L(\mathbb{R}, \mu)$ is contained in an $\mathbb{R}$-mouse.[5] Working in $L(\mathbb{R}, \mu)$, fix a tree T for a universal Σ^2_1 set as before (we may take T to be OD in $L(\mathcal{P}(\mathbb{R}))$). Let

$$\mathbb{D} = \{\langle d_i \mid i < \omega\rangle \mid \forall i (d_i \text{ is a } \Sigma^2_1 \text{ degree and } d_i < d_{i+1})\}.$$

Next, we define a measure ν on $\mathbb{D}$. We say

$$A \in \nu \quad \text{iff} \quad \text{for any } \infty\text{-Borel code} S \text{ for } A,$$
$$\forall^*_\mu \sigma \; L[T, S](\sigma) \vDash \text{“}\mathsf{AD}^+ + \sigma = \mathbb{R} + \exists (\emptyset, U) \in \mathbb{P}_{\Sigma^2_1}\, (\emptyset, U) \Vdash \dot{G} \in \mathcal{A}_S\text{”}.$$

In the definition of ν, $\mathbb{P}_{\Sigma^2_1}$ is the usual Prikry forcing using the Σ^2_1 degrees (see, e.g., Section 6.2 of [3]) and the cone measure in $L[T, S](\sigma)$, $\dot{G}$ is the name for the corresponding Prikry sequence, $\mathcal{A}_S$ is the set of reals coded by S. Note that:

(a) for all set of ordinals S, $\forall^*_\mu \sigma \; L[T, S](\sigma) \vDash \text{“}\mathsf{AD}^+ + \sigma = \mathbb{R}\text{”}$;
(b) whether $A \in \nu$ does not depend on the choice of S;
(c) for $A \subseteq \mathcal{P}_{\omega_1}(\mathbb{R})$, let $A^* = \{d \in \mathbb{D} \mid \cup d \in A\}$[6], then $A \in \mu \Leftrightarrow A^* \in \nu$.

We verify (b). Let S_0, S_1 be ∞-Borel codes for A. Let $T^\infty = \prod_\sigma T/\mu$ and $S^\infty_i = \prod_\sigma S_i/\mu$ be the ultraproducts by μ.

Claim. $L[T^\infty, S^\infty_0](\mathbb{R}) \cap \mathcal{P}(\mathbb{R}) = L[T^\infty, S^\infty_1](\mathbb{R}) \cap \mathcal{P}(\mathbb{R}) = L(\mathbb{R}, \mu) \cap \mathcal{P}(\mathbb{R})$.

[4] MC is the statement that whenever $x, y \in \mathbb{R}$ are such that x is OD_y, then there is a sound mouse $\mathcal{M}$ over y such that $\rho(\mathcal{M}) = \omega$ and $x \in \mathcal{M}$.

[5] $\mathcal{M}$ is an $\mathbb{R}$-mouse if $\mathcal{M}$ is a premouse over $\mathbb{R}$ in the sense of [11], $\mathcal{M}$ is ω-sound, $\rho(\mathcal{M}) = \mathbb{R}$, and the transitive collapse of every countable substructure of $\mathcal{M}$ is $(\omega, 1, \omega_1 + 1)$-iterable.

[6] Say $d = \langle d_i \mid i < \omega\rangle$; then $\cup d = \{x \in \mathbb{R} \mid \exists n \; x \in L[T, d_n]\}$.

https://doi.org/10.1017/jsl.2014.65 Published online by Cambridge University Press

PROOF. To see this, first observe that by MC in $L(\mathcal{P}(\mathbb{R}))$, $\mathcal{P}(\mathbb{R}) = \mathcal{P}(\mathbb{R}) \cap Lp(\mathbb{R})$ by [10, Theorem 17.1];[7] the second observation is by Los, $Lp(\mathbb{R}) = \prod_\sigma Lp(\sigma)/\mu$; the final observation is $\forall^*_\mu \sigma\ L[T, S_0](\sigma) \cap \mathcal{P}(\sigma) = L[T, S_1](\sigma) \cap \mathcal{P}(\sigma) = OD(\sigma) \cap \mathcal{P}(\sigma) = Lp(\sigma) \cap \mathcal{P}(\sigma)$.

To see the final observation, note that for $i \in \{0, 1\}$, $L[T^\infty, S^\infty_i](\mathbb{R}) \cap \mathcal{P}(\mathbb{R}) \subseteq L(\mathbb{R}, \mu) \cap \mathcal{P}(\mathbb{R}) = Lp(\mathbb{R}) \cap \mathcal{P}(\mathbb{R})$, so by Los, $\forall^*_\mu \sigma\ L[T, S_i](\sigma) \cap \mathcal{P}(\sigma) \subseteq Lp(\sigma) \cap \mathcal{P}(\sigma)$. To see the converse, it suffices to prove the following claim, whose proof is based on an unpublished note of J.R. Steel.

SUBCLAIM. In $L(\mathcal{P}(\mathbb{R}))$, there is a real z such that whenever a is countable, transitive, and $z \in a$, then $\mathcal{P}(a) \cap L[T, a] = \mathcal{P}(a) \cap OD(a)$.[8]

PROOF. First we prove that: on a cone of reals z, $\mathbb{R} \cap L[T, z] = \mathbb{R} \cap OD(z)$. To prove this, first let

$$A(z, n, m) \Leftrightarrow \exists y \in \mathbb{R}\ (y \in OD(z) \backslash L[T, z]) \wedge \text{letting } y_z \text{ be the } OD(z)\text{-least such } y, \text{ then } y_z(n) = m.$$

Now it is a basic AD^+ fact that since $\Theta = \theta_0$, there is a real z_0 such that for all z Turing above z_0, $A \cap L[T, z] \in L[T, z]$ (in other words, the (boldface) envelope of Σ^2_1 is $\mathcal{P}(\mathbb{R})$). For any such z, $OD(z) \cap \mathbb{R} = L[T, z] \cap \mathbb{R}$. To see this, if not, then $y_z(n) = m$ if and only if $A(z, n, m)$. So y_z is computable from $A \cap L[T, z]$ so y_z is in $L[T, z]$. This contradicts the definition of y_z.

Take z_0 to be the base of the cone in the above argument. For any countable transitive a such that $z_0 \in a$, a set $b \subseteq a$ is $OD(a)$ just in case for comeager many enumerations g of a (in order type ω), b is $OD(g)$. This and the above argument give the subclaim. ⊣

The subclaim and MC give

$$\forall^*_\mu \sigma\ Lp(\sigma) \cap \mathcal{P}(\sigma) = OD(\sigma) \cap \mathcal{P}(\sigma) \subseteq L[T, S_i](\sigma) \cap \mathcal{P}(\sigma).$$

This completes the proof of the third observation. The three observations give us the claim. ⊣

The claim gives us that the $\mathbb{P}_{\Sigma^2_1}$ forcing relations in these models are the same, in particular, $L[T^\infty, S^\infty_0](\mathbb{R}) \vDash \exists (\emptyset, U) \in \mathbb{P}_{\Sigma^2_1}\ (\emptyset, U) \Vdash \dot{G} \in \mathcal{A}_{S^\infty_0}$ if and only if $L[T^\infty, S^\infty_1](\mathbb{R}) \vDash \exists (\emptyset, U) \in \mathbb{P}_{\Sigma^2_1}\ (\emptyset, U) \Vdash \dot{G} \in \mathcal{A}_{S^\infty_1}$. This gives us (b). (a) follows from the claim and Los theorem.

To see (c), suppose $A \in \mu$. Let S be an ∞-Borel code for A^*. By (a),

$$\forall^*_\mu \sigma\ (\sigma \in A \wedge L[T, S^*](\sigma) \vDash \text{“}\mathsf{AD}^+ + \sigma = \mathbb{R}\text{”}).$$

For any such σ, if d is the sequence of degrees corresponding to a $\mathbb{P}_{\Sigma^2_1}$-generic over $L[T, S^*](\sigma)$, then clearly $\cup d = \sigma \in A$ since d is cofinal in the Σ^2_1 degrees of $L[T, S^*](\sigma)$. This means $d \in A^*$. This gives $A^* \in \nu$. The converse is proved using the proof of the forward direction applied to $\mathcal{P}_{\omega_1}(\mathbb{R}) \backslash A$. This finishes the proof of (c).

Let $\mathbb{P}$ be the usual Prikry forcing using ν (cf. [3]). First let $\nu_1 = \nu$, for $n > 0$, let ν_n be the product measure induced by ν_0 on $\mathbb{D}^{n+1}$; that is $\nu_n(Z) = 1 \Leftrightarrow \forall^*_\nu d_0 \cdots \forall^*_\nu d_n \langle d_0, \ldots, d_n \rangle \in Z$. Conditions in $\mathbb{P}$ are pairs (p, U) where for some

[7] In [11], $Lp(\mathbb{R})$ is denoted $K(\mathbb{R})$ and is the stack of all $\mathbb{R}$-mice.

[8] We remind the reader that T is OD; so $OD(a) = OD(T, a)$.

https://doi.org/10.1017/jsl.2014.65 Published online by Cambridge University Press

$n \in \omega$, $p = \langle \vec{d^i} \mid i \leq n \wedge \vec{d^i} \in \mathbb{D} \wedge \vec{d^i} \in d^{\vec{i+1}}(0)$[9]$\rangle$ and U is such that for all $n < \omega$, $U(n) \subseteq \mathbb{D}^{n+1}$, and $\nu_n(U(n)) = 1$. $(p, U) \leq_{\mathbb{P}} (q, W)$ if p end extends q, say $p = q^\frown r$ for some $r \in \mathbb{D}^n$, and for all k and all $s \in U(k)$, $r^\frown s \in W(n+k)$. $\mathbb{P}$ has the usual Prikry property, that is given any condition (p, U), a term τ, a formula $\varphi(x)$, we can find a $(p, U') \leq_{\mathbb{P}} (p, U)$ such that (p, U') decides the value of $\varphi[\tau]$; furthermore, (p, U') is ordinal definable from p, U, τ (see [13] or Section 6 of [3] for a proof). Let G be $\mathbb{P}$ generic. We identify G with the union of the stems of conditions in G, i.e., G is identified with $\langle \vec{d^i} \mid i < \omega \wedge \exists U(\langle d^j \mid j \leq i\rangle, U) \in G\rangle$. We need some notations before proceeding. We write V for $L(\mathbb{R}, \mu)$ (and use them interchangably); for any $g \in \mathbb{D}$, let $\omega_1^g = \sup_i \omega_1^{L[T^\infty, g(i)]}$ and $\delta(g \restriction i) = \omega_2^{L[T^\infty, g\restriction i]}$ (note that $\delta(g \restriction i)$ doesn't depend on the representatives for the degrees in g). To produce a model with ω^2 Woodin cardinals, we use Theorem 2.5.

For any countable transitive a which admits a well-ordering rudimentary in a and for any real x coding a, let

$$Q_a^x = \mathrm{HOD}^{L[T^\infty, x]}_{T^\infty, a} \restriction (\delta(x) + 1).$$

The expression on the right hand side above stands for $V_{\delta(x)+1} \cap \mathrm{HOD}^{L[T^\infty, x]}_{T^\infty, a}$. Note that Q_a^x only depends on the degree of x; hence for a cone of Σ^2_1-degree e, $Q_a^e = Q_a^x$ for all $x \in a$. Let a be a base of the cone in the subclaim above. We now let

$$Q_0^0 = Q_a^{\vec{d^0}(0)},$$

and

$$\delta_0^0 = \delta(\vec{d^0}(0)).$$

For $i < \omega$, let

$$Q_{i+1}^0 = Q_{Q_i^0}^{\vec{d^0}(i+1)},$$

and

$$\delta_{i+1}^0 = \delta(\vec{d^0}(i+1)).$$

This finishes the first block. Let $Q_\omega^0 = \cup_i Q_i^0$. In general, we let

$$Q_0^{j+1} = Q_{Q_\omega^j}^{\vec{d^{j+1}}(0)},$$

and

$$\delta_0^{j+1} = \delta(\vec{d^{j+1}}(0)).$$

For $i < \omega$, let

$$Q_{i+1}^{j+1} = Q_{Q_i^{j+1}}^{\vec{d^{j+1}}(i+1)},$$

and

$$\delta_{i+1}^{j+1} = \delta(\vec{d^{j+1}}(i+1)).$$

In $V[G]$, let

$$N =_{\mathrm{def}} L[T^\infty, \langle Q_j^i \mid i, j < \omega\rangle]$$

Note that N can be defined in $\mathrm{HOD}^{(V[G], V)}_{\{G\}}$. We claim that

$$N \vDash \delta_j^i \text{ is a Woodin cardinal for all } i, j < \omega.$$

[9] We abuse notation here to mean $\vec{d^i} \in L[T, d^{\vec{i+1}}(0)]$ and is countable there.

https://doi.org/10.1017/jsl.2014.65 Published online by Cambridge University Press

The claim follows from the following observations.

(a) For all $i, j < \omega$, $Q_0^{j+1} \cap \mathcal{P}(\delta_i^j) = Q_i^j \cap \mathcal{P}(\delta_i^j) = Q_{i+1}^j \cap \mathcal{P}(\delta_i^j)$.
(b) For $i, j < \omega$, $N \cap \mathcal{P}(\delta_j^i) = Q_j^i \cap \mathcal{P}(\delta_j^i)$.

The second equality of (a) follows from basic facts about Prikry forcing (see Section 6.2 of [3]). Also from [3], we get $L[T^\infty, Q_\omega^i] \cap \mathcal{P}(\delta_j^i) = Q_j^i \cap \mathcal{P}(\delta_j^i)$ for all $i, j < \omega$.

For the first equality, it's enough to prove: $(\dagger) \equiv$ "for any n, for a cone of d, $\mathcal{P}(Q_\omega^n) \cap L[T^\infty, Q_\omega^n] = \mathcal{P}(Q_\omega^n) \cap L[T^\infty, d]$". $(\dagger)$ easily implies the first equality of (a). To see $(\dagger)$, suppose not. Note that $\mathcal{P}(Q_\omega^n) \cap L[T^\infty, Q_\omega^n] = \mathcal{P}(Q_\omega^n) \cap L[T, Q_\omega^n]$ and $L[T^\infty, d] \cap \mathcal{P}(Q_\omega^n) = L[T, d] \cap \mathcal{P}(Q_\omega^n)$ by Los theorem. Working in $L(\mathcal{P}(\mathbb{R}))$, for a cone of d, let b_d be the least $b \subseteq Q_\omega^n$ in $L[T, d] \backslash L[T, Q_\omega^n]$ (the minimality of b_d is in terms of the canonical well-ordering of $L[T, d]$). Since Q_ω^n is countable, there is a b and a cone of d such that $b = b_d$, so b is $OD_{Q_\omega^n}$. This means $b \in L[T, Q_\omega^n]$ (by the subclaim and the choice of Q_0^0). Contradiction.

Now to see (b), we use the Prikry property of $\mathbb{P}$. Let $A \subseteq \delta_j^i$ be in N. Then A is ordinal definable in $V[G]$ from $\{T^\infty, \langle Q_j^i \mid i, j < \omega\rangle\}$. Let $\dot{Q}$ be the canonical forcing term for $\langle Q_j^i \mid i, j < \omega\rangle$ and $\varphi(v, \hat{t}, \dot{Q})$ be a formula in the forcing language with only v free and $t \in \mathrm{OR}^{<\omega} \cup \{T^\infty\}$ such that φ defines A over $V[G]$ from t and $\langle Q_j^i \mid i, j < \omega\rangle$. Let $(p, U) \in G$ with $\mathrm{dom}(p) > i$. By the fact that δ_j^i is countable, the Prikry property gives a condition $(p, Y) \leq (p, U)$ such that (p, Y) decides $\varphi(\hat{\eta}, \hat{t}, \dot{Q})$ for all $\eta < \delta_j^i$. By density, we may fix such a $(p, Y) \in G$. Letting $n + 1 = \mathrm{dom}(p)$, we claim

$$\eta \in A \Leftrightarrow \exists r \in \mathbb{D}^{\mathrm{dom}(p)+1} \exists X\ (r, X) \Vdash \varphi(\hat{\eta}, \hat{t}, \dot{Q}) \wedge \forall i \leq n \forall j < \omega\ Q_j^i = (Q_j^i)^r,$$

where in the above $(Q_j^i)^r$ is the model Q_j^i defined relative to the sequence of degrees given by r (over the set a specified above). If the equivalence holds, then A is OD from T^∞ and $\langle Q_j^i \mid j < \omega \wedge i \leq n\rangle$. By the proof of (a), we get that $A \in Q_j^i$, which is what we want to prove. We've already shown the $\Rightarrow$ direction. To see the converse, suppose (r, X) is as on the right hand clause but $\eta \notin A$, then we have $(p, Y) \Vdash \neg\varphi(\hat{\eta}, \hat{t}, \dot{Q})$. Letting $Z(n) = X(n) \cap Y(n)$, we have $(r, Z) \leq (r, X)$ and $(p, Z) \leq (p, Y)$. Let $H \subseteq \mathbb{P}$ be V-generic with $(p, Z) \in H$ and $p^\frown\langle e_i \mid i > n\rangle$ be the Prikry sequence determined by H. It's easy to see that $r^\frown\langle e_i \mid i < \omega\rangle$ is a Prikry sequence giving rise to a generic I such that

$$(r, Z) \in I \wedge V[H] = V[I].$$

But then since $(Q_j^i)^r = (Q_j^i)^p$ for all $j < \omega, i \leq n$, $\dot{Q}^H = \dot{Q}^I$; and so both $\varphi(\dot{\eta}, \dot{t}, \dot{Q})$ and its negation hold in $V[H]$. Contradiction.[10]

Letting $\lambda = \sup_{i,j} \delta_j^i$ and $\gamma_i = \sup_{j<\omega} \delta_j^i$, by the construction of N, there is a $H \subseteq Col(\omega, < \lambda)$ generic over N such that $\mathbb{R}_H^* = \mathbb{R}^V$. To see this, it suffices to see that every $x \in \mathbb{R}^V$ is N-generic for some poset in V_λ^N. Pick n such that $x \in L[T^\infty, y]$ for some (any) $y \in \vec{d}^n(0)$. In $L[T^\infty, y]$, x is generic over $\mathrm{HOD}_{T^\infty, \langle Q_j^i \mid i<n \wedge j<\omega\rangle}$ for the Vopenka poset $\mathbb{B}$ (this gives also $\mathbb{B} \in N$). A theorem of Becker and Woodin

[10] We note that there is a canonical name $\dot{N}$ for N and the proof above gives a condition of the form $(0, U)$ forcing that $\dot{N}$ has ω^2 Woodin cardinals.

https://doi.org/10.1017/jsl.2014.65 Published online by Cambridge University Press

states that on a cone of x, $L[T^\infty, x]$ satisfies $2^\alpha = \alpha^+$ for all $\alpha < \omega_1^V$. Since we can work in that cone from the beginning (i.e., can demand $\vec{d}^0(0)$ is in the cone), in $L[T, y]$, $2^\omega = \omega_1$ and $2^{\omega_1} = \omega_2 = \delta_0^n$. Hence in $L[T, y]$, $|\mathbb{B}| = \delta_0^n < \lambda$. Furthermore, since $Q_0^n = V^N_{\delta_0^n+1}$, x is $\mathbb{B}$-generic over M. We're done.

Recall G is the sequence $\langle \vec{d}^i \mid i < \omega\rangle$. For each $i < \omega$, let $\sigma_i = \cup_{\alpha<\gamma_i} \mathbb{R}^{H|\alpha} = \cup \vec{d}^i$.[11] In $N[H]$, let $\mathcal{F}$ be the tail filter defined by the sequence $\langle \sigma_i \mid i < \omega\rangle$. It remains to see that $L(\mathbb{R},\mu) = L(\mathbb{R},\mathcal{F})$ and $\mu \cap L(\mathbb{R},\mu) = \mathcal{F} \cap L(\mathbb{R},\mathcal{F})$. For this it's enough to show $\mu \subseteq \mathcal{F}$. Let $A \in \mu$. Then $A^* = \{d \in \mathbb{D} \mid \cup d \in A\} \in \nu$. Since $\mathbb{P}$ is the Prikry forcing relative to ν, $\exists n \forall m \geq n\ \vec{d}^m \in A^*$; this means $\exists n \forall m \geq n\ \sigma_m \in A$. This implies $A \in \mathcal{F}$. On the other hand, if $A \notin \mu$ then $\nu(A^*) = 0$. This implies $\neg A \in \mathcal{F}$. So $\mu \subseteq \mathcal{F}$. ⊣

Proof of Theorem 1.1. The (1) ⇒ (2) direction follows from Lemmas 2.4 and 2.2. The (2) ⇒ (1) direction follows from Lemma 2.6 ⊣

Proof of Theorem 1.2. Let $N, \lambda, G, \mathcal{F}$ be defined as in the paragraph after the proof of Lemma 2.4. In $N[G]$, let $D = L(\Gamma,\mathbb{R})$[12] where $\Gamma = \{A \subseteq \mathbb{R} \mid A \in N(\mathbb{R}) \wedge L(A,\mathbb{R}) \vDash \mathsf{AD}^+\}$. Woodin has shown that $D \vDash \mathsf{AD}^+$ and $\Gamma = \mathcal{P}(\mathbb{R})^D$ (see [20]). Letting T^∞ be defined as in the proof of Lemma 2.6, we already know $\mathcal{P}(\mathbb{R}) \cap L(\mathbb{R},\mu) = \mathcal{P}(\mathbb{R}) \cap L(T^\infty,\mathbb{R}) \subseteq \Gamma$ and $L(\mathbb{R},\mu) = L(\mathbb{R},\mathcal{F})$ and $\mu \cap L(\mathbb{R},\mu) = \mathcal{F} \cap L(\mathbb{R},\mathcal{F})$. Also, by the proof of Lemma 2.2 and the ⇒ direction of Theorem 1.1, $\mathcal{F} \cap L(\mathbb{R},\mathcal{F}) = \mathcal{C} \cap L(\mathbb{R},\mathcal{F})$ where $\mathcal{C}$ is the club filter on $\mathcal{P}_{\omega_1}(\mathbb{R})$ in $N[G]$.

Suppose $(L(\mathbb{R},\mu),\mu) \vDash \phi$ where ϕ is a Σ_1 statement. Then since Θ is regular in $L(\mathbb{R},\mu)$, there is a $\kappa < \Theta$ such that $(L_\kappa(\mathbb{R},\mu), \mu \cap L_\kappa(\mathbb{R},\mu)) \vDash \phi$. There is a set $B \subseteq \mathbb{R}$ in $L(\mathbb{R},\mu)$ such that B codes the structure $(L_\kappa(\mathbb{R},\mu), \mu \cap L_\kappa(\mathbb{R},\mu))$ and hence there is a φ such that

$$(L(\mathbb{R},\mu),\mu) \vDash \phi \Leftrightarrow (HC, \in, B) \vDash \varphi.$$

Such a B is called a φ-witness as before. We let γ_0 be the least such that $L_{\gamma_0}(\mathbb{R},\mu)$ ordinal defines a φ-witness. By minimizing the ordinal parameters, we assume then that the φ-witness B is definable over $L_{\gamma_0}(\mathbb{R},\mu)$ by (Φ, x) for some $x \in \mathbb{R}$, that is

$$y \in B \Leftrightarrow L_{\gamma_0}(\mathbb{R},\mu) \vDash \Phi[y, x].$$

By the construction of N and the proof of Lemma 2.3, there is $\alpha < \lambda$ and a $B \in Hom^{N[G\restriction\alpha]}_{<\lambda}$ such that

$$(HC^{N[G\restriction\alpha]}, \in, B) \vDash \varphi.$$

But $(HC^{N[G\restriction\alpha]}, \in, B) \prec (HC, \in, B^*)$ where $B^* \in (\underset{\sim}{\delta}^2_1)^{L(\mathbb{R},\mu)}$ is the canonical blowup of B by Lemma 6.3 of [9].[13] This gives us a $\kappa < \underset{\sim}{\delta}^2_1$ such that $(L_\kappa(\mathbb{R},\mu), \mu \cap L_\kappa(\mathbb{R},\mu)) \vDash \phi$.[14] Since ϕ is Σ_1, we have $(L_{\underset{\sim}{\delta}^2_1}(\mathbb{R},\mu), \mu \cap L_{\underset{\sim}{\delta}^2_1}(\mathbb{R},\mu)) \vDash \phi$.

---

[11] $\bigcup \vec{d}^i$ is union of all reals in a degree in $\vec{d}^i$.

[12] D is called the "new derived model" of N at λ.

[13] To see this, first note that $B^* \in L(\mathbb{R},\mu)$. By Theorem 4.3 of [9], B has a $Hom^{N[G\restriction\alpha]}_{<\lambda}$-scale and so does $\neg B$. This fact is projective in B so the structure $(HC, \in, B^*)$ sees that $B^*, \neg B^*$ both have a scale. Hence $B^* \in (\underset{\sim}{\delta}^2_1)^{L(\mathbb{R},\mu)}$.

[14] The proof of Lemma 2.3, in particular, the definition of the formula $\theta(u, v)$ there, tells us that B codes a structure of the form $(L_\kappa(\mathbb{R},\nu),\nu)$ where ν comes from the club filter in $N[G \restriction \alpha]$ and $\kappa < \underset{\sim}{\delta}^2_1$ so in fact $L_\kappa(\mathbb{R},\nu) = L_\kappa(\mathbb{R},\mu)$ and $\mu \cap L(\mathbb{R},\mu) = \nu \cap L(\mathbb{R},\nu)$.

https://doi.org/10.1017/jsl.2014.65 Published online by Cambridge University Press

This finishes the proof of (1) in Theorem 1.2. (2) of Theorem 1.2 is also a corollary of the proof of Lemma 2.6. One first modifies the definition of $\mathbb{P}$ in Lemma 2.6 by redefining the set U in the condition (p, U) to be: $U(2n) \in \nu_0$ and $U(2n+1) \in \nu_1$ for all n where ν_i is defined from μ_i in the exact way that ν is defined from μ in the proof of Lemma 2.6. Everything else in the proof of the lemma stays the same. This implies $L(\mathbb{R}, \mu_0) = L(\mathbb{R}, \mu_1) = L(\mathbb{R}, \mathcal{F})$ and $\mu_0 = \mu_1 = \mathcal{F}$. To see this, just note that since we already know

$$L(\mathbb{R}, \mathcal{F}) \vDash \mathsf{AD}^+ + \mathcal{F} \text{ is a normal fine measure on } \mathcal{P}_{\omega_1}(\mathbb{R}),$$

it suffices to show if $A \in \mathcal{F}$ then $A \in \mu_0$ and $A \in \mu_1$. Suppose there is an $A \in \mathcal{F}$ such that $A \in \mu_0$ and $A \notin \mu_1$ (the cases $A \in \mu_1 \backslash \mu_0$ and $A \notin \mu_0 \cap \mu_1$ are handled similarly). Let

$$A^* = \{d \in \mathbb{D} \mid \cup d \in A\}.$$

Then $A^* \in \nu_0 \backslash \nu_1$. For any condition (p, U), just shrink U to U^* by setting $U^*(2n) = U(2n) \cap A^*$ and $U^*(2n+1) = U(2n+1) \cap \neg A^*$. Then $(p, U^*) \Vdash A \notin \mathcal{F}$. Contradiction. This finishes the proof of Theorem 1.2. ⊣

§3. The HOD analysis. Throughout this section, we assume $L(\mathbb{R}, \mu) \vDash \mathsf{AD}^+$. The following theorem is due to Woodin.

THEOREM 3.1. *Suppose $L(\mathbb{R}, \mu) \vDash \mathsf{AD}^+ + \mu$ is a normal fine measure on $\mathcal{P}_{\omega_1}(\mathbb{R})$. Then in $L(\mathbb{R}, \mu)$, there is a set $A \subseteq \Theta$ such that* $\mathrm{HOD} = L[A]$.

PROOF. Working in $L(\mathbb{R}, \mu)$, let $N = L(\mathcal{P}(\mathbb{R}))$. Note that $\Theta^N = \Theta$ and $N \vDash \mathsf{AD}^+ + \Theta = \theta_0$ (see [17]). By general AD^+ theory,

1. $\mathrm{HOD}^N = L[B]$ for some $B \subseteq \Theta$ in HOD^N;
2. $\mathrm{HOD}^N[x] = \mathrm{HOD}^N_x$ for any $x \in \mathbb{R}$.

Let $\delta = \underset{\sim}{\delta}^2_1$. Since $\mu \cap L_\delta(\mathbb{R})[\mu]$ is the club filter, $N|\delta = L_\delta(\mathbb{R})[\mu]$ and hence HOD^N and HOD agree up to δ by Σ_1-reflection. Again, by general AD^+ theory, δ is strong to Θ via embeddings given by measures (see [3]) and these measures are unique (and hence OD) in N, hence HOD^N and HOD agree up to Θ. The same conclusion holds for HOD^N_x and HOD_x.[15] This is key to our proof.

Let $j : \mathrm{HOD} \to M$ be the ultrapower embedding given by μ using all functions in $L(\mathbb{R}, \mu)$. j is definable from μ. By Theorem 1.2, μ is unique hence j is OD. Similarly, μ also induces an embedding $j_x : HOD_x \to M_x$ for all $x \in \mathbb{R}$. Note that $\mathrm{HOD}^N[x] = \mathrm{HOD}^N[G_x]$ for a generic G_x for the Vopenka algebra whose elements are OD ∞-Borel codes. By (2) and the fact that $\mathrm{HOD}[x]^N|\Theta = \mathrm{HOD}[x]|\Theta$, j's restriction on bounded subsets of Θ can compute j_x's restriction on bounded subsets of Θ.

CLAIM. $L(\mathbb{R}, \mu) = L[\mathrm{HOD}^N, j \restriction \Theta](\mathbb{R}) = L[A](\mathbb{R})$ *for some* $A \subseteq \mathrm{OR}$ in HOD.[16]

PROOF. The second equality is clear since $\mathrm{HOD}^N = L[B]$ for some $B \subseteq \Theta$ so now we prove the first equality. First it's easy to see that

$$L[\mathrm{HOD}^{\mathrm{N}}, \mathrm{j} \restriction \Theta](\mathbb{R}) = \mathrm{L}[\mathrm{HOD}^{\mathrm{N}}_{\mathrm{x}}, \mathrm{j} \restriction \Theta](\mathbb{R}) = \mathrm{L}[\mathrm{HOD}^{\mathrm{N}}[\mathrm{x}], \mathrm{j} \restriction \Theta](\mathbb{R}). \qquad (*)$$

[15] This has a consequence that Mouse Capturing holds in $L(\mathbb{R}, \mu)$ since Mouse Capturing holds in N.
[16] By $j \restriction \Theta$ we mean the set of (a, γ) such that $a \in V^{\mathrm{HOD}}_\Theta$ and $\gamma \in j(a)$.

https://doi.org/10.1017/jsl.2014.65 Published online by Cambridge University Press

Let $X \subseteq \mathcal{P}_{\omega_1}(\mathbb{R})$. Note that $X \in N$. To see whether X is in μ, let S be an ∞-Borel code for X. S is a bounded subset of Θ. First suppose S is OD in N. So $X \in \mu$ if and only if whenever $g \subseteq Col(\omega, \mathbb{R})$ is generic over $L[\mathrm{HOD}^N, j \restriction \Theta](\mathbb{R})$, in $L[\mathrm{HOD}^N, j \restriction \Theta](\mathbb{R})[g]$, $\mathbb{R}$ is in the set with code $j(S)$. The case where S is OD^N_x for some $x \in \mathbb{R}$ can be handled by using $(*)$. This means $L[HOD^N, j \restriction \Theta](\mathbb{R})$ can compute μ by consulting the homogeneous forcing $Col(\omega, \mathbb{R})$; this gives us the first equality. $\dashv$

Pick a large γ and consider the elementary substructure Z of $L_\gamma[\mathrm{HOD}^N, j \restriction \Theta]$ consisting of elements definable in $L(\mathbb{R}, \mu)$ from $\{\mathrm{HOD}^N, j\}$, reals, and ordinals less than Θ. Hence Z is OD and has size at most Θ. Let j^* be the transitive collapse of j. Note that

$$\mathrm{HOD}^N = L[B]$$

for some $B \subseteq \Theta$ and since $\Theta \subseteq Z$, B collapses to itself. Hence there is a set $A \subseteq \Theta$ in HOD such that $L[\mathrm{HOD}^N, j^*] \subseteq L[A]$ and it's easy to see that $L(\mathbb{R}, \mu) = L[\mathrm{HOD}^N, j^*](\mathbb{R}) = L[A](\mathbb{R})$. Now since

$$V_\Theta^{L[\mathrm{HOD^N}, \mathrm{j^*}]} = V_\Theta^{\mathrm{HOD^N}} = V_\Theta^{\mathrm{HOD}},$$

there is a Θ-c.c. forcing $\mathbb{P}$ ($\mathbb{P}$ is a variation of the Vopenka algebra) such that we have $L[A] \subseteq \mathrm{HOD} \subseteq L[A](\mathbb{R}) = L(\mathbb{R}, \mu)$ and $L(\mathbb{R}, \mu)$ is the symmetric part of $L[A][g]$ where $g \subseteq \mathbb{P} \in L[A]$ is generic over HOD (such a g exists). This implies HOD$= L[A]$ hence completes our proof of the theorem. $\dashv$

We further assume μ comes from the club filter in V, $\mathcal{M}^\sharp_{\omega^2}$ exists and has unique $(\omega, \omega_1, \omega_1 + 1)$ iteration strategy in all generic extensions of V.[17] We'll show how to get rid of these assumptions later on. We first show how to iterate $\mathcal{M}_{\omega^2}$ to realize μ as the tail filter.

LEMMA 3.2. *There is an iterate $\mathcal{N}$ of $\mathcal{M}_{\omega^2}$ such that letting λ be the limit of $\mathcal{N}$'s Woodin cardinals, $\mathbb{R}$ can be realized as the symmetric reals over $\mathcal{N}$ at λ and letting $\mathcal{F}$ be the tail filter over $\mathcal{N}$ at λ, $L(\mathbb{R}, \mu) = L(\mathbb{R}, \mathcal{F})$.*

PROOF. Let δ_i be the sup of the first ωi Woodin cardinals of $\mathcal{M}_{\omega^2}$ and $\gamma = \sup_i \delta_i$. Let $\xi \geq \omega_1$ be such that $H(\xi) \vDash \mathsf{ZFC}^-$. In $V^{Col(\omega, H(\xi))}$, let $\langle X_i \mid i < \omega \rangle$ be an increasing and cofinal chain of countable (in V) elementary substructures of $H(\xi)$ and $\sigma_i = \mathbb{R} \cap X_i$. To construct the $\mathcal{N}$ as in the statement of the lemma, we do an $\mathbb{R}$-genericity iteration (in $V^{Col(\omega, H(\xi))}$) as follows. Let $\mathcal{P}_0 = \mathcal{M}^\sharp_{\omega^2}$ and assume $\mathcal{P}_0 \in X_0$. For $i > 0$, let $\mathcal{P}_i$ be the result of iterating $\mathcal{P}_{i-1}$ in X_{i-1} in the window between the $\omega(i-1)^{th}$ and ωi^{th} Woodin cardinals of $\mathcal{P}_{i-1}$ to make σ_{i-1} generic. We can make sure that each finite stage of the iteration is in X_{i-1}. Let $\mathcal{P}_\omega$ be obtained from the direct limit of the $\mathcal{P}_i$'s and iterating the top extender out of the universe. Let λ be the limit of Woodin cardinals in $\mathcal{P}_\omega$. It's clear that there is a $G \subseteq Col(\omega, < \lambda)$ generic over $\mathcal{P}_\omega$ such that $\mathbb{R} =_{def} \mathbb{R}^V$ is the symmetric reals over $\mathcal{P}_\omega$ and $L(\mathbb{R}, \mu)$ is in $\mathcal{P}_\omega[G]$. Let $\mathcal{F}$ be the tail filter on $\mathcal{P}_{\omega_1}(\mathbb{R})$ defined over $\mathcal{P}_\omega[G]$. By Section 2, $L(\mathbb{R}, \mathcal{F}) \vDash \mathcal{F}$ is a normal fine measure on $\mathcal{P}_{\omega_1}(\mathbb{R})$.

[17] In fact, it's enough to assume $\mathcal{M}^\sharp_{\omega^2}$ to be iterable in $V^{Col(\omega, \mathcal{P}(\mathbb{R}))}$.

https://doi.org/10.1017/jsl.2014.65 Published online by Cambridge University Press

We want to show $L(\mathbb{R},\mu)=L(\mathbb{R},\mathcal{F})$. To show this, it's enough to see that if $A\subseteq \mathcal{P}_{\omega_1}(\mathbb{R})$ is in $L(\mathbb{R},\mu)$ and A is a club (i.e., $A\in\mu$) then $A\in\mathcal{F}$. Let $\pi:\mathbb{R}^{<\omega}\to\mathbb{R}\in V$ witness that A is a club. By the choice of the X_i's, there is an n such that for all $m\geq n$, $\pi\in X_m$ and hence $\pi''\sigma_m^{<\omega}\subseteq\sigma_m$. This shows $A\in\mathcal{F}$. This in turns implies $L(\mathbb{R},\mu)=L(\mathbb{R},\mathcal{F})$ and $\mathcal{F}\cap L(\mathbb{R},\mathcal{F})=\mu\cap L(\mathbb{R},\mu)$. ⊣

We fix some notation. For a nondropping iterate $\mathcal{P}$ of $\mathcal{M}_{\omega^2}$, let $\gamma_i^{\mathcal{P}}$ be the supremum of the first $\omega(i+1)$ Woodin cardinals of $\mathcal{P}$ and $\lambda^{\mathcal{P}}=\sup_{i<\omega}\gamma_i$. From this point on to the end of the section, we assume the reader has in hands a copy of [13]. Our construction follows closely that paper. There's no point in rewritting every detail there.

Let $\mathcal{M}_\infty^+$ be the direct limit of all nondropping iterates (via countable stacks of countable normal trees) $\mathcal{P}$ of $\mathcal{M}_{\omega^2}$ below the first Woodin cardinal and $\mathcal{H}^+$ be the corresponding direct limit system. By definition, $\mathcal{H}^+$ is countably directed and hence $\mathcal{M}_\infty^+$ is well-founded. We'll define a direct limit system $\mathcal{H}$ in $L(\mathbb{R},\mu)$ that approximates $\mathcal{H}^+$. Working in $L(\mathbb{R},\mu)$, we say $\mathcal{P}$ is *suitable* if

1. $\mathcal{P}$ has only one Woodin cardinal $\delta^{\mathcal{P}}$;
2. it is full (with respect to mice), that is for all $\xi<o(\mathcal{P})$ such that ξ is a cutpoint of $\mathcal{P}$, $Lp(\mathcal{P}|\xi)\lhd\mathcal{P}$ and for all $\xi\neq\delta^{\mathcal{P}}$, $Lp(\mathcal{P}|\xi)\models\xi$ is not Woodin and $Lp(\mathcal{P}|\xi)\in\mathcal{P}$;
3. $\mathcal{P}=Lp_\omega(\mathcal{P}|\delta^{\mathcal{P}})$.

The following definition comes from Definition 6.21 in [13].

Definition 3.3. Working in $L(\mathbb{R},\mu)$, we let $\mathcal{O}$ be the collection of all functions f such that f is an ordinal definable function with domain the set of all countable, suitable $\mathcal{P}$, and $\forall\mathcal{P}\in\mathrm{dom}(f)(f(\mathcal{P})\subseteq\delta^{\mathcal{P}})$.

Definition 3.4. Suppose $\vec{f}\in\mathcal{O}^{<\omega}$, $\mathcal{P}$ is suitable, and $\mathrm{dom}(\vec{f})=n$. Let

$$\gamma_{(\mathcal{P},\vec{f})}=\sup\{Hull^{\mathcal{P}}(\vec{f}(0)(\mathcal{P}),\dots,\vec{f}(n-1)(\mathcal{P}))\cap\delta^{\mathcal{P}}\},$$

and

$$H_{(\mathcal{P},\vec{f})}=Hull^{\mathcal{P}}(\gamma_{(\mathcal{P},\vec{f})}\cup\{\vec{f}(0)(\mathcal{P}),\dots,\vec{f}(n-1)(\mathcal{P})\}).$$

We refer to reader to Section 6.3 of [13] for the definitions of $\vec{f}$-iterability, strong $\vec{f}$-iterability. The only difference between our situation and the situation in [13] is that our notions of "suitable", "short", "maximal", "short tree iterable" etc. are relative to the pointclass $(\Sigma^2_1)^{L(\mathbb{R},\mu)}$ instead of $(\Sigma^2_1)^{L(\mathbb{R})}$ as in [13].

Now, let $(\mathcal{P},\vec{f})\in\mathcal{H}$ if $\mathcal{P}$ is strongly $\vec{f}$-iterable. The ordering on $\mathcal{H}$ is defined as follows:

$$(\mathcal{P},\vec{f})\leq_{\mathcal{H}}(\mathcal{Q},\vec{g})\Leftrightarrow\vec{f}\subseteq\vec{g}\wedge\mathcal{Q}\text{ is a psuedo-iterate of }\mathcal{P}.^{18}$$

Note that if $(\mathcal{P},\vec{f})\leq_{\mathcal{H}}(\mathcal{Q},\vec{g})$ then there is a natural embedding $\pi_{(\mathcal{P},\vec{f}),(\mathcal{Q},\vec{q})}:H_{\mathcal{P},\vec{f}}\to H_{\mathcal{Q},\vec{g}}$. We need to see that $\mathcal{H}\neq\emptyset$.

Lemma 3.5. *Let $\vec{f}\in\mathcal{O}^{<\omega}$. Then there is a $\mathcal{P}$ such that $(\mathcal{P},\vec{f})\in\mathcal{H}$.*

[18] See definition 6.20 of [13] for the definition of psuedo-iterate.

https://doi.org/10.1017/jsl.2014.65 Published online by Cambridge University Press

PROOF SKETCH. For simplicity, assume $\mathrm{dom}(\vec{f}) = 1$. The proof of this lemma is just like the proof of Theorem 6.29 in [13]. We only highlight the key changes that make that proof work here.

First let $v, \mathbb{P}$ be as in the proof of Lemma 2.6. Let a be a countable transitive self-wellordered set and x be a real that codes a. We need to modify the $\mathcal{Q}_a^x$ defined in the proof of Lemma 2.6. Fix a coding of relativized premice by reals and write $\mathcal{P}_z$ for the premouse coded by z. Then let

$$\mathcal{F}_a^x = \{\mathcal{P}_z \mid z \leq_T x \text{ and } \mathcal{P}_z \text{ is a suitable premouse over } a \text{ and } \mathcal{P}_z \text{ is short-tree iterable}\}.$$

Let

$$\mathcal{Q}_a^x = Lp(\mathcal{Q}_a^{x,-}),$$

where $\mathcal{Q}_a^{x,-}$ is the direct limit of the simultaneous comparison and $\{y \mid y \leq_T x\}$-genericity iteration of all $\mathcal{P} \in \mathcal{F}_a^x$. The definition of $\mathcal{Q}_a^x$ comes from Section 6.6 of [13]. As in the proof of Lemma 2.6, we have:

1. letting $\langle \vec{d}^i \mid i < \omega \rangle$ be the generic sequence for $\mathbb{P}$ and $\langle \mathcal{Q}_j^i \mid i, j < \omega \rangle$ be the sequence of models associated to $\langle \vec{d}^i \mid i < \omega \rangle$ as defined in the proof of Lemma 2.6, we have that the model $N = L[T^\infty, \mathcal{M}^{\langle \vec{d}^i \rangle_i}] \vDash$ "there are ω^2 Woodin cardinals", where $\mathcal{M}^{\langle \vec{d}^i \rangle_i} = L[\cup_i \cup_j \mathcal{Q}_j^i]$;
2. letting λ be the sup of the Woodin cardinals of N, there is a $G \subseteq Col(\omega, < \lambda)$, G is N-generic such that letting $\mathbb{R}_G^*$ be the symmetric reals of $N[G]$ and $\mathcal{F}$ be the tail filter defined over $N[G]$, then $L(\mathcal{R}_G^*, \mathcal{F}) = L(\mathbb{R}, \mu)$ and $\mathcal{F} \cap L(\mathbb{R}, \mu) = \mu$.

The second key point is that whenever $\mathcal{P} \in \mathcal{H}^+$, we can then iterate $\mathcal{P}$ to $\mathcal{Q}$ (above any Woodin cardinal of $\mathcal{P}$) so that $\mathbb{R}^V$ can be realized as the symmetric reals for some $G \subseteq Col(\omega, < \delta_{\omega^2}^{\mathcal{Q}})$ and $L(\mathbb{R}, \mu) = L(\mathbb{R}, \mathcal{F})$ and $\mu \cap L(\mathbb{R}, \mu) = \mathcal{F} \cap L(\mathbb{R}, \mu)$, where $\mathcal{F}$ is the tail filter defined over $\mathcal{Q}[G]$. This is proved in Lemma 3.2.

We leave it to the reader to check that the proof of Theorem 6.29 of [13] goes through for our situation. This completes our sketch. ⊣

REMARK. The lemma above obviously shows $\mathcal{H} \neq \emptyset$. Its proof also shows for any $\vec{f} \in \mathcal{O}^{<\omega}$ and any $(\mathcal{P}, \vec{g}) \in \mathcal{H}$, there is a $\vec{g}$-iterate $\mathcal{Q}$ of $\mathcal{P}$ such that $\mathcal{Q}$ is $(\vec{f} \cup \vec{g})$-strongly iterable.

Now we outline the proof that $\mathcal{M}_\infty^+ \subseteq \mathrm{HOD}^{L(\mathbb{R},\mu)}$. We follow the proof in Section 6.7 of [13]. Suppose $\mathcal{P}$ is suitable and $s \in [\mathrm{OR}]^{<\omega}$, let $\mathcal{L}_{\mathcal{P},s}$ be the language of set theory expanded by constant symbols c_x for each $x \in \mathcal{P}|\delta^{\mathcal{P}} \cup \{\mathcal{P}\}$ and d_x for each x in the range of s. Since s is finite, we can fix a coding of the syntax of $\mathcal{L}_{\mathcal{P},s}$ such that it is definable over $\mathcal{P}|\delta^{\mathcal{P}}$ and the map $x \mapsto c_x$ is definable over $\mathcal{P}|\delta^{\mathcal{P}}$. We continue to use $\mathbb{P}$ to denote the Prikry forcing in Lemma 2.6.

DEFINITION 3.6. Let $\mathcal{P}$ be suitable and $s = \{\alpha_1, \ldots, \alpha_n\}$. We set

$$T_s(\mathcal{P}) = \{\phi \in \mathcal{L}_{\mathcal{P},s} \mid \exists p \in \mathbb{P}\ (p = (\emptyset, X) \wedge p \Vdash (\mathcal{M}^{\vec{d}_{\dot{G}}}, \alpha_1, \ldots, \alpha_n, x)_{x \in \mathcal{P}|\delta^{\mathcal{P}}} \vDash \phi)\}.$$

In the above definition, $\mathcal{M}^{\vec{d}_{\dot{G}}}$ is the canonical name for the model $\mathcal{M}^{\langle \vec{d}^i \rangle_i}$ defined in Lemma 3.5 where $\langle \vec{d}^i \rangle_i$ is the Prikry sequence given by a generic $G \subseteq \mathbb{P}$. Note that $T_s(\mathcal{P})$ is a complete, consistent theory of $\mathcal{L}_{\mathcal{P},s}$ and if $s \subseteq t$, we can think of

https://doi.org/10.1017/jsl.2014.65 Published online by Cambridge University Press

$T_s(\mathcal{P})$ as a subtheory of $T_t(\mathcal{P})$ in a natural way (after appropriately identifying the constant symbols of one with those of the other). Furthermore, $T_s \in \mathcal{O}$ for any $s \in [\mathrm{OR}]^{<\omega}$.

Let $\mathcal{N}_\infty$ be the direct limit of $\mathcal{H}$ under maps $\pi_{(\mathcal{P},\vec{f}),(\mathcal{Q},\vec{g})}$ for $(\mathcal{P},\vec{f}) \leq_{\mathcal{H}} (\mathcal{Q},\vec{g})$. Let $\pi_{(\mathcal{P},\vec{f}),\infty} : H_{\mathcal{P},\vec{f}} \to \mathcal{N}_\infty$ be the direct limit map. For each $s \in [\mathrm{OR}]^{<\omega}$ and $\mathcal{P}$ which is strongly T_s-iterable, we let

$$T_s^* = \pi_{(\mathcal{P},T_s),\infty}(T_s(\mathcal{P})).$$

Again, $s \subseteq t$ implies $T_s^* \subseteq T_t^*$, so we let

$$T^* = \bigcup\{T_s^* \mid s \in [\mathrm{OR}]^{<\omega}\}.$$

We have that T^* is a complete, consistent, and Skolemized[19] theory of $\mathcal{L}$, where $\mathcal{L} = \bigcup\{\mathcal{L}_{\mathcal{N}_\infty,s} \mid s \in [\mathrm{OR}]^{<\omega}\}$. We note that T^* is definable in $L(\mathbb{R},\mu)$ because the map $s \mapsto T_s^*$ is definable in $L(\mathbb{R},\mu)$.

Let $\mathcal{A}$ be the unique pointwise definable $\mathcal{L}$-structure such that $\mathcal{A} \models T^*$. We show $\mathcal{A}$ is wellfounded and let $\mathcal{N}_\infty^+$ be the transitive collapse of $\mathcal{A}$, restricted to the language of premice.

Lemma 3.7. $\mathcal{N}_\infty^+ = \mathcal{M}_\infty^+$

Proof sketch. We sketch the proof which completely mirrors the proof of Lemma 6.51 in [13]. Let Σ be the iteration strategy of $\mathcal{M}_{\omega^2}$ and $\Sigma_{\mathcal{P}}$ be the tail of Σ for a Σ-iterate $\mathcal{P}$ of $\mathcal{M}_{\omega^2}$. We will also use $\langle \delta_\alpha^{\mathcal{P}} \mid \alpha < \omega^2\rangle$ to denote the Woodin cardinals of a Σ-iterate $\mathcal{P}$ of $\mathcal{M}_{\omega^2}$. We write $\mathcal{P}^- = \mathcal{P}|((\delta_0^{\mathcal{P}})^{+\omega})^{\mathcal{P}}$. Working in $V^{Col(\omega,\mathbb{R})}$, we define sequences $\langle \mathcal{N}_k \mid k < \omega\rangle$, $\langle \mathcal{N}_k^\omega \mid k < \omega\rangle$, $\langle j_{k,l} \mid k \leq l \leq \omega\rangle$, $\langle i_k \mid k < \omega\rangle$, $\langle G_k \mid k < \omega\rangle$, and $\langle j_{k,l}^\omega \mid k \leq l \leq \omega\rangle$ such that

(a) $\mathcal{N}_k \in \mathcal{H}^+$ for all k;
(b) for all k, $\mathcal{N}_{k+1}$ is a $\Sigma_{\mathcal{N}_k}$-iterate of $\mathcal{N}_k$ (below the first Woodin cardinal of $\mathcal{N}_k$) and the corresponding iteration map is $j_{k,k+1}$;
(c) the $\mathcal{N}_k$'s are cofinal in $\mathcal{H}^+$;
(d) $i_k : \mathcal{N}_k \to \mathcal{N}_k^\omega$ is an iteration map according to $\Sigma_{\mathcal{N}_k}$ with critical point $> \delta_0^{\mathcal{N}_k}$;
(e) G_k is generic over $\mathcal{N}_k^\omega$ for the symmetric collapse up to the sup of its Woodins and $\mathbb{R}^*_{G_k} = \mathbb{R}^V$;
(f) $\mathcal{N}_k^\omega = \mathcal{M}^{\langle \vec{d}^i\rangle_i}$ for some $\langle \vec{e}^i\rangle_i$ which is $\mathbb{P}$-generic over $L(\mathbb{R},\mu)$ such that $(\mathcal{N}_k^\omega)^-$ is coded by a real in $\vec{e}^0(0)$;
(g) $j_{k,k+1}^\omega : \mathcal{N}_k^\omega \to \mathcal{N}_{k+1}^\omega$ is the iteration map;
(h) for $k < l$, $j_{k,l}^\omega \circ i_k = i_l \circ j_{k,l}$, where $j_{k,l} : \mathcal{N}_k \to \mathcal{N}_l$ and $j_{k,l}^\omega : \mathcal{N}_k^\omega \to \mathcal{N}_l^\omega$ are natural maps;
(i) $j_{k,k+1}|\mathcal{N}_k^- = j_{k,k+1}^\omega|(\mathcal{N}_k^\omega)^-$;
(j) the direct limit $\mathcal{N}_\omega^\omega$ of the $\mathcal{N}_k^\omega$ under maps $j_{k,l}^\omega$'s embeds into a $\Sigma_{\mathcal{M}_\infty^+}$-iterate of $\mathcal{M}_\infty^+$;
(k) for each $s \in [\mathrm{OR}]^{<\omega}$, for all sufficiently large k,

$$\mathcal{N}_k^\omega \models \phi[x,s] \Leftrightarrow \exists p \in \mathbb{P}\ (p = (\emptyset, X) \wedge p \Vdash (\mathcal{M}^{\vec{d}_{\dot{G}}} \models \phi[x,s]),$$

for $x \in \mathcal{N}_k^\omega|\delta_0^{\mathcal{N}_k^\omega}$.

[19] This is because of the Prikry property of $\mathbb{P}$.

https://doi.org/10.1017/jsl.2014.65 Published online by Cambridge University Press

Everything except for (f) is as in the proof of Lemma 6.51 of [13]. To see (f), fix a $k < \omega$. We fix a Prikry sequence $\langle \vec{d}^i \rangle_i$ such that $(\mathcal{N}_k^\omega)^-$ is coded into $\vec{d}^0(0)$ and letting $\sigma_i = \{y \in \mathbb{R}^V \mid y \text{ is recursive in } \vec{d}^i(j) \text{ for some } j < \omega\}$, then for each i, σ_i is closed under the iteration strategy $\Sigma_{\mathcal{N}_k}$ (this can be done in V). We then (inductively) for all i, construct a sequence $\langle \vec{e}^i \mid i < \omega \rangle$ such that $\vec{e}^i$ is a Prikry generic subsequence of $\vec{d}^i$ such that $M^{\langle \vec{e}^i \rangle_i}$ is an iterate of $\mathcal{N}_k$ (see Lemma 6.49 of [13]). The sequence $\langle \vec{e}^i \rangle_i$ satisfies (f) for $\mathcal{N}_k^\omega$.

Having constructed the above objects, the proof of Lemma 6.51 in [13] adapts here to give an isomorphism between $\mathcal{A}$ (viewed as a structure for the language of premice) and $\mathcal{M}_\infty^+$. The isomorphism is the unique extension to all of $\mathcal{A}$ of the map σ, where $\sigma(c_x^{\mathcal{A}}) = x$ (for $x \in \mathcal{M}_\infty^+ | \delta_0^{\mathcal{M}_\infty^+}$) and $\sigma(d_\alpha^{\mathcal{A}}) = j_{k,\omega}^\omega(\alpha)$ for k large enough such that $j_{l,l+1}^\omega(\alpha) = \alpha$ for all $l \geq k$. This completes our sketch. ⊣

Now we continue with the sketch of the proof that $\mathrm{HOD}^{L(\mathbb{R},\mu)}$ is a strategy mouse in the presence of $\mathcal{M}_{\omega^2}^\sharp$. Let λ_∞ be the supremum of the Woodin cardinals of $\mathcal{M}_\infty^+$. Let $\mathbb{R}^*$ be the symmetric reals given by an M_∞^+ generic $G \subseteq Col(\omega, < \lambda_\infty)$ and $\mathcal{F}^*$ be the corresponding tail filter defined in $\mathcal{M}_\infty^+[G]$. Since $L(\mathbb{R}^*, \mathcal{F}^*) \equiv L(\mathbb{R},\mu)$, $L(\mathbb{R}^*, \mathcal{G}^*)$ has its own version of $\mathcal{H}$ and $\mathcal{N}_\infty^+$, so we let

$$\mathcal{H}^* = \mathcal{H}^{L(\mathbb{R}^*,\mathcal{F}^*)} \text{ and } (\mathcal{N}_\infty^+)^* = (\mathcal{N}_\infty^+)^{L(\mathbb{R}^*,\mathcal{F}^*)}.$$

Let Λ be the restriction of $\Sigma_{\mathcal{M}_\infty^+}$ to stacks $\vec{\mathcal{T}} \in \mathcal{M}_\infty^+ | \lambda_\infty$, where

- $\vec{\mathcal{T}}$ is based on $\mathcal{M}_\infty^+ | \delta_0^{\mathcal{M}_\infty^+}$;
- $L(\mathbb{R}^*, \mathcal{F}^*) \vDash \vec{\mathcal{T}}$ is a finite full stack.[20]

We show $L[\mathcal{M}_\infty^+, \Lambda] = \mathrm{HOD}^{L(\mathbb{R},\mu)}$ through a sequence of lemmas. For an ordinal α, put

$$\alpha^* = d_\alpha^{\mathcal{A}},$$

and for $s = \{\alpha_1, \ldots, \alpha_n\}$ a finite set of ordinals, put

$$s^* = \{\alpha_1^*, \ldots, \alpha_n^*\}.$$

Lemma 3.8 (Derived model resemblance). *Let $(\mathcal{P}, \vec{f}) \in \mathcal{H}$ and $\overline{\eta} < \gamma_{(\mathcal{P},\vec{f})}$, and $\eta = \pi_{(\mathcal{P},\vec{f}),\infty}(\overline{\eta})$. Let $s \in [\mathrm{OR}]^{<\omega}$, and $\phi(v_0, v_1, v_2)$ be a formula in the language of set theory; then the following are equivalent*

(a) $L(\mathbb{R}^*, \mathcal{F}^*) \vDash \phi[\mathcal{M}_\infty, \eta, s^*]$;
(b) $L(\mathbb{R},\mu) \vDash$ *"there is an $(\mathcal{R}, \vec{f}) \geq_{\mathcal{F}} (\mathcal{P}, \vec{f})$ such that whenever $(\mathcal{Q}, \vec{f}) \geq_{\mathcal{H}} (\mathcal{R}, \vec{f})$, then $\phi(\mathcal{Q}, \pi_{(\mathcal{P},\vec{f}),(\mathcal{Q},\vec{f})}(\overline{\eta}), s)$"*.

The proof of this lemma is almost exactly like the proof of Lemma 6.54 of [13], so we omit it. The only difference is in Lemma 6.54 of [13], the proof of Lemma 6.51 of [13] is used, here we use that of Lemma 3.7.

Lemma 3.9. *Λ is definable over $L(\mathbb{R},\mu)$, and hence $L[\mathcal{M}_\infty^+, \Lambda] \subseteq \mathrm{HOD}^{L(\mathbb{R},\mu)}$*

Proof. Suppose $f \in \mathcal{O}$ is definable in $L(\mathbb{R},\mu)$ by a formula ψ and $s \in [\mathrm{OR}]^{<\omega}$, then we let $f^* \in \mathcal{O}^{L(\mathbb{R}^*,\mathcal{F}^*)}$ be definable in $L(\mathbb{R}^*, \mathcal{F}^*)$ from ψ and s^*.

[20] See Definition 6.20 of [13] for the precise definition of finite full stacks.

https://doi.org/10.1017/jsl.2014.65 Published online by Cambridge University Press

SUBLEMMA 3.10. *Let $\vec{\mathcal{T}}$ be a finite full stack on $\mathcal{M}_\infty^+|\delta_0^{\mathcal{M}_\infty^+}$ in $L(\mathbb{R}^*, \mathcal{F}^*)$ and let $\vec{b} = \Sigma_{\mathcal{M}_\infty^+}(\vec{\mathcal{T}})$. Then $\vec{b}$ respects f^*, for all $f \in \mathcal{O}$.*

The proof of Sublemma 3.10 is just like that of Claim 6.57 in [13] (with appropriate use of the proof of Lemma 3.7). Sublemma 3.10 implies $\mathcal{M}_\infty$ that is strongly f^*-iterable in $L(\mathbb{R}^*, \mathcal{F}^*)$ for all $f \in \mathcal{O}$. Sublemma 3.10 also gives the following.

SUBLEMMA 3.11. *Suppose $\mathcal{Q}$ is a psuedo-iterate*[21] *of $\mathcal{M}_\infty$ and $\mathcal{T}$ is a maximal tree on $\mathcal{Q}$ in the sense of $L(\mathbb{R}^*, \mathcal{F}^*)$. Let $b = \Lambda(\mathcal{T})$; then for all $\eta < \delta^{\mathcal{Q}}$, the following are equivalent*:

(a) $i_b^{\mathcal{T}}(\eta) = \xi$;
(b) *there is some $f \in \mathcal{O}$ such that $\eta < \gamma_{(\mathcal{Q}, f^*)}$ and exists some branch choice*[22] *of $\mathcal{T}$ that respects f^* and $i_c^{\mathcal{T}}(\eta) = \xi$.*

Since the $\gamma_{(\mathcal{Q}, f^*)}$'s sup up to $\delta^{\mathcal{Q}}$ and i_b is continuous at $\delta^{\mathcal{Q}}$, clause (b) defines Λ over $L(\mathbb{R}, \mu)$. ⊣

We have an iteration map

$$\pi_\infty : \mathcal{N}_\infty \to \mathcal{N}_\infty^*$$

which is definable over $L(\mathbb{R}^*, \mathcal{F}^*)$ by the equality

$$\pi_\infty = \cup_{f \in \mathcal{O}} \pi^{\mathcal{H}^*}_{(\mathcal{N}_\infty, f^*), \infty}.$$

By Boolean comparison, π_∞ is definable over $L[\mathcal{M}_\infty^+, \Lambda]$. This implies $\mathcal{N}_\infty^*$ is the direct limit of all Λ-iterates of $\mathcal{N}_\infty$ which belong to $\mathcal{M}_\infty^+$ and π_∞ is the canonical map into the direct limit. Lemma 3.8 also gives us the following.

LEMMA 3.12. *For all $\eta < \delta_0^{\mathcal{M}_\infty^+}$, $\pi_\infty(\eta) = \eta^*$.*

Finally, we have

THEOREM 3.13. *Suppose $\mathcal{M}_{\omega^2}^\sharp$ exists and is $(\omega, \mathrm{OR}, \mathrm{OR})$-iterable. Suppose μ is the club filter on $\mathcal{P}_{\omega_1}(\mathbb{R})$ and $L(\mathbb{R}, \mu) \vDash \mathsf{AD}^+ + \mu$ is a normal fine measure on $\mathcal{P}_{\omega_1}(\mathbb{R})$. Then the following models are equal*:

1. $\mathrm{HOD}^{L(\mathbb{R}, \mu)}$,
2. $L[\mathcal{M}_\infty^+, \pi_\infty]$,
3. $L[\mathcal{M}_\infty^+, \Lambda]$.

PROOF. Since $\pi_\infty \in L[\mathcal{M}_\infty^+, \Lambda]$, $L[\mathcal{M}_\infty^+, \pi_\infty] \subseteq L[\mathcal{M}_\infty^+, \Lambda]$. Lemma 3.9 implies $L[\mathcal{M}_\infty^+, \Lambda] \subseteq \mathrm{HOD}^{L(\mathbb{R}, \mu)}$. It remains to show $\mathrm{HOD}^{L(\mathbb{R}, \mu)} \subseteq L[\mathcal{M}_\infty, \pi_\infty]$. By Theorem 3.1, in $L(\mathbb{R}, \mu)$, there is some $A \subseteq \Theta$ such that $\mathrm{HOD} = L[A]$. Let ϕ define A. By Lemma 3.8

$$\alpha \in A \Leftrightarrow L[\mathcal{M}_\infty^+, \pi_\infty] \vDash \mathcal{M}_\infty^+ \vDash (1 \Vdash L(\mathbb{R}^*, \mathcal{F}^*) \vDash \phi[\alpha^*]).$$

By Lemma 3.12, $\alpha^* = \pi_\infty(\alpha)$ and hence the above equivalence defines A over $L[\mathcal{M}_\infty^+, \pi_\infty]$. This completes the proof of the theorem. ⊣

We now describe how to compute HOD just assuming $V = L(\mathbb{R}, \mu)$ satisfying AD^+. Let $\mathcal{H}$ be as above. The idea is that we use Σ_1 reflection to reflect a "bad" statement φ (like "$\mathcal{N}_\infty^+$ is illfounded" or "$\mathrm{HOD} \neq L(\mathcal{N}_\infty^+, \Lambda)$") to a level $L_\kappa(\mathbb{R}, \mu)$

[21] See Definition 6.13 of [13].
[22] See Definition 6.23 of [13].

https://doi.org/10.1017/jsl.2014.65 Published online by Cambridge University Press

where $\kappa < \underset{\sim}{\delta}^2_1$ (i.e., we have that $L_\kappa(\mathbb{R},\mu) \vDash \varphi$). But then since $\mu \cap L_\kappa(\mathbb{R},\mu)$ comes from the club filter, all we need to compute HOD in $L_\kappa(\mathbb{R},\mu)$ is to construct a mouse $\mathcal{N}$ related to N just like $M^\sharp_{\omega^2}$ related to $L(\mathbb{R},\mu)$. Once the mouse $\mathcal{N}$ is constructed, we successfully compute HOD of $L_\kappa(\mathbb{R},\mu)$ and hence show that $L_\kappa(\mathbb{R},\mu) \vDash \neg\varphi$. This gives us a contradiction.

We now proceed to construct $\mathcal{N}$. To be concrete, we fix a "bad" statement φ (like "HOD is illfounded") and let $N = L_\kappa(\mathbb{R},\mu)$ be least such that $N \vDash (T)$ where $(T) \equiv$ "$\mathsf{MC} + \mathsf{AD}^+ + \mathsf{DC} + \mathsf{ZF}^- + \Theta = \theta_0 + \varphi$". Let $\Gamma^* = (\Sigma^2_1)^N$, $\Phi = \mathcal{P}(\mathbb{R})^N$ and U be the universal Φ-set. We have that Γ^* is a good pointclass and $Env(\underset{\sim}{\Gamma^*}) = \Phi$ by closure of N. Let $\vec{B} = \langle B_i \mid i < \omega\rangle$ be a sjs sealing $Env(\underset{\sim}{\Gamma^*})$ with each $B_i \in N$ and $B_0 = U$. Such a $\vec{B}$ exists (see Section 4.1 of [18]).

Because MC holds and $\Phi \subsetneq \underset{\sim}{\delta}^2_1$, there is a real x such that there is a sound mouse $\mathcal{M}$ over x such that $\rho(\mathcal{M}) = x$ and $\mathcal{M}$ doesn't have an iteration strategy in N. Fix then such an $(x,\mathcal{M})$ and let Σ be the strategy of $\mathcal{M}$. Let $\Gamma \subsetneq \underset{\sim}{\Delta}^2_1$ be a good pointclass such that $Code(\Sigma), \vec{B}, U, U^c \in \underset{\sim}{\delta}_\Gamma$. By Theorem 10.3 in [10], there is a z such that $(\mathcal{N}^*_z, \delta_z, \Sigma_z)$ Suslin captures $Code(\Sigma), \vec{B}, U, U^c$ and $\mathcal{N}^*_z$ is coarse mouse with iteration strategy $\Sigma_z \in \underset{\sim}{\delta}^2_1$ and δ_z is the unique Woodin cardinal of $\mathcal{N}^*_z$.

Because $\vec{B}$ is Suslin captured by $\mathcal{N}^*_z$, we have $(\delta^+_z)^{\mathcal{N}^*_z}$-complementing trees $T, S \in \mathcal{N}^{*}_z$[23] with the property that for any Σ_z-iterate N^* of $\mathcal{N}^*_z$ such that the iteration map $i : \mathcal{N}^*_z \to N^*$ exists, for any $<$-$i((\delta^+_z)^{\mathcal{N}^*_z})$-generic g over N^*, $p[i(T)] \cap N^*[g] = \vec{B} \cap N^*[g] = \mathbb{R}^{N^*[g]} \backslash p[i(S)]$. Let κ be the least cardinal of $\mathcal{N}^*_z$ which, in $\mathcal{N}^*_z$ is $< \delta_z$-strong.

Claim 1. $\mathcal{N}^*_z \vDash$ "*κ is a limit of points η such that $Lp^{\Gamma^*}(\mathcal{N}^*_z|\eta) \vDash$ "η is Woodin"*.

Proof. The proof is an easy reflection argument. Let $\lambda = \delta^+_z$ and let $\pi : M \to \mathcal{N}^*_z|\lambda$ be an elementary substructure such that

1. $T, S \in ran(\pi)$,
2. if $\mathrm{cp}(\pi) = \eta$ then $V_\eta^{\mathcal{N}^*_z} \subseteq M$, $\pi(\eta) = \delta_z$ and $\eta > \kappa$.

By elementarity, we have that $M \vDash$ "η is Woodin". Letting $\pi^{-1}(\langle T, S\rangle) = \langle \bar{T}, \bar{S}\rangle$, we have that $(\bar{T}, \bar{S})$ Suslin captures $\vec{B}$ over M at η. This implies that M is Φ-full and in particular, $Lp^{\Gamma^*}(\mathcal{N}^*_z|\eta) \in M$. Therefore, $Lp^{\Gamma^*}(\mathcal{N}^*_z|\eta) \vDash$ "η is Woodin". The claim then follows by a standard argument. ⊣

Let now $\langle \eta_i : i < \omega^2\rangle$ be the first ω^2 points $< \kappa$ such that for every $i < \omega$, $Lp^{\Gamma^*}(\mathcal{N}^*_z|\eta_i) \vDash$ "η_i is Woodin". Let now $\langle \mathcal{N}_i : i < \omega^2\rangle$ be a sequence constructed according to the following rules:

1. $\mathcal{N}_0 = L[\vec{E}]^{\mathcal{N}^*_z|\eta_0}$,
2. if i is limit, $\mathcal{N}'_i = \cup_{j<i}\mathcal{N}_i$ and $\mathcal{N}_i = (L[\vec{E}][\mathcal{N}'_i])^{\mathcal{N}^*_z|\eta_i}$,
3. $\mathcal{N}_{i+1} = (L[\vec{E}][\mathcal{N}_i])^{\mathcal{N}^*_z|\eta_{i+1}}$.

Let $\mathcal{N}_{\omega^2} = \cup_{i<\omega^2}\mathcal{N}_i$.

Claim 2. *For every $i < \omega^2$, $\mathcal{N}_{\omega^2} \vDash$ "η_i is Woodin" and $\mathcal{N}_{\omega^2}|(\eta^+_i)^{\mathcal{N}_\omega} = Lp^{\Gamma^*}(\mathcal{N}_i)$.*

[23] This means that whenever g is $< (\delta^+_z)^{\mathcal{N}^*_z}$-generic over $\mathcal{N}^*_z$, then in $\mathcal{N}^*_z[g]$, $p[T]$ and $p[S]$ project to complements.

https://doi.org/10.1017/jsl.2014.65 Published online by Cambridge University Press

PROOF. It is enough to show that

1. $\mathcal{N}_{i+1} \vDash$ "η_i is Woodin",
2. $\mathcal{N}_i = V_{\eta_i}^{\mathcal{N}_{i+1}}$,
3. $\mathcal{N}_{i+1}|(\eta_i^+)^{\mathcal{N}_{i+1}} = Lp^{\Gamma^*}(\mathcal{N}_i)$,
4. if i is limit, then $\mathcal{N}_i|((\sup_{j<i}\eta_j^+)^{\mathcal{N}_i}) = Lp^{\Gamma^*}(\mathcal{N}_i')$.

To show 1–4, it is enough to show that if $\mathcal{W} \trianglelefteq \mathcal{N}_{i+1}$ is such that $\rho_\omega(W) \leq \eta_i$ or if i is limit and $\mathcal{W} \triangleleft \mathcal{N}_i$ is such that $\rho_\omega(W) \leq \sup_{j<i}\eta_j$ then the fragment of $\mathcal{W}$'s iteration strategy which acts on trees above η_i ($\sup_{j<i}\eta_j$ respectively) is in Γ^*. Suppose first that i is a successor and $\mathcal{W} \trianglelefteq \mathcal{N}_{i+1}$ is such that $\rho_\omega(W) \leq \eta_i$. Let ξ be such that if $\mathcal{S}$ is the ξth model of the full background construction producing $\mathcal{N}_{i+1}$ then $\mathbb{C}(\mathcal{S})$[24] $= \mathcal{W}$. Let $\pi : \mathcal{W} \to \mathcal{S}$ be the core map. The iteration strategy of $\mathcal{W}$ is the π-pullback of the iteration strategy of $\mathcal{S}$. Let then $\nu < \eta_{i+1}$ be such that $\mathcal{S}$ is the ξth model of the full background construction of $\mathcal{N}_x^*|\nu$. To determine the complexity of the induced strategy of $\mathcal{S}$ it is enough to determine the strategy of $\mathcal{N}_x^*|\nu$ which acts on nondropping stacks that are completely above η_i. Now, notice that by the choice of η_{i+1}, for any nondropping tree $\mathcal{T}$ on $\mathcal{N}_x^*|\nu$ which is above η_i and is of limit length, if $b = \Sigma(\mathcal{T})$ then $\mathcal{Q}(b,\mathcal{T})$ exists and $\mathcal{Q}(b,\mathcal{T})$ has no overlaps, and $\mathcal{Q}(b,\mathcal{T}) \trianglelefteq Lp^{\Gamma^*}(\mathcal{M}(\mathcal{T}))$. This observation and the fact that Γ^* is closed under real quantifiers indeed show that the fragment of the iteration strategy of $\mathcal{N}_x^*|\nu$ that acts on nondropping stack that are above η_i is in Γ^*. Hence, the strategy of $\mathcal{W}$ is in Γ^*.

Suppose $i < \omega^2$ is limit and (1)–(4) are satisfied for all $j < i$. We first claim that the induced strategy $\Sigma_{\mathcal{N}_i'}$ from Σ_z is Γ^*-fullness preserving: suppose $k : \mathcal{N}_i' \to \mathcal{P}$ is according to $\Sigma_{\mathcal{N}_i'}$ then $\mathcal{P}$ is Γ^*-i-suitable, that is

- $\langle k(\eta_j) \mid j < i\rangle$ are the only Woodin cardinals of $\mathcal{P}$;
- for any cut point ξ of $\mathcal{P}$, $Lp^{\Gamma^*}(\mathcal{P}|\xi) \triangleleft \mathcal{P}$ and for any $\xi \neq i(\eta_j)$ for any $j < i$, $Lp^{\Gamma^*}(\mathcal{P}|\xi) \vDash \xi$ is not Woodin.

First we see that $\mathcal{N}_i'$ is Γ^*-i-suitable. We show for instance if $\eta < \eta_0$ then $C_{\Gamma^*}(\mathcal{N}_i'|\eta) \vDash$ "η is not Woodin" (the rest of the verification is similar). Otherwise, $\mathcal{N}_i'|\eta$ is the η-th model in the $L[E]$-construction of $\mathcal{N}_z^*$ and $L[T_{\Gamma^*}, \mathcal{N}_i'|\eta] \vDash$ "η is Woodin", where T_{Γ^*} is the tree projecting to the Γ^*-universal set. We also get that $L[T_{\Gamma^*}, \mathcal{N}_i'|\eta] \cap V_\eta = \mathcal{N}_i'|\eta$ and $V_\eta^{\mathcal{N}_z^*}$ is generic over $L[T_{\Gamma^*}, \mathcal{N}_i'|\eta]$ for $\mathbb{B}_\eta$, the η-generic extender algebra at η. $\mathbb{B}_\eta$ is η-cc, so every $f : \eta \to \eta$ in $L[T_{\Gamma^*}, \mathcal{N}_i'|\eta][V_\eta^{\mathcal{N}_z^*}]$ is bounded by a function $g : \eta \to \eta$ in $L[T_{\Gamma^*}, \mathcal{N}_i'|\eta]$. Furthermore, if E witnesses the Woodin property for g in $L[T_{\Gamma^*}, \mathcal{N}_i'|\eta]$ and $\nu(E)$ is a cardinal in $L[T_{\Gamma^*}, \mathcal{N}_i'|\eta]$ then the background extender E^* witnesses the Woodin property for f in $L[T_{\Gamma^*}, \mathcal{N}_i'|\eta][V_\eta^{\mathcal{N}_z^*}]$ (note also $E \restriction \nu(E) = E^* \restriction \nu(E)$). So η is Woodin in $L[T_{\Gamma^*}, V_\eta^{\mathcal{N}_z^*}]$. By the minimality of η_0, $\eta = \eta_0$. Contradiction. The proof works also for any η_j.

Now let k be as in the claim. Let $k^* : \mathcal{N}_z^* \to N^*$ be the map coming from resurrecting the tree giving rise to k. Let $\sigma : \mathcal{P} \to k^*(\mathcal{N}_i')$ be the resurrection map. Since $\mathcal{N}_z^*, N^*$ have absolute definitions of Γ^*, $k^*(\mathcal{N}_i')$ is Γ^*-i-suitable. This and the fact that σ has in its range all the term relations for $\vec{B}$, we get that $\mathcal{P}$ is Γ^*-i-suitable.

[24] $\mathbb{C}(\mathcal{S})$ denotes the core of $\mathcal{S}$.

https://doi.org/10.1017/jsl.2014.65 Published online by Cambridge University Press

The argument in Lemma 3.7 that an iterate of $\mathcal{M}_{\omega^2}$ extends a Prikry generic and the fact that $\Sigma_{\mathcal{N}_i'}$ is Γ^*-fullness preserving show that $\mathcal{W}$ cannot project across $\sup_{j<i}\eta_j$ and that $\mathcal{W} \lhd Lp^{\Gamma^*}(\mathcal{N}_i')$. This completes the proof of the claim. ⊣

Working in $L(\mathbb{R},\mu)$, we now claim that there is $\mathcal{W} \trianglelefteq Lp(\mathcal{N}_{\omega^2})$ such that $\rho(W) < \eta_{\omega^2}$. To see this suppose not. It follows from MC that $Lp(\mathcal{N}_{\omega^2})$ is Σ^2_1-full. We then have that x is generic over $Lp(\mathcal{N}_{\omega^2})$ at the extender algebra of $\mathcal{N}_{\omega^2}$ at η_0. Because $Lp(\mathcal{N}_{\omega^2})[x]$ is Σ^2_1-full, we have that $\mathcal{M} \in Lp(\mathcal{N}_{\omega^2})[x]$ and $Lp(\mathcal{N}_{\omega^2})[x] \vDash$ "$\mathcal{M}$ is η_{ω^2}-iterable" by fullness of $Lp(\mathcal{N}_{\omega^2})[x]$. Let $\mathcal{S} = (L[\vec{E}][x])^{\mathcal{N}_{\omega^2}[x]|\eta_2}$ where the extenders used have critical point $> \eta_0$. Then working in $\mathcal{N}_{\omega^2}[x]$ we can compare $\mathcal{M}$ with $\mathcal{S}$. Using standard arguments, we get that $\mathcal{S}$ side doesn't move and by universality, $\mathcal{M}$ side has to come short (see [6]). This in fact means that $\mathcal{M} \trianglelefteq \mathcal{S}$. But the same argument used in the proof of Claim 2 shows that every $\mathcal{K} \trianglelefteq \mathcal{S}$ has an iteration strategy in Γ^*, contradiction!

Let $\eta_{\omega^2} = \sup_{i<\omega^2}\eta_i$ and $\mathcal{W} \trianglelefteq Lp(\mathcal{N}_{\omega^2})$ be least such that $\rho_\omega(\mathcal{W}) < \eta_{\omega^2}$.We can show the following.

LEMMA 3.14. *$\mathcal{W} = \mathcal{J}_{\xi+1}(\mathcal{N}_{\omega^2})$ where ξ is least such that for some τ, $\mathcal{J}_\xi(\mathcal{N}_{\omega^2}) \vDash$ "$\mathsf{ZF}^- + \tau$ is a limit of Woodin cardinals $+(T)$ holds in my derived model below τ".*[25]

Since the proof of this lemma is almost the same as that of Claim 7.5 in [13], we will not give it here. However, we have a few remarks regarding the proof:

- we typically replace N by a countable transitive $\overline{N}$ elementarily embeddable into N since the strategy of $\mathcal{W}$ is not known to extend to $V^{Col(\omega,\mathbb{R})}$. Having said this, we will confuse our N with its countable copy.
- We can then do an $\mathbb{R}^N$-iteration of $\mathcal{W}$ to "line up" its iterate with a $\mathbb{P}^N$-generic.

Asides from these remarks, everything else can just be transferred straightforwardly from the proof of Lemma 7.5 in [13] to the proof of Lemma 3.14. Now we just let $\mathcal{N}$ be the pointwise definable hull of $\mathcal{W}|\xi$. Letting $\mathcal{N}$'s unique iteration strategy be Λ, we can show Λ is Φ-fullness preserving and for any $\vec{f} \in (\mathcal{O}^{<\omega})^N$, there is a strongly $\vec{f}$-iterable, N-suitable $\mathcal{P}$ (in fact, $\mathcal{P} = \mathcal{Q}^-$ for some Λ-iterate $\mathcal{Q}$ of $\mathcal{N}$). We leave the rest of the details to the reader.

§4. Further applications. We first prove a series of lemmas which imply Theorem 1.3. For each $\sigma \in \mathcal{P}_{\omega_1}(\mathbb{R})$, let

$$M_\sigma = \mathrm{HOD}^{(\mathrm{L}(\mathbb{R},\mu),\mu)}_{\sigma\cup\{\sigma\}}.$$

Suppose G is a $\mathbb{P}_{\max}$ generic over $L(\mathbb{R},\mu)$, where

$$L(\mathbb{R},\mu) \vDash \text{"}\mathsf{AD}^+ + \mu \text{ is a normal fine measure on } \mathcal{P}_{\omega_1}(\mathbb{R})\text{"}.$$

Note that $L(\mathbb{R},\mu)[G] \vDash \mathsf{ZFC}$ since $\mathbb{P}_{\max}$ wellorders the reals. In $L(\mathbb{R},\mu)[G]$, let

$$\mathcal{I} = \{A \mid \exists\langle A_x \mid x \in \mathbb{R}\rangle(A \subseteq \nabla_{x\in\mathbb{R}}A_x \wedge \forall x\ (\mu(A_x) = 0 \text{ or } A_x = \neg S))\},$$

where $S = \{\sigma \in \mathcal{P}_{\omega_1}(\mathbb{R}) \mid G \cap \sigma \text{ is } \mathbb{P}_{\max} \restriction \sigma\text{-generic over } M_\sigma\}$. It's clear that in $L(\mathbb{R},\mu)[G]$, $\mathcal{I}$ is a normal fine ideal. Let $\mathcal{F}$ be the dual filter of $\mathcal{I}$.

[25] Here "derived model" means the model $L(\mathbb{R}^*,\mathcal{F}^*)$ where $\mathbb{R}^*$ is the symmetric reals for the Levy collapse at τ and $\mathcal{F}^*$ is the corresponding tail filter.

https://doi.org/10.1017/jsl.2014.65 Published online by Cambridge University Press

LEMMA 4.1. *Let* $\mathcal{I}^- = \{A \mid \exists\langle A_x \mid x \in \mathbb{R}\rangle(A \subseteq \nabla_{x\in\mathbb{R}} A_x \wedge \forall x\ \mu(A_x) = 0)\}$. *Let* $\mathcal{F}^-$ *be the dual filter of* $\mathcal{I}^-$. *Suppose* $A \in \mathcal{F}^-$. *Then* $\exists B, C$ *such that* $\mu(B) = 1$ *and* C *is a club in* $L(\mathbb{R}, \mu)[G]$ *such that* $B \cap C \subseteq A$.

PROOF. Suppose $1 \Vdash_{\mathbb{P}_{\max}} \tau : \mathbb{R} \to \mu$ witnesses $\{\sigma \mid \forall x \in \sigma\ \sigma \in \tau(x)\} \in \mathcal{F}^-$. For each $x \in \mathbb{R}$. let $D_x = \{p \mid p \parallel \tau(x)\}$. It's easy to see that D_x is dense for each x. Furthermore,

$$\forall^*_\mu \sigma \forall x \in \sigma(D_x \cap \sigma \text{ is dense in } \mathbb{P}_{\max} \restriction \sigma \wedge \{q \in D_x \cap \sigma \mid q \Vdash \sigma \in \tau(x)\} \text{ is dense.})$$

For otherwise, $\exists x, q \forall^*_\mu \sigma\ x \in \sigma \wedge q \in D_x \cap \sigma \wedge q \Vdash \sigma \notin \tau(x)$. This contradicts that $q \Vdash \tau(x) \in \mu$. Let B be the set of σ having the property displayed above. $\mu(B) = 1$.

Let $A \subseteq \mathbb{R}$ code the function $x \mapsto D_x$ and let G be a $\mathbb{P}_{\max}$-generic over $L(\mathbb{R}, \mu)$. Hence $D = \{\sigma \mid \forall x \in \sigma\ \sigma \in \tau_G(x)\} \in \mathcal{F}^-$. Let $C = \{\sigma \mid (\sigma, A \cap \sigma, G \cap \sigma) \prec (\mathbb{R}, A, G)\}$. Hence C is a club in $L(\mathbb{R}, \mu)[G]$ and $B \cap C \subseteq D$. ⊣

LEMMA 4.2. *Let* $\mathcal{I}^-, \mathcal{F}^-$ *be as in Lemma* 4.1. *Then* $S \notin \mathcal{I}^-$.

PROOF. Suppose not. Then $\neg S \in \mathcal{F}^-$. The following is a Σ_1-statement (with predicate μ) that $L(\mathbb{R}, \mu)[G]$ satisfies:

$$\exists B, C(\mu(B) = 1 \wedge C \text{ is a club } \wedge \forall \sigma(\sigma \in B \cap C \Rightarrow \exists D \subseteq \mathbb{P}_{\max}(M_\sigma \vDash \text{“}D \text{ is dense”} \wedge G \cap D = \emptyset))).$$

By part (1) of Theorem 1.2 and the fact that $\mathbb{P}_{\max}$ is a forcing of size $\mathbb{R}$, $L_{\underset{\sim}{\delta}^2_1}(\mathbb{R}, \mu)[G]$ satisfies the same statement. Here μ coincides with the club measure and hence $L_{\underset{\sim}{\delta}^2_1}(\mathbb{R}, \mu)[G] \vDash$ “$\neg S$ contains a club”. Let $\mathcal{C}$ be a club of elementary substructures X_σ containing everything relevant (and a pair of complementing trees for the universal Σ^2_1 set). Then it's easy to see that $\mathcal{C}^* \subseteq S$ where $\mathcal{C}^* = \{\sigma \mid \sigma = \mathbb{R} \cap X_\sigma \wedge X_\sigma \in \mathcal{C}\}$. This is a contradiction. ⊣

The above lemmas say that $\mathcal{I}$ strictly contains $\mathcal{I}^-$, i.e., S adds nontrivial information to $\mathcal{I}^-$. We now proceed to characterize $\mathcal{I}$-positive sets in terms of the $\mathbb{P}_{\max}$ forcing relation over $L(\mathbb{R}, \mu)$.

LEMMA 4.3. *Suppose* $p \in \mathbb{P}_{\max}$ *and* τ *is a* $\mathbb{P}_{\max}$ *term for a subset of* $\mathcal{P}_{\omega_1}(\mathbb{R})$ *in generic extensions of* $L(\mathbb{R}, \mu)$. *Then the following is true in* $L(\mathbb{R}, \mu)$.

$$p \Vdash_{\mathbb{P}_{\max}} \tau \textit{ is } \mathcal{I}\textit{-positive} \Leftrightarrow \forall^*_\mu \sigma\ \forall^* g \subseteq \mathbb{P}_{\max} \restriction \sigma\ (p \in g \Rightarrow \exists q < g\ q \Vdash_{\mathbb{P}_{\max}} \sigma \in \tau).$$

PROOF. Some explanations about the notation in the lemma are in order. “$\forall^* g \subseteq \mathbb{P}_{\max} \restriction \sigma$” means “for comeager many filters g over $\mathbb{P}_{\max} \restriction \sigma$”; “$\exists^* g \subseteq \mathbb{P}_{\max} \restriction \sigma$” means “for nonmeager many filters g over $\mathbb{P}_{\max} \restriction \sigma$”. These category quantifiers make sense because σ is countable. Also we only force with $\mathbb{P}_{\max}$ here so we'll write “$\Vdash$” for “$\Vdash_{\mathbb{P}_{\max}}$” and “$p < q$” for “$p <_{\mathbb{P}_{\max}} q$”. Finally, “$q < g$” means “$\forall r \in g\ q < r$”.

CLAIM. *Suppose in* $L(\mathbb{R}, \mu)$, $\forall \sigma\ X_\sigma$ *is comeager in* $\mathbb{P}_{\max} \restriction \sigma$. *Then* $\forall^*_\mu \sigma \forall G_\sigma (G_\sigma$ *is* $\mathbb{P}_{\max} \restriction \sigma$*-generic over* $M_\sigma \Rightarrow G_\sigma \in X_\sigma)$.

PROOF. Suppose $\sigma \mapsto X_\sigma$ is $OD_{\mu,x}$ for some $x \in \mathbb{R}$. Let $A = \{y \in \mathbb{R} \mid y$ codes (σ, g) where $g \in X_\sigma\}$. Hence A is $OD_{\mu,x}$. Let S be an $OD_{\mu,x}$ ∞-Borel code for A and $\mathcal{A}_S$ be the set of reals coded by S. Hence, $\forall^*_\mu \sigma\ S \in M_\sigma$.

https://doi.org/10.1017/jsl.2014.65 Published online by Cambridge University Press

For each such σ, let $G_\sigma \in X_\sigma$ be M_σ-generic and H be $M_\sigma[G_\sigma]$-generic for $Col(\omega, \sigma)$. Then

$$M_\sigma[G_\sigma][H] \vDash (\sigma, G_\sigma) \in \mathcal{A}_S.$$

In the above, note that we use $S \in M_\sigma$. Also no $p \in \mathbb{P}_{\max} \restriction \sigma$ can force $(\sigma, \dot{G}) \notin \mathcal{A}_S$. Hence we're done. ⊣

Suppose the conclusion of the lemma is false. There are two directions to take care of.

Case 1. $p \Vdash \tau$ is $\mathcal{I}$-positive but $\forall^*_\mu \sigma \exists^* g\ (p \in q \wedge \forall q < g\ q \Vdash \sigma \notin \tau)$.

Extending p if necessary and using normality, we may assume $\forall^*_\mu \sigma \forall^* g (p \in g \wedge \forall q < g\ q \vDash \sigma \notin \tau)$. Let T be the set of such σ. Let G be a $\mathbb{P}_{\max}$ generic and $p \in G$. By the claim and the fact that $S \in \mathcal{F}$, $\tau_G \cap S \cap T \neq \emptyset$. So let $\sigma \in \tau_G \cap S \cap T$ such that $p \in G \cap \sigma$. Then $G \cap \sigma$ is M_σ-generic and $\forall q < G \cap \sigma\ q \Vdash \sigma \notin \tau$. But $\exists q < G \cap \sigma$ such that $q \in G$ by density. This implies $\sigma \notin \tau_G$. Contradiction.

Case 2. $p \Vdash \tau \in \mathcal{I}$ and $\forall^*_\mu \sigma \forall^* g\ (p \in g \Rightarrow \exists q < g\ q \Vdash \sigma \in \tau)$.

Let T be the set of σ as above. Let G be $\mathbb{P}_{\max}$ generic containing p. Hence $T \in \mathcal{F}$. Let $\sigma \in T \cap S \cap \neg \tau_G$ and $p \in G \cap \sigma$. By density, $\exists q < G \cap \sigma\ q \in G \wedge q \Vdash \sigma \in \tau$. Hence $\sigma \in \tau_G$. Contradiction. ⊣

Now suppose $\dot{f}$ is a $\mathbb{P}_{\max}$ name for a function from an $\mathcal{I}$-positive set into OR and let τ be a name for $\mathrm{dom}(f)$ and for simplicity suppose $\emptyset \Vdash \tau$ is $\mathcal{I}$-positive $\wedge\ \dot{f} : \tau \to \check{\mathrm{OR}}$. Let $F : \mathcal{P}_{\omega_1}(\mathbb{R}) \to \mathrm{OR} \cup \{\infty\}$ be defined as follows:

$$F(\sigma) = \alpha_\sigma \text{ where } \alpha_\sigma \text{ is the least } \alpha \text{ such that}$$
$$\forall^* g \subseteq \mathbb{P}_{\max} \restriction \sigma\ (g \text{ is } M_\sigma\text{-generic} \Rightarrow \exists q < g\ q \Vdash \check{\sigma} \in \tau \wedge \dot{f}(\check{\sigma}) = \check{\alpha}),$$
$$\text{if } \alpha \text{ exists, and } \infty \text{ otherwise.}$$

Clearly, $F \in L(\mathbb{R}, \mu)$ and by the fact that τ_G is $\mathcal{I}$-positive and a standard application of Baire category theorem,[26] $\forall^*_\mu \sigma\ F(\sigma) \neq \infty$.

Lemma 4.4. *Suppose $\dot{f}, \tau, F$ are as above. Suppose G is a $\mathbb{P}_{\max}$ generic over $L(\mathbb{R}, \mu)$. Then in $L(\mathbb{R}, \mu)[G]$, $\{\sigma \mid F(\sigma) = f(\sigma)\}$ is $\mathcal{I}$-positive.*

Proof. Suppose not; assume $p \Vdash \tau' = \{\sigma \mid F(\check{\sigma}) = \dot{f}(\check{\sigma})\} \in \mathcal{I}$. Using Lemma 4.3, we get

$$\forall^*_\mu \sigma\ \exists^* g \subseteq \mathbb{P}_{\max} \restriction \sigma\ (p \in g \wedge \forall q < g\ q \Vdash \sigma \notin \tau'). \tag{4.1}$$

Using the Baire category theorem, we get from 4.1

$$\forall^*_\mu \sigma\ \exists p > q_\sigma \in \sigma\ \forall^* q_\sigma \in g \subseteq \mathbb{P}_{\max} \restriction \sigma\ \forall r < g\ r \Vdash \sigma \notin \tau'. \tag{4.2}$$

Now using normality of μ, we "freeze out" the q_σ's

$$\exists q < p\ \forall^*_\mu \sigma\ \forall^* q \in g \subseteq \mathbb{P}_{\max} \restriction \sigma\ \forall r < g\ r \Vdash \sigma \notin \tau'. \tag{4.3}$$

From 4.2 and 4.3, we get

$$\exists q < p\ \forall^*_\mu \sigma\ \forall^* q \in g \subseteq \mathbb{P}_{\max} \restriction \sigma\ (g \text{ is } M_\sigma\text{-generic } \Rightarrow \forall r < g\ r \Vdash F(\check{\sigma}) \neq \dot{f}(\check{\sigma})). \tag{4.4}$$

[26] More precisely, we use the fact that if $F : A \to \mathrm{OR}$ is a function on a comeager set A then F is constant on some comeager subset of A.

https://doi.org/10.1017/jsl.2014.65 Published online by Cambridge University Press

We get a contradiction from 4.4 as follows. Fix a σ in the μ-measure one set from 4.4 such that $F(\sigma) = \alpha_\sigma \neq \infty$. For the chosen σ, fix a g in the set described in 4.4 as well as in the set described in the definition of $F(\sigma)$. Then by 4.4, $\forall r < g$, $r \Vdash F(\sigma) = \alpha_\sigma \neq \dot{f}(\check{\sigma})$ but by the definition of $F(\sigma)$, $\exists r < g\ r \Vdash \dot{f}(\check{\sigma}) = \alpha_\sigma$. Contradiction. ⊣

Proof of Theorem 1.3. Working in $L(\mathbb{R}, \mu)[G]$, let $H \subseteq \mathcal{I}^+$ be generic. We show that (1)–(3) hold. Let $A \subseteq \mathbb{R}$ be OD_x for some $x \in \mathbb{R}$. By countable closure and homogeneity of $\mathbb{P}_{\max}$, $x \in L(\mathbb{R}, \mu)$ and hence $A \in L(\mathbb{R}, \mu)$. Since $\mathcal{F} \restriction L(\mathbb{R}, \mu) = \mu$, we obtain (1).[27] Lemma 4.4 implies $\forall s \in \mathrm{OR}^\omega\ j_H \restriction \mathrm{HOD}_s \in V$ and is independent of H. To see this, note that $s \in L(\mathbb{R}, \mu)$ as $\mathbb{P}_{\max}$ is countably closed and $L(\mathbb{R}, \mu) \vDash \mathrm{DC}$; furthermore, by homogeneity of $\mathbb{P}_{\max}$, $\mathrm{HOD}_s \subseteq \mathrm{HOD}_s^{L(\mathbb{R},\mu)}$ and there is a bijection between OR and HOD_s in $L(\mathbb{R}, \mu)$. So Lemma 4.4 applies to functions $f : S \to \mathrm{HOD}_s$ where S is $\mathcal{I}$-positive. This implies $j_H \restriction \mathrm{HOD}_s = j_\mu \restriction \mathrm{HOD}_s$, which also shows (2).

To show $j_H \restriction \mathrm{HOD}_{\mathcal{I}}$ is independent of H, first note that $\mathcal{F}$ is generated by μ and $\mathcal{A} =_{\mathrm{def}} \{T \subseteq \mathcal{P}_{\omega_1}(\mathbb{R}) \mid \exists C (C \text{ is a club and } T \cap C = S \cap C\}$, where S is defined at the beginning of the section in relation to the definition of $\mathcal{I}$. Note that $\mathcal{A}$ is definable in $L(\mathbb{R}, \mu)[G]$ (from no parameters). To see this, suppose G_0, G_1 are two $\mathbb{P}_{\max}$ generics (in $L(\mathbb{R}, \mu)[G]$) and let S_{G_i} be defined relative to G_i ($i \in \{0, 1\}$) the same way S is defined relative to G. Also let $A_{G_i} \subseteq \omega_1$ be the generating set for G_i. Let $p \in G_0 \cap G_1$ and $a_0, a_1 \in \mathcal{P}(\omega_1)^p$ be such that $j_i(a_i) = A_i$ where j_i are unique iteration maps of p. The proof of homogeneity of $\mathbb{P}_{\max}$ gives a bijection π from $\{q \mid q < p\}$ to itself. It's easy to see that

$$C = \{\sigma \mid (\sigma, \mathbb{P}_{\max} \restriction \sigma, \pi \restriction \sigma) \prec (\mathbb{R}, \mathbb{P}_{\max}, \pi)\},$$

is club and $S_{G_0} \cap C = S_{G_1} \cap C$. By homogeneity of $\mathbb{P}_{\max}$, there is a bijection (definable over) $L(\mathbb{R}, \mu)$ from OR onto $\mathrm{HOD}_{\mathcal{I}}$. So the ultraproduct $[\sigma \mapsto \mathrm{HOD}_{\mathcal{I}}]_H$ using functions in $L(\mathbb{R}, \mu)[G]$ is just $[\sigma \mapsto \mathrm{HOD}_{\mathcal{I}}]_\mu$ using functions in $L(\mathbb{R}, \mu)$.

Finally, to see $\mathrm{HOD}^{M_0}_{\mathbb{R}^V \cup \{\mathbb{R}^V\}} = \mathrm{HOD}^{M_1}_{\mathbb{R}^V \cup \{\mathbb{R}^V\}} \in V$, note that for any generic H, letting $V = L(\mathbb{R}, \mu)[G]$, $\mathrm{HOD}^{Ult(V,H)}_{\mathbb{R}^V \cup \{\mathbb{R}^V\}}$ is represented by $\sigma \mapsto \mathrm{HOD}^V_{\sigma \cup \{\sigma\}}$. Let f be such that $\mathrm{dom}(f) = S$ where S is $\mathcal{I}$-positive and $\forall \sigma \in S, f(\sigma) \in \mathrm{HOD}^V_{\sigma \cup \{\sigma\}}$. By normality, shrinking S if necessary, we may assume $\exists x \in \mathbb{R} \forall \sigma \in S, f(\sigma) \in \mathrm{HOD}^V_{\{x, \sigma\}}$ and Lemma 4.4 can be applied to this f. We finished the proof of Theorem 1.3. ⊣

Proof of Theorem 1.4. Let $\mathcal{I}$ be as in the hypothesis of the theorem. Since we're shooting for a model of the form $L(\mathbb{R}, \mu)$, we may as well assume there is no model M containing $\mathbb{R} \cup \mathrm{OR}$ such that $M \vDash \mathsf{AD}^+ + \Theta > \theta_0$; the existence of such an M gives a model of ZFC+ there are ω^2 Woodin cardinals, which in turns gives a model of the form $L(\mathbb{R}, \mu)$ satisfying the conclusion of the theorem.

By arguments in [18] (see in particular Section 4.6), the existence of a normal fine ideal $\mathcal{I}$ on $\mathcal{P}_{\omega_1}(\mathbb{R})$ such that $\mathcal{I}$ is precipituous and for all generics $G_0, G_1 \subseteq \mathcal{I}^+$, $s \in \mathrm{OR}^\omega$, $j_{G_0} \restriction \mathrm{HOD}_s = j_{G_1} \restriction \mathrm{HOD}_s \in V$ and $\mathrm{HOD}^{Ult(V,G_0)}_{\mathbb{R}^V \cup \{\mathbb{R}^V\}} = \mathrm{HOD}^{Ult(V,G_1)}_{\mathbb{R}^V \cup \{\mathbb{R}^V\}} \in V$

[27] The proof of (1) in fact shows more. It shows that if $A \subseteq \mathbb{R}$ is OD_s for some $s \in \mathrm{OR}^\omega$, then $A \in \mathcal{F}$ or $\mathbb{R} \backslash A \in \mathcal{F}$.

https://doi.org/10.1017/jsl.2014.65 Published online by Cambridge University Press

implies that AD^+ holds in $Lp(\mathbb{R})$. Let $M = Lp(\mathbb{R}) \vDash \mathsf{AD}^+$. Let $\mathcal{F}$ be the hod direct limit system in M, and $\mathcal{M}_\infty$ be the direct limit of $\mathcal{F}$ in M (see [7] or [18] for the full definition of $\mathcal{F}$). Fix a generic $G \subseteq \mathcal{I}^+$ and let $j = j_G$ be the generic embedding. To prove the theorem, we consider two cases.

CASE 1. $\Theta^M < \mathfrak{c}^+$.

We first observe that the argument in Chapter 5 of [18] for getting a strategy with branch condensation from $\mathcal{I}$ being strong and $j_H \restriction \mathrm{HOD}_{\{s,\mathcal{I}\}}$ being independent of V-generic $H \subset \mathcal{I}^+$ for any $s \in \mathrm{OR}^\omega$ can be used in our situation. Here are the two key points. The hypothesis of Case 1 replaces the strength of the ideal, which is used in showing Θ^M is countable in $j(M)$ and $j \restriction \mathcal{M}_\infty \in \mathrm{Ult}(V,G)$ and is countable there. The hypothesis $j_H \restriction \mathrm{HOD}_{\{s,\mathcal{I}\}} \in V$ being independent of V-generic $H \subset \mathcal{I}^+$ for any $s \in \mathrm{OR}^\omega$ is used in getting a strategy with branch condensation (see [2]), and a model N containing $\mathbb{R} \cup \mathrm{OR}$ such that $N \vDash \mathsf{AD}^+ + \Theta > \theta_0$.

Working over N, by a similar reasoning as in the first paragraph of this section, we obtain the desired model $L(\mathbb{R},\mu)$. This finishes the proof of the theorem in Case 1.

CASE 2. $\Theta^M \geq \mathfrak{c}^+$

Recall that $\mathcal{F}$ is the dual filter to $\mathcal{I}$. Let $\mu = \mathcal{F} \cap M$. First we observe by (1) that μ is a normal fine measure on $\mathcal{P}_{\omega_1}(\mathbb{R})^M$. Next, we need to see that μ doesn't construct sets of reals beyond M. This is the content of the next claim.

CLAIM. $L(\mathbb{R},\mu) \subseteq M$.[28]

PROOF. We first prove the following subclaim.

SUBCLAIM. μ is amenable to M in that if $\langle A_x \mid x \in \mathbb{R} \wedge A_x \in \mathcal{P}(\mathcal{P}_{\omega_1}(\mathbb{R}))^M \rangle \in M$ then $\langle A_x \mid x \in \mathbb{R} \wedge \mu(A_x) = 1 \rangle \in M$.

PROOF. Fix a sequence $\mathcal{C} = \langle A_x \mid x \in \mathbb{R} \wedge A_x \in \mathcal{P}(\mathcal{P}_{\omega_1}(\mathbb{R}))^M \rangle \in M$ and fix an ∞-Borel code S for the sequence. Let T be the tree for a universal $(\Sigma^2_1)^M$ set. We may assume $S \in OD^M$ and is a bounded subset of Θ^M. We also assume S codes T. Let $\mathcal{A}_S$ be the set coded by S over any model containing S. By MC and the definition of T, S in M, it's easy to see that in M,

$$\forall^*_\mu \sigma (\mathcal{P}(\sigma) \cap L(S,\sigma) = \mathcal{P}(\sigma) \cap L(T,\sigma) = \mathcal{P}(\sigma) \cap Lp(\sigma)).$$

Let $S^* = [\sigma \mapsto S]_\mu$ and $T^* = [\sigma \mapsto T]_\mu$ where the ultraproducts are taken with functions in M. Now, S^*, T^* may not be in M but

$$\mathcal{P}(\mathbb{R}) \cap L(S^*,\mathbb{R}) = \mathcal{P}(\mathbb{R}) \cap L(T^*,\mathbb{R}) = \mathcal{P}(\mathbb{R})^M.$$

This implies $\mathcal{C} \in L(S^*,\mathbb{R})$. For each $x \in \mathbb{R}$,

$$\begin{aligned} A_x \in \mu &\Leftrightarrow (\forall^*_\mu \sigma)(\sigma \in A_x \cap \mathcal{P}_{\omega_1}(\sigma)) \\ &\Leftrightarrow (\forall^*_\mu \sigma)(L(S,\sigma) \vDash \emptyset \Vdash_{Col(\omega,\sigma)} \sigma \in (\mathcal{A}_S)_x \\ &\Leftrightarrow L(S^*,\mathbb{R}) \vDash \emptyset \Vdash_{Col(\omega,\mathbb{R})} \mathbb{R} \in (\mathcal{A}^*_S)_x. \end{aligned}$$

The above shows $\mu \restriction \mathcal{C} \in L(S^*,\mathbb{R})$. Since $\mu \restriction \mathcal{C}$ can be coded by a set of reals in $L(S^*,\mathbb{R})$, $\mu \restriction \mathcal{C} \in M$. This finishes the proof of the claim. ⊣

Using the subclaim, we finish the proof of the claim as follows. Suppose α is least such that $\exists A \subseteq \mathcal{P}_{\omega_1}(\mathbb{R})\ A \in L_{\alpha+1}(\mathbb{R})[\mu] \setminus L_\alpha(\mathbb{R})[\mu]$ and $A \notin M$. By properties of

[28] We just need from the claim that $\mathcal{P}(\mathbb{R}) \cap L(\mathbb{R},\mu) \subset M$.

https://doi.org/10.1017/jsl.2014.65 Published online by Cambridge University Press

α and condensation of μ, there is a definable over $L_\alpha(\mathbb{R})[\mu]$ surjection of $\mathbb{R}$ onto $L_\alpha(\mathbb{R})[\mu]$. This implies $\alpha < \mathfrak{c}^+$. Also by minimality of α, $\mathcal{P}(\mathbb{R}) \cap L_\alpha(\mathbb{R})[\mu] \subseteq M$.

Now, if $\mathcal{P}(\mathbb{R}) \cap L_\alpha(\mathbb{R})[\mu] \subsetneq \mathcal{P}(\mathbb{R})^M$, then the subclaim gives us $\mu \cap L_\alpha(\mathbb{R})[\mu] \in M$ which implies $A \in M$. Contradiction. So we may assume $\mathcal{P}(\mathbb{R}) \cap L_\alpha(\mathbb{R})[\mu] = \mathcal{P}(\mathbb{R})^M$. This means $\Theta^{L_\alpha(\mathbb{R})[\mu]} = \Theta^M \geq \mathfrak{c}^+$. This contradicts the fact that $\alpha < \mathfrak{c}^+$. $\dashv$

The claim implies $L(\mathbb{R}, \mu) \vDash \mathsf{AD} + \mu$ is a normal fine measure on $\mathcal{P}_{\omega_1}(\mathbb{R})$. This finishes the proof of the theorem. $\dashv$

§5. Open problems and questions. We list some open problems and questions related to models of the form $L(\mathbb{R}, \mu)$. In Theorem 1.2, we prove the internal uniqueness of μ inside $L(\mathbb{R}, \mu)$. It's natural to ask whether $L(\mathbb{R}, \mu)$ is unique externally.

QUESTION 5.1. *Suppose μ_0, μ_1 are filters on $\mathcal{P}_{\omega_1}(\mathbb{R})$ such that for $i \in \{0, 1\}$, $L(\mathbb{R}, \mu_i) \vDash$ "$\mathsf{AD}^+ + \mu_i$ is a normal fine measure on $\mathcal{P}_{\omega_1}(\mathbb{R})$". Must $L(\mathbb{R}, \mu_0) = L(\mathbb{R}, \mu_1)$ and $\mu_0 \cap L(\mathbb{R}, \mu_0) = \mu_1 \cap L(\mathbb{R}, \mu_1)$? What is the consistency strength of having distinct models of $\mathsf{AD}^+ + V = L(\mathbb{R}, \mu)$?*

In [17], it's shown that $L(\mathbb{R}, \mu) \vDash \mathsf{AD}^+$ if and only if $L(\mathbb{R}, \mu) \vDash \Theta > \omega_2$. It's known that the equivalence fails for $L(\mathbb{R})$. However, the following is still open.

OPEN PROBLEM 5.2. *Suppose $L(\mathbb{R}) \vDash \Theta$ is strongly inaccessible.*[29] *Must $L(\mathbb{R}) \vDash \mathsf{AD}^+$?*

A variation of the above that we believe is still open is when we replace the hypothesis "$L(\mathbb{R}) \vDash \Theta$ is inaccessible" by "$\mathrm{HOD}^{L(\mathbb{R})} \vDash \Theta$ is inaccessible (or Woodin)". Finally, with regard to constructing $L(\mathbb{R}, \mu)$ in a core model induction, the following is still open (cf. [14]), where NS is the nonstationary ideal on ω_1.

CONJECTURE 5.3. *The following are equiconsistent.*

1. $\mathsf{ZFC}+$ *there are ω^2 Woodin cardinals.*
2. *NS is saturated and $\mathsf{WRP}^*_{(2)}(\omega_2)$ holds.*
3. *NS is saturated and $\mathsf{SRP}^*_{(2)}(\omega_2)$ holds.*

§6. Acknowledgments. The author would like to thank Hugh Woodin for his many insightful discussions regarding the paper's topic as well as his permission to include the proof of Theorems 1.1 and 1.2 in this paper. Thanks are also in order for Trevor Wilson for many helpful conversations regarding the content of Section 4. We would also like to thank the anonymous referee for pointing out many unclear passages and typos.

REFERENCES

[1] J. CUMMINGS, *Iterated forcing and elementary embeddings*, ***Handbook of Set Theory***, pp. 775–883, 2010.

[2] R. KETCHERSID, ***Toward $AD_{\mathbb{R}}$ from the Continuum Hypothesis and an ω_1-dense ideal***, Ph. D. thesis, Berkeley, 2000.

[3] P. KOELLNER and W. H. WOODIN, *Large cardinals from determinacy*, ***Handbook of Set Theory***, pp. 1951–2119, 2010.

[29] This means Θ is regular and for all $\kappa < \Theta$, there is a surjection from $\mathbb{R}$ onto $\mathcal{P}(\kappa)$ in $L(\mathbb{R})$.

https://doi.org/10.1017/jsl.2014.65 Published online by Cambridge University Press

[4] PAUL B. LARSON, ***The stationary tower: Notes on a course by W. Hugh Woodin***, vol. 32, University Lecture Series, American Mathematical Society, Providence, RI, 2004.

[5] ———, *Forcing over models of determinacy*, ***Handbook of Set Theory***, pp. 2121–2177, 2010.

[6] GRIGOR SARGSYAN, *A tale of hybrid mice*, available at http://math.rutgers.edu/∼gs481/.

[7] RALF SCHINDLER and JOHN R. STEEL, *The core model induction*, available at math.berkeley.edu/∼steel.

[8] R. SOLOVAY, *The independence of DC from AD*, ***Cabal Seminar 76–77***, pp. 171–183. Springer, New York, 1978.

[9] J. R. STEEL, *The derived model theorem*, available at www.math.berkeley.edu/∼ steel.

[10] ———, *Derived models associated to mice*. ***Computational prospects of infinity. Part I. Tutorials***, vol. 14, Lecture Notes Series, Institure of Mathematical Science, National Institure of Singapore, pp. 105–193, World Scientific, Hackensack, NJ, 2008.

[11] ———, *Scales in* $\mathbf{K}(\mathbb{R})$. ***Games, scales, and Suslin cardinals. The Cabal Seminar. Vol. I***, Lecture Notes in Logic, pp. 176–208, vol. 31, Association of Symbolic Logic, Chicago, IL, 2008.

[12] ———, *An outline of inner model theory*, ***Handbook of Set Theory***, pp. 1595–1684, 2010.

[13] JOHN R. STEEL and HUGH W. WOODIN, ***HOD as a core model***, 2012.

[14] JOHN STEEL and STUART ZOBLE, *Determinacy from strong reflection*. ***Transactions of the American Mathematical Society***, vol. 366 (2014), no. 8, pp. 4443–4490.

[15] J. R. STEEL and N. TRANG, AD^+*, derived models, and* σ_1*-reflection*, available at http://math.cmu.edu/∼namtrang. 2011.

[16] N. TRANG, *A hierarchy of measures from* $AD_{\mathbb{R}}$, available at http://math.cmu.edu/∼namtrang. 2013.

[17] ———, *Determinacy in* $L(\mathbb{R}, \mu)$. ***Journal of Mathematical Logic***, vol. 14 (2014), no. 01.

[18] T. M. WILSON, ***Contributions to Descriptive Inner Model Theory***, PhD thesis, University of California, 2012.

[19] W. HUGH WOODIN, ***The axiom of determinacy, forcing axioms, and the nonstationary ideal***, *de Gruyter Series in Logic and its Applications*, vol. 1, Walter de Gruyter & Co., Berlin, 1999.

[20] Y. ZHU, *The derived model theorem II*, available at http://graduate.math.nus.edu.sg/∼g0700513/der.pdf. 2010.

DEPARTMENT OF MATHEMATICAL SCIENCES
CARNEGIE MELLON UNIVERSITY
5000 FORBES AVE
PITTSBURGH, PA 15213, USA
E-mail: namtrang@andrew.cmu.edu

https://doi.org/10.1017/jsl.2014.65 Published online by Cambridge University Press