Mathematika 65 (2019) 1074–1092 © 2019 University College London
doi:10.1112/S0025579319000251

ZEROS OF THE MÖBIUS FUNCTION OF PERMUTATIONS

ROBERT BRIGNALL, VÍT JELÍNEK, JAN KYNČL AND DAVID MARCHANT

Abstract. We show that if a permutation π contains two intervals of length 2, where one interval is an ascent and the other a descent, then the Möbius function $\mu[1,\pi]$ of the interval $[1,\pi]$ is zero. As a consequence, we prove that the proportion of permutations of length n with principal Möbius function equal to zero is asymptotically bounded below by $(1-1/e)^2 \geqslant 0.3995$. This is the first result determining the value of $\mu[1,\pi]$ for an asymptotically positive proportion of permutations π. We further establish other general conditions on a permutation π that ensure $\mu[1,\pi]=0$, including the occurrence in π of any interval of the form $\alpha \oplus 1 \oplus \beta$.

§1. *Introduction.* In this section, we describe our principal results, and give an overview of the previous work in this area. Formal definitions are given in the next section.

Let σ and π be permutations of positive integers. We say that π *contains* σ if there is a subsequence of elements of π that is order-isomorphic to σ. As an example, 3624715 contains 3142 as the subsequences 6275 and 6475. If σ is contained in π, then we write $\sigma \leqslant \pi$.

The set of all permutations is a poset under the partial order given by containment. A closed interval $[\sigma,\pi]$ in a poset is the set defined by $\{\tau : \sigma \leqslant \tau \leqslant \pi\}$, and a half-open interval $[\sigma,\pi)$ is the set $\{\tau : \sigma \leqslant \tau < \pi\}$. The Möbius function of an interval $[\sigma,\pi]$ is defined recursively as follows:

$$\mu[\sigma,\pi] = \begin{cases} 0 & \text{if } \sigma \not\leqslant \pi, \\ 1 & \text{if } \sigma = \pi, \\ -\sum_{\tau \in [\sigma,\pi)} \mu[\sigma,\tau] & \text{otherwise.} \end{cases}$$

From the definition of the Möbius function it follows that if $\sigma < \pi$, then $\sum_{\tau\in[\sigma,\pi]} \mu[\sigma,\tau] = 0$.

In this paper, we are mainly concerned with the *principal Möbius function* of a permutation π, written $\mu[\pi]$, defined by $\mu[\pi] = \mu[1,\pi]$. We focus on the zeros of the principal Möbius function, that is, on the permutations π for which $\mu[\pi]=0$. We show that we can often determine that a permutation π is such a Möbius zero by examining small localities of π. We formalize this idea using

Received 12 October 2018, published online 14 August 2019.
MSC (2010): 05A05 (primary).
V. Jelínek and J. Kynčl were supported by project 16-01602Y of the Czech Science Foundation (GAČR). J. Kynčl was also supported by Charles University project UNCE/SCI/004.

https://doi.org/10.1112/S0025579319000251 Published online by Cambridge University Press

the notion of an "annihilator". Informally, an annihilator is a permutation α such that any permutation π containing an interval copy of α is a Möbius zero. We will describe an infinite family of annihilators.

We will also prove that any permutation containing an increasing as well as a decreasing interval of size 2 is a Möbius zero. Based on this result, we show that the asymptotic proportion of Möbius zeros among the permutations of a given length is at least $(1 - 1/e)^2 \geqslant 0.3995$. This is the first known result determining the values of the principal Möbius function for an asymptotically positive fraction of permutations. We will also demonstrate how our results on the principal Möbius function can be extended to intervals whose lower bound is not 1.

The question of computing the Möbius function in the permutation poset was first raised by Wilf [20]. The first result was by Sagan and Vatter [12], who determined the Möbius function on intervals of layered permutations. Steingrímsson and Tenner [19] found pairs of permutations (σ, π), where $\mu[\sigma, \pi] = 0$.

Burstein, Jelínek, Jelínková and Steingrímsson [7] found a recursion for the Möbius function for sum and skew decomposable permutations. They used this to determine the Möbius function for separable permutations. Their results for sum and skew decomposable permutations implicitly include a result that only concerns small localities, which is that, up to symmetry, if a permutation π of length greater than 2 begins 12, then $\mu[\pi] = 0$.


Smith [14] found an explicit formula for the Möbius function on the interval $[1, \pi]$ for all permutations π with a single descent. Smith's paper includes a lemma stating that if a permutation π contains an interval order-isomorphic to 123, then $\mu[\pi] = 0$. While the result in [7] requires that the permutation starts with a particular sequence, Smith's result is, in some sense, more general, as the critical interval (123) can occur in any position. Smith's lemma may be viewed as the first instance of an annihilator result. Our results on annihilators provide a common generalization of Smith's lemma and the above mentioned result of Burstein *et al* [7].

Smith [15] has explicit expressions for the Möbius function $\mu[\sigma, \pi]$ when σ and π have the same number of descents. In [16], Smith found an expression that determines the Möbius function for all intervals in the poset, although the expression involves a rather complicated double sum, starting with $\sum_{\tau \in [\sigma, \pi)} \mu[\sigma, \tau]$.

Brignall and Marchant [6] showed that if the lower bound of an interval is indecomposable, then the Möbius function depends only on the indecomposable permutations contained in the upper bound, and used this result to find a fast polynomial algorithm for computing $\mu[\pi]$, where π is an increasing oscillation.

§2. *Definitions and notation.* We let $\mathcal{S}_n$ denote the set of permutations of length n. We represent a permutation $\pi \in \mathcal{S}_n$ as a sequence $\pi_1, \pi_2, \ldots, \pi_n$ of integers from the set $[n] = \{1, 2, \ldots, n\}$ in which each element of $[n]$ appears exactly once. We let ϵ denote the unique permutation of length 0.

https://doi.org/10.1112/S0025579319000251 Published online by Cambridge University Press



367249815

Figure 1: A permutation with opposing adjacencies.

A sequence of numbers $a_1, a_2, \ldots, a_n$ is *order-isomorphic* to a sequence b_1, $b_2, \ldots, b_n$ if for every $i, j \in [n]$ we have $a_i < a_j \Leftrightarrow b_i < b_j$. A permutation $\pi \in \mathcal{S}_n$ *contains* a permutation $\sigma \in \mathcal{S}_k$ if π has a subsequence of length k order-isomorphic to σ.

An *interval* of a permutation π is a non-empty set of contiguous indices i, $i+1, \ldots, j$, where the set of values $\{\pi_i, \pi_{i+1}, \ldots, \pi_j\}$ is also contiguous. We say that π has an *interval copy* of a permutation α if it contains an interval of length $|\alpha|$ whose elements form a subsequence order-isomorphic to α.

An *adjacency* in a permutation is an interval of length 2. If a permutation contains a monotonic interval of length 3 or more, then each subinterval of length 2 is an adjacency. As examples, 367249815 has two adjacencies, 67 and 98; and 1432 also has two adjacencies, 43 and 32. If an adjacency is ascending, then it is an *up-adjacency*, otherwise it is a *down-adjacency*.

If a permutation π contains at least one up-adjacency, and at least one down-adjacency, then we say that π has *opposing adjacencies*. An example of a permutation with opposing adjacencies is 367249815, which is shown in Figure 1.

A permutation that does not contain any adjacencies is *adjacency-free*. Some early papers use the term "strongly irreducible" for what we call adjacency-free permutations. See, for example, Atkinson and Stitt [**3**].

Given a permutation σ of length n, and permutations $\alpha_1, \ldots, \alpha_n$, not all of them equal to the empty permutation ϵ, the *inflation* of σ by $\alpha_1, \ldots, \alpha_n$, written as $\sigma[\alpha_1, \ldots, \alpha_n]$, is the permutation obtained by removing the element σ_i if $\alpha_i = \epsilon$, and replacing σ_i with an interval isomorphic to α_i otherwise. Note that this is slightly different to the standard definition of inflation, originally given in Albert and Atkinson [**1**], which does not allow inflation by the empty permutation. As examples, $3624715[1, 12, 1, 1, 21, 1, 1] = 367249815$, and $3624715[\epsilon, 1, 1, \epsilon, 1, \epsilon, 1] = 3142$.

In many cases, we will be interested in permutations where most positions are inflated by the singleton permutation 1. If $\sigma = 3624715$, then we will write $\sigma[1, 12, 1, 1, 21, 1, 1] = 367249815$ as $\sigma_{2,5}[12, 21]$. Formally, $\sigma_{i_1,\ldots,i_k}[\alpha_1, \ldots, \alpha_k]$ is the inflation of σ where σ_{i_j} is inflated by α_j for $j = 1, \ldots, k$, and all other positions of σ are inflated by 1. When using this notation, we always assume that

https://doi.org/10.1112/S0025579319000251 Published online by Cambridge University Press

the indices $i_1, \ldots, i_k$ are distinct; however, we make no assumption about their relative order.

Our aim is to study the Möbius function of the permutation poset, that is, the poset of finite permutations ordered by containment. We are interested in describing general examples of intervals $[\sigma, \pi]$ such that $\mu[\sigma, \pi] = 0$, with particular emphasis on the case $\sigma = 1$. We say that π is a *Möbius zero* (or just *zero*) if $\mu[\pi] = 0$, and we say that π is a σ*-zero* if $\mu[\sigma, \pi] = 0$.

It turns out that many sufficient conditions for π to be a Möbius zero can be stated in terms of inflations. We say that a permutation ϕ is an *annihilator* if every permutation that has an interval copy of ϕ is a Möbius zero; in other words, for every τ and every $i \leqslant |\tau|$ the permutation $\tau_i[\phi]$ is a Möbius zero. More generally, we say that ϕ is a σ*-annihilator* if every permutation with an interval copy of ϕ is a σ-zero.

We say that a pair of permutations ϕ, ψ is an *annihilator pair* if for every permutation τ and every pair of distinct indices $i, j \leqslant |\tau|$, the permutation $\tau_{i,j}[\phi, \psi]$ is a Möbius zero.

Observe that for an annihilator ϕ, any permutation containing an interval copy of ϕ is also an annihilator. Likewise, if ϕ and ψ form an annihilator pair then any permutation containing disjoint interval copies of ϕ and ψ is an annihilator.

As our first main result, presented in §3, we show that the two permutations 12 and 21 are an annihilator pair, or equivalently, any permutation with opposing adjacencies is a Möbius zero. Later, in §5, we use this result to prove that Möbius zeros have asymptotic density at least $(1 - 1/e)^2$.

We also prove that for any two non-empty permutations α and β, the permutation $\alpha \oplus 1 \oplus \beta = 123[\alpha, 1, \beta]$ is an annihilator, and generalize this result to a construction of σ-annihilators for general σ. These results are presented in §4.

Finally, in §6, we give several examples of annihilators and annihilator pairs that do not directly follow from the results in the previous sections.

2.1. *Intervals with vanishing Möbius function.* We will now present several basic facts about the Möbius function, which are valid in an arbitrary finite poset. The first fact is a simple observation following directly from the definition of the Möbius function, and we present it without proof.

Fact 1. Let P be a finite poset with Möbius function μ_P, and let x and y be two elements of P satisfying $\mu_P[x, y] = 0$. Let Q be the poset obtained from P by deleting the element y, and let μ_Q be its Möbius function. Then for every $z \in Q$, we have $\mu_Q[x, z] = \mu_P[x, z]$.

Next, we introduce two types of intervals whose specific structure ensures that their Möbius function is zero.

Let $[x, y]$ be a finite interval in a poset P. We say that $[x, y]$ is *narrow-tipped* if it contains an element z different from x such that $[x, y) = [x, z]$. The element z is then called the *core* of $[x, y]$.

We say that the interval $[x, y]$ is *diamond-tipped* if there are three elements z, z' and w, all different from x, and such that

https://doi.org/10.1112/S0025579319000251 Published online by Cambridge University Press




Figure 2: Examples of narrow-tipped (left) and diamond-tipped (right) posets.

(1) $[x, y) = [x, z] \cup [x, z']$; and
(2) $[x, z] \cap [x, z'] = [x, w]$.

Condition 2 is equivalent to w being the greatest lower bound of z and z' in the interval $[x, y]$. The triple of elements (z, z', w) is again called the *core* of $[x, y]$. Figure 2 shows examples of narrow-tipped and diamond-tipped posets.

Fact 2. Let P be a poset with Möbius function μ_P, and let $[x, y]$ be a finite interval in P. If $[x, y]$ is narrow-tipped or diamond-tipped, then $\mu_P[x, y] = 0$.

Proof. If $[x, y]$ is narrow-tipped with core z, then

$$\mu_P[x, y] = -\sum_{v \in [x,y)} \mu_P[x, v] = -\sum_{v \in [x,z]} \mu_P[x, v] = 0.$$

If $[x, y]$ is diamond-tipped with core (z, z', w) then

$$\begin{aligned}
\mu_P[x, y] &= -\sum_{v \in [x,y)} \mu_P[x, v] \\
&= -\sum_{v \in [x,z] \cup [x,z']} \mu_P[x, v] \\
&= -\sum_{v \in [x,z]} \mu_P[x, v] - \sum_{v \in [x,z']} \mu_P[x, v] + \sum_{v \in [x,z] \cap [x,z']} \mu_P[x, v] \\
&= -\sum_{v \in [x,z]} \mu_P[x, v] - \sum_{v \in [x,z']} \mu_P[x, v] + \sum_{v \in [x,w]} \mu_P[x, v] \\
&= 0.
\end{aligned}$$

□

2.2. *Embeddings.* An *embedding* of a permutation $\sigma \in \mathcal{S}_k$ into a permutation $\pi \in \mathcal{S}_n$ is a function $f : [k] \to [n]$ with the following properties:

- $1 \leqslant f(1) < f(2) < \cdots < f(k) \leqslant n$;
- for any $i, j \in [k]$, we have $\sigma_i < \sigma_j$ if and only if $\pi_{f(i)} < \pi_{f(j)}$.

https://doi.org/10.1112/S0025579319000251 Published online by Cambridge University Press

We let $\mathcal{E}(\sigma, \pi)$ denote the set of embeddings of σ into π, and $E(\sigma, \pi)$ denote the cardinality of $\mathcal{E}(\sigma, \pi)$.

For an embedding f of σ into π, the *image* of f, denoted $\mathrm{Img}(f)$, is the set $\{f(i);\ i \in [k]\}$. In particular, $|\mathrm{Img}(f)| = |\sigma|$. The permutation σ is the *source* of the embedding f, denoted $\mathrm{src}_\pi(f)$. When π is clear from the context (as it usually will be) we write $\mathrm{src}(f)$ instead of $\mathrm{src}_\pi(f)$. Note that for a fixed π, the set $\mathrm{Img}(f)$ determines both f and $\mathrm{src}_\pi(f)$ uniquely.

We say that an embedding f is *even* if the cardinality of $\mathrm{Img}(f)$ is even, otherwise f is *odd*. In our arguments, we will frequently consider *sign-reversing* mappings on sets of embeddings (with different sources), which are mappings that map an odd embedding to an even one and vice versa. A typical example of a sign-reversing mapping is the so-called i-switch, which we now define. For a permutation $\pi \in \mathcal{S}_n$, let $\mathcal{E}(*, \pi)$ be the set $\bigcup_{\sigma \leqslant \pi} \mathcal{E}(\sigma, \pi)$. For an index $i \in [n]$, the *i-switch* of an embedding $f \in \mathcal{E}(*, \pi)$, denoted $\Delta_i(f)$, is the embedding $g \in \mathcal{E}(*, \pi)$ uniquely determined by the following properties:

$$\mathrm{Img}(g) = \mathrm{Img}(f) \cup \{i\} \quad \text{if } i \notin \mathrm{Img}(f),$$
$$\mathrm{Img}(g) = \mathrm{Img}(f) \setminus \{i\} \quad \text{if } i \in \mathrm{Img}(f).$$

For example, consider the permutations $\sigma = 132$ and $\pi = 41253$, and the embedding $f \in \mathcal{E}(\sigma, \pi)$ satisfying $f(1) = 2$, $f(2) = 4$ and $f(3) = 5$. We then have $\mathrm{Img}(f) = \{2, 4, 5\}$. Defining $g = \Delta_3(f)$, we see that $\mathrm{Img}(g) = \{2, 3, 4, 5\}$, and $\mathrm{src}(g)$ is the permutation 1243. Similarly, for $h = \Delta_5(g)$, we have $\mathrm{Img}(h) = \{2, 3, 4\}$ and $\mathrm{src}(h) = 123$.

Note that for any $\pi \in \mathcal{S}_n$ and any $i \in [n]$, the function Δ_i is a sign-reversing involution on the set $\mathcal{E}(*, \pi)$.

Consider, for a given $\pi \in \mathcal{S}_n$, two embeddings $f, g \in \mathcal{E}(*, \pi)$. We say that f *is contained in* g if $\mathrm{Img}(f) \subseteq \mathrm{Img}(g)$. Note that if f is contained in g, then the permutation $\mathrm{src}(f)$ is contained in $\mathrm{src}(g)$, and if a permutation λ is contained in a permutation τ, then any embedding from $\mathcal{E}(\tau, \pi)$ contains at least one embedding from $\mathcal{E}(\lambda, \pi)$. In particular, the mapping $f \mapsto \mathrm{src}(f)$ is a poset homomorphism from the set $\mathcal{E}(*, \pi)$ ordered by containment onto the interval $[\epsilon, \pi]$ in the permutation pattern poset.

2.3. *Möbius function via normal embeddings.* We will now derive a general formula, which will become useful in several subsequent arguments. The formula can be seen as a direct consequence of the well-known Möbius inversion formula. The following form of the Möbius inversion formula can be deduced, for example, from Proposition 3.7.2 in Stanley's book [**18**]. A poset is *locally finite* if each of its intervals is finite.

Fact 3 *(Möbius inversion formula).* Let P be a locally finite poset with maximum element y, let μ be the Möbius function of P, and let $F\colon P \to \mathbb{R}$ be a function. If a function $G\colon P \to \mathbb{R}$ is defined by

$$G(x) = \sum_{z \in [x, y]} F(z),$$

https://doi.org/10.1112/S0025579319000251 Published online by Cambridge University Press

then for every $x \in P$, we have

$$F(x) = \sum_{z \in [x,y]} \mu[x,z] G(z).$$

As a consequence, we obtain the following result.

PROPOSITION 4. *Let σ and π be arbitrary permutations, and let $F\colon [\sigma, \pi] \to \mathbb{R}$ be a function satisfying $F(\pi) = 1$. We then have*

$$\mu[\sigma,\pi] = F(\sigma) - \sum_{\lambda \in [\sigma,\pi)} \mu[\sigma,\lambda] \sum_{\tau \in [\lambda,\pi]} F(\tau). \tag{1}$$

Proof. Fix σ, π and F. For $\lambda \in [\sigma,\pi]$, define $G(\lambda) = \sum_{\tau \in [\lambda,\pi]} F(\tau)$. Using Fact 3 for the poset $P = [\sigma,\pi]$, we obtain

$$F(\sigma) = \sum_{\lambda \in [\sigma,\pi]} \mu[\sigma,\lambda] G(\lambda).$$

Substituting the definition of $G(\lambda)$ into the above identity and noting that $F(\pi) = 1$, we get

$$\begin{aligned} F(\sigma) &= \sum_{\lambda \in [\sigma,\pi]} \mu[\sigma,\lambda] \sum_{\tau \in [\lambda,\pi]} F(\tau) \\ &= \mu[\sigma,\pi] + \sum_{\lambda \in [\sigma,\pi)} \mu[\sigma,\lambda] \sum_{\tau \in [\lambda,\pi]} F(\tau), \end{aligned}$$

from which the proposition follows. □

In our applications, the function $F(\tau)$ will usually be defined in terms of the number of embeddings of τ into π satisfying certain additional conditions. In the literature, there are several definitions of such restricted embeddings, which are usually referred to as normal embeddings.

The notion of normal embedding seems to originate from the work of Björner [4], who defined normal embeddings between words, and showed that in the subword order of words over a finite alphabet, the Möbius function of any interval $[x, y]$ is equal in absolute value to the number of normal embeddings of x into y.

Björner's approach was later extended to the computation of the Möbius function in the composition poset [12], the poset of separable permutations [7], or the poset of permutations with a fixed number of descents [15]. In all these cases, the authors define a notion of "normal" embeddings tailored for their poset, and then express the Möbius function of an interval $[x, y]$ as the sum of weights of the "normal" embeddings of x into y, where each normal embedding has weight 1 or -1.

For general permutations, this simple approach fails, since the Möbius function $\mu[\sigma,\pi]$ is sometimes larger than the number of all embeddings of

https://doi.org/10.1112/S0025579319000251 Published online by Cambridge University Press

σ into π. However, Smith [16] introduced a notion of normal embedding applicable to arbitrary permutations, and proved a formula expressing $\mu[\sigma, \pi]$ as a summation over certain sets of normal embeddings.

For consistency, we adopt the term "normal embedding" in this paper, although in our proofs, we will need to introduce several notions of normality, which are different from each other and from the notions of normality introduced by previous authors. We will always use $\mathcal{NE}(\tau, \pi)$ to denote the set of embeddings of τ into π satisfying the definition of normality used in the given context, and we let $\mathrm{NE}(\tau, \pi)$ be the cardinality of $\mathcal{NE}(\tau, \pi)$.

The next proposition provides a general basis for all our subsequent applications of normal embeddings.

PROPOSITION 5. *Let σ and π be permutations. Suppose that for each $\tau \in [\sigma, \pi]$ we fix a subset $\mathcal{NE}(\tau, \pi)$ of $\mathcal{E}(\tau, \pi)$, with the elements of $\mathcal{NE}(\tau, \pi)$ being referred to as* normal embeddings *of τ into π. Assume that $\mathcal{NE}(\pi, \pi) = \mathcal{E}(\pi, \pi)$, that is, the unique embedding of π into π is normal. For each $\lambda \in [\sigma, \pi)$, define the two sets of embeddings*

$$\mathcal{NE}_\lambda(\mathrm{odd}, \pi) = \bigcup_{\substack{\tau \in [\lambda, \pi] \\ |\tau| \text{ odd}}} \mathcal{NE}(\tau, \pi) \quad \textit{and}$$

$$\mathcal{NE}_\lambda(\mathrm{even}, \pi) = \bigcup_{\substack{\tau \in [\lambda, \pi] \\ |\tau| \text{ even}}} \mathcal{NE}(\tau, \pi).$$

If for every $\lambda \in [\sigma, \pi)$ such that $\mu[\sigma, \lambda] \neq 0$, we have the identity

$$|\mathcal{NE}_\lambda(\mathrm{odd}, \pi)| = |\mathcal{NE}_\lambda(\mathrm{even}, \pi)|, \tag{2}$$

then $\mu[\sigma, \pi] = (-1)^{|\pi|-|\sigma|}\mathrm{NE}(\sigma, \pi)$.

Proof. The trick is to define the function $F(\tau) = (-1)^{|\pi|-|\tau|}\mathrm{NE}(\tau, \pi)$ and apply Proposition 4. This yields

$$\begin{aligned}
\mu[\sigma, \pi] &= F(\sigma) - \sum_{\lambda \in [\sigma, \pi)} \mu[\sigma, \lambda] \sum_{\tau \in [\lambda, \pi]} F(\tau) \\
&= F(\sigma) - \sum_{\lambda \in [\sigma, \pi)} \mu[\sigma, \lambda] \sum_{\tau \in [\lambda, \pi]} (-1)^{|\pi|-|\tau|}\mathrm{NE}(\tau, \pi) \\
&= F(\sigma) - \sum_{\lambda \in [\sigma, \pi)} \mu[\sigma, \lambda](-1)^{|\pi|}\big(|\mathcal{NE}_\lambda(\mathrm{even}, \pi)| - |\mathcal{NE}_\lambda(\mathrm{odd}, \pi)|\big) \\
&= F(\sigma) \\
&= (-1)^{|\pi|-|\sigma|}\mathrm{NE}(\sigma, \pi),
\end{aligned}$$

as claimed. □

We remark that the general formula of Proposition 4 can be useful even in situations where the more restrictive assumptions of Proposition 5 fail. An example of such application of Proposition 4 will appear in an upcoming manuscript [9], which is being prepared in parallel to this publication.

https://doi.org/10.1112/S0025579319000251 Published online by Cambridge University Press

§3. *Permutations with opposing adjacencies.* In this section, we show that if a permutation has opposing adjacencies, then the value of the principal Möbius function is zero.

THEOREM 6. *If π has opposing adjacencies, then $\mu[\pi] = 0$.*

For this theorem, we are able to give two proofs. One of them is based on the notion of diamond-tipped intervals, and the other uses the approach of normal embeddings. As both these approaches will later be adapted to more complicated settings, we find it instructive to include both proofs here.

Proof via diamond-tipped posets. For contradiction, suppose that the theorem fails, and let π be a shortest permutation with opposing adjacencies such that $\mu[\pi] \neq 0$. Since π has opposing adjacencies, there is a permutation τ and indices $i, j \leqslant |\tau|$ such that $\pi = \tau_{i,j}[12, 21]$. Define $\phi = \tau_{i,j}[1, 21]$ and $\phi' = \tau_{i,j}[12, 1]$.

We claim that the interval $[1, \pi]$ can be transformed into a diamond-tipped interval with core (ϕ, ϕ', τ) by deleting a set of Möbius zeros from the interior of $[1, \pi]$. Since by Fact 1, the deletion of Möbius zeros does not affect the value of $\mu[1, \pi]$, and since diamond-tipped intervals have zero Möbius function by Fact 2, this claim will imply that $\mu[1, \pi] = 0$, a contradiction.

To prove the claim, note first that any permutation $\lambda \in [1, \pi)$ with opposing adjacencies is a Möbius zero, since π is a minimal counterexample to the theorem. Choose any $\lambda \in [1, \pi)$. Observe that if λ has no up-adjacency, then $\lambda \leqslant \phi$, and symmetrically, if λ has no down-adjacency, then $\lambda \leqslant \phi'$. Thus, any $\lambda \in [1, \pi)$ not belonging to $[1, \phi] \cup [1, \phi']$ has opposing adjacencies and can be deleted from $[1, \pi]$.

Next, suppose that a permutation λ is in $[1, \phi] \cap [1, \phi']$ but not in $[1, \tau]$. Observe that any permutation in $[1, \phi] \setminus [1, \tau]$ has a down-adjacency, while any permutation in $[1, \phi'] \setminus [1, \tau]$ has an up-adjacency. It follows that λ has opposing adjacencies and can again be deleted from $[1, \pi]$.

After these deletions, the remaining poset is diamond-tipped with core (ϕ, ϕ', τ) as claimed, and hence $\mu[1, \pi] = 0$, a contradiction. □

Proof via normal embeddings. Suppose again that $\pi \in \mathcal{S}_n$ is a shortest counterexample. Suppose that π has an up-adjacency at positions $i, i+1$, and a down-adjacency at positions $j, j+1$. Note that the positions $i, i+1, j$ and $j+1$ are all distinct, and in particular $n \geqslant 4$.

We will say that an embedding $f \in \mathcal{E}(*, \pi)$ is *normal* if $\mathrm{Img}(f)$ is a superset of $[n] \setminus \{i, j\}$. In other words, $\mathrm{Img}(f)$ contains all positions of π with the possible exception of i and j. Thus, there are four normal embeddings.

We will use Proposition 5 with the above notion of normal embeddings and with $\sigma = 1$. Clearly, we have $\mathcal{E}(\pi, \pi) = \mathcal{NE}(\pi, \pi)$. The main task is to verify equation (2), that is, to show that for every $\lambda \in [1, \pi)$ such that $\mu[\lambda] \neq 0$ we have $|\mathcal{NE}_\lambda(\mathrm{odd}, \pi)| = |\mathcal{NE}_\lambda(\mathrm{even}, \pi)|$. To prove this identity, we let $\mathcal{NE}_\lambda(*, \pi)$ denote the set $\mathcal{NE}_\lambda(\mathrm{odd}, \pi) \cup \mathcal{NE}_\lambda(\mathrm{even}, \pi)$, and we will provide a sign-reversing involution on $\mathcal{NE}_\lambda(*, \pi)$.

https://doi.org/10.1112/S0025579319000251 Published online by Cambridge University Press

Choose a $\lambda \in [1, \pi)$ with $\mu[\lambda] \neq 0$. It follows that λ does not have opposing adjacencies, otherwise it would be a counterexample shorter than π. Without loss of generality, assume that λ has no up-adjacency. We will prove that the i-switch operation Δ_i is a sign-reversing involution on $\mathcal{NE}_\lambda(*, \pi)$.

It is clear that Δ_i is sign-reversing. We need to demonstrate that for every $f \in \mathcal{NE}_\lambda(*, \pi)$, the embedding $g = \Delta_i(f)$ is again in $\mathcal{NE}_\lambda(*, \pi)$. It is clear that g is normal. It remains to argue that $\mathrm{src}(g)$ contains λ, or in other words, that there is an embedding of λ into π contained in g. Let h be a (not necessarily normal) embedding of λ into π contained in f. If i is not in $\mathrm{Img}(h)$, then h is also contained in g, and we are done. Suppose now that $i \in \mathrm{Img}(h)$. Then $i + 1 \notin \mathrm{Img}(h)$, because i and $i + 1$ form an up-adjacency in π while λ has no up-adjacency. We modify the embedding h so that the element mapped to i will be mapped to $i + 1$ instead, and the mapping of the remaining elements is unchanged; let h' be the resulting embedding (formally, we have $\Delta_i(\Delta_{i+1}(h)) = h'$). Since i and $i + 1$ form an adjacency in π, we have $\mathrm{src}(h') = \mathrm{src}(h) = \lambda$. Since $i + 1$ is in the image of all normal embeddings, we see that h' is contained in g, and so $g \in \mathcal{NE}_\lambda(*, \pi)$. This shows that Δ_i is the required sign-reversing involution on $\mathcal{NE}_\lambda(*, \pi)$, verifying the assumptions of Proposition 5.

Proposition 5 then gives us that $\mu[1, \pi] = (-1)^{n-1}\mathrm{NE}(1, \pi)$. Since every normal embedding into π contains both $i + 1$ and $j + 1$ in its image, there is clearly no normal embedding of 1 into π and therefore we get $\mu[1, \pi] = 0$. □

§4. *A general construction of σ-annihilators.* Let σ be a fixed non-empty lower bound permutation (the case $\sigma = 1$ being the most interesting). Recall that a permutation ϕ is a *σ-zero* if $\mu[\sigma, \phi] = 0$, and ϕ is a *σ-annihilator* if every permutation with an interval copy of ϕ is a σ-zero. Clearly, any σ-annihilator is also a σ-zero. Our goal in this section is to present a general construction of an infinite family of σ-annihilators.

A permutation ϕ is *σ-narrow* if ϕ contains a permutation ϕ^- of size $|\phi| - 1$ such that every permutation in the set $[1, \phi) \setminus [1, \phi^-]$ is a σ-annihilator. In such a situation, we call ϕ^- a *σ-core of ϕ*.

Note that if ϕ is σ-narrow with σ-core ϕ^-, then the interval $[1, \phi]$ can be transformed into a narrow-tipped interval by a deletion of σ-annihilators. Our first goal is to show that, with a few exceptions, all σ-narrow permutations are σ-annihilators.

PROPOSITION 7. *If a permutation ϕ is σ-narrow with a σ-core ϕ^-, and if σ has no interval copy of ϕ or of ϕ^-, then ϕ is a σ-annihilator.*

Proof. Let ϕ be σ-narrow with a σ-core ϕ^-. Let π be a permutation with an interval copy of ϕ, that is, $\pi = \tau_i[\phi]$ for some τ and i. We show that $\mu[\sigma, \pi] = 0$. We may assume that $\sigma \leqslant \pi$, otherwise $\mu[\sigma, \pi] = 0$ trivially. Let π^- be the permutation $\tau_i[\phi^-]$. Note that $\sigma \neq \pi$ and $\sigma \neq \pi^-$, since σ has no interval copy of ϕ or of ϕ^-.

The key step of the proof is to show that any permutation in $[\sigma, \pi) \setminus [\sigma, \pi^-]$ is a σ-zero. After we have proved this, we may use Fact 1 to remove all such

https://doi.org/10.1112/S0025579319000251 Published online by Cambridge University Press

σ-zeros from the interval $[\sigma, \pi]$ without affecting the value of $\mu[\sigma, \pi]$; note that σ itself is clearly not a σ-zero, so it will not be removed, implying that $\sigma < \pi^-$. After the removal of $[\sigma, \pi) \setminus [\sigma, \pi^-]$, the remainder of the interval $[\sigma, \pi]$ is a narrow-tipped poset with core π^-, yielding $\mu[\sigma, \pi] = 0$ by Fact 2.

Therefore, to prove that $\mu[\sigma, \pi] = 0$ for a particular $\pi = \tau_i[\phi]$, it is enough to show that all the permutations in $[\sigma, \pi) \setminus [\sigma, \pi^-]$ are σ-zeros. We prove this by induction on $|\tau|$.

If $|\tau| = 1$, we have $\pi = \phi$ and $\pi^- = \phi^-$. Then all the permutations in $[1, \pi) \setminus [1, \pi^-]$ are σ-annihilators (and therefore σ-zeros) by definition of σ-narrowness, and in particular, restricting our attention to permutations containing σ, we see that all the permutations in $[\sigma, \pi) \setminus [\sigma, \pi^-]$ are σ-zeros, as claimed.

Suppose that $|\tau| > 1$. Consider a permutation $\gamma \in [\sigma, \pi) \setminus [\sigma, \pi^-]$. Since γ is contained in $\pi = \tau_i[\phi]$, it can be expressed as $\gamma = \tau_j^*[\phi^*]$ for some $\epsilon \leqslant \phi^* \leqslant \phi$ and $1 \leqslant \tau^* \leqslant \tau$, where τ^* has an embedding into τ, which maps j to i. Note that ϕ^* cannot be contained in ϕ^-, because in such a case we would have $\gamma \leqslant \pi^-$. Moreover, if $\phi^* = \phi$, then necessarily $\tau^* < \tau$, and by induction γ is a σ-zero. Finally, if ϕ^* is in $[1, \phi) \setminus [1, \phi^-]$, then ϕ^* is a σ-annihilator by the σ-narrowness of ϕ, and hence γ is a σ-zero. □

With the help of Proposition 7, we can now provide an explicit general construction of σ-annihilators.

PROPOSITION 8. *Let α and β be non-empty permutations. Assume that σ does not contain any interval copy of a permutation of the form $\alpha' \oplus \beta'$ with $1 \leqslant \alpha' \leqslant \alpha$ and $1 \leqslant \beta' \leqslant \beta$ (in particular, σ has no up-adjacency). Then $\alpha \oplus 1 \oplus \beta$ is σ-narrow with σ-core $\alpha \oplus \beta$, and $\alpha \oplus 1 \oplus \beta$ is a σ-annihilator.*

Proof. We proceed by induction on $|\alpha| + |\beta|$. Suppose first that $\alpha = \beta = 1$. Then trivially $\alpha \oplus 1 \oplus \beta = 123$ is σ-narrow with σ-core $\alpha \oplus \beta = 12$, since the set $[1, 123) \setminus [1, 12]$ is empty. Moreover, by assumption, σ has no interval copy of 12, and therefore also no interval copy of 123, hence 123 is a σ-annihilator by Proposition 7.

Suppose now that $|\alpha| + |\beta| > 2$. Define $\phi = \alpha \oplus 1 \oplus \beta$ and $\phi^- = \alpha \oplus \beta$. To prove that ϕ is σ-narrow with σ-core ϕ^-, we will show that any permutation $\gamma \in [1, \phi) \setminus [1, \phi^-]$ is a σ-annihilator. Such a γ has the form $\alpha' \oplus 1 \oplus \beta'$ for some $1 \leqslant \alpha' \leqslant \alpha$ and $1 \leqslant \beta' \leqslant \beta$, with $|\alpha'| + |\beta'| < |\alpha| + |\beta|$; note that we here exclude the cases $\alpha' = \epsilon$ and $\beta' = \epsilon$, because in these cases γ would be contained in ϕ^-. By induction, γ is σ-narrow, with σ-core $\gamma^- = \alpha' \oplus \beta'$. Moreover, σ has no interval isomorphic to γ or γ^-: observe that if σ had an interval isomorphic to γ, it would also have an interval isomorphic to $\alpha' \oplus 1$, which is forbidden by our assumptions on σ. Thus, we may apply Proposition 7 to conclude that γ is a σ-annihilator, and in particular ϕ is σ-narrow with σ-core ϕ^-, as claimed. Proposition 7 then gives us that ϕ is a σ-annihilator. □

Focusing on the special case $\sigma = 1$, which satisfies the assumptions of Proposition 8 trivially, we obtain the following result.

https://doi.org/10.1112/S0025579319000251 Published online by Cambridge University Press

COROLLARY 9. *For any non-empty permutations α and β, the permutation $\alpha \oplus 1 \oplus \beta$ is an annihilator.*

§5. *The density of zeros.* Our goal is to find an asymptotic positive lower bound on the proportion of permutations of length n whose principal Möbius function is zero. The key step is the following lemma.

LEMMA 10. *Let s_n be the number of permutations of size n with opposing adjacencies. Then*

$$\frac{s_n}{n!} = \left(1 - \frac{1}{e}\right)^2 + O\left(\frac{1}{n}\right).$$

Proof. Let a_n be the number of permutations of size n that have no up-adjacency, and let b_n be the number of permutations of size n that have neither an up-adjacency nor a down-adjacency.

The numbers a_n (sequence A000255 in the OEIS [13]) have already been studied by Euler [8], and it is known [11] that they satisfy $a_n/n! = e^{-1} + O(n^{-1})$.

The numbers b_n (sequence A002464 in the OEIS [13]) satisfy the asymptotics $b_n/n! = e^{-2} + O(n^{-1})$, which follows from the results of Kaplansky [10] (see also Albert *et al* [2]).

We may now express the number s_n of permutations with opposing adjacencies by inclusion–exclusion as follows: we subtract from $n!$ the number of permutations having no up-adjacency and the number of permutations having no down-adjacency, and then we add back the number of permutations having no adjacency at all. This yields $s_n = n! - 2a_n + b_n$, from which the lemma follows by the above-mentioned asymptotics of a_n and b_n. □

Combining Theorem 6 with Lemma 10 we obtain the following consequence, which is the main result of this section.

COROLLARY 11. *For a given n and for π a uniformly random permutation of length n, the probability that $\mu[\pi] = 0$ is at least*

$$\left(1 - \frac{1}{e}\right)^2 - O\left(\frac{1}{n}\right).$$

§6. *More complicated examples.* We will now construct several specific examples of annihilators and annihilator pairs, which are not covered by the general results obtained in the previous sections. We begin with a construction of four new annihilator pairs, which we will later use to construct new annihilators.

THEOREM 12. *The two permutations* 213 *and* 2431 *form an annihilator pair.*

Proof. Our proof is based on the concept of normal embeddings and follows a similar structure as the normal embedding proof of Theorem 6.

Suppose for contradiction that there is a permutation π that contains an interval isomorphic to 213 as well as an interval isomorphic to 2431, and that

https://doi.org/10.1112/S0025579319000251 Published online by Cambridge University Press




Figure 3: The intervals 213 and 2431 in Theorem 12. Normal embeddings may omit some of the hollow points.

$\mu[\pi] \neq 0$. Fix a smallest possible π, and let n be its length. Note that an interval isomorphic to 213 is necessarily disjoint from an interval isomorphic to 2431, and in particular, $n \geqslant 7$.

Let i, $i+1$ and $i+2$ be three positions of π containing an interval copy of 213, and let j, $j+1$, $j+2$ and $j+3$ be four positions containing an interval copy of 2431. We will apply the approach of Proposition 5, with $\sigma = 1$. We will say that an embedding $f \in \mathcal{E}(*, \pi)$ is *normal* if $\mathrm{Img}(f)$ is a superset of $[n] \setminus \{i+2, j+2, j+3\}$. Informally, the image of a normal embedding contains all the positions of π, except possibly some of the three positions that correspond to the value 3 of 213 or the values 3 and 1 of 2431 in the chosen interval copies of 213 and 2431, as shown in Figure 3. In particular, there are eight normal embeddings.

We now verify the assumptions of Proposition 5. We obviously have $\mathcal{NE}(\pi, \pi) = \mathcal{E}(\pi, \pi)$. The main task is to verify, for a given $\lambda \in [1, \pi)$ with $\mu[\lambda] \neq 0$, the identity (2) of Proposition 5, that is, the identity $|\mathcal{NE}_\lambda(\mathrm{odd}, \pi)| = |\mathcal{NE}_\lambda(\mathrm{even}, \pi)|$.

Fix a $\lambda \in [1, \pi)$ such that $\mu[\lambda] \neq 0$, and let $\mathcal{NE}_\lambda(*, \pi)$ be the set $\mathcal{NE}_\lambda(\mathrm{odd}, \pi) \cup \mathcal{NE}_\lambda(\mathrm{even}, \pi)$. We will describe a sign-reversing involution Φ_λ on $\mathcal{NE}_\lambda(*, \pi)$. The involution Φ_λ will always be equal to a switch operation Δ_k, where the choice of k will depend on λ.

Suppose first that λ does not contain any down-adjacency. We claim that Δ_{j+2} is an involution on the set $\mathcal{NE}_\lambda(*, \pi)$. To see this, choose $f \in \mathcal{NE}_\lambda(*, \pi)$ and define $g = \Delta_{j+2}(f)$. It is clear that g is a normal embedding.

To prove that g belongs to $\mathcal{NE}_\lambda(*, \pi)$, it remains to show that $\mathrm{src}(g)$ contains λ, or equivalently, that there is an embedding of λ into π that is contained in g. Let h be an embedding of λ into π, which is contained in f. If $j+2 \notin \mathrm{Img}(h)$, then h is also contained in g and we are done.

Suppose then that $j+2 \in \mathrm{Img}(h)$. This means that $j+1$ is not in $\mathrm{Img}(h)$, because π has a down-adjacency at positions $j+1$ and $j+2$, while λ has no down-adjacency. We now modify h in such a way that the element previously mapped to $j+2$ will be mapped to $j+1$, while the mapping of the remaining elements remains unchanged. Let h' be the embedding obtained from h by this modification; formally, we have $h' = \Delta_{j+1}(\Delta_{j+2}(h))$. Since the two elements

https://doi.org/10.1112/S0025579319000251 Published online by Cambridge University Press

π_{j+1} and π_{j+2} form an adjacency, we have $\mathrm{src}(h') = \mathrm{src}(h) = \lambda$. Moreover, h' is contained in g (recall that g is normal, and therefore $\mathrm{Img}(g)$ contains $j+1$). Consequently, g is in $\mathcal{NE}_\lambda(*, \pi)$, as claimed.

We now deal with the case when λ contains a down-adjacency. Since $\mu[\lambda] \neq 0$, it follows by Theorem 6 that λ has no up-adjacency. We distinguish two subcases, depending on whether λ contains an interval copy of 2431.

Suppose that λ contains an interval copy of 2431. We will prove that in this case, Δ_{i+2} is a sign-reversing involution on $\mathcal{NE}_\lambda(*, \pi)$. We begin by observing that λ has no interval copy of 213, otherwise λ would be a counterexample to Theorem 12, contradicting the minimality of π. Fix again an embedding $f \in \mathcal{NE}_\lambda(*, \pi)$, and define $g = \Delta_{i+2}(f)$. As in the previous case, g is clearly normal, and we only need to show that there is an embedding of λ into π contained in g. Let h be an embedding of λ into π contained in f. If $i+2 \notin \mathrm{Img}(h)$, then h is contained in g and we are done, so suppose $i+2 \in \mathrm{Img}(h)$. If at least one of the two positions i and $i+1$ belongs to $\mathrm{Img}(h)$, then λ contains an up-adjacency or an interval copy of 213, contradicting our assumptions. Therefore, we can modify h so that the element mapped to $i+2$ is mapped to i instead, obtaining an embedding of λ contained in g and showing that $g \in \mathcal{NE}_\lambda(*, \pi)$.

Finally, suppose that λ has no interval copy of 2431. In this case, we prove that Δ_{j+3} is the required involution on $\mathcal{NE}_\lambda(*, \pi)$. As in the previous cases, we fix $f \in \mathcal{NE}_\lambda(*, \pi)$, define $g = \Delta_{j+3}(f)$ and let h be an embedding of λ contained in f; we again want to modify h into an embedding λ contained in g. Let α be the subpermutation of λ formed by those positions that are mapped into the set $J = \{j, j+1, j+2, j+3\}$ by h. Recall that the positions in J induce an interval copy of 2431 in π. In particular, $\alpha \leqslant 2431$, and λ has an interval copy of α. We know that $\alpha \neq 2431$, since we assume that λ has no interval copy of 2431. Also, $\alpha \neq 321$, since 321 is an annihilator by Corollary 9, while $\mu[\lambda] \neq 0$. Finally, $\alpha \neq 231$, since λ has no up-adjacency. This implies that $\alpha \leqslant 132$, and we can modify h so that all the positions originally mapped into J will get mapped into $J \setminus \{j+3\}$, obtaining an embedding of λ into π contained in g.

Having thus verified the assumptions of Proposition 5, we can conclude that $\mu[\pi] = (-1)^{|\pi|-1}\mathrm{NE}(1, \pi) = 0$, a contradiction. □

The following three results are proved using similar methods to those used in the proof of Theorem 12.

THEOREM 13. *The permutations* 2143 *and* 2431 *form an annihilator pair.*

THEOREM 14. *The permutations* 312 *and* 23514 *form an annihilator pair.*

THEOREM 15. *The permutations* 25134 *and* 23514 *form an annihilator pair.*

We omit the proofs here, but they can be found in an extended version of this paper [**5**].

With the help of the new annihilator pairs established in Theorems 12 to 15, we are able to present several new examples of annihilators.

https://doi.org/10.1112/S0025579319000251 Published online by Cambridge University Press




Figure 4: The three annihilators from Theorem 16, and the posets of their subpermutations. The figures omit the permutations with opposing adjacencies, as well as the permutations with an interval copy of a permutation of the form $\alpha \oplus 1 \oplus \beta$.

THEOREM 16. *Each of the three permutations* 215463*,* 236145 *and* 214653 *is a Möbius annihilator.*

Proof. We first present the proof for the permutation 215463. Let $\alpha = 215463$, $\beta = \alpha_1[\epsilon] = 14352$, $\beta' = \alpha_6[\epsilon] = 21435$ and $\gamma = \alpha_{1,6}[\epsilon, \epsilon] = 1324$. From Figure 4 (left) we see that, after the removal of the annihilators $\alpha_3[\epsilon]$, $\alpha_4[\epsilon]$ and $\alpha_5[\epsilon]$, the interval $[1, \alpha]$ becomes diamond-tipped with core (β, β', γ). Hence by Facts 1 and 2 we have $\mu[1, \alpha] = 0$.

Let π be a permutation of the form $\tau_i[\alpha]$ for some τ and $i \leqslant |\tau|$. We will show, by induction on $|\tau|$, that π is a zero. The case $|\tau| = 1$ has been proved in the previous paragraph.

Assume that $|\tau| > 1$. We will demonstrate that we can remove some zeros from the interval $[1, \pi]$ to end up with a diamond-tipped interval with core $(\tau_i[\beta], \tau_i[\beta'], \tau_i[\gamma])$. Choose a $\lambda \in [1, \pi)$. We can then write λ as $\lambda = \tau_j^*[\alpha^*]$ for some $\tau^* \leqslant \tau$ and some (possibly empty) $\alpha^* \leqslant \alpha$, where τ^* has an embedding into τ mapping j to i.

If α^* is an annihilator, then λ is a zero and can be removed. If $\alpha^* = \alpha$, then $|\tau^*| < |\tau|$, and by induction, λ is a zero and can be removed. In all the other cases, we have $\alpha^* \leqslant \beta$ or $\alpha^* \leqslant \beta'$, and in particular, λ belongs to $[1, \tau_i[\beta]] \cup [1, \tau_i[\beta']]$.

https://doi.org/10.1112/S0025579319000251 Published online by Cambridge University Press

Suppose now that λ is in $[1, \tau_i[\beta]] \cap [1, \tau_i[\beta']]$ but not in $[1, \tau_i[\gamma]]$. Since $\lambda \leqslant \tau_i[\beta]$, we can write it as $\lambda = \tau_j^L[\beta^L]$, for some $\tau^L \leqslant \tau$ and $\beta^L \leqslant \beta$, where τ^L has an embedding into τ mapping j to i. Since $\lambda \not\leqslant \tau_i[\gamma]$, we know that $\beta^L \not\leqslant \gamma$. This means that $\beta^L \in [1, \beta] \setminus [1, \gamma] = \{14352, 3241, 1342, 231\}$. Similarly, $\lambda \in [1, \tau_i[\beta']] \setminus [1, \tau_i[\gamma]]$ means that λ can be written as $\lambda = \tau_k^R[\beta^R]$, with $\beta^R \in \{21435, 2143\}$. Since λ has an interval copy of β^L as well as an interval copy of β^R, Theorem 6 shows that λ is a zero if $\beta^L \in \{1342, 231\}$, and Theorem 13 shows that λ is a zero if $\beta^L \in \{14352, 3241\}$ (using that 3241 is a diagonal reflection of 2431). Therefore, λ can be removed.

After the removal described above, $[1, \pi]$ is transformed into a diamond-tipped interval, showing that π is a zero.

The arguments for the other two permutations are completely analogous. For 236145 we have $\alpha = 236145$, $\beta = 25134$, $\beta' = 23514$, $\gamma = 2413$, $\beta^L \in \{25134, 1423\}$ and $\beta^R \in \{23514, 2314\}$, and use Theorems 12, 14 and 15. For 214653 we have $\alpha = 214653$, $\beta = 13542$, $\beta' = 2143$, $\gamma = 132$, $\beta^L \in \{13542, 2431, 1342, 231\}$ and $\beta^R \in \{2143, 213\}$, and use Theorems 6, 12 and 13. □

The annihilator 215463 of Theorem 16 can be written as a sum of two intervals, namely $215463 = 21 \oplus 3241$. One might wonder whether the two summands are in fact an annihilator pair. This, however, is not the case, as evidenced by the permutation $32417685 = 3241 \oplus 3241$, which is not a Möbius zero. An analogous example applies to $214653 = 21 \oplus 2431$.

In the proof of Theorem 16, it was crucial that for each $\alpha \in \{215463, 236145, 214653\}$, the interval $[1, \alpha]$ becomes diamond-tipped after the removal of some annihilators. However, this property alone is not sufficient to make a permutation α an annihilator. Consider, for instance, the permutation $\alpha = 214635$. We may routinely check that by removing some annihilators, the interval $[1, \alpha]$ can be made diamond-tipped with core ($\beta = 13524$, $\beta' = 21435$, $\gamma = 1324$). This implies that α is a Möbius zero by Facts 1 and 2; however, it does not imply that α is an annihilator. In fact, α is not an annihilator, as demonstrated by the permutation

$$\begin{aligned}\pi &= 582741936_{2,4,5}[\beta, \alpha, \beta'] \\ &= 9, 17, 19, 21, 18, 20, 2, 12, 11, 14, 16, 13, 15, 5, 4, 7, 6, 8, 1, 22, 3, 10,\end{aligned}$$

whose principal Möbius function is 1, not 0. This example also shows that not all Möbius zeros are annihilators.

In fact, among permutations of size at most 6, there are up to symmetry four Möbius zeros that are not annihilators. Apart from the permutation 214635 pointed out above, there are these three more examples: 235614, 254613 and 465213. To see that these three permutations are not annihilators, it suffices to check that for any $\alpha \in \{235614, 254613, 465213\}$, the permutation $24153_2[\alpha]$ has a non-zero principal Möbius function. We verified, with the help of a computer, that all the Möbius zeros of size at most 6 that are not symmetries of the four examples above can be shown to be annihilators by our results. This data is available at https://iuuk.mff.cuni.cz/∼jelinek/mf/zeros.txt.

https://doi.org/10.1112/S0025579319000251 Published online by Cambridge University Press

Table 1: The density of Möbius zeros among permutations of length n, with $n = 1, \ldots, 13$.

n	d_n	n	d_n
1	0.0000	8	0.5942
2	0.0000	9	0.6019
3	0.3333	10	0.6040
4	0.4167	11	0.6034
5	0.4833	12	0.6021
6	0.5361	13	0.6006
7	0.5742		

§7. *Concluding remarks.* Given Theorem 6, it is natural to wonder if we can find a similar result that applies to a permutation with multiple adjacencies, but no opposing adjacencies. One difficulty here is that there are permutations that have multiple adjacencies, and do not have opposing adjacencies, where the principal Möbius function value is non-zero. As an example, any permutation $\pi = 2, 1, 4, 3, \ldots, 2k, 2k-1 = \bigoplus^k 21$ has $\mu[\pi] = -1$ by the results of Burstein *et al* [**7**, Corollary 3].

Let d_n be the "density of zeros" of the Möbius function, that is, the probability that $\mu[\pi] = 0$ for a uniformly random permutation π of size n. The asymptotic behaviour of d_n is still elusive.

Problem 17. Does the limit $\lim_{n\to\infty} d_n$ exist? And if it does, what is its value?

Corollary 11 implies that $\liminf_{n\to\infty} d_n \geqslant (1 - 1/e)^2 \geqslant 0.3995$. We have no upper bound on d_n apart from the trivial bound $d_n \leqslant 1$, but computational data suggest that simple permutations very often (though not always) have a non-zero principal Möbius function, where a permutation π is *simple* if all its intervals have size 1 or $|\pi|$. Since a random permutation is simple with probability approaching $1/e^2$ [**2**], this would suggest that $\limsup_{n\to\infty} d_n$ is at most $1 - 1/e^2 \approx 0.8647$.

Table 1 lists the values of d_n for $n = 1, \ldots, 13$. The values are based on data supplied by Smith [**17**] for $1 \leqslant n \leqslant 9$, and calculations performed by the fourth author. Data files with the values of the principal Möbius function for all permutations of length 12 or less are available from https://doi.org/10.21954/ou.rd.7171997.v2. Based on this somewhat limited numeric evidence, we make the following conjecture.

CONJECTURE 18. *The values d_n are bounded from above by 0.6040.*

It is natural to look for further ways to identify Möbius zeros and Möbius annihilators. Characterizing all the Möbius zeros would be an ambitious goal, since $\mu[\pi]$ might be zero as a result of "accidental" cancellations with no deeper structural significance for π.

https://doi.org/10.1112/S0025579319000251 Published online by Cambridge University Press

An *annihilator multiset* is a multiset of permutations $A = \{\alpha_1, \ldots, \alpha_n\}$ such that any permutation π that contains disjoint interval copies of the permutations $\alpha_1, \ldots, \alpha_n$ has $\mu[\pi] = 0$.

If $A = \{\alpha_1, \ldots, \alpha_n\}$ and $B = \{\beta_1, \ldots, \beta_m\}$ are annihilator multisets, then we say that A *contains* B if $A \neq B$ and we can find the elements of B as disjoint interval copies in the elements of A. An annihilator multiset A is *minimal* if there is no annihilator multiset contained in A.

Using [7, Corollary 3], which implies $\mu[\pi] = \mu[\pi \oplus \pi]$ for $\pi \neq 1$, it is simple to show that the permutations in a minimal annihilator multiset are, in fact, all distinct, and so we can refer to *minimal annihilator sets* of permutations.

Problem 19. Which permutations are Möbius annihilators? Are there infinitely many minimal annihilator sets that contain just one element, and are not of the form $\alpha \oplus 1 \oplus \beta$?

It seems likely to us that the proofs of Theorems 12–15 might be extended to give several more annihilator pairs, such as (312, 235614). However, we do not see any general pattern in these examples yet.

Problem 20. Are there infinitely many minimal annihilator sets with two elements?

Problem 21. Are there any minimal annihilator sets with more than two elements?

Acknowledgments. We wish to thank Ida Kantor and Martin Tancer for helpful comments. We would also like to thank the anonymous reviewer for their feedback and suggestions.

References

1. M. H. Albert and M. D. Atkinson, Simple permutations and pattern restricted permutations. *Discrete Math.* **300**(1-3) (2005), 1–15.
2. M. H. Albert, M. D. Atkinson and M. Klazar, The enumeration of simple permutations. *J. Integer Seq.* **6**(4) (2003), Article 03.4.4.
3. M. D. Atkinson and T. Stitt, Restricted permutations and the wreath product. *Discrete Math.* **259**(1–3) (2002), 19–36.
4. A. Björner, The Möbius function of subword order. In *Invariant Theory and Tableaux (Minneapolis, MN, 1988)* (*IMA Vol. Math. Appl.,* **19**), Springer (New York, 1990), 118–124.
5. R. Brignall, V. Jelínek, J. Kynčl and D. Marchant, Zeros of the Möbius function of permutations. *Preprint*, 2018, arXiv:org/abs/1810.05449v1.
6. R. Brignall and D. Marchant, The Möbius function of permutations with an indecomposable lower bound. *Discrete Math.* **341**(5) (2018), 1380–1391.
7. A. Burstein, V. Jelínek, E. Jelínková and E. Steingrímsson, The Möbius function of separable and decomposable permutations. *J. Combin. Theory Ser. A* **118**(8) (2011), 2346–2364.
8. L. Euler, Recherches sur une nouvelle espèce des quarrés magiques. *Verhandelingen uitgegeven door het zeeuwsch Genootschap der Wetenschappen te Vlissingen* **9** (1782), 85–239.
9. V. Jelínek, I. Kantor, J. Kynčl and M. Tancer, On the growth of the Möbius function of permutations. *J. Combin. Theory, Ser. A* **169** (2020),105121.
10. I. Kaplansky, The asymptotic distribution of runs of consecutive elements. *Ann. Math. Stat.* **16**(2) (1945), 200–203.

https://doi.org/10.1112/S0025579319000251 Published online by Cambridge University Press

11. M. Rumney and E. J. F. Primrose, A sequence connected with the sub-factorial sequence. *Gaz. Math.* **52**(382) (1968), 381–382.
12. B. E. Sagan and V. Vatter, The Möbius function of a composition poset. *J. Algebraic Combin.* **24**(2) (2006), 117–136.
13. N. J. A. Sloane, The on-line encyclopedia of integer sequences, published electronically at http://oeis.org/.
14. J. P. Smith, On the Möbius function of permutations with one descent. *Electron. J. Combin.* **21**(2) (2014), 19 Paper 2.11.
15. J. P. Smith, Intervals of permutations with a fixed number of descents are shellable. *Discrete Math.* **339**(1) (2016), 118–126.
16. J. P. Smith, A formula for the Möbius function of the permutation poset based on a topological decomposition. *Adv. Appl. Math.* **91** (2017), 98–114.
17. J. P. Smith, private correspondence, 2018.
18. R. P. Stanley, *Enumerative Combinatorics*, 2nd edn., Vol. 1 (*Cambridge Studies in Advanced Mathematics* **49**), Cambridge University Press (Cambridge, 2012).
19. E. Steingrímsson and B. E. Tenner, The Möbius function of the permutation pattern poset. *J. Comb.* **1**(1) (2010), 39–52.
20. H. S. Wilf, The patterns of permutations. *Discrete Math.* **257**(2-3) (2002), 575–583.

Robert Brignall,
School of Mathematics and Statistics,
The Open University,
Milton Keynes, MK7 6AA,
U.K.
E-mail: robert.brignall@open.ac.uk

Vít Jelínek,
Computer Science Institute,
Faculty of Mathematics and Physics,
Charles University, Prague,
Czech Republic
E-mail: jelinek@iuuk.mff.cuni.cz

Jan Kynčl,
Department of Applied Mathematics,
Faculty of Mathematics and Physics,
Charles University, Prague,
Czech Republic
E-mail: kyncl@kam.mff.cuni.cz

David Marchant,
School of Mathematics and Statistics,
The Open University,
Milton Keynes, MK7 6AA,
U.K.
E-mail: david.marchant@open.ac.uk

https://doi.org/10.1112/S0025579319000251 Published online by Cambridge University Press