Ergod. Th. & Dynam. Sys. (2006), **26**, 1339–1362
doi:10.1017/S0143385706000393
© *2006 Cambridge University Press*
Printed in the United Kingdom

Lorentzian foliations on 3-manifolds

C. BOUBEL†, P. MOUNOUD‡ and C. TARQUINI†

† *École Normale Supérieure de Lyon, Unité de Mathématiques Pures et Appliquées, 46 allée d'Italie, F-69364 Lyon Cedex 07, France*
(e-mail: {charles.boubel, cedric.tarquini}@ens-lyon.fr)
‡ *Université Bordeaux 1, Laboratoire Bordelais d'Analyse et Géométrie, 351 cours de la Libération, F-33405 Talence Cedex, France*
(e-mail: pierre.mounoud@math.u-bordeaux1.fr)

(*Received* 18 *January* 2006 *and accepted in revised form* 18 *April* 2006)

Abstract. We study transversely Lorentzian foliations of closed 3-manifolds. We classify them under a completeness hypothesis and deduce the dual classification of codimension-one geodesically complete timelike totally geodesic foliations. Also we provide an example of a Lorentzian foliation of a compact 3-manifold which is equicontinuous on a *proper* dense open subset of the manifold, and hence not transversely complete.

1. *Introduction*

In this article we study the foliations admitting a transverse Lorentzian metric. It is the Lorentzian analogue of the better-known Riemannian foliations and they still have a rigid transverse geometry. We recall that a transverse (pseudo-)Riemannian metric of a foliation $\mathcal{F}$ is a (pseudo-)Riemannian metric on the normal bundle $\nu(\mathcal{F}) = TM/T\mathcal{F}$, invariant under the flow of any vector field tangent to $\mathcal{F}$. We send the reader to the book [**14**] of Molino for a general theory in the Riemannian case and to §2 here for a precise definition. The Lorentzian foliations are thus the foliations admitting a transverse pseudo-Riemannian metric of index 1.

One of the main differences between pseudo-Riemannian (and in particular Lorentzian) and Riemannian geometry is that a pseudo-Riemannian metric may admit non-equicontinuous sequences of isometries. In the same way, the holonomy pseudogroup of a transversely pseudo-Riemannian foliation may be non-equicontinuous. For example, on the compact 3-manifolds, Ghys used that the Anosov flows having smooth weak stable and unstable foliations and preserving a volume form are transversely Lorentzian to classify them, see [**8**].

Nevertheless, on compact manifolds, we may hope that equicontinuity fails in very special situations. Indeed, it is so for sequences of isometries of a compact pseudo-Riemannian manifold: such a sequence, either is equicontinuous, or has nowhere-bounded

https://doi.org/10.1017/S0143385706000393 Published online by Cambridge University Press

derivative, see [**27**]. Moreover, an equicontinuous group of pseudo-Riemannian isometries preserves a Riemannian metric; this is mainly the Ascoli theorem. The Lorentzian foliations that were already known satisfied also such an alternative 'everywhere or nowhere equicontinuous': they were either Anosov or Riemannian.

However, it turns out that this alternative is false for Lorentzian foliations. In §6 we give an example of a Lorentzian foliation which is equicontinuous on a dense proper open subset. However, adding a 'completeness' hypothesis, called transverse completeness, see Definition 2.4 here, this alternative is still true. Using Molino's theory and the classification of the Lorentzian Anosov flows, given by Ghys in [**8**], we prove the following theorem.

THEOREM 4.1. *Up to finite cover, a one-dimensional transversely complete Lorentzian foliation on a compact 3-manifold is either smoothly equivalent to the foliation generated by an algebraic Anosov flow or Riemannian.*

Then using Carrière's classification [**3**] of the Riemannian flows on compact 3-manifolds, we give a detailed panorama of the last case.

As a natural corollary of Theorem 4.1, we classify the codimension-one foliations of Lorentzian compact 3-manifolds, the leaves of which are *timelike*, totally geodesic and geodesically complete for a Lorentzian metric. Indeed the integral foliation of the orthogonal distribution of those foliations are transversely complete Lorentzian foliations, see Propositions 2.8 and 5.3. We obtain the following.

THEOREM 5.15. *Up to a finite cover the timelike geodesically complete totally geodesic codimension-one foliations of the closed three-dimensional manifolds are:*

(1) *the foliations of the circle bundles over the torus, transverse to the fibres;*
(2) *the foliations on $\mathbb{T}^3_A$ with a compact leaf and with neither Reeb nor type II components.*

We send the reader to §5 for the definition of a type II component and to §2.2 for a reminder of that of $\mathbb{T}^3_A$. We make precise that, on $\mathbb{T}^3_A$, the foliations with neither Reeb nor type II components are exactly the taut foliations.

1.1. *Structure of the article.* In §2 we introduce the notion of transversely complete pseudo-Riemannian foliation. In §3 we prove the alternative 'Riemannian or Anosov' for transversely complete Lorentzian foliations and in §4 we give the classification: Theorem 4.1 and its detailed version, Theorem 4.2.

Section 5 deals with timelike totally geodesic foliations and Theorem 5.15 is proved. In §6, we give examples of non-transversely complete Lorentzian foliations.

1.2. *Smoothness of the objects.* All manifolds involved here are without boundary and all differential objects are of class C^2. Notice that in the C^1 framework it is known that many transversely Lorentzian flows exist. Consider the numerous Anosov flows preserving a volume form and having weak stable and unstable distributions of class C^1 but not of class C^2.

https://doi.org/10.1017/S0143385706000393 Published online by Cambridge University Press

2. *Definitions of completeness*

2.1. *The definitions.* Let $\mathcal{F}$ be a foliation of codimension q on a manifold M. We will say that $\mathcal{F}$ is transversely Lorentzian if it admits a so-called transverse Lorentzian metric. One can define this transverse structure using a holonomy pseudogroup H of $\mathcal{F}$ acting on some transverse manifold T and a Lorentzian metric g_T on T such that each element of H preserves g_T. However, we thought that a definition in terms of a bundle-like metric would be clear enough. We recall the definitions of a bundle-like metric, a basic function, form or vector field, and transverse parallelism, for a foliation $\mathcal{F}$ on a manifold M (see, for instance, [**14**]).

(1) A real function of M is basic (or foliated) for $\mathcal{F}$ if it is constant along the leaves.

(2) A vector field Y on M is basic (or foliated) for $\mathcal{F}$ if for all vector field X tangent to $\mathcal{F}$ the vector field $[X, Y]$ is tangent to $\mathcal{F}$. This is equivalent to asking that the flow of Y preserve the foliation.

(3) A 1-form α on M is basic (or foliated) for $\mathcal{F}$ if for any vector field X tangent to $\mathcal{F}$ we have $\alpha(X) = 0$ and $i_X\, d\alpha = 0$ (or equivalently if $\alpha(X) = 0$ and $L_X\alpha = 0$ for L_X the Lie derivative (by X). Notice that this constraint is stronger than that of point 2 for vector fields. Indeed, a foliation $\mathcal{F}$ always admits non-zero basic vector fields (though not interesting ones: the fields tangent to $\mathcal{F}$), while a foliation may admit no non-zero basic forms.

(4) A transverse parallelism of $\mathcal{F}$ is a q-tuple of basic vector fields $(X_1, \ldots, X_q)$ spanning a complement of $T\mathcal{F}$ at every point of M. In general, a foliation $\mathcal{F}$ does not admit any parallelism. If it admits one, it is said to be transversely parallelizable.

(5) A (pseudo-)Riemannian metric g on M is called $\mathcal{F}$-bundle-like if its restriction to the tangent bundle $T\mathcal{F}$ of $\mathcal{F}$ is non-degenerate and if, for any vector field X tangent to $\mathcal{F}$,

$$(L_X g)_{|T\mathcal{F}^\perp} = 0.$$

Notice that the integral curves of a nowhere lightlike and nowhere vanishing Killing vector field give a particular case of foliation having a bundle-like metric.

Definition 2.1. We will say that $\mathcal{F}$ is *transversely Lorentzian* if there exists an $\mathcal{F}$-bundle-like metric g which induces on the normal bundle $\nu(\mathcal{F}) = TM/T\mathcal{F}$ a Lorentzian metric.

We denote by $\mathrm{GL}(M, \mathcal{F})$ the transverse frame bundle of $\mathcal{F}$; it is a $\mathrm{GL}(q)$-principal bundle. We can lift the foliation $\mathcal{F}$ on a foliation $\mathcal{F}^1$ on $\mathrm{GL}(M, \mathcal{F})$ which has the same dimension but is of course of codimension $q + q^2$. Saying that $\mathcal{F}$ is transversely Lorentzian is equivalent to saying that $\mathrm{GL}(M, \mathcal{F})$ has an $\mathcal{F}^1$-foliated reduction to the Lorentzian group $\mathrm{O}(1, q-1)$, i.e. a reduction to the group $\mathrm{O}(1, q-1)$ of $\mathrm{GL}(M, \mathcal{F})$ which is $\mathcal{F}^1$-saturated. We will denote by $\mathrm{Lor}(\mathcal{F})$ this reduction.

CLAIM 2.2. *The transverse Lorentzian structure of $\mathcal{F}$ gives a natural transverse parallelism of $\mathcal{F}^1$ on* $\mathrm{GL}(M, \mathcal{F})$*, and hence on* $\mathrm{Lor}(\mathcal{F})$.

This follows from the existence of the Levi-Civita connection of the Lorentzian metric and from the application of the Molino theory for Riemannian foliations [**14**]. The construction is also recalled in the proof of our Proposition 2.8. Following Molino we introduce the following.

https://doi.org/10.1017/S0143385706000393 Published online by Cambridge University Press

Definition 2.3. Let $\mathcal{G}$ be a transversely parallelizable foliation of the manifold N and let $L_c(\mathcal{G})$ be the set of complete vector fields of N basic for the foliation $\mathcal{G}$. We will say that $\mathcal{G}$ is a *complete* transversely parallelizable foliation if for all points x in N the set $L_c(\mathcal{G})(x) = \{X(x), X \in L_c(\mathcal{G})\}$ spans a complement to $T\mathcal{G}$.

Definition 2.4. We will say that a Lorentzian foliation $\mathcal{F}$ is *transversely complete* if the foliation $\mathcal{F}^1$ on $\mathrm{Lor}(\mathcal{F})$ is a complete transversely parallelizable foliation.

Remark 2.5. The last definition can be equivalently stated with $\mathrm{GL}(M, \mathcal{F})$ instead of $\mathrm{Lor}(\mathcal{F})$. Indeed for all points $z \in \mathrm{GL}(M, \mathcal{F})$ and all $h \in \mathrm{GL}(q)$ let us denote by $R_h(z) = z.h$ the right action of $\mathrm{GL}(q)$ on $\mathrm{GL}(M, \mathcal{F})$. Then for each vector Z in $\mathrm{gl}(q)$, the Lie algebra of $\mathrm{GL}(q)$, we have a natural complete vector field on $\mathrm{GL}(M, \mathcal{F})$ tangent to the fibres of $\pi : \mathrm{GL}(M, \mathcal{F}) \to M$ given by

$$\widetilde{Z}(z) = \left.\frac{d}{dt}\right|_{t=0} z.\exp(tZ).$$

For every element z of $\mathrm{GL}(M, \mathcal{F})$, this identifies canonically $\mathrm{gl}(q)$ with the tangent space of the fibre of π at z. So the transverse completeness of a Lorentzian foliation depends on the existence of complete vector fields in $L_c(\mathcal{F}^1)$, transverse to $\mathcal{F}^1$ and to those fibres.

This remark implies also that Riemannian foliations are transversely complete since in this case $\mathrm{GL}(M, \mathcal{F})$ has an $\mathcal{F}^1$-foliated reduction to the group $\mathrm{O}(q)$ which is compact. A well-known property of transversely parallelizable complete foliations is the following (see [**14**]).

PROPOSITION 2.6. *Let $(N, \mathcal{G})$ be a transversely parallelizable complete foliation on a connected manifold N. Then the group* $\mathrm{Diff}(N, \mathcal{G})$ *of diffeomorphisms of N keeping $\mathcal{G}$ invariant (i.e. all elements of* $\mathrm{Diff}(N)$ *sending leaves of $\mathcal{G}$ onto leaves of $\mathcal{G}$), acts transitively on N.*

The proof follows from the fact that the complete parallelism ensures that the orbits of $\mathrm{Diff}(N, \mathcal{G})$ are open subsets of N. Thus there is only one orbit.

For Riemannian $\mathcal{F}$-bundle-like metrics, a geodesic starting orthogonally to $T\mathcal{F}$ remains orthogonal to it for all time, see [**4**]. This still holds for pseudo-Riemannian metrics. So speaking of the geodesic completeness of the orthogonal distribution $T\mathcal{F}^\perp$ makes sense.

Definition 2.7. A Lorentzian foliation $\mathcal{F}$ will be called *geodesically complete* if it admits a bundle-like Lorentzian metric for which the orthogonal distribution $T\mathcal{F}^\perp$ is geodesically complete.

This gives another notion of completeness. Be careful that the existence of an $\mathcal{F}$-bundle-like Lorentzian geodesically complete metric does not imply that all $\mathcal{F}$-bundle-like Lorentzian metrics are geodesically complete (see Proposition 5.16 in §5). The following proposition connects those definitions of completeness.

PROPOSITION 2.8. *A geodesically complete Lorentzian foliation is transversely complete in the sense of Definition 2.4.*

https://doi.org/10.1017/S0143385706000393 Published online by Cambridge University Press

Proof. A stronger statement actually holds: $(\mathrm{GL}(M,\mathcal{F}),\mathcal{F}^1)$ admits a q-tuple of basic, complete vector fields which are everywhere independent and transverse to the fibres and to the leaves of $\mathcal{F}^1$.

This is a foliated version of a classical result, [**12**, Proposition 6.5, p. 140]. Remark 2.5 implies that dealing with vector fields transverse to the fibres of $\pi : \mathrm{GL}(M,\mathcal{F}) \to M$ is sufficient. In the non-foliated case, i.e. here if $\dim\mathcal{F} = 0$, the metric on M provides a canonical (Levi-Civita) geometric connection of the fibre bundle. If $\dim\mathcal{F} \geq 1$, we deal with the principal bundle $\pi : \mathrm{GL}(M,\mathcal{F}) \to M$ of transverse frames to $\mathcal{F}$. We have to show that the bundle-like metric similarly provides a basic canonical supplement H to $T\mathcal{F}^1 \oplus \ker d\pi$ in $T\mathrm{GL}(M,\mathcal{F})$. Then, a remark and the proof of [**12**] conclude. So let us take $\mathcal{F}$ a geodesically complete Lorentzian foliation of codimension q and build H.

For all $z \in \mathrm{GL}(M,\mathcal{F})$, z is a transverse frame at $x = \pi(z)$, that is to say z is a linear isomorphism from $\mathbb{R}^q$ to $\nu_x(\mathcal{F})$. Let X be a tangent vector to M at x orthogonal to $\mathcal{F}$. We can find a C^1 curve $\gamma :]-\varepsilon,\varepsilon[\to M$ orthogonal to $\mathcal{F}$ and such that $\gamma(0) = x$ and $\gamma'(0) = X$. We denote by τ_γ the parallel transport along γ. Thus for each t in $]-\varepsilon,\varepsilon[$, the linear isomorphism $\tau_\gamma(t) \circ z$ is a transverse frame over $\gamma(t)$. Now we can set out

$$s_{x,z}(X) = \left.\frac{d}{dt}\right|_{t=0} \tau_\gamma(t) \circ z.$$

It is well defined since τ_γ depends smoothly on t and its 1-jet depends only on x, X and z. We set H_z to be $s_{x,z}(T\mathcal{F}^\perp)$. From the definition it satisfies $H_{z.h} = d_z R_h(H_z)$ for all $h \in \mathrm{GL}_q(\mathbb{R})$ (where R_h is the right translation by h). And we also have $d_z\pi(s_{x,z}(X)) = X$. Therefore, H_z is right invariant and transverse to the fibres and to the foliation $\mathcal{F}^1$ (the lift of $\mathcal{F}$ on $\mathrm{GL}(M,\mathcal{F})$). We can define a geometric connection P_z at z by setting $P_z = H_z \oplus T_z\mathcal{F}^1$.

Now 'standard' vector fields $(Z_i)_{i=1}^q$ can be built, using the distribution H and the fundamental form θ, see [**12**, Proposition 2.2, p. 119]. Recall that in our foliated case, θ is defined by $\theta_z(Z) = z^{-1}(\overline{d_z\pi(Z)})$ where $X \mapsto \overline{X}$ denotes the projection from TM to $\nu(\mathcal{F})$. Then for each i, the field Z_i is basic for $\mathcal{F}^1$ since so is θ and so is H. By [**12**, Proposition 6.3, p. 139], its integral curves are mapped by π onto geodesics of M, thus are complete by assumption. □

The reciprocal of Proposition 2.8 is false: the next section provides a counter-example to it.

2.2. *An example of a transversely complete but not geodesically complete Lorentzian foliation.*

PROPOSITION 2.9. *Let $\mathcal{F}$ be the strongly stable (or unstable) foliation of the Anosov flow obtained by suspending a hyperbolic diffeomorphism of the 2-torus. Then $\mathcal{F}$ is transversely Lorentzian complete in the sense of Definition 2.4, and not geodesically complete in the sense of Definition 2.7.*

Proof. The manifold obtained by this suspension is classically denoted by $\mathbb{T}^3_A$, with $A \in \mathrm{SL}_2(\mathbb{Z})$ the linear hyperbolic automorphism of $\mathbb{T}^2$ to which the initial diffeomorphism is isotopic. It is well known that $\mathcal{F}$ is a transversely affine Lie foliation, see [**3**].

https://doi.org/10.1017/S0143385706000393 Published online by Cambridge University Press

Hence it is transversely parallelizable and therefore both Riemannian and Lorentzian. It follows from Remark 2.5 that $\mathcal{F}$ is transversely complete. We are going to prove that $\mathcal{F}$ admits no geodesically complete bundle-like Lorentzian metric.

Let $\widetilde{\mathcal{F}}$ be the lift of $\mathcal{F}$ to the universal cover of $\mathbb{T}^3_A$. The leaves space of $\widetilde{\mathcal{F}}$ is diffeomorphic to $\mathbb{R}^2$. The fundamental group of $\mathbb{T}^3_A$, denoted by Γ, acts on this space. This action can be seen as generated by the transformations

$$(x, y) \mapsto (x + \tau, y) \quad \text{and} \quad (x, y) \mapsto (\lambda x, y + 1),$$

where τ belongs to a *dense* subgroup of $\mathbb{R}$ and λ is an irrational number greater than 1. A transverse metric for $\mathcal{F}$ is just a metric on $\mathbb{R}^2$ invariant by this action. We are going to prove that such metrics are never geodesically complete. As a first example we can notice that this action can be seen as the action of a subgroup of the group of isometries of a half Minkowski plane (with lightlike boundary) on it. This is certainly not a geodesically complete metric.

Unfortunately the space of transverse metrics of $\mathcal{F}$ is big, indeed $\mathcal{F}$ is transversely parallelizable and admits non-constant basic functions. To achieve the proof, we find incomplete lightlike geodesics, using a lemma of Carrière and Rozoy.

LEMMA 2.10. [**5**, p. 229] *Let $\mathcal{F}$ be one of the two lightlike foliations of a Lorentzian surface (up to a possible 2-cover, it is well defined) and F a compact leaf of $\mathcal{F}$. Suppose c is attractive (or repulsive, the sense of the parametrization has no meaning) on at least one of its sides. Then almost every leaf of $\mathcal{F}$ attracted by F is an incomplete lightlike geodesic.*

Let g be a Lorentzian metric on $\mathbb{R}^2$ invariant by the action of Γ. This metric has two transverse lightlike geodesic foliations, $\mathcal{G}^1$ and $\mathcal{G}^2$. Those foliations are invariant under Γ.

Let us first suppose that one of the lightlike foliations of g, for example $\mathcal{G}^1$, is everywhere transverse to the foliation $\mathcal{G}^0$ the leaves of which are the $\mathbb{R} \times \{y\}$. Note that this is in particular the case if one of those foliations is $\mathcal{G}^0$. This foliation is determined by one of its leaves, the other being deduced by the translations $(t, 0)$, for any $t \in \mathbb{R}$. It means that there exists a real number α such that for any $x \in \mathbb{R}$ the leaf of $\mathcal{G}^1$ through $(x, 0)$ contains the point $(x + \alpha, 1)$. Consequently the leaf containing $(\alpha/(\lambda - 1), 0)$ is fixed by the transformation $\gamma : (x, y) \mapsto (\lambda x, y + 1)$. Let us consider $\langle\gamma\rangle$, the group generated by γ. It acts isometrically, freely and properly on $\mathbb{R}^2$. Thus $\mathbb{R}^2/\langle\gamma\rangle$ is a cylinder endowed with the quotient metric $\bar{g}$. The foliation $\mathcal{G}^1$ gives a lightlike foliation $\overline{\mathcal{G}^1}$ which has a compact attractive leaf. By Lemma 2.10, $\bar{g}$, and therefore g, is not geodesically complete.

We suppose now that $\mathcal{G}^1$ is somewhere but not everywhere tangent to $\mathcal{G}^0$. Let U be the open set where $\mathcal{G}^1$ is transverse to $\mathcal{G}^0$ and V one connected component of U; as all the foliations are invariant by the translations $(x, y) \mapsto (x + \xi, y)$, $\xi \in \mathbb{R}$, so is V. Thus its boundary ∂V is a union of horizontal lines, leaves of $\mathcal{G}^0$. Take G^0 one of them; the foliation $\mathcal{G}^1$ is tangent to G^0 at every point of it, so by the uniqueness of the solutions of the ordinary differential equation defining $\mathcal{G}^1$, G^0 is a leaf of $\mathcal{G}^1$. Let $\langle\gamma'\rangle$ be the group generated by a translation $\gamma' : (x, y) \mapsto (x + \tau, y)$ belonging to Γ. In $\mathbb{R}^2/\langle\gamma'\rangle$, G^0 becomes a compact lightlike geodesic, which is necessarily attractive on the side of $V/\langle\gamma'\rangle$. Lemma 2.10 concludes again.

https://doi.org/10.1017/S0143385706000393 Published online by Cambridge University Press

We have proved that $\mathbb{R}^2$ has no Γ-invariant geodesically complete Lorentzian metric and therefore that $\mathcal{F}$ is not a geodesically complete Lorentzian foliation. □

3. *The alternative: Riemannian or Anosov*

THEOREM 3.1. *Let $\mathcal{F}$ be a one-dimensional complete transversely Lorentzian foliation on a compact connected manifold M of dimension three. Then either $\mathcal{F}$ is Riemannian (and thus transversely parallelizable) or up to a finite cover (in fact a 2-, 4- or 8-cover) the foliation $\mathcal{F}$ is given by the orbits of an Anosov flow.*

Proof. Up to a 2-cover, we suppose that $\mathcal{F}$ is orientable, and then parametrized by some flow φ^t. Both isotropic directions E_1 and E_2 of the Lorentzian metric span each fibre of the normal bundle $\nu(\mathcal{F}) = TM/T\mathcal{F}$. Necessarily, $d\varphi^t$ preserves or swaps them. Again up to a possible 2- of 4-cover (as O(1, 1) has four connected components) we also assume that the bundles E_1 and E_2 are *each one* preserved by $d\varphi^t$, and preserved with an orientation, i.e. they admit non-vanishing sections s_1 and s_2. We define the *deformation cocycle* $u(x, t)$ by

$$\varphi^t_*(s_1(x)) = e^{u(x,t)} s_1(\varphi^t(x)) \quad \text{and} \quad \varphi^t_*(s_2(x)) = e^{-u(x,t)} s_2(\varphi^t(x)),$$

where φ^t_* denotes the action of φ^t on $\nu(\mathcal{F})$, and where u satisfies

$$u(x, s + t) = u(x, t) + u(\varphi^t(x), s).$$

The leaves of $\mathcal{F}^1$ are given by the action of φ^t on $\mathrm{Lor}(\mathcal{F})$. More precisely the action of φ^t on M gives rise to an action on $\nu(\mathcal{F})$ and then to an action on $\mathrm{GL}(M, \mathcal{F})$ which stabilizes $\mathrm{Lor}(\mathcal{F})$. We will denote by φ^t_* this action.

The foliation $\mathcal{F}^1$ on $\mathrm{Lor}(\mathcal{F})$ is by Claim 2.2 a complete parallelizable foliation. Therefore, by Proposition 2.6, either all leaves of $\mathcal{F}^1$ are precompact in $\mathrm{Lor}(\mathcal{F})$ or no leaf is.

First case. The precompactness of leaves of $\mathcal{F}^1$ implies the completeness, see [**24**]. We then apply [**25**, Theorem 2] of Wolak, which shows that the foliation $\mathcal{F}$ is Riemannian. Normalizing s_1 and s_2 by a bundle-like Riemannian metric gives a transverse parallelism for $\mathcal{F}$.

Second case. We will show that the flow φ^t is Anosov with E_1 and E_2 (the projections on $\nu(\mathcal{F})$ of) its weak stable and unstable bundles. The idea is to show that there exists a leaf of $\mathcal{F}^1$ closed in $\mathrm{Lor}(\mathcal{F})$. Using Proposition 2.6 this will imply that every leaf of $\mathcal{F}^1$ is closed and using the structure theorem of Molino (see Theorem 3.5 below) the flow φ will be quasi-Anosov, hence Anosov by a result of Mañé.

We will say that a flow φ^t on a compact C^∞ manifold M is quasi-Anosov over a saturated subset K (or that K is quasi-Anosov for φ^t) if for every vector $v \in \nu(\mathcal{F})$ based on a point of K, $v \neq 0$, the set $\{\|\varphi^t_*(v)\|, t \in \mathbb{R}\}$ is unbounded. Here φ_* is the flow on $\nu(\mathcal{F})$ induced by the differential of φ^t and the foliation $\mathcal{F}$ is given by the orbits of φ. We will use the following proposition which is a weak version of a theorem of Mañé [**13**].

PROPOSITION 3.2. *Let K be a saturated connected compact subset of M such that φ is quasi-Anosov over K. Then K is hyperbolic for φ.*

https://doi.org/10.1017/S0143385706000393 Published online by Cambridge University Press

As Barbot told us, the arguments of [**13**] are remarkably simplified here. So they are worth being reproduced to give the following, elementary proof. We thank Barbot for it.

Proof. By assumption, no φ_*^t-orbit is bounded. Therefore, for any x in K, there is a real number $T(x)$ such that $u(x, T(x))$ is bigger than 1 (apply φ_*^t to $s_1(x)$). Applying similarly φ_*^t to $s_2(x)$, we find a $T'(x)$ such that $u(x, T'(x))$ is smaller than -1. By continuity of u and compactness of K we deduce the existence of some positive real number T such that:

$$\text{for all } x \in K, \quad \sup_{|t|\leq T} u(x,t) \geq 1 \quad \text{and} \quad \inf_{|t|\leq T} u(x,t) \leq -1.$$

Let us set $C = \sup_{0\leq t\leq T, x\in K} |u(x,t)|$.

LEMMA 3.3. *For any real positive numbers $0 \leq s \leq t$, and any point x of K, we have*

$$\min(-C, -C + u(x,t)) \leq u(x,s) \leq \max(C, u(x,t) + C).$$

Proof. Fix $t > 0$, $x \in K$ and denote by τ a time between 0 and t such that $u(x,\tau)$ is equal to $\sup_{0\leq s\leq t} u(x,s)$. By definition of T, the segment $[\tau - T, \tau + T]$ cannot be contained in $[0, t]$. Hence either $0 \leq \tau < T$, in which case we have $u(x,s) \leq u(x,\tau) \leq C$ for every $0 \leq s \leq t$, or $0 \leq t-\tau < T$, in which case $u(x,s) \leq u(x,\tau) = u(x,t)-u(\varphi^\tau(x), t-\tau) \leq u(x,t) + C$.

The proof of the lower bound is similar. □

At every point x of K, we obtain the following dichotomy:

– either $u(x,t)$ tends to $+\infty$ when t tends to $+\infty$;

– or there is some increasing sequence t_n converging to $+\infty$ for which $u(x,t_n)$ is uniformly bounded from above. By Lemma 3.3, in this case all the $u(x,s)$, for $0 \leq s$, are bounded from above.

Similarly, the $u(x,s)$, for $0 \leq s$, are bounded from below or converge to $-\infty$. Symmetrically, we obtain similar statements for t converging to $-\infty$.

Since the norm of $e^{u(x,t)}s_1(x)$ cannot be bounded from above for all times t, $u(x,t)$ must tend to $+\infty$ for t converging to $+\infty$ or $-\infty$. Furthermore, in the first case, $u(x,t)$ tends to $-\infty$ for negative t, and in the other case, $u(x,t)$ tends to $+\infty$ for negative t. (Hint: consider $e^{-u(x,t)}s_2(x)$.)

In other words, there is an $\varepsilon(x) = \pm 1$ such that $\lim_{t\to+\infty} \varepsilon(x)u(x,t) = +\infty$. By compactness of K, there is a positive real T' such that $\varepsilon(x)u(x,T') \geq 2$ for every x. It follows, by connectedness of K, that $\varepsilon(x)$ does not depend on x; let us say that $\varepsilon(x) = 1$ everywhere.

LEMMA 3.4. *If there exist two real numbers T' and a such that $\inf_{x\in K} u(x,T') \geq a > 0$, then K is hyperbolic for φ^t.*

Proof. Possibly reversing the time we can suppose that T' is positive.

https://doi.org/10.1017/S0143385706000393 Published online by Cambridge University Press

Let $C = \sup_{0\le t\le T', x\in K} |u(x,t)|$, it is a finite constant. For any positive t and any point $x \in K$, considering the unique integer n such that $nT' \le t < (n+1)T'$, we observe that

$$\begin{aligned} u(x,t) &= u(x,T') + u(\varphi^{T'}(x), T') + \cdots + u(\varphi^{nT'}(x), t - nT') \\ &\ge a(n-1) - C \\ &\ge a\frac{t}{T'} - C \\ &\ge a't + b' \quad \text{for some constants } a', b' \text{ with } a' > 0. \end{aligned}$$

Then, at x, vectors in E_1 are exponentially expanded in the future and exponentially contracted in the past. The fact that φ is transversely Lorentzian proves that, at x, elements of E_2 are exponentially contracted in the future and are exponentially expanded in the past.

It is well known that such a behaviour implies the Anosov property, but since some readers may be unfamiliar with this notion, we provide the proof here.

Let E_{11}, respectively E_{22}, be the pull-back of E_1, E_2, by the projection $TM \to \nu(\mathcal{F})$. The exponential behaviour of φ^t_* on E_1, E_2 implies the hyperbolic property as soon as there are, over K, some $D\varphi^t$-invariant one-dimensional subbundle E^{uu}, respectively E^{ss}, of E_{11}, respectively E_{22}, transverse to $\mathcal{F}$. To exhibit such invariant subbundles, for example E^{uu}, identify the space $\mathcal{E}$ of one-dimensional subbundles of E_{11} complementary to $\langle X\rangle = T\mathcal{F}$ (X is the vector field of the flow), with the space of continuous functions f on K *via* some splitting $E_{11} = \langle X\rangle \oplus \langle \eta\rangle$, where η is some vector field: f defines a subbundle by $E_f(x) = \langle f(x)X + \eta\rangle$.

The action of $D\varphi^t$ on E_{11} is defined by functions $u(x,t)$ and $\alpha(x,t)$ such that

$$\begin{aligned} D\varphi^t(x, \eta(x)) &= e^{u(x,t)}\eta(\varphi^t(x)) + \alpha(x,t)X(\varphi^t(x)) \\ D\varphi^t(x, X(x)) &= X(\varphi^t(x)). \end{aligned}$$

The natural action induced by $D\varphi^t$ on $\mathcal{E}$ defines a one-parameter group of transformations $\mathcal{A}_t$ of $C^0(K)$ given by

$$\mathcal{A}_t(f)(x) = \exp(-u(\varphi^{-t}(x), t))(f(\varphi^{-t}(x)) + \alpha(\varphi^{-t}(x), t)).$$

Since $u(x,T) \ge a > 0$, the operator $\mathcal{A}_T$ is contracting on $C^0(K)$ equipped with the sup-norm. It admits a unique fixed point f_0. Since all the $\mathcal{A}_t$ commute with $\mathcal{A}_T$, f_0 must be a common fixed point for all the $\mathcal{A}_t$. Then, E_{f_0} is the required invariant bundle. □

This lemma concludes the proof of Proposition 3.2. □

For the end of the proof we need the following.

THEOREM 3.5. (Molino [**14**, Theorem 4.2′, p. 134]) *Let $\mathcal{G}$ be a complete transversely parallelizable foliation on a connected manifold N. Then N is the total space of a fibre bundle $p : N \to W$ on some manifold W and the closures of the leaves of $\mathcal{G}$ are the fibres of p. Moreover, on each fibre of p, $\mathcal{G}$ is a transversely Lie foliation with dense leaves.*

Remark 3.6. It follows from Molino's proof of this theorem that the developing maps given by the transverse Lie structures on the closures of the leaves of $\mathcal{G}$ are complete, i.e. are fibre bundles.

https://doi.org/10.1017/S0143385706000393 Published online by Cambridge University Press

COROLLARY 3.7. *If there exists a closed non-compact leaf in* $\mathrm{Lor}(\mathcal{F})$*, then* φ *is Anosov.*

Proof. By Theorem 3.5 applied to $(\mathrm{Lor}(\mathcal{F}), \mathcal{F}^1)$, all leaves of $\mathcal{F}^1$ are the fibres of a fibre bundle $p : \mathrm{Lor}(\mathcal{F}) \to W$. As the foliation is given by a flow, this fibre bundle becomes an $\mathbb{R}$-principal bundle. The application $s : M \to \mathrm{Lor}(\mathcal{F}), x \mapsto (s_1(x), s_2(x))$ is a non-basic section of the principal bundle $\pi : \mathrm{Lor}(\mathcal{F}) \to M$. Using the continuity of the action of $\mathbb{R}$ (by φ^t_*) on $\mathrm{Lor}(\mathcal{F})$, the triviality of the fibre bundle $p : \mathrm{Lor}(\mathcal{F}) \to W$ (since $\mathbb{R}$ is contractible) and the compactness of $s(M)$, we prove that there exists a real number $T > 0$ such that $\varphi^T_*(s(M)) \cap s(M) = \emptyset$. Then by definition of T, $\inf_{x\in M} |u(x, T)| = a > 0$. By continuity we can suppose that $a = \inf_{x\in M} u(x, T)$. Applying Lemma 3.4 with $K = M$ we see that φ^t is Anosov. □

Finally, let L^1 be a leaf of $\mathcal{F}^1$. Theorem 3.5 and Remark 3.6 enable us to apply [**7**, Proposition 7-3] to $\mathcal{F}^1$ restricted to $\overline{L^1}$, which ensures that $\overline{L^1}$ is either compact or diffeomorphic to $\mathbb{R}$. As all leaves of $\mathcal{F}^1$ are supposed to be non-precompact, $L^1 = \overline{L^1} \cong \mathbb{R}$ and by Corollary 3.7 the flow φ is Anosov. □

4. *Classification*

4.1. *Anosov flows preserving a volume form are algebraic, after Ghys.* In [**8**], Ghys proves that on compact 3-manifolds and up to conjugation by a C^∞-diffeomorphism, the Anosov flows the weak stable and unstable distributions of which are C^∞ and which preserve a volume form (notice that those properties do not depend on their parametrization, and so are properties of the foliation) are exactly the '*algebraic Anosov flows*' i.e. the finite coverings and the finite quotients of the following:

- the geodesic flows of the unit tangent bundle of the hyperbolic compact surfaces;
- the suspensions of the linear hyperbolic diffeomorphisms A of the 2-torus (those give rise to a manifold denoted by $\mathbb{T}^3_A$).

Notice that both are transversely Lorentzian—the first ones with constant non-vanishing transverse curvature (positive or negative, it amounts to the same as the signature of the metric is (1, 1), 'neutral'), the second ones with a flat Lorentzian metric.

THEOREM 4.1. *Up to finite cover, a one-dimensional transversely complete Lorentzian foliation on a compact 3-manifold is either smoothly equivalent to an algebraic Anosov flow or Riemannian.*

Proof. The only thing we still have to prove is the completeness of an algebraic Anosov flow in the sense of Definition 2.4. This follows by Proposition 2.8. □

4.2. *With Carrière's work: the detailed panorama.* Finally, using Carrière's classification, provided in [**3**], of the Riemannian flows on the compact 3-manifolds, we detail the description of the foliations of Theorem 3.1, in the case when they are at once Lorentzian and Riemannian. This provides the following statement.

THEOREM 4.2. *Up to conjugation by a diffeomorphism, the one-dimensional transversely complete Lorentzian foliations of the compact 3-manifolds are:*

https://doi.org/10.1017/S0143385706000393 Published online by Cambridge University Press

(a) *up to a possible 2-, 4- or 8-covering, the algebraic Anosov flows on* $\mathbb{T}^3_A$ *and on the unit tangent bundle of the compact hyperbolic surfaces;*

(b) *the simultaneously Riemannian and Lorentzian ones. Up to a possible 2- or 4-covering, those are the parallelizable foliations of the compact 3-manifolds, which are, up to a possible orientation 2-covering:*
   (i) *the linear flows on a 3-torus,*
   (ii) *the flows defined by the fibres of a circle bundle over the 2-torus,*
   (iii) *the flows defined by the stable (respectively unstable) direction of the suspension of a linear hyperbolic diffeomorphism of the 2-torus.*

Of course, categories (b) (i) and (b) (ii) have a non-empty intersection.

Proof We first recall that a transversely Lorentzian and Riemannian foliation on a compact manifold is transversely complete Lorentzian (see Remark 2.5). Moreover, up to a possible 2- or 4-covering, foliations of codimension two and at once Riemannian and Lorentzian are parallelizable. Indeed, the bundle of transverse frames which are Riemannian orthonormal and Lorentzian orthogonal is a covering with fibre $\mathrm{O}(2, \mathbb{R}) \cap \mathrm{O}(1, 1) \simeq (\mathbb{Z}/2\mathbb{Z})^2$. As in Claim 2.2, the lift of $\mathcal{F}$ to this bundle is transversely parallelizable. Conversely, a parallelizable foliation preserves many pseudo-Riemannian or Riemannian metrics: take a transverse parallelism and say it is a (pseudo-)orthonormal basis.

Now we are left with classifying the parallelizable one-dimensional foliations. Taking again, if necessary, a 2-cover we can assume the foliation to be orientable. A dimension-one orientable foliation is usually called a flow yet without considering any parametrization. Instead of simply using Carrière's classification of the Riemannian flows on the compact 3-manifolds (see [**3**, part III A]), we pick up the relevant parts of his proof to achieve the desired list. Here is Carrière's ground theorem, from which follows his classification.

THEOREM 4.3. (Carrière [**3**, part II.C, Theorems 2 and 3]) *The closures of the leaves of a Riemannian flow* $\mathcal{F}$ *are tori, on which* $\mathcal{F}$ *induces a dense linear flow.*

Remark. In the very special case of a parallelizable flow, in which we are interested, this result follows immediately from (a more complete version of) the Molino Theorem 3.5 and from a result of Carrière's thesis (see [**3**, part II.C, Theorem 1]).

So let $\mathcal{F}$ be a parallelizable flow on a compact 3-manifold. After the Molino Theorem 3.5, the closure of its leaves have a common dimension and are fibres of a fibration on some compact manifold W. After Theorem 4.3, these fibres are tori and the study splits into three cases according to their dimension.

(1) There is a dense leaf of $\mathcal{F}$. Then $\mathcal{F}$ is a dense linear flow on the 3-torus. Conversely, such a flow is immediately parallelizable. This is the case (b) (i) of Theorem 4.2.

(2) All the leaves of $\mathcal{F}$ are closed. Then M is a circle bundle over a closed surface W. Since $\mathcal{F}$ is transversely Lorentzian, W admits some Lorentzian metric and thus it must be a 2-torus ($\mathcal{F}$ is transversely orientable). In this case, the leaves of $\mathcal{F}$ are defined by the fibres of a locally trivial fibration by circles over the 2-torus. Conversely, the foliation defined by such a fibration is automatically parallelizable. This is the case (b) (ii) of Theorem 4.2.

https://doi.org/10.1017/S0143385706000393 Published online by Cambridge University Press

(3) The closure of any leaf of $\mathcal{F}$ is a 2-torus. Then M is a torus bundle over a compact 1-manifold, thus over the circle. Such a bundle is obtained by considering the product $\mathbb{T}^2 \times [0, 1]$ and gluing the tori $\mathbb{T}^2 \times \{0\}$ and $\mathbb{T}^2 \times \{1\}$ by some diffeomorphism φ of $\mathbb{T}^2$. On each fibre $\mathbb{T}^2 \times \{s\}$, the flow $\mathcal{F}$ induces a linear flow of irrational direction Δ_s. But since Δ_s must be continuous in s, this direction is actually constant; we denote it by Δ. Thus φ lets the direction Δ be invariant. By [**10**, Lemma I.B.5], φ is in fact isotopic to a linear diffeomorphism A of $\mathbb{T}^2$, through an isotopy φ_t such that for all $t \in [0, 1]$, φ_t preserves the direction Δ. Our flow $\mathcal{F}$ is then smoothly conjugated to the flow induced by the translation of direction Δ on the manifold $\mathbb{T}^3_A$. Now a matrix A of $\mathrm{SL}_2(\mathbb{Z})$ preserving an irrational direction is either the identity or a hyperbolic matrix.

If $A = \mathrm{Id}$ we are again in the case (b) (i) of Theorem 4.2. Otherwise, Δ is the stable or the unstable direction of A and we are in the case (b) (iii). It remains to check that any flow of that last type is parallelizable. In fact, it is even transversely a Lie flow, modelled on the affine group $\mathrm{AG}(1, \mathbb{R})$ (see, for instance, [**3**, I.D. Example 6]). Thus any left-invariant frame on $\mathrm{AG}(1, \mathbb{R})$ yields a transversely invariant framefield. □

4.3. *Isometric foliations.* Among transversely Lorentzian one-dimensional foliations there are, in particular, those given by nowhere-vanishing spacelike Killing fields: the invariant metric of the manifold gives an invariant transverse metric on $(T\mathcal{F})^{\perp}$, canonically identified with $\nu(\mathcal{F})$. So let us define the following.

Definition 4.4. A transversely (pseudo-)Riemannian flow is called *isometric* if it admits a Killing parametrization for some bundle-like metric on the manifold.

We have the following natural characterization, see [**3**, Proposition III.B.1].

PROPOSITION 4.5. *A transversely pseudo-Riemannian flow is isometric if and only if it admits a parametrization which preserves a codimension-one transverse distribution.*

Moreover, an algebraic Anosov flow has smooth strongly stable and unstable distributions. Hence it preserves the distribution spanned by them and it is isometric. This proposition shows also that a flow which is both Lorentzian and Riemannian is Lorentzian isometric if and only if it is Riemannian isometric. Thus, thanks to [**3**, Corollary III.B.4], we see that the only non-isometric transversely complete Lorentzian flows on compact 3-manifolds are of type (b) (iii). Actually the nowhere-vanishing spacelike Killing fields of three-dimensional manifolds have been classified by Zeghib in [**27**]. It turns out that they all generate transversely complete Lorentzian foliations. It means that the examples given in § 6 are not isometric.

5. *The geodesically complete totally geodesic timelike foliations in dimension three*

Let us first specify what is a totally geodesible foliation of a pseudo-Riemannian manifold.

Definition 5.1. Let $\mathcal{G}$ be a codimension-one foliation on a manifold M. The foliation $\mathcal{G}$ is said to be *geodesible* if there exists a pseudo-Riemannian metric for which the leaves of $\mathcal{G}$ are totally geodesic submanifolds, i.e. any geodesic of the metric starting tangentially to a leaf stays in the leaf.

https://doi.org/10.1017/S0143385706000393 Published online by Cambridge University Press

In Riemannian geometry it is well known that a transversely oriented codimension-one foliation is geodesible if and only if it is transverse to a Riemannian flow. It is the starting point of the classification given by Carrière and Ghys in [**4**] for three-dimensional manifolds and in [**7**] in any dimension. We will see that the Lorentzian analogue is less general.

The signature of the restriction of the metric to some totally geodesic, connected submanifold N is constant, as any tangent space T_xN is stable by parallel transport along any path drawn on N. Thus, there exist three types of totally geodesic connected hypersurfaces in a Lorentzian manifold. They can be timelike, spacelike or lightlike if this restriction is respectively Lorentzian, Riemannian or degenerate. Hence we can talk about spacelike, timelike and lightlike totally geodesic foliations if all their leaves are of this type. (In general however, there may be leaves of each type, see [**26**] and [**15**].) The spacelike case is exactly the same as when M is Riemannian and is classified. The lightlike one has been principally studied by Zeghib in [**28**] (see also [**15**]) but a classification is only conjectured. In the general case a classification is out of reach for the moment (cf. [**15**]). Here we will look at the timelike case and to begin this study we specify the definition.

Definition 5.2. Let $\mathcal{G}$ be a codimension-one foliation on a manifold M. The foliation $\mathcal{G}$ is said to be *timelike geodesible* if there exists a Lorentzian metric g for which the leaves of $\mathcal{G}$ are totally geodesic, timelike submanifolds.

We state the following fact; its proof is exactly the same as the Riemannian one (see [**4**]).

PROPOSITION 5.3. *A smooth transversely oriented codimension-one foliation is timelike (respectively spacelike) totally geodesible if and only if it is transverse to a transversely Lorentzian (respectively Riemannian) flow.*

Or more precisely we have the following.

PROPOSITION 5.4. *Let (M, g) be a Lorentzian manifold. A smooth codimension-one timelike (or spacelike) foliation $\mathcal{G}$ is totally geodesic if and only if g is bundlelike for the foliation generated by its orthogonal distribution.*

As we have said, we restrict ourselves to smooth *timelike* totally geodesic foliations. They are not the only ones but they still represent an important and interesting family. Moreover, they are definitely related to Lorentzian flows. We further restrict ourselves to the geodesically complete case, i.e. when the geodesics tangent to the foliation are all complete. Thanks to Propositions 2.8 and 5.3 and Theorem 4.2 we can state the following.

COROLLARY 5.5. *Let $\mathcal{G}$ be a complete totally geodesic timelike codimension-one foliation on a closed 3-manifold M. Then, up to finite cover, one of the following situations occurs.*

(1) *The manifold M is a circle bundle over the torus and $\mathcal{G}$ is transverse to the fibres.*
(2) *The foliation $\mathcal{G}$ is transverse to the strong (un)stable direction of a flow defined by the suspension of a linear hyperbolic diffeomorphism of the 2-torus.*
(3) *The foliation $\mathcal{G}$ is transverse to an algebraic Anosov flow.*

Proof. By Proposition 5.3 our foliations are transverse to a Lorentzian flow; moreover, this flow is necessarily transversely complete by Proposition 2.8. So Theorem 4.2 gives

https://doi.org/10.1017/S0143385706000393 Published online by Cambridge University Press

more or less the list. The only thing to notice is that if $\mathcal{G}$ is transverse to a translation flow on a torus with irrational slope it is also transverse to a translation flow with rational slope. □

There exist foliations satisfying condition (1), (2) or (3); they are actually all timelike geodesible but it is not true that they can all be realized as *geodesically complete* timelike totally geodesic foliations. The first and second cases correspond to both spacelike and timelike totally geodesible foliations. Thus they have been studied by Carrière and Ghys in [**4**]. In the first case the foliations are given by the suspensions of pairs of commuting diffeomorphisms of the circle. We will see that they can be realized as geodesically complete timelike totally geodesic foliations. We will not try to classify them. In the second case, Carrière and Ghys showed that the foliations have no compact leaves and therefore, using [**10**], are diffeomorphically conjugated to the weak stable or unstable foliation of the Anosov flow. We have seen in §2.2 that even if they are transversely complete, the strongly stable and unstable foliations of the Anosov flow obtained by suspension do not have any geodesically complete bundle-like metrics. This construction will not give geodesically complete timelike totally geodesic foliations.

We are going to describe the foliations transverse to an algebraic Anosov flow, i.e. either to the geodesic flow of a hyperbolic surface Σ, and the ambient manifold is the unitary tangent bundle $T_1\Sigma$, or to the flow given by the suspension of a hyperbolic diffeomorphism of the torus A, and the ambient manifold is the torus bundle over the circle with monodromy A classically denoted by $\mathbb{T}^3_A$. Of course those flows are just considered as oriented one-dimensional foliations. In what follows $\mathcal{F}$ will denote one of the algebraic Anosov flows and $\mathcal{G}$ will denote a codimension-one foliation transverse to $\mathcal{F}$.

We are going to prove the following proposition.

PROPOSITION 5.6. *A codimension-one foliation $\mathcal{G}$ is transverse to an algebraic Anosov flow if and only if it is one of the following.*

(1) *It is conjugated to the weak stable or unstable foliation of an algebraic Anosov flow.*
(2) *$M = \mathbb{T}^3_A$ and $\mathcal{G}$ has a compact leaf and is without Reeb or type II components.*

To be able to give the definition of a type II component, we have to recall the construction of the two fundamental classes of examples of codimension-one foliations on $\mathbb{T}^2 \times [0, 1]$ tangent to the boundary (cf. the work of Moussu and Roussarie [**16**]).

We start with a foliation on the annulus $\mathbb{S}^1 \times [0, 1]$, invariant by rotation, tangent to the boundary and without compact leaves in the interior. There are two possibilities: this foliation is given by the suspension of a diffeomorphism of the interval without fixed point in the interior, or it is a Reeb component. Now let us consider the product foliation on the annulus times $[0, 1]$ and at last let us glue both sides of this product after some rotation. We obtain in this way two families of foliations on $\mathbb{T}^2 \times [0, 1]$ tangent to the boundary and without compact leaves in the interior. According to the choice of the rotation the non-compact leaves are all cylinders or all planes. Those foliations can also be obtained by the following 1-forms on $\mathbb{T}^2 \times [0, 1]$:

$$\Omega_1 = dr + \varphi(r)\omega \quad \text{and} \quad \Omega_2 = (1 - 2r)\, dr + \varphi(r)\omega,$$

https://doi.org/10.1017/S0143385706000393 Published online by Cambridge University Press

where $r \in [0, 1]$, ω is a linear form on the torus and φ is a smooth function such that $\varphi(0) = \varphi(1) = 0$ and $\varphi(r) > 0$ if $0 < r < 1$. So we denote them by $\mathcal{G}([\omega], 1)$ and $\mathcal{G}([\omega], 2)$.

We define another class of foliations. We start from the foliation $\mathcal{G}([\omega_0], 2)$ where the form ω_0 defines a trivial foliation by circles on $\mathbb{T}^2$ (i.e. where the rotation is the identity and of course the leaves of $\mathcal{G}([\omega_0], 2)$ are cylinders). There exists an orientation-preserving diffeomorphism of order two, without fixed points which permutes the connected components of the boundary and preserves the foliation. When we take the quotient we obtain a foliation on a manifold with boundary $\mathbb{T}^2$ foliated by cylinders accumulating on the boundary. In fact the manifold is the non-trivial fibre bundle over the Klein bottle by $[0, 1]$ (see [**21**] for details). Such a foliation will be called a cylindrical component.

Now we can set the following definition.

Definition 5.7. A foliation will be called a *component of type II* if it is topologically conjugated to a foliation $\mathcal{G}([\omega], 2)$ (i.e. obtained from a Reeb component on an annulus) or to a cylindrical component.

This definition is very close to the definition on [**21**, p. 104] but we do not ask the foliations to be with cylindrical leaves.

Definition 5.8. An embedded surface S of a 3-manifold M, not diffeomorphic to a 2-sphere, will be called *incompressible* if the map $\pi_1(S) \to \pi_1(M)$ is injective.

We recall that the Novikov theorem says that a codimension-one foliation on a compact 3-manifold has a compressible leaf if and only if it contains a Reeb component. Now we can start the proof of Proposition 5.6.

Proof of Proposition 5.6. Let us consider a foliation $\mathcal{G}$ transverse to an algebraic Anosov flow and prove that it is in case (1) or (2). We have seen in §4.1 that a transverse metric preserved by an algebraic Anosov flow $\mathcal{F}$ has constant curvature—it comes from the fact that $\mathcal{F}$ has a dense orbit. If $\mathcal{F}$ is a suspension flow the transverse metric has to be flat and if $\mathcal{F}$ is a geodesic flow the metric is with non-zero constant curvature (as seen in §4.1, its sign has no meaning on a two-dimensional Lorentzian manifold).

Moreover, if L is a compact leaf of $\mathcal{G}$, it is Lorentzian, so has to be a torus or a Klein bottle. But by the Gauss–Bonnet–Avez theorem (see [**1**]) it cannot be endowed with a metric with non-vanishing curvature. Hence a foliation transverse to the geodesic flow has no compact leaves. Consequently, we can use the work of Ghys [**9**] and see that $\mathcal{G}$ is then conjugated to the weak stable foliation of $\mathcal{F}$.

We still have to consider the suspension case. If $\mathcal{G}$ has no compact leaves then according to [**10**] it is conjugated to either the weak stable or unstable foliation of $\mathcal{F}$. Suppose now that the foliation has a compact leaf L.

FACT 5.9. *The foliation $\mathcal{G}$ contains neither Reeb nor type II components.*

Proof The first point is to know if a compact leaf L can separate the manifold, i.e. is $\mathbb{T}^3_A \setminus L$ connected?

https://doi.org/10.1017/S0143385706000393 Published online by Cambridge University Press

LEMMA 5.10. *A compact leaf L of $\mathcal{G}$ cannot separate $\mathbb{T}^3_A$.*

Proof. Suppose that L separates the manifold into two manifolds M_1 and M_2. As the Anosov flow is a flow, thus oriented, all its orbits cutting L go, for example, from M_1 to M_2. Thus any orbit of $\mathcal{F}$ cutting L is trapped in M_2. This is in contradiction to the fact that an algebraic Anosov flow possesses a dense set of closed orbits. □

Lemma 5.10 implies that $\mathcal{G}$ has no Reeb or cylindrical components. We still have to prove that $\mathcal{G}$ does not contain some component topologically conjugated to some $\mathcal{G}([\omega], 2)$. Taking again a possible finite cover we assume the Anosov flow transversely oriented. As the orientation of the compact leaves of such a foliation are opposite, such a component admits no transverse path joining its two compact leaves. As an algebraic Anosov flow has a dense leaf it cannot be transverse to it. This proves Fact 5.9. □

Remark 5.11. Fact 5.9 actually just says that the foliation $\mathcal{G}$ is *taut*, i.e. every compact leaf admits a closed transversal. Moreover, a foliation is taut if and only if there exists a metric (*a priori* Riemannian) for which the leaves are minimal surfaces (see [**22**]). Hence Riemannian totally geodesic foliations are taut but it can be verified that timelike totally geodesic foliations are also taut and Fact 5.9 is true for these foliations.

We are done with the direct part of Proposition 5.6. Let us now consider a foliation $\mathcal{G}$ as in point (1) or (2) of Proposition 5.6. Taking a possible 2-cover we assume that both $\mathcal{F}$ and $\mathcal{G}$ are oriented.

If $\mathcal{G}$ is as in point (1), we notice that the pull-back of the Anosov flow by the flow of a strong unstable (respectively stable) vector field is transverse to it and of course still Anosov.

If $\mathcal{G}$ is as in point (2), it is a foliation on $\mathbb{T}^3_A$ without Reeb or cylindrical components but with a compact leaf; it follows from [**11**, Theorem 2.1] that the compact leaf cannot separate the manifold. Hence if we cut along it we obtain a foliation on a manifold with incompressible—by the Novikov theorem and as $\mathcal{G}$ contains no Reeb component—non-connected boundary, i.e. $\mathbb{T}^2 \times [0, 1]$, according to [**6**, Theorem 4-2].

As the leaves are incompressible tori, we may reproduce the beginning of the proof of [**16**, Theorem 2], and see that $\mathcal{G}$ is isotopic to a foliation $\mathcal{G}'$ whose compact leaves are fibres of the fibration of $\mathbb{T}^3_A$ on $\mathbb{S}^1$. We use now [**16**, Theorem 1] in the following form.

THEOREM 5.12. (Moussu and Roussarie) *Let $\mathcal{G}$ be a foliation on $\mathbb{T}^2 \times [0, 1]$ tangent to the boundary and without compact leaves in the interior. Then there exists a homeomorphism of $\mathbb{T}^2 \times [0, 1]$ fixing the boundary which sends the foliation $\mathcal{G}$ to a model foliation $\mathcal{G}([\omega], 1)$ or $\mathcal{G}([\omega], 2)$.*

If follows from this theorem that there exists a homeomorphism of $\mathbb{T}^3_A$ which sends $\mathcal{G}$ transversely to the natural Anosov flow of $\mathbb{T}^3_A$. As being transverse is an open property and as for three-dimensional manifolds (see [**17**]) the set of diffeomorphisms is dense inside the set of homeomorphisms, we see that there is actually a diffeomorphism sending $\mathcal{G}$ transversely to the Anosov flow. This completes the proof of Proposition 5.6. □

We are going to prove now that the stable or unstable foliation $\mathcal{G}$ of an algebraic Anosov flow is never timelike totally geodesic and geodesically complete.

https://doi.org/10.1017/S0143385706000393 Published online by Cambridge University Press

According to Corollary 5.5 and the result of §2.2, the only case left we have to deal with is when $\mathcal{G}^{\perp}$ is an algebraic Anosov foliation we denote by $\mathcal{F}$. As the weak (un)stable foliations of two algebraic Anosov foliations on some manifold M are conjugated, we take φ a diffeomorphism of M such that $\mathcal{G} = \varphi(\mathcal{G}')$ with $\mathcal{G}'$ the weak (un)stable foliation of $\mathcal{F}$. The foliation $\mathcal{F}$ has a unique transverse Lorentzian (G, X) structure. Indeed, the lightlike foliations of $\mathcal{F}$ are its weak stable and unstable foliations: this determines the conformal class of the transverse Lorentzian metric. Moreover, $\mathcal{F}$ has dense leaves, so this metric is unique up to a scalar factor and has constant curvature. The transverse model space X is then the Minkowski plane or the universal cover of the two-dimensional de Sitter space. It means that the lift of $\mathcal{F}$ to the universal cover $\widetilde{M}$ of M is given by the preimages of a submersion δ from $\widetilde{M}$ to X called the developing map. As $\mathcal{F}$ is conjugated to an algebraic Anosov flow and as its transverse (G, X) structure is unique, this structure is conjugated to that of the algebraic flow, which is complete (see [**8**]), that is to say, δ is a locally trivial fibration with fibre $\mathbb{R}$.

The restriction of δ to the lift $\widetilde{L}$ of a leaf of $\mathcal{G}$ gives it a (G, X)-structure. If $\mathcal{G}$ is geodesically complete then the restriction of δ to $\widetilde{L}$ is a diffeomorphism. It means that the couple $(\mathcal{F}, \mathcal{G})$ must lift to the universal cover as a product foliation. We are going to prove that this last point is not possible.

As $\mathcal{G} = \varphi(\mathcal{G}')$ with $\mathcal{G}'$ the weak (un)stable foliation of $\mathcal{F}$, $\mathcal{G}$ has a cylindrical leaf $L_0 = \varphi(L_0')$. The lift $\widetilde{L_0'}$ of L_0' to $\widetilde{M}$ is a plane containing the lift of a compact leaf of $\mathcal{F}$, so is invariant by an element $\gamma \in \pi_1(M)$ sent by the holonomy representation ρ to an isometry $\rho(\gamma)$ of X having a fixed point. Now the lift $\widetilde{L_0}$ of L_0 on $\widetilde{M}$ is fixed by $\widetilde{\varphi}^{-1} \circ \gamma \circ \widetilde{\varphi} = \gamma' \in \pi_4(M)$, as the lift $\widetilde{\varphi}$ of φ to $\widetilde{M}$ normalizes the action of $\pi_1(M)$. Moreover, as $\gamma' \notin Z(\pi_1(M))$, $p(\gamma'))$ fixes a point of X. Eventually, as $\delta_{|\widetilde{L_0}} : \widetilde{L_0} \to X$ conjugates the action of γ' and of $\rho(\gamma')$, γ' fixes a point in $\widetilde{L_0} \subset \widetilde{M}$, which is impossible. It means that $\mathcal{G}$ cannot be timelike totally geodesic and geodesically complete.

The last point is to prove that the other cases are geodesically complete. This is done by noticing that, in those cases, the couple $(\mathcal{G}, \mathcal{F})$ lifts to the universal cover as a product and that $\mathcal{F}$ possesses a Lorentzian and geodesically complete transverse (G, X)-structure.

Remark 5.13. Actually those two conditions are necessary and sufficient. The main step to prove it is the following theorem proved in [**20**] by Ponge and Rieckzigel.

THEOREM 5.14. (Ponge and Rieckzigel) *If a timelike totally geodesic codimension-one foliation $\mathcal{G}$ is geodesically complete then the pair of foliations $\mathcal{G}$ and $\mathcal{G}^{\perp}$ lifts to the universal cover as a product.*

In particular this theorem says that the orthogonal foliation of a geodesically complete timelike codimension-one foliation admits a *complete* Lorentzian (G, X)-transverse structure. To get a reciprocal we have to ask the Lorentzian manifold X to be geodesically complete.

Setting together what we just showed with Corollary 5.5 and Proposition 5.6 we obtain the following theorem.

THEOREM 5.15. *Up to a finite cover the geodesically complete timelike geodesible codimension-one foliations of the closed three-dimensional manifolds are:*

https://doi.org/10.1017/S0143385706000393 Published online by Cambridge University Press

(1) *the foliations of the circle bundles over the torus, transverse to the fibres;*
(2) *the foliations on $\mathbb{T}^3_A$ with a compact leaf but without Reeb or type II components.*

By the way we showed also the following little result, announced after Definition 2.7.

PROPOSITION 5.16. *The algebraic Anosov flow $\mathcal{F}$ of $\mathbb{T}^3_A$ admits some bundle-like Lorentzian metrics such that $T\mathcal{F}^\perp$ is geodesically complete and some others such that it is not.*

Remarks. (1) In the second case the proof gives a description of the foliations. They contain a countable number of $\mathcal{G}([\omega], 1)$ components (with possibly different ω's) and a closed set of toral leaves. Moreover, up to a finite cover, circle bundles over the torus are also torus bundles over the circle. Hence in the first case of Theorem 5.15, the foliations having a compact leaf do have the same description but we have to consider also the foliations by planes and by cylinders.

(2) As we saw, a transversely Lorentzian flow $\mathcal{F}$ on a 3-manifold M is always (after a possible 2-cover) given by the intersection of two transverse codimension-one foliations $\mathcal{G}_1$ and $\mathcal{G}_2$. Hence a Lorentzian totally geodesic foliation $\mathcal{G}$ together with those two foliations form what is called a total foliation: $(\mathcal{G}, \mathcal{G}_1, \mathcal{G}_2)$, i.e. at any $p \in M$, $(T_p\mathcal{G})^\perp \oplus (T_p\mathcal{G}_1)^\perp \oplus (T_p\mathcal{G}_2)^\perp = T^*_pM$. What about the converse? Actually, we can give a Lorentzian geometric interpretation for $\mathcal{G}$ to be part of a total foliation of M. We recall that a diffeomorphism of a Lorentzian manifold (M, g) is conformal if and only if it preserves the lightcone of g. Therefore, up to a 2-cover, we see that a dimension-one foliation on a 3-manifold is transversely conformally Lorentzian if and only if it is given by the intersection of two transverse codimension-one foliations. As a codimension-one foliation is umbilical (i.e. its leaves are umbilic for a metric of the manifold) if and only if it is transverse to a transversely conformal flow (adapt the argument given in [**4**, Proposition I.3], we have that a codimension-one foliation $\mathcal{G}$ is umbilical with timelike leaves for some Lorentzian metric if and only if there exist two codimension-one foliations $\mathcal{G}_1$ and $\mathcal{G}_2$ such that $(\mathcal{G}, \mathcal{G}_1, \mathcal{G}_2)$ is a total foliation of M.

(3) There exists other timelike totally geodesic foliations on closed 3-manifolds: the foliations transverse to Lorentzian flows which are not transversely complete. We will give such an example below. It is not clear for the moment if it will be possible to classify them. If we remove the hypothesis of completeness, the only thing we know is that the foliations are taut, which is a weaker property. But it has interesting consequences. For example, as a taut foliation has no Reeb components we know that the manifold is covered by $\mathbb{R}^3$. We can also notice that the taut foliations on torus bundles over the circle are the foliations of Corollary 5.5. Hence they are all timelike totally geodesic. It means that we know all the timelike totally geodesic foliations of those manifolds.

6. *Non-transversely complete Lorentzian flows*

6.1. *The examples.* We are going to give the simplest example we know of a non-transversely complete Lorentzian flow. We will shortly say how to produce new examples from this one. We first construct it as a transversely conformally Lorentzian flat foliation and show that it is Lorentzian. For a clearer exposition, we replaced, when it was possible,

https://doi.org/10.1017/S0143385706000393 Published online by Cambridge University Press

technical details by drawings. A more precise and systematic study will be done in a forthcoming paper.

The torus $\mathbb{R}\mathrm{P}^1 \times \mathbb{R}\mathrm{P}^1$ endowed with the metric $d\theta\, d\varphi$ is called the Einstein torus and will be denoted by Ein_2. Its main particularity is the size of its conformal group: it is a finite index extension of $\mathrm{Diff}(\mathbb{S}^1) \times \mathrm{Diff}(\mathbb{S}^1)$. This implies that being only a transversely conformally Lorentzian flat codimension-two foliation is too general for us.

But the topological group $\mathrm{Diff}(\mathbb{S}^1) \times \mathrm{Diff}(\mathbb{S}^1)$ contains $\mathrm{PSL}(2,\mathbb{R}) \times \mathrm{PSL}(2,\mathbb{R})$ as a Lie subgroup. Thus we will rather construct our example among the foliations admitting a transverse $(\mathrm{PSL}(2,\mathbb{R}) \times \mathrm{PSL}(2,\mathbb{R}), \mathbb{R}\mathrm{P}^1 \times \mathbb{R}\mathrm{P}^1)$-structure. We will call them *transversely Einstein* foliations. We recall that such foliations are defined by an equivariant submersion D, called the developing map, from the universal cover of the ambient manifold M to the Einstein torus. Moreover, this submersion must be equivariant, that is there exists a morphism ρ from the fundamental group of M to $\mathrm{PSL}(2,\mathbb{R}) \times \mathrm{PSL}(2,\mathbb{R})$ such that $D \circ \gamma = \rho(\gamma) \circ D$ for any γ in the fundamental group. Sometimes the universal cover is replaced by an intermediate cover called the holonomy cover, which will be the case here.

Let us begin the construction. We denote by Σ_2 the compact orientable surface of genus two. In order to find a submersion from $\Sigma_2 \times \mathbb{R}$ to the Einstein torus, we first consider a projection $P = (P_1, P_2)$ from Σ_2 to Ein_2. We can easily find P such that the image of Σ_2 is given by Figure 1 and that any point in the interior of the image is regular (and has two preimages). We consider now a hyperbolic element A of $\mathrm{PSL}(2,\mathbb{R})$. There exists an element a of $\mathfrak{sl}(2,\mathbb{R})$ such that $A = \exp a$. In a well-chosen parametrization of $\mathbb{R}\mathrm{P}^1 \times \mathbb{R}\mathrm{P}^1$, and denoting by A_+ and A_-, respectively, the attractive and repulsive points at A, the orbits of $\exp(ta) \times \exp(-ta)$ on Ein_2 are the dotted curves of Figure 2. We can now define our submersion:

$$\begin{array}{rccc} & \Sigma_2 \times \mathbb{R} & \longrightarrow & \mathrm{Ein}_2 \\ D: & (x,t) & \mapsto & (\exp(ta)(P_1(x)), \exp(-ta)(P_2(x))). \end{array}$$

It is a submersion at any point (x,t) with $P(x)$ in the interior of $P(\Sigma_2)$, the grey zone in Figure 1. It is also a submersion if $P(x)$ is in the one-dimensional manifold $\mathcal{C}$ forming the boundary of $P(\Sigma_2)$. Indeed, on the one hand, $dD(x,t).(T_x\Sigma_2 \times \{0\}) = T_{P(x)}\mathcal{C}$, and on the other hand, as the orbits of $\exp(ta) \times \exp(-ta)$ are transverse to $\mathcal{C}$, $dD(x,t).(\{0\} \times \mathbb{R})$ is transverse to $T_{P(x)}\mathcal{C}$, this gives result. The transformation $(x,t) \mapsto (x, t+1)$ is conjugated by D to the transformation (A, A^{-1}) which lies in $\mathrm{PSL}(2,\mathbb{R}) \times \mathrm{PSL}(2,\mathbb{R})$. Hence the fibres of D go down to $\Sigma_2 \times \mathbb{R}/\mathbb{Z} \simeq \Sigma_2 \times \mathbb{S}^1$ and define on it a transversely Einstein foliation.

The image of D, denoted by U, is Ein_2 deprived of the 'vertical' circle $\{A_-\} \times \mathbb{R}\mathrm{P}^1$, containing the repulsive point $q = (A_-, A_+)$, and of the attractive point $p = (A_+, A_-)$. In order to prove that D actually defines a Lorentzian foliation we have to find a metric on U preserved by (A, A^{-1}). This will be done by finding U_A, the biggest subset of Ein_2 on which (A, A^{-1}) acts isometrically and proving that U is included in U_A. The points p and q are the attractive and repulsive fixed points of (A, A^{-1}) so they do not belong to U_A.

FACT 6.1. *Let A be a hyperbolic element of* $\mathrm{PSL}(2,\mathbb{R})$ *and let p and q be the attractive and repulsive fixed points of (A, A^{-1}) acting on $\mathbb{R}\mathrm{P}^1 \times \mathbb{R}\mathrm{P}^1$. There exists a Lorentzian*

https://doi.org/10.1017/S0143385706000393 Published online by Cambridge University Press




FIGURE 1. $P(\Sigma_2)$ in $\mathbb{R}P^1 \times \mathbb{R}P^1 = \mathrm{Ein}_2$.




FIGURE 2. The $P(B_i)$ are given for correspondence with Figure 3.

metric (in the conformal class of the Einstein metric) on $\mathbb{R}\mathrm{P}^1 \times \mathbb{R}\mathrm{P}^1 \setminus \{p, q\}$ *preserved by* (A, A^{-1}).

Proof. Let us first consider the following map:

$$\begin{array}{rccc} & \mathbb{R}^2 & \longrightarrow & \mathbb{R}/\pi\mathbb{Z} \times \mathbb{R}/\pi\mathbb{Z} \\ \psi: & (x, y) & \mapsto & (\arctan x, \arctan y). \end{array}$$

It is a diffeomorphism on its image and provides a conformal embedding of the Minkowski 2-space into the Einstein torus minus two lightlike circles. More precisely,

$$(\psi^{-1})^* \, dx \, dy = \frac{d\theta \, d\varphi}{\cos^2\theta \cos^2\varphi}.$$

Without any loss of generality we can suppose that there exists $\lambda \notin \{0, 1\}$ such that

$$f_A(x, y) = (\psi^{-1} \circ (A, A^{-1}) \circ \psi)(x, y) = \left(\lambda x, \frac{1}{\lambda} y\right)$$

https://doi.org/10.1017/S0143385706000393 Published online by Cambridge University Press

and so leaves the metric $dx\,dy$ invariant. It is easy to see that f_A leaves invariant any metric of the kind $\alpha(xy)\,dx\,dy$, where $\alpha : \mathbb{R} \to \mathbb{R}$ does not vanish. In particular f_A leaves invariant the metric $g_0 = 1/(1+x^2y^2)\,dx\,dy$, but

$$(\psi^{-1})^* \frac{1}{1+x^2y^2}\,dx\,dy = \frac{d\theta\,d\varphi}{\cos^2\theta\cos^2\varphi + \sin^2\theta\sin^2\varphi}.$$

This metric, *a priori* only defined on $\psi(\mathbb{R}^2)$, extends to $\mathbb{R}\mathrm{P}^1 \times \mathbb{R}\mathrm{P}^1 \setminus \{p, q\}$. Moreover, as $\psi(\mathbb{R}^2)$ is dense in $\mathbb{R}\mathrm{P}^1 \times \mathbb{R}\mathrm{P}^1 \setminus \{p, q\}$ the metric is preserved by (A, A^{-1}). □

Clearly U is included in U_A, hence D defines a Lorentzian foliation on $\Sigma_2 \times \mathbb{S}^1$. We denote it by $\mathcal{F}$. Let us remark that $\mathcal{F}$ has the following interesting properties: it is Lorentzian, transversely Einstein and radial (i.e. its holonomy fixes a point).

Let us tell now how to make more complicated examples from this one. First of all we can take $\hat{\Pi} : \widehat{\Sigma} \to \Sigma_2$ a finite cover and compose D by $\hat{\Pi}$. But we can do better: let us take θ a closed 1-form on $\widehat{\Sigma}$. Its pull-back $\widetilde{\Pi}^*\theta$ on the universal cover $\widetilde{\Sigma}$ is exact and therefore there exists a function Θ such that it is equal to $d\Theta$. We can replace the map D by D_θ defined by

$$D_\theta : \begin{array}{ccc} \widetilde{\Sigma} \times \mathbb{R} & \longrightarrow & \mathrm{Ein}_2 \\ (x,t) & \mapsto & (\exp((t+\Theta(x))a)(P_1 \circ \widetilde{\Pi}(x)), \exp(-(t+\Theta(x))a)(P_2 \circ \widetilde{\Pi}(x))). \end{array}$$

Those maps define a whole family of radial, transversely Einstein, Lorentzian foliations $\mathcal{F}_\theta$ on $\widehat{\Sigma}$. They depend on the cohomology class of θ. Indeed, if $\theta' = \theta + df$ with f a function of $\widehat{\Sigma}$, the change of variable $t' = t + f(x)$ conjugates the foliations. It will be shown in a further work that this family contains a lot of examples not conjugated with each other.

Let us describe now the dynamics of $\mathcal{F}$. On Figure 3, we draw the picture of $\mathcal{F}$ restricted to $W \times \mathbb{S}^1$, where W is a maximal open set of Σ_2 on which the projection P is a diffeomorphism. The set W is diffeomorphic to a disc with two holes and $\Sigma_2 \times \mathbb{S}^1$ is actually given by two copies of $W \times \mathbb{S}^1$ glued along their boundaries $B_1 \times \mathbb{S}^1$, $B_2 \times \mathbb{S}^1$, $B_3 \times \mathbb{S}^1$. We note that the B_i's are circles. On Figure 3, the vertical segments represent circles. We see that $\mathcal{F}$ restricted to $W \times \mathbb{S}^1$ has only one closed leaf with non-trivial holonomy. On $\Sigma_2 \times \mathbb{S}^1$ we have two interesting tori T_1 and T_2 given by the gluing of the dotted annuli of the picture, equal to $P^{-1}(\{A_+\} \times \mathbb{R}\mathrm{P}^1) \times \mathbb{S}^1$ and $P^{-1}(\mathbb{R}\mathrm{P}^1 \times \{A_+\}) \times \mathbb{S}^1$. Restricted to each of these tori $\mathcal{F}$ is a Reeb foliation with two closed leaves. Finally, we can see that the leaves of $\mathcal{F}$ not included in $T_1 \cup T_2$ are compact. As equicontinuity is a delicate notion for foliations, admitting slightly different, and non-equivalent, definitions, we give the definition we consider.

Definition 6.2. Let $\mathcal{F}$ be a one-dimensional foliation on a compact manifold M. Taking possibly a 2-cover of M, the leaves of $\mathcal{F}$ are the orbits of a flow φ without fixed point. We say that $\mathcal{F}$ is *(uniformly) equicontinuous* on an open $\mathcal{F}$-saturated subset U if $\{(\varphi_t)_{|U}, t \in \mathbb{R}\}$ is an (uniformly) equicontinuous set of diffeomorphisms of U.

The foliation $\mathcal{F}$ restricted to $(\Sigma_2 \times \mathbb{S}^1) \setminus (T_1 \cup T_2)$ is transversely parallelizable and so in particular equicontinuous. Moreover, $(\Sigma_2 \times \mathbb{S}^1) \setminus (T_1 \cup T_2)$ is an union of $\mathcal{F}$-saturated open subsets, precompact in it, and on which $\mathcal{F}$ is uniformly equicontinuous.

https://doi.org/10.1017/S0143385706000393 Published online by Cambridge University Press




FIGURE 3. The dynamics of $\mathcal{F}$.

Conversely on the T_i's the foliation is nowhere uniformly equicontinuous. So the open dense subset $(\Sigma_2 \times \mathbb{S}^1) \setminus (T_1 \cup T_2)$ is the domain of equicontinuity of $\mathcal{F}$. This kind of behaviour is not possible for an isometric flow (see for example [**27**, §3]); therefore this foliation is not isometric. Actually the isometric Lorentzian foliations of 3-manifolds are known, see [**27**], and are all transversely complete. We do not give any proof that the dynamics of $\mathcal{F}$ is as drawn in the picture, but it can be checked by the reader.

To conclude we construct a codimension-one foliation transverse to $\mathcal{F}$. We consider first the codimension-one foliation $\mathcal{G}_0$ of $W \times \mathbb{S}^1$ whose leaves are the $W \times \{t\}$. This foliation is transverse to $\mathcal{F}$ except along the $B_i \times \mathbb{S}^1$'s where it is everywhere tangent. Then we spin $\mathcal{G}_0$ along the three tori of the boundary (see [**2**, p. 84], for example). We do this spinning upwards along $B_3 \times \mathbb{S}^1$ and downwards along $B_1 \times \mathbb{S}^1$ and $B_2 \times \mathbb{S}^1$ in order to remain transverse to $\mathcal{F}$. We obtain a foliation $\mathcal{G}_1$ on $W \times \mathbb{S}^1$, tangent to the boundary, everywhere transverse to $\mathcal{F}$, and whose holonomy along the toral leaves has a trivial Taylor development. Consequently it is possible to glue two copies of $(W \times \mathbb{S}^1, \mathcal{G}_1)$ along their boundary. The obtained foliation $\mathcal{G}$ is a smooth codimension-one foliation, transverse to $\mathcal{F}$, with three toral leaves on $\Sigma_2 \times \mathbb{S}^1$. According to Proposition 5.3 there exists a Lorentzian metric on $\Sigma_2 \times \mathbb{S}^1$ for which $\mathcal{G}$ is timelike totally geodesic. This is the example announced at the end of §5. As the orthogonal of $\mathcal{G}$ is not transversely complete, its leaves are not geodesically complete. Actually we can see that the lightlike foliations induced on the three compact leaves have both compact and non-compact leaves and thus are geodesically incomplete by [**5**] (see Lemma 2.10 here). A similar construction is possible for the foliations $\mathcal{F}_\theta$.

6.2. *A result without any completeness assumption.* Up to a finite cover, a codimension-two Lorentzian foliation is given by the intersection of two codimension-one foliations, called its lightlike foliations. In the three-dimensional case, it is natural to wonder if those lightlike foliations may contain Reeb components. We prove the following.

https://doi.org/10.1017/S0143385706000393 Published online by Cambridge University Press

PROPOSITION 6.3. *Let M be a closed 3-manifold. Let $\mathcal{F}$ be a foliation of M given by the intersection of two codimension-one foliations $\mathcal{G}_1$ and $\mathcal{G}_2$. If $\mathcal{G}_1$ contains a Reeb component then $\mathcal{F}$ has no transverse volume form and therefore is not Lorentzian.*

Proof. This is a direct consequence of the results of Tamura and Sato [**23**] about foliations transverse to a Reeb component. They proved that such a foliation must contain a half Reeb component. It is not hard to see that it implies that $\mathcal{F}$ has an attractive leaf and therefore no transverse volume form. □

Let us remark that this result cannot be extended to taut foliations: the lightlike foliations of the transversely Einstein example given above are not taut: they have a separating toral leaf. Anyway, thanks to the classical theorems of Novikov (see [**18**]) and Palmeira (see [**19**]), we deduce the following theorem.

THEOREM 6.4. *Let M be a three-dimensional closed manifold. If M possesses a Lorentzian foliation $\mathcal{F}$ then its universal cover $\widetilde{M}$ is diffeomorphic to $\mathbb{R}^3$. Moreover, the leaves of $\widetilde{\mathcal{F}}$, the lift of $\mathcal{F}$ to $\widetilde{M}$, are all lines.*

Proof. We know that the lightlike foliations of $\mathcal{F}$ have no Reeb components; then, by Novikov's theorem, we deduce that they have no vanishing cycles or equivalently that their lift to the universal cover as foliations by planes. Palmeira's theorem affirms then that the universal cover of M is diffeomorphic to $\mathbb{R}^3$. To conclude we observe that $\widetilde{\mathcal{F}}$ is tangent to a foliation by planes and therefore its leaves are lines. □

Acknowledgements. The idea of this work was given us by a weekly workshop about conformally Lorentzian foliations at the École Normale Supérieure de Lyon in March–May 2003, together with Charles Frances, Thierry Barbot and Abdelghani Zeghib. We thank them for their contributions to this workshop, where we all learnt good mathematics. We also hail the referee for his perspicacity and thank him for his help in shortening some proofs.

REFERENCES

[1] A. Avez. Formule de Gauss–Bonnet–Chern en métrique de signature quelconque. *C. R. Acad. Sci. Paris* **255** (1962), 2049–2051.

[2] A. Candel and L. Conlon. *Foliations I (Graduate Studies in Mathematics, 23)*. American Mathematical Society, Providence, RI, 2000.

[3] Y. Carrière. Flots riemanniens (Structure transverse des feuilletages, Toulouse 1982). *Astérisque* **116** (1984), 31–52.

[4] Y. Carriére and É. Ghys. Feuilletages totalement géodesiques. *An. Acad. Brasil. Ciênc.* **53**(3) (1981), 427–432.

[5] Y. Carrière and L. Rozoy. Complètude des mètriques lorentziennes de T^2 et difféormorphismes du cercle. *Bol. Soc. Brasil. Mat. (N.S.)* **25**(2) (1994), 223–235.

[6] B. Evans and L. Moser. Solvable fundamental groups of compact 3-manifolds. *Trans. Amer. Math. Soc.* **168** (1972), 189–210.

[7] É. Ghys. Classification des feuilletages totalement géodésiques de codimension un. *Comment. Math. Helv.* **58** (1983), 543–572.

https://doi.org/10.1017/S0143385706000393 Published online by Cambridge University Press

[8] É. Ghys. Dèformations de flots d'Anosov et de groupes fuchsiens. *Ann. Inst. Fourier (Grenoble)* **42**(1–2) (1992), 209–247.
[9] É. Ghys. Rigidité différentiable des groupes fuchsiens. *Publ. Math. Inst. Hautes Études Sci.* **78** (1993), 164–185.
[10] É. Ghys and V. Sergiescu. Stabilité et conjugaison différentiable pour certains feuilletages. *Topology* **19** (1980), 179–197.
[11] S. Goodman. On the structure of foliated 3-manifolds separated by a compact leaf. *Invent. Math.* **39** (1977), 213–221.
[12] S. Kobayashi and K. Nomizu. *Foundations of Differential Geometry*. Vol. 1. Wiley Interscience, New York, 1963.
[13] R. Mañe. Quasi-Anosov diffeomorphisms and hyperbolic manifolds. *Trans. Amer. Math. Soc.* **229** (1977), 351–370.
[14] P. Molino. *Riemannian Foliations (Progress in Mathematics, 73)*. Birkhauser, Boston, MA, 1988.
[15] P. Mounoud. Feuilletages totalement géodésiques, flots riemanniens, et variétés de Seifert. *Ann. Inst. Fourier (Grenoble)* **55**(4) (2005), 1411–1438.
[16] R. Moussu and R. Roussarie. Relations de conjugaison et de cobordisme entre certains feuilletages. *Publ. Math. Inst. Hautes Études Sci.* **43** (1973), 143–168.
[17] J. Munkres. Obstructions to the smoothing of piecewise-differentiable homeomorphisms. *Ann. of Math.* **72** (1960), 521–554.
[18] S. P. Novikov. Topology of foliations. *Trans. Moscow Math. Soc.* **14** (1965), 248–278.
[19] C. F. B. Palmeira. Open manifolds foliated by planes. *Ann. of Math.* (2) **107**(1) (1978), 109–131.
[20] R. Ponge and H. Reckziegel. Twisted products in pseudo-Riemannian geometry. *Geom. Dedicata* **48**(1) (1993), 15–25.
[21] R. Roussarie. Plongements dans les variétés feuilletées et classification des feuilletages sans holonomies. *Publ. Math. Inst. Hautes Études Sci.* **43** (1973), 101–141.
[22] D. Sullivan. A homological characterization of foliations consisting of minimal surfaces. *Comment. Math. Helv.* **54**(2) (1979), 218–223.
[23] I. Tamura and A. Sato. On transverse foliations. *Publ. Math. Inst. Hautes Études Sci.* 54 (1981), 205–235.
[24] C. Tarquini. Feuilletages de type fini compact. *C. R. Math. Acad. Sci. Paris* **339**(3) (2004), 209–214.
[25] R. A. Wolak. Some remarks on equicontinuous foliations. *Ann. Univ. Sci. Budapest. Eötvös Sect. Math.* **41** (1999), 13–21.
[26] K. Yokumoto. Examples of Lorentzian geodesible foliations of closed three-manifolds having Heegaard splittings of genus one. *Tohoku Math. J. (2)* **56**(3) (2004), 423–443.
[27] A. Zeghib. Killing fields in compact Lorentz 3-manifolds. *J. Differential. Geom.* **43**(4) (1996), 859–894.
[28] A. Zeghib. Geodesic foliations in Lorentz 3-manifolds. *Comment. Math. Helv.* **74** (1999), 1–21.

https://doi.org/10.1017/S0143385706000393 Published online by Cambridge University Press