Bull. Aust. Math. Soc. **111** (2025), 497–504
doi:10.1017/S0004972724000819

NONMONOGENITY OF NUMBER FIELDS DEFINED BY TRUNCATED EXPONENTIAL POLYNOMIALS

ANUJ JAKHAR

(Received 9 June 2024; accepted 30 June 2024; first published online 8 October 2024)

Dedicated to Professor Sudesh K. Khanduja on her 74th birthday

Abstract

Let p be a prime number. Let $n \geq 2$ be an integer given by $n = p^{m_1} + p^{m_2} + \cdots + p^{m_r}$, where $0 \leq m_1 < m_2 < \cdots < m_r$ are integers. Let $a_0, a_1, \ldots, a_{n-1}$ be integers not divisible by p. Let $K = \mathbb{Q}(\theta)$ be an algebraic number field with $\theta \in \mathbb{C}$ a root of an irreducible polynomial $f(x) = \sum_{i=0}^{n-1} a_i x^i/i! + x^n/n!$ over the field $\mathbb{Q}$ of rationals. We prove that p divides the common index divisor of K if and only if $r > p$. In particular, if $r > p$, then K is always nonmonogenic. As an application, we show that if $n \geq 3$ is an odd integer such that $n - 1 \neq 2^s$ for $s \in \mathbb{Z}$ and K is a number field generated by a root of a truncated exponential Taylor polynomial of degree n, then K is always nonmonogenic.

2020 *Mathematics subject classification*: primary 11T24.

Keywords and phrases: monogenity, nonmonogenity, Newton polygon, power basis.

1. Introduction

Let $K = \mathbb{Q}(\theta)$ be an algebraic number field with θ in the ring $\mathbb{Z}_K$ of algebraic integers of K. Let $f(x)$ be the minimal polynomial of θ having degree n over the field $\mathbb{Q}$ of rational numbers. It is well known that $\mathbb{Z}_K$ is a free abelian group of rank n. A number field K is said to be monogenic if there exists some $\beta \in \mathbb{Z}_K$ such that $\mathbb{Z}_K = \mathbb{Z}[\beta]$. In this case, $\{1, \beta, \ldots, \beta^{n-1}\}$ is an integral basis of K; such an integral basis of K is called a power integral basis or briefly a power basis of K. If K does not possess any power basis, we say that K is nonmonogenic. Quadratic and cyclotomic fields are monogenic. In algebraic number theory, it is important to know whether a number field is monogenic or not. The first example of a nonmonogenic number field was given by Dedekind in 1878; he proved that the cubic field $\mathbb{Q}(\eta)$ is not monogenic when η is a root of the polynomial $x^3 - x^2 - 2x - 8$ (see [15, page 64]). The problems of

The author is thankful to the Indian Institute of Technology, Madras for NFIG grant RF/22-23/1035/MA/NFIG/009034.

© The Author(s), 2024. Published by Cambridge University Press on behalf of Australian Mathematical Publishing Association Inc.

https://doi.org/10.1017/S0004972724000819 Published online by Cambridge University Press

testing the monogenity of number fields and constructing power integral bases have been intensively studied (see [7] for an overview of the latest developments).

Throughout this paper, $\operatorname{ind}\theta$ denotes the index of the subgroup $\mathbb{Z}[\theta]$ in $\mathbb{Z}_K$ and $i(K)$ stands for the index of the field K defined by $i(K) = \gcd\{\operatorname{ind}\alpha \mid K = \mathbb{Q}(\alpha) \text{ and } \alpha \in \mathbb{Z}_K\}$. A prime number p dividing $i(K)$ is called a prime common index divisor of K. Note that if K is monogenic, then $i(K) = 1$. Therefore, a number field having a prime common index divisor is nonmonogenic. However, there exist nonmonogenic number fields having $i(K) = 1$, for example, $K = \mathbb{Q}(\sqrt[3]{175})$ is not monogenic and has $i(K) = 1$. Nakahara [14] studied the index of noncyclic but abelian biquadratic number fields. Gaál *et al.* [8] characterised the field indices of biquadratic number fields having Galois group V_4. Ahmad *et al.* [1, 2] proved that for a square free integer m not congruent to $\pm 1 \bmod 9$, a pure field $\mathbb{Q}(m^{1/6})$ having degree 6 over $\mathbb{Q}$ is monogenic when $m \equiv 2$ or $3 \bmod 4$ and it is nonmonogenic when $m \equiv 1 \bmod 4$. Gaál and Remete [9] studied monogenity of number fields of the type $\mathbb{Q}(m^{1/n})$ where $3 \leq n \leq 9$ and m is square free. Gaál [6] and Jakhar and Kaur [10] studied monogenity of number fields defined by some sextic irreducible trinomials.

Let $a_0, \ldots, a_{n-1}$ be integers. It is known that the polynomial

$$f(x) = a_0 + a_1 x + a_2 \frac{x^2}{2!} + \cdots + a_{n-1}\frac{x^{n-1}}{(n-1)!} + \frac{x^n}{n!} \tag{1.1}$$

of degree n is irreducible over $\mathbb{Q}$ if one of the following conditions is satisfied:

(1) $\gcd(a_0, n!) = 1$ (see [5, 16]);
(2) $\gcd(a_0 a_1 \cdots a_{n-1}, n) = 1$ (see [11, Theorem 1.2]).

Let p be a prime number. Let $n \geq 2$ be an integer given by $n = p^{m_1} + p^{m_2} + \cdots + p^{m_r}$, where $0 \leq m_1 < m_2 < \cdots < m_r$ are integers. Let $K = \mathbb{Q}(\theta)$ with θ a root of an irreducible polynomial $f(x)$ over $\mathbb{Q}$, where $f(x)$ is given by (1.1) and $a_0, \ldots, a_{n-1}$ are integers not divisible by p. We provide necessary and sufficient conditions so that $p \mid i(K)$ for $n \geq 2$. As an application, we give a family of number fields which are nonmonogenic. Precisely stated, we prove the following result.

THEOREM 1.1. *Let p be a prime number. Let $n \geq 2$ be an integer given by $n = p^{m_1} + p^{m_2} + \cdots + p^{m_r}$, where $0 \leq m_1 < m_2 < \cdots < m_r$ are integers. Let $a_0, a_1, \ldots, a_{n-1}$ be integers not divisible by p. Let $K = \mathbb{Q}(\theta)$ be an algebraic number field with θ a root of an irreducible polynomial $f(x) = x^n + n!\sum_{i=0}^{n-1} a_i x^i/i!$ over $\mathbb{Q}$. Then:*

(1) *$p\mathbb{Z}_K = \wp_1^{e_1} \cdots \wp_r^{e_r}$, where the $\wp_i$ are distinct prime ideals lying above the prime p with index of ramification $e_i = p^{m_i}$ and residual degree one for each i;*
(2) *p divides $i(K)$ if and only if $r > p$.*

In particular, if $r > p$, then K is always nonmonogenic.

The following corollary is an immediate consequence of the theorem.

https://doi.org/10.1017/S0004972724000819 Published online by Cambridge University Press

COROLLARY 1.2. *Let $n \geq 2$ be an integer with 2-adic expansion $n = 2^{m_1} + 2^{m_2} + \cdots + 2^{m_r}$, where $0 \leq m_1 < m_2 < \cdots < m_r$. Let $a_0, a_1, \ldots, a_{n-1}$ be odd integers. Let $K = \mathbb{Q}(\theta)$ be an algebraic number field with θ a root of an irreducible polynomial $f(x) = x^n + n! \sum_{i=0}^{n-1} a_i x^i/i!$ over $\mathbb{Q}$. If $r > 2$, then K is nonmonogenic.*

As an application of this corollary, we obtain the following result.

COROLLARY 1.3. *Let $n \geq 2$ be an integer with 2-adic expansion $n = 2^{m_1} + 2^{m_2} + \cdots + 2^{m_r}$, where $0 \leq m_1 < m_2 < \cdots < m_r$. Let $K = \mathbb{Q}(\theta)$ be an algebraic number field with θ a root of a truncated exponential Taylor polynomial $f(x) = 1 + x + x^2/2! + \cdots + x^n/n!$. Assume that $r \geq 3$. Then K is always nonmonogenic.*

EXAMPLE 1.4. This example provides a family of nonmonogenic algebraic number fields. Let $n \geq 3$ be an odd integer such that $n - 1 \neq 2^s$ for any $s \in \boldsymbol{N}$. If $K = \mathbb{Q}(\theta)$ is an algebraic number field with $\theta \in \mathbb{C}$ a root of $f(x) = \sum_{i=0}^{n} x^i/i!$, then K is nonmonogenic by Corollary 1.3.

REMARK 1.5. If we take $r < 3$, then K can be monogenic. For example, consider $n = 3$, $r = 2$ and $f(x) = x^3 + 3x^2 + 6x + 6$ in Corollary 1.3. It can be easily checked that the discriminant of $f(x)$ is $-2^3 \cdot 3^3$. Let $K = \mathbb{Q}(\theta)$ with θ a root of $f(x)$. Since $f(x)$ is an Eisenstein polynomial with respect to 3, in view of a basic result [12, Theorem 2.18], we see that $3 \nmid [\mathbb{Z}_K : \mathbb{Z}[\theta]]$. Further note that $f(x) \equiv x^2(x + 1) \pmod 2$. Hence, using Dedekind's criterion [12, page 78], it is easy to see that $2 \nmid [\mathbb{Z}_K : \mathbb{Z}[\theta]]$. Therefore, in view of the formula $D_f = [\mathbb{Z}_K : \mathbb{Z}[\theta]]^2 d_K$, where D_f denotes the discriminant of the polynomial $f(x)$ and d_K denotes the discriminant of K, it follows that $\mathbb{Z}_K = \mathbb{Z}[\theta]$. Hence, K is monogenic.

2. Preliminary results

Let $K = \mathbb{Q}(\theta)$ be an algebraic number field with θ a root of a monic irreducible polynomial $f(x)$ belonging to $\mathbb{Z}[x]$. In what follows, $\mathbb{Z}_K$ stands for the ring of algebraic integers of K. For a rational prime p, let $\mathbb{F}_p$ be the finite field with p elements and $\mathbb{Z}_p$ denote the ring of p-adic integers. Throughout the paper, $f(x) \to \overline{f(x)}$ stands for the canonical homomorphism from $\mathbb{Z}_p[x]$ onto $\mathbb{F}_p[x]$. For a prime p and a nonzero m belonging to the ring $\mathbb{Z}_p$ of p-adic integers, $v_p(m)$ denotes the highest power of p dividing m. The following lemma will play an important role in the proof of the theorem.

LEMMA 2.1 [15, Theorem 4.34]. *Let K be an algebraic number field and p be a rational prime. Then p is a prime common index divisor of K if and only if for some positive integer h, the number of distinct prime ideals of $\mathbb{Z}_K$ lying above p having residual degree h is greater than the number of monic irreducible polynomials of degree h in $\mathbb{F}_p[x]$.*

The following simple result will also be used. Its proof is omitted.

https://doi.org/10.1017/S0004972724000819 Published online by Cambridge University Press

LEMMA 2.2. *Let p be a prime number. If $n = c_0 + c_1 p + \cdots + c_r p^r$ is the representation of the positive integer n in base p with $0 \le c_i < p$ for each i, then*

$$v_p(n!) = \frac{n - (c_0 + c_1 + \cdots + c_r)}{p - 1}.$$

3. A short introduction to prime ideal factorisation based on Newton polygons

In 1894, Hensel developed a powerful approach for finding prime ideals of $\mathbb{Z}_K$ over a rational prime p. He showed that for every prime p, the prime ideals of $\mathbb{Z}_K$ lying above p are in one-to-one correspondence with monic irreducible factors of $f(x)$ in $\mathbb{Q}_p[x]$. Newton polygons are very helpful for finding the factors of $f(x)$ in $\mathbb{Q}_p[x]$. This is a standard method which is rather technical but efficient to apply. Therefore, we first introduce the notion of Gauss valuation and ϕ-Newton polygon, where $\phi(x)$ belonging to $\mathbb{Z}_p[x]$ is a monic polynomial with $\overline{\phi}(x)$ irreducible over $\mathbb{F}_p$.

DEFINITION 3.1. The Gauss valuation of the field $\mathbb{Q}_p(x)$ of rational functions in an indeterminate x extends the valuation v_p of $\mathbb{Q}_p$ and is defined on $\mathbb{Q}_p[x]$ by

$$v_{p,x}(a_0 + a_1 x + a_2 x^2 + \cdots + a_s x^s) = \min_{1 \le i \le s}\{v_p(a_i)\}, \quad a_i \in \mathbb{Q}_p.$$

DEFINITION 3.2. Let p be a rational prime. Let $\phi(x) \in \mathbb{Z}_p[x]$ be a monic polynomial which is irreducible modulo p and $f(x) \in \mathbb{Z}_p[x]$ be a monic polynomial not divisible by $\phi(x)$. Let $\sum_{i=0}^{n} a_i(x)\phi(x)^i$, with $\deg a_i(x) < \deg \phi(x)$, $a_n(x) \ne 0$, be the $\phi(x)$-expansion of $f(x)$ obtained by dividing $f(x)$ by the successive powers of $\phi(x)$. Let P_i stand for the point in the plane having coordinates $(i, v_{p,x}(a_{n-i}(x)))$ when $a_{n-i}(x) \ne 0$, $0 \le i \le n$. Let μ_{ij} denote the slope of the line joining the point P_i to P_j if $a_{n-i}(x)a_{n-j}(x) \ne 0$. Let i_1 be the largest positive index not exceeding n such that

$$\mu_{0i_1} = \min\{\mu_{0j} \mid 0 < j \le n,\ a_{n-j}(x) \ne 0\}.$$

If $i_1 < n$, let i_2 be the largest index such that $i_1 < i_2 \le n$ with

$$\mu_{i_1 i_2} = \min\{\mu_{i_1 j} \mid i_1 < j \le n,\ a_{n-j}(x) \ne 0\},$$

and so on. The ϕ-Newton polygon of $f(x)$ with respect to p is the polygonal path having segments $P_0P_{i_1}, P_{i_1}P_{i_2}, \ldots, P_{i_{k-1}}P_{i_k}$ with $i_k = n$. These segments are called the edges of the ϕ-Newton polygon and their slopes form a strictly increasing sequence; these slopes are nonnegative as $f(x)$ is a monic polynomial with coefficients in $\mathbb{Z}_p$.

DEFINITION 3.3. Let $\phi(x) \in \mathbb{Z}_p[x]$ be a monic polynomial which is irreducible modulo a rational prime p having a root α in the algebraic closure $\overline{\mathbb{Q}}_p$ of $\mathbb{Q}_p$. Let $f(x) \in \mathbb{Z}_p[x]$ be a monic polynomial not divisible by $\phi(x)$ whose $\phi(x)$-expansion is given by $\phi(x)^n + a_{n-1}(x)\phi(x)^{n-1} + \cdots + a_0(x)$ and such that $\overline{f}(x)$ is a power of $\overline{\phi}(x)$. Suppose that the ϕ-Newton polygon of $f(x)$ with respect to p consists of a single edge, say S, having positive slope l/e with l, e coprime, that is,

$$\min\left\{\frac{v_{p,x}(a_{n-i}(x))}{i} \;\middle|\; 1 \le i \le n\right\} = \frac{v_{p,x}(a_0(x))}{n} = \frac{l}{e},$$

https://doi.org/10.1017/S0004972724000819 Published online by Cambridge University Press

so that n is divisible by e, say $n = et$, and $v_{p,x}(a_{n-ej}(x)) \geq lj$ with $1 \leq j \leq t$. Thus, the polynomial $b_j(x) := a_{n-ej}(x)/p^{lj}$ has coefficients in $\mathbb{Z}_p$ and $b_j(\alpha) \in \mathbb{Z}_p[\alpha]$ for $1 \leq j \leq t$. The polynomial $T(Y)$ in the indeterminate Y defined by $T(Y) = Y^t + \sum_{j=1}^{t} \overline{b_j}(\overline{\alpha})Y^{t-j}$ with coefficients in $\mathbb{F}_p[\overline{\alpha}] \cong \mathbb{F}_p[x]/\langle \phi(x) \rangle$ is called the residual polynomial of $f(x)$ with respect to (ϕ, S).

The following weaker version of the theorem of the product, originally due to Ore, will be used in the proof of main result (see [4, Theorem 1.5], [13, Theorem 1.1]).

THEOREM 3.4. *Let $\phi(x) \in \mathbb{Z}_p[x]$ be a monic polynomial which is irreducible modulo a rational prime p having a root α in the algebraic closure $\overline{\mathbb{Q}}_p$ of $\mathbb{Q}_p$. Let $g(x) \in \mathbb{Z}_p[x]$ be a monic polynomial not divisible by $\phi(x)$ whose $\phi(x)$-expansion is given by $\phi(x)^n + a_{n-1}(x)\phi(x)^{n-1} + \cdots + a_0(x)$ and such that $\overline{f}(x)$ is a power of $\overline{\phi}(x)$. Suppose that the ϕ-Newton polygon of $g(x)$ with respect to the prime p has k edges $S_1, \ldots, S_k$ having slopes $\lambda_1 < \cdots < \lambda_k$. Then:*

(1) *$g(x) = g_1(x) \cdots g_k(x)$, where each $g_i(x) \in \mathbb{Z}_p[x]$ is a monic polynomial of degree $\ell_i \deg(\phi(x))$ and whose ϕ-Newton polygon has a single edge, say S_i', which is a translate of S_i such that ℓ_i is the length of the horizontal projection of S_i;*
(2) *the residual polynomial $T_i(Y) \in \mathbb{F}_p[\overline{\alpha}][Y]$ of $g_i(x)$ with respect to (ϕ, S_i') has degree ℓ_i/e_i, where e_i is the smallest positive integer such that $e_i\lambda_i \in \mathbb{Z}$.*

The next definition extends the notion of residual polynomial to more general polynomials $f(x)$.

DEFINITION 3.5. Let $p, \phi(x), \alpha$ be as in Definition 3.3. Let $g(x) \in \mathbb{Z}_p[x]$ be a monic polynomial not divisible by $\phi(x)$ such that $\overline{g}(x)$ is a power of $\overline{\phi}(x)$. Let $\lambda_1 < \cdots < \lambda_k$ be the slopes of the edges of the ϕ-Newton polygon of $g(x)$ and S_i denote the edge with slope λ_i. In view of Theorem 3.4, we can write $g(x) = g_1(x) \cdots g_k(x)$, where the ϕ-Newton polygon of $g_i(x) \in \mathbb{Z}_p[x]$ has a single edge, say S_i', which is a translate of S_i. Let $T_i(Y)$ belonging to $\mathbb{F}_p[\overline{\alpha}][Y]$ denote the residual polynomial of $g_i(x)$ with respect to (ϕ, S_i') as in Definition 3.3. For convenience, the polynomial $T_i(Y)$ will be referred to as the residual polynomial of $g(x)$ with respect to (ϕ, S_i). The polynomial $g(x)$ is said to be p-regular with respect to ϕ if none of the polynomials $T_i(Y)$ has a repeated root in the algebraic closure of $\mathbb{F}_p$, $1 \leq i \leq k$. In general, if $f(x)$ belonging to $\mathbb{Z}_p[x]$ is a monic polynomial and $\overline{f}(x) = \overline{\phi}_1(x)^{e_1} \cdots \overline{\phi}_r(x)^{e_r}$ is its factorisation modulo p into irreducible polynomials with each $\phi_i(x)$ belonging to $\mathbb{Z}_p[x]$ monic and $e_i > 0$, then by Hensel's lemma [3, Ch. 4, Section 3], there exist monic polynomials $f_1(x), \ldots, f_r(x)$ belonging to $\mathbb{Z}_p[x]$ such that $f(x) = f_1(x) \cdots f_r(x)$ and $\overline{f}_i(x) = \overline{\phi}_i(x)^{e_i}$ for each i. The polynomial $f(x)$ is said to be p-regular (with respect to $\phi_1, \ldots, \phi_r$) if each $f_i(x)$ is p-regular with respect to ϕ_i.

We provide a simple example of a p-regular polynomial with respect to any monic polynomial $\phi(x) \in \mathbb{Z}[x]$ which is irreducible modulo a prime p.

https://doi.org/10.1017/S0004972724000819 Published online by Cambridge University Press

EXAMPLE 3.6. If $p, \phi(x)$ are as above and $g(x) \neq \phi(x)$ belonging to $\mathbb{Z}_p[x]$ is a monic polynomial with $\overline{g}(x) = \overline{\phi}(x)$, then the ϕ-Newton polygon of $g(x)$ with respect to p is a line segment S joining the point $(0, 0)$ with $(1, b)$ for some $b > 0$. Consequently, the polynomial associated to $g(x)$ with respect to (ϕ, S) is linear and $g(x)$ is p-regular with respect to ϕ.

To determine the number of distinct prime ideals of $\mathbb{Z}_K$ lying above a rational prime p, we will use the following theorem which is a weaker version of [13, Theorem 1.2].

THEOREM 3.7. *Let $L = \mathbb{Q}(\xi)$ be an algebraic number field with ξ satisfying an irreducible polynomial $g(x) \in \mathbb{Z}[x]$ and p be a rational prime. Let $\overline{\phi}_1(x)^{e_1} \cdots \overline{\phi}_r(x)^{e_r}$ be the factorisation of $g(x)$ modulo p into powers of distinct irreducible polynomials over $\mathbb{F}_p$ with each $\phi_i(x) \neq g(x)$ belonging to $\mathbb{Z}[x]$ monic. Suppose that the ϕ_i-Newton polygon of $g(x)$ has k_i edges, say S_{ij}, having slopes $\lambda_{ij} = l_{ij}/e_{ij}$ with $\gcd(l_{ij}, e_{ij}) = 1$ for $1 \leq j \leq k_i$. If $T_{ij}(Y) = \prod_{s=1}^{s_{ij}} U_{ijs}(Y)$ is the factorisation of the residual polynomial $T_{ij}(Y)$ into distinct irreducible factors over $\mathbb{F}_p$ with respect to (ϕ_i, S_{ij}) for $1 \leq j \leq k_i$, then*

$$p\mathbb{Z}_L = \prod_{i=1}^{r} \prod_{j=1}^{k_i} \prod_{s=1}^{s_{ij}} \mathfrak{p}_{ijs}^{e_{ij}},$$

where $\mathfrak{p}_{ijs}$ are distinct prime ideals of $\mathbb{Z}_L$ having residual degree $\deg \phi_i(x) \cdot \deg U_{ijs}(Y)$.

4. Proof of Theorem 1.1

PROOF. Observe that $p \leq n$. We first show that x is the only repeated factor of $f(x)$ modulo p. If $p \mid n$, then clearly $f(x) \equiv x^n \pmod{p}$. If $p \nmid n$, then assume that $j, 0 \leq j \leq n-2$, is the smallest index such that p divides $n-j$. Keeping in mind that $p \nmid a_i$, we see that $f(x)$ is congruent to

$$\begin{aligned} &x^n + na_{n-1}x^{n-1} + \cdots + a_{n-j}\frac{n!}{(n-j)!}x^{n-j} \\ &\quad \equiv x^{n-j}\left(x^j + na_{n-1}x^{j-1} + \cdots + a_{n-j}\frac{n!}{(n-j)!}\right) \bmod p. \end{aligned}$$

Note that $p \nmid j$. Otherwise, if $p \mid j$, then since $p \mid (n-j)$, we have $p \mid n$, which is a contradiction. Hence, the polynomial $x^j + \overline{n}\overline{a}_{n-1}x^{j-1} + \cdots + \overline{a}_{n-j}\overline{n!/(n-j)!}$ belonging to $\mathbb{Z}/p\mathbb{Z}[x]$ is a separable polynomial. It follows that x is the only repeated factor of $f(x)$ modulo p.

Now we show that $f(x)$ is p-regular with respect to $\phi(x) = x$. Recall that $p \nmid a_i$. By the definition of the p-Newton polygon, we see that it will be the polygonal path formed by the lower edges along the convex hull of the points of the set S defined by

$$S = \left\{ \left(i, v_p\left(\frac{n!}{(n-i)!}\right)\right) \,\middle|\, 0 \leq i \leq n \right\}.$$

https://doi.org/10.1017/S0004972724000819 Published online by Cambridge University Press

By hypothesis, $n = p^{m_1} + p^{m_2} + \cdots + p^{m_r}$, where $0 \le m_1 < m_2 < \cdots < m_r$. Let ℓ_i denote the integer

$$\ell_i = p^{m_1} + \cdots + p^{m_i}, \quad 1 \le i \le r.$$

Set $\ell_0 = 0$. As in [5], using Lemma 2.2 and keeping in mind that $v_p(a_i) = 0$ for each i, it can be easily checked that the p-Newton polygon of $f(x)$ consists of r edges, and the ith edge is the line segment having vertices $(\ell_{i-1}, v_p(n!/(n-\ell_{i-1})!))$ and $(\ell_i, v_p(n!/(n-\ell_i)!))$. So by Lemma 2.2, the slope λ_i of the ith edge of the p-Newton polygon of $f(x)$ is

$$\lambda_i = \frac{-v_p((n-\ell_i)!) + v_p((n-\ell_{i-1})!)}{\ell_i - \ell_{i-1}} = \frac{\ell_i - \ell_{i-1} - 1}{(\ell_i - \ell_{i-1})(p-1)} = \frac{p^{m_i} - 1}{p^{m_i}(p-1)}.$$

Observe that $f(x)$ can have an edge with slope zero if and only if $m_1 = 0$. Also, m_1 can be zero only when $p \nmid n$. Therefore, in view of Hensel's lemma and Theorem 3.4, we can write $f(x) = g_1(x) \cdots g_r(x)$, where $g_i(x) \in \mathbb{Z}_p[x]$ has degree $\ell_i - \ell_{i-1} = p^{m_i}$ and the p-Newton polygon of $g_i(x)$ has a single edge, say S_i, with slope λ_i. When $\lambda_i > 0$, the polynomial, say $T_i(y) \in \mathbb{F}_p[y]$, associated to $g_i(x)$ with respect to (x, S_i) is linear. Hence, $f(x)$ is p-regular with respect to $\phi(x) = x$. So, by Theorem 3.7,

$$p\mathbb{Z}_K = \wp_1^{e_1} \cdots \wp_r^{e_r},$$

where the $\wp_i$ are distinct prime ideals lying above prime p with index of ramification $e_i = p^{m_i}$ and residual degree one for each i. Hence, by Lemma 2.1, $p \mid i(K)$ if and only if $r > p$. This completes the proof of the theorem. □

Acknowledgement

The author appreciates the anonymous referee's suggestions, which have enhanced the quality of this paper.

References

[1] S. Ahmad, T. Nakahara and A. Hameed, 'On certain pure sextic fields related to a problem of Hasse', *Internat. J. Algebra Comput.* **26**(3) (2016), 577–583.
[2] S. Ahmad, T. Nakahara and S. M. Husnine, 'Power integral basis for certain pure sextic fields', *Int. J. Number Theory* **10**(8) (2014), 2257–2265.
[3] Z. I. Borevich and I. R. Shafarevich, *Number Theory* (Academic Press, New York, 1966).
[4] S. D. Cohen, A. Movahhedi and A. Salinier, 'Factorization over local fields and the irreducibility of generalized difference polynomials', *Mathematika* **47** (2000), 173–196.
[5] R. F. Coleman, 'On the Galois groups of the exponential Taylor polynomials', *Enseign. Math.* **33** (1987), 183–189.
[6] I. Gaál, 'An experiment on the monogenity of a family of trinomials', *JP J. Algebra Number Theory Appl.* **51**(1) (2021), 97–111.
[7] I. Gaál, 'Monogenity and power integral bases: recent developments', *Axioms* **13** (2024), Article no. 429.
[8] I. Gaál, A. Pethö and M. Pohst, 'On the indices of biquadratic number fields having Galois group V_4', *Arch. Math.* **57** (1991), 357–361.

https://doi.org/10.1017/S0004972724000819 Published online by Cambridge University Press

[9] I. Gaál and L. Remete, 'Power integral bases and monogenity of pure fields', *J. Number Theory* **173** (2017), 129–146.
[10] A. Jakhar and S. Kaur, 'A note on non-monogenity of number fields arising from sextic trinomials', *Quaest. Math.* **46** (2023), 833–840.
[11] A. Jindal and S. K. Khanduja, 'An extension of Schur's irreducibility result', Preprint, 2023, arXiv:2305.04781.
[12] S. K. Khanduja, *A Textbook of Algebraic Number Theory*, UNITEXT Series, 135 (Springer, Singapore, 2022).
[13] S. K. Khanduja and S. Kumar, 'On prolongations of valuations via Newton polygons and liftings of polynomials', *J. Pure Appl. Algebra* **216** (2012), 2648–2656.
[14] T. Nakahara, 'On the indices and integral bases of non-cyclic but abelian biquadratic fields', *Arch. Math.* **41** (1983), 504–508
[15] W. Narkiewicz, *Elementary and Analytic Theory of Algebraic Numbers*, 3rd edn, Springer Monographs in Mathematics (Springer-Verlag, Berlin, 2004).
[16] I. Schur, 'Einige sätze über primzahlen mit anwendungen auf irreduzibilitätsfragen I', *Sitzungsber. Preussischen Akad. Wiss. Phys.-Math. Kl.* **14** (1929), 125–136.

ANUJ JAKHAR, Indian Institute of Technology (IIT) Madras, Chennai, India
e-mail: anujjakhar@iitm.ac.in, anujiisermohali@gmail.com

https://doi.org/10.1017/S0004972724000819 Published online by Cambridge University Press